Title: Diet-induced changes to host gut microbiota is linked to foraging innovation in a wild bird

Authors: Gabrielle L. Davidson[a,b,*] , Niamh Wiley[c,d], Amy C. Cooke[a], Crystal N. Johnson[c,d], Fiona
Fouhy[c,d], Michael S. Reichert[a, e], Iván de la Hera[a, e], Jodie M.S. Crane[a,f], Ipek G. Kulahci[a], R. Paul
Ross[c,d], Catherine Stanton[c,d], John L. Quinn[a,*]

* Correspondence: Gabrielle Davidson email: gd339@cam.ac.uk; John Quinn email: j.quinn@ucc.ie.

Tel: +44(0)1223 747321

a. School of Biological, Earth and Environmental Sciences, Distillery Fields, North Mall,
8   University College Cork, Cork, Ireland.
b. Department of Psychology, Downing Street, University of Cambridge, Cambridge, UK.
c. APC Microbiome Ireland, University College Cork, Cork, Ireland.
d. Teagasc Food Research Centre, Moorepark, Fermoy, Ireland.
e. Department of Integrative Biology, Oklahoma State University, USA
f. Kākāpō Recovery Programme, Department of Conservation, 7th Floor, 33 Don Street,
Invercargill 9810, New Zealand.

Keywords: gut microbiome, gut-brain axis, diet, innovation, problem solving performance, great tits,
foraging, plasticity.

ABSTRACT: The microbial community in the gut is influenced by environmental factors, especially diet, which can moderate host behaviour through the microbiome-gut-brain axis. However, the ecological relevance of microbiome-mediated behavioural plasticity in wild animals is unknown. We presented wild-caught great tits (*Parus major*) with a problem-solving task and showed that performance was weakly associated with variation in the gut microbiome. We then manipulated the gut microbiome by feeding birds one of two diets that differed in their relative levels of fat, protein and fibre content: an insect diet (low content), or a seed diet (high content). Microbial communities were less diverse among individuals given the insect compared to those on the seed diet. Individuals were less likely to problem-solve after being given the insect diet, and the same microbiota metrics that were altered as a consequence of diet were also those that correlated with variation in problem solving performance. Although the effect on problem-solving behaviour could have been caused by motivational or nutritional differences between our treatments, our results nevertheless support the idea that dietary induced changes in the gut microbiota could be an important mechanism underlying individual behavioural plasticity in wild populations.

INTRODUCTION

The enteric microbial community, frequently referred to as the gut microbiome, is an important ecosystem that contributes to host behaviour [1,2]. Recent evidence points to a bidirectional communication link between the host's brain and their own gut microbiome, known as the microbiome-gut-brain axis [3,4]. Experimental alteration of the gut microbiome can impact important behaviours and cognition, including learning, memory, anxiety, activity levels and social interactions [3,5-8], as well as cause changes to neurogenesis [9] and protein expression in the brain [5,7,10]. Neurotransmitters and short chain fatty acids released by microbes act as signals that can be communicated to the brain [reviewed in 1] and the extent to which signalling occurs is dependent on the microbial taxa present [11]. Evidence of the microbiome-gut-brain axis affecting behaviour is limited to experiments on model laboratory animals and to correlational studies on mental health in humans [1,4]. Little is known about whether these findings can be applied to natural populations, where microbiome-host interactions are predicted to have important effects on traits that directly impact animal fitness [12], such as cognition [13] and foraging behaviour [14].

Several environmental factors have been shown to contribute to enteric microbial community composition in wild vertebrates, including geographical location [15,16], habitat characteristics [17,18], and seasonality [19-21]. These temporal and spatial differences in the microbiome, between and within populations, also correlate with variation in diet [19,22]. Reports of direct effects of dietary manipulations on the gut microbiota in birds to date have been largely restricted to poultry [23], and a recent study in wild-caught house sparrows demonstrated diversity and family-level changes to microbiota as a consequence of dietary manipulation [24]. Changes in diet can also lead to a significant alteration to the gut microbiome within 24 hours in humans [25] where changes are dependent on dietary features such as protein levels [25,26], fat levels [27], and whether plants are

present in the diet [28,29].Youngblut et al. [30] found diet explained a high proportion of variation in gut microbiota across vertebrate species including a subset of avian species. Hird et al [31] also showed that variation in gut microbiome across species was explained by diet in neotropical birds. . Thus, microbiome plasticity should be particularly important in wild populations [12], many of which are subject to regular changes in food availability [e.g. 32], and in which individuals typically differ in their foraging strategies [33].

Studies in model laboratory organisms show that phenotypic plasticity in cognitive performance can occur in parallel with gut microbiome alterations caused by diet [8,34]. For example mice fed beef-chow had a higher microbial diversity than those fed on normal chow (diets which differed in nutritional, amino acid and fatty acid content) and showed improved working and reference memory [34]. Observational studies also point to a relationship between diet, cognition and mood in humans [35,36], effects that may be mediated by the host gut microbiome [e.g. 36]. Although some empirical studies have been published on lab model systems, there is now a need for manipulative experiments using model ecological organisms to understand how dietary induced changes in the microbiome might affect functionally significant behaviours in the wild.

The great tit (*Parus major*) is a highly innovative species that has long been a model organism in field studies on behaviour, ecology and evolution. It is also ideal for short-term laboratory studies because they adapt well to temporary captivity [37,38; G. Davidson personal observation]. 'Problem solving performance' (PSP), in which individuals must solve a technical problem to obtain a reward, is often described as being underpinned by cognitive mechanisms [39]. In wild great tits, PSP differed consistently over time and across different tasks among individuals [40], and has been linked to a range of behaviours and fitness-related traits in the great tit, both in the wild and in the laboratory [37,38,41]. Quantitative genetic analysis in one population also showed that PSP was driven by positive correlations with favourable ecological conditions during the nestling stage, including habitat characteristics strongly linked to diet [41]. Innovativeness is also positively correlated with dietary breadth across bird species [42] because innovations such as PSP may increase their access to a wide range of resources [43]. If problem solvers have a wider dietary breadth, then we predicted that they would have a more diverse gut microbiome than non-problem solvers [12], although the direction of causality can also be in the opposite direction since the microbiome can also affect behaviour directly [1]. Thus given the links between diet, the microbiome and behaviour found in laboratory studies [34,44], the possible links with i) diet diversity and PSP, and ii) habitat quality and PSP, leads to the expectation of a positive correlation between PSP and alpha diversity in the great tit.

Here we tested whether habitat and diet predicted gut microbiota metrics in wild-caught great tits, and whether natural and manipulated gut microbiome profiles correlated with PSP. First we sampled birds from both rural and urban areas, known to differ in their quality as habitat for great tits [e.g. 45], and therefore we expected host gut microbiota may differ in diversity, community structure (i.e. beta diversity) and/or taxa-level abundance [e.g. 46], though our replication at the site level was limited. In

line with our hypotheses above, we predicted a positive correlation between PSP and natural
microbiome alpha diversity, although we also tested for differences in microbial community structure
similarity/dissimilarity (i.e. beta diversity) and taxa-level relative abundance between solvers and non-
solvers. Second, we manipulated the diet of wild-caught great tits to test whether microbiota
community structures became more dissimilar, and whether phylum-level and genus-level taxa
changed within and between treatment groups. Given the seed diet was higher in fat, protein and fibre
content than the insect only diet, we expected these individuals to have higher alpha diversity, higher
Firmicutes and lower Bacteroidetes if similar dietary effects occur in great tits as have been reported
in humans [e.g 26]. Third, we tested whether a change in the gut microbiome as a consequence of the
manipulated diet influenced PSP by presenting these birds with the same task again. We predicted that
if the gut microbiome influenced host behaviour, rather than the dietary treatment itself doing so, then
a change in behaviour should specifically be associated with the same metrics of the gut microbiome
as those that were changed as a result of dietary manipulation.

RESULTS

Sequencing summary

We collected fresh faecal samples within one hour of arrival in captivity, and at the end of the
experimental treatments, and we quantified the gut microbial community profiles using 16S RNA
sequencing. We obtained 4,109,746 sequences distributed along 2306 OTUs with an average 67,373 $\pm$
4,873 se sequences per sample (min. 16,126 and max. 180,09)7. A total of 314 OTUs from 192
genera, 116 family and 19 phyla were present in the data, following filtering steps.

Relative abundance, diet and microbial diversity

The most prominent phyla across all samples were as follows (mean percentage relative abundance
$\pm$SE): Proteobacteria (55.3%$\pm$4.0); Cyanobacteria (14.8%$\pm$2.6); Firmicutes (10.2%$\pm$2.1); Tenericutes
(9.0%$\pm$2.3); Actinobacteria (4.0%$\pm$1) and Bacteroidetes (2.0%$\pm$0.5). Mean relative abundance of most
abundance phyla across treatment groups is depicted in Figure 1 and Table 1, and all phyla in
supplementary (Fig S1). Proteobacteria increased significantly over the course of captivity following
the insect diet (z=2.02, p=0.04), but not in the seed diet (z=-0.28, p=0.78), and tended to be higher in
adults than juveniles (z=1.84, p=0.07). Bacteroidetes (natural log (Ln)-transformed) increased
following the insect diet (t=0.22, p=0.03), but not the seed diet (t=1.01, p=0.32), and tended to be
higher in urban compared to rural habitats (t=1.80, p=0.08). Actinobacteria was significantly higher in
birds from urban habitats (t= 2.47, p=0.02). There was no significant effect of diet, habitat, sex or age
on Firmicutes (Ln-transformed) or Tenericutes (Ln-transformed) (Table 2). There were no significant
interactions between diet and habitat for all models. Figure 1 depicts relative abundance across dietary
treatments. Full model outputs are provided in Table S2, supplementary.
127

Significant differences in genus-level abundance attributed to dietary treatments were found for 22
genera. Birds given the insect diet showed a decrease in *Devosia, Rickettsiella, Sphingomonas,*
Pantoea, Arthrobacter, Brevibacterium, Brachybacterium, Clostridium and *Carnobacterium,* and an
increase in *Candidatus, Methylobacterium.* Birds given the seed diet showed a decrease in
Cronobacter and *Serratia*, and an increase in *Microbacterium.* Birds in both dietary groups showed a
decrease in *Bradyrhizobium, Staphylococcus* and *Rahnella*, and an increase in *Lactobacillus, Bacillus,*
Ureaplasma, Delftia, Flavobacterium, Streptococcus and *Rhodococcus* (Table S3, supplementary).
135
There was a significant decrease in alpha diversity, specifically Chao1 (square root transformed)
following the insect diet (t=-2.51, p=0.02), but not the seed diet (t=-0.07, p=0.94) (Figure 2a). A
similar pattern was found in Shannon's index, although the effect of the insect diet was marginally
non-significant (insect diet: t=-2.02, p=0.06; seed diet: t=0.01, p=0.99) (Table 2, Figure 2b). Number
of observed OTUs (Ln-transformed) did not differ significantly across treatments (insect diet t=-1.74,
p=0.10; seed diet t=0.43, p=0.67) (Figure 2c). There was no significant effect of sex, age, or habitat
type on alpha diversity. There was no significant interaction between diet and habitat. (Table S4,
supplementary). ADONIS tests showed dietary treatment significantly influenced beta diversity across
all three metrics (Table 2, Figure 3a and Table S5). Post-hoc tests with Bonferroni correction indicate
that the differences were between day 1 and the post-diet insect group (R2=0.07, p<0.01), and
between the two post-diet groups (R2=0.10, p=0.03). Beta diversity also significantly differed
between habitat types (Table 2, Table S5).

Problem solving, diet and microbial diversity

The innovation task was solved 17 times, by 15 different individuals across both trial days, with two
individuals solving both before and after the dietary treatment. Twelve birds solved on day 1 (4 lever,
8 door), 5 birds in the seed group solved (2 lever, 3 door) and none of the birds from the insect group
solved. Beta diversity prior to dietary manipulation tended to differ between problem solvers and non-
problem solvers (unweighted unifrac distances R = 0.05, p=0.07). All other metrics of natural
variation in the gut microbiome were not significantly associated with PSP (Table S4). However, PSP
across both trials, when controlling for dietary treatment and repeated measures, was positively
correlated with alpha diversity (Shannon: z=2.22, p=0.03; Chao1 z=2.13, p=0.04, observed OTUs
z=1.96; p=0.06) (Table S2, Figure 4a,b,c), and beta diversity (unweighted unifrac: R2=0.02, p=0.04)
(Table 2, Table S5, Figure 3b). Phylum-level and genus-level relative abundance was not associated
with PSP.

Birds assigned to the seed diet solved more than the birds assigned to the insect diet (z=2.22, p=0.03),
and birds tended to be more likely to solve on day 1 than day 12 (z=1.93, p=0.054), though this effect
was likely driven by birds assigned to the insect diet (Figure 5). There was a tendency for juveniles to
solve more than adults (z=1.94, p=0.053). Neither the interactions between habitat, nor between
experiment day and diet were significant (Table S2, supplementary). PSP was not attributed to

differences in motivation to consume the food reward as all birds were equally likely to consume the
same reward when it was made freely available to them, regardless of dietary treatment and
experiment day (diet $z=1.18$, $p=0.24$, experiment day $z=0.85$, $p=0.40$). To rule out effects of
glucocorticoids on cognitive performance [47], we found that performance was not attributed to
individual baseline or stress-induced blood-circulating corticosterone levels measured as part of a
separate experiment (in prep, preliminary analysis available on request).

DISCUSSION

We demonstrate that an experimentally induced dietary change caused significant alterations to the
gut microbiome diversity, and to phylum- and genus-level abundance in a wild bird species, which in
turn was correlated with reduced innovative behaviour. To our knowledge, this is the first
experimental evidence from a non-model laboratory organism that gross microbiota composition is
associated with a behavioural trait, in this case PSP. We discuss the extent to which our results may
point to a causal link between microbiota and behaviour, and their broader relevance to foraging
ecology, the microbiome-gut-brain axis, and environment-behaviour interactions.

Seasonal and geographic differences in gut microbial communities in wild mammals have been
attributed to changes in food availability [19-21]. In a population of wild birds temporarily taken into
captivity, we show that changes in microbial community composition are sensitive to dietary changes,
independent of other factors that may differ with seasonality and impact on gut microbiota, such as
hormonal differences [48], because the changes here were recorded under controlled conditions over a
two week period. We show that phylum-level and genus-level diversity changes to the gut
microbiome were caused by the diet, which may be attributed to differences in relative protein, fat and
fibre content [25,26]. At the phylum-level, we found higher abundance of Bacteroidetes present in
subjects with lower fat, protein and fibre intake, a finding in line with Clarke et al [26]; in contrast to
that study, however, we did not find a decrease in Firmicutes, and found an increase in Proteobacteria.
These differences in dietary effects on the gut microbiota may be due to differences in the
microbiome between birds and mammals [49], and the type of fat and/or fibre [reviewed in 50]. At the
genus level, the insect diet caused a reduction in *Clostridum spp.* – taxa that have previously been
reported to be associated with host health in Eastern bluebird chicks (*Sialia sialis*) supplemented with
mealworms [51], and in poultry fed diets that differed in animal protein and in non-digestible non-
starch polysaccharides [23].

Birds in the insect diet showed a decrease in Chao1 index, a measure of diversity which estimates taxa
abundance and rare taxa missed from undersampling [52]. This result may be because the insect diet
itself was less diverse than the seed diet. While birds in both the seed and insect diets showed both
decreases and increases in genus-level abundance, only birds given the insect diet showed significant
changes in diversity and phylum-level abundance. This could perhaps be explained because our birds
had already been taking seeds at the feeders we used to lure them for capture, and because great tits

consume a high proportion of plant-based foods in the winter [53]. In urban and rural house sparrows, microbiota changes as a result of dietary manipulation were greatest when fed on diets that contrasted from their place of origin [24]. In the current study, the use of garden feeders in both urban and rural environments may also explain why there were no rural versus urban habitat differences in alpha diversity. There was, however, a significant difference in beta diversity as well as a higher proportion of Actinobacteria in urban birds compared to rural birds, similar metrics to those that have previously been shown to be related to urban environments in house sparrows (*Passer domesticus*) [46]. While our experiment in captivity aimed to mimic variation in individual food consumption, or in seasonal food availability, in the wild longitudinal sampling of individuals across seasons would be necessary to confirm whether similar microbial taxa changes would occur under natural conditions, particularly given that the invertebrate species accessible in the wild would differ from those provided in our experiment. Whatever the reason for limited effects of habitat on the microbiome, our manipulation showed clearly that diet does modify the microbiome in this model species in field ecology, and that more generally environmental drivers of variation in diet may play a significant role in driving these effects.

Dietary manipulation in this experiment affected not just the gut microbiome but was also correlated to the likelihood of solving. Those on the insect diet had reduced microbiome diversity and were less likely to solve, suggesting a potential causal link whereby the dietary induced reduction in the microbiome may have impeded PSP via the microbiome-gut-brain axis. Our results showed that the indices of microbial community diversity that decreased as a consequence of diet (i.e. Chao1 index, beta diversity) were the same metrics that were associated with variation in PSP. Proteobacteria and Bacteroidetes increased following an insect diet, but these two phyla were not associated with PSP, nor were the genus-level microbial taxa that were altered as a consequence of diet, suggesting that the microbial community structure as a whole may be important for regulating behaviour. Moreover, our results suggest that less abundant taxa may contribute this relationship as two of the metrics that were significantly associated with PSP are those that are most sensitive to differences in low-abundance features (i.e. Chao1 index and unweighted unifrac), although the significant relationship with Shannon's diversity also suggests OTU evenness is an important predictor.

Motivation can influence PSP [reviewed in 54] and an insect diet may have decreased motivation to engage in the problem solving task baited with an insect reward. However, the insect diet did not influence the birds' motivation to consume the freely available wax worm, irrespective of dietary treatment; and birds solved on day one when other insects (i.e. mealworms) were freely available, indicating that wax worms are a highly-valued and preferred food reward. Furthermore the links between PSP and the microbiota were significant, even after controlling for dietary manipulation as a fixed effect (see Table S2). Nevertheless, we cannot discount the possibility that birds were motivated to meet specific caloric or nutrient requirements due to differences in diets, and further manipulative experiments, such as microbial transplantation experiments [55] in the absence of dietary differences

would provide further support that the microbiome itself caused the change in problem solving
performance.

How the gut microbiome impacts behaviour via the gut-brain axis may be attributed to the metabolic
functions of the microbial community itself [e.g. 11], or derived from the diets of the host [e.g.
reviewed in 56]. Studies have attempted to disentangle nutritional and microbial effects on behaviour
by depleting the original microbes of the hosts and re-introducing specific bacterial organisms, or
transplanting gut microbiomes between hosts [e.g. 44,57]. And while we did not manipulate
microbiome directly owing to logistical and ethical constraints, the aim of our study was to test diet-
microbiome-behaviour relationships within an ecologically relevant context that would translate to
wild animals in their natural environment. To control for nutritional deficiencies that may have an
impact on behaviour independent of microbiome, we provided vitamin supplements, though other
nutritional differences cannot be discounted. Fat content and fibre content was five-fold and three-fold
higher in the seed diet compared to the insect diet, respectively. Mice fed on high fat diets, or given
microbiome transplantations from obese donors show poorer cognitive performance than control mice
[8,44]; whereas, our study showed that birds fed the lower-fat diet (i.e. insect) had poorer PSP.
Having a higher proportion of fibre present in the seed diet may have offset any negative effects of a
high-fat diet. Non-digestible carbohydrates are fermented by gut microbes in the large intestines,
promote the growth of microbial organisms and can have positive effects on cognition and behaviour
in mammals [reviewed in 58]. Metabolomics profiling would be an informative future endeavour to
provide a functional assessment of microbial products such as short chain fatty acids involved in gut-
brain axis communication [reviewed in 11,58,59] to pinpoint further the mechanisms underlying
phenotypic plasticity in innovative behaviour.

Contrary to our predictions, we did not find that individual variation in PSP was positively associated
with natural variation in microbial alpha diversity, as measured on the first day in captivity and before
dietary manipulation. Instead, beta diversity was dissimilar between solvers and non-solvers, although
this result was marginally non-significant and correlational. One reason for the lack of an effect is that
all of the birds were feeding on the same food at the feeders, and individual variation in foraging on
other food sources may simply not have been pronounced until we manipulated the diet in captivity.
Further manipulative investigations are needed to pinpoint causal directions of these relationships
under natural conditions, in particular whether innovative behaviour leads to variation in microbial
diversity through food access, or indeed whether innovation arises because of microbial diversity
caused by some other mechanism, such as developmental effects related to habitat quality, as
previously shown in this species [41]. Together our findings lend support to the hypothesis that the
gut microbiome, innovation and diet are interlinked, and more generally that the microbiome may be
an important source of variation in ecologically important behaviours, especially those linked to diet
and foraging.

MATERIALS AND METHODS

Subjects

Thirty six great tits were captured between January and March 2017 across four sites. Two sites were
within Cork city (urban), 1.6 km apart (51°54'32.4"N 8°28'12.0"W and 51°53'56.1"N 8°29'15.4"W),
and two were in deciduous woodlands (rural) 23 km apart (51˚43'08.4"N 8˚35'52.8"W and
51˚51'37.5"N 8˚47'01.1"W), and located least 23km from the urban sites. Birds were lured using
feeders containing sunflower seeds and peanuts. All birds were banded with rings issued by the
British Trust of Ornithology for individual identification. Birds were sexed (male/female) and aged
(adult/juvenile) from plumage characteristics [60]. Upon capture, birds were transported to the aviary
facilities at University College Cork and singly-housed in wire cages ($45 \times 50 \times 60$ cm) containing
two wooden perches.

Faecal sampling

Faecal samples were collected within 1 hour of arrival into the aviary, and again on Day 13 of
captivity. A clean sheet of brown paper was placed on the floor of each cage for faecal collection.
Paper was used in order to soak liquid urea away from the faecal matter as urea can act as a
downstream inhibitor to amplification [61]. Using sterile inoculation loops, we transferred the faecal
matter into tubes containing 1ml of 100% ethanol and stored tubes at -20 degrees Celsius.

Dietary manipulation

From day 2-13 of captivity, birds were given one of two different dietary treatments designed to
reflect ecological variation seen in the wild, for example changes in the availability of seed or animal
food sources [53,62], or perhaps reflecting potential individual differences in dietary specialisations
[63]: 1) Seed, n = 17; and 2) insect diet, n = 19. The insect diet consisted of wax moth larvae (*Achroia*
grisella) and mealworm larvae (*Tenebrio molitor*). Notably, these two diets differed in the relative
content of protein, fat and fibres (Table S1), features of diets that have previously been associated
with differences in host microbiota in mammals [50]. Mealworms were provided ad libitum, and five
wax worms were provided each morning and each evening (except during the problem solving task).
The seed diet consisted of sunflower hearts, peanuts and suet. We provided birds in the seed diet five
mealworms and one wax worm on day seven of captivity for welfare reasons as we did not want to
deprive the birds of a preferred food item as preference for mealworms/wax worms was observed in
our population (pers. obs), and on the basis of other work [e.g. 64]. To limit more general nutritional
deficiencies, all birds received vitamin powder mixed with their food and drops mixed in their water
(AviMix®). Birds were assigned to treatment groups randomly, counterbalanced for age and sex.

Problem solving assay

To quantify individual foraging innovation, naïve birds were presented with a novel problem solving
foraging task, derived from a similar foraging task [37,38,41]. The birds were given the task overnight

from one hour before sunset to two hours after sunrise, once on Day 1 of captivity and once on Day

12 of captivity. Due to the length of the trial, birds were not food deprived for welfare reasons. During

the first trial, all birds had access to mealworms, peanuts and sunflower hearts ad libitum. During the

second trial, birds had access to their assigned diets ad libitum. During both trials, wax worms, a

highly preferred food reward [40,65; G. Davidson personal observation], were placed inside a

transparent Perspex tube 16cm (height) x 5cm (width). The worms could be accessed by solving at

least one of three solutions: 1) by pulling a lever to drop a platform holding a worm; 2) by pushing a

door to the side; and 3) by pulling a string attached to one of the worms from the top of the tube. By

having multiple access possibilities, PSP could be assessed without limiting solutions to one particular

motor action that may be more feasible in some individuals over others. While we report the numbers

of each solving method in the PSP results, there was not sufficient power to analyse these specific

differences statistically. Moreover, our aviary setup did not facilitate video recording, therefore PSP

was recorded as a binary variable (i.e. solved versus not solved).At the start of the problem solving

assay, a freely available wax worm was placed outside, at the base of the problem solving device to

measure birds' motivation to approach the apparatus and consume the wax worm. Wax worms were

otherwise not provided when the problem solving task was presented. One bird died of unknown

causes following 10 days of captivity and therefore only data from Day 1 were included for this bird.

Microbiome analysis

Microbial DNA was extracted using the Qiagen QIAamp DNA Stool Kit, following the "Isolation of

DNA from Stool for Pathogen Detection" protocol with modifications described in Zeale *et al*. [66] to

increase DNA yield and remove excess inhibitors expected to be present in the uric acid of bird faeces

[but see 67 where they found no evidence of uric acid in feacal matter from a subset of avian species].

A 0.10 - 0.20 g aliquot of each faecal sample was added to the kit, alongside a negative control.

The V3-V4 variable region of the 16S rRNA gene was amplified from the DNA extracts using the 16S

metagenomic sequencing library protocol (Illumina) as described in Fouhy *et al* [68]. In the current

study, each PCR reaction contained 23 µl DNA template, 1 µl forward primer (10 µM), 1 µl reverse

primer (10 µM) and 25 µl 2X Kapa HiFi Hotstart ready mix (Roche, Ireland), to a final volume of 50

µl. Two negative controls were run in parallel – one from the DNA extraction and one containing

PCR water instead of DNA template. Of the bird samples, ten failed to amplify and were not pooled

for sequencing (Day 1: n = 1 (seed), Day 13: n = 3 (seed), 6 (insect)). Successful PCR products were

cleaned using AMPure XP magnetic bead based purification (Labplan, Dublin, Ireland). Samples were

sequenced at the Teagasc Sequencing Centre on the MiSeq sequencing platform, using a 2 x 300 cycle

kit, following standard Illumina sequencing protocols.

Three hundred base pair paired-end reads were assembled using FLASH (FLASH: fast length

adjustment of short reads to improve genome assemblies). Further processing of paired-end reads

including quality filtering based on a quality score of > 25 and removal of mismatched barcodes and

sequences below length thresholds was completed using QIIME (Quantitative Insights Into Microbial

Ecology) version 1.9.0 [52]. Denoising, chimera detection and clustering into operational taxonomic

units (OTUs) (97% identity, following Konstantinidis & Tiedje [69]) were performed using

USEARCH version 7 (64-bit). OTUs were aligned using PyNAST (PyNAST: python nearest

alignment space termination; a flexible tool for aligning sequences to a template alignment) and

taxonomy was assigned using BLAST against the SILVA SSURef database release version 123.

Alpha diversities (observed OTUs, Chao1 and Shannon's index) were generated from all OTUs in

QIIME.

Statistical analyses

The QIIME files (OTU table, taxonomic table, phylogenetic tree file) and metadata files were

analysed using phyloseq (McMurdie & Holmes 2013) in R Statistical Software (R Development Core

Team 2011). Singletons and taxa present at <0.005% were removed [70]. We applied cumulative sum

scaling normalisation to standardise library size across samples for beta diversity analysis. We

included all taxa in the dataset, most notably those that are often viewed as 'dietary contaminants' (i.e.

microbes originating from ingested food, such as cyanobacteria present in plants). This is because not

all cyanobacteria are dietary contaminants (Di Rienzi *et al*. 2013) and because our study specifically

tested differences between diets. Thereby removal of one so-called 'dietary contaminant' specific to

plants (i.e. cyanobacteria), but none specific to insects (e.g. insect microbiome) could systematically

bias our results. We also note that the non-sterilised water was an additional source of dietary

contaminants, although this would not systematically differ between dietary treatment groups.

Linear Mixed Models (LMMs) and Generalized Linear Mixed Models (GLMMs) were run using lme4

[71], and where relevant, p-values were obtained using lmerTest [72] in R version 3.5.2 [73]. Terms

with $p < 0.1$ were sequentially removed from the models, starting with interaction terms. Site ID and

Bird ID were included as random effects. Response variables were transformed where necessary to

meet assumptions of normality (Natural log (Ln) or square root transformed), and therefore all models

were run with a Gaussian distribution unless stated otherwise.

Dietary effects on microbiome

The effect of diet on the relative abundance (i.e. percentage abundance relative to all other phyla) was

tested for Firmicutes, Proteobacteria, Bacteroidetes, Tenericutes and Actinobacteria as these were the

most abundant phyla. Proteobacteria was modelled as a proportion and run with a binomial

distribution, as it could not be transformed to fit a normal distribution. Three different measures of

alpha diversity were used as response variables to test the effect of diet on the microbiome

community: Shannon's index, total observed OTUs, and Chao1. The above gut microbiome metrics

did not differ between pre-assigned dietary groups, therefore we used diet as a three level factor

accounting for both experimental date as a fixed effect: Diet (pre-dietary assignment (day 1), insect

post-diet (day 13) and seed post-diet (day 13)). Habitat (rural, urban), age (juvenile, adult [60]) and

sex were included as fixed effects. To avoid overparameterization of models, we tested the interaction

between diet and habitat only, as we expected inherent differences in diets between birds from the two

habitats. We also investigated the effect of diet on genus-level relative abundance with mixed

regression models using the web-based software Calypso (version 8.72) [74]. ID was included as a

random term and significant taxa were adjusted by a false discovery rate (FDR) whereby p-values of

less than 0.05 were considered statistically significant.

Dietary effects on problem solving performance

PSP was modelled as a binomial distribution (solved vs not solved) with diet (seed, insect),

experiment day (day 1, day 12), habitat (rural, urban), age (juvenile, adult) and sex included as fixed

effects. We included interactions between diet and habitat type and diet and experiment day. To test

whether diet affected motivation, we ran a binomial GLMM with consumption of the freely available

wax worm (Yes/No) as the response variable, assuming that birds who took the waxworm were more

motivated to solve than those who were not. The fixed and random effects were included as described

for the problem solving analysis above.

Relationship between problem solving and microbiome

We tested whether natural variation in the gut microbiome correlated with PSP on Day 1. We tested

for associations between microbial community and PSP as a binomial response variable (solved vs not

solved) in GLMMs for each microbial community measurement (the top five phyla, and the three

measures of alpha diversity), and included habitat, sex and age as fixed effects and site as a random

effect. Beta diversity was calculated in four ways: Jaccard distance matrix, and phylogenetic weighted

(accounting for relative abundance of taxa) and unweighted (presence/absence of taxa) unifrac

distance matrices. Each matrix was analysed using permutational multivariate analysis of variance

(ADONIS) with 1000 permutations. We tested for homogeneity of dispersion, and found there was no

difference in dispersion between groups. We also investigated the relationship between problem

solving and relative abundance at the genus-level as described above. To test whether gut microbiome

alteration as a consequence of dietary manipulation caused a change in PSP, we ran the same analyses

described above, but included data from Day 12 and diet as a fixed effect with bird ID as a random

term.

Research and Animal Ethics: This study was conducted under licences from the Health Products

Regulatory Authority (AE19130_P017), The National Parks and Wildlife Services (C11/2017) and

the British Trust for Ornithology, and permission from Coillte Forestry and private landowners. The

research project received ethical approval from the Animal Welfare Body at University College Cork,

and was in accordance with the ASAB (Association for the Study of Animal Behaviour) Guidelines

for the Treatment of Animals in Behavioural Research and Teaching.

Acknowledgements: James Nichols, Ally Phillimore and Sarah Knowles for helpful discussions on
molecular methodology; Shane Somers for advice on statistics; Jennifer Coomes for assistance with
fieldwork; Dr. Paul Cotter, Dr. Fiona Crispie and Ms. Laura Finnegan from the Teagasc Sequencing
facility for their role in relation to the 16S rRNA sequencing. Funding for GLD, ACC, MSR, IGK,
IDH and JMSC from the European Research Council under the European Union's Horizon 2020
Programme (FP7/2007-2013)/ERC Consolidator Grant "Evoecocog" Project No. 617509, awarded to
JLQ. This work was funded in part by Science Foundation Ireland through APC Microbiome Ireland.

Data accessibility. Our data will be deposited at Dryad, and sequence data uploaded to a repository
such as NCBI Sequence Read Archive, if the manuscript is published.

Authors contributions: GLD, ACC and JLQ designed the experiment with input from CS and RPR.
GLD and ACC ran the experiment with assistance from MSR, IGK, IDH and JMSC. GLD, NW and
CNJ carried out DNA extraction and library prep. FF and GLD analysed data. GLD wrote the
manuscript with input from all authors. All authors gave final approval for publication.

Additional Information: The authors have no competing interests.

431   Reference List

1. Cryan J.F. and Dinan T.G. (2012). Mind-altering microorganisms: the impact of the gut
microbiota on brain and behaviour. Nature Reviews Neuroscience *13*: 701.

2. Sherwin E., Bordenstein S.R., Quinn J.L., Dinan T.G., and Cryan J.F. (2019). Microbiota and
the social brain. Science *366*: eaar2016.

3. Diaz Heijtz R., Wang S., Anuar F., Qian Y., Bjorkholm B., Samuelsson A., Hibberd M.L.,
Forssberg H., and Pettersson S. (2011). Normal gut microbiota modulates brain development
and behavior. Proc Natl Acad Sci USA *108*: 3047-3052.

4. Foster J.A. and McVey Neufeld K.A. (2013). Gut-brain axis: how the microbiome influences
anxiety and depression. Trends in Neurosciences *36*: 305-312.

5. Clarke G., Grenham S., Scully P., Fitzgerald P., Moloney R.D., Shanahan F., Dinan T.G., and
Cryan J.F. (2012). The microbiome-gut-brain axis during early life regulates the hippocampal
serotonergic system in a sex-dependent manner. Molecular Psychiatry *18*: 666.

6. Desbonnet L., Clarke G., Shanahan F., Dinan T.G., and Cryan J.F. (2014). Microbiota is
essential for social development in the mouse. Molecular Psychiatry *19*: 146-148.

7. Hoban A.E., Stilling R.M., Moloney G., Shanahan F., Dinan T.G., Clarke G., and Cryan J.F.
(2017). The microbiome regulates amygdala-dependent fear recall. Molecular Psychiatry.

8.   Magnusson K.R., Hauck L., Jeffrey B.M., Elias V., Humphrey A., Nath R., Perrone A., and
Bermudez L.E. (2015). Relationships between diet-related changes in the gut microbiome
and cognitive flexibility. Neuroscience *300*: 128-140.

9.   Ogbonnaya E.S., Clarke G., Shanahan F., Dinan T.G., Cryan J.F., and O'Leary O.F. (2015). Adult
Hippocampal Neurogenesis Is Regulated by the Microbiome. Biological Psychiatry *78*: e7-e9.

10. Gareau M.G., Wine E., Rodrigues D.M., Cho J.H., Whary M.T., Philpott D.J., MacQueen G.,
and Sherman P.M. (2011). Bacterial infection causes stress-induced memory dysfunction in
mice. Gut *60*: 307.

11. Stilling R.M., van de Wouw M., Clarke G., Stanton C., Dinan T.G., and Cryan J.F. (2016). The
neuropharmacology of butyrate: The bread and butter of the microbiota-gut-brain axis?
Neurochemistry International *99*: 110-132.

12. Davidson G.L., Cooke A.C., Johnson C.N., and Quinn J.L. (2018). The gut microbiome as a
driver of individual variation in cognition and functional behaviour. Philos Trans R Soc Lond B
Biol Sci *373*: 20170286.

13. Morand-Ferron J., Cole E.F., and Quinn J.L. (2016). Studying the evolutionary ecology of
cognition in the wild: a review of practical and conceptual challenges. Biol Rev *91*: 367-389.

14. Stephens D.W. and Krebs J.K. (1986). Foraging Theory. (Princeton, New Jersey: Princeton
University Press).

15. De Filippo C., Cavalieri D., Di Paola M., Ramazzotti M., Poullet J.B., Massart S., Collini S.,
Pieraccini G., and Lionetti P. (2010). Impact of diet in shaping gut microbiota revealed by a
comparative study in children from Europe and rural Africa. Proc Natl Acad Sci USA *107*:
14691.

16. Gillingham M.A.F., Bechet A., Cezilly F., Wilhelm K., Rendon-Martos M., Borghesi F., Nissardi
S., Baccetti N., Azafzaf H., Menke S. et al. (2019). Offspring Microbiomes Differ Across
Breeding Sites in a Panmictic Species. Front Microbiol *10*: 35.

17. Costa S., Lopes I., Proenca D.N., Ribeiro R., and Morais P.V. (2016). Diversity of cutaneous
microbiome of Pelophylax perezi populations inhabiting different environments. Science of
The Total Environment *572*: 995-1004.

18. Knutie S.A., Chaves J.A., and Gotanda K.M. (2019). Human activity can influence the gut
microbiota of Darwin's finches in the Galapagos Islands. Mol Ecol *28*: 2441-2450.

19. Amato K.R., Leigh S.R., Kent A., Mackie R.I., Yeoman C.J., Stumpf R.M., Wilson B.A., Nelson
K.E., White B.A., and Garber P.A. (2015). The Gut Microbiota Appears to Compensate for
Seasonal Diet Variation in the Wild Black Howler Monkey (Alouatta pigra). Microbial Ecology
69: 434-443.

20. Hicks A.L., Lee K.J., Couto-Rodriguez M., Patel J., Sinha R., Guo C., Olson S.H., Seimon A.,
Seimon T.A., Ondzie A.U. et al. (2018). Gut microbiomes of wild great apes fluctuate
seasonally in response to diet. Nature Communications *9*: 1786.

21. Maurice C.F., CL Knowles S., Ladau J., Pollard K.S., Fenton A., Pedersen A.B., and Turnbaugh
P.J. (2015). Marked seasonal variation in the wild mouse gut microbiota. The Isme Journal *9*:
2423.

22.　Lozupone C.A., Stombaugh J.I., Gordon J.I., Jansson J.K., and Knight R. (2012). Diversity,
stability and resilience of the human gut microbiota. Nature *489*: 220.

23.　Pan D. and Yu Z. (2014). Intestinal microbiome of poultry and its interaction with host and
diet. Gut Microbes *5*: 108-119.

24.　Teyssier A., Matthysen E., Hudin N.S., de Neve L., White J.I., and Lens L. (2020). Diet
contributes to urban-induced alterations in gut microbiota: experimental evidence from a
wild passerine. Proc Biol Sci *287*: 20192182.

25.　David L.A., Maurice C.F., Carmody R.N., Gootenberg D.B., Button J.E., Wolfe B.E., Ling A.V.,
Devlin A.S., Varma Y., Fischbach M.A. et al. (2014). Diet rapidly and reproducibly alters the
human gut microbiome. Nature *505*.

26.　Clarke S.F., Murphy E.F., Sullivan O., Lucey A.J., Humphreys M., Hogan A., Hayes P., Reilly M.,
Jeffery I.B., Wood-Martin R. et al. (2014). Exercise and associated dietary extremes impact
on gut microbial diversity. Gut *63*: 1913.

27.　Fava F., Gitau R., Griffin B.A., Gibson G.R., Tuohy K.M., and Lovegrove J.A. (2013). The type
and quantity of dietary fat and carbohydrate alter faecal microbiome and short-chain fatty
acid excretion in a metabolic syndrome at-risk population. International Journal of Obesity
37: 216-223.

28.　Wu G.D., Compher C., Chen E.Z., Smith S.A., Shah R.D., Bittinger K., Chehoud C., Albenberg
L.G., Nessel L., Gilroy E. et al. (2016). Comparative metabolomics in vegans and omnivores
reveal constraints on diet-dependent gut microbiota metabolite production. Gut *65*: 63.

29.　Zimmer J., Lange B., Frick J.S., Sauer H., Zimmermann K., Schwiertz A., Rusch K.,
Klosterhalfen S., and Enck P. (2012). A vegan or vegetarian diet substantially alters the
human colonic faecal microbiota. European Journal of Clinical Nutrition *66*: 53-60.

30.　Youngblut N.D., Reischer G.H., Walters W., Schuster N., Walzer C., Stalder G., Ley R.E., and
Farnleitner A.H. (2019). Host diet and evolutionary history explain different aspects of gut
microbiome diversity among vertebrate clades. Nature Communications *10*: 2200.

31.　Hird S.M., Sanchez C., Carstens B.C., and Brumfield R.T. (2015). Comparative Gut Microbiota
of 59 Neotropical Bird Species. Front Microbiol *6*: 1403.

32.　Kartzinel T.R., Hsing J.C., Musili P.M., Brown B.R.P., and Pringle R.M. (2019). Covariation of
diet and gut microbiome in African megafauna. Proc Natl Acad Sci USA 201905666.

33.　Bolnick D., Svanback R., Fordyce J., Yang L., Davis J., Hulsey C., and Forister M. (2003). The
Ecology of Individuals: Incidence and Implications of Individual Specialization. The American
Naturalist *161*: 1-28.

34.　Li W., Dowd S.E., Scurlock B., Acosta-Martinez V., and Lyte M. (2009). Memory and learning
behavior in mice is temporally associated with diet-induced alterations in gut bacteria.
Physiology & Behavior *96*: 557-567.

35.　Psaltopoulou T., Sergentanis T.N., Panagiotakos D.B., Sergentanis I.N., Kosti R., and Scarmeas
N. (2013). Mediterranean diet, stroke, cognitive impairment, and depression: A meta-
analysis. Ann Neurol. *74*: 580-591.

36.   Carlson A.L., Xia K., Azcarate-Peril M.A., Goldman B.D., Ahn M., Styner M.A., Thompson A.L.,
529   Geng X., Gilmore J.H., and Knickmeyer R.C. (2018). Infant Gut Microbiome Associated With
530   Cognitive-Development. Biological Psychiatry *83*: 148-159.

531   37.   Dunn J.C., Cole E.F., and Quinn J.L. (2011). Personality and parasites: sex-dependent
532   associations between avian malaria infection and multiple behavioural traits. Behavioral
533   Ecology and Sociobiology *65*: 1459-1471.

534   38.   Cole E., Morand-Ferron J., Hinks A., and Quinn J. (2012). Cognitive Ability Influences
535   Reproductive Life History Variation in the Wild. Current Biology *22*: 1808-1812.

536   39.   Seed A. and Mayer C. (2017). Problem solving. In APA handbook of comparative psychology:
537   Perception, learning, and cognition, Vol. 2, (Washington, DC, US: American Psychological
538   Association), pp. 601-625.

539   40.   Cole E.F., Cram D.L., and Quinn J.L. (2011). Individual variation in spontaneous problem-
540   solving performance among wild great tits. Animal Behaviour *81*: 491-498.

541   41.   Quinn J.L., Cole E.F., Reed T.E., and Morand-Ferron J. (2016). Environmental and genetic
542   determinants of innovativeness in a natural population of birds. Philos Trans R Soc Lond B
543   Biol Sci *371*: 20150184.

544   42.   Ducatez S., Clavel J., and Lefebvre L. (2015). Ecological generalism and behavioural
545   innovation in birds: technical intelligence or the simple incorporation of new foods? J Anim
546   Ecol *84*: 79-89.

547   43.   Reader S.M. and MacDonald K. (2003). Environmental variability and primate behavioural
548   flexibility. In Animal innovation, (New York, NY, US: Oxford University Press), pp. 83-116.

549   44.   Bruce-Keller A.J., Salbaum J.M., Luo M., Blanchard E., IV, Taylor C.M., Welsh D.A., and
550   Berthoud H.R. (2015). Obese-type Gut Microbiota Induce Neurobehavioral Changes in the
551   Absence of Obesity. Biological Psychiatry *77*: 607-615.

552   45.   Biard C., Brischoux F., Meill+¿re A.e., Michaud B., Nivi+¿re M., Ruault S.p., Vaugoyeau M.,
553   and Angelier F.d.r. (2017). Growing in Cities: An Urban Penalty for Wild Birds? A Study of
554   Phenotypic Differences between Urban and Rural Great Tit Chicks (Parus major). Frontiers in
555   Ecology and Evolution *5*: 79.

556   46.   Teyssier A., Rouffaer L.O., Saleh Hudin N., Strubbe D., Matthysen E., Lens L., and White J.I.
557   (2018). Inside the guts of the city: Urban-induced alterations of the gut microbiota in a wild
558   passerine. Science of The Total Environment *612*: 1276-1286.

559   47.   Lupien S.J., McEwen B.S., Gunnar M.R., and Heim C. (2009). Effects of stress throughout the
560   lifespan on the brain, behaviour and cognition. Nature Reviews Neuroscience *10*: 434.

561   48.   Escallon C., Belden L.K., and Moore I.T. (2019). The Cloacal Microbiome Changes with the
562   Breeding Season in a Wild Bird. Integrative Organismal Biology *1*.

563   49.   Waite D.W. and Taylor M.W. (2014). Characterizing the avian gut microbiota: membership,
564   driving influences, and potential function. Front Microbiol *5*: 223.

50. Singh R.K., Chang H.W., Yan D., Lee K.M., Ucmak D., Wong K., Abrouk M., Farahnik B.,
Nakamura M., Zhu T.H. et al. (2017). Influence of diet on the gut microbiome and
implications for human health. Journal of Translational Medicine *15*: 73.

51. Knutie S.A. (2020). Food supplementation affects gut microbiota and immunological
resistance to parasites in a wild bird species. J Appl Ecol *57*: 536-547.

52. Caporaso J.G., Kuczynski J., Stombaugh J., Bittinger K., Bushman F.D., Costello E.K., Fierer N.,
Pe+¦a A.G., Goodrich J.K., Gordon J.I. et al. (2010). QIIME allows analysis of high-throughput
community sequencing data. Nat Methods *7*.

53. Vel'ky M., Kanuch P., and Kristin A. (2011). Food composition of wintering great tits (Parus
major): habitat and seasonal aspects. 2011 *60*: 228-236.

54. Griffin A.S. and Guez D. (2014). Innovation and problem solving: A review of common
mechanisms. Behavioural Processes *109*: 121-134.

55. Greyson-Gaito C.J., Bartley T.J., Cottenie K., Jarvis W.M.C., Newman A.E.M., and Stothart
M.R. (2020). Into the wild: microbiome transplant studies need broader ecological reality.
Proc Biol Sci *287*: 20192834.

56. Roager H.M. and Dragsted L.O. (2019). Diet-derived microbial metabolites in health and
disease. Nutr Bull *44*: 216-227.

57. Mohle L., Mattei D., Heimesaat M., Bereswill S., Fischer A., Alutis M., French T.,
Hambardzumyan D., Matzinger P., Dunay I. et al. (2016). Ly6C-hi- Monocytes Provide a Link
between Antibiotic-Induced Changes in Gut Microbiota and Adult Hippocampal
Neurogenesis. Cell Reports *15*: 1945-1956.

58. Cryan J.F., O'Riordan K.J., Cowan C.S.M., Sandhu K.V., Bastiaanssen T.F.S., Boehme M.,
Codagnone M.G., Cussotto S., Fulling C., Golubeva A.V. et al. (2019). The Microbiota-Gut-
Brain Axis. Physiological Reviews *99*: 1877-2013.

59. Heintz-Buschart A. and Wilmes P. (2018). Human Gut Microbiome: Function Matters. Trends
in Microbiology *26*: 563-574.

60. Svensson L. (1992). Identification guide to European passerines. (Thetford, United Kingdom:
British Trust for Ornithology).

61. Khan G., Kangro H.O., Coates P.J., and Heath R.B. (1991). Inhibitory effects of urine on the
polymerase chain reaction for cytomegalovirus DNA. J Clin Pathol *44*: 360-365.

62. Perrins C. (1991). Tits and their caterpillar food supply. Ibis *133*: 49-54.

63. Serrano-Davies E., O'Shea W., and Quinn J.L. (2017). Individual foraging preferences are
linked to innovativeness and personality in the great tit. Behavioral Ecology and Sociobiology
71: 161.

64. Aplin L.M., Sheldon B.C., and McElreath R. (2017). Conformity does not perpetuate
suboptimal traditions in a wild population of songbirds. Proc Natl Acad Sci USA *114*: 7830.

65.   O'Shea W., Serrano-Davies E., Quinn J.L., and Pruitt J. (2017). Do personality and
602   innovativeness influence competitive ability? An experimental test in the great tit.
603   Behavioral Ecology *28*: 1435-1444.

604   66.   Zeale M.R.K., Butlin R.K., Barker G.L.A., Lees D.C., and JONES G.A.R.E. (2011). Taxon-specific
605   PCR for DNA barcoding arthropod prey in bat faeces. Mol Ecol Resour *11*: 236-244.

606   67.   Crouch N.M.A., Lynch V.M., and Clarke J.A. (2019). A re-evaluation of the chemical
607   composition of avian urinary excreta. Journal of Ornithology.

608   68.   Fouhy F., Watkins C., Hill C.J., O'Shea C.A., Nagle B., Dempsey E.M., O'Toole P.W., Ross R.P.,
609   Ryan C.A., and Stanton C. (2019). Perinatal factors affect the gut microbiota up to four years
610   after birth. Nature Communications *10*: 1517.

611   69.   Konstantinidis K.T. and Tiedje J.M. (2005). Genomic insights that advance the species
612   definition for prokaryotes. Proc Natl Acad Sci U S A *2005/02/08*: 2567-2572.

613   70.   Bokulich N.A., Subramanian S., Faith J.J., Gevers D., Gordon J.I., Knight R., Mills D.A., and
614   Caporaso J.G. (2013). Quality-filtering vastly improves diversity estimates from Illumina
615   amplicon sequencing. Nat Methods *2012/12/02*: 57-59.

616   71.   Bates D., Maechler M., Bolker B., and Walker S. (2014). lme4: Linear mixed-effects models
617   using Eigen and S4.).

618   72.   Kuznetsova A., Brockhoff P.B., and Christensen R.H.B. (2017). lmerTest Package: Tests in
619   Linear Mixed Effects Models. Journal of Statistical Software; Vol 1, Issue 13 (2017).

620   73.   R Development Core Team. (2011). R: a language and environment for statistical computing.
621   (Vienna, Austria: R Foundation for Statistical Computing).

622   74.   Zakrzewski M., Proietti C., Ellis J.J., Hasan S., Brion M.J., Berger B., and Krause L. (2017).
623   Calypso: a user-friendly web-server for mining and visualizing microbiome-environment
624   interactions. Bioinformatics *33*: 782-783.

625
626   Table 1. Means and se of most abundance phyla across treatment groups

Phylum	Day 1	Insect diet	Seed diet
Proteobacteria	51.0%±5.3	76.0±7.9	53.8%±8.3
Cyanobacteria	21.0%±4.0	0.3%±0.1	14.5%±4.6
Firmicutes	9.4%±2.0	11%±7.1	10.4%±4.5
Tenericutes	7.8%±3.3	7.0±3.1	15.8%±5.8
Actinobacteria	3.5%±1.3	2.1%±1.0	1.3%±0.3
Bacteroidetes	0.5%±0.1	2.3%±0.8	1.6%±1.1
Others	6.6%±0.2	1.4%±0.1	2.6%±1.1

628   Table 2. Summary of the relationships between microbiota and host traits. Significant relationships
629   are shown as either positive (+), negative (-) or as p-values (i.e. beta-diversity where differences do
630   not represent directionality). NS = Non-significant.

Microbiota metrics	Diet	PSP	Habitat	Sex	Age

Protebacteria	(+) insect	NS	NS	NS	NS
Firmicutes	NS	NS	NS	NS	NS
Bacteroidetes	(+) insect	NS	NS	NS	NS
Actinobacteria	NS	NS	(+) urban	NS	NS
Tenericutes	NS	NS	NS	NS	NS
Chao1	(-) insect	(+)	NS	NS	NS
Shannon's index	NS	(+)	NS	NS	NS
Observed OTUs	NS	(-)	NS	NS	NS
Jaccard	p <0.01	NS	p<0.01	NS	NS
Weighted unifrac	p <0.01	NS	p=0.01	NS	p<0.05
unweighted unifrac	p <0.01	p<0.05	NS	NS	NS

Figure 1. Relative abundance of top seven phyla across dietary treatments. Proteobacteria and Bacteroidetes significantly increased following the insect diet, but not the seed diet.

Figure 2. Alpha diversity for birds pre-dietary assignment (Day 1), and following the insect diet (ID)
and the seed diet (SD) for a) Chao1 (sqrt), b) Shannon's index, c) Observed OTUs (Ln). Coloured
points denote individual data points, black points and line denote mean and ± SE. p-values represent
the comparison between post-insect diet group and day 1.

Figure 3. Nonmetric Multidimensional Scaling (NMDS) ordination plots based on (a) weighted
unifrac distances of diet and experiment day, and (b) unweighted unifrac distances of problem solving
performance. Ellipses represent standard deviations around the centroids of the groups. Numbers in
brackets refer to the variance explained by NDMS axes.

Figure 4. PSP-alpha diversity relationships in a) Chao1 (sqrt), b) Shannon's index, c) Observed OTUs
(Ln) including data points from both day 1 and day 12. Coloured points denote individual data points,
black points and line denote mean and ± SE.

Figure 5. Problem solving performance (PSP) as a measure of innovation. Number of individuals that solved (dark grey) and number of birds that did not solve (light grey) pre- and post- dietary treatments.

Legends:

Table 1. Means and se of most abundance phyla across treatment groups

Table 2. Summary of the relationships between microbiota and host traits. Significant relationships are shown as either positive (+), negative (-) or as p-values (i.e. beta-diversity where differences do not represent directionality). NS = Non-significant.

Figure 1. Relative abundance of top seven phyla across dietary treatments. Proteobacteria and Bacteroidetes significantly increased following the insect diet, but not the seed diet.

Figure 2. Alpha diversity for birds pre-dietary assignment (Day 1), and following the insect diet (ID) and the seed diet (SD) for a) Chao1 (sqrt), b) Shannon's index, c) Observed OTUs (Ln). Coloured points denote individual data points, black points and line denote mean and ± SE. p-values represent the comparison between post-insect diet group and day 1.

Figure 3. Nonmetric Multidimensional Scaling (NMDS) ordination plots based on (a) weighted unifrac distances of diet and experiment day, and (b) unweighted unifrac distances of problem solving performance. Ellipses represent standard deviations around the centroids of the groups. Numbers in brackets refer to the variance explained by NDMS axes.

Figure 4. PSP-alpha diversity relationships in a) Chao1 (sqrt), b) Shannon's index, c) Observed OTUs (Ln) including data points from both day 1 and day 12. Coloured points denote individual data points, black points and line denote mean and ± SE.

672   Figure 5. Problem solving performance (PSP) as a measure of innovation. Number of individuals that
673   solved (dark grey) and number of birds that did not solve (light grey) pre- and post- dietary
674   treatments.