Phasic and tonic serotonin modulate alarm reactions and post-exposure behavior in zebrafish

Monica Lima-Maximino[1], Maryana Pereira Pyterson[2], Rhayra Xavier do Carmo Silva[2,3], Gabriela Cristini Vidal Gomes[2], Sueslene Prado Rocha[1], Anderson Manoel Herculano[4], Denis Broock Rosemberg[5], Caio Maximino[2*]

1 – Laboratório de Neurofarmacologia e Biofísica, Centro de Ciências Biológicas e da Saúde, Departamento de Morfologia e Ciências Fisiológicas, Universidade do Estado do Pará, Campus VIII, Marabá/PA, Brazil

2 – Laboratório de Neurociências e Comportamento "Frederico Guilherme Graeff", Faculdade de Psicologia, Universidade Federal do Sul e Sudeste do Pará, Marabá/PA, Brazil

3 – Programa de Pós-Graduação em Neurociências e Biologia Celular, Instituto de Ciências Biológicas, Universidade Federal do Pará, Belém/PA, Brazil

4 – Laboratório de Neurofarmacologia Experimental, Instituto de Ciências Biológicas, Universidade Federal do Pará, Belém/PA, Brazil

5 – Laboratório de Neuropsicobiologia Experimental, Departamento de Bioquímica e Biologia Molecular, Centro de Ciências Naturais e Exatas, Universidade Federal de Santa Maria, Santa Maria/RS, Brazil

* Corresponding author

Caio Maximino

Laboratório de Neurociências e Comportamento "Frederico Guilherme Graeff", Faculdade de Psicologia, Universidade Federal do Sul e Sudeste do Pará, Unidade III

Av. dos Ipês, S/N, s/ CEP, Bairro Cidade Jardim, Marabá/PA, Brazil

cmaximino@unifesspa.edu.br

List of abbreviations

27	4-OH quinoline	4-hydroxyquinoline
28	5-HT	Serotonin
29	5,7-DHT	5,7-dihydroxytryptamine
30	ANOVA	Analysis of variance
31	CAS	Conspecific alarm substance
32	CONCEA	Conselho Nacional de Controle de Experimentação Animal
33	CTRL	Control groups
34	IACUC	Institutional Animal Care and Use Committee
35	Ibama	Instituto Brasileiro do Meio Ambiente e dos Recursos Naturais Renováveis
36	IV	Independent variable
37	MAD	Median absolute difference
38	pCPA	*para*-chlorophenylalanine
39	PI	Principal Investigator
40	ppm	Parts per million
41	UEPA	Universidade do Estado do Pará
42	WAY 100,635	*N*-[2-[4-(2-Methoxyphenyl)-1-piperazinyl]ethyl]-*N*-(2-
43		pyridyl)cyclohexanecarboxamide
44	zMAO	Zebrafish monoamine oxidase

Abstract

Current theories on the role of serotonin (5-HT) in vertebrate defensive behavior suggest that this monoamine increases anxiety but decreases fear, by acting at different levels of the neuroaxis. This paradoxical, dual role of 5-HT suggests that a serotonergic tone inhibits fear responses, while an acute increase in 5-HT would produce anxiety-like behavior. However, so far no evidence for a serotonergic tone has been found. Using zebrafish alarm responses, we investigate the participation of phasic and tonic 5-HT levels in fear-like behavior, as well as in behavior after stimulation. Conspecific alarm substance (CAS) increased bottom-dwelling and erratic swimming, and animals transferred to a novel environment after CAS exposure (post-exposure behavior) showed increased bottom-dwelling and freezing. Clonazepam blocked CAS effects during and after exposure. Acute fluoxetine dose-dependently decreased fear-like behavior, but increased post-exposure freezing. Metergoline had no effect on fear-like behavior, but blocked the effects of CAS on post-exposure behavior; similar effects were observed with pCPA. Finally, CAS was shown to decrease the activity of monoamine oxidase in the zebrafish brain after exposure. These results suggest that phasic and tonic serotonin encode an aversive expectation value, switching behavior towards cautious exploration/risk assessment/anxiety when the aversive stimulus is no longer present.

Keywords: Serotonin; Fear; Alarm substance; Zebrafish; Panic

1. Introduction

The neurocircuitry of defensive reactions involves regulation by a plethora of neuromodulators, including monoamines and peptides (Maximino 2012). In vertebrates, the monoamine serotonin (5-HT) is produced in specific brain nuclei, including the raphe, and is thought to inhibit fear/escape responses to proximate threat by acting on more caudal structures of the aversive brain system (Paul *et al.* 2014; Deakin and Graeff 1991; Maximino 2012). This response appears to be dependent on the specific brain region in which serotonin acts, as well as on the receptor that is activated. For example, in the rodent periaqueductal gray, the activation of 5-HT_{1A} and 5-HT_2-type receptors inhibit fear responses, while in amygdaloid nuclei the activation of 5-HT_2- and 5-HT_3-type receptors increase anxiety-like responses (Guimarães *et al.* 2008; Paul *et al.* 2014; Hale and Lowry 2011). There is also evidence for a serotonergic "tone" inhibiting anxiety, since antagonists usually inhibit anxiety-like responses in animal models; however, antagonists do not appear to modify fear-like responses, suggesting that phasic, not tonic, serotonin is involved in fear. For example, 5-HT levels do not change in the basolateral amygdala or in the dorsal periaqueductal gray during chemical stimulation of this latter structure in rats (Zanoveli *et al.* 2009), a manipulation that induces panic-like responses (Brandão *et al.* 2008). This suggests that the inhibitory role of serotonin in fear functions as a "switching" signal: as the threatening stimulus ceases, serotonin is released, inhibiting the fear reactions that are now non-adaptive, and initiating careful exploration and risk assessment responses to ensure that the threat is actually over.

In non-mammalian vertebrates, including teleost fish, 5-HT is produced in additional brain regions, including pretectal and hypothalamic populations (Herculano and Maximino 2014). There is some evidence that 5-HTergic neurons innervate areas of the teleostean brain which participate in defensive behavior, including prosencephalic and mesencephalic regions (do Carmo Silva *et al.* 2018a). A role for 5-HT in modulating fish defensive behavior has been demonstrated before: in zebrafish, 5-HT_{1A} and 5-HT_{1B} receptor antagonists decrease anxiety-like behavior (Maximino *et al.* 2013; Nowicki *et al.* 2014; Herculano *et al.* 2015; Maximino *et al.* 2015), while 5-HT_2- and 5-HT_3-

type antagonists increase it (Nowicki *et al.* 2014). Little is known, however, of the modulation of

fear-like responses. Microinjection of the serotonergic neurotoxin 5,7-dihydroxytryptamine (5,7-

DHT) in the telencephalon of zebrafish, destroying most serotonergic innervation in regions

associated with aversive learning, impairs the acquisition of active avoidance (Amo *et al.* 2014),

suggesting that serotonin encodes an aversive expectation value.

In zebrafish and other Actinopterygian fish, specialized club cells in the skin produce a

substance (conspecific alarm substance, CAS) that, when the skin is damaged, is dispersed in the

water, signaling to conspecifics a potential threat (von Frisch 1941; Hüttel 1941; von Frisch 1938;

Døving and Lastein 2009; Maximino *et al.* 2019). CAS induces defensive behavior in zebrafish,

including increased bottom-dwelling, erratic swimming, and freezing (Maximino *et al.* 2019; Egan

et al. 2009; Speedie and Gerlai 2008; Maximino *et al.* 2014). These responses have been exploited

as a model system to study fear in more basal vertebrates (Maximino *et al.* 2019; Jesuthasan and

Mathuru 2008).

The serotonergic system has also been implicated in some of these behavioral functions.

CAS increases extracellular serotonin levels (Maximino *et al.* 2014) and inhibits monoamine

oxidase activity (Quadros *et al.* 2018) in the zebrafish brain after exposure. Zebrafish exposed to

CAS show increased anxiety-like behavior in the light/dark test after exposure (i.e., when the

substance is no longer present), an effect that is blocked by fluoxetine but not by the 5-HT$_{1A}$

receptor antagonist WAY 100,635 (Maximino *et al.* 2014). Interestingly, WAY 100,635 blocked the

analgesic effects of CAS in zebrafish (Maximino *et al.* 2014), suggesting that this receptor

participates in some, but not all, neurobehavioral responses to threatening stimuli. While WAY

100,635 was not able to alter anxiety-like behavior *after* exposure, the drug blocked the increased

geotaxis *during* CAS exposure, both in the first minutes of exposure and in the last minutes (Nathan

et al. 2015). Blocking 5-HT$_2$-type receptors with methysergide did not affect these responses,

except at a sedative dose (Nathan *et al.* 2015). These results are difficult to interpret, but suggest

that CAS increases serotonergic activity after exposure, and that a serotonergic tone on the 5-HT_{1A}

receptor is involved in behavioral switching after exposure – that is, when the threat is no longer

present, and risk assessment begins; whether this is true for behavioral responses *during* exposure –

that is, when the threat is present – is unknown.

The present paper investigated whether phasic and tonic serotonin participates in the alarm

response in zebrafish during and after exposure. Our results reinforce the idea that behavior during

CAS exposure is qualitatively different from behavior after CAS exposure. We also show that

behavior in both contexts are differentially sensitive to clonazepam, a high potency benzodizepine

commonly used in the clinical management of panic disorder (Caldirola *et al.* 2016). Increasing

serotonin levels by treating zebrafish with acute fluoxetine blocked the effects of CAS during and

after exposure, but blocking serotonin receptors with metergoline, or blocking serotonin synthesis

with *p*CPA, produced an effect only after exposure. Finally, we show that CAS inhibited

monoamine oxidase activity in the brain. Results are discussed in terms of the putative role of

serotonin in an homeostatic "neurobehavioral switch" in the absence of threat after predatory risk.

2. Methods

2.1. Animals, housing, and baseline conditions

435 zebrafish (*Danio rerio*) from the longfin phenotype were used in the present

experiments; details for sample size calculations can be found on each experimental section, below.

Outbred populations were used due to their increased genetic variability, decreasing the effects of

random genetic drift that could lead to the development of uniquely heritable traits (Parra *et al.*

2009; Speedie and Gerlai 2008). The populations used in the present experiments are expected to

better represent the natural populations in the wild. Animals were bought from a commercial vendor

(Fernando Peixes, Belém/PA) and arrived in the laboratory with an approximate age of 3 months

(standard length = 13.2 ± 1.4 mm), and were quarantined for two weeks; the experiment began

when animals had an approximate age of 4 months (standard length = 23.0 ± 3.2 mm). Animals

were kept in mixed-sex tanks during acclimation, with an approximate ratio of 50-50 males to

females (confirmed by body morphology). The breeder was licensed for aquaculture under Ibama's

(Instituto Brasileiro do Meio Ambiente e dos Recursos Naturais Renováveis) Resolution 95/1993.

Animals were group-housed in 40 L tanks, with a maximum density of 25 fish per tank, for at least

2 weeks before experiments begun. Tanks were filled with non-chlorinated water at room

temperature (28 °C) and a pH of 7.0-8.0. Lighting was provided by fluorescent lamps in a cycle of

14-10 hours (LD), according to standards of care for zebrafish (Lawrence, 2007). Water quality

parameters were as follows: pH 7.0-8.0; hardness 100-150 mg/L $CaCO_3$; dissolved oxygen 7.5-8.0

mg/L; ammonia and nitrite < 0.001 ppm. Potential suffering of animals was minimized by

controlling for the aforementioned environmental variables and scoring humane endpoints (clinical

signs, behavioral changes, bacteriological status), following Brazilian legislation (Conselho

Nacional de Controle de Experimentação Animal - CONCEA 2017). Animals were used for only

one experiment and in a single behavioral test, to reduce interference from apparatus exposure.

Experiments were approved by UEPA's IACUC under protocol 06/18.

2.2. Alarm substance extraction

CAS was extracted at a ratio of 1 donor fish for 10 ml distilled water. A detailed protocol for

extraction can be found at protocols.io (Silva, Rocha, Lima-Maximino, & Maximino, 2018;

https://dx.doi.org/10.17504/protocols.io.tr3em8n). Briefly, a donor fish was cold-anesthetized and

euthanized, and 15 shallow cuts were made on the side of its trunk to lesion club cells. The cuts

were washed with 10 mL distilled water, and 7 mL of the eluate was reserved as 1 unit CAS.

2.3. General experimental design

After the onset of drug effects (see details below for each drug), animals were individually transferred to a 1.5 L tank (12 cm X 12 cm x C cm, w X l X h), filled with system water, and left to acclimate for 3 min. Filming was started, and animals were exposed to either 7 mL distilled water (CTRL groups) or 7 mL (1 unit) alarm substance (CAS groups). Exposure was made by slowly pouring the substance on the water from the top. Animals were then left undisturbed as filming continued for 6 min; this was termed "alarm reaction". The animal was then transferred to a tank with 500 mL mineral water for a 1 min "washout" period, to remove potential residues from the alarm substance. After this period, the animal was transferred to a 5 L tank (A cm X 24 cm X 22 cm, w X l X h) and freely explored for 6 min, during which its behavior was recorded; this was termed "post-exposure behavior" (Figure 1A). Tanks for both stages were differently shaped to increase the novelty of the second environment, a variable that is important to induce an anxiety-like "diving" response in animals not exposed to CAS (Bencan *et al.* 2009). In all experiments, the following variables were recorded:

- Time spent on the bottom third of the tank (s) [Primary outcome]
- Time spent on the top third of the tank (s) [Secondary outcome]
- Absolute turn angle (equivalent to erratic swimming) [Secondary outcome]
- Freezing: duration of complete movement cessation, defined as speed lower than 0.5 cm/s. [Secondary outcome]
- Swimming speed (cm/s) [Secondary outcome]

Variables were extracted by automated video tracking, using the software TheRealFishTracker v. 0.4.0 (http://www.dgp.toronto.edu/~mccrae/projects/FishTracker/), running on a Windows platform. Animals were randomly allocated to groups using a random number generator (http://www.jerrydallal.com/random/random_block_size_r.htm), with each subject randomized to a single treatment using random permuted blocks. One PI attributed a random letter

to treatment (e.g., "A" for CTRL, "B" for CAS) and a random integer for drug dose (e.g., "1" for 1

mg/kg, "2" for 0 mg/kg [vehicle]), and combinations for letters and integers were randomized. For

each experiment, animals were treated and tested in the order of allocation (i.e., randomly). In all

experiments, experimenters and data analysts were blinded to drugs and treatment by using coded

vials (with the same code used for randomization); blinding was removed only after data analysis.

Experiments were always run between 08:00AM and 02:00 PM. After experiments, animals were

sacrificed by prolonged bath in ice-cold water (< 12 ℃), followed by spinal transection (Matthews

and Varga 2011).

2.4. Quality control

Exclusion criteria: With the exception of outlier exclusion (described in 2.5.3), no

exclusion criteria were predetermined.

Behavioral data: Quality control of samples was maintained by periodic assessment of

water quality and health parameters. All experimenters were trained in the behavioral methods

before experiments; training included observation of all experiments by a PI (CM or MGL) on at

least two occasions. After these observations, each trainee performed two mock experiments, on a

single subject each, while being observed by the PI. All protocols were reviewed by all PIs, and are

publicly available. Behavioral records were reviewed by at least one PI for administration/scoring

accuracy, in order to ensure adherence to protocols and consistency across tests.

Biochemical data: All experimenters were trained in the analytical method before

experiments. Quality control was achieved periodically using Levey-Jennings charts for known

concentrations of kynuramine, adopting a 1_{2S} rule.

2.5. Experiments 1-4: Effects of clonazepam and serotonergic drugs on alarm reaction and post-exposure behavior

2.5.1. Sample size calculations

Sample size calculations were based on a power analysis for a 2-way ANOVA with interaction effects, with $\alpha = 0.05$, $\beta = 0.8$, and expected effect size $f = 0.25$ for each independent variable (IV); Effect sizes used for estimating sample sizes were based on the range of effects observed after pharmacological manipulations on zebrafish anxiety-like behavior in a metanalysis (Kysil et al., 2017). Sample size of 15 animals/group was established. Thus, a total of 270 animals were used for experiments, and another 135 animals were used to produce CAS. The distribution of samples through groups can be found in Figure S1.

2.5.2. Drugs and treatments

Clonazepam (CAS #1622-61-3) was bought from Roche on 2018, and dissolved in Cortland's salt solution (NaCl 124.1 mM, KCl 5.1 mM, Na_2HPO_4 2.9 mM, $MgSO_4$ 1.9 mM, $CaCl_2$ 1.4 mM, $NaHCO_3$ 11.9 mM, Polyvinylpyrrolidone 4%, 1,000 USP units Heparin; Wolf 1963). Clonazepam, as a high-potency benzodiazepine widely used in treating panic disorder (Cloos 2005; Caldirola *et al.* 2017), is expected to decrease fear-like responses to CAS, and therefore used as a positive control. Fluoxetine is expected to acutely increase serotonin levels in the synapse (Figure 1B). Fluoxetine hydrochloride (CAS #54910-89-3) was bought from Libbs on 2017, and dissolved in Cortland's salt solution. Metergoline is expected to block 5-HT receptors from the $5-HT_1$, $5-HT_2$, and $5-HT_7$ families (Figure 1C). Metergoline (CAS #17692-51-2) was bought from Virbac on 2017, and dissolved in Cortland's salt solution. pCPA is expected block 5-HT synthesis, therefore greatly reducing serotonergic tone on all receptors (Figure 1D). 4-chloro-DL-phenylalanine (pCPA; CAS #7424-00-2) was bought from Sigma-Aldrich (C6506) on 2018, and dissolved in 10% DMSO. For Experiment 1, animals were injected intraperitoneally with either vehicle (Cortland's salt solution) or clonazepam (0.05 mg/kg; Maximino, Silva, Gouveia Jr., & Herculano, 2011). For Experiment 2,

animals were injected intraperitoneally with vehicle (Cortland's salt solution) or fluoxetine (2.5 or

25 /kg; Maximino et al. 2014). For Experiment 3, animals were injected intraperitoneally with

vehicle (Cortland's salt solution) or metergoline (1 mg/kg; Pimentel et al. 2019). For Experiment 4,

animals were injected intraperitoneally with either vehicle (DMSO) or pCPA (one injection of 150

mg/kg/day for 2 days, followed by 24 h without treatment; Curzon et al. 1978). Injections were

made according to the protocol proposed by Kinkel et al. (2010); briefly, animals were cold-

anesthetized and transferred to a sponge-based surgical bed, in which injection was made. Injections

were made using a microsyringe (Hamilton® 701N syringe, needle size 26 gauge at cone tip), with

total volumes of injection ranging from 4.81 to 5.05 µL. Cold-anesthesia has been shown to

produce satisfactory results in zebrafish, with faster recovery and less animal loss than commonly

used anesthetics such as MS-222 (Matthews and Varga 2011). The sponge allowed gill perfusion to

be kept, minimizing suffering. 20 min after recovery, animals were subjected to CAS or water.

2.5.3. Statistical analysis

Outliers were removed based on median absolute differences (MADs), using time on bottom

as main endpoint; values were removed when they were higher or lower than 3 MADs around the

median (Leys *et al.* 2013), and the number of outliers was reported in the results. Differences

between groups were analyzed using two-way analyses of variance (ANOVAs) with robust

estimators on Huber's M-estimators, using the R package 'rcompanion' (Mangiafico 2017;

https://cran.r-project.org/package=rcompanion). Normality was not assumed, and thus no specific

test for normality was performed; however, this type of analysis is resistant to deviations from the

assumptions of the traditional ordinary-least-squares ANOVA, and are robust to outliers, thus being

insensitive to distributional assumptions (such as normality)(Huber 1981). Behavioral variables

were included as outcomes, with treatment and drug used as independent variables; interaction

between IVs was assessed as the most important predictor. P-values were adjusted for the false
discovery rate.

2.6. Experiment 5: Effects of CAS on monoamine oxidase activity

2.6. Experiment 5: Effects of CAS on monoamine oxidase activity
2.6.1. Sample size
Based on a power analysis for two-sample unpaired t-test. $\alpha = 0.05$, power $= 0.8$, and
expected effect size $d = 1.5$, a sample size of 10 animals/group was established. Thus, a total of 20
animals were used for experiments, and 10 more used to produce CAS. The distribution of samples
through groups can be found in Figure S1.

2.6.2. Methods
z-MAO activity was determined as reported previously (Quadros *et al.* 2018). Two zebrafish
brains were pooled per sample and homogenized in 0.5 mL of buffer solution containing 16.8 mM
Na_2HPO_4 and 10.6 mM KH_2PO_4, pH 7.4, isotonized with sucrose. Samples (n = 10 per group) were
centrifuged at 1.000 x g for 5 min, and the supernatants were kept on ice for the experiments.
Protein samples (approximately 100 µg) were mixed with 460 µL of assay buffer (168 mM
Na_2HPO_4 and 10.6 mM KH_2PO_4, pH 7.4, isotonized with KCl) and preincubated at 37°C for 5 min.
The reaction started by adding 110 µM kynuramine hydrobromide in a final volume of 700 µL, and
was stopped 30 min later with 300 µL 10% trichloroacetic acid. Reaction products were further
centrifuged at 16.000 x g for 5 min and supernatants (800 µL) were mixed with 1M NaOH (1 mL).
Fluorescence was measured using excitation at 315 nm and emission at 380 nm. Product formation
(4-hydroxyquinoline) was estimated and enzyme activity was expressed as expressed as nmol 4-OH
quinoline/min/mg protein.

2.6.3. Statistical analysis

Data were analyzed with an asymptotic general independence test, using the R package 'coin' (Hothorn et al. 2006; https://cran.r-project.org/package=coin).

2.7. Open science practices

Experiments were formally preregistered at Open Science Framework (https://doi.org/10.17605/OSF.IO/QM3PX). Data packages and analysis scripts for all experiments can be found at a GitHub repository (https://github.com/lanec-unifesspa/5-HT-CAS). Preprints for the manuscript can be found at bioRxiv (https://doi.org/10.1101/827824).

2.7.1. Changes from pre-registration

During pre-registration, we proposed to use manual recording of behavioral variables. During experiments, we decided to use automated tracking, due to the higher availability of open source software; automated tracking allows for a better reproducibility and precision in measures but, as a trade-off, some measurements were not possible. In the present experiments, the software TheRealFishTracker (v. 0.4.0, for Windows; http://www.dgp.toronto.edu/~mccrae/projects/FishTracker/) was used. Moreover, during pre-registration we proposed to also analyze melanophore responses to CAS; this data will appear in a separate paper, on behavioral and physiological aspects of the alarm reaction, and the data is available at a GitHub repository (https://github.com/lanec-unifesspa/5-HT-CAS/tree/master/data/melanophore). Finally, the following doses were changed from pre-registration: clonazepam was reduced to 0.05 mg/kg, to avoid unwanted sedation; a second dose of fluoxetine was added to approach the range in which fluoxetine blocks fear conditioning at a high shock intensity (Santos *et al.* 2006). Due to a problem with solubility, the pCPA dose was reduced to 150 mg/kg, a dose that has been shown to reduce serotonin levels in the rat brain by about 50%, and to block the release of 5-HT elicited by electrical stimulation of the raphe (Curzon *et al.* 1978).

3. Results

3.1. Experiment 1

No outliers were removed in Experiment 1. During CAS exposure, significant effects of treatment ($p = 0.006$), dose ($p = 0.044$, and interaction ($p = 0.006$) were found for time on top (Figure 2A); however, post-hoc tests failed to find differences between groups (all $p > 0.6$). Significant effects of treatment ($p = 0.0044$), dose ($p = 0.0034$), and interaction ($p = 0.0044$) were found for time on bottom (Figure 2B); CAS increased time on bottom (adjusted $p = 0.05$), and CLZ blocked this effect. Main effects of treatment ($p = 0.0042$) and dose ($p = 0.0038$), as well as an interaction effect ($p = 0.0042$), were found for absolute turn angle (Figure 2C); CAS increased absolute turn angle (adjusted $p = 4 \cdot 10^{-5}$), and CLZ blocked this effect. Main effects of treatment ($p = 0.0028$) and dose ($p = 0.0036$), as well as an interaction effect ($p = 0.0028$), were found for freezing (Figure 2D); again, CAS increased freezing (adjusted $p = 0.0243$), and CLZ blocked this effect. Main effects were found for treatment ($p = 0.0024$) and dose ($p = 0.0048$); however, post-hoc tests failed to uncover differences between groups (Figure 2E).

After CAS exposure, significant main effects of treatment ($p = 0.0048$) and dose ($p = 0.0042$), as well as an interaction effect ($p = 0.0038$), were found for time on top (Figure 3A); CAS decreased time on top ($p = 0.0482$), and CLZ partially blocked this effect. While main effects of treatment ($p = 0.0054$) and dose ($p = 0.0046$), as well as an interaction effect ($p = 0.0044$), were found for time on bottom (Figure 3B), post-hoc tests failed to find significant differences between groups. No main effects ($p > 0.18$), no an interaction effect ($p = 0.366$) were found for absolute turn angle (Figure 3C). A main effect of treatment ($p = 0.0008$) and drug ($p = 7.43 \cdot 10^{-5}$), as well as an interaction effect ($p = 2.37 \cdot 10^{-6}$), were found for freezing (Figure 3D); CAS increased freezing (adjusted $p = 0.0005$), and CLZ blocked this effect. No main effects were found for swimming

333 speed ($p > 0.08$), but an interaction was found ($p = 0.0304$); however, post-hoc tests faild to find

334 significant differences between groups (Figure 3E).

3.2. Experiment 2

One outlier was removed from the group exposed to CAS and treated with 2.5 mg/kg

fluoxetine. During CAS exposure, significant effects of treatment ($p = 0.0458$), dose ($p < 0.0001$),

and interaction ($p < 0.0001$) were found for time on top. Fluoxetine alone increased time on top at

2.5 mg/kg (adjusted $p = 3.218 \cdot 10^{-5}$ vs. control), CAS decreased it (adjusted $p = 0.01132$), and

fluoxetine blocked the effect of CAS at both doses (Figure 4A). Main effects of treatment ($p =$

0.004) and dose ($p < 0.0001$), but no interaction ($p = 0.6152$), were found for time on bottom (Figure

4B); CAS increased time on bottom (adjusted $p < 0.0236$), fluoxetine decreased it at both doses

(adjusted $p < 0.01$), and fluoxetine blocked the effect of CAS at the highest dose. A main effect of

treatment ($p = 0.0436$), but not dose ($p = 0.1102$) nor interaction ($p = 0.1148$), was found for

absolute turn angle (Figure 4C); post-hoc tests suggested that CAS increased absolute turn angle

(adjusted $p = 0.0061$ vs. control), and fluoxetine partially (2.5 mg/kg) or fully (25 mg/kg) blocked

this effect. A main effect of treatment ($p < 0.0001$) and dose ($p = 0.0004$), as well as an interaction

effect ($p = 0.00002$), were found for freezing (Figure 4D); CAS increased freezing (adjusted $p =$

0.0058), and both doses partially blocked this effect. Finally, no effect was found on swimming

speed ($p > 0.39$; Figure 4E).

Significant effects of treatment ($p = 0.0072$) and interaction ($p = 0.0196$) were found for

time on top after exposure (Figure 5A). Post-hoc pairwise permutation tests found a difference

between control and CAS-exposed animals (adjusted $p = 1.194 \cdot 10^{-5}$), an effect that was not

blocked by fluoxetine. Similarly, main effects of treatment ($p = 0.0002$), but not a drug ($p = 0.2914$)

nor an interaction effect ($p = 0.5878$) were found for time on bottom (Figure 5B), with a significant

increase in CAS-exposed animals (all adjusted $p < 0.001$). No effects were found for erratic

swimming ($p > 0.7$; Figure 5C). Significant treatment ($p < 0.0001$), dose ($p = 0.025$), and interaction effects ($p = 0.0014$), were found for freezing (Figure 5D), with CAS increasing freezing at all drug treatments, and the highest fluoxetine dose potentiating this effect. No effects were found for swimming speed ($p > 0.24$; Figure 5E).

3.3 Experiment 3

One outlier was removed from the group exposed to water and treated with vehicle. During CAS exposure, significant effects of treatment ($p < 0.0001$), but not metergoline ($p = 0.6682$) or interaction ($p = 0.5162$), were found for time on top (Figure 6A); post-hoc comparisons suggested that CAS decreased time on top (adjusted $p = 1.977 \cdot 10^{-6}$), but metergoline did not block effect. Significant effects of treatment ($p = 0.0432$), but not metergoline ($p = 0.9518$) nor interaction ($p = 0.4174$) were found for time on bottom (Figure 6B), with CAS increasing time on bottom (adjusted $p = 0.01472$) and no effect of metergoline. Significant effects of treatment ($p = 0.0002$), but not metergoline ($p = 0.6496$) nor interaction ($p = 0.1814$), were found for absolute turn angle (Figure 6C), with CAS increasing absolute turn angle (adjusted $p = 0.02627$) and metergoline having no effect. Significant effects of treatment ($p < 0.0001$), but not metergoline ($p = 0.462$) nor interaction ($p = 0.1922$), were found for freezing (Figure 6D); post-hoc tests found significant differences between CAS-exposed animals and controls treated with vehicle (adjusted $p = 0.02073$), but metergoline did not block the effects of CAS on freezing. No effects of treatment ($p = $, metergoline, and interaction (all $p > 0.1$) were found for swimming speed (Figure 6E).

After CAS exposure, no main nor interaction effects were found for time on top (all $p > 0.13$; Figure 7A). A dose ($p = 0.0156$) and an interaction ($p = 0.047$) effects were found for time on bottom, but a treatment effect was not found ($p = 0.827$); CAS increased time on bottom (adjusted $p = 0.0292$), an effect that was decreased by metergoline (Figure 7B). No main nor interaction effects were found for absolute turn angle ($p > 0.33$; Figure 7C). A main effect of treatment ($p = 0.0036$),

383   but not drug ($p = 0.4182$), nor interaction ($p = 0.1508$), was found for freezing (Figure 7D); CAS

384   increased freezing (adjusted $p = 0.0075$), and metergoline partially blocked this effect. No main

385   effects were found for swimming speed ($p > 0.27$), but an interaction effect was found ($p = 0.026$);

386   however, post-hoc tests failed to find significant differences between groups.

3.4. Experiment 4

Three outliers were removed from the group exposed to water and treated with vehicle, two

from the group exposed to water and treated with pCPA, one from the group exposed to CAS and

treated with vehicle, and two from the group exposed to CAS and treated with pCPA. During CAS

exposure, a main effect of treatment ($p < 0.0001$), but no drug ($p = 0.801$) nor interaction effects (p

$= 0.5386$) were found for time on top (Figure 8A); CAS decreased time on top on both vehicle- and

pCPA-injected animals (both ajusted $p > 0.0025$). Likewise, a main effect of treatment ($p = 0.0022$),

but no drug ($p = 0.6496$) nor an interaction effects ($p = 0.1166$), were found for time on bottom

(Figure 8B); CAS increased time on bottom on both vehicle- and pCPA-injected animals (both

adjusted $p = 0.05$). A main effect of treatment ($p = 0.0002$), but not an effect of drug ($p = 0.8162$)

nor interaction ($p = 0.5612$), was found for absolute turn angle (Figure 8C); CAS increased absolute

turn angle on both vehicle- and pCPA-injected animals (both adjusted $p < 0.002$). A main effect of

treatment ($p < 0.0001$), but not an effect of drug ($p = 0.6822$) nor interaction ($p = 0.4156$), was

found for freezing (Figure 8D); CAS increased freezing on both vehicle- and pCPA-injected

animals (both adjusted $p < 0.018$). No main or interaction effects were found for speed (all $p > 0.11$;

Figure 8E).

--Figure 8---

After CAS exposure, no main (all $p > 0.08$) or interaction ($p = 0.1472$) effects were found

for time on top (Figure 9A). A main effect of drug ($p < 0.0001$), but not of treatment ($p = 0.4514$),

nor an interaction effect ($p = 0.5436$), was found for time on bottom (Figure 9B); pCPA decreased

time on bottom at both controls and CAS-exposed animals (all adjusted $p < 0.001$). No main (all p

> 0.3) nor interaction ($p = 0.6958$) effects were found for absolute turn angle (Figure 9C). A main

effect of treatment ($p = 0.0108$), but not drug ($p = 0.1892$) nor interaction ($p = 0.2576$), was found

for freezing (Figure 9D); CAS increased freezing (adjusted $p = 0.02278$), an effect that was blocked

by pCPA (ajusted $p = 0.01453$). No main (all $p > 0.19$) nor interaction ($p = 0.0954$) effects were

found for swimming speed (Figure 9E).

--Figure 9---

3.5. Experiment 5

420   No outliers were removed. After CAS exposure, zMAO activity was reduced in the brain (Z

421    $= 3.205$, $p = 0.00135$; Figure 10).

4. Discussion

The present work attempted to clarify the role of phasic and tonic serotonin in the alarm

reaction of zebrafish during (fear-like behavior) and after (recovery) exposure. We found that

clonazepam decreased fear-like behavior, as well as post-exposure behavior, suggesting a good

predictive validity of the assay. Moreover, acute fluoxetine dose-dependently decreased fear-like

behavior, but increased freezing post-exposure. Metergoline had no effect on fear-like behavior, but

blocked the effects of conspecific alarm substance (CAS) on post-exposure behavior; similar effects

were observed with pCPA. Finally, CAS was shown to decrease the activity of monoamine oxidase

in the zebrafish brain after exposure.

4.1. Behavior during and after CAS exposure

In zebrafish, reported behavioral effects of CAS vary widely as a function of timing,

extraction method, and whether animals are tested alone or in groups (Maximino *et al.* 2019). When

animals are exposed and/or tested alone, as in the present experiments, bottom-dwelling, freezing,

and erratic swimming consistently increases *during* exposure (Eachus *et al.* 2017; Nathan *et al.*

2015; Ogawa *et al.* 2014; Maximino *et al.* 2014), but effects *after* exposure are less clear (Quadros

et al. 2016; Schirmer *et al.* 2013; Nathan *et al.* 2015; Egan *et al.* 2009). In the present experiments,

CAS consistently increased bottom-dwelling and erratic swimming during exposure, while after

exposure bottom-dwelling and freezing were increased. The first effects were blocked by treatment

with the panicolytic drug clonazepam, which nonetheless had a very mild effect on post-exposure

behavior. Thus, two components can be elicited by CAS: the first, dominated by erratic swimming,

occurs when the substance is present, and the second, dominated by freezing, occurs when the

concentrations of CAS decrease.

Observing the context in which defensive behavior, and not only the response topography, is

important to understand the function of a specific response. The context in which CAS elicits alarm

reactions is akin to a circa-strike defensive situation (Maximino *et al.* 2019), therefore producing

freezing and escape reactions that are fear-like; when CAS signals decrease, however (i.e., *after*

exposure in the present experiments), the context is akin to a post-encounter defensive situation,

eliciting avoidance and freezing behavior (see Perusini and Fanselow 2015 for a discussion on

predatory imminence, defensive reactions, and fear vs. anxiety). In these contexts, increases in

freezing, for example, can be interpreted as representing two different functions: to escape detection

by predators in the first case, and to allow careful vigilance, in the second case.

The differences in behavior during and after exposure are reminiscent of the different types

of freezing elicited during and after electrical stimulation of the periaqueductal gray in rodents

(Brandão *et al.* 2008). The increased erratic swimming observed during CAS exposure suggest

escape and/or avoidance attempts, while the increased freezing observed after CAS exposure

suggest a role in risk assessment. The effects of clonazepam also suggest different neurobehavioral

states: this drug usually decreases panic attacks, but has small effects on generalized anxiety in

human clinical settings (Caldirola *et al.* 2016; Cloos 2005). These results imply good predictive

validity, suggesting that behavior during and after CAS can be used to study fear- vs. anxiety-like

effects.

4.2. Role of phasic serotonin on CAS effects

Fluoxetine dose-dependently blocked the alarm reaction (fear-like behavior during CAS

exposure) and post-exposure behavior in zebrafish. These results are similar to what was previously

observed in the light/dark test, in which fluoxetine (2.5 mg/kg) blocked post-exposure effects on

scototaxis, freezing, and erratic swimming (Maximino *et al.* 2014). While the role of phasic

increases in serotonergic signaling on acute fear-like responses in zebrafish has not been previously

investigated, a higher dose (10 mg/kg) blocked the alarm reaction (i.e., during exposure) in the

piauçu *Leporinus macrocephalus* (Barbosa *et al.* 2012). This phasic role of serotonin is likely

highly conserved, as serotonin has been shown to decrease responses to aversive odors in

Caenorhabditis elegans (Li *et al.* 2012; Harris *et al.* 2009). The effects of fluoxetine strongly

suggest that serotonin phasically inhibits fear-like behavior in zebrafish, acting as a switch towards

risk assessment.

Phasic serotonin has been proposed to modulate fear-like behavior in mammals before

(Zangrossi Jr *et al.* 2001; Paul *et al.* 2014; Guimarães *et al.* 2010); the Deakin/Graeff theory

suggests a "panic inhibition system" (Paul *et al.* 2014; Silva *et al.* 2019) that inhibits behavioral

and sympathoexcitatory responses to these stimuli, and is mediated by serotonergic signaling. The

theory proposes a "dual role" for serotonin, increasing anxiety-like responses and inhibiting fear-

like responses. A similar mechanism has been proposed for zebrafish based on data regarding

serotonergic drugs in anxiety-like behavior (Herculano and Maximino 2014). We propose that, at

least in zebrafish, phasic serotonin does not physiologically inhibits fear responses; instead, the

inhibitory role of serotonin in fear functions as a "neurobehavioral switch": as the threatening

stimulus ceases, serotonin is released, inhibiting the fear reactions that are now non-adaptive, and

initiating careful exploration and risk assessment responses to ensure that the threat is actually over

(Figure 11). This is consistent with expectancy value theories, in which serotonin signals represent

the expectation of risk/threatening outcomes (aversive expectation values), from which appropriate

behavioral strategies can be selected (Amo *et al.* 2014; Cools *et al.* 2011).

A role for the serotonin transporter has also been proposed for the selection of behavior at

different levels of a threat imminence continuum: animals with lower expression of the transporter

are more cautious and readily show defensive responses under distal threat, while animals with high

expression show more defensive responses under proximal threat (Kroes *et al.* 2019). While the

expression levels of the serotonin transporter are more associated with controlling serotonergic

tone, this protein has been also shown to mediate the increases in serotonin levels after CAS

exposure in zebrafish (Maximino *et al.* 2014), suggesting a participation also in phasic signals.

Whether serotonin transporter expression levels are associated with the alarm reaction and/or post-

exposure behavior in zebrafish is still unknown.

4.3. Is there a tonic inhibition of fear-like responses in zebrafish?

The hypothesis that serotonin functions as a "neurobehavioral switch" signal in zebrafish

aversive behavior would be strengthened if decreasing the effects of serotonin on its receptors

inhibited post-exposure behavior. Indeed, metergoline, which non-specifically blocks 5-HT_1, 5-HT_2,

and 5-HT$_7$ receptors, had no effect on the alarm reaction, but blocked the post-exposure effects of

CAS on bottom-dwelling and homebase use; no effect was observed during exposure, suggesting

that fear-like responses are not under tonic inhibition. pCPA had similar effects. Nathan et al. (2015)

observed that blocking 5-HT$_{1A}$ and 5-HT$_2$ receptors potentiates freezing and bottom-dwelling both

in the initial moments of exposure and in a "recovery period"; however, during the recovery period

animals were still exposed to CAS, the doses which produced effect in Nathan et al. (2015) were

higher than reported in other experiments with zebrafish (Maximino *et al.* 2013), and important

controls were lacking, making comparison of results difficult.

Further support for this hypothesis is lent by sophisticated experiments made by Amo et al.

(2014) using the serotonergic neurotoxin 5,7-DHT. Injection of this neurotoxin in the telencephalon

destroyed most serotonergic fibers projecting to it, and led to an inability to learn an active

avoidance contingency (Amo *et al.* 2014), suggesting that serotonergic signaling in the

telencephalon represents an aversive expectation value. Amo et al. (2014) demonstrated that this

circuitry is under the control of projections from the ventral habenula which are not necessary for

classical fear conditioning. This suggests that this habenula-raphe-telencephalon pathways do not

simply process a fear response, but instead represents expectations values that can be used to inhibit

fear when threat is no longer present.

A caveat of the results from metergoline and pCPA experiments is that these drugs also

affect other neurotransmitter systems. Metergoline also acts as a non-selective dopamine receptor

antagonist (https://www.ebi.ac.uk/chebi/searchId.do?chebiId=CHEBI:64216); although its affinity

for 5-HT$_2$ receptors is ~25 times higher than for the dopamine D$_2$ receptor, the affinity for 5-HT$_1$

receptors is comparable to D$_2$ receptors (Dukhovich *et al.* 2004). While pCPA has been reported to

produce a selective effect on serotonin levels in zebrafish (Sallinen *et al.* 2009), not altering levels

of catecholamines, at the moment we cannot discard the possibility that the treatment used in the

present article were not due to changes in these systems. We cannot discard, then, the participation

of catecholamines along with serotonin in the effects of these drugs on post-exposure behavior.

In addition to the effects of the manipulations of the serotonergic system on post-exposure

behavior, zMAO activity has been shown to be decreased after CAS exposure, which would

increase serotonin levels at this moment. Previously, CAS has been shown to increase 5-HT levels

in the extracellular fluid of the zebrafish brain 20 min after CAS stress (Maximino *et al.* 2014), and

repeated (7 day) exposure to CAS decreases the mRNA levels of the serotonergic genes *pet1* and

slc6a4a (serotonin transporter)(Ogawa *et al.* 2014). These results suggest that CAS increases

serotonergic activity after the stimulus is no longer present, but it is not known whether CAS does

so during exposure.

540 4.4. Which receptors are involved?

The present work did not investigate specific receptors which are involved in the alarm

reaction in zebrafish. However, a role for 5-HT_1-, 5-HT_2-, and 5-HT_7-like receptors is suggested by

the effects of metergoline. The 5-HT_{1A} receptor antagonist WAY 100,635 has been previously

shown to block fear-induced analgesia elicited by CAS, but not the increase in anxiety-like behavior

in the light/dark test (Maximino *et al.* 2014). At higher doses, however, WAY 100,635 potentiated

the effects of CAS both in the early responses (0-8 min) and in the late phase (8-13 min) (Nathan *et*

al. 2015). These contradictory results can be explained by differences in exposure methods, as well

as differences in behavioral scoring techniques. Methysergide, which non-selectively blocks 5-

HT_{2A}, 5-HT_{2B}, and 5-HT_{2C} receptors, also potentiate the effects of CAS at both times (Nathan *et al.*

2015). However, 5-HT_{1A}, 5-HT_{2A}, and 5-HT_{2C} receptors in the dorsolateral periaqueductal gray

area have been shown to phasically inhibit escape/fear responses in rats (Soares and Zangrossi Jr

2004). While currently it is unknown whether the griseum centrale, the teleostean homolog of the

periaqueductal gray area, is involved in fear responses in zebrafish or not, its anatomical position

and hodology suggests so (Maximino *et al.* 2019; do Carmo Silva *et al.* 2018a). Thus, 5-HT$_{1A}$ and

5-HT$_2$-like receptors appear to be involved in phasic inhibition of fear-like behavior, but so far

evidence for a tonic inhibition is lacking.

558 5. Conclusion

The present experiments evidenced two qualitatively different stages of the alarm reaction in

zebrafish, one in the presence of the alarm substance, and another when it is no longer present, both

sensitive to clonazepam. Results from biochemistry and pharmacological manipulations suggest

that phasic and tonic serotonin acts as a neurobehavioral switch towards cautious exploration/risk

assessment/anxiety when the aversive stimulus is no longer present. These results refine previous

theories on the role of serotonin in anxiety and fear, suggesting new avenues of research.

566 Acknowledgments

567   This work was supported by a Conselho Nacional de Desenvolvimento Científico e

568 Tecnológico/CNPq grant to C.M. (#400726/2016-5). M.P.P. was the recipient of a CNPq

569 undergraduate scholarship. D.B.R. is the recipient of a CNPq research productivity grant

570 (#307595/2015-3). The funders had no influence on the study design, the collection, analysis and

571 interpretation of data, as well as writing and submission of this manuscript.

573 References

Amo R., Fredes F., Kinoshita M., Aoki R., Aizawa H., Agetsuma M., Aoki T., et al. (2014) The
habenulo-raphe serotonergic circuit encodes an aversive expectation value essential for
adaptive active avoidance of danger. *Neuron* **84**, 1034–1048.

Barbosa A., Alves F. L., Fim Pereira A. de S., Ide L. M., Hoffmann A. (2012) Behavioral
characterization of the alarm reaction and anxiolytic-like effect of acute treatment with
fluoxetine in piauçu fish. *Physiol. Behav.* **105**, 784–790.

Bencan Z., Sledge D., Levin E. D. (2009) Buspirone, chlordiazepoxide and diazepam effects in a
zebrafish model of anxiety. *Pharmacol. Biochem. Behav.* **94**, 75–80.

Brandão M. L., Zanoveli J. M., Ruiz-Martinez R. C., Oliveira L. C., Landeira-Fernandez J. (2008)
Different patterns of freezing behavior organized in the periaqueductal gray of rats:
Association with different types of anxiety. *Behav. Brain Res.* **188**, 1–13.

Caldirola D., Grassi M., Alciati A., Riva A., Sangiorgio E., Daccò S., Perna G. (2016) Personalized
medicine in panic disorder: Where are we now? A meta-regression analysis. *Pers. Med.*
Psychiatry **1**–**2**, 26–38.

Caldirola D., Grassi M., Alciati A., Riva A., Sangiorgio E., Daccò S., Perna G. (2017) Personalized
medicine in panic disorder: where are we now? A meta-regression analysis. *Pers. Med.*
Psychiatry **1**–**2**, 26–38.

Carmo Silva R. X. do, Lima-Maximino M. G., Maximino C. (2018a) The aversive brain system of
teleosts: Implications for neuroscience and biological psychiatry. *Neurosci. Biobehav. Rev.* **95**,
123–135.

Carmo Silva R. X. do, Rocha S. P., Lima-Maximino M. G., Maximino C. (2018b) *Extraction of*
alarm substance in zebrafish.

Cloos J.-M. (2005) The treatment of panic disorder. *Curr. Opin. Psychiatry* **18**, 45–50.

Conselho Nacional de Controle de Experimentação Animal - CONCEA (2017) *Diretriz brasileira*
para o cuidado e a utilização de animais para fins científicos e didáticos - DBCA. Anexo I.
Peixes mantidos em instalações de instituições de ensino ou pesquisa científica. Brasil.

Cools R., Nakamura K., Daw N. D. (2011) Serotonin and dopamine: Unifying affective,
activational, and decision functions. *Neuropsychopharmacology* **36**, 98–113.

Curzon G., Fernando J. C. R., Marsden C. A. (1978) 5-hydroxytryptamine: The effects of impaired
synthesis on its metabolism and release in rat. *Br. J. Pharmacol.* **63**, 627–634.

Deakin J. F. W., Graeff F. G. (1991) 5-HT and mechanisms of defense. *J. Psychopharmacol.* **5**, 305–
315.

Døving K. B., Lastein S. (2009) The alarm reaction in fishes - Odorants, modulations of responses,
neural pathways. *Ann. N. Y. Acad. Sci.* **1170**, 413–423.

Dukhovich F. S., Darkhovskii M. B., Gorbatova E. N., Kurochkin V. K. (2004) *Molecular*
recognition: Pharmacological aspects. Nova Science, New York, NY.

Eachus H., Bright C., Cunliffe V. T., Placzek M., Wood J. D., Watt P. J. (2017) *Disrupted-in-*
Schizophrenia-1 is essential for normal hypothalamic-pituitary-interrenal (HPI) axis function.
Hum. Mol. Genet. **26**, 1992–2005.

Egan R. J., Bergner C. L., Hart P. C., Cachat J. M., Canavello P. R., Elegante M. F., Elkhayat S. I.,
et al. (2009) Understanding behavioral and physiological phenotypes of stress and anxiety in
zebrafish. *Behav. Brain Res.* **205**, 38–44.

Frisch K. von (1938) Zur psychologie des Fische-Schwarmes. *Naturwissenschaften* **26**, 601–606.

Frisch K. von (1941) Über einen Schreckstoff der fischhaut und seine biologische Bedeutung. *Z.*
Vgl. Physiol. **29**, 46–145.

Guimarães F. S., Carobrez A. P., Graeff F. G. (2008) Modulation of anxiety behaviors by 5-HT-
interacting drugs, in *Handb. Anxiety Fear*, (Blanchard R. J., Blanchard D. C., Griebel G., Nutt
D. J., eds), pp. 241–268. Elsevier B. V., Amsterdam.

Guimarães F. S., Zangrossi Jr H., Del-Ben C. M., Graeff F. G. (2010) Serotonin in panic and anxiety
disorders, in *Handb. Behav. Neurobiol. Serotonin*, (Müller C., Jacobs B., eds), pp. 667–685.
Elsevier B. V., Amsterdam.

Hale M. W., Lowry C. A. (2011) Functional topography of midbrain and pontine serotonergic
systems: Implications for synaptic regulation of serotonergic circuits. *Psychopharmacology*
(Berl). **213**, 243–264.

Harris G. P., Hapiak V. M., Wragg R. T., Miller S. B., Hughes L. J., Hobson R. J., Steven R.,
Bamber B., Komuniecki R. W. (2009) Three Distinct Amine Receptors Operating at Different
Levels within the Locomotory Circuit Are Each Essential for the Serotonergic Modulation of
Chemosensation in *Caenorhabditis elegans. J. Neurosci.* **29**, 1446–1456.

Herculano A. M., Maximino C. (2014) Serotonergic modulation of zebrafish behavior: Towards a
paradox. *Prog. Neuropsychopharmacol. Biol. Psychiatry* **55**, 50–66.

Herculano A. M., Puty B., Miranda V., Lima M. G., Maximino C. (2015) Interactions between
serotonin and glutamate-nitric oxide pathways in zebrafish scototaxis. *Pharmacol. Biochem.*
Behav. **129**, 97–104.

Hothorn T., Hornik K., Wiel M. A. van de, Zeileis A. (2006) A Lego system for conditional
inference. *Am. Statician* **60**, 257–263.

Huber P. J. (1981) *Robust statistics*. John Wiley & Sons, Inc, New York, NY.

Hüttel R. (1941) Die chemische Untersuchung des Schreckstoffes aus Elritzenhaut.
Naturwissenschaften **29**, 333–334.

Jesuthasan S. J., Mathuru A. S. (2008) The alarm response in zebrafish: Innate fear in a vertebrate
genetic model. *J. Neurogenet.* **22**, 211–229.

Kinkel M. D., Eames S. C., Philipson L. H., Prince V. E. (2010) Intraperitoneal injection into adult
zebrafish. *J. Vis. Exp.* **42**, 2126.

Kroes M. C., Henckens M. J., Homberg J. R. (2019) How serotonin transporter gene variance
affects defensive behaviours along the threat imminence continuum. *Curr. Opin. Behav. Sci.*
26, 25–31.

Leys C., Ley C., Klein O., Bernard P., Licata L. (2013) Detecting outliers: Do not use standard
deviation around the mean, use absolute deviation around the median. *J. Exp. Soc. Psychol.* **49**,
764–766.

Li Z., Li Y., Yi Y., Huang W., Yang S., Niu W., Zhang L., et al. (2012) Dissecting a central flip-flop
circuit that integrates contradictory sensory cues in *C. elegans* feeding regulation. *Nat.*
Commun. **3**, 776.

Mangiafico S. (2017) *rcompanion: Functions to Support Extension Education Program Evaluation.*

Matthews M., Varga Z. M. (2011) Anesthesia and euthanasia in Zebrafish. *ILAR J.* **53**, 192–204.

Maximino C. (2012) *Serotonin and anxiety. Neuroanatomical, pharmacological, and functional*
aspects. Springer, New York, NY.

Maximino C., Carmo Silva R. X. do, Santos Campos K. dos, Oliveira J. S. de, Rocha S. P., Pyterson
M. P., Santos Souza D. P. dos, et al. (2019) Sensory ecology of ostariophysan alarm
substances. *J. Fish Biol.* **95**, 274–286.

Maximino C., Lima M. G., Batista E. de J. O., Oliveira K. R. H. M., Herculano A. M. (2015)
Interaction between 5-HT1B receptors and nitric oxide in zebrafish responses to novelty.
Neurosci. Lett. **588**, 54–56.

Maximino C., Lima M. G., Costa C. C., Guedes I. M. L., Herculano A. M. (2014) Fluoxetine and
WAY 100,635 dissociate increases in scototaxis and analgesia induced by conspecific alarm
substance in zebrafish (Danio rerio Hamilton 1822). *Pharmacol. Biochem. Behav.* **124C**, 425–
433.

Maximino C., Puty B., Benzecry R., Araújo J., Lima M. G., Jesus Oliveira Batista E. de, Renata de
Matos Oliveira K., et al. (2013) Role of serotonin in zebrafish (*Danio rerio*) anxiety:
Relationship with serotonin levels and effect of buspirone, WAY 100635, SB 224289,
fluoxetine and para-chlorophenylalanine (pCPA) in two behavioral models.
Neuropharmacology **71**, 83–97.

Maximino C., Silva A. W. B. da, Gouveia Jr. A., Herculano A. M. (2011) Pharmacological analysis
of zebrafish (*Danio rerio*) scototaxis. *Prog. Neuropsychopharmacol. Biol. Psychiatry* **35**, 624–
631.

Nathan F. M., Ogawa S., Parhar I. S. (2015) Kisspeptin1 modulates odorant-evoked fear response
via two serotonin receptor subtypes (5-HT1A and 5-HT2) in zebrafish. *J. Neurochem.* **133**,
870–878.

Nowicki M., Tran S., Muraleetharan A., Markovic S., Gerlai R. (2014) Serotonin antagonists induce
anxiolytic and anxiogenic-like behavior in zebrafish in a receptor-subtype dependent manner.
Pharmacol. Biochem. Behav. **126**, 170–180.

Ogawa S., Nathan F. M., Parhar I. S. (2014) Habenular kisspeptin modulates fear in the zebrafish.
Proc. Natl. Acad. Sci. **111**, 3841–3846.

Parra K. V, Adrian Jr J. C., Gerlai R. (2009) The synthetic substance hypoxanthine 3-N-oxide elicits
alarm reactions in zebrafish (*Danio rerio*). *Behav. Brain Res*. **205**, 336–341.

Paul E. D., Johnson P. L., Shekhar A., Lowry C. A. (2014) The Deakin/Graeff hypothesis: Focus on
serotonergic inhibition of panic. *Neurosci. Biobehav. Rev*. **46**, 379–96.

Perusini J. N., Fanselow M. S. (2015) Neurobehavioral perspectives on the distinction between fear
and anxiety. *Learn. Mem*. **22**, 417–425.

Pimentel A., Carvalho T. dos S., Lima F., Lima-Maximino M., Soares M. C., Maximino C. (2019)
Conditional approach as cooperation in predator inspection: a role for serotonin? *Behav. Brain*
Res. **365**, 164–169.

Quadros V. A., Costa F. V., Canzian J., Nogueira C. W., Rosemberg D. B. (2018) Modulatory role of
conspecific alarm substance on aggression and brain monoamine oxidase activity in two
zebrafish populations. *Prog. Neuro-Psychopharmacology Biol. Psychiatry* **86**, 322–330.

Quadros V. A., Silveira A., Giuliani G. S., Didonet F., SIlveira A. S., Nunes M. E., Silva T. O., Loro
V. L., Rosemberg D. B. (2016) Strain- and context-dependent behavioural responses of acute
alarm substance exposure in zebrafish. *Behav. Processes* **122**, 1–11.

Sallinen V., Sundvik M., Reenilä I., Peitsaro N., Khrustalyov D., Anichtchik O., Toleikyte G.,
Kaslin J., Panula P. (2009) Hyperserotonergic phenotype after monoamine oxidase inhibition
in larval zebrafish. *J. Neurochem*. **109**, 403–415.

Santos J. M., Martinez R. C. R., Brandão M. L. (2006) Effects of acute and subchronic treatments
with fluoxetine and desipramine on the memory of fear in moderate and high-intensity
contextual conditioning. *Eur. J. Pharmacol*. **542**, 121–128.

Schirmer A., Jesuthasan S., Mathuru A. S. (2013) Tactile stimulation reduces fear in fish. *Front.*
Behav. Neurosci. **7**, Article 167.

Silva R. X. do C., Rocha S. P., Herculano A. M., Maximino M. L., Maximino C. (2019) Animal
models for panic disorder. *Psychol. Neurosci.*

Soares V. de P., Zangrossi Jr H. (2004) Involvement of 5-HT1A and 5-HT2 receptors of the dorsal
periaqueductal gray in the regulation of the defensive behaviors generated by the elevated T-
maze. *Brain Res. Bull*. **64**, 181–188.

Speedie N., Gerlai R. (2008) Alarm substance induced behavioral responses in zebrafish (*Danio*
rerio). *Behav. Brain Res*. **188**, 168–177.

Wolf K. (1963) Physiological salines for freshwater teleosts. *Prog. Fish-Culture* **25**, 135–140.

Zangrossi Jr H., Viana M. de B., Zanoveli J., Bueno C., Nogueira R. L., Graeff F. G. (2001)
Serotonergic regulation of inhibitory avoidance and one-way escape in the rat elevated T-maze.
Neurosci. Biobehav. Rev. **25**, 637–645.

Zanoveli J. M., Carvalho M. C., Cunha J. M., Brandão M. L. (2009) Extracellular serotonin level in
the basolateral nucleus of the amygdala and dorsal periaqueductal gray under unconditioned and
conditioned fear states: An in vivo microdialysis study. *Brain Res.* **1294**, 106–115.
722

Figure captions

Figure 1 – (A) Time-course of the experiments, with behavioral observations during two 6-min blocks, "CAS exposure" and "Post exposure", separated with a 1-min washout period. The boxes in red indicate the moment that the observation is made in each block. (B-D) Representation of the synaptic effects of the pharmacological manipulations: acute fluoxetine (B) is expected to increase synaptic and extra-synaptic serotonin levels, while metergoline (C) is expected to block receptors from the $5-HT_1$, $5-HT_2$, and $5-HT_7$ family; pCPA (D) is expected to decrease synaptic and extra-synaptic serotonin levels.

Figure 2 – Clonazepam (0.05 mg/kg) blocks all CAS-elicited increases in defensive behvavior during exposure. (A) Time spent on the top third of the tank; (B) Time spent on the bottom third of the tank; (C) Absolute turn angle; (D) Freezing; (E) Swimming speed. Different letters represent statistical differences at the $p < 0.05$ level; similar letters indicate lack of statistically significant differences. Data are presented as individual data points (dots) superimposed over the median ± interquartile ranges. CTRL = controls (water-exposed animals); CAS = conspecific alarm substance. Final sample sizes: CTRL + VEH: n = 15 animals; CTRL + CLZ: n = 15 animals; CAS + VEH: n = 17 animals; CAS + CLZ: n = 15 animals.

Figure 3 – Clonazepam (0.05 mg/kg) blocks only the CAS-elicited increases in bottom-dwelling after exposure. (A) Time spent on the top third of the tank; (B) Time spent on the bottom third of the tank; (C) Absolute turn angle; (D) Freezing; (E) Swimming speed. Different letters represent statistical differences at the $p < 0.05$ level; similar letters indicate lack of statistically significant differences. Data are presented as individual data points (dots) superimposed over the median ± interquartile ranges. CTRL = controls (water-exposed animals); CAS = conspecific alarm substance. Final sample sizes: CTRL + VEH: n = 15 animals; CTRL + CLZ: n = 15 animals; CAS + VEH: n = 17 animals; CAS + CLZ: n = 15 animals.

Figure 4 – **Acute fluoxetine dose-dependently blocks all the CAS-elicited increases in defensive behavior during exposure.** (A) Time spent on the top third of the tank; (B) Time spent on the bottom third of the tank; (C) Absolute turn angle; (D) Freezing; (E) Swimming speed. Different letters represent statistical differences at the $p < 0.05$ level; similar letters indicate lack of statistically significant differences. Data are presented as individual data points (dots) superimposed over the median ± interquartile ranges. CTRL = controls (water-exposed animals); CAS = conspecific alarm substance. Final sample sizes: CTRL + 0 mg/kg: n = 15 animals; CTRL + 2.5 mg/kg: n = 15 animals; CTRL + 25 mg/kg: n = 15 animals; CAS + 0 mg/kg: n = 15 animals; CAS + 2.5 mg/kg: n = 14 animals; CAS + 25 mg/kg: n = 15 animals.

Figure 5 – **Acute fluoxetine dose-dependently blocks only the CAS-elicited increases in bottom-dwelling after exposure.** (A) Time spent on the top third of the tank; (B) Time spent on the bottom third of the tank; (C) Absolute turn angle; (D) Freezing; (E) Swimming speed. Different letters represent statistical differences at the $p < 0.05$ level; similar letters indicate lack of statistically significant differences. Data are presented as individual data points (dots) superimposed over the median ± interquartile ranges. CTRL = controls (water-exposed animals); CAS = conspecific alarm substance. CTRL = controls (water-exposed animals); CAS = conspecific alarm substance. Final sample sizes: CTRL + 0 mg/kg: n = 15 animals; CTRL + 2.5 mg/kg: n = 15 animals; CTRL + 25 mg/kg: n = 15 animals; CAS + 0 mg/kg: n = 15 animals; CAS + 2.5 mg/kg: n = 14 animals; CAS + 25 mg/kg: n = 15 animals.

Figure 6 – **Metergoline does not block the CAS-elicited increases in defensive behavior after exposure.** (A) Time spent on the top third of the tank; (B) Time spent on the bottom third of the tank; (C) Absolute turn angle; (D) Freezing; (E) Swimming speed. Different letters represent

statistical differences at the $p < 0.05$ level; similar letters indicate lack of statistically significant

differences. Data are presented as individual data points (dots) superimposed over the median ±

interquartile ranges. CTRL = controls (water-exposed animals); CAS = conspecific alarm substance.

CTRL = controls (water-exposed animals); CAS = conspecific alarm substance. Final sample sizes:

CTRL + VEH: n = 14 animals; CTRL + MET: n = 15 animals; CAS + VEH: n = 15 animals; CAS +

MET: n = 15 animals.

Figure 7 – Metergoline blocks the CAS-elicited increases in bottom-dwellig and freezing after

exposure. (A) Time spent on the top third of the tank; (B) Time spent on the bottom third of the

tank; (C) Absolute turn angle; (D) Freezing; (E) Swimming speed. Different letters represent

statistical differences at the $p < 0.05$ level; similar letters indicate lack of statistically significant

differences. Data are presented as individual data points (dots) superimposed over the median ±

interquartile ranges. CTRL = controls (water-exposed animals); CAS = conspecific alarm substance.

Final sample sizes: CTRL + VEH: n = 14 animals; CTRL + MET: n = 15 animals; CAS + VEH: n =

15 animals; CAS + MET: n = 15 animals.

Figure 8 – pCPA does not block the CAS-elicited increases in defensive behavior during

exposure. (A) Time spent on the top third of the tank; (B) Time spent on the bottom third of the

tank; (C) Absolute turn angle; (D) Freezing; (E) Swimming speed. Different letters represent

statistical differences at the $p < 0.05$ level; similar letters indicate lack of statistically significant

differences. Data are presented as individual data points (dots) superimposed over the median ±

interquartile ranges. CTRL = controls (water-exposed animals); CAS = conspecific alarm substance.

Final sample sizes: CTRL + VEH: n = 12 animals; CTRL + pCPA: n = 13 animals; CAS + VEH: n

= 14 animals; CAS + pCPA: n = 13 animals.

Figure 9 – **pCPA blocks the CAS-elicited increases in bottom-dwellig and freezing after**

exposure. (A) Time spent on the top third of the tank; (B) Time spent on the bottom third of the

tank; (C) Absolute turn angle; (D) Freezing; (E) Swimming speed. Different letters represent

statistical differences at the $p < 0.05$ level; similar letters indicate lack of statistically significant

differences. Data are presented as individual data points (dots) superimposed over the median $\pm$

interquartile ranges. VEH = Vehicle (10% DMSO); pCPA = *para*-chlorophenylalanine; CTRL =

controls (water-exposed animals); CAS = conspecific alarm substance. Final sample sizes: CTRL +

VEH: n = 12 animals; CTRL + pCPA: n = 13 animals; CAS + VEH: n = 14 animals; CAS + pCPA:

n = 13 animals.

Figure 10 – Conspecific alarm substance (CAS) reduces the activity of monoamine oxidase in

the brain after exposure. Different letters represent statistical differences at the $p < 0.05$ level.

Data are presented as individual data points (dots) superimposed over the median $\pm$ interquartile

ranges. CTRL = controls (water-exposed animals); CAS = conspecific alarm substance. Final

sample sizes: CTRL: n = 10 animals; CAS: n = 10 animals.

Figure 11 – Hypothetical mechanism of the serotonergic signaling in zebrafish defensive

behavior during and after exposure to conspecific alarm substance. CAS elicits responses

dominated by erratic swimming, which decreases as the substance's concentrations decline. After

CAS exposure, the behavioral response is dominated by freezing. Serotonin shifts responding from

the first to the second (represented by the purple arrow, as well as by the arrows connecting the

raphe to the "switch" green boxes), putatively by switching control from the mesencephalic

aversive circuit ("switch OFF") to the prosencephalic aversive circuit ("switch ON").