1

ASSESSMENT OF ANESTHESIA ON PHYSIOLOGICAL STABILITY AND BOLD SIGNAL RELIABILITY

DURING VISUAL OR ACOUSTIC STIMULATION IN THE CAT

Alexandra Levine[2,3†], Benson Li[1†], Paisley Barnes[2‡], Stephen G. Lomber[1-4‡], and Blake E. Butler[*1,3,4]

[1]Department of Psychology

[2]Department of Physiology & Pharmacology

[3]Brain and Mind Institute

[4]National Centre for Audiology

[†]Authors made equal contributions

University of Western Ontario

London, Ontario, Canada N6A 5B7

[‡]Present address:

Department of Physiology

McGill University

Montréal, Québec, Canada H3G 1Y6

[*]**Correspondence should be addressed to:**

Dr. Blake Butler

Email: bbutler9@uwo.ca

Phone: 519-661-2111 x85831

Abstract:

Background: Neuroimaging methods including fMRI provide powerful tools to observe whole-brain functional networks. This is particularly powerful in animal models, allowing these networks to be probed using complementary methods. However, most animals must be anesthetized for neuroimaging, giving rise to complications resulting from anesthetic effects on the animal's physiological and neurological functions. For example, an established protocol for feline neuroimaging involves co-administration of ketamine and isoflurane – the latter of which is known to suppress cortical function.

New Method: Here, we compare this established protocol to alfaxalone, a single-agent anesthetic for functional neuroimaging. We first compare the two in a controlled environment to assess relative safety and to measure physiological stability over an extended time window. We then compare patterns of auditory and visually-evoked activity measured at 7T to assess mean signal strength and between-subjects signal variability.

Results in Comparison with Existing Methods: We show that alfaxalone results in more stable heart and respiratory rates over the 120 minutes testing period, with evidence of smaller between measurements variability in respiratory rates within this time window, when compared to ketamine plus isoflurane. Moreover, we demonstrate that both agents evoke similar mean BOLD signals across animals, but that alfaxalone elicits more consistent BOLD activity in response to sound stimuli across all ROIs observed.

Conclusions: Alfaxalone is observed to be more physiologically stable, evoking a more consistent BOLD signal across animals than the co-administration of ketamine and isoflurane. Thus, an alfaxalone-based protocol may represent a better approach for neuroimaging in animal models requiring anesthesia.

Keywords: anesthesia, fMRI, physiological stability, animal models, stimulus-evoked, BOLD

1. Introduction

Experiments in animal models continue to be critical for understanding the structure and function of the brain. For decades, cats have been successfully used as an animal model to study sensory systems, largely due to the remarkable similarities that they share with human cell types, neural pathways, and cytoarchitecture (Blake, 1979). Electrophysiological (Hubel and Wiesel, 1962; Liu et al., 2010), neuroanatomical (Lomber et al., 1995; Wong et al., 2015; Butler et al., 2016), behavioral (Heffner and Heffner, 1988; Wong et al., 2018), and imaging studies (Brown et al., 2014; Butler et al., 2015; Stolzberg et al., 2018) undertaken in the cat have played a critical role in advancing our knowledge of neural processing within visual and auditory systems, and the interactions between the two.

Recently, there has been increased interest in non-invasive neuroimaging methods like functional magnetic resonance imaging (fMRI) for the study of sensory system function in animal models. This approach offers several advantages, including the ability to examine perception at the whole-brain level, and the ability to undertake longitudinal, within-animal studies of sensory system development (which in turn aids in reducing the number of animals required to power meaningful comparisons). This offers a distinct advantage over other methods such as electrophysiological studies, which are highly invasive and are limited in their capacity to evaluate processes occurring over spatially disparate neural networks. fMRI measures changes in the ratio of oxygenated to deoxygenated blood, or blood-oxygen-level-dependent (BOLD) signals (Ogawa et al., 1990). An increase in this BOLD signal is thought to reflect increased neuronal activity compared to a baseline measurement (Buxton and Frank, 1997; Logothetis et al., 2001; Ferris et al., 2006). Moreover, by measuring the temporal coherence of the BOLD signal across spatially disparate areas of the brain, it is possible to estimate the degree to which these areas are functionally connected into networks that support perception and associated behaviors.

Modern fMRI scanners can provide spatial resolution with 1 mm precision, but the accuracy of these measures depends critically on minimizing subject movement within the scanner. While most human participants can be instructed to remain still during imaging sessions, this is not possible in most other animals, and meaningful measurements must thus

be taken under anesthesia. Anesthetic agents help minimize potential stress and fluctuations in

behavior that can affect the quality of data retrieved. However, the use of anesthetics during

fMRI necessitates due consideration be given to the effects of the drugs themselves (Ueki et al.,

1992; Biermann et al., 2012), as these agents can influence neurovasculature by changing

cerebral blood flow, blood volume, and rate of oxygen metabolism (Gao et al., 2017), and can

suppress neuronal activity by reducing excitatory synaptic transmission or increasing inhibitory

transmission (Richards, 1983). Further complications may include physiological variability based

on the drug type, concentration, and route of administration (Peng et al., 2010; Nagore et al.,

2013; Aksenov et al., 2015; Ros et al., 2017). Therefore, there is a need to establish a robust and

reliable anesthetic protocol that will facilitate bridging the gap between animal and human

neuronal organization and function.

Fortunately, decades of electrophysiological work in the cat has revealed a great deal

about the effects of different anesthetic agents on recorded neural function. A common

protocol involves the continuous infusion of ketamine alongside other agents such as sodium

pentobarbital, xylazine, or diazepam to induce and maintain anesthesia (e.g. Heil and Irvine,

1998; Miller et al., 2002; Pienkowski and Eggermont, 2009). This protocol has evolved over

time; for example, early studies found that ketamine infusion reduces spontaneous and peak

firing rates in auditory cortex (Zurita et al., 1994), possibly due to altered sensory perception

and reduced cortical glucose metabolism (Crosby et al., 1982; Oye et al., 1992). To reduce the

amount of ketamine required for anesthesia (and reduce these suppressive effects,

accordingly), Jezzard et al. (1997) proposed to pre-medicate with ketamine but maintain

sedation with isoflurane (1-2%, gas). However, other studies showed that isoflurane redirected

cerebral blood flow, and reduced neural activity recorded in various visual brain regions by up

to 50% (Harel et al., 2002; Olman et al., 2003; White & Alkire, 2002). Isoflurane has recently

been shown to suppress resting-state connectivity in the primary somatosensory cortex of non-

human primates as well (Wu et al., 2016). Therefore, when establishing the protocol for initial

fMRI experiments in the cat, Brown et al. (2013) reduced isoflurane concentrations to the

minimum level required to maintain sedation (0.4-0.5%) and supplemented with a continuous

rate infusion of ketamine (0.6-0.75 mg/kg/hr). Under this protocol, the authors were able to

record BOLD signal changes up to 6% in some but not all auditory regions. The protocol was

used in subsequent auditory-evoked studies (Hall et al., 2014; Butler et al., 2015) and in an

examination of resting-state connectivity using fMRI (Stolzberg et al., 2018).

In spite of successive revisions, the combination of isoflurane and ketamine is known to

result in widespread cortical deactivation in other animals (e.g. Hodkinson et al., 2012).

Moreover, these effects appear to differ by brain region/sensory modality such that the co-

administration of ketamine and isoflurane may limit the ability to study sensory processes

beyond audition (Oye et al., 1992; Ries & Puil, 1999; Höflich et al., 2017). Thus, there remains a

need to develop an anesthetic protocol that can induce and maintain a light anesthetic plane

sufficient to suppress movement, without drastic reductions in cortical activity across multiple

brain regions.

Several potential agents were considered in the current study, but many were excluded

because their mechanism of action was deemed not conducive for measuring BOLD signals

(Table 1). Others were eliminated in consultation with veterinary care staff due to concerns

with respect to physiological effects. As a result, alfaxalone was considered to be the strongest

candidate protocol to serve as an alternative to the coadministration of ketamine and

isoflurane as a primary anesthetic agent for fMRI. Alfaxalone: i) has dose-dependent effects on

cardiovasculature, respiration, neuronal activity, and neuromusculature (Warne et al., 2015;

Whittem et al., 2008; Muir et al., 2009; Taboada and Murison, 2010; Baldy-Moulinier et al.,

1975) which may allow for more predictable changes in BOLD response; ii) has been found to

sufficiently maintain a stable level of anesthesia for up to 2-hours as a stand-alone agent

(Tamura et al., 2015; Deutsch et al., 2017) which in our experience is the typical amount of time

required for neuroimaging in cats; and iii) has been successfully administered intravenously to

maintain anesthesia in cats during surgical procedures in our own laboratory, and by others

(Beths et al., 2014; Nagakubo et al., 2017) with minimal physiological side effects.

Table 1. Studies examining cardiovascular, respiratory, and neural effects of anesthetic agents.

Anesthetic	Primary Mechanism	First Author and Year	Reason for Exclusion
Pentobarbital	GABA$_A$ Agonist	Kaitin (1985)	Suppressed cortical activity
		Morin-Surun et al. (1984)	Inhibition of respiratory neurons
Propofol	GABA$_A$ Agonist	Lahti et al. (1999)	Decreased BOLD signal intensity
		Bonhomme et al. (2000)	Reduced cerebral blood flow
		Dueck et al. (2005)	Decreased BOLD signal intensity
Butorphanol	Opioid (κ-type)	Paddleford (1999)	Long-acting (up to 4-hours)
Fentanyl	Opioid (μ-type)	Peng et al. (2010)	Suppressed cortical activity
		Freeman et al. (1967)	Reduced cortical blood flow
Dexmedetomidine	Adrenergic (α$_2$) Agonist	Fukuda et al., 2013	Reduced cerebral blood flow

Here, we compare an alfaxalone (Alf) protocol with a previously established protocol consisting of a combination of isoflurane and ketamine (Iso+Ket), providing detailed measures of physiological stability as well as evoked activity in cats during fMRI. The investigation is separated into two parts; in the first study, we evaluate the physiological stability of each protocol in an operating suite. A light anesthetic plane was maintained for a minimum of 2-hours while cats were exposed to mock scanner noises at 90 dB and vital signs (heart rate, respiratory rate, end-tidal CO_2, blood pressure, etc.) were recorded. In the second study, both protocols were employed in the scanner while BOLD signal changes were recorded in response to visual and auditory stimuli. This study is the first to evaluate different anesthetic protocols in cats by directly comparing BOLD signal responses. Quantification of these signals will provide insight into the contribution of different anesthetics on neural activity, and potentially offer an alternative option to the combination of isoflurane and ketamine.

2. Methods

2.1 Animals

Two healthy adult domestic short-hair cats were used in study one, and a total of 12 cats were compared in the second study. Animals were born to pregnant queens obtained from a commercial laboratory animal breeding facility (Liberty Labs, Waverly, NY), and were housed as a clowder. Normal hearing status was confirmed at approximately 3 months of age using auditory brainstem responses. All procedures were conducted in accordance with the Canadian Council on Animal Care's Guide to the Care and Use of Experimental Animals and were approved by the University of Western Ontario Animal Use Subcommittee of the University Council on Animal Care.

2.2 Study 1: Physiological Stability during Anesthesia

2.2.1 Anesthesia

The first study was conducted in a surgical suite in order to evaluate the safety and stability of the selected protocols. In the alfaxalone protocol, the animal was first pre-medicated with dexdomitor (0.04 mg/kg, i.m.) prior to catheter placement. Sedation was confirmed after 10 minutes by the absence of a paw-pinch reflex. Ophthalmic ointment was applied to prevent drying of the eyes, body temperature was maintained at 37°C using a circulating warm water pad, and an indwelling 22g catheter was placed in the cephalic vein to facilitate maintenance of anesthesia. A bolus dose of alfaxalone (0.3-0.5 ml, i.v.) was administered to achieve deeper anesthesia and the animal's larynx was sprayed with xylocaine prior to intubation. The animal was placed in a sternal position on the surgical table, and anesthesia was maintained through continuous infusion of alfaxalone (7 mg/kg/hr, i.v.), while 100% oxygen was provided at a rate of 1.0L/min. Finally, a bolus dose of atipamezole (0.27 ml, i.m.) was administered to reverse any residual effects of the dexdomitor.

For the ketamine plus isoflurane protocol, animals were pre-medicated with a combination of dexdomitor (0.022 mg/kg, i.m.), ketamine (4 mg/kg, i.m.), and acepromazine (0.05 mg/kg, i.m.). Sedation was confirmed, the animal's core temperature was maintained, ophthalmic ointment was applied, and a catheter was placed for anesthetic maintenance as

above. The animal was placed in a sternal position on the surgical table, and a continuous

infusion of ketamine (5 ml/kg/hr, i.v.), combined with gaseous isoflurane (0.5% in oxygen

provided at a rate of 1.0L/min) was used to maintain anesthesia. The reversal of dexdomitor

was not necessary in this protocol as premedication volume was lower and consequently would

not be expected to have effects lasting into the experimental period. Approximately 60 minutes

into the session, the rate of ketamine infusion was increased to 6.25 ml/kg/hr (i.v.) and

isoflurane was reduced to 0.25% in order to mimic the protocol developed previously for

imaging, in which these changes are required prior to functional image acquisition to optimize

BOLD signal.

At the end of each session, anesthesia was discontinued, and animals were monitored

until they recovered fully from anesthetic effects. Animals anesthetized with alfaxalone

received a bolus dose of butorphanol (0.2 mg/kg, s.c.; opioid analgesic) to counteract

hyperkinesia, a side-effect commonly observed during post-anesthetic recovery from prolonged

IV administration of alfaxalone in cats (Whittem et al., 2008). The intubation tube was removed

when the animal exhibited a gag reflex and increased jaw tone, and following recovery, the

indwelling catheter was removed and the animal was returned to their clowder. Each agent was

tested twice in each animal for a total of 4 sessions per agent.

2.2.2 Data Recording

To mimic conditions in the scanner, the animal was presented with previously recorded

scanner noise through foam insert earbuds (Sensimetric S14) at 90 dB SPL for the duration of

experimental sessions. Each agent's ability to induce anesthesia, maintain a lightly sedated

state for 2-hours, and to allow for uneventful recovery was noted. Anesthetic and physiological

stability was evaluated by monitoring and recording parameters including autonomic reflexes

(e.g. paw-pinch, gag, palpebral) and vital signs (e.g. heart rate, end-tidal CO_2, respiratory rate,

peripheral capillary oxygen saturation, blood pressure, and mean arterial pressure) in 5-minute

intervals.

2.3 Study 2: fMRI

The second study sought to compare the auditory- and visually- evoked BOLD signals recorded while animals were anesthetized with each candidate agent. A group of 6 cats were scanned while anesthetized with alfaxalone, and results were compared to a group of 6 sex- and age-matched animals scanned previously using the exact same equipment and experimental procedure.

2.3.1 Animal Preparation and Anesthesia

For both the ketamine plus isoflurane and alfaxalone protocols, anesthesia was induced and maintained as described for Study 1 above. Once anesthetized, the animal was placed in a sternal position within a custom-built Plexiglass sled. Phenylephrine hydrochloride and atropine sulfate ophthalmic solutions were applied to both eyes to dilate the pupils and retract the nictitating membranes. Lubricated contact lenses were placed in both eyes (a blackout lens in the left eye, and a clear lens in the right eye). This permitted visual stimuli to be brought into focus on the retina and have visual signals preferentially sent to the left hemisphere. MRI-compatible foam insert earphones (Sensitmetrics S14) were inserted in each ear to allow for the presentation of auditory stimuli, and the animal's head was stabilized within a custom 8-channel radio-frequency (RF) coil. Vital signs (heart rate, respiratory rate, end-tidal CO_2, inspiratory CO_2, percent oxygen saturation, systolic/diastolic and mean blood pressure, and rectal body temperature) were monitored throughout the scanning session. At the conclusion of the imaging session, anesthesia was discontinued and animals were recovered as outlined in Study 1 above.

2.3.2 Stimuli

Visual stimuli were generated with PsychoPy (Peirce, 2007; 2009) and presented through a Dell laptop to an Avotec SV-6011 Rear Projector. From their sternal position within the bore of the magnet, the animals eyes were located approximately 75 cm from an acrylic screen (H = 14.5 cm, W = 19cm), which was viewed through a custom-built mirrored periscope. The stimulus extended 14.5 visual degrees horizontally and 11 degrees vertically and consisted of a black and white flickering checkerboard (8 ring-segments of 16 wedges) on a grey

background (100% luminance contrast, 50% luminance background), counter-phase flickering at

5Hz. The stimulus was arranged in a simple ON/OFF block design, where the OFF block

consisted of a blank grey screen, each block lasting 30s. The animal's gaze was assessed visually

through the scanner bore before the acrylic screen was placed at the end of the bore.

Auditory stimuli were generated using Audacity® recording and editing software

(Audacity Team, 2019), and consisted of a 30s stimulus consisting of 400ms broadband noises

separated by 100ms silent gaps. This stimulus was arranged in an ON/OFF block design, where

the OFF block consisted of a 30s period of silence. Sounds were presented diotically from a Dell

laptop through an external Roland Corporation soundcard (24-bit/96 kHz; Model UA- 25EX), a

PylePro power amplifier (Model PCAU11), and Sensimetrics MRI-compatible ear inserts (Model

S14). All stimuli were calibrated to 80-90 dB SPL using an ear simulator (Bruel & Kjaer, Model

#4157).

2.3.3 Scanning Parameters

Data were collected using an ultra-high-field 7T Siemens MRI human head-only scanner

located at the Centre for Functional and Metabolic Mapping at the Robarts Research Institute

operating at a 350 mT/m/s slew rate. An automated 3D mapping procedure (Klassen & Menon,

2004) was used to optimize the magnetic field (B_0 shimming) over the specific volume of

interest.

High-resolution structural T1-weighted MP2RAGE images were acquired prior to

functional scanning with the following parameters: isotropic voxel size of $0.5mm^3$, 80 slices,

FoV=96mm, TE=3.95ms, TR= 6000ms, TI =800ms, and a flip angle of 4. Functional images were

acquired over the whole brain in axial orientation with a single shot echo-planar imaging (EPI)

acquisition with grappa acceleration and the following parameters: isotropic voxels $1mm^3$, 38

slices (interleaved), FoV=72mm, TE=22.0ms, TR= 2000ms and a flip angle of 60 degrees. Each

functional scan (visual- and auditory-evoked) lasted six minutes, and consisted of alternating 30

second blocks of stimulus and baseline conditions.

2.3.4 Image Analysis

T1-weighted structural images were processed with a combined approach of automated

and manual processing. The structural images were skull-stripped with use of MRIcron (NITRC;

Rorden & Brett, 2000) and FSLmath functions (FMRIB's toolbox, Oxford, UK; Smith et al., 2004,

http://fsl.fmrib.ox.ac.uk/fsl/fslwiki/).

First-level statistical analysis of each animal's functional data was carried out using FEAT

processing in FSL (Woolrich et al., 2001). The functional images were skull stripped using FSL's

Brain Extraction Tool (BET). Preprocessing began with the removal of the first 2 volumes (4s) of

the scan to allow for the scanner magnetic field to stabilize and reach magnetic saturation.

The following were also applied: motion correction (MCFLIRT; though the movement is

nearly non-existent during these procedures), spatial smoothing (Gaussian, FWHM, 2mm) and a

temporal high-pass filter cut off (0.01Hz/60s). First-level general linear model analysis (FILM)

was then carried out, where regressors for each condition-block were convolved with a gamma

hemodynamic response function (phase = 0, standard deviation = 3s, mean lag = 6s; the BOLD

signal time course in cats has been shown to closely resemble that observed in humans and

non-human primates [Brown et al., 2013]).

Each individual EPI sequence underwent time series pre-whitening (Smith et al., 2004),

allowing us to carry through contrasts for higher-level analysis to test for group effects;

individual animal GLM results were co-registered to the coordinate space of the high-resolution

structural image for each participant using FMRIBs Linear Image Registration Tool (FLIRT;

Jenkinson et al., 2002). Further analysis compared differences in average BOLD signal change

under each anesthetic agent across all voxels within a given region of interest. Visual ROIs

included primary visual cortical areas 17, 18 and 19, as well as the lateral geniculate nuclei

(LGN) of the thalamus. Auditory ROIs included primary (A1) and second (A2) auditory cortex

and the medial geniculate nuclei (MGN) of the thalamus.

Mean BOLD signal changes (relative to baseline) evoked by visual and auditory stimuli

were extracted for each ROI of each animal, using FEATquery (FMRIB toolbox in FSL). To carry

this out, FEATquery takes each participant's high-resolution structural scan and co-registers it

to the feline template space (CATLAS, Stolzberg et al., 2017) using FLIRT multi-registration, in

which all the predefined functional ROIs are defined (Fig. 1).

Figure 1. Visual and auditory regions of interest (ROIs) used in analysis displayed in
feline template space (CATLAS; Stolzberg et al., 2017). Visual ROIs (LGN, 17/18/19)
contralateral to the open eye were isolated for analysis.

3. Results

3.1 Study 1 - Physiology

Two animals were observed two times under each anesthetic protocol in the operating

suite to evaluate the stability of heart and respiratory rates across anesthetics. These

physiological measures are commonly yoked; in practice, respiratory rate is used as an indicator

of anesthetic depth where high rates are more representative of an awake state (Myles, 2007).

The data from the present evaluation, averaged across animals and runs for each anesthetic

agent tested, are presented in Figure 2A. It is apparent that both measures exhibit more

stability across the duration of testing under alfaxalone than under ketamine plus isoflurane. To

examine the variability within a test session, the change in each measure between successive

timepoints (i.e. the change in heart/respiratory rate across each 5-minute interval) was

calculated; these values are presented in Figure 2B. On this time scale, changes in heart rate

were very similar across anesthetics; however, the changes in respiratory rate under alfaxalone

appear smaller, suggesting more within-session stability under this anesthetic when compared

to coadministration of ketamine and isoflurane.

Figure 2. A) Normalized respiratory rates (breaths per min) and heart rates (beats
per min) under alfaxalone and co-administered ketamine and isoflurane. Data were
normalized to the mean rate for an individual run, and then normalized values were
averaged across animals and runs. Shaded regions represent the standard error of
the mean. The first 10 minutes of each measurement period were omitted from
analysis to account for setup of monitoring/recording devices. B) A histogram
representing change magnitude in successive measurements of respiratory and
heart rate under each anesthetic tested. Lower values indicate relative stability in
the variable measured, while larger values indicate increased variability over time.

It is important to note that increased variability under ketamine plus isoflurane is not

entirely due to the drugs per se; imaging under this protocol requires that the rate of ketamine

infusion be increased and the concentration of isoflurane reduced approximately 60 minutes

into a testing session in order to acquire functional images with measurable BOLD signal (Brown

et al., 2013; Stolzberg et al., 2018). As a result, physiological measures such as heart rate and

respiratory rate often change dramatically at this point in time (observable in Figure 2A). As

described above, these shifts indicate drift towards a lighter anesthetic plane, as is necessary to

provide increased BOLD signal in the presence of isoflurane. However, this also increases the

risk of the animal becoming alert, which should be avoided. In contrast, the rate of alfaxalone

infusion can remain unchanged for the duration of the session, resulting in more stable vital

signs throughout.

3.2 Study 2 - fMRI

3.2.1 BOLD signal strength

To examine whether a global difference in BOLD signal amplitude exists between

anesthetics, normalized percent BOLD signal changes (hereafter referred to as BOLD signals)

were extracted from all selected ROIs, in each animal.

As an initial test, a mixed ANOVA was performed on these BOLD signals, where

anesthetic protocol (ketamine plus isolflurane/alfaxalone) was treated as a between-subjects

factor, and within-subject factors included two stimulus types (auditory/visual) and seven ROIs

(auditory cortical areas A1 and A2, the MGB, visual cortical areas 17, 18, and 19, and the LGN).

A significant interaction was observed between stimulus type and ROI (F[11, 110] = 13.505, $p <$

0.001) demonstrating that across anesthetic protocols, the BOLD signal observed in a given ROI

depended on the nature of the stimulus presented. The comparison of between-subjects

effects across all regions and conditions revealed no significant effect of anesthetic protocol

(F[1, 10] = 1.108, p = 0.337). Overall, these results indicate that, as expected, auditory and

visual stimuli evoke different patterns of activity across sensory brain regions. Moreover, the

patterns of evoked activity are similar across the anesthetics tested. While this suggests that

both alfaxalone and co-administered ketamine and isoflurane may be appropriate for imaging

experiments, this analysis provides little insight into the consistency and stability of the BOLD

signal measured under each. We thus conducted planned post-hoc tests investigating the

variability of these signals.

3.2.2 BOLD signal variability

Figure 3 shows the BOLD signals recorded in each ROI broken down by stimulus type

(auditory, visual) and anesthetic protocol for each animal tested. Doing so allows BOLD signal

variability to be observed across all ROIs (bilateral MGN, A1, & A2; LGN, 17, 18, & 19

contralateral to the opened eye) and individual subjects. Across regions of interest, between-

subjects variability in the BOLD signal evoked by stimuli to which a region is typically responsive

(i.e. the signal recorded in A1/A2 in response to sound) was greater under ketamine plus

isoflurane than under alfaxalone. Further, signals recorded in response to stimuli to which an

ROI is *not* typically responsive (i.e. the signal recorded in areas 17/18/19 in response to sound),

were also far more variable under ketamine plus isoflurane. This latter analysis provides a

measure of signal variability in the absence of evoked activity (i.e. a measure of background

activity under a given anesthetic agent).

Figure 3 Individual z-scores of BOLD signal changes within all regions of interest in
response to visual (a & b) and auditory (c & d) stimulation under coadministration
of ketamine and isoflurane (a & c) or alfaxalone (b & d). Data are thresholded using

clusters determined by Z>2.3 and a corrected cluster significance threshold of
p=0.05 (Worsley, 2001).
375

To quantify variability in more detail, the range of the normalized BOLD signal across

animals for each region of interest and anesthetic protocol is presented in Figure 4. With

respect to visually-evoked activity, the between-subjects variability in BOLD activity was highly

similar across all ROIs. In response to sound, the signal recorded was less variable under

alfaxalone than under co-administered ketamine and isoflurane across all ROIs. Thus, while

both anesthetics evoke similar mean BOLD signal amplitudes (as evidenced by the absence of a

statistically significant effect of anesthetic protocol, as described above), between-subjects

variability is decreased under alfaxalone, notably in response to sounds.

384
385   **Figure 4** Range (max-min) of z-scored percent BOLD signal change across animals for
386   each region of interest studied, in response to auditory and visual stimuli under
387   alfaxalone (blue) and co-administered ketamine and isoflurane (red).
388

4. Discussion

This study was undertaken to compare and contrast potential anesthetic protocols for

functional neuroimaging with a focus on 1) physiological stability and 2) optimizing BOLD signal

across stimulus modalities. While all anesthetics affect neural function, their use remains

necessary in most animal models to minimize movement and stress, and to ensure animal

safety within the magnet. Thus, it is important to establish a regime that strikes a balance

between achieving and maintaining safe and stable anesthesia, while optimizing cortical

activity. Previous studies have used a variety of anesthetic agents in an attempt to optimize

responses; one common protocol involves the co-administration of ketamine and isoflurane–

an approach that has been shown to allow for the observation of sound-evoked BOLD activity

(Brown et al., 2013; Hall et al., 2014; Butler et al., 2015; Stolzberg et al., 2018). However, this

combination can produce physiological instability and has suppressive effects on cortical

activity (Zurita et al., 1994; Harel et al., 2002; Olman et al., 2003; Hodkinson et al., 2012). It

therefore remains important to explore alternative and perhaps more reliable protocols. Here,

we provide physiological and neuroimaging evidence that supports the use of alfaxalone as a

stable and consistent anesthetic agent for neuroimaging.

4.1 Anesthesia and Physiology

The primary goal of the physiological pilot described above, was to determine the safety

and stability of candidate protocols using heart and respiratory rates as indicators. Importantly,

these measures have been shown to reflect anesthetic depth (Musizza & Ribaric, 2010; Thomas

& Lerche, 2011). Moreover, fluctuations in either heart or respiratory rate have been shown to

interfere with neural signals measured by fMRI (Gao et al., 2017; Birn et al., 2003; Abbott et al.,

2005; Kastrup et al., 1999). In the present study, both heart and respiratory rates appear more

stable across a 120-minute testing duration under alfaxalone than under co-administered

ketamine and isoflurane. In part, this reflects a simultaneous increase in ketamine infusion rate

and decrease in isoflurane concentration approximately 60 minutes into an imaging session

that is necessary to allow for BOLD signal visualization (Brown et al., 2013; Hall et al., 2014;

Butler et al., 2015; Stolzberg et al., 2018). Thus, this previously established protocol includes a

trade-off between measurable BOLD responses and increased risk of the animal becoming alert

in the scanner. Conversely, the alfaxalone protocol described here maintains a constant

infusion rate for the duration of the experimental session, and thus results in only small-scale

changes in heart and respiratory rate over time, consistent with previous examinations of the

anesthetic properties of alfaxalone for other applications (Beths et al., 2014; Muir et al., 2009).

In addition to long-term stability, respiratory rate under alfaxalone was also shown to be more

stable across shorter duration measurement intervals (5 min; Figure 2B). Both protocols

examined attained safe levels of anesthetic depth for functional imaging; however, alfaxalone

resulted in more physiological stability across individuals and sessions when compared to the

co-administration of isoflurane and ketamine.

4.2 Anesthesia and BOLD

Having demonstrated the safety of both candidate protocols, the imaging experiment

(Study 2) sought to compare and contrast BOLD signal changes evoked in auditory and visual

thalamic and cortical regions of interest under each anesthetic. To the best of our knowledge,

this is the first study to provide such a comparison. Here, we demonstrate that both protocols

facilitate comparable mean levels of overall evoked BOLD activity across ROIs. These results are

in accordance with similar mechanisms of action between alfaxalone and isoflurane (Lambert et

al., 2003; Nakahiro et al., 1999). Examining these patterns of evoked activity in more detail,

more reliable and consistent neural responses were observed under alfaxalone, evidenced by

decreased BOLD signal variability across animals (Figures 3 & 4). This is highly important, as

fMRI analyses often involve averaging or subtractive computations across blocks of data

acquired over the duration of an imaging session. That the activity recorded under co-

administered ketamine and isoflurane is highly variable within a given ROI means these

between-block contrasts may be particularly susceptible to anesthetic effects. Interestingly,

signals measured in brain regions not traditionally associated with a particular stimulus

modality (e.g. BOLD signal estimates from primary visual cortex in response to auditory

stimulation) were more variable under co-administered ketamine and isoflurane than under

alfaxalone, suggesting signal variability associated with the former extends well beyond effects

on stimulus-evoked activity. This activity may reflect between-subjects differences in non-

selective suppression observed under isoflurane (Wu et al., 2016) or the potentially dissociative

effects of ketamine (Abel et al., 2003) – either of which presents a challenge to the

interpretation of stimulus-evoked signals. Increased consistency between individual animals

and ROIs under alfaxalone suggests that it may be better suited for fMRI studies than the co-

administration of isoflurane and ketamine.

5. Conclusion

452   Anesthetics are important to many experimental approaches in animal research. However, without good, consistently applied protocols for neuroimaging, findings from these studies remain difficult to consolidate, and the degree to which they can be generalized to understand the brain in its natural neural state remains unclear. The present study sought to compare alfaxalone as a novel imaging anesthetic, to the previously established protocol for feline functional imaging studies that relied upon the co-administration of ketamine and isoflurane (Brown et al., 2013; Hall et al., 2014; Butler et al., 2015; Stolzberg et al., 2018). We provide evidence that alfaxalone is more physiologically stable and allows for the observation of more consistent patterns of BOLD activity. Additionally, there are some practical implications that favor the use of alfaxalone: 1) a single-agent protocol is easier to maintain over extended testing, is preferred by veterinary staff, and reduces concerns related to drug interactions; 2) unlike ketamine, alfaxalone in not a controlled substance, and is thus easier to obtain and store; and 3) because anesthetic depth is more consistent across the testing session under alfaxalone, the duration of testing is more predictable and concerns related to the animal waking up within the magnet are reduced. For all of these reasons, we therefore propose that alfaxalone may be a superior anesthetic agent for the safe and reliable collection of fMRI data.

6. Acknowledgments

470   We would like to thank Stephen Gordon and Alessandra Sacco for their assistance with animal care and data collection, and Trevor Szekeres for technical assistance. This research was undertaken thanks to funding from the Canada First Research Excellence Fund awarded to BrainsCAN at Western University, the Natural Sciences and Engineering Research Council of Canada, Canadian Institutes of Health Research, and the Canada Foundation for Innovation.

7. References

Abbott, D.F., Opdam, H.I., Briellmann, R.S., & Jackson, G.D. (2005). Brief breath holding may confound functional magnetic resonance imaging studies. *Human Brain Mapping*, *24*(4), 284-290.

Abel, K. M., Allin, M. P., Kucharska-Pietura, K., Andrew, C., Williams, S., David, A. S., & Phillips, M. L. (2003). Ketamine and fMRI BOLD signal: distinguishing between effects mediated by change in blood flow versus change in cognitive state. Human brain mapping, 18(2), 135-145.

Aksenov, D.P., Limin, L., Miller, M.J., Iordanescu, G., & Wyrwicz, A.M. (2015). Effects of anesthesia on BOLD signal and neuronal activity in the somatosensory cortex. *Journal of Cerebral Blood Flow and Metabolism*, *35*(11), 1819-1826.

Audacity Team (2019). Audacity®: Free audio editor and recorder [computer application]. Version 2.3.1 retrieved May 6, 2019 from https://audacityteam.org/.

Baldy-Moulinier, M., Basset-Lehmann, J., & Passonaut, P. (1975). Effects of combination alfaxalone and alfadolone, anesthetic derivatives of pregnenedione, on cerebral hemodynamics in cats. *Comptes Rendus Des Seances de la Societe de Biologie et de Ses Filiales*, *169*(1), 126-131.

Beths, T., Touzot-Jourde, G., Musk, G., & Pasloske, K. (2014). Clinical evaluation of alfaxalone to induce and maintain anesthesia in cats undergoing neutering procedures. *Journal of Feline Medicine and Surgery*, *16*(8), 609-615.

Biermann, K., Hungerbühler, S., Mischke, R., & Kästner, S.B.R. (2012). Sedative, cardiovascular, haematologic and biochemical effects of four different drug combinations administered intramuscularly in cats. *Veterinary Anaesthesia and Analgesia*, *39*(2), 137-150.

Birn, R.M., Smith, M.A., Jones, T.B., & Bandettini, P.A. (2008). The respiration response function: the temporal dynamics of fMRI signal fluctuations related to changes in respiration. *NeuroImage*, *40*(2), 644-654.

Blake, R. (1979). The visual system of the cat. *Perception and Psychophysics*, *26*(6), 423-448.

Bonhomme, V., Fiset, P., Meuret, P., Backman, S., Plourde, G., Paus, T., Bushnell, M.C., & Evans, A.C. (2000). Propofol anesthesia and cerebral blood flow changes elicited by vibrotactile

stimulation: a positron emission tomography study. *Journal of Neurophysiology*, *85*(3),

1299-1308.

Brown, T.A., Joanisse, M.F., Gati, J.S., Hughes, S.M., Nixon, P.L., Menon, R.S., & Lomber, S.G.

(2013). Characterization of the blood-oxygen level-dependent (BOLD) response in cat

auditory cortex using high-field fMRI. *NeuroImage*, *64*, 458-465.

Brown, T.A., Gati, J.S., Hughes, S.M., Nixon, P.L., Menon, R.S., & Lomber S.G. (2014). Functional

imaging of auditory cortex in adult cats using high-field fMRI. *Journal of Visualized*

Experiments, (84), e50872, doi:10.3791/50872.

Butler, B.E., Chabot, N., & Lomber, S.G. (2016). Quantifying and comparing the pattern of

thalamic and cortical projections to the posterior auditory field in hearing and deaf cats.

Journal of Comparative Neurology, *524*(15), 3042-3063.

Butler, B.E., Hall, A.J., & Lomber, S.G. (2015). High-field functional imaging of pitch processing in

auditory cortex of the cat. *PLoS One*, *10*(7), e0134362.

https://doi.org/10.1371/journal.pone.0134362.

Buxton, R.B. & Frank, L.R. (1997). A model for the coupling between cerebral blood flow and

oxygen metabolism during neural stimulation. *Journal of Cerebral Blood Flow and*

Metabolism, *17*(1), 64-72.

Crosby, G., Crane, A.M., & Sokoloff, L. (1982). Local changes in cerebral glucose utilization

during ketamine anesthesia. *Anesthesiology*, *56*(6), 437-443.

Deutsch, J., Jolliffe, C., Archer, E., & Leece, E.A. (2017). Intramuscular injection of alfaxalone in

combination with butorphanol for sedation in cats. *Veterinary Anaesthesia and*

Analgesia, *44*(4), 794-802.

Dueck, M.H., Petzke, F., Gerbershagen, H.J., Paul, M., Hebelmann, V., Girnus, R., … Boerner, U.

(2005). Propofol attenuates responses of the auditory cortex to acoustic stimulation in a

dose-dependent manner: a fMRI study. *Acta Anaesthesiologica Scandinavica*, *49*(6),

784-791.

Ferris, C.F., Febo, M., Luo, F., Schmidt, K., Brevard, M., Harder, J.A., … King, J.A. (2006).

Functional magnetic resonance imaging in conscious animals: a new tool in behavioral

neuroscience research. *Journal of Neuroendocrinology*, *18*(5), 307-318.

Freeman, J., & Ingvar, H. (1967). Effects of fentanyl on cerebral cortical blood flow and EEG in
the cat. *Acta Anaesthesiology Scandinavica*, *11*(4), 381-391.

Fukuda, M., Vazquez, A.L., Zong, X., & Kim, S.G. (2013). Effects of the alpha 2-adrenergic
receptor agonist dexmedetomidine on neural, vascular and BOLD fMRI responses in the
somatosensory cortex. *European Journal of Neuroscience*, *37*(1), 80-95.

Gao, Y.R., Ma, D., Zhang, Q., Winder, A.T., Liang, Z., Antinori, L., Drew, P.J., & Zhang, N. (2017).
Time to wake up: Studying neurovascular coupling and brain-wide circuit function in the
un-anesthetized animal. *NeuroImage*, *153*, 382-398.

Hall, A.J., Brown, T.A., Grahn, J.A., Gati, J.S., Nixon, P.L., Hughes, S.M., Menon, R.S., & Lomber,
S.G. (2014). There's more than one way to scan a cat: Imaging cat auditory cortex with
high-field fMRI using continuous or sparse sampling. *Journal of Neuroscience Methods*,
224, 96-106.

Harel, N., Lee, S.P., Nagaoka, T., Kim, D.S., & Kim, S.G. (2002). Origin of negative blood
oxygenation level-dependent MRI signals. *Journal of Cerebral Blood Flow and*
Metabolism, *22*(8), 908-917.

Heffner, R.S., & Heffner, H.E. (1988). Sound localization acuity in cat: Effect of azimuth, signal
duration, and test procedure. *Hearing Research*, *36*(2-3), 221-232.

Heil, P., & Irvine, D.R. (1998). Functional specialization in auditory cortex: responses to
frequency-modulated stimuli in cat's posterior auditory field. *Journal of*
Neurophysiology, *79*(6), 3041-3059.

Hodkinson, D.J., de Groote, C., McKie, S., William Deakin, J.F., & Williams, S.R. (2012).
Differential effects on anaesthesia on the phMRI response to acute ketamine challenge.
British Journal of Medicine and Medical Research, 2(3), 373-385.

Höflich, A., Hahn, A., Küblböck, M., Kranz, G.S., Vanicek, T., Ganger, S., Spies, M.,
Windischberger, C., Kasper, S., Winkler, D. & Lanzenberger, R (2017). Ketamine-
dependent neuronal activation in healthy volunteers. Brain Structure and Function,
222(3), 1533-1542.

Hubel, D.H., & Wiesel, T.N. (1962). Receptive fields, binocular interaction and functional
architecture in the cat's visual cortex. *Journal of Physiology*, *160*(1), 106-154.

Jenkinson, M., Bannister, P., Brady, M., & Smith, S. (2002). Improved optimization for the robust

and accurate linear registration and motion correction of brain images. *NeuroImage*,

17(2), 825-841.

Jezzard, P., Rauschecker, J.P., & Malonek, D. (1997). An in vivo model for functional MRI in cat

visual cortex. *Magnetic Resonance in Medicine*, *38*(5), 699-705.

Kaitin, K.I. (1985). Effects of thalidomide and pentobarbital on neuronal activity in the preoptic

area during sleep and wakefulness in the cat. *Psychopharmacology*, *85*(1), 47-50.

Kastrup, A., Liu, T.Q., Glover, G.H., & Moseley, M.E. (1999). Cerebral blood flow-related signal

changes during breath-holding. *American Journal of Neuroradiology*, *20*(7), 1233-1238.

Klassen, L.M., & Menon, R.S. (2004). Robust automated shimming technique using arbitrary

mapping acquisition parameters (RASTAMAP). *Magnetic Resonance in Medicine*, *51*(5),

881-887.

Lahti, K.M., Ferris, C.F., Li, F., Sotak, C.H., & King, J.A. (1999). Comparison of evoked cortical

activity in conscious and propofol-anesthetized rats using functional MRI. *Magnetic*

Resonance in Medicine, *41*(2), 412-416.

Lambert, J.J., Belelli, D., Peden, D.R., Vardy, A.W., & Peters, J.A. (2003). Neurosteroid

modulation of $GABA_A$ receptors. *Progress in Neurobiology*, *71*(1), 67-80.

Liu, Y., Qin, L., Zhang, X., Dong, C., & Sato, Y. (2010). Neural correlates of auditory temporal-

interval discrimination in cats. *Behavioural Brain Research*, *215*(1), 28-38.

Logothetis, N.K., Pauls, J., Augath, M., Trinath, T., & Oeltermann, A. (2001). Neurophysiological

investigation of the basis of the fMRI signal. *Nature*, *412*(6843), 150-157.

Lomber, S.G., MacNeil M.A., & Payne, B.R. (1995). Amplification of thalamic projections to

middle suprasylvian cortex following ablation of immature primary visual cortex in the

cat. *Cerebral Cortex*, *5*(2),166-191.

Miller, L.M., Escabi, M.A., Read, H.L., & Schreiner, C.E. (2002). Spectrotemporal receptive fields

in the lemniscal auditory thalamus and cortex. *Journal of Neurophysiology*, *87*(1), 516-

527.

Morin-Surun, M.P., Champagnat, J., Denavit-Saubie, M., & Moyanova, S. (1984). The effects of

acetylcholine on bulbar respiratory related neurons: Consequences of anaesthesia by

pentobarbital. *Archives of Pharmacology*, *325*(3), 205-208.

Muir, W., Lerche, P., Wiese, A., Nelson, L., Pasloske, K., & Whittem, T. (2009). The

cardiorespiratory and anesthetic effect of clinical and supraclinical doses of alfaxalone in

cats. *Veterinary Anaesthesia and Analgesia*, *36*(1), 42-54.

Musizza, B., & Ribaric, S. (2010). Monitoring the depth of anaesthesia. *Sensors (Basel,*

Switzerland), *10*(12), 10896-10935.

Myles, P.S. (2007). Prevention of awareness during anaesthesia. *Best Practice and Research:*

Clinical Anaesthesiology, *21*(3), 345-355.

Nagakubo, D., Hamamoto, Y., Hasegawa, D., Kamata, M., Iizuka, T., Muta, K., Fujita, N.,

Nakagawa, T. & Nishimura, R. (2017). Functional MRI-based identification of brain

regions activated by mechanical noxious stimulation and modulatory effect of

remifentanil in cats. Research in Veterinary Science, 114, 444-449.

Nagore, L., Soler, C., Gil, L., Serra, I., Soler, G., & Redondo, J.I. (2013). Sedative effects of

dexmedetomidine, dexmedetomidine-pethidine and dexmedetomidine-butorphanol in

cats. *Journal of Veterinary Pharmacology and Therapeutics*, *36*(3), 222-228.

Nakahiro, M., Yeh, J.Z., Brunner, E. & Narahashi, T. (1999). General anesthetics modulate GABA

receptor channel complex in rat dorsal root ganglion neurons. *Federation of American*

Societies for Experimental Biology, *3*(7), 1850-1854.

Ogawa, S., Lee, T.M., Kay, A.R., & Tank, D.W. (1990). Brain magnetic resonance imaging with

contrast dependent on blood oxygenation. Proceedings of the National Academy of

Sciences (USA), *87*(24), 9868-9872.

Olman, C., Ronen, I., Ugurbil, K., Kim, D.S. (2003). Retinotopic mapping in cat visual cortex using

high-field functional magnetic resonance imaging. *Journal of Neuroscience Methods*,

131(1-2), 161-170.

Oye, I., Paulsen, O., & Maurset, A. (1992). Effects of ketamine on sensory perception: evidence

for a role of N-methyl D-aspartate. *Journal of Pharmacological and Experimental*

Therapeutics, *260*(3), 1209-1213.

Paddleford, R.R. (1999). Manual of small animal anesthesia. Philadelphia, Pennsylvania: W.B.

Saunders Company.

Peirce, J.W. (2007). Psychophysics software in Python. *Journal of Neuroscience Methods*, *162*(1-

2), 8-13. doi:10.1016/j.jneumeth.2006.11.017.

Peirce J.W. (2009). Generating stimuli for neuroscience using PsychoPy. *Frontiers in*

Neuroinformatics, *2*(10), 1-8. doi:10.3389/neuro.11.010.2008.

Peng, Y.Z., Li, X.X., & Wang, Y.W. (2010). Effects of parecoxib and fentanyl on nociception-

induced cortical activity. *Molecular Pain*, *6*(3), doi: 10.1186/1744-8069-6-3.

Pienkowski, M., & Eggermont, J.J. (2009). Long-term, partially-reversible reorganization of

frequency tuning in mature cat primary auditory cortex can be induced by passive

exposure to moderate-level sounds. *Hearing Research*, *257*(1-2), 24-40.

Richards, C.D. (1983). Actions of general anaesthetics on synaptic transmission in the CNS.

British Journal of Anaesthesia, *55*(3), 201-207.

Ries, C.R., & Puil, E. (1999). Mechanism of anesthesia revealed by shunting actions of isoflurane

on thalamocortical neurons. *Journal of Neurophysiology*, *81*(4), 1795-1801.

Rorden, C., & Brett, M. (2000). Stereotaxic display of brain lesions. *Behavioral Neurology*, *12*(4),

191-200.

Ros, C., Soler, C., & Mateo, A.G.G.C. (2017). Comparison of brainstem auditory evoked

responses during sevoflurane or alfaxalone anaesthesia in adult cats. *Veterinary*

Anaesthesia and Analgesia, *44*(5), 1085-1090.

Smith, S.M., Jenkinson, M., Woolrich, M.W., Beckmann, C.F., Behrens, T.E.J., Johansen-Berg, H.,

… Matthews, P.M. (2004). Advances in functional and structural MR image analysis and

implementation as FSL. *NeuroImage*, *23*(S1), 208-219.

Stolzberg, D., Wong, C., Blake, B.E., & Lomber, S.G. (2017). Catlas: A magnetic resonance

imaging-based three-dimensional cortical atlas and tissue probability maps for the

domestic cat (Felis Catus). *Journal of Comparative Neurology*, *525*(15), 3190-3206.

Stolzberg, D., Blake, B.E., & Lomber, S.G. (2018). Effects of neonatal deafness on resting-state

functional network connectivity. *NeuroImage*, *165*, 69-82.

Taboada, F.M., & Murison, P.J. (2010). Induction of anesthesia with alfaxalone or propofol

before isoflurane maintenance in cats. *Veterinary Records*, *167*(3), 85-89.

Tamura, J., Ishizuka, T., Fukui, S., Oyama, N., Kawase, K., Itami, T., … Yamashita, K. (2015).

Sedative effects of intramuscular alfaxalone administered to cats. *Journal of Veterinary*

Medical Science, *77*(8), 874-904.

Thomas, J.A., & Lerche, P. (2011). *Anesthesia and analgesia for veterinary technicians*. St. Louis:

Elsevier.

Ueki, M., Mies, G., & Hossmann, K.A. (1992). Effects of alpha-chloralose, halothane,

pentobarbital and nitrous oxide anesthesia on metabolic coupling in somatosensory

cortex of rat. *Anaesthesiologica Scandinavica*, *36*(4), 318-322.

Warne, L.N., Beths, T., Whittem, T., Carter, J.E., & Bauquier, S.H. (2015). A review of the

pharmacology and clinical application of alfaxalone in cats. *The Veterinary Journal*, *203*,

141-148.

White, N.S., & Alkire, M.T. (2003). Impaired thalamocortical connectivity in humans during

general-anesthesia-induced unconsciousness. *NeuroImage*, *19*(2), 402-411.

Whittem, T., Pasloske, K.S., Heit, M.C., & Ransinghe, M.G. (2008). The pharmacokinetics and

pharmacodynamics of alfaxalone in cats after single and multiple intravenous

administration of Alfaxan at clinical and supraclinical doses. *Journal of Veterinary*

Pharmacology, *31*(6), 571-579.

Woolrich, M.W., Ripley, B.D., Brady, M., & Smith, S.M. (2001). Temporal autocorrelation in

univariate linear modeling of fMRI data. *NeuroImage, 14*(6) , 1370-1386.

Wong, C., Chabot, N., Kok, M.A., & Lomber, S.G. (2015). Amplified somatosensory and visual

cortical projections to a core auditory area, the anterior auditory field, following early-

and late-onset deafness: Modified projections to AAF following deafness. *Journal of*

Comparative Neurology, *523*(13), 1925-1947.

Wong, C., Pearson, K.G., & Lomber, S.G. (2018). Contributions of parietal cortex to the working

memory of an obstacle acquired visually or tactilely in the locomoting cat. *Cerebral*

Cortex, *28*(9), 3143-3158.

Worsley, K.J. Statistical analysis of activation images. (2001) In: Jezzard, P., Matthews, P.M., &

Smith, S.M. (2002*). Functional MRI: an introduction to methods*. Oxford: Oxford

University Press.

Wu, T.L., Mishra, A., Wang, F., Yang, P.F., Gore, J.C., & Chen, L.M. (2016). Effects of isoflurane

anesthesia on resting-state fMRI signals and functional connectivity within primary

somatosensory cortex of monkeys. *Brain and Behavior*, *6*(12), 2016.

Zurita, P., Villa, A.E.P., de Ribaupierre, Y., de Ribaupierre, F., Rouiller, E.M. (1994). Changes of

single unit activity in the cat auditory thalamus and cortex associated with different

anesthetic conditions. *Neuroscience Research*, *19*(3), 303-316.

683
684