Divergent density feedback control of migratory predator recovery following sex-biased

perturbations

Daisuke Goto[1,2*], Martin J. Hamel[3], Mark A. Pegg[3], Jeremy J. Hammen[4], Matthew L. Rugg[5], and

Valery E. Forbes[2,6]

[1]Institute of Marine Research/Havforskningsinstituttet, Postboks 1870, Nordnes, 5817 Bergen,

Norway

[2]School of Biological Sciences, Manter Hall, University of Nebraska-Lincoln, Lincoln,

Nebraska, 68588 USA

[3]School of Natural Resources, Hardin Hall, 3310 Holdrege Street, University of Nebraska-

Lincoln, Lincoln, Nebraska 68583-0973 USA

[4]U.S. Fish and Wildlife Conservation Office, 101 Park Deville Drive, Suite A, Columbia,

Missouri 65203 USA

[5]Montana Fish, Wildlife & Parks, 907 North Kendrick Avenue, Glendive, Montana 59330 USA

[6]Department of Ecology, Evolution and Behavior, University of Minnesota, 123 Snyder Hall,

1475 Gortner Ave, St. Paul, Minnesota 55108 USA

*Corresponding author Email: daisuke.goto2@gmail.com

Abstract

Uncertainty in risks posed by emerging stressors such as synthetic hormones impedes

conservation efforts for threatened vertebrate populations. Synthetic hormones often induce sex-

biased perturbations in exposed animals by disrupting gonad development and early life-history

stage transitions, potentially diminishing per capita reproductive output of depleted populations

and, in turn, being manifest as Allee effects. We use a spatially-explicit biophysical model to

evaluate how sex-biased perturbation in life history traits of individuals (maternal investment in

egg production and male-skewed sex allocation in offspring) modulates density feedback control

of year class strength and recovery trajectories of a long-lived, migratory fish–shovelnose

sturgeon (*Scaphirhynchus platorynchus*)–under spatially and temporally dynamic synthetic

androgen exposure and habitat conditions. Simulations show that reduced efficiency of maternal

investment in gonad development prolonged maturation time, increased the probability of

skipped spawning, and, in turn, gradually shrunk spawner abundance, weakening year class

strength. However, positive density feedback quickly disappeared (no Allee effect) once the

exposure ceased. By contrast, responses to the demographic perturbation manifested as strong

positive density feedback; an abrupt shift in year class strength and spawner abundance followed

after more than two decades owing to persistent negative population growth (a strong Allee

effect), reaching an alternate state without any sign of recovery. When combined with the

energetic perturbation, positive density feedback of the demographic perturbation was dampened

as extended maturation time reduced the frequency of producing male-biased offspring, allowing

the population to maintain positive growth rate (a weak Allee effect) and gradually recover. The

emergent patterns in long-term population projections illustrate that sex-biased perturbation in

life history traits of individuals can interactively regulate the strength of density feedback in

depleted populations such as *Scaphirhynchus* sturgeon to further diminish reproductive capacity

and abundance, posing increasingly greater conservation challenges in chemically altered

riverscapes.

KEYWORDS

1. INTRODUCTION

Human activities have uniquely shaped life history strategies of wild animal populations by imposing unnatural stress for centuries (Crain, Kroeker & Halpern 2008; Hendry, Farrugia & Kinnison 2008). The symptoms of human-induced stressors, in turn, manifest in life history traits varyingly among individuals (Marr *et al.* 2006) over time and space (Hope 2005; Maxwell *et al.* 2013). As stress becomes intensified, these sources of intraspecific variation in fitness may emerge as a directional shift in demographic rates (Wedekind 2017) as abundance declines. Positive density feedback following these demographic shifts may further diminish per capita reproductive capacity (a demographic Allee effect, Stephens, Sutherland & Freckleton 1999; Gregory *et al.* 2010), posing a threat to persistence of wild populations (Coltman *et al.* 2003).

One emerging stressor with such effects is synthetic chemicals, which have been increasingly detected in aquatic systems around the globe (Bernhardt, Rosi & Gessner 2017). A rising demand for human food production has increased agricultural use of synthetic chemicals (Tilman 1999; Tilman *et al.* 2001), posing a greater risk to freshwater vertebrates (Bernhardt, Rosi & Gessner 2017). Synthetic hormones–often used as growth-promoters for farmed animals–are especially potent and often induce sex-biased perturbation, impairing development and reproduction by disrupting energy allocation and maturation (del Carmen Alvarez & Fuiman 2005). Because maternal investment in egg production incurs greater energetic costs than somatic growth, females may more likely experience reproductive impairment than males (Arukwe & Goksøyr 1998). In pre-spawning females, even low-dose (often below detection) exposure to synthetic hormones can disrupt maturation processes such as vitellogenin production–a critical step in egg production (Arcand Hoy & Benson 1998; Nash *et al.* 2004), which can lead to skipped spawning in some fishes (Rideout, Rose & Burton 2005). Chemically

induced inefficiency in egg production may, thus, act density-dependently especially in

threatened and endangered populations as it reduces the number of females contributing to a

reproductive event (known as a 'mate-finding' Allee effect, Gascoigne *et al.* 2009).

Synthetic hormones can also irreversibly perturb sex determination in exposed animals,

skewing sex allocation in offspring (Mills & Chichester 2005). Resulting skewed adult sex ratio

may manifest as a mate-finding Allee effect (Gascoigne *et al.* 2009) by impairing reproduction

density-dependently (e.g., a decline in mating success), impeding recovery and posing extinction

risk for depleted populations as abundance declines further. Such human-mediated modification

in demographic traits has been reported in exploited populations with sex-biased mortality

(Molnar *et al.* 2007; Grüebler *et al.* 2008). However, it remains unclear how positive density-

dependent processes regulating fitness (also known as 'component' Allee effects, Stephens,

Sutherland & Freckleton 1999) manifested from modified traits such as skewed sex ratio and

delayed maturation act together to influence per capita reproductive output and population

recovery in nature.

We here present a simulation study on how sex-biased, density-dependent perturbation in life

history traits can emerge and modify recovery potential of a long-lived, migratory fish exposed

to low-level synthetic androgens in a large river. Migratory animals are exposed to spatially and

temporally dynamic flow-dependent exposure regimes as the chemicals move throughout

networks of streams and rivers (O'Brien *et al.* 2016); variable exposure and subsequent

symptoms among exposed individuals can obscure a causal link, if any, to population-level

responses, delaying management actions (Peterson *et al.* 2003). We test varying exposure and

effect scenarios on the Missouri River shovelnose sturgeon (*Scaphirhynchus platorynchus*, Fig.

1a) as an illustrative case study. Population sizes of *Scaphirhynchus* sturgeon have declined in

many of their endemic habitats, and some are facing potential extirpation owing to extensive

habitat modification (e.g., impoundments) (Bramblett & White 2001) that disrupt reproductive

activities including seasonal spawning migration and aggregation. Some of the Missouri River

populations have shown signs of unsuccessful spawning (DeLonay *et al.* 2009a) and year class

failure owing to low larval fish survival (Hrabik *et al.* 2007)–a critical process for population

persistence (DeLonay *et al.* 2009a). Because female *Scaphirhynchus* sturgeon mature later,

spawn less frequently, and live longer than males (DeLonay *et al.* 2009a), female-biased

disruption in reproductive traits likely regulate the strength of, if any, Allee effects. We use a

spatially-explicit individual-based biophysical model that accounts for sex-specific life history

traits to evaluate how sex-biased perturbation in energetics (reduced maternal investment in egg

production) and demographics (male-skewed sex allocation in offspring)–two commonly

reported symptoms of synthetic androgens–disrupts density-dependent reproductive processes to

control year class success and population recovery under spatially and temporally dynamic

exposure and habitat conditions.

2. MATERIALS AND METHODS

2.1. Study system

The study system is the lower 162 km of the Platte River (Nebraska, USA), one of the largest

tributaries of the Missouri River basin (Fig. 2 and Appendix 1: Fig. A1), providing spawning and

nursing habitats essential for threatened populations such as *Scaphirhynchus* sturgeon (Peters &

Parham 2008). Owing to its proximity to wastewater treatment plants and agricultural fields,

however, the lower Platte River is also one of the impaired water bodies in the region (NDEQ

2004). Fish that inhabit this river and its tributaries, such as *Scaphirhynchus* sturgeon, are

exposed to elevated concentrations of synthetic hormones widely used at concentrated cattle

feedlots near the lower Platte River watershed (Huntzinger & Ellis 1993) and exhibit symptoms

of endocrine disruption (e.g., intersexuality and atresia, DeLonay *et al.* 2009a).

2.2. Sturgeon biophysical model

We use a spatially explicit individual-based biophysical model of shovelnose sturgoen

previously developed for the lower Platte River popuatlion (Fig. 1b); the model is fully described

in Goto *et al.* (2015) and Goto *et al.* (2018). The following briefly describes the model according

to the Overview, Design concepts, and Details (ODD) protocol (Grimm *et al.* 2010).

Purpose. The model is designed to uncover mechanisms underlying early life-history processes

modified by natural and human-induced stressors (e.g., synthetic chemicals), and their

contributions to population persistence in a spatially and temporally dynamic environment.

Design concepts. 1) *Basic principles*: The model primarily hinges on energetics principles to

understand how maternal and environmental influences modulate year class variability through

energy acquisition and allocation (Houde 1989); 2) *Emergence*: Demographic traits and spatial

distribution of the population emerge from individual-level processes in response to habitat and

indirect intraspecific competition. 3) *Sensing*: Sturgeon sense gradients of habitat conditions and

prey density over neighboring cells; 4) *Interaction*: Sturgeon indirectly compete with each other

for food in the same cell (density dependence is incorporated in directly *Foraging* and indirectly

in *Growth,* Koch *et al.* 2009); 5) *Stochasticity*: Sturgeon's abilities to sense habitat conditions,

capture prey, decide when to spawn, and survive are stochastic because of inherent uncertainty

and randomness; and 6) *Observation*: The model stores sturgeon state variables daily to compute

demographic and life history trait values.

In this study, we evaluated sex-biased effects of synthetic androgen perturbation in maternal

investment in gonad development and first-year sex allocation. The following briefly describes

the essential features of the model relevant to this study.

Entities, state variables, and scales. The model simulates sturgeon population dynamics with

two types of entities; 1) river and 2) sturgeon. The model river consists of 162 equally divided

longitudinal rectangular segments (i.e., 1 river km each) with six state variables; daily, depth-

averaged discharge ($m^3 \cdot s^{-1}$), mean depth (gage height, m), channel width (m), benthic

invertebrate prey density ($g \cdot m^{-2}$), water temperature (°C), and photoperiod (hrs). Discharge,

depth, width, and prey density are spatially explicit, whereas temperature and photoperiod are

spatially uniform. The model keeps track of 11 state variables, body length (mm), storage mass

(g), structural mass (g), gonadal mass (g), the number of individuals in each super-individual,

maturity status, reproductive status, physiological condition (see *Growth*), sex, age (years), and

longitudinal location in the river (rkm), of the entire life cycle (Fig. 1b). Early life stages

(fertilized eggs, yolk-sac larvae, and feeding larvae, and age-0 juveniles) are represented as

super-individuals (Scheffer *et al.* 1995), whereas age 1+ juveniles and adults are represented as

individuals. Each super-individual is a collection of individuals ("cohorts") from the same

mother.

Process overview and scheduling. The model simulates the population and updates state

variables at discrete daily time steps over years and generations. The habitat conditions

(including prey production) are first updated, and then the model evaluates actions by sturgeon in

the following order; (i) *spawning,* (ii) *hatching and larval development,* (iii) *movement,* (iv)

foraging, (v) *growth*, and (vi) *mortality* (Fig. 1b). The model updates state variables at the end of

each action. At the beginning of each year, maturity is evaluated, and if mature sturgeon are

healthy (see *Growth* below), they enter the reproductive cycle. Spawning normally occurs from

late spring to early summer in the study system.

Submodels. The model consists of one benthic prey submodel and six sturgeon submodels. In

this study, simulations were designed to evaluate the effects of synthetic androgen perturbations

in vitellogenin production in adult females and sex allocation in offspring (see *Synthetic*

androgen effects below for detailed description) through the following processes in the

submodels:

a) Benthic prey production is computed using a mass-balance model of standing biomass,

daily production, and net migration from neighboring cells as described by Rashleigh &

Grossman (2005) with modifications described in Goto *et al.* (2018). Prey biomass density in

each cell is initialized on January 1st of each simulation year with a mean of 18.8 (wet g ·

m^{-2}) + 2.2RN, where RN is a random number drawn from the normal distribution (Whiles

and Goldowitz, 2005).

b) *Embryonic and yolk-sac larval development* is computed using cumulative fractional

temperature-dependent nonlinear functions (Goto *et al.* 2018) fitted to data reported by

Wang, Binkowski & Doroshov (1985). Hatched larvae are assumed to use their yolk sac for

~10 days (at 17–18 ℃) before exogenous feeding.

c) *Foraging* is computed using a modified version of the Beddington–DeAngelis functional

response model (Beddington 1975; DeAngelis, Goldstein & O'Neill 1975), a function of

encounter rate, handling time, size-dependent prey selectivity, and prey and predator

densities. The exact prey length is randomly assigned using a uniform distribution (between

minimum and maximum of each size class) each day (Goto *et al.* 2018).

d) *Growth* is simulated by the bioenergetics model (Hanson *et al.* 1997) modified to

represent the assimilated energy allocated to storage (reversible) and structure (irreversible)

(Gurney *et al.* 2003). When structure is less than optimal storage, feeding larvae and

juveniles allocate surplus net energy to storage first until optimal storage is reached (Goto *et*

al. 2018). Growth in length is then calculated using modified length–mass functions, where

length increases only when structural mass increases. When the assimilated energy is less

than the energy expended through metabolism, mass is lost only from storage without any

change in structure and length (Goto *et al.* 2018). Sturgeon mature based on sex-specific

probability, which is length-dependent functions derived from field observations of the Platte

River population (Goto *et al.* 2018). Reproductive adults further allocate energy to gonad

incrementally between spawning events. Adults initiate the reproductive cycle only when

their physiological condition is above a threshold based on relative storage mass, current

storage/optimal storage (Goto *et al.* 2018). Non-reproductive adults allocate energy only to

structure and storage. The amount of daily energy allocated to gonad ($J \cdot d^{-1}$) by healthy

adults (based on their physiological condition) is computed by dividing expected gonad

energy content on the spawning day by the minimal spawning interval (365 days for males

and 1095 days for females) plus the earliest expected spawning day (day of the year).

Expected gonad energy is computed by an allometric function derived by calibration for the

Platte River population. The amount of daily energy allocated to gonad is further adjusted

when the physiological condition is below the threshold (Goto *et al.* 2018). While the earliest

expected spawning day occurs in late March (when daylight hours >12 hours), the realized

spawning day is determined by discharge rate, temperature, and relative gonad mass.

e) *Spawning* is simulated as a stochastic event and occurs only when spawning cues–water

temperature, river discharge rate, daylight hours, and gonad development (gonadosomatic

index or GSI), concurrently meet the thresholds (Delonay *et al.* 2016). Female

Scaphirhynchus sturgeon spawn at highly variable river discharge rates among Lower

Missouri River tributaries (Delonay *et al.* 2016). Because our sturgeon model showed little

sensitivity to this parameter (Goto *et al.* 2015), a minimum threshold is set at 141.6 m$^3 \cdot$ s^{-1},

which is expected to reach roughly 50% connectivity (migratory pathway connectedness) in

the lower Platte River (Peters & Parham 2008) and directly influences the sturgeon

movement and subsequently habitat conditions for other individual-level actions. A

minimum GSI to initiate spawning is set at 0.18 for females and 0.03 for males based on field

observations (Wildhaber *et al.* 2007a). Water temperature suitable for female *Scaphirhynchus*

sturgeon spawning is set the range at 12.6–24.9℃ based on field observations (DeLonay *et*

al. 2009b). Fecundity is calculated by dividing gonad energy on the spawning day by single

egg energy, assuming constant egg quality (45.72 J $\cdot$ egg^{-1}, USDA 2001). After spawning,

fish lose all gonad mass, and their physiological condition and total mass are updated; fish do

not re-enter the reproductive cycle until the physiological condition meets the threshold.

When either temperature or discharge rate does not meet the thresholds, fish skip spawning.

We assume that when the physiological condition does not meet the threshold to remain in

the reproductive cycle, gonadal energy is reabsorbed and reallocated to storage.

f) *Movement* is tracked in the longitudinal direction only. The planktonic life stage fish (eggs

and yolk-sac larvae) move by passive drift with water currents (Braaten *et al.* 2008). Feeding

larvae (post-settlement fish, Kynard, Henyey & Horgan 2002), juveniles, and adults move in

the direction with probability of attaining higher habitat quality based on discharge rate

(Goto *et al.* 2018) using the empirical models developed for the lower Platte River

population (Peters & Parham 2008). Habitat suitability is based on relative areas of exposed

sandbars, open water, and shallow sandbar complexes. The new realized location is then

determined using fish swimming speed and water velocity (Goto *et al.* 2018); the maximum

distance that fish can travel daily in this study is thus limited within three (above or below

the current cell) neighboring cells (or 3.0 rkm).

g) *Mortality* results from thermal stress (eggs and yolk-sac larvae), predation (eggs, larvae,

and juveniles), starvation, and recreational fishing (age 3+). We assume that a constant

proportion of eggs and yolk-sac larvae die when water temperature rises above 24 °C

(Kappenman, Webb & Greenwood 2013). Starvation mortality depends on relative storage

mass; starvation mortality is evaluated only when a proportion of storage mass relative to

total mass drops below the threshold (optimal storage, Goto *et al.* 2018). Fishing mortality

was derived from annual creel surveys in the Platte River (Peters & Parham 2008). Realized

predation, starvation, and fishing mortality are a stochastic event. Realized mortality of eggs

and larvae is computed by drawing a random number from the binomial distribution with

mortality rates as the probability and the number of individuals represented by each SI_{yoy} as

the number of trials. For age 1+ fish, each fish dies when a random number drawn from a

uniform distribution is below the probability set above.

Input data. Model inputs include water temperature (°C), mean gauge height (depth, m),

channel width (m), discharge rate ($m^3 \cdot s^{-1}$), and photoperiod (hrs). Water temperature, gauge

height, channel width, and discharge rate (1995–2011, Appendix 1: Fig. A and Goto et al. 2018)

are derived from the USGS National Water Information System (http://waterdata.usgs.gov/nwis).

Further details on derivation of physical and hydrological input data are described in Goto *et al.*

(2018).

Initialization. We initialized simulations with 60 000 age 1+ individuals on January 1st. The

initial location of each sturgeon was randomly assigned to a cell with non-extreme water velocity

($\leq$0.6 m $\cdot$ s^{-1}) and depth ($\geq$0.5 m). The initial age structure follows a Poisson distribution with

mean age of 7, which is truncated at a maximum age of 16 based on field observations (Hamel *et*

al. 2015), and the initial sex ratio is 1:1. Sex is randomly assigned to each fish, assuming a

binomial distribution. For age 1+ sturgeon, initial lengths-at-age were computed using the von

Bertalanffy growth function (Ricker 1975) parameterized for the lower Platte River population

(Goto *et al.* 2018). Once spawned, fertilized eggs were initialized as super-individuals; the

number of super-individuals is the number of mothers, and the number of individuals in each

super-individual is the total number of eggs spawned by each mother. After settlement, feeding

larvae were initialized as an individual with an initial total length of 15.6 (mm) + 0.84RN

(Braaten *et al.* 2008), where RN is a random number drawn from the normal distribution; these

fish were subsequently simulated with individual-level processes (e.g., growth) and traits (e.g.,

body size).

2.3. Simulation scenarios for sex-biased purtubations by synthetic androgens

We ran 57-year (27-year exposure + 30-year recovery) simulations for the following four

scenarios to assess population performance under two types of perturbation in life history traits

of individuals induced by low-level synthetic androgen exposure, energetics and demorgraphics;

1) No sex-biased purtubation–baseline;

2) Synthetic androgens perturb maternal energetic investment in egg production by imposing

reduction in vitellogenin production rate;

3) Synthetic androgens perturb sex ratio in offspring by imposing male-skewed sex allocation

at recruitment; and

4) Synthetic androgens perturb both maternal investment and sex allocation. We assumed no

direct mortality caused by synthetic androgen exposure. Further, we assumed that: a) synthetic

androgens are released between May 1st and October 31st (a six-month growing season); b)

androgens affect only fish located below the confluence with the Elkhorn River (~51 rkm from

Plattsmouth, NE, Fig 2), where a number of cattle feedlots are located; and c) because the

androgen concentration varies with stream flow rate, the realized exposure level for individual

fish is spatially and temporally dynamic, and is thus adjusted based on flow rate in a grid cell in

which the given fish is located.

Energetic perturbation. We explicitly evaluated the mechanism underlying synthetic androgen-

induced reduction in egg production by simulating the dose-dependent reduction of vitellogenin

production in individual fish based on an exponential decay function (Ankley *et al.* 2003).

Vitellogenin (VTG) concentration (mg $\cdot$ ml^{-1}) is converted to relative gonad mass (GSI) using an

empirically derived linear function based on field survey data on the Missouri River shovelnose

sturgeon reported by Wildhaber *et al.* (2007b) (Appendix 1: Fig. A2) as GSI = 0.984VTG +

35.103 (R^2 = 0.99). Reduction in daily energy allocation to gonad (g $\cdot$ d^{-1}) in exposed females

was then computed based on somatic mass on the given day.

Because we lack information on a quantitative relationship between synthetic androgen

exposure and *Scaphirhynchus* sturgeon vitelogenin production, we tested the hypothetical

scenarios of 15%, 25%, and 35% reduction in daily vitellogenin production (Appendix 1: Fig.

A2) in fish that are located at the Elkhorn River–Platte River confluence; these reduction rates in

vitelogenin production approximately corresspond to the responses in fish exposed to

environmentally realistic, low level sysnthetic androgens (e.g., $<\sim15$ ng $\cdot$ l^{-1} 17β-trenbolone, a

synthetic androgen commonly used in cattle feedlots, Ankley *et al.* 2003). For fish that are

located in cells below the confluence, the reduction rates were proportionally adjusted by the

difference in flow rate (e.g., if flow rates at the confluence and in the cell where a fish is located

are 100 and 120 m^3 $\cdot$ s^{-1}, respectively, the realized reduction rate for this fish is 12.5% under the

15% reduction scenario). Because the model sturgeon cannot detect low-level synthetic

androgens in water, we assumed that their movement and energy acquisition (foraging) vary

independently of synthetic androgens.

Demographic perturbation. We implemented synthetic androgen-induced skewness in sex

allocation by proportionally reducing the probability of being female. We tested the hypothetical

scenarios of 20%, 60%, and 80% reduction in probability of being females for post-hatch fish

that remain in the cell at or below the Elkhorn River–Platte River confluence; these rates

approximately corresspond to the responses in fish exposed to environmentally realistic

concentrations (e.g., $<\sim15$ ng $\cdot$ l^{-1} 17β-trenbolone, Morthorst, Holbech & Bjerregaard 2010). We

assumed that the probability of being female incrementally declines by remaining in the

exposure area each day during the annual six-month exposure period (i.e., 0.11%, 0.33%, and

0.44%, respectively). For fish in cells below the confluence, the probability was proportionally

adjusted by the difference in flow rates (as described above).

For all the scenarios, we used hydrological and temperature inputs empirically derived from

the lower Platte River field surveys during 1995–2011 (Goto *et al.* 2018), from which we

randomly selected each simulation year; we assumed no systematic change or shift in

hydrological and thermal conditions in this study. Further, we assumed no interannual change in

turbidity, channel geomorphology, and photoperiod. We ran ten simulations for each exposure

318 scenario and computed summary statistics and % deviations from the baseline simulations in

319 demographic and life-history traits that contribute to per capita reproductive output; adult sex

320 ratio, relative reproductive female number (the proportion of adult females in the reproductive

321 cycle), reproductive female number, spawner biomass, and recruit number.

322 3. RESULTS

323 3.1. Baseline sturgeon population dynamics

324  Under the baseline conditions, the sturgeon population remained stable over 57 years (Fig. 3a)

325 with positive density dependence in per capita population growth rate (Fig. 3b). The population

326 maintained, on average, ~32% of adults being female (Appendix 1: Fig A3a), adult females

327 allocated, on average, ~9.4 % of total mass to gonad while they were in the reproductive cycle

328 (Appendix 1: Fig. A3b); ~22% of reproductive females (Appendix 1: Fig. A3c) eventually

329 spawned (= ~15.8 metric ton spawner biomass, Appendix 1: Fig A3d), producing, on average, 23

330 million eggs (or ~33000 eggs spawner^{-1}) annually (Appendix 1: Fig. A3e, and Table S1 and S2).

331 Most spawning and subsequent larval settlement occurred below the confluence of the Elkhorn

332 River and the Platte River (112 rkm) in late April to early May (Table S1 and S2); on average,

333 0.11% of first-year fish survived (Table S1 and S2) with ~7100 recruits per year (Appendix 1:

334 Fig. A3f). Although the recruit numbers varied highly among years, a non-linear (hump-shaped)

335 density-dependent relationship emerged between spawners and recruits (Appendix 1: Fig. A3g).

336 3.2. Recovery from reduced vitellogenin production efficiency

337  Simulated reduction in vitellogenin production rate lowered reproductive capacities of the

338 sturgeon population exposed to synthetic androgens (Fig. 4a). The proportion of energy allocated

339 to gonad mass (10%) or storage mass (24%) in reproductive females under exposure did not

340 differ from those under baseline (Table S1). However, the number of adult females in the

reproductive cycle declined by 7.5–34% during exposure (Fig. 4a). Further, spawner biomass

and recruit numbers declined, especially when exposed to moderate and high levels of the

perturbation (6.8–25% and 23–38%, respectively, Fig. 4a), which were linearly correlated with

relative changes in reproductive female abundance (Fig. 4b). Lowered reproductive capacities

and year class resulted in an 8.6–43% reduction (relative to baseline) in population biomass (Fig.

3a) with positive density dependence in growth rate (Fig. 3b) during exposure.

Reproductive females, spawners, and recruits of the population with moderate exposures

returned to the baseline levels (Fig. 4a); similar linear relationships among the reproductive traits

remained during recovery (Fig. 4b). Population size fully recovered (i.e., mean population size

returns to the baseline level) within ~five years (Fig. 3a) and density dependence in per capita

growth rate virtually disappeared (Fig. 3b). However, the reproductive females, spawners, and

recruits of the population with high exposure continued to decline (Fig. 4a), ultimately reducing

mean population biomass by up to 83% before showing a sign of recovery (~year 42, Fig. 3a).

3.3. Recovery from skewed recruit sex ratio

Sturgeon population biomass abruptly declined by up to 67% after ~year 22 (Fig. 3a), and per

capita growth rate continued to decline with biomass (Fig. 3b) under all scenarios of simulated

male-skewed sex allocation during the first year of life. Skewed sex ratios in recruits were

reflected in adults within ~6 years with relative female numbers (the number of adult females

entering the reproductive cycle) declining, on average, by 6.3–25% during exposure; mean

reproductive female abundance declined by 19–21% (Fig. 4c). Subsequently, mean spawner

biomass declined by up to 29% under high exposure (Fig. 4c). By contrast, mean recruit number

declined under low and moderate exposure (by 15% and 23%, respectively), but differed little

from the baseline population under high exposure (Fig. 4c). These changes in reproductive traits

were, however, nonlinearly covaried with sex ratio skewness (Fig. 4d).

During recovery, the moderate and high exposure population biomass reached a stable state

with less than 20% of the baseline population (Fig. 3a). Despite large year classes in early

recovery years (year 31–33), up to more than 85% of adults were males under high exposure

(Fig. 4c). On average, sex ratios under all the exposure scenarios remained skewed (by additional

9.3–16%) toward males with high interannual variation (Fig. 4c); sex ratio skewness did not

show exposure-dependence and covary nonlinearly with reproductive traits (Fig. 4d). Although

the low-exposure population fully recovered, the moderate and high exposure populations

continued to have declining reproductive female abundance, spawner biomass, and recruit

number (Fig. 4c), and, in turn, per capita growth rate continued to show positive density

dependence and remained, on average, negative (Fig. 3b).

3.4. Recovery from combined effects of reduced vitellogenin production and skewed sex ratio

Concurrently imposing reduction in vitellogenin production and skewed sex ratios modulated

the sturgeon population dynamics additively under low exposure, but antagonistically under

moderate and high exposure. Population biomass declined within ~5 years and remained stable

and then declined further in the last five years (Fig. 3a). During exposure, adult sex ratio differed

little under low exposure, but skewness toward males increased by 11% under medium and high

exposure (Fig. 5a). Further, reproductive female abundance, spawner biomass, and recruit

number declined by ~31%, ~23%, and ~31% (respectively) under low and moderate exposures,

and declined by 46%, 29%, and 47% (respectively) under high exposure (Fig. 5a), resulting in

positive density dependence in per capita growth rate (Fig. 3b). Reproductive output (recruit

number and spawner biomass) covaried nonlinearly with changes in relative reproductive

number because of amplified responses under low exposure (Fig. 5b). By contrast, the

relationships between reproductive traits (reproductive female number, spawner biomass, and

recruit number) and adult sex ratio skewness abruptly shifted under high exposure (Fig. 5c).

Population biomass under low and moderate exposures recovered within ~25 years, whereas

population biomass under high exposure reached a stable state in ~22 years at ~44% lower than

the baseline population (Fig. 3a). Although per capita growth rate continued to show positive

density dependence, it remained, on average, positive during recovery (Fig. 3b). Adult sex ratios

remained skewed toward males under all the exposure scenarios (Fig. 5a). Mean reproductive

female number, spawner biomass, and recruit number (except for reproductive females under

low exposure) recovered after low and moderate exposures but remained low under high

exposure (Fig. 5a). The reproductive output covaried linearly with changes in relative adult

female number (Fig. 5b); by contrast, the dampened responses in the reproductive traits during

exposure also led to full recovery under medium as well as low (except for reproductive female

number) exposures (Fig. 5b,c).

4. DISCUSSION

Simulations revealed plausible mechanisms involving chemically-induced, sex-biased

perturbations in life history traits of long-lived, migratory vertebrates such as *Scaphirhynchus*

sturgeon; these perturbations can emerge as vastly different response and recovery patterns–

being manifest as differed strengths of demographic Allee effects (Hutchings 2015)–in a

spatially and temporally dynamic environment. We find that lowered vitellogenin production rate

prolongs maturation period and incurs skipped spawning (a relative decline in reproductive

female abundance), and, in turn, reduces spawner abundance, ultimately weakening year class

strength. Positive density feedback in per capita growth rate, however, disappeared (no Allee

effect) during recovery. Further, the energetic perturbation operates as exposure-dependent

processes–the higher the exposure level, the more severe the damage, thereby resulting in longer

recovery time. By contrast, male-skewed sex allocation (an absolute decline in reproductive

female abundance) emerges as delayed (by more than two decades) population-level responses,

followed by an abrupt shift in year class strength and population trajectory under higher

exposures, ultimately reaching an alternative state without any sign of recovery owing to

persistent negative population growth (a strong Allee effect, Stephens, Sutherland & Freckleton

1999; Gascoigne *et al.* 2009). When combined with the energetic perturbation, however, impacts

of the demographic perturbation are dampened as extended maturation time reduces the

frequency of producing male-biased offspring, allowing the population to maintain positive

growth rate (a weak Allee effect) and gradually recover. The emergent patterns in long-term

population projections illustrate that sex-biased perturbation in life history traits of individuals

can amplify or dampen Allee effects in depleted populations such as *Scaphirhynchus* sturgeon to

further diminish reproductive capacity and abundance, posing conservation challenges in

chemically altered riverscapes.

Sex-biased differences in energy allocation to gonadal maturation often determine life-history

strategies such as foraging and mate selection, shaping sexual dimorphism in animal populations

(Huse 1998; Rennie *et al.* 2008). Delayed gonad maturation in energetically perturbed females of

our model sturgeon population prolongs the spawning cycle instead of reducing fecundity.

Because sturgeon spend several years to build gonad energy prior to spawning (a capital

spawner, McBride *et al.* 2015), slowed female gonad maturation and longer spawning cycle

delay spawning timing. Although sturgeon are iteroparous, because of relatively infrequent

spawning and long maturation time, delayed or skipped spawning events can reduce individual

lifetime (in our simulations, some individuals spawned only once) and per capita annual

reproductive output owing to fewer females contributing to spawning events (a reproductive

Allee effect, Gascoigne *et al.* 2009). Inefficient energy use triggered by synthetic hormones in

threatened animal populations with long maturation time can, thus, prolong population recovery

time. Our simulations, however, suggest that the energetic perturbation also triggers plastic

responses. Although reproductive and spawning females decline by up to 25%, the population

eventually recovers partly because of limited positive density feedback in per capita growth rate

following synthetic androgen exposure, indicating that reproductive capacity diminished during

exposure can be restored if energy allocation to gonad resumes.

Delayed spawning events may, however, increase the probability of year class failures owing

to environmental stochasticity. We find that exposed (energetically challenged) female sturgeon

are more likely to abort the reproductive cycle, as they divert energy to meet metabolic needs

when poor habitat conditions persist. Although evaluating energetic tradeoffs between migration

and fecundity is beyond the scope of our study, animals with extensive spawning migration such

as *Scaphirhynchus* sturgeon likely skip spawning or produce offspring with lower survival rate

owing to reduced energy reserves (Glebe & Leggett 1981). Sufficient energy reserves in eggs are

vital to larval development and survival; maternal investment in egg production determines the

nutritional status of post-hatching larvae (Kjesbu *et al.* 1996). Our model assumes that individual

eggs do not vary in energy content with synthetic androgen exposure; however, energetic

tradeoffs in maternal investment in egg production and offspring survival can arise and vary

greatly among taxa (Jönsson 1997; McBride *et al.* 2015) and thus need further investigation for

Scaphirhynchus sturgeon.

Our study shows that recovery from demographic perturbation can pose greater conservation

challenges than energetic perturbation. We find that chemically-induced, male-skewed adult sex

ratio can incur a lasting mate-finding Allee effect (Gascoigne *et al.* 2009) in long-lived

populations such as *Scaphirhynchus* sturgeon, severely diminishing their reproductive fitness and

recovery potential. Although skewed sex ratio has been widely reported for a variety of

vertebrate taxa (Donald 2007), causes of the skewness vary; it can result from sex-specific

phenotypic differences in physiology, dispersal, and natural or human-induced mortality

(Weimerskirch, Lallemand & Martin 2005; Goto 2009; Goto & Wallace 2010; Pen *et al.* 2010).

In our baseline population, adult sex ratios are skewed toward males (~1:2.2), primarily resulting

from 1) a longer maturation time for females than males and 2) more energy investment to

reproduction in females, resulting in higher probability of starvation mortality in poor

environment years. Simulations show that further deviation (by 13–23%) in adult sex ratio

skewness result in long-lasting threats to year class success and population viability.

Consequences of skewed adult sex ratio may not immediately emerge for long-lived species

(Holmes & York 2003; Lawson *et al.* 2010) such as *Scaphirhynchus* sturgeon. Once the

skewness reaches a tipping point, however, it often drastically reduces recovery potential

(Gascoigne *et al.* 2009) as it may stabilize in an alternative state (low abundance) facilitated by

positive feedback (Van Nes & Scheffer 2007; Takimoto 2009; Dai *et al.* 2012). Although

spawning aggregation formed by the Missouri River populations of *Scaphirhynchus* sturgeon

may, to some extent, mitigate the component (mate-finding) Allee effect (Stephens, Sutherland

& Freckleton 1999; Gascoigne *et al.* 2009), the threshold of this effect (female shortage) to avoid

demographic shifts and facilitate recovery of these populations needs further investigation.

Finally, our study demonstrates that two or more component Allee effects can act interactively

to dampen (or amplify) density feedback in per capita reproductive output and promote (or

prevent) recovery in masculinized populations. Simulations show that by reallocating gonad

energy to growth and postponing spawning events, surviving adult female sturgeon mitigate the

strong Allee effect of skewed sex ratio by reducing the overall frequency of producing male-

skewed offspring during exposure. In conservation planning for long-lived migratory species

such as *Scaphirhynchus* sturgeon, thus, to prevent the shifts and minimize recovery cost, it is

critical to identify early warnings that signal impending demographic shifts; the threshold of

skewness in adult sex ratio that leads to demographic Allee effects can be a sensitive metric for

assessing viability of threatened populations (Gascoigne *et al.* 2009; Ancona *et al.* 2017).

Long-term, low-level exposure to stressors such as synthetic hormones often obscures

causative relationships (Windsor, Ormerod & Tyler 2018). Establishing ecologically relevant

indicators such as the thresholds of Allee effects (where the shift occurs from positive to

negative per capita growth) resulting from modified life history traits would guide the risk

assessment of synthetic chemicals and habitat conservation planning for threatened populations.

Further, given the time lags across biological hierarchy, continued biological monitoring of

individual-level traits would help detect early signals that may indicate population-level changes

such as the strength of Allee effects. Although our study was designed to evaluate consequences

of low-level exposure, our findings also indicate larger implications of among-individual

variation for population persistence and recovery of threatened and endangered species.

Population models widely used in risk assessments often assume no among-individual variation

in a population such as sex-, size-, or age- dependent life history traits (e.g., age at maturity).

These models may not only gloss over individual variation in life history traits that may

contribute to population viability, but also under- or over- estimate the strength of density

feedback, and, in turn, the probability of extinction risk.

Large-river migratory populations such as *Scaphirhynchus* sturgeon experience multiple

synthetic chemicals over agriculturally rich landscapes in regions such as the Midwestern United

States (Welcomme 1995). Further confounding factors in conservation and management efforts

can arise in hydrologically altered systems such as the Missouri River. Precipitation, for

example, can regulate the amount of agrichemicals released into rivers. Assessing the likelihood

of ecological risks of agrichemicals for long-lived species requires the development of trait-

based predictive models using anticipated scenarios. With expected trends in rising climate

variability, which can have considerable effects on river hydrology and water chemistry, these

forecasting tools are critical for fisheries and water resource management in chemically altered

ecosystems.

AUTHORS' CONTRIBUTIONS

DG, VEF, and MAP conceived the ideas; MJH, JJH and MLR collected and managed the data;

DG led the writing of the manuscript. All authors contributed to revisions and gave final

approval for publication.

ACKNOWLEDGEMENTS

DATA AVAILABILITY STATEMENT

Model input data on water temperature, gauge height, channel width, and discharge rate used are

publicly available at the United States Geological Survey (USGS) National Water Information

System (http://waterdata.usgs.gov/nwis).

REFERENCES

Ancona, S., Dénes, F.V., Krüger, O., Székely, T. & Beissinger, S.R. (2017) Estimating adult sex

ratios in nature. *Philosophical Transactions of the Royal Society B: Biological Sciences,* **372,**

20160313.

Ankley, G.T., Jensen, K.M., Makynen, E.A., Kahl, M.D., Korte, J.J., Hornung, M.W., Henry,

T.R., Denny, J.S., Leino, R.L. & Wilson, V.S. (2003) Effects of the androgenic growth

promoter 17-β-trenbolone on fecundity and reproductive endocrinology of the fathead

minnow. *Environmental Toxicology and Chemistry,* **22,** 1350-1360.

Arcand Hoy, L.D. & Benson, W.H. (1998) Fish reproduction: an ecologically relevant indicator

of endocrine disruption. *Environmental Toxicology and Chemistry,* **17,** 49-57.

Arukwe, A. & Goksøyr, A. (1998) Xenobiotics, xenoestrogens and reproduction disturbances in

fish. *Sarsia,* **83,** 225-241.

Beddington, J. (1975) Mutual interference between parasites or predators and its effect on

searching efficiency. *Journal of Animal Ecology***,** 331-340.

Bernhardt, E.S., Rosi, E.J. & Gessner, M.O. (2017) Synthetic chemicals as agents of global

change. *Frontiers in Ecology and the Environment,* **15,** 84-90.

Braaten, P.J., Fuller, D.B., Holte, L.D., Lott, R.D., Viste, W., Brandt, T.F. & Legare, R.G. (2008)

Drift dynamics of larval pallid sturgeon and shovelnose sturgeon in a natural side channel of

the upper Missouri River, Montana. *North American Journal of Fisheries Management,* **28,**

808-826.

Bramblett, R.G. & White, R.G. (2001) Habitat use and movements of pallid and shovelnose

sturgeon in the Yellowstone and Missouri rivers in Montana and North Dakota. *Transactions*

of the American Fisheries Society, **130,** 1006-1025.

Coltman, D.W., O'Donoghue, P., Jorgenson, J.T., Hogg, J.T., Strobeck, C. & Festa-Bianchet, M.

(2003) Undesirable evolutionary consequences of trophy hunting. *Nature,* **426,** 655-658.

Crain, C.M., Kroeker, K. & Halpern, B.S. (2008) Interactive and cumulative effects of multiple

human stressors in marine systems. *Ecology Letters,* **11,** 1304-1315.

Dai, L., Vorselen, D., Korolev, K.S. & Gore, J. (2012) Generic indicators for loss of resilience

before a tipping point leading to population collapse. *science,* **336,** 1175-1177.

DeAngelis, D., Goldstein, R. & O'Neill, R. (1975) A model for tropic interaction. *Ecology*, 881-

892.

del Carmen Alvarez, M. & Fuiman, L.A. (2005) Environmental levels of atrazine and its

degradation products impair survival skills and growth of red drum larvae. *Aquatic*

Toxicology, **74,** 229-241.

Delonay, A.J., Chojnacki, K.A., Jacobson, R.B., Albers, J.L., Braaten, P.J., Bulliner, E.A.,

Elliott, C.M., Erwin, S.O., Fuller, D.B., Haas, J.D., Ladd, H.L.A., Mestl, G.E., Papoulias,

D.M. & Wildhaber, M.L. (2016) Ecological requirements for pallid sturgeon reproduction and

recruitment in the Missouri River—A synthesis of science, 2005 to 2012. *Scientific*

Investigations Report, pp. 242. Reston, VA.

DeLonay, A.J., Jacobson, R.B., Papoulias, D.M., Simpkins, D.G., Wildhaber, M.L., Reuter, J.M.,

Bonnot, T.W., Chojnacki, K.A., Korschgen, C.E. & Mestl, G.E. (2009a) Ecological

requirements for pallid sturgeon reproduction and recruitment in the Lower Missouri River: a

research synthesis 2005-08.

DeLonay, A.J., Jacobson, R.B., Papoulias, D.M., Simpkins, D.G., Wildhaber, M.L., Reuter, J.M.,

Bonnot, T.W., Chojnacki, K.A., Korschgen, C.E., Mestl, G.E. & Mac, M.J. (2009b)

Ecological requirements for pallid sturgeon reproduction and recruitment in the Lower

Missouri River: A research synthesis 2005–08. U.S. Geological Survey Scientific

Investigations Report 2009–5201.

Donald, P.F. (2007) Adult sex ratios in wild bird populations. *Ibis,* **149,** 671-692.

Gascoigne, J., Berec, L., Gregory, S. & Courchamp, F. (2009) Dangerously few liaisons: a

review of mate-finding Allee effects. *Population Ecology,* **51,** 355-372.

Glebe, B. & Leggett, W. (1981) Temporal, intra-population differences in energy allocation and

use by American shad (*Alosa sapidissima*) during the spawning migration. *Canadian Journal*

of Fisheries and Aquatic Sciences, **38,** 795-805.

Goto, D. (2009) Impacts of habitat degradation on *Fundulus heteroclitus* (Linnaeus) in urban

tidal salt marshes in New York. Ph.D. Dissertation, City University of New York, New York

NY.

Goto, D., Hamel, M.J., Hammen, J.J., Rugg, M.L., Pegg, M.A. & Forbes, V.E. (2015)

Spatiotemporal variation in flow-dependent recruitment of long-lived riverine fish: Model

development and evaluation. *Ecological Modelling,* **296,** 79-92.

Goto, D., Hamel, M.J., Pegg, M.A., Hammen, J.J., Rugg, M.L. & Forbes, V.E. (2018) Spatially

dynamic maternal control of migratory fish recruitment pulses triggered by shifting seasonal

cues. *Marine and Freshwater Research,* **69,** 942-961.

Goto, D. & Wallace, W.G. (2010) Bioenergetic responses of a benthic forage fish (Fundulus

heteroclitus) to habitat degradation and altered prey community in polluted salt marshes.

Canadian Journal of Fisheries and Aquatic Sciences, **67,** 1566-1584.

Gregory, S.D., Bradshaw, C.J., Brook, B.W. & Courchamp, F. (2010) Limited evidence for the

demographic Allee effect from numerous species across taxa. *Ecology,* **91,** 2151-2161.

Grimm, V., Berger, U., DeAngelis, D.L., Polhill, J.G., Giske, J. & Railsback, S.F. (2010) The

ODD protocol: a review and first update. *Ecological Modelling,* **221,** 2760-2768.

Grüebler, M.U., Schuler, H., Müller, M., Spaar, R., Horch, P. & Naef-Daenzer, B. (2008) Female

biased mortality caused by anthropogenic nest loss contributes to population decline and adult

sex ratio of a meadow bird. *Biological Conservation,* **141,** 3040-3049.

Gurney, W.S.C., Jones, W., Veitch, A.R. & Nisbet, R.M. (2003) Resource allocation,

hyperphagia, and compensatory growth in juveniles. *Ecology,* **84,** 2777-2787.

Hanson, P.C., Johnson, T.B., Schindler, D.E. & Kitchell, J.F. (1997) Fish bioenergetics 3.0.

University of Wisconsin Sea Grant Institute, Madison, WI.

Hendry, A.P., Farrugia, T.J. & Kinnison, M.T. (2008) Human influences on rates of phenotypic

change in wild animal populations. *Molecular Ecology,* **17,** 20-29.

Holmes, E. & York, A. (2003) Using age structure to detect impacts on threatened populations: a

case study with Steller sea lions. *Conservation Biology,* **17,** 1794-1806.

Hope, B.K. (2005) Performing spatially and temporally explicit ecological exposure assessments

involving multiple stressors. *Human and Ecological Risk Assessment,* **11,** 539-565.

Houde, E.D. (1989) Comparative growth, mortality, and energetics of marine fish larvae:

temperature and implied latitudinal effects. *Fishery Bulletin,* **87,** 471-495.

Hrabik, R., Herzog, D., Ostendorf, D. & Petersen, M. (2007) Larvae provide first evidence of

successful reproduction by pallid sturgeon, Scaphirhynchus albus, in the Mississippi River.

Journal of Applied Ichthyology, **23,** 436-443.

Huntzinger, T.L. & Ellis, M.J. (1993) Central Nebraska river basins, Nebraska. *JAWRA Journal*

of the American Water Resources Association, 29, 533-574.

Huse, G. (1998) Sex-specific life history strategies in capelin (*Mallotus villosus*)? *Canadian*

Journal of Fisheries and Aquatic Sciences, 55, 631-638.

Hutchings, J.A. (2015) Thresholds for impaired species recovery. *Proceedings of the Royal*

Society B: Biological Sciences, 282, 20150654.

Jönsson, K.I. (1997) Capital and income breeding as alternative tactics of resource use in

reproduction. *Oikos*, 57-66.

Kappenman, K.M., Webb, M.A.H. & Greenwood, M. (2013) The effect of temperature on

embryo survival and development in pallid sturgeon *Scaphirhynchus albus* (Forbes &

Richardson 1905) and shovelnose sturgeon *S. platorynchus* (Rafinesque, 1820). *Journal of*

Applied Ichthyology, 29, 1193-1203.

Kjesbu, O.S., Solemdal, P., Bratland, P. & Fonn, M. (1996) Variation in annual egg production

in individual captive Atlantic cod (*Gadus morhua*). *Canadian Journal of Fisheries and*

Aquatic Sciences, 53, 610-620.

Kynard, B., Henyey, E. & Horgan, M. (2002) Ontogenetic behavior, migration, and social

behavior of pallid sturgeon, Scaphirhynchus albus, and shovelnose sturgeon, S. platorynchus,

with notes on the adaptive significance of body color. *Environmental Biology of Fishes, 63,*

389-403.

Lawson, D.M., Regan, H.M., Zedler, P.H. & Franklin, J. (2010) Cumulative effects of land use,

altered fire regime and climate change on persistence of Ceanothus verrucosus, a rare, fire-

dependent plant species. *Global Change Biology, 16,* 2518-2529.

Marr, A., Arcese, P., Hochachka, W., Reid, J.M. & Keller, L. (2006) Interactive effects of

environmental stress and inbreeding on reproductive traits in a wild bird population. *Journal*

of Animal Ecology, **75,** 1406-1415.

Maxwell, S.M., Hazen, E.L., Bograd, S.J., Halpern, B.S., Breed, G.A., Nickel, B., Teutschel,

N.M., Crowder, L.B., Benson, S. & Dutton, P.H. (2013) Cumulative human impacts on

marine predators. *Nature Communications,* **4,** 2688.

McBride, R.S., Somarakis, S., Fitzhugh, G.R., Albert, A., Yaragina, N.A., Wuenschel, M.J.,

Alonso-Fernández, A. & Basilone, G. (2015) Energy acquisition and allocation to egg

production in relation to fish reproductive strategies. *Fish and Fisheries,* **16,** 23-57.

Mills, L.J. & Chichester, C. (2005) Review of evidence: Are endocrine-disrupting chemicals in

the aquatic environment impacting fish populations? *Science of the Total Environment,* **343,**

1-34.

Molnar, P.K., Derocher, A.E., Lewis, M.A. & Taylor, M.K. (2007) Modelling the mating system

of polar bears: a mechanistic approach to the Allee effect. *Proceedings of the Royal Society B:*

Biological Sciences, **275,** 217-226.

Morthorst, J.E., Holbech, H. & Bjerregaard, P. (2010) Trenbolone causes irreversible

masculinization of zebrafish at environmentally relevant concentrations. *Aquatic Toxicology,*

98, 336-343.

Nash, J.P., Kime, D.E., Van der Ven, L.T.M., Wester, P.W., Brion, F., Maack, G., Stahlschmidt-

Allner, P. & Tyler, C.R. (2004) Long-term exposure to environmental concentrations of the

pharmaceutical ethynylestradiol causes reproductive failure in fish. *Environmental Health*

Perspectives, **112,** 1725.

NDEQ (2004) 2004 surface water quality integrated report. Water Quality Division. pp. 118 pp.

O'Brien, D., Lewis, S., Davis, A., Gallen, C., Smith, R., Turner, R., Warne, M., Turner, S.,

Caswell, S. & Mueller, J.F. (2016) Spatial and temporal variability in pesticide exposure

downstream of a heavily irrigated cropping area: application of different monitoring

techniques. *Journal of Agricultural and Food Chemistry,* **64,** 3975-3989.

Pen, I., Uller, T., Feldmeyer, B., Harts, A., While, G.M. & Wapstra, E. (2010) Climate-driven

population divergence in sex-determining systems. *Nature,* **468,** 436.

Peters, E.J. & Parham, J.E. (2008) Ecology and management of sturgeon in the lower Platte

River, Nebraska. *Nebraska Technical Series.*

Peterson, C.H., Rice, S.D., Short, J.W., Esler, D., Bodkin, J.L., Ballachey, B.E. & Irons, D.B.

(2003) Long-term ecosystem response to the Exxon Valdez oil spill. *Science,* **302,** 2082.

Rashleigh, B. & Grossman, G.D. (2005) An individual-based simulation model for mottled

sculpin (Cottus bairdi) in a southern Appalachian stream. *Ecological Modelling,* **187,** 247-

258.

Rennie, M.D., Purchase, C.F., Lester, N., Collins, N.C., Shuter, B.J. & Abrams, P.A. (2008)

Lazy males? Bioenergetic differences in energy acquisition and metabolism help to explain

sexual size dimorphism in percids. *Journal of Animal Ecology,* **77,** 916-926.

Ricker, W.E. (1975) *Computation and interpretation of biological statistics for fish populations.*

Fisheries Research Board of Canada, Ottawa, Ontario.

Rideout, R.M., Rose, G.A. & Burton, M.P. (2005) Skipped spawning in female iteroparous

fishes. *Fish and Fisheries,* **6,** 50-72.

Scheffer, M., Baveco, J., DeAngelis, D., Rose, K. & Van Nes, E. (1995) Super-individuals a

simple solution for modelling large populations on an individual basis. *Ecological Modelling,*

80, 161-170.

Stephens, P.A., Sutherland, W.J. & Freckleton, R.P. (1999) What is the Allee effect? *Oikos*, 185-

190.

Takimoto, G. (2009) Early warning signals of demographic regime shifts in invading

populations. *Population Ecology,* **51,** 419-426.

Tilman, D. (1999) Global environmental impacts of agricultural expansion: the need for

sustainable and efficient practices. *Proceedings of the National Academy of Sciences,* **96,**

5995-6000.

Tilman, D., Fargione, J., Wolff, B., D'antonio, C., Dobson, A., Howarth, R., Schindler, D.,

Schlesinger, W.H., Simberloff, D. & Swackhamer, D. (2001) Forecasting agriculturally driven

global environmental change. *Science,* **292,** 281-284.

USDA (2001) USDA (U.S. Department of Agriculture, Agricultural Research) Nutrient

Database for Standard Reference. http://www.nal.usda.gov/fnic/foodcomp.

Van Nes, E.H. & Scheffer, M. (2007) Slow recovery from perturbations as a generic indicator of

a nearby catastrophic shift. *The American Naturalist,* **169,** 738-747.

Wang, Y.L., Binkowski, F.P. & Doroshov, S.I. (1985) Effect of temperature on early

development of white and lake sturgeon, *Acipenser transmontanus* and *A. fulvescens*.

Environmental Biology of Fishes, **14,** 43-50.

Wedekind, C. (2017) Demographic and genetic consequences of disturbed sex determination.

Philosophical Transactions of the Royal Society B: Biological Sciences, **372,** 20160326.

Weimerskirch, H., Lallemand, J. & Martin, J. (2005) Population sex ratio variation in a

monogamous long-lived bird, the wandering albatross. *Journal of Animal Ecology,* **74,** 285-

291.

Welcomme, R.L. (1995) Relationships between fisheries and the integrity of river systems.

Regulated Rivers: Research & Management, **11,** 121-136.

Wildhaber, M., Papoulias, D., DeLonay, A., Tillitt, D., Bryan, J. & Annis, M. (2007a) Physical

and hormonal examination of Missouri River shovelnose sturgeon reproductive stage: a

reference guide. *Journal of Applied Ichthyology,* **23,** 382-401.

Wildhaber, M.L., Papoulias, D.M., DeLonay, A.J., Tillitt, D.E., Bryan, J.L. & Annis, M.L.

(2007b) Physical and hormonal examination of Missouri River shovelnose sturgeon

reproductive stage: a reference guide. *Journal of Applied Ichthyology,* **23,** 382-401.

Windsor, F.M., Ormerod, S.J. & Tyler, C.R. (2018) Endocrine disruption in aquatic systems: up-

scaling research to address ecological consequences. *Biological Reviews,* **93,** 626-641.

Figure legends

Figure 1. Shovelnose sturgeon *Scaphirhynchus platorynchus*. a) adult shovelnose sturgeon; b) a schematic of the spatially explicit individual-based biophysical model used in this study.

Figure 2. Map of the study system, lower Platte River, Nebraska, USA.

Figure 3. Population dynamics and density dependence in population growth of shovelnose sturgeon population. a) population projections during exposure and recovery under four synthetic androgen scenarios: a baseline scenario; an energetic perturbation (reduced maternal investment in gonad development) scenario with three exposure levels (shown as biomass relative to the baseline); a demographic perturbation (male-biased sex allocation in offspring) scenario; and a combined scenario of energetic and demographic perturbations. b) relationships between population biomass and per capita population growth rate, $\ln(N_{t+1}/N_t)$, during exposure (baseline: $y = 0.32x - 1.3$, $r = 0.26$; energetic perturbation: $y = 0.24x - 1.0$, $r = 0.26$; demographic perturbation: $y = 0.29x - 1.2$, $r = 0.28$; and combined: $y = 0.20x - 0.84$, $r = 0.25$) and recovery (baseline: $y = 0.49x - 2.0$, $r = 0.43$; energetic perturbation: $y = 0.0098x - 0.031$ $r = 0.023$; demographic perturbation: $y = 0.083x - 0.33$, $r = 0.13$; and combined: $y = 0.072x - 0.25$, $r = 0.11$).

Figure 4. Deviation (%) compared to the baseline scenario in life history traits of shovelnose sturgeon populations. a) proportion of mature females in the reproductive cycle, recruit number, reproductive female number, and spawner biomass under energetic perturbation scenarios; b) relationship between relative reproductive female number and spawner biomass or recruit number during exposure and exposure under energetic perturbation scenarios; c) adult sex ratio, recruit number, reproductive female number, and spawner biomass under demographic perturbation scenarios, d) relationship between adult sex ratio and reproductive female number,

spawner biomass, or recruit number during exposure and recovery under demographic perturbation scenarios. In a and c, black circles indicate annual mean values of life history traits. In b and d, data points show mean values from low, moderate, and high androgen exposure levels.

Figure 5. Deviation (%) compared to the baseline scenario in life history traits of shovelnose sturgeon populations under combined scenario of energetic and demographic perturbations. a) adult sex ratio, proportion of adult females in the reproductive cycle, recruit number, reproductive female number, and spawner biomass; b) relationship between relative reproductive female number and spawner biomass or recruit number during exposure and recovery, and c) relationship between adult sex ratio and reproductive female number, spawner biomass, or recruit number during exposure and recovery. In a, black circles indicate annual mean values of life history traits. In b and c, data points show mean values from low, moderate, and high androgen exposure levels.

FIGURE LEGENDS FOR APPENDIX 1

Figure A1. Daily mean river discharge rates measured at five USGS gaging stations (Duncan–station #06774000, North Bend–#06796000, Leshara–#06796500, Ashland–#06801000, and Louisville–#06805500) during 1995–2011.

Figure A2. a) Relationship between energy allocation to female gonad development and vitellogenin computed from field estimates of vitellogenin concentrations and gonadosomatic index in pre-spawning female shovelnose sturgeon reported in Wildhaber et al. (2007); and b) simulation scenarios tested for the maternal investment in egg production (energetic perturbation). In a, Stage I, II, III, IV, and V represent immature, developing, vitellogenesis, pre-spawning, spawning, and spent (post-spawning) of the female gonad development (Wildhaber et al. 2007). In b, the baseline relationship is based on data reported in Wildhaber et al. (2007).

Figure A3. Temporal patterns of demographic and reproductive traits of simulated shovelnose sturgeon populations under the baseline scenario – a) adult sex ratio, b) female number, c) relative female gonad mass, d) fecundity, e) spawner biomass, f) recruit number, and g) a stock-recruit relation. The sturgeon model used for the baseline simulation has been calibrated and validated with field survey data on the lower Platte River population (Goto *et al.* 2015).

REFERENCES FOR APPENDIX 1

Goto, D., Hamel, M.J., Hammen, J.J., Rugg, M.L., Pegg, M.A. & Forbes, V.E. (2015) Spatiotemporal variation in flow-dependent recruitment of long-lived riverine fish: Model development and evaluation. *Ecological modelling,* **296,** 79-92.

Wildhaber, M.L., Papoulias, D.M., DeLonay, A.J., Tillitt, D.E., Bryan, J.L. & Annis, M.L. (2007) Physical and hormonal examination of Missouri River shovelnose sturgeon reproductive stage: a reference guide. *Journal of Applied Ichthyology,* **23,** 382-401.

a

biomass (mt)

baseline

time (year)

low ─── medium ······ high ─── ─

exposure   recovery

energetic perturbation

deviation (%)

demographic perturbation

combined

time (year)

b

baseline ─── energetic perturbation ─── demographic perturbation ─── combined ───

exposure   recovery

per capita population growth ln(N_{t+1}/N_t)

$\log_{10}$ population biomass

Supporting Information

Table S1. Summary of exposure period (27 years) model outputs (mean ± sd) from the shovelnose sturgeon spatially explicit individual-based model simulations for energetic (reduced energy allocation to female gonad development) and demographic (pre-recruit sex ratio skewness) perturbation scenarios. Vt_low, Vt_mid, and Vt_high indicate vitellogenin reduction in adult females exposed to low-, mid-, and high- level synthetic androgen. SS_low, SS_mid, and SS_high indicate skewed sex ratio in larvae exposed to low-, mid-, and high- level synthetic androgen. Numbers indicates mean values computed across years.

model output	baseline	Vt_low	Vt_mid	Vt_high	SS_low	SS_mid	SS_high	Vt + SS_low	Vt + SS_mid	Vt + SS_high
adult female										
sex ratio (proportion of female)	0.31±0.15	0.32±0.13	0.32±0.13	0.32±0.13	0.29±0.14	0.27±0.13	0.24±0.11	0.31±0.12	0.27±0.11	0.27±0.12
mean age of adult female	6.88±0.56	6.86±0.52	6.85±0.52	6.86±0.52	6.90±0.57	6.92±0.56	6.89±0.55	6.88±0.53	6.87±0.54	6.86±0.52
reproductive female abundance	3119.0±2260.5	2819.4±1979.2	2505.4±1782.5	2002.4±1550.5	2643.6±2023.5	2532.6±1908.1	2475.2±1836.0	2138.0±1596.5	2092.3±1542.9	1648.4±1407.2
mean relative storage mass of reproductive female	0.24±0.04	0.24±0.04	0.24±0.04	0.24±0.04	0.24±0.04	0.24±0.04	0.24±0.04	0.24±0.04	0.24±0.04	0.24±0.03
mean relative storage mass of non-reproductive	0.25±0.02	0.24±0.05	0.24±0.02	0.25±0.02	0.24±0.04	0.25±0.03	0.24±0.04	0.24±0.03	0.24±0.02	0.24±0.03
mean relative gonad mass of reproductive female	0.10±0.03	0.10±0.03	0.10±0.03	0.10±0.03	0.10±0.03	0.10±0.03	0.10±0.03	0.10±0.03	0.10±0.03	0.10±0.03
number of spawner	701.6±708.3	714.3±670.4	680.1±.670.7	467.1±572.5	604.4±646.7	600.0±653.5	577.6±645.5	500.8±593.8	480.1±577.6	406.2±558.2
mean spawning location (rkm)	103.9±.12.9	103.7±.13.2	100.1±13.9	99.0±.13.9	103.4±.11.3	103.9±15.7	104.0±.15.2	102.2±.17.1	100.4±.15.5	124.8±.14.0
mean spawning time (day of year)	117.6±.12.1	116.0±.10.6	115.3±10.6	113.7±10.6	117.3±.14.3	118.5±12.6	118.5±.11.6	115.6±.11.0	114.1±.12.7	113.5±.11.5
mean egg number per female	33369.5±2697.1	33622.5±2570.8	33965.5±2571.0	34667.3±2882.0	32899.3±3189.0	33074.0±3232.3	33023.7±3173.8	33551.3±3217.0	33815.4±3145.3	34326.9±3423.3
mean spawning interval (years)	2.81±0.52	2.91±0.53	2.88±0.54	3.01±0.55	2.91±0.66	2.89±0.55	2.77±0.54	3.04±0.69	3.00±0.55	3.99±0.55
early life stages										
total egg number (millions)	22.7±22.1	20.1±20.6	18.1±18.9	15.2±16.9	19.8±20.1	19.5±20.1	22.5±19.9	16.2±17.6	15.6±17.2	17.7±18.4.
mean larval settlement location (rkm)	100.2±20.9	97.7±16.6	92.7±15.4	92.4±17.3	100.3±22.4	97.9±21.1	99.1±19.2	97.4±15.9	99.8±17.8	92.8±17.9
mean first-year survival rate (%)	0.11±0.22	0.11±0.18	0.085±0.14	0.074±0.12	0.063±0.097	0.054±0.095	0.057±0.10	0.057±0.092	0.069±0.12	0.041±0.083
total age 0 recruit number	8323.5±16732.8	7750.3±14896.6	5782.8±11599.4	3479.3±7535.8	6579.9±12886.2	5812.1±11200.5	8050.7±15081.8	3463.1±7271.0	4192.3±9077.9	3656.4±7152.8
age 0 recruit sex ratio (proportion of female)	0.57±0.22	0.56±0.19	0.48±0.13	0.52±0.22	0.47±0.20	0.44±0.18	0.38±0.19	0.48±0.23	0.55±0.17	0.52±0.25

Table S2. Summary of recovery period (30 years) model outputs (mean ± sd) from the shovelnose sturgeon spatially explicit individual-based model simulations for energetic (reduced energy allocation to female gonad development) and demographic (skewed sex ratio) perturbation scenarios. Vt_low, Vt_mid, and Vt_high indicate vitellogenin reduction in adult females exposed to low-, mid-, and high- level synthetic androgen. SS_low, SS_mid, and SS_high indicate skewed sex ratio in larvae exposed to low-, mid-, and high- level synthetic androgen.

model output	baseline	Vt_low	Vt_mid	Vt_high	SS_low	SS_mid	SS_high	Vt + SS_low	Vt + SS_mid	Vt + SS_high
adult female										
sex ratio (proportion of female)	0.31±0.11	0.31±0.09	0.31±0.09	0.31±0.08	0.31±0.11	0.29±0.09	0.28±0.13	0.30±0.09	0.28±0.09	0.29±0.11
mean age of adult female	6.69±0.17	6.71±0.23	6.67±0.21	6.51±0.64	6.70±0.31	6.67±0.24	6.80±0.47	6.64±0.28	6.67±0.20	6.55±0.13
reproductive female abundance	2282.5±1024.0	2958.4±1212.7	2044.6±831.4	837.8±426.5	2087.0±1034.9	822.3±404.0	1133.6±1052.6	1458.2±713.4	1867.9±1183.2	1022.3±516.7
mean relative storage mass of reproductive female	0.25±0.02	0.25±0.02	0.25±0.02	0.26±0.02	0.25±0.02	0.25±0.02	0.25±0.02	0.25±0.01	0.25±0.02	0.26±0.01
mean relative storage mass of non-reproductive	0.24±0.01	0.24±0.01	0.24±0.01	0.23±0.02	0.23±0.04	0.22±0.04	0.22±0.04	0.23±0.02	0.23±0.02	0.23±0.03
mean relative gonad mass of reproductive female	0.09±0.01	0.09±0.01	0.09±0.01	0.09±0.01	0.09±0.02	0.09±0.01	0.09±0.01	0.09±0.01	0.09±0.01	0.09±0.01
number of spawner	436.0±356.5	504.6±459.0	355.9±335.3	145.9±122.4	357.5±258.6	137.6±114.4	209.9±291.4	256.8±172.1	346.8±341.3	178.4±133.2
mean spawning location (rkm)	102.1±.6.3	100.6±6.2	102.6±.7.9	102.7±.6.8	103.4±.11.3	104.1±7.8	102.3±.7.0	104.0±6.8	101.8±.7.8	102.6±7.3
mean spawning time (day of year)	119.0±.5.6	121.5±6.4	120.0±7.6	120.7±6.0	118.8±9.6	119.7±6.6	118.3±8.2	114.8±.6.4	119.4±7.0	119.0±8.4
mean egg number per female	33886.9±1414.8	33499.4±1177.1	33925.7±1221.6	33833.5±1285.7	34562.2±1880.3	34199.4±1487.3	34551.2±1463.8	34768.0±1365.8	34104.0±1659.9	34905.2±1858.5
mean spawning interval (years)	3.18±0.36	3.08±0.43	3.24±0.42	3.21±0.47	3.27±0.38	3.24±0.36	3.09±0.37	3.31±0.31	3.18±0.36	3.22±0.40
early life stages										
total egg number (millions)	14.7±12.1	17.2±11.9	12.1±9.03	4.87±3.99	12.1±8.48	4.71±3.94	8.39±12.4	8.85±5.88	11.8±11.6	5. 90±5.35
mean larval settlement location (rkm)	96.8±8.9	95.9±10.0	97.6±10.9	94.6±9.5	97.7±10.8	99.0±11.2	98.2±13.7	99.1±10.3	97.6±9.0	95.9±10.8
mean first-year survival rate (%)	0.11±0.19	0.13±0.17	0.11±0.15	0.13±0.17	0.076±0.011	0.13±0.22	0.076±0.15	0.11±0.18	0.28±0.99	0.20±0.24
total age 0 recruit number	9340.3± 15274.4	11045.1±16848.3	6584.9±10952.2	3191.5±4420.9	5497.5±10621.3	2761.4±4475.5	2030.62±4863.7	3920.8±5213.8	7666.8±14721.8	4188.1±5734.9
age 0 recruit sex ratio (proportion of female)	0.56±0.16	0.52±0.10	0.51±0.11	0.51±0.16	0.51±0.20	0.53±0.18	0.52±0.22	042±0.17	0.47±0.17	0.53±0.23