Divergent population recovery trajectories following sex-biased perturbation in life history traits

Daisuke Goto[1,2,*], Martin J. Hamel[3], Jeremy J. Hammen[4], Matthew L. Rugg[5], Mark A. Pegg[3],

and Valery E. Forbes[6]

[1]Institute of Marine Research/Havforskningsinstituttet, Postboks 1870, Nordnes, 5817 Bergen,

Norway

[2]School of Biological Sciences, Manter Hall, University of Nebraska-Lincoln, Lincoln,

Nebraska, 68588 USA

[3]School of Natural Resources, Hardin Hall, 3310 Holdrege Street, University of Nebraska-

Lincoln, Lincoln, Nebraska 68583-0973 USA

[4]U.S. Fish and Wildlife Conservation Office, 101 Park Deville Drive, Suite A, Columbia,

Missouri 65203 USA

[5]Montana Fish, Wildlife & Parks, 907 North Kendrick Avenue, Glendive, Montana 59330 USA

[6]Department of Ecology, Evolution and Behavior, University of Minnesota, 123 Snyder Hall,

1475 Gortner Ave, St. Paul, Minnesota 55108 USA

*Corresponding author Email: daisuke.goto2@gmail.com

Abstract

Uncertainty in risks posed by emerging stressors such as synthetic hormones impedes the

management and conservation of threatened and endangered long-lived vertebrate populations.

Synthetic hormones often induce sex-biased perturbations in exposed animals by disrupting

gonad development and early life-history stage transitions, threatening population persistence.

We use a spatially-explicit biophysical model to evaluate how sex-biased perturbations in life

history traits of individuals (maternal investment in egg production and sex determination)

modulate year class strength and population recovery trajectories of a long-lived, migratory fish–

shovelnose sturgeon (*Scaphirhynchus platorynchus*)–under spatially and temporally dynamic

synthetic androgen exposure. Simulations revealed that reduced efficiency of maternal energetic

investment in gonad development prolonged maturation time, increased the probability of

skipped spawning, and, in turn, gradually shrunk spawner biomass, ultimately weakening year

class strength and extending recovery time at higher exposure. Demographic perturbation in

individuals emerged as vastly different population projection and recovery trajectories: the

responses to skewed sex allocation were delayed at the population level by more than two

decades, followed by an abrupt shift in year class strength and population size, ultimately

reaching an alternate state without any sign of recovery. When energetics and demographics

were both perturbed, they acted antagonistically, allowing the population to gradually recover

without reaching a stable state. The emergent patterns in long-term population projections

suggest the complexity of cumulative effects of sex-biased perturbation in life history traits of

individuals exposed to low-level (often below detection) synthetic hormones, which can be

further obscured by lagged responses and environmental stochasticity. Our study illustrates that

the recovery from skewed sex allocation can pose greater conservation challenges than

diminished maternal investment in egg production–highlighting the importance of population

monitoring for restoring depleted long-lived populations such as *Scaphirhynchus* sturgeon in

ever-changing large rivers.

Keywords: early life history; endocrine disruptors; life history traits; endangered species; agent-

based modeling; spatially explicit models; conservation

1. INTRODUCTION

Humans activities have uniquely shaped life history strategies of wild populations of animals by imposing unnatural stress for centuries (Crain, Kroeker & Halpern 2008; Hendry, Farrugia & Kinnison 2008). The symptoms of human-induced stressors, in turn, manifest in life history traits varyingly among individuals (Marr *et al.* 2006) over time and space (Hope 2005; Maxwell *et al.* 2013). Sexual dimorphism, for example, is prevalent in natural populations (Fairbairn 1997), often leading to sex-biased pressure (Coltman *et al.* 2003). As stress becomes intensified, these sources of intraspecific variation in fitness may emerge as a directional shift in demographic rates (Wedekind 2017), diminishing reproductive output (Knell & Martínez-Ruiz 2017). Because of sex-dependent differential investment in reproduction, sex-biased stresses may pose a unpredictable threat to persistence of wild populations (Coltman *et al.* 2003).

One emerging stressor with such effects is synthetic chemicals, which have been increasingly detected in aquatic systems around the globe (Bernhardt, Rosi & Gessner 2017). A rising demand for human food production has increased agricultural use of synthetic chemicals (Tilman 1999; Tilman *et al.* 2001), posing a greater risk to freshwater vertebrates (Bernhardt, Rosi & Gessner 2017). Synthetic hormones–often used as growth-promoters for farmed animals–are especially potent, impairing fish development and reproduction by disrupting energy allocation and maturation (del Carmen Alvarez & Fuiman 2005). These indirect processes thus stunt somatic and gonadal growth (Arcand Hoy & Benson 1998). In pre-spawning females, even low-dose (often below detection) exposure to synthetic hormones can disrupt maturation processes such as vitellogenin production–a critical step in egg production (Arcand Hoy & Benson 1998; Nash *et al.* 2004). Because maternal investment in egg production incur greater energetic costs than somatic growth, females may more likely experience reproductive impairment than males

from chemical perturbation in gonad development (Arukwe & Goksøyr 1998). Further, synthetic

hormones can irreversibly perturb sex determination, skewing sex allocation in offspring (Mills

& Chichester 2005). Modified demographic traits such as skewed sex ratio have been reported in

wild animal populations with sex-biased mortality (McTavish, Stech & Stay 1998). However, it

remains unclear how skewed sex allocation in early life stages resulting from exposure to

synthetic hormones influence population persistence and recovery in nature (Ankley *et al.* 2008).

We here present a simulation study on how long-term sex-biased perturbations in life history

traits can emerge and modify recovery potential of a migratory fish exposed to low-level

synthetic androgens in a large river. Migratory animals are exposed to spatially and temporally

dynamic flow-dependent exposure regimes as the chemicals move throughout networks of

streams and rivers (O'Brien *et al.* 2016); variable exposure and subsequent symptoms among

exposed individuals can obscure a causal link, if any, to population-level responses, delaying

management actions (Peterson *et al.* 2003). We test varying exposure and effect scenarios on the

Missouri River shovelnose sturgeon (*Scaphirhynchus platorynchus*) as an illustrative case study.

Population sizes of *Scaphirhynchus* sturgeon have declined in many of their endemic habitats,

and some are facing potential extirpation owing to extensive habitat modification (e.g.,

impoundments) (Bramblett & White 2001). Further, some of the Missouri River populations

have shown signs of unsuccessful spawning (DeLonay *et al.* 2009a) and year class failure owing

to low larval fish survival (Hrabik *et al.* 2007)–a threat to population persistence (DeLonay *et al.*

2009a). We use a spatially-explicit individual-based biophysical model to evaluate how modified

energetics (reduced maternal investment in egg production) and demographics (skewed sex

allocation)–two commonly reported symptoms of synthetic androgens–influence year class

success, and population persistence and recovery of sturgeon.

2. MATERIALS AND METHODS

2.1. Study system

The study system is the lower 162 km of the Platte River (Nebraska, USA), one of the largest

tributaries of the Missouri River basin (Fig. 1a and S1), providing spawning and nursing habitats

essential for threatened and endategered fish populations such as *Scaphirhynchus* sturgeon

(Peters & Parham 2008). Due to its proximity to wastewater treatment plants and agricultural

fields, however, the lower Platte River is also one of the impaired water bodies in the region

(NDEQ 2004). Fish that inhabit this river and its tributaries such as *Scaphirhynchus* sturgeon are

exposed to elevated concentrations of synthetic hormones widely used at concentrated cattle

feedlots near the lower Platte River watershed (Huntzinger & Ellis 1993) and exhibit symptoms

of endocrine disruption (e.g., intersexuality and atresia, DeLonay *et al.* 2009a).

2.2. Sturgeon biophysical model

We use a spatially explicit individual-based biophysical model of shovelnsoe sturgoen

previously developed for the Lower Platte River popuatlion; the model is fully described in Goto

et al. (2015) and Goto *et al.* (2018). The model is designed to uncover ecological mechanisms

underlying early life-history processes modified by natural and human-induced stressors, and

their contributions to population persistence under spatially and temporally dynamic habitat

conditions. The model primarily hinges on energetics principles to understand how maternal and

environmental influences modulate year class variability through energy acquisition and

allocation (Houde 1989). In this study, we evaluated sex-biased effects of synthetic androgen

perturbation in maternal investment in gonad development and first-year sex allocation (Fig. 1b

and S2). The following briefly describes the essetial features of the model relevant to this study.

The model river consists of 162 equally divided longitudinal rectangular segments (i.e., 1 rkm

each). Model inputs include water temperature (°C), mean gauge height (depth, m), channel

width (m), discharge rate ($m^3 s^{-1}$), and photoperiod (hrs). Water temperature, gauge height,

channel width, and discharge rate (1995–2011) were derived from the USGS National Water

Information System (http://waterdata.usgs.gov/nwis). Further details on derivation of physical

and hydrological input data are described in Goto *et al.* (2018).

The sturgeon model keeps track of state variables of the entire life cycle (Fig. 1b). Early life

stages (fertilized eggs, yolk-sac larvae, and feeding larvae, and age-0 juveniles) are represented

as super-individuals (Scheffer *et al.* 1995), whereas age 1+ juveniles and adults are represented

as individuals. Each super-individual is a collection of individuals ("cohorts") from the same

mother. We initialized simulations with 60 000 age 1+ individuals on January 1st. The initial

location of each sturgeon was randomly assigned to a cell with non-extreme water velocity ($\leq$0.6

$m s^{-1}$) and depth ($\geq$0.5 m). The initial age structure follows a Poisson distribution with mean age

of 7, which is truncated at a maximum age of 16, and the initial sex ratio is 1:1. Sex was

randomly assigned to each fish, assuming a binomial distribution. For age 1+ sturgeon, initial

lengths-at-age were computed using the von Bertalanffy growth function (Ricker 1975)

parameterized for the Platte River population (Goto *et al.* 2018). Once spawned, fertilized eggs

were initialized as super-individuals; the number of super-individuals is the number of mothers,

and the number of individuals in each super-individual is the total number of eggs spawned by

each mother. After settlement, feeding larvae were initialized with individual-level processes

(e.g., growth) and traits (e.g., body size).

The model simulates the population and updates state variables at discrete daily time steps over

years and generations. The habitat conditions (including prey production) are first updated, and

then the model evaluates actions by sturgeon in the following order; (i) *spawning,* (ii) *hatching*

and larval development, (iii) *movement,* (iv) *foraging,* (v) *growth*, and (vi) *mortality* (Fig. 1b).

Density dependence is incorporated in directly foraging and indirectly in growth. Sturgeon's

abilities to sense habitat conditions and prey density, capture prey, decide when to spawn, and

survive are partially stochastic because of inherent uncertainty and randomness.

The model consists of one benthic prey submodel and six sturgeon submodels (Goto *et al.*

2018). Benthic prey production is computed using a mass-balance model described by Rashleigh

& Grossman (2005) with modifications described in Goto *et al.* (2018). The early stages of larval

development are computed using temperature-dependent nonlinear functions (Goto *et al.* 2018).

Foraging is computed using a modified version of the Beddington–DeAngelis functional

response model (Beddington 1975; DeAngelis, Goldstein & O'neill 1975). Growth is simulated

by the bioenergetics model (Hanson *et al.* 1997) modified so that the assimilated energy is

allocated to storage (reversible) and structure (irreversible) (Gurney *et al.* 2003). Sturgeon

mature based on sex-specific probability, which is length-dependent functions parameterized for

the Platte River population (Goto *et al.* 2018). Reproductive adults further allocate energy to

gonad incrementally between spawning events. Adults initiate the reproductive cycle only when

their physiological condition is above a threshold based on relative storage mass. The amount of

daily energy allocated to gonad is further adjusted when the physiological condition is below the

threshold (Goto *et al.* 2018). Spawning is simulated as a stochastic event and occurs only when

spawning cues–water temperature, river discharge rate, daylight hours, and gonad development

(gonadosomatic index or GSI), concurrently meet the thresholds (Delonay *et al.* 2016). Water

temperature suitable for the female *Scaphirhynchus* sturgeon spawning ranges is set at 12.6–

24.9°C (DeLonay *et al.* 2009b). A minimum threshold for river discharge rate was set at 141.6

$m^3\ s^{-1}$, which is expected to reach roughly 50% connectivity (migratory pathway connectedness)

in the Lower Platte River (Peters & Parham 2008) and directly influences movement and

subsequently habitat conditions for other individual-level actions. A minimum GSI to initiate

spawning is set at 0.18 for females and 0.03 for males (Wildhaber *et al.* 2007a). After spawning,

fish lose all gonad mass, and their physiological condition and total mass are updated; fish do not

re-enter the reproductive cycle until the physiological condition meets the threshold. When either

temperature or discharge rate does not meet the thresholds, fish skip spawning. We assume that

when the physiological condition does not meet the threshold to remain in the reproductive

cycle, gonadal energy is reabsorbed and reallocated to storage. Movement is tracked in the

longitudinal direction only. The planktonic life stage fish (eggs and yolk-sac larvae) move by

passive drift with water currents (Braaten *et al.* 2008). Feeding larvae, juveniles, and adults move

in the direction with probability of attaining higher habitat quality using the empirical models

developed for the Lower Platte River population (Peters & Parham 2008). Mortality results from

thermal stress (eggs and yolk-sac larvae), predation (eggs, larvae, and juveniles), starvation, and

recreational fishing (age 3+). We assume that a constant proportion of eggs and yolk-sac larvae

die when water temperature rises above 24 °C (Kappenman, Webb & Greenwood 2013).

Realized predation, starvation, and fishing mortality are a stochastic event. Starvation mortality

depends on relative storage mass (Goto *et al.* 2018).

2.3. Simulation scenarios for sex-biased purtubations by synthetic androgens

We ran 57-year (27-year exposure + 30-year recovery) simulations for the following four

scenarios to assess the population performance under two types of perturbations in life history

traits of individuals induced by low-level synthetic androgens, energetics and demorgraphics;

1) No sex-biased purtubation–baseline;

2) Synthetic androgens perturb maternal energetic investment in egg production by imposing

reduction in vitellogenin production rate;

3) Synthetic androgens perturb sex allocation by imposing skewness in sex ratio at recruitment;

and

4) Synthetic androgens perturb both maternal energetic investment and sex allocation. We

assumed no direct mortality caused by synthetic androgen exposure. Further, we assumed that: a)

synthetic androgens are released between May 1st and October 31st (a six-month growing

season); b) androgens affect only fish located below the confluence with the Elkhorn River (~51

rkm from Plattsmouth, NE, Fig 1a), where a number of cattle feedlots are located; and c) because

the androgen concentration varies with stream flow rate, the realized exposure level for

individual fish is spatially and temporally dynamic, and is thus adusted based on flow rate in a

grid cell in which the given fish is located.

Energetic perturbation. We explicitly evaluated the mechanism underlying synthetic androgen-

induced reduction in egg production by simulating the dose-dependent reduction of vitellogenin

production in individual fish based on an exponential decay function (Ankley *et al.* 2003).

Vitellogenin (VTG) concentration (mg ml^{-1}) is converted to relative gonad mass (GSI) using an

empirically derived linear function based on field survey data on the Missouri River shovelnose

sturgeon reported by Wildhaber *et al.* (2007b) (Fig. S2) as GSI = 0.984VTG + 35.103. Reduction

in daily energy allocation to gonad (g d^{-1}) in exposed females was then computed based on

somatic mass on the given day.

We tested the scenarios of 15%, 25%, and 35% reduction in daily vitellogenin production in

fish that are located at the Elkhorn River–Platte River confluence; these reduction rates in

vitelogenin production approximately corresspond to the responses in fish exposed to

environmentally realistic, low level sysnthetic androgens (e.g., <~15 ng l^{-1} 17β-trenbolone, a

synthetic androgen commonly used in cattle feedlots, Ankley *et al.* 2003). For fish that are

located in cells below the confluence, the reduction rates were proportionally adjusted by the

difference in flow rate (e.g., if flow rates at the confluence and in the cell where a fish is located

are 100 and 120 m^3 s^{-1}, respectively, the realized reduction rate for this fish is 12.5% under the

15% reduction scenario). Because the model sturgeon cannot detect low-level synthetic

androgens in water, we assumed that their movement and energy acquisition (foraging) vary

independently of synthetic androgens.

Demographic perturbation. We implemented synthetic androgen-induced skewness in sex

allocation by proportionally reducing the probability of being female. We tested the scenarios of

20%, 60%, and 80% reduction in probability of being females for post-hatch fish that remain in

the cell at the Elkhorn River–Platte River confluence; these rates approximately corresspond to

the responses in fish exposed to environmentally realistic concentrations (e.g., <~15 ng l^{-1} 17β-

trenbolone, Morthorst, Holbech & Bjerregaard 2010). We assumed that the probability of being

female incrementally declines by remaining in the exposure area each day (i.e., 0.11%, 0.33%,

and 0.44%, respectively). For fish in cells below the confluence, the probability was

proportionally adjuted by the difference in flow rates (as above).

For all the scenarios, we used hydrological and temperature inputs empirically derived from

the lower Platte River field surveys during 1995–2011 (Goto *et al.* 2018), from which we

randomly selected each year; we assumed no systematic change or shift in hydrological and

thermal conditions in this study. Further, we assumed no change in turbidity, channel

geomorphology, and photoperiod. We ran ten simulations for each exposure scenario and

computed summary statistics and % deviations in demographic and life-history traits from the

baseline simulations.

3. RESULTS

3.1. Baseline sturgeon population dynamics

Under the baseline conditions, the sturgeon population remained stable over 57 years (Fig. 2

and S3). The population maintained, on average, ~32% of adults being female, ~15.8 metric ton

spawner biomass, and ~7100 recruits (Fig. S3, Table S1 and S2). Adult females allocated, on

average, ~9.4 % of total mass to gonad while they were in the reproductive cycle; ~22% of

reproductive females eventually spawned, producing, on average, 23 million eggs (or ~33000

eggs spawner^{-1}) annually (Table S1 and S2). Most spawning and subsequent larval settlement

occurred below the confluence of the Elkhorn River and the Platte River (112 rkm) in late April

to early May (Table S1 and S2). Although the recruit numbers highly varied among years, a non-

linear (hump-shaped) density-dependent relationship emerged between spawners and recruits

(Fig. S3); on average, 0.11% of first-year fish survived (Table S1 and S2).

3.2. Recovery from reduced vitellogenin production efficiency

Simulated reduction in vitellogenin production rate gradually lowered reproductive capacities

and, in turn, biomass production of the sturgeon population exposed to synthetic androgens (Fig.

2 and 3). The proportion of energy allocated to gonad mass (10%) or storage mass (24%) in

reproductive females under exposure did not differ from those under baseline (Table S1).

However, the number of adult females in the reproductive cycle declined by 7.5–34% during

exposure (Fig. 4a,b). Further, spawner biomass and recruit numbers declined especially when

exposed to moderate and high levels of the perturbation (6.8–25% and 23–38%, respectively,

Fig. 3b, c), which were linearly correlated with relative changes in reproductive female

abundance (Fig. 3e). Lowered reproductive capacities and year class resulted in 8.6–43%

reduction (relative to baseline) in population biomass during exposure (Fig. 2b and 3d).

Similar relationships remained during recovery; reproductive females, spawners, and recruits

of the population with moderate exposures returned to the baseline levels with population size

fully recovered (i.e., mean population size returns to the baseline level) within five years (Fig. 2b

and 3), whereas the reproductive females, spawners, and recruits of the population with high

exposure continued to decline, ultimately reducing mean population biomass by up to 83%

before showing a sign of recovery after 15 years (~year 42, Fig. 2b).

3.3. Recovery from skewed recruit sex ratio

Sturgeon spawner biomass abruptly declined by up to 67% by ~year 22 under all scenarios of

simulated skewed sex ratios during the first year of life (Fig. 2c). Skewed sex ratios in recruits

were reflected in adults within ~6 years with relative female numbers declining, on average, by

6.3–25% during exposure (Fig. 3g). Subsequently, the number of adult females entering the

reproductive cycle and spawner biomass both started declining by ~year 8. During exposure,

mean reproductive female abundance declined by 19–21%; mean spawner biomass declined by

up to 29% under high exposure (Fig. 3h and 3i). In contrast, mean recruit number declined under

low and moderate exposure (by 15% and 23%, respectively), but differed little from the baseline

population under high exposure (Fig. 3j).

During recovery, mean sex ratios under all the exposure scenarios remained slightly (by 9.3–

16%) skewed toward males with high interannual variation (Fig. 3g). However, only the low-

exposure population fully recovered, whereas the moderate and high exposure populations

continued to show declining reproductive female abundance, spawner biomass, and recruit

number (Fig. 3). Despite large year class in early recovery years (year 31–33), more than 85% of

adults were males under high exposure and did not contribute to spawner biomass (Fig. 3). The

moderate and high exposure spawner biomass ultimately reached a stable state with less than

20% of the baseline population (Fig. 2c).

3.4. Recovery from combined effects of reduced vitellogenin production and skewed sex ratio

Concurrently simulating reduction in vitellogenin production and skewed sex ratios modulated

the sturgeon population dynamics additively under low exposure, but antagonistically under

moderate and high exposure (Fig. 2d). Spawner biomass declined within 5 years and remained

relatively stable and then declined further during the last five years (Fig. 2d). During exposure,

adult sex ratio differed little under low exposure, but skewness toward males increased by 11%

under medium and high exposure (Fig. 4a). Further, reproductive female abundance, spawner

biomass, and recruit number declined by 31%, 23%, and 31% (respectively) under low and

moderate exposure, and declined by 46%, 29%, and 47% (respectively) under high exposure

(Fig. 4b-d).

During recovery, adult sex ratios remained skewed toward males (by ~11%) under all the

exposure scenarios (Fig. 4a). Mean reproductive female number, spawner biomass, and recruit

number (except for reproductive females with low exposure) recovered after low and moderate

exposures but remained low under high exposure (Fig. 4b-d). Spawner biomass under low and

moderate exposure recovered within ~25 years of recovery, whereas the biomass of the

population under high exposure reached a stable state in ~22 years at ~44% lower than the

baseline population (Fig. 2d).

4. DISCUSSION

Simulations revealed that energetic (gonad maturation) and demographic (sex allocation)

perturbations in life history traits of long-lived, migratory freshwater vertebrates such as

Scaphirhynchus sturgeon can emerge as vastly different response and recovery patterns in

population projection in a spatially and temporally dynamic environment. We find that lowered

vitellogenin production rate prolongs maturation and, in turn, reduces spawner biomass,

ultimately weakening year class strength. Further, the energetic perturbation operates as

exposure-dependent processes–the higher the exposure level, the more severe the impacts,

thereby resulting in longer recovery time. In contrast, skewed sex allocation emerges as delayed

(by more than two decades) population-level responses, followed by an abrupt shift in year class

strength and population trajectory under higher exposures, ultimately reaching an alternative

(stable) state without any sign of recovery. When energetics and demographics are concurrently

perturbed, however, they act antagonistically as the exposure level increases, allowing it to

gradually recover without reaching a stable state. These contrasting emergent patterns at the

population level highlight the complexity of cumulative effects of stochastic perturbations in life

history traits by long-term, low-level exposure to stressors, posing challenges to conservation

efforts to restore the depleted vertebrate population.

Delayed gonad maturation in energetically perturbed females prolongs the spawning cycle of

our model sturgeon population instead of reducing fecundity. Sex-biased differences in energy

allocation to gonadal maturation often determine life-history strategies such as foraging and mate

selection, shaping sexual dimorphism in animal populations (Huse 1998; Rennie *et al.* 2008). For

a capital spawner such as sturgeon, slowed female gonad maturation delays spawning timing.

Although sturgeon are iteroparous, because of infrequent spawning and long maturation time,

delayed spawning events can reduce lifetime reproductive output (our simulations show that

some individuals spawned only once). Inefficient energy use triggered by synthetic hormones in

animals with long maturation time may thus threaten population persistence. Our simulations,

however, also suggest that the energetic perturbation triggers plastic responses. Although

reproductive and spawning females decline by up to 25%, the population eventually recovers,

indicating the reproductive capacity diminished during androgen exposure can be restored if

energy allocation to gonad resumes.

Delayed spawning events may also increase the probability of year class failures owing to

environmental stochasticity. We find that exposed (energetically challenged) female sturgeon are

more likely to abort the reproductive cycle as they divert energy to meet metabolic needs when

poor environmental conditions persist. Although energetic tradeoffs between migration and

fecundity are beyond the scope of our study, animals with long spawning migration such as

sturgeon likely skip spawning or produce offspring with lower survival rate owing to reduced

energy reserves (Glebe & Leggett 1981). Sufficient energy reserves in eggs are vital to larval

development and survival; maternal investment in egg production determines the nutritional

status of post-hatching larvae (Kjesbu *et al.* 1996). Our model assumes that individual eggs do

not vary in energy content; however, energetic tradeoffs in maternal investment in egg

production and offspring survival can arise and vary greatly among taxa (Jönsson 1997; McBride

et al. 2015) and thus need further investigation for *Scaphirhynchus* sturgeon.

Our study suggests that the recovery from demographic perturbation can pose greater

conservation challenges than energetic perturbation. We find that a long-lived animal population

such as sturgeon experiencing skewed sex allocation can remain in a small population size for

long time. Although skewed sex ratio has been widely reported for a variety of vertebrate taxa

(Donald 2007), causes of the skewness vary; it can result from phenotypic sex-biased differences

in physiology, dispersal, and (natural or human-induced) mortality (Weimerskirch, Lallemand &

Martin 2005; Pen *et al.* 2010). In our baseline model population, adult sex ratios are skewed

toward males (~1:2.2), primarily resulting from 1) a longer maturation time for females than

males and 2) more energy investment for reproduction in females, resulting in higher probability

of starvation in poor years. Simulations, however, show that further deviation (by 13–23%) in

adult sex ratio skewness result in long-lasting threats to year class success and population

viability. Consequences of skewed adult sex ratio may not immediately emerge for long-lived

species (Holmes & York 2003; Lawson *et al.* 2010). Once the skewness reaches a tipping point

(Scheffer *et al.* 2009), however, it often becomes difficult for a population to recover (Scheffer *et*

al. 2001) as it may stabilize in an alternative state facilitated by a positive feedback (Van Nes &

Scheffer 2007; Dai *et al.* 2012). In conservation planning for long-lived migratory species such

as sturgeon, thus, to prevent the shifts and minimize recovery cost, it is critical to identify early

warnings that signal impending demographic shifts; skewness in adult sex ratio can be a sensitive

metric for assessing viability of stressed populations (Ancona *et al.* 2017).

Long-term, low-level exposure to stressors such as synthetic hormones often obscures

causative relationships (Windsor, Ormerod & Tyler 2018). Establishing ecologically relevant

indicators such as deviation in adult sex ratio and vitellogenin production rate would guide the

risk assessment of synthetic hormones and habitat conservation planning for depleted

populations. Further, given potential time lags across biological hierarchy, continued biological

monitoring of individual-level traits would help detect early signals that may indicate population-

level changes. Although our study was designed to evaluate consequences of low-level exposure,

our findings indicate larger implications of among-individual variation in sex-biased

perturbations for population persistence and recovery of threatened and endangered species.

Population models widely used in risk assessments often assume no individual-level variation in

each size or age group. These models may not only gloss over individual variation in fertility and

age at maturity that may contribute to population viability, but also under- or over- estimate

effects of synthetic hormones and other stressors (Jobling & Tyler 2003).

Large-river migratory populations such as *Scaphirhynchus* sturgeon likely experience multiple

synthetic hormones over agriculturally rich landscapes in regions such as the Midwestern United

States (Welcomme 1995). Further confounding factors in conservation and management efforts

can arise in hydrologically altered systems such as the Platte River. Precipitation can affect the

amount of agrichemicals released into rivers. Assessing the likelihood of ecological risks of

agrichemicals for long-lived species requires the development of trait-based predictive models

using anticipated scenarios. With expected trends in climate variability, which can have

considerable effects on river hydrology and water chemistry, these forecasting tools are critical

for fisheries and water resource management.

AUTHORS' CONTRIBUTIONS

DG, VEF, and MAP conceived the ideas; MJH, JJH and MLR collected and managed the data;

DG led the writing of the manuscript. All authors contributed to revisions and gave final

approval for publication.

ACKNOWLEDGEMENTS

REFERENCES

Ancona, S., Dénes, F.V., Krüger, O., Székely, T. & Beissinger, S.R. (2017) Estimating adult sex

ratios in nature. *Philosophical Transactions of the Royal Society B: Biological Sciences,*

372, 20160313.

Ankley, G.T., Jensen, K.M., Makynen, E.A., Kahl, M.D., Korte, J.J., Hornung, M.W., Henry,

T.R., Denny, J.S., Leino, R.L. & Wilson, V.S. (2003) Effects of the androgenic growth

promoter 17-β-trenbolone on fecundity and reproductive endocrinology of the fathead

minnow. *Environmental Toxicology and Chemistry,* **22,** 1350-1360.

Ankley, G.T., Miller, D.H., Jensen, K.M., Villeneuve, D.L. & Martinovic, D. (2008)

Relationship of plasma sex steroid concentrations in female fathead minnows to

reproductive success and population status. *Aquatic Toxicology,* **88,** 69-74.

Arcand Hoy, L.D. & Benson, W.H. (1998) Fish reproduction: an ecologically relevant indicator

of endocrine disruption. *Environmental Toxicology and Chemistry,* **17,** 49-57.

Arukwe, A. & Goksøyr, A. (1998) Xenobiotics, xenoestrogens and reproduction disturbances in

fish. *Sarsia,* **83,** 225-241.

Beddington, J. (1975) Mutual interference between parasites or predators and its effect on

searching efficiency. *Journal of Animal Ecology***,** 331-340.

Bernhardt, E.S., Rosi, E.J. & Gessner, M.O. (2017) Synthetic chemicals as agents of global

change. *Frontiers in Ecology and the Environment,* **15,** 84-90.

Braaten, P.J., Fuller, D.B., Holte, L.D., Lott, R.D., Viste, W., Brandt, T.F. & Legare, R.G. (2008)

Drift dynamics of larval pallid sturgeon and shovelnose sturgeon in a natural side channel

of the upper Missouri River, Montana. *North American Journal of Fisheries*

Management, **28,** 808-826.

Bramblett, R.G. & White, R.G. (2001) Habitat use and movements of pallid and shovelnose

sturgeon in the Yellowstone and Missouri rivers in Montana and North Dakota.

Transactions of the American Fisheries Society, **130,** 1006-1025.

Coltman, D.W., O'Donoghue, P., Jorgenson, J.T., Hogg, J.T., Strobeck, C. & Festa-Bianchet, M.

(2003) Undesirable evolutionary consequences of trophy hunting. *Nature,* **426,** 655-658.

Crain, C.M., Kroeker, K. & Halpern, B.S. (2008) Interactive and cumulative effects of multiple

human stressors in marine systems. *Ecology Letters,* **11,** 1304-1315.

Dai, L., Vorselen, D., Korolev, K.S. & Gore, J. (2012) Generic indicators for loss of resilience

before a tipping point leading to population collapse. *science,* **336,** 1175-1177.

DeAngelis, D., Goldstein, R. & O'neill, R. (1975) A model for tropic interaction. *Ecology***, 881-

892.

del Carmen Alvarez, M. & Fuiman, L.A. (2005) Environmental levels of atrazine and its

degradation products impair survival skills and growth of red drum larvae. *Aquatic

Toxicology,* **74,** 229-241.

Delonay, A.J., Chojnacki, K.A., Jacobson, R.B., Albers, J.L., Braaten, P.J., Bulliner, E.A.,

Elliott, C.M., Erwin, S.O., Fuller, D.B., Haas, J.D., Ladd, H.L.A., Mestl, G.E., Papoulias,

D.M. & Wildhaber, M.L. (2016) Ecological requirements for pallid sturgeon

reproduction and recruitment in the Missouri River—A synthesis of science, 2005 to

2012. *Scientific Investigations Report*, pp. 242. Reston, VA.

DeLonay, A.J., Jacobson, R.B., Papoulias, D.M., Simpkins, D.G., Wildhaber, M.L., Reuter, J.M.,

Bonnot, T.W., Chojnacki, K.A., Korschgen, C.E. & Mestl, G.E. (2009a) Ecological

requirements for pallid sturgeon reproduction and recruitment in the Lower Missouri

River: a research synthesis 2005-08.

DeLonay, A.J., Jacobson, R.B., Papoulias, D.M., Simpkins, D.G., Wildhaber, M.L., Reuter, J.M.,

Bonnot, T.W., Chojnacki, K.A., Korschgen, C.E., Mestl, G.E. & Mac, M.J. (2009b)

Ecological requirements for pallid sturgeon reproduction and recruitment in the Lower

Missouri River: A research synthesis 2005–08. U.S. Geological Survey Scientific

Investigations Report 2009–5201.

Donald, P.F. (2007) Adult sex ratios in wild bird populations. *Ibis,* **149,** 671-692.

Fairbairn, D.J. (1997) Allometry for sexual size dimorphism: pattern and process in the

coevolution of body size in males and females. *Annual review of Ecology and*

Systematics, **28,** 659-687.

Glebe, B. & Leggett, W. (1981) Temporal, intra-population differences in energy allocation and

use by American shad (*Alosa sapidissima*) during the spawning migration. *Canadian*

Journal of Fisheries and Aquatic Sciences, **38,** 795-805.

Goto, D., Hamel, M.J., Hammen, J.J., Rugg, M.L., Pegg, M.A. & Forbes, V.E. (2015)

Spatiotemporal variation in flow-dependent recruitment of long-lived riverine fish:

Model development and evaluation. *Ecological modelling,* **296,** 79-92.

Goto, D., Hamel, M.J., Pegg, M.A., Hammen, J.J., Rugg, M.L. & Forbes, V.E. (2018) Spatially

dynamic maternal control of migratory fish recruitment pulses triggered by shifting

seasonal cues. *Marine and Freshwater Research,* **69,** 942-961.

Gurney, W.S.C., Jones, W., Veitch, A.R. & Nisbet, R.M. (2003) Resource allocation,

hyperphagia, and compensatory growth in juveniles. *Ecology,* **84,** 2777-2787.

Hanson, P.C., Johnson, T.B., Schindler, D.E. & Kitchell, J.F. (1997) Fish bioenergetics 3.0.

University of Wisconsin Sea Grant Institute, Madison, WI.

Hendry, A.P., Farrugia, T.J. & Kinnison, M.T. (2008) Human influences on rates of phenotypic

change in wild animal populations. *Molecular ecology,* **17,** 20-29.

Holmes, E. & York, A. (2003) Using age structure to detect impacts on threatened populations: a

case study with Steller sea lions. *Conservation Biology,* **17,** 1794-1806.

Hope, B.K. (2005) Performing spatially and temporally explicit ecological exposure assessments

involving multiple stressors. *Human and Ecological Risk Assessment,* **11,** 539-565.

Houde, E.D. (1989) Comparative growth, mortality, and energetics of marine fish larvae:

temperature and implied latitudinal effects. *Fishery Bulletin,* **87,** 471-495.

Hrabik, R., Herzog, D., Ostendorf, D. & Petersen, M. (2007) Larvae provide first evidence of

successful reproduction by pallid sturgeon, Scaphirhynchus albus, in the Mississippi

River. *Journal of Applied Ichthyology,* **23,** 436-443.

Huntzinger, T.L. & Ellis, M.J. (1993) Central Nebraska river basins, Nebraska. *JAWRA Journal*

of the American Water Resources Association, **29,** 533-574.

Huse, G. (1998) Sex-specific life history strategies in capelin (*Mallotus villosus*)? *Canadian*

Journal of Fisheries and Aquatic Sciences, **55,** 631-638.

Jobling, S. & Tyler, C.R. (2003) Topic 4.3: Endocrine disruption in wild freshwater fish. *Pure*

and applied chemistry, **75,** 2219-2234.

Jönsson, K.I. (1997) Capital and income breeding as alternative tactics of resource use in

reproduction. *Oikos*, 57-66.

Kappenman, K.M., Webb, M.A.H. & Greenwood, M. (2013) The effect of temperature on

embryo survival and development in pallid sturgeon *Scaphirhynchus albus* (Forbes &

Richardson 1905) and shovelnose sturgeon *S. platorynchus* (Rafinesque, 1820). *Journal*

of Applied Ichthyology, **29,** 1193-1203.

Kjesbu, O.S., Solemdal, P., Bratland, P. & Fonn, M. (1996) Variation in annual egg production

in individual captive Atlantic cod (*Gadus morhua*). *Canadian Journal of Fisheries and*

Aquatic Sciences, **53,** 610-620.

Knell, R.J. & Martínez-Ruiz, C. (2017) Selective harvest focused on sexual signal traits can lead

to extinction under directional environmental change. *Proceedings of the Royal Society*

B: Biological Sciences, **284,** 20171788.

Lawson, D.M., Regan, H.M., Zedler, P.H. & Franklin, J. (2010) Cumulative effects of land use,

altered fire regime and climate change on persistence of Ceanothus verrucosus, a rare,

fire-dependent plant species. *Global Change Biology,* **16,** 2518-2529.

Marr, A., Arcese, P., Hochachka, W., Reid, J.M. & Keller, L. (2006) Interactive effects of

environmental stress and inbreeding on reproductive traits in a wild bird population.

Journal of Animal Ecology, **75,** 1406-1415.

Maxwell, S.M., Hazen, E.L., Bograd, S.J., Halpern, B.S., Breed, G.A., Nickel, B., Teutschel,

N.M., Crowder, L.B., Benson, S. & Dutton, P.H. (2013) Cumulative human impacts on

marine predators. *Nature communications,* **4,** 2688.

McBride, R.S., Somarakis, S., Fitzhugh, G.R., Albert, A., Yaragina, N.A., Wuenschel, M.J.,

Alonso-Fernández, A. & Basilone, G. (2015) Energy acquisition and allocation to egg

production in relation to fish reproductive strategies. *Fish and Fisheries,* **16,** 23-57.

McTavish, K., Stech, H. & Stay, F. (1998) A modeling framework for exploring the population

level effects of endocrine disruptors. *Environmental Toxicology and Chemistry,* **17,** 58-

67.

Mills, L.J. & Chichester, C. (2005) Review of evidence: Are endocrine-disrupting chemicals in

the aquatic environment impacting fish populations? *Science of the total environment,*

343, 1-34.

Morthorst, J.E., Holbech, H. & Bjerregaard, P. (2010) Trenbolone causes irreversible

masculinization of zebrafish at environmentally relevant concentrations. *Aquatic*

Toxicology, **98,** 336-343.

Nash, J.P., Kime, D.E., Van der Ven, L.T.M., Wester, P.W., Brion, F., Maack, G., Stahlschmidt-

Allner, P. & Tyler, C.R. (2004) Long-term exposure to environmental concentrations of

the pharmaceutical ethynylestradiol causes reproductive failure in fish. *Environmental*

Health Perspectives, **112,** 1725.

NDEQ (2004) 2004 surface water quality integrated report. Water Quality Division. pp. 118 pp.

O'Brien, D., Lewis, S., Davis, A., Gallen, C., Smith, R., Turner, R., Warne, M., Turner, S.,

Caswell, S. & Mueller, J.F. (2016) Spatial and temporal variability in pesticide exposure

downstream of a heavily irrigated cropping area: application of different monitoring

techniques. *Journal of Agricultural and Food Chemistry,* **64,** 3975-3989.

Pen, I., Uller, T., Feldmeyer, B., Harts, A., While, G.M. & Wapstra, E. (2010) Climate-driven

population divergence in sex-determining systems. *Nature,* **468,** 436.

Peters, E.J. & Parham, J.E. (2008) Ecology and management of sturgeon in the lower Platte

River, Nebraska. *Nebraska Technical Series*.

Peterson, C.H., Rice, S.D., Short, J.W., Esler, D., Bodkin, J.L., Ballachey, B.E. & Irons, D.B.

(2003) Long-term ecosystem response to the Exxon Valdez oil spill. *Science,* **302,** 2082.

Rashleigh, B. & Grossman, G.D. (2005) An individual-based simulation model for mottled

sculpin (*Cottus bairdi*) in a southern Appalachian stream. *Ecological Modelling,* **187,**

247-258.

Rennie, M.D., Purchase, C.F., Lester, N., Collins, N.C., Shuter, B.J. & Abrams, P.A. (2008)

Lazy males? Bioenergetic differences in energy acquisition and metabolism help to

explain sexual size dimorphism in percids. *Journal of Animal Ecology,* **77,** 916-926.

Ricker, W.E. (1975) *Computation and interpretation of biological statistics for fish populations*.

Fisheries Research Board of Canada, Ottawa, Ontario.

Scheffer, M., Bascompte, J., Brock, W.A., Brovkin, V., Carpenter, S.R., Dakos, V., Held, H.,

Van Nes, E.H., Rietkerk, M. & Sugihara, G.J.N. (2009) Early-warning signals for critical

transitions. **461,** 53.

Scheffer, M., Baveco, J., DeAngelis, D., Rose, K. & Van Nes, E. (1995) Super-individuals a

simple solution for modelling large populations on an individual basis. *Ecological*

modelling, **80,** 161-170.

Scheffer, M., Carpenter, S., Foley, J.A., Folke, C. & Walker, B. (2001) Catastrophic shifts in

ecosystems. *Nature,* **413,** 591.

Tilman, D. (1999) Global environmental impacts of agricultural expansion: the need for

sustainable and efficient practices. *Proceedings of the National Academy of Sciences,* **96,**

5995-6000.

Tilman, D., Fargione, J., Wolff, B., D'antonio, C., Dobson, A., Howarth, R., Schindler, D.,

Schlesinger, W.H., Simberloff, D. & Swackhamer, D. (2001) Forecasting agriculturally

driven global environmental change. *science,* **292,** 281-284.

Van Nes, E.H. & Scheffer, M. (2007) Slow recovery from perturbations as a generic indicator of

a nearby catastrophic shift. *The American Naturalist,* **169,** 738-747.

Wedekind, C. (2017) Demographic and genetic consequences of disturbed sex determination.

Philosophical Transactions of the Royal Society B: Biological Sciences, **372,** 20160326.

Weimerskirch, H., Lallemand, J. & Martin, J. (2005) Population sex ratio variation in a

monogamous long-lived bird, the wandering albatross. *Journal of Animal Ecology,* **74,**

285-291.

Welcomme, R.L. (1995) Relationships between fisheries and the integrity of river systems.

Regulated Rivers: Research & Management, **11,** 121-136.

Wildhaber, M., Papoulias, D., DeLonay, A., Tillitt, D., Bryan, J. & Annis, M. (2007a) Physical

and hormonal examination of Missouri River shovelnose sturgeon reproductive stage: a

reference guide. *Journal of Applied Ichthyology,* **23,** 382-401.

Wildhaber, M.L., Papoulias, D.M., DeLonay, A.J., Tillitt, D.E., Bryan, J.L. & Annis, M.L.

(2007b) Physical and hormonal examination of Missouri River shovelnose sturgeon

reproductive stage: a reference guide. *Journal of Applied Ichthyology,* **23,** 382-401.

Windsor, F.M., Ormerod, S.J. & Tyler, C.R. (2018) Endocrine disruption in aquatic systems: up-

scaling research to address ecological consequences. *Biological Reviews,* **93,** 626-641.

Figures

(a)

(b)

Fig. 1. Study system and model structure. a) a map of the study system, Lower Platte River (Nebraska–NE, USA) and b) a schematic of the spatially explicit individual-based biophysical model for shovelnose sturgeon.

Fig. 2. Shovelnose sturgeon population projections during exposure and recovery under synthetic androgen scenarios – a) a baseline scenario, b) an energetic perturbation (reduced maternal investment in gonad development) scenario with three exposure levels (shown as biomass relative to the baseline scenario); c) a demographic perturbation (pre-recruit sex ratio skewness) scenario with three exposure levels; and d) a combined scenario of energetic and demographic perturbations with three exposure levels. Red solid, green dashed, and blue dotted lines indicate low, medium and high androgen exposure.

Fig. 3. Changes (%, compared to the baseline scenario) in life history traits of shovelnose sturgeon populations– under energetic perturbation scenarios: a) proportion of mature females in the reproductive cycle, b) reproductive female number, c) spawner biomass, d) recruit number, e) relationship between reproductive female number and spawner biomass or recruit number during exposure, f) relationship between reproductive female number and spawner biomass or recruit number during recovery; and demographic perturbation scenarios: g) adult sex ratio, h) reproductive female number, i) spawner biomass, j) recruit number, k) relationship between adult sex ratio and reproductive female number, spawner biomass, or recruit number during exposure, and l) relationship between adult sex ratio and reproductive female number, spawner biomass, or recruit number during recovery. In, a–d and g–j, red solid lines in box plots indicate mean values and black solid lines show 25th, 50th, and 75th percentile (bottom to top). In e, f, k, and l, data points show mean values from low, moderate, and high androgen exposure levels.

Fig. 4. Changes (%, compared to the baseline scenario) in life history traits of shovelnose sturgeon populations under combined scenario of energetic and demographic perturbations – a) proportion of mature females in the reproductive cycle, b) adult sex ratio, c) reproductive female number, d) spawner biomass, e) recruit number, f) relationship between reproductive female number and spawner biomass or recruit number during exposure, g) relationship between reproductive female number and spawner biomass or recruit number during recovery, h) relationship between adult sex ratio and reproductive female number, spawner biomass, or recruit number during exposure, and i) relationship between adult sex ratio and reproductive female number, spawner biomass, or recruit number during recovery. In, a–e, red solid lines in box plots show mean values and black solid lines indicate 25th, 50th, and 75th percentile (bottom to top). In f–i, data points show mean values from low, moderate, and high androgen exposure levels.