Low-disturbance Farming Regenerates Healthy Deep Soil towards Sustainable

Agriculture

Fangbo Deng[a, b, c#], Hongjun Wang[d#], Hongtu Xie[a, c#], Xuelian Bao[a, c], Hongbo He[a, c],

Xudong Zhang[a, c], Chao Liang[a, c*]

[a] Key Laboratory of Forest Ecology and Management, Institute of Applied Ecology, Chinese

Academy of Sciences, Shenyang 110016, China

[b] University of Chinese Academy of Sciences, Beijing 100049, China

[c] Key Lab of Conservation Tillage & Ecological Agriculture, Liaoning Province, Institute of

Applied Ecology, Chinese Academy of Sciences, Shenyang 110016, China

[d] Duke University Wetland Center, Nicholas School of the Environment, Duke University, Durham,

NC 27708, USA

[*]Corresponding author

Email: cliang823@gmail.com

[#]These authors contributed equally to this work.

Abstract

Intensive conventional farming has degraded farmland topsoil and seriously threaten food and environment security globally. Although low-disturbance practices have been widely adapted to restore soil health, whether this measure in a long run can potentially recover the critical deep soil to meet sustainable intensification of crop production are still unclear. Here we compared soil microbiome, physicochemical parameters along 3-m deep soil profiles, and crop yield in Northeast China subjected to ten years of farming practices at 3 levels of disturbance, including conventional tillage (CT), no-tillage without stover mulching (NTNS), and no-tillage with stover mulching (NTSM). We found that low-disturbance practices (NTNS and NTSM) promoted the ability of the deep soil to retain water, nitrogen and salt-extractable organic, regenerated whole-soil microbial diversity and metabolic function, improved topsoil organic carbon stock and corn yield in the drought year, showed the potential to reduce energy consumption and greenhouse gas emissions, thus regenerating highly efficient, sustainable agriculture.

Keywords: sustainable agriculture, no-tillage, stover mulching, microbial diversity and function, deep soil.

Introduction

Since the Industrial Revolution, the rate of soil carbon loss has increased dramatically, resulting in a global carbon debt due to agriculture of 116 Pg carbon for the top 2 m of soil(Sanderman *et al.*, 2017). The loss of soil carbon in farmlands has not only changed global climate but also produced catastrophic cascade impacts on global food and environment security, as soil carbon is the cornerstone for healthy and productive soil that will be needed to sustainably feed 10 billion people in 2050 (United Nations, World Population Prospects 2019). It is well known that intensive conventional farms with high energy inputs (chemical fertilizers) and disturbance (e.g. tillage/compaction, burn/remove stover) have caused a series of environmental issues, like water pollution, biodiversity loss, freshwater depletion and climate change(Congreves *et al.*, 2015). Even worse, increasing the amount of chemical fertilizer is unlikely to continue the increase in quantity and quality of food products worldwide(Vitousek *et al.*, 2009), while the options to expand the farmland area at the expense of nature and biodiversity that already under pressure is limited(Springmann *et al.*, 2018). Moreover, the topsoil disturbance, tillage in particular, prevents root growth into deeper soil(Kemper *et al.*, 2011), thus critically minimizing the exploitation of deep soil profile(Billings *et al.*, 2018) and reducing crop's nutrient using efficiency and its resilience to drought.

Since the 1970s, low-disturbance practices (e.g. reduced tillage, no-tillage and stover mulching) have been gradually applied to restore soil health and reduce non-point source pollution(Salem). Growing evidence shows that no-tillage and stover mulching boosted top-soil organic carbon (SOC)(Blanco-Canqui and Lal, 2008; Liu *et al.*, 2014), increased soil aggregate stability(Song *et al.*, 2019) and reduced soil erosion and surface runoff(Prosdocimi *et al.*, 2016). All these benefits from

low-disturbance tillage practices are tied with complex microbial processes that

contribute to soil structure formation, modulate the biogeochemistry cycling of

nutrients and drive soil carbon transformation and stabilization(Schimel and Schaeffer,

2012; Bardgett and Van Der Putten, 2014; Liang *et al.*, 2017). Microorganisms also

respond quickly to changes in soil environment conditions, due to their high surface to

volume ratio, which can provide an early signal in soil improvement or degradation,

thus has been recommend as a key indicator of soil health(Nielsen *et al.*, 2002).

However, the effect of tillage practices on soil microbial communities are complex

and diverse(Helgason *et al.*, 2010; Sun *et al.*, 2018; Li *et al.*, 2020; Rincon-Florez *et

al.*, 2020) and most studies by now mainly focused on farmland topsoil or soils within

1-m depth(Hartman *et al.*, 2018; Nevins *et al.*, 2018; Alahmad *et al.*, 2019), and soil

below 1 meter, which contains more carbon than the topsoil(Jobbágy and Jackson,

2000) and belongs to Earth's Critical zone, was often overlooked. However, microbes

inhabiting in the deep soils (> 1 m) may substantially impact long-term carbon

sequestration, mineral weathering and crop production(Eilers *et al.*, 2012;

Sagova-Mareckova *et al.*, 2016; Pries *et al.*, 2017), and play important roles in

bridging aboveground vegetation with parent soils and even acts as an essential buffer

protecting underground water(Chorover *et al.*, 2007). Moreover, studies indicate

many crops' roots (depending on the species and management) can penetrate over one

meter in depth(Stone *et al.*, 2001; Thorup-Kristensen, 2006), which means they can

potentially forage for nutrient and water in deep soil and impact deep-soil microbial

community(Pierret *et al.*, 2016; Thorup-Kristensen *et al.*, 2020). More importantly,

tillage practices significantly influence root growth(Qin *et al.*, 2004; Kemper *et al.*,

2011; Nunes *et al.*, 2019), the change of surface soil carbon by land use or agricultural

practice influenced the carbon dynamic in deep soil layers has also been

observed(Fontaine *et al.*, 2007; Tautges *et al.*, 2019; Dal Ferro *et al.*, 2020). We thus

hypothesized that: 1) different tillage practices will affect the vertical distribution of

the microbial community by altering the spatial distribution of nutrients and dissolved

organic carbon, thereby changing the ecosystem function; 2) lowest disturbance

practice—no-tillage with 100% stover mulching as a nature-based solution, close to

undisturbed natural ecosystem, would regenerate healthy deep-soil with highly

diversified and functional microbes over time toward a highly resilient, sustainable

agricultural ecosystem.

Recent research shows that corn belts in the U.S.A., western Europe, and China

have experienced the most soil carbon loss globally(Sanderman *et al.*, 2017). The

corn belt in Northeast China is considered as the "breadbasket" of the country, having

the largest grain production and overlapping with the most fertile Mollisol region that

sustains 3% of population in the world(Liu *et al.*, 2010), accounting for over 30% of

corn production of China(Liu *et al.*, 2012). Here, a 10-year manipulative experiment

was conducted at a temperate corn farm in Northeast China, investigating farming

practices with three levels of disturbance: high disturbance—conventional tillage

(CT), low disturbance—no-tillage without stover mulching (NTNS) and lowest

disturbance—no-tillage with 100% stover mulching (NTSM) (described in the

Methods and Fig. S1). We compared soil physicochemical properties, fine-root

associated organic carbon, and microbial communities of the 3-m soil profiles at the

end of dormant season (legacy effects of nutrients) after the 10-year manipulation, and

multi-year corn yield as well. We aimed at testing our hypotheses and providing a

general picture of soil microbial community in the entire soil profiles, which are

important for comprehensive recognize the impact of low-disturbing tillage practices

on ecosystem function and developing effective strategies to ensure the sustainable

development of agriculture.

Results

Soil properties and corn yield

The SOC, TN and C/N ratio substantially decreased from the soil surface to

around 150 cm depths and then remained unchanged within 150-300 cm (Fig. 1). The

NTSM slightly increased SOC, TN and C/N ratio at 0-20 cm soil layers compared

with the NTNS and the CT (Fig. 1 and Table S1). The NTSM and NTNS reduced soil

pH in surface and deeper layers (Fig. 1d) and increased soil moisture at surface layers

(0-60 cm) (Fig. 1e). In the CT plots, soil NO_3^--N concentration first decreased and

then increased remarkably, ranged from 4.19 to 23.32 mg kg^{-1} (Fig. 1i). However,

under the NTNS and NTSM treatments, soil NO_3^--N decreased significantly at 0-40

cm then increased to the maximum at 120-150 cm. Interestingly, above 120-150 cm

layer, NO_3^--N was significantly higher with low-disturbance practices than

conventional tillage, while the soil below 150 cm under low-disturbance practices had

much lower NO_3^--N compared to conventional tillage (Fig. 1i). The NTNS plots

contained much higher amounts of ammonium than the CT and the NTSM plots (Fig.

1h). Soil salt-extractable organic carbon (SEOC)—a proxy for biotically-derived

organic acid declined from the surface to 40-60 cm and then increase to its peak at

60-90 cm under CT, at 90-120 cm under NTNS and at 120-150 cm under NTSM (Fig.

1b). The NTSM increased the SEOC concentration at almost all soil layers compared

with the CT and the NTNS (Fig. 1b), in which at the surface and 120-150 cm depth

the contents of SEOC with NTSM were twice higher than CT. The relative

contributions of SEOC to SOC (SEOC/SOC) in the NTSM were also always higher

than in the CT and NTNS (Fig. 1c). Based on the estimated root depths and soil bulk

density, total soil inorganic nitrogen available for the coming growing season in the

NTSM and the NTNS was approximate 427.34 and 352.34 kg ha^{-1}, respectively, while

only 179.63 kg ha^{-1} in conventional tillage.

The mean annual corn yield (2013-2016) in the NTSM is 13416.8 kg/ha, which is

much higher than the CT and NTNS (Fig. 2), particularly during the drought year of

2015, with only 409.6 mm of rainfall during the growing season (about 100 mm lower

than the mean rainfall), while the corn yield in NTSM is 36.4% and 22.3% higher

than the CT and NTNS, respectively (Fig. 2).

Microbial diversity, composition, and structure

The microbial richness (Chao1), observed number of species (Observed-species)

and diversity (Shannon-Index) first increased within 0-20 cm and decreased from 20

to 90 cm, then increased hereafter (Fig. 3). The low-disturbance practices significantly

increased Chao1, Observed-species and Shannon-Index, particularly in 0-40 cm soil

depths (Fig. 3). There were 54 microbial phyla across all soil samples. The dominant

phyla (relative abundance > 1% across all soil samples) were Proteobacteria,

Actinobacteria, Chloroflexi, Acidobacteria, Nitrospirae, Gemmatimonadetes,

Planctomycetes, and these phyla accounted for 60-91% of the total microbial

abundances in the whole soil profile (Fig. S2a). Bacteroidetes, Verrucomicrobia,

Latescibacteria, Parcubacteria, Firmicutes, Microgenomates and Saccharibacteria

were less dominant (relative abundance > 0.1% across all soil samples) but were still

found across all soil samples (Fig. S2a). Although no difference in the composition of

dominant phyla among treatments was found, there were more non-dominant phyla

with higher relative abundance in low disturbance practices than conventional tillage

practice (Fig. S2b).

Indicator analysis identified 16 and 51 clearly classified genera (relative

abundances > 0.005%) in the NTNS and the NTSM plots, respectively, while no

indicator genera were found in the conventional tillage plots (Fig. 4 and Table S2).

The indicator genera in the NTNS plots belonged to Proteobacteria, Actinobacteria,

Chloroflexi, Gemmatimonadetes and Planctomycetes, and most of them appeared in

the surface soil (0-20 cm) with only 1 genus below 150 cm. Importantly, more extra

indicator genera — including Bacteroidetes, Acidobacteria, Deferribacteres,

Firmicutes, Verrucomicrobia, Chlorobi and Spirochaetae — existed in the NTSM

plots, in which under 150 cm we observed 7 genera (Fig. 4 and Table S2).

Microbial community structures were visualized by Non-metric multidimensional

scaling (MDS) and tested by PERMANOVA based on Bray–Curtis. The microbial

communities among treatments in the root zones were marginally different

(PERMANOVA p=0.08); however, below the root zone they differed distinctively

(PERMANOVA p=0.02). The disturbance practices influenced the vertical

distribution dissimilarity in microbial community structure (Fig. 5). Three clusters —

0-10 cm and 10-20 cm, 20-150 cm and 150-300 cm — were observed in the CT plots

(PERMANOVA-F=9.57, p=0.0001) (Fig. 5). In the NTNS plots, 0-10 cm formed an

independent cluster, while other soil depths showed some separation (e.g. 20-120 cm

were separated from 150-300 cm soil depths by axis 1); however, Bray-Curtis

distances between adjacent depths were too close to be separated

(PERMANOVA-F=8.18, p=0.0001) (Fig. 5). The NTSM treatment clustered 0-10 cm

and 10-20 cm together, 120-150 cm, 150-200 cm, 200-250 cm and 250-300 cm

separately, and the other depths show some separations as well

(PERMANOVA-F=11.32, p=0.0001) (Fig. 5).

Predicted Ecological functions of microbial communities

According to the results of microbial diversity, composition and structure, the

metabolic capabilities of microbial community in the whole 3-m soil profiles were

predicted using Tax4Fun (Fig. S3). Results showed that low-disturbance practices

significantly increased the abundance of predicted functions related to carbohydrate

metabolism, nucleotide metabolism, glycan biosynthesis and metabolism, lipid

metabolism and metabolism related to cofactors and vitamins (Fig. S3a). Moreover,

the relative abundances of genes encoding for assimilatory nitrate reduction in

low-disturbance practices were higher than that in conventional tillage practice (Fig.

S4). The results suggested that in low disturbance practices, microbial community

prefer to convert the nitrate/nitrite to ammonia. We then further assessed the impact of

stover mulching on functional profiles (Fig. S3b). The extended error bar plot shows

that the NTNS enriched the abundance of amino acid metabolism and lipid

metabolism, while the NTSM enriched the functions associatedto energy metabolism,

carbohydrate metabolism, biosynthesis of secondary metabolites, glycan biosynthesis

and metabolism as well as metabolism of cofactors and vitamins (Fig. S3b).

Relationships between microbial communities and soil properties

Forward selection in Redundancy analysis (RDA) revealed that soil depth

(pseudo-F=48, p= 0.002), SOC (pseudo-F=11.5, p= 0.002), SM (pseudo-F=3.4, p=

0.012), soil pH (pseudo-F=2.3, p=0.018) and soil NH_4^+-N (pseudo-F=2.7, p= 0.026)

significantly affected the vertical distribution of microbial communities (Fig. S5).

Furthermore, the soil properties that regulated the distribution of soil microbes were

different under different disturbance practices. Under the CT treatment, soil microbial

community was mainly affected by soil NH_4^+-N (pseudo-F=4, p= 0.002) and soil

NO_3^--N (pseudo-F=2.3, p= 0.012) that mainly came from applied fertilizer (Fig. 6).

The microbial community positively correlated to soil NH_4^+-N in the 0-20 cm soil, to

soil NO_3^--N negatively within 20-150 cm, while to soil NO_3^--N positively after 150

cm (Fig. 6). Under the NTNS treatment, soil pH (pseudo-F=3.7, p=0.004) constrained the distribution of the microbial community, in which strong negative correlations occurred in 0-10 cm soil and a positive correlation in 90-150 cm (Fig. 6). Under the NTSM treatment, soil TN (pseudo-F=11, p=0.002), SM (pseudo-F=2.6, p=0.004) and C/N ratio (pseudo-F=1.8, p=0.016) significantly influenced the soil microbial community separation (Fig. 6). In general, the microbes positively correlated with the soil TN and C/N ratio in the surface soil layers (0-40 cm) and with SM in the middle layers (40-150 cm), while they were mainly influenced by depth in the deeper soil (150-300 cm) (Fig. 6).

Root depth among tillage practices

Currently the measurement of fine roots with diameter less than 0.2 mm are still technically difficult and their high turnover rates in-situ make the measurement even more complex(Pierret *et al.*, 2016). Instead of detecting fine roots directly, we measured SEOC—a proxy for biotically-derived organic acid, which is a sensitive signal of root density and could be an indicator of root depth(Billings *et al.*, 2018). We also did a literature review on whether no-tillage and straw mulching extend root depth in other crops. According to the change of SEOC with soil depth under different disturbing practices(Fig. 1b), we estimate that corn roots reached up to 60-90 cm, 90-120 cm and 120-150 cm under the CT, the NTNS and the NTSM, respectively, which is in line with reported corn root depths (~150 cm) (Canadell *et al.*, 1996; Kemper *et al.*, 2011).According the literature review, we can't find root-depth studies including both no-tillage and straw mulching. However, no-tillage as a conservative management increases root depth by 23 cm, 39 cm and 14 cm in corn, wheat, and sunflower, respectively (Fig. S6).

Discussion

No-tillage practices promote soil nutrient, water holding capacities and corn yield

The changes of SEOC with soil depth and the literature review of root depth among tillage indicated that no-tillage could promote root growth into deep soil, up to 150 cm in the NTSM. The root exudates with various organic acid and dead roots likely contributed to the lower soil pH in the NTSM, as shown by a significant negative relationship between SEOC and pH (r=–0.678, p<0.05). Despite function predication analyses have various uncertainty, the increased dissimilatory and assimilatory nitrate reduction genes suggested that low disturbance practices tend to convert nitrate to ammonium (Fig. S4), while lower ammonium in NTSM may be due to ammonium subsequently incorporated into biomass, as hints by the increase in genes related to biosynthesis metabolism and glycometabolism (Fig. S3). Moreover, maize stover or increased fine roots in deeper soil in NTSM provided labile carbon (Fig. 1b) to remove leaked nitrate through denitrification in deeper soil (below 1.5 m), where higher relative abundance of the denitrification bacteria (*Pseudomonas* and *Caldithrix*)(Koike and Hattori, 1975; Miroshnichenko *et al.*, 2003) (Fig. 4 and Table S2) were detected. However, shallower roots in the CT treatment can't provide enough labile carbon to remove extra soil NO_3^--N in deep soil, thus causing nitrite accumulation and leaching into deeper soil layers. The amount of inorganic nitrogen accumulated in the root zones under NTSM (427.34 kg ha^{-1}) likely could provide plenty of nitrogen for corn growth in the coming growing season (Fig. 2), based on the removed nitrogen in the grain (~200 kg ha^{-1}). Additionally, in line with many studies that show stover mulching reduces water evaporation and surface runoff and increase soil moisture in top soils(De Vita *et al.*, 2007; Prosdocimi *et al.*, 2016), we

found that the topsoil moisture was significantly higher in the NTSM than in the CT

plots (Fig. 1e) when the precipitation during the entire dormant period was only 38.5

mm. More importantly, the topsoil (0-20 cm) SOC stock in NTSM (18.5 Mg/ha) were

significantly higher than in CT (15.8 Mg/ha) due to slightly higher bulk density in

NTSM. Therefore, no-tillage with stover mulching not only increased SOC stock, the

holding capacities for nutrients and water, thus reducing energy input to farm, but also

tended to reduce the risk of nitrate leaching into groundwater. And more importantly,

the healthy deep soil in turn raises corn production and promote the crop resistance to

drought (Fig. 2). All these are critical to the development of sustainable agriculture

and the associated ecosystems.

No-tillage with stover mulching promotes microbial diversity, richness, and

ecological function contributing to sustainable farming

Under the CT treatment, tillage heavily disturbed the topsoil and liberated occluded

organic materials. Microbes tended to rapidly use available nutrients in the plowed

layer (e.g. NH_4^+-N)(Ramirez-Villanueva *et al.*, 2015), thereby causing the reduction

of microbial metabolic diversity (Fig. S3a). Then, the resistance of the soil to stress or

disturbance may also decrease(Kremen, 2005). In deeper soil layers, due to shallower

roots, NO_3^--N could quickly move downward and accumulate in deeper soil (Fig. 1i),

which not only contaminated the underground water but also limited the activity of

non-dominant microbes with important ecological functions, as no indicator genera

were identified for each soil depth in CT treatment (Fig. 4 and Table S2). Because the

microbial communities were closely associated with inorganic nitrogen, the microbes

under CT were mainly influenced by added chemical fertilizer(Wood *et al.*, 2015).

Although the dominant microbial communities in CT were similar to those in the

NTNS and NTSM, the loss of function resulted from the difference of non-dominant

microbes, indicating that the soil under CT had degraded.

Under the NTNS treatment, soil pH was the major edaphic factor affecting the

microbial community and the indicator genera (Fig. 6 and Table S3). The lower soil

pH possibly was caused by deeper roots as shown by higher SEOC that is generally

positively related to root density(Billings *et al.*, 2018). Soil pH is often observed as a

major factor determining the microbial composition and structure in natural

ecosystems(Lauber *et al.*, 2009; Zhalnina *et al.*, 2015), as microbes often show a

narrow tolerance to soil pH. In addition, soil pH regulates the availability of nutrient

and mitigate ion toxicity(Lauber *et al.*, 2009; Rousk *et al.*, 2010; Zhalnina *et al.*,

2015). Under NTNS, soil pH and depth explained 35% distribution of the microbial

community (Fig. 6). We speculated that other edaphic factors (e.g. salinity and iron)

directly or indirectly related to soil pH and SEOC also influenced the changes in the

microbial community.

Under NTSM treatment, TN and C/N significantly correlated with soil microbial

community due to the high C/N ratio of stover or roots (Fig. 6). Prior studies have

reported that, following maize stover mulching, more organic N, amino acid N, and

amino sugar N were observed in soil(Liu *et al.*, 2016; Lu *et al.*, 2018), combining

with higher genes related to amino acid metabolism and carbohydrate metabolism

from function predication analyses (Fig. S3), jointly hint that NTSM increased the

retention time of nitrogen, hence meeting the nutrient requirement of corn growth and

reducing nitrate loss to underground water. The increased available nitrogen, labile

carbon and water in deep soil under NTSM can increase the resilience and resistance

of maize to disturbances with higher grain production (Fig. 2). Zhang et al.(Zhang *et*

al., 2013a) also observed litter-covered soil showed greater resistance to heating and

copper addition due to the changes in soil properties and microbial community

structure. Resistance to disturbance or stresses is the nature of a healthy soil and is

essential for maintaining ecosystem functions, such as decomposing organic

matter(Kibblewhite *et al.*, 2008; Zhang *et al.*, 2013a). Under the NTSM treatment, the

microorganisms associated with the degradation of relatively stable carbon

compounds, such as Planctomycetes and Verrucomicrobia (Table S4)(Herlemann *et*

al., 2013; Erbilgin *et al.*, 2014) as well as the indicator *Cellulomonas* and

Azospirillum (Fig. 4 and Table S2) with the function of cellulose

decomposition(Pathma *et al.*; Halsall and Goodchild, 1986) were increased. The

predicted functional profiles related to energy metabolism (Carbon fixation pathways

in prokaryotes), carbohydrate metabolism (TCA cycle, amino sugar, nucleotide sugar,

galactose, fructose), biosynthesis of secondary metabolites (Carotenoid and Betalain)

and glycan biosynthesis were increased, suggesting a higher metabolic activity and a

change in substrate quality (Fig. S3). In addition, stover mulching also increased the

ecological filter function of soil depth for selecting microbial communities as more

indicator genera of each soil depths were identified under NTSM compared to NTNS

and CT practices (Fig. 4 and Table S2). And these indicators residing at different soil

depths might enhance the anti-disturbance ability of NTSM. For example,

denitrification bacteria *Caldithrix* and *Pseudomonas*(Koike and Hattori, 1975;

Miroshnichenko *et al.*, 2003) were the indicator genera of 150-200 cm and 250-300

cm, respectively (Fig. 4 and Table S2), which might explain the low nitrate in the

deep soil in NTSM. *Ignavibacteria* and *Spirochaeta,* the indicator genera of deep soil,

have the ability to grow under the conditions of strictly anaerobic(Iino *et al.*, 2010)

and severely limited nutrients(Terracciano and Canale-Parola, 1984), respectively.

Surface indicator genera belonging to Bacteroidetes might have the ability to degrade

organic matter that is difficult to decompose(Thomas *et al.*, 2011).

Implications for climate change and food security

It was observed that about 179.63, 352.34 and 427.34 kg ha^{-1} inorganic N were kept in the root-zone soil in the CT, NTNS and NTSM, respectively. Generally, corn roots reach their maximum depth at the silking stage(Archontoulis and Licht, 2017), which is also the time when the heaviest rainfall occurs in northeastern China. We therefore expect that the available N kept in the root zone would be utilized by crops in the coming growing season before the storm leach the nitrogen down to ground water, which means that fertilizer N could be cut to meet crop growth in, at least, Northeast China and also prevent reactive N losses. Since the nitrogen use efficiency (NUE) of maize system under the conventional management is 51% in northeast China (NUE is defined as the efficiency of fertilizer N transferring to harvested crop N)(Zhang *et al.*, 2019). Then, we conservatively calculate the required fertilizer N in the next year based on two assumptions: 1) the NUE of soil available N in root zone is equal to that NUE of applied fertilizer N, both of them are 50%; 2) the mineralized N during the coming growing season is neglected. Thus, N supply requirement = Fertilizer N×NUE + N in root zone ×NUE + Stover-N, where Stover-N for NTSM is 60 kg ha^{-1}. We estimated the N requirement for each disturbance practice by multiplying grain yield by grain N concentration (1.4%)(Zhang *et al.*, 2019) plus multiplying stover yield by stover N concentration (0.8%)(Izewska and Woloszyk, 2015). For CT, NTNS and NTSM, the mean annual corn yields were 10946.74, 12487.81 and 13416.81 kg ha^{-1}, and the stover yields were 966.67, 10083.33 and 10833.33 kg ha^{-1}, respectively. Thus, the N requirements were 230.6, 255.5 and 274.5 kg ha^{-1} for CT, NTNS and NTSM, respectively. Therefore, the theoretically conservative amounts of fertilizer N in the coming growing season are 281.6, 158.7 and 1.7 kg ha^{-1} for CT, NTNS and NTSM, respectively. No fertilizer-N is needed to apply without reducing corn yield in the

NTSM plot. Compared to CT, the NTNS and NTSM could at least save respectively

about 122.9 and 281.6 kg ha^{-1} N-fertilizer. For every kilogram of fertilizer-N produced

and used on cropland, up to 87.9 MJ of energy is consumed(Kennedy, 2000) and 13.5

kg of CO_2-equivalent (eq) (CO_2-eq) is emitted(Zhang *et al.*, 2013b). Hence, totally

24,752.6 MJ of energy consumption could be reduced and 3,801.6 kg CO_2-eq

emission could be cut per hectare cornland in Northeast China at least by using

NTSM tillage practice. If this could be applied to all maize farmland in Northeast

China (13,000,000 ha, Source: China Statistics Yearbook 2018), 0.3 EJ of energy

could be saved and 49.4 Mt of CO_2-eq could be reduced. Based on the average annual

energy consumption for households of China in 2017 (15 EJ, China Statistics

Yearbook 2018) and CO_2 emissions (9,839 Mt, Global Carbon Atlas), the NTSM

practice in corn farming of Northeast China has the potential to save 2% of household

energy and to reduce 0.5% of CO_2 emissions each year in the whole country.

Our results, particularly higher SEOC content, microbial diversity, and indicator

genera in the NTSM deep soil compared with CT, clearly showed that

low-disturbance practices can affect agricultural resource in deeper soil over time.

This indicates that crops, like corn in this study, under appropriate management can

ultimately explore nutrients and water from deeper soil, thus not only increasing the

volume of soil (almost double in this study) exploited without reclaiming more natural

land areas but also reducing nutrient loss into ground water. Meanwhile, the input of

labile carbon including SEOC into deeper soil in the NTSM provides essential energy

and nutrients to microbes and gradually shape highly diversified and functional

microbial communities in the deep soil over time, and hence improve the

self-sustaining ability of farmland in the face of climate change. The improvements of

microbial communities in the deep soil (1-3 m) at the end of dormant season in our

study provide evidence for the first time that a nature-based management in farmland

is conducive to deep-soil health for sustainable farm in a long run. Although many

scientists have realized the importance of deep rooting for sustainable intensification

of crop production(Pierret *et al.*, 2016; Thorup-Kristensen *et al.*, 2020), the deep-root

studies are still rare due to technological bottleneck. According the literature review,

no-tillage extend root depth in other crops, like corn, wheat and sunflower. While

some studies also show no effects or even reduce root depth(Dwyer *et al.*, 1996;

Izumi *et al.*, 2004), the possible reason is the legacy effects of tillage, for example,

long-term no-tillage leads to soil compaction, and the soil system might be still in its

transition stage. Coupled with our results, low-disturbance practices as a nature-based

agricultural management likely can develop a deeper root system to explore more

resource in deep soil to sustain food production, while without threatening

environmental security and reclaiming more lands.

Materials and Methods

Site description and soil sampling

The field experiment was established in 2007 at the Lishu Conservation Tillage

Research and Development Station of the Chinese Academy of Sciences in Jilin

province, Northeast China (43.19° N, 124.14° E). The region has a humid continental

climate with a mean annual temperature of 6.9 °C and the mean annual precipitation

of 614 mm. The soils are classified in the Mollisol order (Black Soil in Chinese Soil

Classification) with a clay loam texture(IUSS Working Group, 2006). The site has

been continuously planted with maize since 2007. We set up an experiment by a

randomized complete block design with four replicates and five treatments. Each plot

area was $261m^2$ (8.7×30m). The five treatments included conventional tillage

408 (moldboard plowing to a depth around 30 cm and removed the stover), no-tillage (no

409 soil disturbance and direct seeding), and no-tillage with three-level stover mulching

410 (33%, 67% and 100% newly produced maize stover were evenly spread over the soil

411 surface each fall) (Fig. S1). For each treatment, slow-release fertilizer was applied at

412 one time when sowing, which was equal to 240 kg/ha N; 47 kg/ha P; 90 kg/ha K. The

413 rainfall data were obtained from local meteorological administration. The grain yield

414 was estimated by manually harvesting 20 m^2 area, randomly taken from each plot.

415 In this experiment, in order to reduce the damage to the plots and reduce costs, 3

416 plots were randomly taken from each treatment including conventional tillage (CT),

417 no-tillage without stover mulching (NTNS), no-tillage with 100% stover coverage

418 (NTSM) as three comparative practices. In April 2017, triplicate soil cores (0-300 cm)

419 were collected from each plot at the end of dormant season. After removing surface

420 stover, we took soil cores by a stainless-steel hand auger and sliced each into ten

421 layers: 0-10 cm, 10-20 cm, 20-40 cm, 40-60 cm, 60-90 cm, 90-120 cm, 120-150 cm,

422 150-200 cm, 200-250 cm, 250-300 cm. In total, 90 soil samples were collected and

423 transported to the laboratory within 3 hours, then passed through a 2-mm sieve. All

424 visible roots, crop residues and stones were removed. Each soil sample was divided

425 into three subsamples: one subsample for DNA extraction and soil salt-extractable

426 organic carbon (SEOC) measurement that was immediately placed into a polyethylene

427 plastic bag and stored at -80 ℃, one for chemical measurements including ammonium

428 nitrogen (NH_4^+-N) and nitrate nitrogen (NO_3^--N) (within one day), and the remaining

429 one was air dried for other soil physicochemical properties.

Soil properties

431 Soil total nitrogen (TN) content was measured by an Element analyzer Vario EL

432 III (Elementar Analysensysteme GmbH, Hanau, Germany). Soil organic carbon (SOC)

was converted from soil organic matter that was measured by potassium dichromate

oxidation(Nelson and Sommers, 1982). Soil pH was measured in deionized free-CO_2

water (1:2.5 w/v). Gravimetric soil moisture was determined by oven-drying fresh soil

to a constant weight at 105 °C. Soil NH_4^+-N and NO_3^--N were extracted from fresh

soil by 2 M KCl and measured by a continuous flow analytical system (AA3, SEAI,

Germany). To reflect soil soluble, exchangeable, mineral-bound OC, SEOC was

extracted from the frozen soil samples with 0.5 M K_2SO_4 (1:5 w/v)(Jones and Willett,

2006; Toosi *et al.*, 2012).

DNA extraction, PCR amplification and pyrosequencing

Soil DNA was extracted from the frozen soil samples (0.5 g wet weight) by using

MoBio PowerSoil DNA isolation kit (MoBio Laboratories, Carlsbad, CA, USA)

following the instructions of the manufacturer. The quality of DNA was determined

by 1% agarose gel electrophoresis. The V3–V4 region of the bacterial 16S rRNA gene

was amplified by PCR using the primers 338F and 806R with barcode for Illumina

MiSeq sequencing. PCR was performed in a total volume of 50 μl containing 30 ng

DNA as a template, 20 mol of each primer, 10mM dNTPs, 5μl Pyrobest buffer (10×)

and 0.3 U of Pyrobest polymerase (Takara Code: DR005A). Each sample was

amplified for three replicates. The PCR products from the same sample were pooled,

checked by 2% agarose gel electrophoresis and were then purified using

AxyPrepDNA agarose purification kit (AXYGEN). Finally, purified PCR products

were sequenced on an Illumina MiSeq platform PE300 sequencer (Illumina, USA).

The raw sequence data were further analyzed by the following protocol.

Low-quality sequences with an average quality score of less than 20 were filtered by

employing Trimmomatic(Bolger *et al.*, 2014). The FLASH software was used to

merge overlapping ends and treat them as single-end reads(Derakhshani *et al.*, 2016).

The non-amplified region sequences, chimeras and shorter tags were also removed

using Usearch and Mothur(Mysara *et al.*, 2016). The resulting high-quality sequences

were clustered into Operational Taxonomic Units (OTUs) at 97% sequence similarity

using Usearch (Version 8.1.1861 http://www.drive5.com/usearch/). OTUs were then

classified against the Silva (Release119 http://www.arb-silva.de) database and the

taxonomic information of each OTU representative sequence was annotated using the

RDP Classifier(Wang *et al.*, 2007). A total of 3,255,693 high-quality reads were

obtained from all soil samples, which were clustered into 9,573 unique OTUs at a 97%

sequence similarity. The Good's coverage of all the samples ranged from 0.93 to 0.98,

which indicates an adequate level of sequencing to identify the majority of diversity

in the samples.

Root depth data

Plant roots, fine roots in particular, release large amounts of labile organic

carbon(Dakora and Phillips, 2002) that are essential for healthy microorganisms in

soil. Generally, most studies considered that fine roots are roots < 2mm in diameter,

while roots <0.2 mm in diameter can contribute to >50 % of the overall root length

and play a major role in releasing root exudates and absorbing nutrients and

waters(Pierret *et al.*, 2016). Currently the measurement of fine roots with diameter

less than 0.2 mm are still technically difficult and their high turnover rates *in-situ*

make the measurement even more complex(Pierret *et al.*, 2016). To minimize the

root-turnover effects to the most degree, we collected soil samples at the end of

dormant season, which can likely mirror the long-term legacy effects of our practices.

Also instead of detecting fine roots directly, we measured soil salt-extractable organic

carbon (SEOC)—a proxy for biotically-derived organic acid, which is a sensitive

signal of root density and could be an indicator of root depth(Billings *et al.*, 2018).

We also did a literature review on whether no-tillage and straw mulching extend root

depth in other crops. We collected root depth of main crops data from literature

published in 1996-2019 using the Web of Science resource and only the data with

more than two tillage managements are retained. Finally, a total of 87 observations

were selected (Table S5).

Statistical analyses

Soil properties were analyzed and plotted using Sigmaplot 12.5 software. Alpha

diversity indices were calculated in Qiime (version v.1.8) and used to reflect the

diversity and richness of the microbial community in different samples. The relative

abundances of individual phyla in different samples were computed by R packages.

The indicator analysis based on genera-specific to each soil depth was conducted

using indicspecies package of R with 9999 permutations, and the P-values were

corrected for multiple testing using qvalue package of R(Cáceres and Legendre, 2009).

Functional profiles of the microbial community were predicted by Tax4fun (an

open-source package in R)(Aßhauer *et al.*, 2015) and further statistical analysis was

conducted by STAMP using Welch's t-test (Parks *et al.*, 2014). Non-metric

multidimensional scaling (MDS) was performed by "vegan" package of R to describe

differences in microbial community structure among samples. Permutational

multivariate analysis of variance (PERMANOVA) was employed on Bray-Curtis

distances to test the differences in soil microbial communities among various sample

groups. The redundancy analysis (RDA, Canoco 5 software) were conducted to

identify the correlations between microbial community composition and

environmental variables. ANOVA were conducted by SPSS Version 22. Percentage

data were transformed using arcsine square root function before ANOVA test. All

statistical tests were significant at $p \leq 0.05$.

Sequence availability: All sequencing data that support the findings of this study

have been deposited in the National Center for Biotechnology Information

(https://www.ncbi.nlm.nih.gov/), in the Sequence Read Archive (SRA) database

(BioProject number: PRJNA488172).

Acknowledgements

We would like to thank Dr. William H. Schlesinger at the Cary Institute of Ecosystem

Studies for his comments and Dr. Randy Neighbarger at Duke University for language

editing. This work was supported by the "National Key R&D Program" (No.

2016YFD0800103, No. 2016YFD0200307) and the National Natural Science

Foundation of China (grant number, 41671297). We would like to thank Pengshuai

Shao, Xuesong Ma and many individuals for assistance with sample collection,

processing and analysis.

Author Contributions

H.X., XZ and C.L. designed the experiment, F.D. did field and lab measurements,

F.D., H.W. and C.L analyzed data and wrote the manuscript, and all the authors

discussed results and commented on the manuscript.

Declaration of Interests

The authors declare no competing interests.

Supplementary information

Supplemental information may be found online in the Supplemental Information

section at the end of the article.

References

Alahmad, A., Decocq, G., Spicher, F., Kheirbeik, L., Kobaissi, A., Tetu, T., Dubois, F.,

Duclercq, J., 2019. Cover crops in arable lands increase functional complementarity and

redundancy of bacterial communities. J. Appl. Ecol. 56, 651-664.

Archontoulis, S., Licht, M.A., 2017. How Fast and Deep do Corn Roots Grow in Iowa?
Integrated Crop Management News, 2442.

Aßhauer, K.P., Wemheuer, B., Daniel, R., Meinicke, P., 2015. Tax4Fun: predicting functional
profiles from metagenomic 16S rRNA data. Bioinformatics 31, 2882-2884.

Bardgett, R.D., Van Der Putten, W.H., 2014. Belowground biodiversity and ecosystem
functioning. Nature 515, 505-511.

Billings, S.A., Hirmas, D., Sullivan, P.L., Lehmeier, C.A., Bagchi, S., Min, K., Brecheisen, Z.,
Hauser, E., Stair, R., Flournoy, R., 2018. Loss of deep roots limits biogenic agents of soil
development that are only partially restored by decades of forest regeneration. Elem Sci
Anth 6, 34.

Blanco-Canqui, H., Lal, R., 2008. No-tillage and soil-profile carbon sequestration: An
on-farm assessment. Soil Sci. Soc. Am. J. 72, 693-701.

Bolger, A.M., Lohse, M., Usadel, B., 2014. Trimmomatic: a flexible trimmer for Illumina
sequence data. Bioinformatics 30, 2114-2120.

Cáceres, M.D., Legendre, P., 2009. Associations between species and groups of sites: indices
and statistical inference. Ecology 90, 3566-3574.

Canadell, J., Jackson, R.B., Ehleringer, J.B., Mooney, H.A., Sala, O.E., Schulze, E.-D., 1996.
Maximum rooting depth of vegetation types at the global scale. Oecologia 108, 583-595.

Chorover, J., Kretzschmar, R., Garcia-Pichel, F., Sparks, D.L., 2007. Soil biogeochemical
processes within the critical zone. Elements 3, 321-326.

Congreves, K., Hayes, A., Verhallen, E., Van Eerd, L., 2015. Long-term impact of tillage and
crop rotation on soil health at four temperate agroecosystems. Soil Tillage Res. 152,
17-28.

Dakora, F.D., Phillips, D.A., 2002. Root exudates as mediators of mineral acquisition in
low-nutrient environments. Food Security in Nutrient-Stressed Environments: Exploiting
Plants' Genetic Capabilities. Springer, pp. 201-213.

Dal Ferro, N., Piccoli, I., Berti, A., Polese, R., Morari, F., 2020. Organic carbon storage
potential in deep agricultural soil layers: Evidence from long-term experiments in
northeast Italy. Agric., Ecosyst. Environ. 300, 106967.

De Vita, P., Di Paolo, E., Fecondo, G., Di Fonzo, N., Pisante, M., 2007. No-tillage and
conventional tillage effects on durum wheat yield, grain quality and soil moisture content
in southern Italy. Soil Tillage Res. 92, 69-78.

Derakhshani, H., Tun, H.M., Khafipour, E., 2016. An extended single-index multiplexed 16S
rRNA sequencing for microbial community analysis on MiSeq illumina platforms. J.
Basic Microbiol. 56, 321-326.

Dwyer, L., Ma, B., Stewart, D., Hayhoe, H., Balchin, D., Culley, J., McGovern, M., 1996.
Root mass distribution under conventional and conservation tillage. Can. J. Soil Sci. 76,
23-28.

Eilers, K.G., Debenport, S., Anderson, S., Fierer, N., 2012. Digging deeper to find unique
microbial communities: the strong effect of depth on the structure of bacterial and
archaeal communities in soil. Soil Biol. Biochem. 50, 58-65.

Erbilgin, O., McDonald, K.L., Kerfeld, C.A., 2014. Characterization of a planctomycetal
organelle: a novel bacterial microcompartment for the aerobic degradation of plant
saccharides. Appl. Environ. Microbiol. 80, 2193-2205.

Fontaine, S., Barot, S., Barré, P., Bdioui, N., Mary, B., Rumpel, C., 2007. Stability of organic
carbon in deep soil layers controlled by fresh carbon supply. Nature 450, 277-280.

Halsall, D.M., Goodchild, D.J., 1986. Nitrogen fixation associated with development and
localization of mixed populations of Cellulomonas sp. and Azospirillum brasilense
grown on cellulose or wheat straw. Appl. Environ. Microbiol. 51, 849-854.

Hartman, K., van der Heijden, M.G., Wittwer, R.A., Banerjee, S., Walser, J.-C., Schlaeppi, K.,
2018. Cropping practices manipulate abundance patterns of root and soil microbiome
members paving the way to smart farming. Microbiome 6, 1-14.

Helgason, B., Walley, F., Germida, J., 2010. Long-term no-till management affects microbial
biomass but not community composition in Canadian prairie agroecosytems. Soil Biol.
Biochem. 42, 2192-2202.

Herlemann, D.P., Lundin, D., Labrenz, M., Jürgens, K., Zheng, Z., Aspeborg, H., Andersson,
A.F., 2013. Metagenomic de novo assembly of an aquatic representative of the
verrucomicrobial class Spartobacteria. MBio 4, e00569-00512.

Iino, T., Mori, K., Uchino, Y., Nakagawa, T., Harayama, S., Suzuki, K.-i., 2010.
Ignavibacterium album gen. nov., sp. nov., a moderately thermophilic anaerobic
bacterium isolated from microbial mats at a terrestrial hot spring and proposal of
Ignavibacteria classis nov., for a novel lineage at the periphery of green sulfur bacteria.
Int. J. Syst. Evol. Microbiol. 60, 1376-1382.

IUSS Working Group, W., 2006. World reference base for soil resources. World Soil
Resources Report 103, FAO, Rome.

Izewska, A., Woloszyk, C., 2015. Yields of grain and straw, their content and ionic
proportions of macroelements in maize fertilized with ash from municipal sewage sludge
combustion. J Elementology 20, 319-329.

Izumi, Y., Uchida, K., Iijima, M., 2004. Crop production in successive wheat-soybean rotation
with no-tillage practice in relation to the root system development. Plant production
science 7, 329-336.

Jobbágy, E.G., Jackson, R.B., 2000. The vertical distribution of soil organic carbon and its
relation to climate and vegetation. Ecol. Appl. 10, 423-436.

Jones, D., Willett, V., 2006. Experimental evaluation of methods to quantify dissolved organic
nitrogen (DON) and dissolved organic carbon (DOC) in soil. Soil Biol. Biochem. 38,
991-999.

Kemper, W.D., Schneider, N.N., Sinclair, T.R., 2011. No-till can increase earthworm
populations and rooting depths. J. Soil Water Conserv. 66, 13A-17A.

Kennedy, S., 2000. Energy use in American agriculture. Sustainable energy term paper 5,
1-26.

Kibblewhite, M., Ritz, K., Swift, M., 2008. Soil health in agricultural systems. Philos Trans R
Soc Lond B Biol Sci 363, 685-701.

Koike, I., Hattori, A., 1975. Growth yield of a denitrifying bacterium, Pseudomonas
denitrificans, under aerobic and denitrifying conditions. Microbiology 88, 1-10.

Kremen, C., 2005. Managing ecosystem services: what do we need to know about their
ecology? Ecol. Lett. 8, 468-479.

Lauber, C.L., Hamady, M., Knight, R., Fierer, N., 2009. Pyrosequencing-based assessment of
soil pH as a predictor of soil bacterial community structure at the continental scale. Appl.
Environ. Microbiol. 75, 5111-5120.

Li, Y., Zhang, Q., Cai, Y., Yang, Q., Chang, S.X., 2020. Minimum tillage and residue retention
increase soil microbial population size and diversity: Implications for conservation
tillage. Sci. Total Environ. 716, 137164.

Liang, C., Schimel, J.P., Jastrow, J.D., 2017. The importance of anabolism in microbial
control over soil carbon storage. Nat. Microbiol. 2, 17105.

Liu, C., Lu, M., Cui, J., Li, B., Fang, C., 2014. Effects of straw carbon input on carbon
dynamics in agricultural soils: a meta-analysis. Glob. Chang. Biol. 20, 1366-1381.

Liu, X., Hu, G., He, H., Liang, C., Zhang, W., Bai, Z., Wu, Y., Lin, G., Zhang, X., 2016.
Linking microbial immobilization of fertilizer nitrogen to in situ turnover of soil
microbial residues in an agro-ecosystem. Agric., Ecosyst. Environ. 229, 40-47.

Liu, X., Zhang, X., Wang, Y., Sui, Y., Zhang, S., Herbert, S., Ding, G., 2010. Soil degradation:
a problem threatening the sustainable development of agriculture in Northeast China.
Plant Soil Environ. 56, 87-97.

Liu, Z., Yang, X., Hubbard, K.G., Lin, X., 2012. Maize potential yields and yield gaps in the
changing climate of northeast China. Glob. Chang. Biol. 18, 3441-3454.

Lu, C., Chen, H., Teng, Z., Yuan, L., Ma, J., He, H., Chen, X., Zhang, X., Shi, Y., 2018.
Effects of N fertilization and maize straw on the dynamics of soil organic N and amino
acid N derived from fertilizer N as indicated by 15 N labeling. Geoderma 321, 118-126.

Miroshnichenko, M.L., Kostrikina, N.A., Chernyh, N.A., Pimenov, N.V., Tourova, T.P.,
Antipov, A.N., Spring, S., Stackebrandt, E., Bonch-Osmolovskaya, E.A., 2003.
Caldithrix abyssi gen. nov., sp. nov., a nitrate-reducing, thermophilic, anaerobic
bacterium isolated from a Mid-Atlantic Ridge hydrothermal vent, represents a novel
bacterial lineage. Int. J. Syst. Evol. Microbiol. 53, 323-329.

Mysara, M., Leys, N., Raes, J., Monsieurs, P., 2016. IPED: a highly efficient denoising tool
for Illumina MiSeq Paired-end 16S rRNA gene amplicon sequencing data. BMC
Bioinformatics 17, 192.

Nelson, D., Sommers, L.E., 1982. Total carbon, organic carbon, and organic matter. In: Page,
A., Miller, R., Keeney, D. (Eds.), *Methods of soil analysis,Part 2* (2nd). . American
Society of Agronomy and Soil Science Society of America, Madison, WI, pp. 539-579.

Nevins, C.J., Nakatsu, C., Armstrong, S., 2018. Characterization of microbial community
response to cover crop residue decomposition. Soil Biol. Biochem. 127, 39-49.

Nielsen, M.N., Winding, A., Binnerup, S., 2002. Microorganisms as indicators of soil health.
National Environmental Research Institute, Denmark.

Nunes, M.R., Karlen, D.L., Denardin, J.E., Cambardella, C.A., 2019. Corn root and soil
health indicator response to no-till production practices. Agric., Ecosyst. Environ. 285,
106607.

Parks, D.H., Tyson, G.W., Hugenholtz, P., Beiko, R.G., 2014. STAMP: statistical analysis of
taxonomic and functional profiles. Bioinformatics 30, 3123-3124.

Pathma, J., Raman, G., Sakthivel, N., Microbiome of Rhizospheric Soil and Vermicompost
and Their Applications in Soil Fertility, Pest and Pathogen Management for Sustainable
Agriculture. 189-210 (Springer, Singapore, 2019).

Pierret, A., Maeght, J.-L., Clément, C., Montoroi, J.-P., Hartmann, C., Gonkhamdee, S., 2016.
Understanding deep roots and their functions in ecosystems: an advocacy for more
unconventional research. Ann. Bot. 118, 621-635.

Pries, C.E.H., Castanha, C., Porras, R., Torn, M., 2017. The whole-soil carbon flux in
response to warming. Science 355, 1420-1423.

Prosdocimi, M., Jordán, A., Tarolli, P., Keesstra, S., Novara, A., Cerdà, A., 2016. The
immediate effectiveness of barley straw mulch in reducing soil erodibility and surface
runoff generation in Mediterranean vineyards. Sci. Total Environ. 547, 323-330.

Qin, R., Stamp, P., Richner, W., 2004. Impact of tillage on root systems of winter wheat.
Agron. J. 96, 1523-1530.

Ramirez-Villanueva, D.A., Bello-López, J.M., Navarro-Noya, Y.E., Luna-Guido, M., Verhulst,
N., Govaerts, B., Dendooven, L., 2015. Bacterial community structure in maize residue
amended soil with contrasting management practices. Appl. Soil Ecol. 90, 49-59.

Rincon-Florez, V.A., Carvalhais, L.C., Dang, Y.P., Crawford, M.H., Schenk, P.M., Dennis,
P.G., 2020. Significant effects on soil microbial communities were not detected after
strategic tillage following 44 years of conventional or no-tillage management.
Pedobiologia, 150640.

Rousk, J., Bååth, E., Brookes, P.C., Lauber, C.L., Lozupone, C., Caporaso, J.G., Knight, R.,
Fierer, N., 2010. Soil bacterial and fungal communities across a pH gradient in an arable
soil. ISME J. 4, 1340.

Sagova-Mareckova, M., Zadorova, T., Penizek, V., Omelka, M., Tejnecky, V., Pruchova, P.,
Chuman, T., Drabek, O., Buresova, A., Vanek, A., 2016. The structure of bacterial
communities along two vertical profiles of a deep colluvial soil. Soil Biol. Biochem. 101,
65-73.

Salem, M.A., Economics of reduced tillage technology on soil conservation and risk analysis
for Eastern Oklahoma farmers. (Oklahoma State University, 1983).

Sanderman, J., Hengl, T., Fiske, G.J., 2017. Soil carbon debt of 12,000 years of human land
use. Proc. Natl. Acad. Sci. U.S.A. 114, 9575-9580.

Schimel, J., Schaeffer, S.M., 2012. Microbial control over carbon cycling in soil. Front.
Microbiol. 3, 348.

Song, K., Zheng, X., Lv, W., Qin, Q., Sun, L., Zhang, H., Xue, Y., 2019. Effects of tillage and
straw return on water-stable aggregates, carbon stabilization and crop yield in an
estuarine alluvial soil. Sci. Rep. 9, 4586.

Springmann, M., Clark, M., Mason-D'Croz, D., Wiebe, K., Bodirsky, B.L., Lassaletta, L., De
Vries, W., Vermeulen, S.J., Herrero, M., Carlson, K.M., 2018. Options for keeping the
food system within environmental limits. Nature 562, 519-525.

Stone, L.R., Goodrum, D.E., Jaafar, M.N., Khan, A.H., 2001. Rooting front and water
depletion depths in grain sorghum and sunflower. Agron. J. 93, 1105-1110.

Sun, R., Li, W., Dong, W., Tian, Y., Hu, C., Liu, B., 2018. Tillage changes vertical distribution
of soil bacterial and fungal communities. Front. Microbiol. 9, 699.

Tautges, N.E., Chiartas, J.L., Gaudin, A.C., O'Geen, A.T., Herrera, I., Scow, K.M., 2019.
Deep soil inventories reveal that impacts of cover crops and compost on soil carbon
sequestration differ in surface and subsurface soils. Glob. Chang. Biol. 25, 3753-3766.

Terracciano, J., Canale-Parola, E., 1984. Enhancement of chemotaxis in Spirochaeta aurantia
grown under conditions of nutrient limitation. J. Bacteriol. 159, 173-178.

Thomas, F., Hehemann, J.-H., Rebuffet, E., Czjzek, M., Michel, G., 2011. Environmental and
gut bacteroidetes: the food connection. Front. Microbiol. 2, 93.

Thorup-Kristensen, K., 2006. Effect of deep and shallow root systems on the dynamics of soil
inorganic N during 3-year crop rotations. Plant Soil 288, 233-248.

Thorup-Kristensen, K., Halberg, N., Nicolaisen, M., Olesen, J.E., Crews, T.E., Hinsinger, P.,
Kirkegaard, J., Pierret, A., Dresbøll, D.B., 2020. Digging Deeper for Agricultural
Resources, the Value of Deep Rooting. Trends Plant Sci. 25, 406-417.

Toosi, E.R., Castellano, M.J., Singer, J.W., Mitchell, D.C., 2012. Differences in soluble
organic matter after 23 years of contrasting soil management. Soil Sci. Soc. Am. J. 76,
628-637.

Vitousek, P.M., Naylor, R., Crews, T., David, M.B., Drinkwater, L., Holland, E., Johnes, P.,
Katzenberger, J., Martinelli, L., Matson, P., 2009. Nutrient imbalances in agricultural
development. Science 324, 1519-1520.

Wang, Q., Garrity, G.M., Tiedje, J.M., Cole, J.R., 2007. Naive Bayesian classifier for rapid
assignment of rRNA sequences into the new bacterial taxonomy. Appl. Environ.
Microbiol. 73, 5261-5267.

Wood, S.A., Bradford, M.A., Gilbert, J.A., McGuire, K.L., Palm, C.A., Tully, K.L., Zhou, J.,
Naeem, S., 2015. Agricultural intensification and the functional capacity of soil microbes
on smallholder African farms. J. Appl. Ecol. 52, 744-752.

Zhalnina, K., Dias, R., de Quadros, P.D., Davis-Richardson, A., Camargo, F.A., Clark, I.M.,
McGrath, S.P., Hirsch, P.R., Triplett, E.W., 2015. Soil pH determines microbial diversity
and composition in the park grass experiment. Microb. Ecol. 69, 395-406.

Zhang, B., Wang, H., Yao, S., Bi, L., 2013a. Litter quantity confers soil functional resilience
through mediating soil biophysical habitat and microbial community structure on an
eroded bare land restored with mono Pinus massoniana. Soil Biol. Biochem. 57,
556-567.

Zhang, C., Ju, X., Powlson, D.S., Oenema, O., Smith, P., 2019. Nitrogen surplus benchmarks
for controlling N pollution in the main cropping systems of China. Environ. Sci. Technol.
53, 6678-6687.

Zhang, W.-f., Dou, Z.-x., He, P., Ju, X.-T., Powlson, D., Chadwick, D., Norse, D., Lu, Y.-L.,
Zhang, Y., Wu, L., 2013b. New technologies reduce greenhouse gas emissions from
nitrogenous fertilizer in China. Proc. Natl. Acad. Sci. U.S.A. 110, 8375-8380.
741

 Figure legends

Fig. 1. Soil properties (mean± SE, n = 3) along soil depth under different

practices. a, SOC = soil organic carbon; **b**, SEOC = salt-extractable organic carbon; **c**,

SEOC/SOC = ratio of SEOC to SOC; **d**, soil pH; **e**, SM = soil moisture; **f**, TN = total

nitrogen content; **g**, C/N = ratio of SOC to TN; **h**, NH_4^+-N = ammonium nitrogen; **i**,

NO_3^--N = nitrate nitrogen; **j**, Inorganic N = NH_4^+-N + NO_3^--N. Error bars indicate

standard errors (n = 3). Different letters indicate significant differences (P < 0.05)

among disturbing practices.

Fig. 2. Corn yield (line+ symbol) and annual rainfall during growing and

dormant seasons (bar) under different disturbance practices during 2013-2016.

Error bars indicate standard errors (n = 3 or 4), different letters indicate significant

differences at P < 0.05. Different corn seeds were used among years.

Fig. 3. Microbial richness (Chao1) observed number of species

(Observed_species) and diversity (Shannon_Index) in the CT, NTNS and NTSM

plots. Error bars indicate standard errors (n = 3). Different letters indicate significant

differences (P < 0.05) among disturbing practices.

Fig. 4. Indicator genera significantly (q < 0.1) associated with tillage practices.

The size of each circle represents the indicator value of a specific genus in the

different soil depths. The color indicates the relative abundance of each indicator

genus. Taxonomic information, indicator values, P-values, and q-values of all

indicator genera are given in Table S2. Zero indicator genera were identified in CT

treatment.

Fig. 5. Non-metric multidimensional scaling (MDS) ordination of soil microbial

community structures based on Bray-Curtis distances among soil depths at

different agricultural disturbance practices. Permutational multivariate analysis of

variance (PERMANOVA) revealed that the overall microbial community structures

among soil depth were significantly different at each disturbance practice. Circles,

triangles and squares represent CT, NTNS and NTSM, respectively.

Fig. 6. Redundancy analysis (RDA) of the soil microbial community originating

from microbial phyla constrained by soil properties under different agricultural

practices. Only soil variables that significantly explained variability in microbial

community structure in the forward selection procedure were selected to the

ordination (arrows). TN, total nitrogen content; C/N, a ratio of carbon to nitrogen

content; NH_4^+-N, ammonium nitrogen; NO_3^--N, nitrate nitrogen ; SM, soil moisture.

Circles, triangles and squares represent CT, NTNS and NTSM, respectively.

Fig. 1

782   **Fig. 2**

786   **Fig. 3**

790 **Fig. 4**

794   **Fig. 5**

797 **Fig. 6**

798
799