Low-disturbance farming regenerates healthy critical zone towards sustainable agriculture

Fangbo Deng[1,2#], Hongjun Wang[3#], Hongtu Xie[1#], Xuelian Bao[1], Hongbo He[1], Xudong Zhang[1], Chao Liang[1]*

[1]Key Laboratory of Forest Ecology and Management, Institute of Applied Ecology, Chinese Academy of Sciences, Shenyang 110016, China

[2]University of Chinese Academy of Sciences, Beijing 100049, China

[3]Duke University Wetland Center, Nicholas School of the Environment, Duke University, Durham, NC 27708, USA

[#] These authors contributed equally to this work.

Intensive conventional farming has degraded soil quality in farmlands and other

ecosystems globally. Although low-disturbance practices have been widely

adapted to restore soil health and save energy, the underlying mechanisms

associated with farm sustainability are still unclear. Here, we compared soil

microbiome, physiochemical parameters along 3-m deep soil profiles, and crop

yield in Northeast China subjected to ten years of farming practices at 3 levels of

disturbance, including conventional tillage (CT), no-tillage without stover

mulching (NTNS), and no-tillage with stover mulching (NTSM). We found that

low-disturbance practices (NTNS and NTSM) promoted the ability of the soil to

retain water and nitrogen, regenerated whole-soil microbial diversity and

function, and significantly improved corn yield during the drought year. This

study implies that the NTSM practice could potentially cut fertilizer-N input by

281.6 kg/ha to corn farmland in China, which may consequently reduce China's

total greenhouse gas emissions by 1.6% and save about 6.7% households energy

while without reducing corn production.

Since the Industrial Revolution, the rate of soil carbon loss has increased

dramatically, resulting in a global carbon debt due to agriculture of 116 Pg carbon for

the top 2 m of soil[1]. The loss of carbon in farmlands has not only changed global

climate but also produced catastrophic cascade impacts on global food security, as soil

carbon is the cornerstone for healthy and productive soil that will be needed to feed

10 billion people in 2050 (United Nations, World Population Prospects 2019). It is

well known that intensive conventional farms with high inputs and disturbance (e.g.

chemical fertilizers, tillage/compaction, burn/remove stover) have caused soil carbon

loss with a series of environmental issues[2]. Even worse, increasing the amount of

chemical fertilizer is unlikely to continue the increase in quantity and quality of food

products worldwide[3]. Moreover, tillage particularly prevents root growth into deeper

soil[4], thus affecting mineral weathering in deep soil[5] and reducing crop resilience to

drought.

Since the 1970s, low-disturbance conservative practices (e.g. reduced tillage,

no-tillage and stover mulching) have been gradually applied to restore soil health and

reduce non-point source pollution[6]. Growing evidence shows that no-tillage and

stover mulching boosted top-soil organic carbon (SOC)[7, 8, 9], increased soil

aggregate[10] and reduced soil erosion and surface runoff[11]. All these benefits from

low-disturbance practices are tied with complex microbial processes that interact with

crops and drive soil carbon transformation and stabilization[12]. However, most studies

only focused on topsoil or soils within 1-m depth[13, 14, 15]. Soil below 1 meter, which

belongs to Earth's Critical zone, was often overlooked despite that some crops have

roots over 1 meter deep and microbes in the soils (> 1 m) may substantially impact

long-term carbon sequestration, mineral weathering and crop production[16, 17, 18].

Furthermore, deep roots influence material cycles in surficial soil and microbes

inhabiting in the deep soils, which plays important roles in bridging aboveground

vegetation with parent soils and even acts as an essential buffer protecting

underground water[19]. To completely evaluate the impact of low-disturbance practices

on agro-ecosystem, an urgent need is to test the changes of microbial communities in

deep soil[20]. Since microbial activities in deep soil are generally limited by the

availability of labile carbon, we hypothesized that low-disturbance practices are

conducive to deeper root growth, which could provide labile carbon and nutrients[4]

and accelerate mineral weathering[5], thus substantially influencing the microbial

composition in the deep soils and in turn the sustainability of whole ecosystem.

Recent research shows that corn belts in the U.S.A., western Europe, and China

have experienced the most soil carbon loss globally[1]. In China, this region is

considered as the "breadbasket" of the country, having the largest grain production

and overlapping with the most fertile Mollisol region that sustains 3% of population

in the world[21], accounting for 30% of corn production of China[22]. Here, a 10-year

manipulative experiment was conducted at a temperate corn farm on Mollisol in

Northeast China, investigating farming practices with three levels of disturbance:

high-disturbance—conventional tillage (CT), low disturbance—no-tillage without

stover mulching (NTNS) and no-tillage with 100% stover mulching (NTSM). We

compared corn yield, soil properties and microbial communities of the 3-m soil

profiles at the end of entire 10 years. We aimed at testing our main hypothesis that the

lowest disturbance practice—no-tillage with 100% stover mulching, similar to

undisturbed natural ecosystem, would regenerate microbial diversity and function

toward a high-resilient natural ecosystem.

Results

Soil properties and corn yield. Soil properties varied significantly among

disturbance practices and at different soil depths (Supplementary Table 1). The SOC,

TN and C/N ratio substantially decreased from the soil surface to around 150 cm

depths and then remained unchanged within 150-300 cm (Fig. 1). The NTSM slightly

increased SOC, TN and C/N ratio at 0-20 cm soil layers compared with the NTNS and

the CT (Fig. 1 and Supplementary Table 1). The no-tillage practices of NTSM and

NTNS reduced soil pH in surface and deeper layers (Fig. 1d) and increased soil

moisture at surface layers (0-60 cm) (Fig. 1e). In the CT plots, soil NO_3^--N

concentration first decreased and then increased remarkably, ranged from 4.19 to

23.32 mg kg^{-1} (Fig. 1i). However, under the NTNS and NTSM treatments, soil

NO_3^--N decreased significantly at 0-40 cm then increased to the maximum at 120-150

cm depth. Interestingly, above 120-150 cm layer, NO_3^--N was significantly higher

with low-disturbance practices than conventional tillage, while the soil below 150 cm

under low-disturbance practices had much lower NO_3^--N compared to conventional

tillage (Fig. 1i). The NTNS plots contained much higher amounts of ammonium than

the CT and the NTSM plots (Fig. 1h). Soil salt-extractable organic carbon (SEOC), as

an organic acid proxy, is positively associated with root density and can be an

indicator of root depth in deep soil[5]. The SEOC declined from the surface to 40-60 cm

and then increase to its peak at 60-90 cm under CT, at 90-120 cm under NTNS and at

120-150 cm under NTSM (Fig. 1b). Hence, we estimate that corn roots reached up to

60-90 cm, 90-120 cm and 120-150 cm under the CT, the NTNS and the NTSM,

respectively, which is in line with reported corn root depths (~150 cm)[4, 23]. The NTSM

increased the SEOC concentration at almost all soil layers compared with the CT and

the NTNS (Fig. 1b), in which at the surface and 120-150 cm depth the contents of

SEOC with NTSM were twice higher than CT. The increased SEOC in deep soils

under NTSM reduced soil pH as shown by a significant negative relationship between

SEOC and pH (r=0.678, p<0.05). The relative contributions of SEOC to SOC

(SEOC/SOC) in the NTSM were also always higher than in the CT and the NTNS

(Fig. 1c). Based on the estimated root depths, total soil inorganic nitrogen available

for the coming growing season in the NTSM and the NTNS was 427.34 and 352.34

kg ha^{-1}, respectively, while only 179.63 kg ha^{-1} in conventional tillage.

The mean annual corn yield (2013-2016) in the NTSM is 13416.8 kg/ha, which is

much higher than the CT and NTNS (Fig. 2), particularly during the drought year of

2015, with only 409.6 mm of rainfall during the growing season (about 100 mm lower

than the mean rainfall), while the corn yield in NTSM is 36.4% and 22.3% higher

than the CT and NTNS, respectively (Fig. 2).

Microbial diversity, composition, and structure. The microbial richness (Chao1),

observed number of species (Observed-species) and diversity (Shannon-Index) first

increased within 0-20 cm and decreased from 20 to 90 cm, then increased hereafter

(Fig. 3). The low-disturbance practices significantly increased Chao1,

Observed-species and Shannon-Index, particularly in 0-40 cm soil depths (Fig. 3).

There were 54 microbial phyla across all soil samples. The dominant phyla (relative

abundance > 1% across all soil samples) were Proteobacteria, Actinobacteria,

Chloroflexi, Acidobacteria, Nitrospirae, Gemmatimonadetes, Planctomycetes, and

these phyla accounted for 60-91% of the total microbial abundances in the whole soil

profile (Supplementary Fig. 1a). Bacteroidetes, Verrucomicrobia, Latescibacteria,

Parcubacteria, Firmicutes, Microgenomates and Saccharibacteria were less dominant

(relative abundance > 0.1% across all soil samples) but were still found across all soil

samples (Supplementary Fig. 1a). Although no difference in the composition of

dominant phyla among treatments were found, there are more non-dominant phyla

with higher relative abundance in low disturbance practices than conventional tillage

practice (Supplementary Fig. 1b).

Indicator analysis identified 16 and 51 clearly classified genera (relative

abundances > 0.005%) in the NTNS and the NTSM plots, respectively, while no

indicator genera were found in the conventional tillage plots (Fig. 4 and

Supplementary Table 2). The indicator genera in the NTNS plots belonged to

Proteobacteria, Actinobacteria, Chloroflexi, Gemmatimonadetes and Planctomycetes,

and most of them appeared in the surface soil (0-20 cm) with only 1 genus below 150

cm. Importantly, more extra indicator genera — including Bacteroidetes,

Acidobacteria, Deferribacteres, Firmicutes, Verrucomicrobia, Chlorobi and

Spirochaetae — existed in the NTSM plots, in which under 150 cm we observed 7

genera (Fig. 4 and Supplementary Table 2).

Microbial community structures were visualized by Non-metric multidimensional

scaling (MDS) and tested by Permutational multivariate analysis of variance

(PERMANOVA) based on Bray–Curtis. The microbial communities among

treatments in the root zones were marginally different (PERMANOVA p=0.08);

however, below the root zone they differed distinctively (PERMANOVA p=0.02). The

disturbance practices influenced the vertical distribution dissimilarity in microbial

community structure (Fig. 5). Three clusters — 0-10 cm and 10-20 cm, 20-150 cm

and 150-300 cm — were observed in the CT plots (PERMANOVA-F=9.57, p=0.0001)

(Fig. 5). In the NTNS plots, 0-10 cm formed an independent cluster, while other soil

depths showed some separation (e.g. 20-120 cm were separated from 150-300 cm soil

depths by axis 1); however, Bray-Curtis distances between adjacent depths were too

close to be separated (PERMANOVA-F=8.18, p=0.0001) (Fig. 5). The NTSM

treatment clustered 0-10 cm and 10-20 cm together, 120-150 cm, 150-200 cm,

200-250 cm and 250-300 cm separately, and the other depths show some separations

as well (PERMANOVA-F=11.32, p=0.0001) (Fig. 5).

Predicted Ecological functions of microbial communities. According to the results

of microbial diversity, composition and structure, the metabolic capabilities of

microbial community in the whole 3-m soil profiles were predicted using Tax4Fun

(Supplementary Fig. 2). Results showed that low-disturbance practices significantly

increased the abundance of predicted functions related to carbohydrate metabolism,

nucleotide metabolism, glycan biosynthesis and metabolism, lipid metabolism and

metabolism related to cofactors and vitamins (Supplementary Fig. 2a). Moreover, the

relative abundances of genes encoding for assimilatory nitrate reduction in

low-disturbance practices were higher than that in conventional tillage practice

(Supplementary Fig. 3). The results suggested that in low disturbance practices,

microbial community prefer to convert the nitrate/nitrite to ammonia. We then further

assessed the impact of stover mulching on functional profiles (Supplementary Fig. 2b).

The extended error bar plot shows that the NTNS enriched the abundance of amino

acid metabolism and lipid metabolism, while the NTSM enriched the functions

associated to energy metabolism, carbohydrate metabolism, biosynthesis of secondary

metabolites, glycan biosynthesis and metabolism as well as metabolism of cofactors

and vitamins (Supplementary Fig. 2b).

Relationships between microbial communities and soil properties. Forward

selection in Redundancy analysis (RDA) revealed that soil depth (pseudo-F=48, p=

0.002), SOC (pseudo-F=11.5, p= 0.002), SM (pseudo-F=3.4, p= 0.012), soil pH

(pseudo-F=2.3, p=0.018) and soil NH_4^+-N (pseudo-F=2.7, p= 0.026) significantly

affected the vertical distribution of microbial communities (Supplementary Fig. 4).

Furthermore, the soil properties that regulated the distribution of soil microbes were

different under different disturbance practices. Under the CT treatment, soil microbial

community was mainly affected by soil NH_4^+-N (pseudo-F=4, p= 0.002) and soil

NO_3^--N (pseudo-F=2.3, p= 0.012) that mainly came from applied fertilizer (Fig. 6).

The microbial community positively correlated to soil NH_4^+-N in the 0-20 cm soil, to

soil NO_3^--N negatively within 20-150 cm, while to soil NO_3^--N positively after 150

cm (Fig. 6). Under the NTNS treatment, soil pH (pseudo-F=3.7, p=0.004) constrained

the distribution of the microbial community, in which strong negative correlations

occurred in 0-10 cm soil and a positive correlation in 90-150 cm (Fig. 6). Under the

NTSM treatment, soil TN (pseudo-F=11, p=0.002), SM (pseudo-F=2.6, p=0.004) and

C/N ratio (pseudo-F=1.8, p=0.016) significantly influenced the soil microbial

community separation (Fig. 6). In general, the microbes positively correlated with the

soil TN and C/N ratio in the surface soil layers (0-40 cm) and with SM in the middle

layers (40-150 cm), while they were mainly influenced by depth in the deeper soil

(150-300 cm) (Fig. 6).

Discussion

No-tillage practices promote soil health and corn yield. No-tillage promotes root

growth into deep soil, up to 150 cm in the NTSM. The root exudates with various

organic acid and dead roots likely contributed to the lower soil pH and higher SEOC

in the NTNS and the NTSM, which in turn increased mineral weathering[5] and

diversified the microbial communities with multi-ecological functions. The increased

fine roots in deeper soil retained more nutrients including nutrients in dead roots and

converting nitrate to more stable ammonium (Supplementary Fig. 3) and also

provided labile carbon (Fig. 1b) to remove leaked nitrate through denitrification in

deeper soil (below 1.5 m), as higher relative abundance of the denitrification bacteria

(*Pseudomonas* and *Caldithrix*)[24, 25] (Fig. 4 and Table. S2) and denitrification genes

(Supplementary Fig. 3) were detected in low disturbance practices — particularly in

no-tillage with stover mulching. However, shallower roots in the CT treatment can't

provide enough labile carbon to remove extra soil NO_3^--N in deep soil, thus causing

nitrite accumulation and leaching into deeper soil layers. The amount of inorganic

nitrogen accumulated in the root zones under NTSM (427.34 kg ha^{-1}) likely could

provide plenty of nitrogen for corn growth in the coming growing season (Fig. 2),

based on the removed nitrogen in the grain (~200 kg ha^{-1}). Additionally, in line with

many studies that show stover mulching reduces water evaporation and surface runoff

and increase soil moisture in top soils[11, 26], we found that the soil moisture was

significantly higher in the NTSM than in the CT plots. Therefore, no-tillage with

stover mulching not only restores soil health by increasing the holding capacities for

nutrients and water, thus reducing energy input to farm, but also tended to reduce the

risk of nitrate leaching to groundwater. And more importantly, the healthy soil in turn

raises corn production and promote the crop resistance to drought (Fig. 2). All these

are critical to the development of sustainable agriculture and the associated

ecosystems.

No-tillage with stover mulching promotes microbial diversity, richness, and

ecological function contributing to sustainable farming. Under the CT treatment,

tillage heavily disturbed the topsoil and liberated occluded organic materials.

Microbes tended to rapidly use available nutrients in the plowed layer (e.g. NH_4^+-N)[27],

thereby causing the reduction of microbial metabolic diversity (Supplementary Fig.

2a). Then, the resistance of the soil to stress or disturbance may also decrease[28]. In

deeper soil layers, due to shallower roots, NO_3^--N could quickly move downward and

accumulate in deeper soil (Fig. 1i), which not only contaminated the underground

water but also limited the activity of non-dominant microbes with important

ecological functions, as no indicator genera were identified for each soil depth in CT

treatment (Fig. 4 and Supplementary Table 2). Because the microbial communities

were closely associated with inorganic nitrogen, the microbes under CT were mainly

influenced by added chemical fertilizer. Although the dominant microbial

communities in CT were similar to those in the NTNS and NTSM, the loss of function

resulted from the difference of non-dominant microbes, indicating that the soil under

CT had degraded.

Under the NTNS treatment, soil pH was the major edaphic factor affecting the

microbial community and the indicator genera (Fig. 6 and Supplementary Table 2).

The lower soil pH possibly was caused by deeper roots as shown by higher SEOC that

is generally positively related to root density[5]. Soil pH is often observed as a major

factor determining the microbial composition and structure in natural ecosystems[29, 30],

as microbes often show a narrow tolerance to soil pH. In addition, soil pH regulates

the availability of nutrient and mitigate ion toxicity[29, 30, 31]. Under NTNS, soil pH and

depth only explained 35% distribution of the microbial community (Fig. 6). We

speculated that other edaphic factors (e.g. salinity and iron) directly or indirectly

related to soil pH and SEOC also influenced the changes in the microbial community.

Under NTSM treatment, TN and C/N significantly correlated with soil microbial

community due to the high C/N ratio of stover and roots (Fig. 6). Prior studies have

reported that, following maize stover mulching, more organic N, amino acid N, and

amino sugar N were observed in soil[32, 33], which increased the retention time of

nitrogen, hence meeting the nutrient requirement of corn growth and reducing nitrate

loss to underground water. The increased available nitrogen, labile carbon and water

in deep soil under NTSM can increase the resilience and resistance of maize to

disturbances with higher grain production (Fig. 2). Zhang et al.[34] also observed

litter-covered soil showed greater resistance to heating and copper addition due to the

changes in soil properties and microbial community structure. Resistance to

disturbance or stresses is the nature of a healthy soil and is essential for maintaining

ecosystem functions, such as decomposing organic matter [34, 35]. Under the NTSM

treatment, the microorganisms associated with the degradation of relatively stable

carbon compounds, such as Planctomycetes and Verrucomicrobia (Supplementary

Table 3)[36, 37] as well as the indicator *Cellulomonas* and *Azospirillum* (Fig. 4 and

Supplementary Table 2) with the function of cellulose decomposition[38, 39] were

increased. The predicted functional profiles related to energy metabolism (Carbon

fixation pathways in prokaryotes), carbohydrate metabolism (TCA cycle, amino sugar,

nucleotide sugar, galactose, fructose), biosynthesis of secondary metabolites

(Carotenoid and Betalain) and glycan biosynthesis were increased, suggesting a

higher metabolic activity and a change in substrate quality (Supplement Fig. 2). In

addition, stover mulching also increased the ecological filter function of soil depth for

selecting microbial communities as more indicator genera of each soil depths were

identified under NTSM compared to NTNS and CT practices (Fig. 4 and

Supplementary Table 2). And these indicators residing at different soil depths might

enhance the anti-disturbance ability of NTSM. For example, denitrification bacteria

Caldithrix and *Pseudomonas*[24, 25] were the indicator genera of 150-200 cm and

250-300 cm, respectively (Fig. 4 and Supplementary Table 2), which might explain

the low nitrate in the deep soil in NTSM. *Ignavibacteria* and *Spirochaeta,* the

indicator genera of deep soil, have the ability to grow under the conditions of strictly

anaerobic[40] and severely limited nutrients[41], respectively. Surface indicator genera

belonging to Bacteroidetes might have the ability to degrade organic matter that is

difficult to decompose[42].

Implications for climate change. It was observed that about 179.63, 352.34 and

427.34 kg ha^{-1} inorganic N were kept in the root-zone soil in the CT, NTNS and

NTSM, respectively. Generally, corn roots reach their maximum depth at the silking

stage[43], which is also the time when the heaviest rainfall occurs in northeastern China.

We therefore expect that the available N kept in the root zone would be utilized by

crops in the coming growing season, which means that fertilizer N could be cut to

meet crop growth and also prevent reactive N losses. Since the nitrogen use efficiency

(NUE) of maize system under the conventional management is 51% in northeast

China (NUE is defined as the efficiency of fertilizer N transferring to harvested crop

N)[44]. Then, we   conservatively calculate the required fertilizer N in the next year

based on two assumptions: 1) the NUE of soil available N in root zone is equal to that

NUE of applied fertilizer N, both of them are 50%; 2) the mineralized N during the

coming growing season is neglected. Thus, N supply requirement = Fertilizer N×NUE

+ N in root zone ×NUE + Stover-N, where Stover-N for NTSM is 60 kg ha^{-1}. We

estimated the N requirement for each disturbance practice by multiplying grain yield

by grain N concentration (1.4%)[44] plus multiplying stover yield by stover N

concentration (0.8%)[45]. For CT, NTNS and NTSM, the mean annual corn yields were

10946.74, 12487.81 and 13416.81 kg ha^{-1}, and the stover yields were 966.67,

10083.33 and 10833.33 kg ha^{-1}, respectively. Thus, the N requirements were 230.6,

255.5 and 274.5 kg ha^{-1} for CT, NTNS and NTSM, respectively. Therefore, the

theoretical amounts of fertilizer N in the coming growing season are 281.6, 158.7 and

1.7 kg ha^{-1} for CT, NTNS and NTSM, respectively. No fertilizer-N is needed to apply

without reducing corn yield in the NTSM plot. Compared to CT, the NTNS and

NTSM could save respectively about 122.9 and 281.6 kg ha^{-1} N-fertilizer. For every

kilogram of fertilizer-N produced and used on cropland, up to 87.9 MJ of energy is

consumed[46] and 13.5 kg of CO_2-equivalent (eq) (CO_2-eq) is emitted[47]. Hence, totally

24,752.6 MJ of energy consumption could be reduced and 3,801.6 kg CO_2-eq

emission could be cut per hectare cornland by using NTSM tillage practice.

Multiplying the area of all maize farmland in China (42,000,000 ha, Source: China

Statistics Yearbook 2018), 1.0 EJ of energy could be saved and 159.7 Mt of CO_2-eq

could be reduced. Based on the average annual energy consumption for households of

China in 2017 (15 EJ, China Statistics Yearbook 2018) and CO_2 emissions (9,839 Mt,

Global Carbon Atlas), the NTSM practice in corn farming has the potential to save 6.7%

of household energy and to reduce 1.6% of CO_2 emissions each year in China.

Taken together, we provide new evidence that low-disturbance practice promotes

deep-soil stability to cope with environmental stress through increasing water and

nutrient holding capacity, microbial richness, microbial diversity and ecological

functions. According to ecological theory[48, 49], microbial community assembly in the

CT treatment was mainly based on deterministic processes and significantly

influenced by environmental stress and fertilizer nitrogen. Stover mulching might

alter these processes through deeper roots affecting the vertical heterogeneity in

resource availability[4]. When energy resources are richer in the soil, environmental

stress tend to alleviate[50], and higher biodiversity was caused due to more stochastic

processes introduced in community assembly[51]. Moreover, low disturbance practice

also showed the potential to increase the maize yield (Fig. 2), save energy, and

decrease the risk of groundwater leaching and greenhouse gas emissions. In view of

the importance of microbial community assembly in predicting ecosystem service

functions[52, 53], our results demonstrated that the lowest disturbance-practice —

no-tillage with stover mulching increases the sustainability of agro-ecosystems.

Material and methods

Site description and soil sampling. The field experiment was established in 2007 at the Lishu Conservation Tillage Research and Development Station of the Chinese Academy of Sciences in Jilin province, Northeast China (43.19° N, 124.14° E). The region has a humid continental climate with a mean annual temperature of 6.9 °C and the mean annual precipitation of 614 mm. The soils are classified in the Mollisol order (Black Soil in Chinese Soil Classification) with a clay loam texture[54]. The site has been continuously planted with maize since 2007. We set up an experiment by a randomized complete block design with four replicates and five treatments. Each plot area was $261m^2$ (8.7×30m). The five treatments included conventional tillage (moldboard plowing to a depth around 30 cm), no-tillage (no soil disturbance and direct seeding), and no-tillage with three-level stover mulching (33%, 67% and 100% newly produced maize stover were evenly spread over the soil surface each fall). For each treatment, slow-release fertilizer was applied at one time when sowing, which was equal to 240 kg/ha N; 47 kg/ha P; 90 kg/ha K. The rainfall data were obtained from local meteorological admistration. The grain yield was estimated by manually harvesting 20 m^2 area, randomly taken from each plot.

In this experiment, in order to reduce the damage to the plots and reduce costs, 3 plots were randomly taken from each treatment including conventional tillage (CT), no-tillage without stover mulching (NTNS), no-tillage with 100% stover coverage (NTSM) as three comparative practices. In April 2017, triplicate soil cores (0-300 cm) were collected from each plot at the dormant season. After removing surface stover, we took soil cores by a stainless-steel hand auger and sliced each into ten layers: 0-10 cm, 10-20 cm, 20-40 cm, 40-60 cm, 60-90 cm, 90-120 cm, 120-150 cm, 150-200 cm, 200-250 cm, 250-300 cm. In total, 90 soil samples were collected and transported to

the laboratory within 3 hours, then passed through a 2-mm sieve. All visible roots,

crop residues and stones were removed. Each soil sample was divided into three

subsamples: one subsample for DNA extraction and soil salt-extractable organic

carbon (SEOC) measurement that was immediately placed into a polyethylene plastic

bag and stored at -80 °C, one for chemical measurements including ammonium

nitrogen (NH_4^+-N) and nitrate nitrogen (NO_3^--N) (within one day), and the remaining

one was air dried for other soil physicochemical properties.

Soil properties. Soil total nitrogen (TN) content was measured by an Element

analyzer Vario EL III (Elementar Analysensysteme GmbH, Hanau, Germany). Soil

organic carbon (SOC) was converted from soil organic matter (SOM) that was

measured by potassium dichromate oxidation[55]. Soil pH was measured in deionized

free-CO_2 water (1:2.5 w/v). Gravimetric soil moisture was determined by oven-drying

fresh soil to a constant weight at 105 °C. Soil NH_4^+-N and NO_3^--N were extracted

from fresh soil by 2 M KCl and measured by a continuous flow analytical system

(AA3, SEAI, Germany). To reflect soil soluble, exchangeable, mineral-bound OC,

soil salt-extractable organic carbon (SEOC) was extracted from the frozen soil

samples with 0.5 M K_2SO_4 (1:5 w/v)[4, 56, 57].

DNA extraction, PCR amplification and pyrosequencing. Soil DNA was extracted

from the frozen soil samples (0.5 g wet weight) by using MoBio PowerSoil DNA

isolation kit (MoBio Laboratories, Carlsbad, CA, USA) following the instructions of

the manufacturer. The quality of DNA was determined by 1% agarose gel

electrophoresis. The V3–V4 region of the bacterial 16S rRNA gene was amplified by

PCR using the primers 338F and 806R with barcode for Illumina MiSeq sequencing.

PCR was performed in a total volume of 50 μl containing 30 ng DNA as a template,

20 mol of each primer, 10mM dNTPs, 5μl 10× Pyrobest buffer and 0.3 U of Pyrobest

polymerase (Takara Code: DR005A). The PCR cycle conditions were as follows:

initial denaturation at 95°C for 5 min followed by 26 cycles of denaturation at 95°C

for 45s, annealing at 50°C for the 30s, and extension at 72°C for 45s, with a final

extension at 72°C for 10 min. Each sample was amplified for three replicates. The

PCR products from the same sample were pooled, checked by 2% agarose gel

electrophoresis and were then purified using AxyPrepDNA agarose purification kit

(AXYGEN). Finally, purified PCR products were sequenced on an Illumina MiSeq

platform PE300 sequencer (Illumina, USA).

The raw sequence data were further analyzed by the following protocol.

Low-quality sequences with an average quality score of less than 20 were filtered by

employing Trimmomatic[58]. The FLASH software was used to merge overlapping ends

and treat them as single-end reads[59]. The non-amplified region sequences, chimeras

and shorter tags were also removed using Usearch and Mothur[60, 61]. The resulting

high-quality sequences were clustered into Operational Taxonomic Units (OTUs) at

97% sequence similarity using Usearch (Version 8.1.1861

http://www.drive5.com/usearch/) (Edgar, 2013). OTUs were then classified against

the Silva (Release119 http://www.arb-silva.de) database and the taxonomic

information of each OTU representative sequence was annotated using the RDP

Classifier[62, 63, 64]. A total of 3,255,693 high-quality reads were obtained from all soil

samples, which were clustered into 9,573 unique OTUs at a 97% sequence similarity.

The Good's coverage of all the samples ranged from 0.93 to 0.98, which indicates an

adequate level of sequencing to identify the majority of diversity in the samples.

Statistical analyses. Soil properties were analyzed and plotted using Sigmaplot 12.5

software. Alpha diversity indices were calculated in Qiime (version v.1.8) and used to

reflect the diversity and richness of the microbial community in different samples.

The relative abundances of individual phyla in different samples were computed by R

packages. The indicator analysis based on genera-specific to each soil depth was

conducted using indicspecies package of R with 9999 permutations, and the P-values

were corrected for multiple testing using qvalue package of R[13, 65]. Functional profiles

of the microbial community were predicted by Tax4fun (an open-source package in

R)[66] and further statistical analysis was conducted by STAMP using Welch's t-test[67].

Non-metric multidimensional scaling (MDS) was performed by "vegan" package of R

to describe differences in microbial community structure among samples.

Permutational multivariate analysis of variance (PERMANOVA) was employed on

Bray-Curtis distances to test the differences in soil microbial communities among

various sample groups. The Redundancy analysis (RDA, Canoco 5 software) were

conducted to identify the correlations between microbial community composition and

environmental variables. One-way and two-way ANOVA tests were conducted by

SPSS Version 22. Percentage data were transformed using arcsine square root

function before ANOVA test. All statistical tests were significant at $p \leq 0.05$.

Data availability. All sequencing data that support the findings of this study have

been deposited in the National Center for Biotechnology Information

(https://www.ncbi.nlm.nih.gov/), in the Sequence Read Archive (SRA) database

(BioProject number: PRJNA488172). All other relevant data are available from the

corresponding author on request.

References

References

1. Sanderman J, Hengl T, Fiske GJ. Soil carbon debt of 12,000 years of human land use.

Proc Natl Acad Sci USA **114**, 9575-9580 (2017).

2. Congreves K, Hayes A, Verhallen E, Van Eerd L. Long-term impact of tillage and crop

rotation on soil health at four temperate agroecosystems. *Soil Tillage Res* **152**, 17-28

(2015).

3. Vitousek PM*, et al.* Nutrient imbalances in agricultural development. *Science* **324**,

1519-1520 (2009).

4. Kemper WD, Schneider NN, Sinclair TR. No-till can increase earthworm populations

and rooting depths. *J Soil Water Conserv* **66**, 13A-17A (2011).

5. Billings SA*, et al.* Loss of deep roots limits biogenic agents of soil development that

are only partially restored by decades of forest regeneration. *Elem Sci Anth* **6**, 34

(2018).

6. Salem MA. Economics of reduced tillage technology on soil conservation and risk

analysis for Eastern Oklahoma farmers. *(Oklahoma State University, 1983)*.

7. Blanco-Canqui H, Lal R. No-tillage and soil-profile carbon sequestration: An on-farm

assessment. *Soil Sci Soc Am J* **72**, 693-701 (2008).

8. Liu C, Lu M, Cui J, Li B, Fang C. Effects of straw carbon input on carbon dynamics in

agricultural soils: a meta-analysis. *Glob Chang Biol* **20**, 1366-1381 (2014).

9. Lu FEI*, et al.* Soil carbon sequestrations by nitrogen fertilizer application, straw return

and no-tillage in China's cropland. *Glob Chang Biol* **15**, 281-305 (2009).

10. Song K*, et al.* Effects of tillage and straw return on water-stable aggregates, carbon

stabilization and crop yield in an estuarine alluvial soil. *Sci Rep* **9**, 4586 (2019).

11.   Prosdocimi M, Jordán A, Tarolli P, Keesstra S, Novara A, Cerdà A. The immediate

effectiveness of barley straw mulch in reducing soil erodibility and surface runoff

generation in Mediterranean vineyards. *Sci Total Environ* **547**, 323-330 (2016).

12.   Liang C, Schimel JP, Jastrow JD. The importance of anabolism in microbial control

over soil carbon storage. *Nat Microbiol* **2**, 17105 (2017).

13.   Zhang B, *et al.* Soil depth and crop determinants of bacterial communities under ten

biofuel cropping systems. *Soil Biol Biochem* **112**, 140-152 (2017).

14.   Nevins CJ, Nakatsu C, Armstrong S. Characterization of microbial community

response to cover crop residue decomposition. *Soil Biol Biochem* **127**, 39-49 (2018).

15.   Schmidt R, Gravuer K, Bossange AV, Mitchell J, Scow K. Long-term use of cover

crops and no-till shift soil microbial community life strategies in agricultural soil. *PloS*

ONE **13**, e0192953 (2018).

16.   Eilers KG, Debenport S, Anderson S, Fierer N. Digging deeper to find unique microbial

communities: the strong effect of depth on the structure of bacterial and archaeal

communities in soil. *Soil Biol Biochem* **50**, 58-65 (2012).

17.   Sagova-Mareckova M, *et al.* The structure of bacterial communities along two vertical

profiles of a deep colluvial soil. *Soil Biol Biochem* **101**, 65-73 (2016).

18.   Pries CEH, Castanha C, Porras R, Torn M. The whole-soil carbon flux in response to

warming. *Science* **355**, 1420-1423 (2017).

19.   Chorover J, Kretzschmar R, Garcia-Pichel F, Sparks DL. Soil biogeochemical

processes within the critical zone. *Elements* **3**, 321-326 (2007).

464 20. Richter DdB, Yaalon DH. " The Changing Model of Soil" revisited. *Soil Sci Soc Am J*

465   **76**, 766-778 (2012).

466 21. Liu X, *et al.* Soil degradation: a problem threatening the sustainable development of

467   agriculture in Northeast China. *Plant Soil Environ* **56**, 87-97 (2010).

468 22. Liu Z, Yang X, Hubbard KG, Lin X. Maize potential yields and yield gaps in the

469   changing climate of northeast China. *Glob Chang Biol* **18**, 3441-3454 (2012).

470 23. Canadell J, Jackson R, Ehleringer J, Mooney H, Sala O, Schulze E-D. Maximum

471   rooting depth of vegetation types at the global scale. *Oecologia* **108**, 583-595 (1996).

472 24. Koike I, Hattori A. Growth yield of a denitrifying bacterium, Pseudomonas denitrificans,

473   under aerobic and denitrifying conditions. *Microbiology* **88**, 1-10 (1975).

474 25. Miroshnichenko ML, *et al.* Caldithrix abyssi gen. nov., sp. nov., a nitrate-reducing,

475   thermophilic, anaerobic bacterium isolated from a Mid-Atlantic Ridge hydrothermal

476   vent, represents a novel bacterial lineage. *Int J Syst Evol Microbiol* **53**, 323-329

477   (2003).

478 26. De Vita P, Di Paolo E, Fecondo G, Di Fonzo N, Pisante M. No-tillage and conventional

479   tillage effects on durum wheat yield, grain quality and soil moisture content in southern

480   Italy. *Soil Tillage Res* **92**, 69-78 (2007).

481 27. Ramirez-Villanueva DA, *et al.* Bacterial community structure in maize residue

482   amended soil with contrasting management practices. *Appl Soil Ecol* **90**, 49-59

483   (2015).

484 28. Kremen C. Managing ecosystem services: what do we need to know about their

485   ecology? *Ecol Lett* **8**, 468-479 (2005).

486   29.   Zhalnina K, *et al.* Soil pH determines microbial diversity and composition in the park

487   grass experiment. *Microb Ecol* **69**, 395-406 (2015).

488   30.   Lauber CL, Hamady M, Knight R, Fierer N. Pyrosequencing-based assessment of soil

489   pH as a predictor of soil bacterial community structure at the continental scale. *Appl*

490   *Environ Microbiol* **75**, 5111-5120 (2009).

491   31.   Rousk J, *et al.* Soil bacterial and fungal communities across a pH gradient in an arable

492   soil. *ISME J* **4**, 1340 (2010).

493   32.   Lu C, *et al.* Effects of N fertilization and maize straw on the dynamics of soil organic N

494   and amino acid N derived from fertilizer N as indicated by 15 N labeling. *Geoderma*

495   **321**, 118-126 (2018).

496   33.   Liu X, *et al.* Linking microbial immobilization of fertilizer nitrogen to in situ turnover of

497   soil microbial residues in an agro-ecosystem. *Agric, Ecosyst Environ* **229**, 40-47

498   (2016).

499   34.   Zhang B, Wang H, Yao S, Bi L. Litter quantity confers soil functional resilience through

500   mediating soil biophysical habitat and microbial community structure on an eroded

501   bare land restored with mono Pinus massoniana. *Soil Biol Biochem* **57**, 556-567

502   (2013).

503   35.   Kibblewhite M, Ritz K, Swift M. Soil health in agricultural systems. *Philos Trans R Soc*

504   *Lond B Biol Sci* **363**, 685-701 (2008).

505   36.   Erbilgin O, McDonald KL, Kerfeld CA. Characterization of a planctomycetal organelle:

506   a novel bacterial microcompartment for the aerobic degradation of plant saccharides.

507   *Appl Environ Microbiol* **80**, 2193-2205 (2014).

37.   Herlemann DP, *et al.* Metagenomic de novo assembly of an aquatic representative of

the verrucomicrobial class Spartobacteria. *MBio* **4**, e00569-00512 (2013).

38.   Halsall DM, Goodchild DJ. Nitrogen fixation associated with development and

localization of mixed populations of Cellulomonas sp. and Azospirillum brasilense

grown on cellulose or wheat straw. *Appl Environ Microbiol* **51**, 849-854 (1986).

39.   Pathma J, Raman G, Sakthivel N. Microbiome of Rhizospheric Soil and Vermicompost

and Their Applications in Soil Fertility, Pest and Pathogen Management for

Sustainable Agriculture. *189-210 (Springer, Singapore, 2019).*

40.   Iino T, Mori K, Uchino Y, Nakagawa T, Harayama S, Suzuki K-i. Ignavibacterium

album gen. nov., sp. nov., a moderately thermophilic anaerobic bacterium isolated

from microbial mats at a terrestrial hot spring and proposal of Ignavibacteria classis

nov., for a novel lineage at the periphery of green sulfur bacteria. *Int J Syst Evol*

Microbiol **60**, 1376-1382 (2010).

41.   Terracciano J, Canale-Parola E. Enhancement of chemotaxis in Spirochaeta aurantia

grown under conditions of nutrient limitation. *J Bacteriol* **159**, 173-178 (1984).

42.   Thomas F, Hehemann J-H, Rebuffet E, Czjzek M, Michel G. Environmental and gut

bacteroidetes: the food connection. *Front Microbiol* **2**, 93 (2011).

43.   Archontoulis S, Licht MA. How Fast and Deep do Corn Roots Grow in Iowa? ,

(2017).

44.   Zhang C, Ju X, Powlson DS, Oenema O, Smith P. Nitrogen surplus benchmarks for

controlling N pollution in the main cropping systems of China. *Environ Sci Technol* **53**,

6678-6687 (2019).

45. Izewska A, Woloszyk C. Yields of grain and straw, their content and ionic proportions

of macroelements in maize fertilized with ash from municipal sewage sludge

combustion. *J Elementology* **20**, 319-329 (2015).

46. Kennedy S. Energy use in American agriculture. *Sustainable energy term paper* **5**,

1-26 (2000).

47. Zhang W-f, *et al.* New technologies reduce greenhouse gas emissions from

nitrogenous fertilizer in China. *Proc Natl Acad Sci USA* **110**, 8375-8380 (2013).

48. Goss-Souza D, Mendes LW, Borges CD, Baretta D, Tsai SM, Rodrigues JL. Soil

microbial community dynamics and assembly under long-term land use change.

FEMS Microbiol Ecol **93**, fix109 (2017).

49. Stegen JC, Lin X, Konopka AE, Fredrickson JK. Stochastic and deterministic

assembly processes in subsurface microbial communities. *ISME J* **6**, 1653 (2012).

50. Feng Y, Guo Z, Zhong L, Zhao F, Zhang J, Lin X. Balanced fertilization decreases

environmental filtering on soil bacterial community assemblage in north China. *Front*

Microbiol **8**, 2376 (2017).

51. Chase JM. Stochastic community assembly causes higher biodiversity in more

productive environments. *Science* **328**, 1388-1391 (2010).

52. Ferrenberg S, *et al.* Changes in assembly processes in soil bacterial communities

following a wildfire disturbance. *ISME J* **7**, 1102 (2013).

53. García-Orenes F, Morugán-Coronado A, Zornoza R, Scow K. Changes in soil

microbial community structure influenced by agricultural management practices in a

Mediterranean agro-ecosystem. *PloS ONE* **8**, e80522 (2013).

54.   IUSS Working Group W. World reference base for soil resources. *World Soil*

Resources Report **103**, FAO, Rome (2006).

55.   Nelson D, Sommers LE. Total carbon, organic carbon, and organic matter 1. *Methods*

of soil analysis Part 2 Chemical and microbiological properties, 539-579 (American

Society of Agronomy, Soil Science Society of America, 1982).

56.   Canadell J, Jackson RB, Ehleringer JB, Mooney HA, Sala OE, Schulze E-D. Maximum

rooting depth of vegetation types at the global scale. *Oecologia* **108**, 583-595 (1996).

57.   Rousk J, Baath E. Fungal and bacterial growth in soil with plant materials of different

C/N ratios. *FEMS Microbiol Ecol* **62**, 258-267 (2007).

58.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence

data. *Bioinformatics* **30**, 2114-2120 (2014).

59.   Derakhshani H, Tun HM, Khafipour E. An extended single-index multiplexed 16S

rRNA sequencing for microbial community analysis on MiSeq illumina platforms. *J*

Basic Microbiol **56**, 321-326 (2016).

60.   Mysara M, Leys N, Raes J, Monsieurs P. IPED: a highly efficient denoising tool for

Illumina MiSeq Paired-end 16S rRNA gene amplicon sequencing data. *BMC*

bioinformatics **17**, 192 (2016).

61.   Kozich JJ, Westcott SL, Baxter NT, Highlander SK, Schloss PD. Development of a

dual-index sequencing strategy and curation pipeline for analyzing amplicon

sequence data on the MiSeq Illumina sequencing platform. *Appl Environ Microbiol* **79**,

5112-5120 (2013).

62.   Quast C, *et al.* The SILVA ribosomal RNA gene database project: improved data

574   processing and web-based tools. *Nucleic acids research* **41**, D590-D596 (2012).

575   63.   Wang Q, Garrity GM, Tiedje JM, Cole JR. Naive Bayesian classifier for rapid

576   assignment of rRNA sequences into the new bacterial taxonomy. *Appl Environ*

577   *Microbiol* **73**, 5261-5267 (2007).

578   64.   Werner JJ, *et al.* Impact of training sets on classification of high-throughput bacterial

579   16s rRNA gene surveys. *ISME J* **6**, 94 (2012).

580   65.   Jiménez-Bueno N, *et al.* Bacterial indicator taxa in soils under different long-term

581   agricultural management. *J Appl Microbiol* **120**, 921-933 (2016).

582   66.   Aßhauer KP, Wemheuer B, Daniel R, Meinicke P. Tax4Fun: predicting functional

583   profiles from metagenomic 16S rRNA data. *Bioinformatics* **31**, 2882-2884 (2015).

584   67.   Parks DH, Tyson GW, Hugenholtz P, Beiko RG. STAMP: statistical analysis of

585   taxonomic and functional profiles. *Bioinformatics* **30**, 3123-3124 (2014).

Acknowledgments

We would like to thank Dr. William H. Schlesinger at the Cary Institute of Ecosystem

Studies for his comments and Dr. Randy Neighbarger at Duke University for language

editing. This work was supported by the "National Key R&D Program" (No.

2016YFD0800103, 2016YFD0200307) and the National Natural Science Foundation

of China (grant number, 41671297). We would like to thank Pengshuai Shao,

Xuesong Ma and many individuals for assistance with sample collection, processing

and analysis.

Author Contributions

H.X. and C.L. designed the experiment, F.D. measured soil samples, F.D., H.W. and

C.L analyzed data and wrote the manuscript, and all the authors discussed results and

commented on the manuscript.

Additional information

Supplementary information is available for this paper at ……

Competing interests

The authors declare no competing interests.

Correspondence and requests for materials should be addressed to C.L.
(cliang823@gmail.com)

Figure 1. Soil properties (mean±SE, n = 3) along soil depth under different practices.

SOC = soil organic carbon, SEOC = salt-extractable organic carbon, SEOC/SOC =

ratio of SEOC to SOC, SM = soil moisture, TN = total nitrogen content, C/N = ratio

of SOC to TN, NH_4^+-N = ammonium nitrogen, NO_3^--N = nitrate nitrogen, Inorganic N

= NH_4^+-N + NO_3^--N.

617 **Figure 2.** Corn yield (line+symbol) and annual rainfall during growing and dormant

618 seasons (bar) under different disturbance practices during 2013-2016. Error bars

619 indicate standard errors (n = 3 or 4), different letters indicate significant differences at

620 $P < 0.05$.

Figure 3. Microbial richness (Chao1), observed number of species (Observed_species) and diversity (Shannon_Index) in the CT (conventional tillage), NTNS (no-tillage without stover mulching) and NTSM (no-tillage with stover mulching) plots. Error bars indicate standard deviation (n = 3).

Figure 4. Indicator genera significantly (q < 0.1) associated with tillage practices. The

size of each circle represents the indicator value of a specific genus in the different

soil depths. The color indicates the relative abundance of each indicator genus.

Taxonomic information, indicator values, P-values, and q-values of all indicator

genera are given in Supplementary Table 2. Zero indicator genera were identified in

CT treatment.

Figure 5. Non-metric multidimensional scaling (MDS) ordination of soil microbial

community structures based on Bray-Curtis distances among soil depths at different

agricultural disturbance practices. Permutational multivariate analysis of variance

(PERMANOVA) revealed that the overall microbial community structures among soil

depth were significantly different at each disturbance practice. Circles, triangles and

squares represent CT (conventional tillage), NTNS (no-tillage without stover

mulching) and NTSM (no-tillage with stover mulching), respectively.

Figure 6. Redundancy analysis (RDA) of the soil microbial community originating

from microbial phyla constrained by soil properties under different agricultural

practices. Only soil variables that significantly explained variability in microbial

community structure in the forward selection procedure were selected to the

ordination (arrows). TN, total nitrogen content; C/N, a ratio of carbon to nitrogen

content; NH_4^+-N, ammonium nitrogen; NO_3^--N, nitrate nitrogen ; SM, soil moisture.

Circles, triangles and squares represent CT (conventional tillage), NTNS (no-tillage

without stover mulching) and NTSM (no-tillage with stover mulching), respectively.