Low-disturbance Farming Regenerates Healthy Deep Soil towards Sustainable Agriculture: Evidence from Long-term No-tillage with Stover Mulching

Fangbo Deng[1, 2#], Hongjun Wang[3#], Hongtu Xie[1#], Xuelian Bao[1], Hongbo He[1], Xudong Zhang[1], Chao Liang[1*]

[1] Key Laboratory of Forest Ecology and Management, Institute of Applied Ecology, Chinese Academy of Sciences, Shenyang 110016, China

[2] University of Chinese Academy of Sciences, Beijing 100049, China

[3] Duke University Wetland Center, Nicholas School of the Environment, Duke University, Durham, NC 27708, USA

[#] These authors contributed equally to this work.

[*] Corresponding author: Chao Liang (Email: cliang823@gmail.com)

Summary

Intensive conventional farming has degraded farmland topsoil and seriously threaten food

and environment security globally. Although low-disturbance practices have been widely

adapted to restore soil health, whether this measure in a long run can potentially recover the

critical deep soil to meet sustainable intensification of crop production are still unclear. Here

we compared soil microbiome, physicochemical parameters along 3-m deep soil profiles, and

crop yield in Northeast China subjected to ten years of farming practices at 3 levels of

disturbance, including conventional tillage (CT), no-tillage without stover mulching (NTNS),

and no-tillage with stover mulching (NTSM). We found that low-disturbance practices

(NTNS and NTSM) promoted the ability of the deep soil to retain water, nitrogen and

salt-extractable organic, regenerated whole-soil microbial diversity and metabolic function,

improved topsoil organic carbon stock and corn yield in the drought year, showed the

potential to reduce energy consumption and greenhouse gas emissions, thus regenerating

highly efficient, sustainable agriculture.

Since the Industrial Revolution, the rate of soil carbon loss has increased dramatically,

resulting in a global carbon debt due to agriculture of 116 Pg carbon for the top 2 m of soil[1].

The loss of soil carbon in farmlands has not only changed global climate but also produced

catastrophic cascade impacts on global food and environment security, as soil carbon is the

cornerstone for healthy and productive soil that will be needed to sustainably feed 10 billion

people in 2050 (United Nations, World Population Prospects 2019). It is well known that

intensive conventional farms with high energy inputs (chemical fertilizers) and disturbance

(e.g. tillage/compaction, burn/remove stover) have caused a series of environmental issues,

like water pollution, biodiversity loss, freshwater depletion and climate change[2]. Even worse,

increasing the amount of chemical fertilizer is unlikely to continue the increase in quantity

and quality of food products worldwide[3], while the options to expand the farmland area at the

expense of nature and biodiversity that already under pressure is limited[4]. Moreover, the

topsoil disturbance, tillage in particular, prevents root growth into deeper soil[5], thus critically

minimizing the exploitation of deep soil profile[6] and reducing crop's nutrient using efficiency

and its resilience to drought.

Since the 1970s, low-disturbance practices (e.g. reduced tillage, no-tillage and stover

mulching) have been gradually applied to restore soil health and reduce non-point source

pollution[7]. Growing evidence shows that no-tillage and stover mulching boosted top-soil

organic carbon (SOC)[8,9], increased soil aggregate stability[10] and reduced soil erosion and

surface runoff[11]. All these benefits from low-disturbance tillage practices are tied with

complex microbial processes that contribute to soil structure formation, modulate the

biogeochemistry cycling of nutrients and drive soil carbon transformation and

stabilization[12-14]. Microorganisms also respond quickly to changes in soil environment

conditions, due to their high surface to volume ratio, which can provide an early signal in soil

improvement or degradation, thus has been recommend as a key indicator of soil health[15].

However, the effect of tillage practices on soil microbial communities are complex and

diverse[16-19] and most studies by now mainly focused on farmland topsoil or soils within 1-m

depth[20-22], and soil below 1 meter, which contains more carbon than the topsoil[23] and belongs

to Earth's Critical zone, was often overlooked. However, microbes inhabiting in the deep

soils (> 1 m) may substantially impact long-term carbon sequestration, mineral weathering

and crop production[24-26], and play important roles in bridging aboveground vegetation with

parent soils and even acts as an essential buffer protecting underground water[27]. Moreover,

studies indicate many crops' roots (depending on the species and management) can penetrate

over one meter in depth[28,29], which means they can potentially forage for nutrient and water

in deep soil and impact deep-soil microbial community[30,31]. More importantly, tillage

practices significantly influence root growth[5,32,33], the change of surface soil carbon by land

use or agricultural practice influenced the carbon dynamic in deep soil layers has also been

observed[34-36]. We thus hypothesized that: 1) different tillage practices will affect the vertical

distribution of the microbial community by altering the spatial distribution of nutrients and

dissolved organic carbon, thereby changing the ecosystem function; 2) lowest disturbance

practice—no-tillage with 100% stover mulching as a nature-based solution, close to

undisturbed natural ecosystem, would regenerate healthy deep-soil with highly diversified

and functional microbes over time toward a highly resilient, sustainable agricultural

ecosystem.

Recent research shows that corn belts in the U.S.A., western Europe, and China have

experienced the most soil carbon loss globally[1]. The corn belt in Northeast China is

considered as the "breadbasket" of the country, having the largest grain production and

overlapping with the most fertile Mollisol region that sustains 3% of population in the

world[37], accounting for over 30% of corn production of China[38]. Here, a 10-year

manipulative experiment was conducted at a temperate corn farm in Northeast China,

investigating   farming   practices   with   three   levels   of   disturbance: high

disturbance—conventional tillage (CT), low disturbance—no-tillage without stover mulching

(NTNS) and lowest disturbance—no-tillage with 100% stover mulching (NTSM). We

compared soil physicochemical properties, fine-root associated organic carbon, and microbial

communities of the 3-m soil profiles at the end of dormant season (legacy effects of nutrients)

after the 10-year manipulation, and multi-year corn yield as well. We aimed at testing our

hypotheses and providing a general picture of soil microbial community in the entire soil

profiles, which are important for comprehensive recognize the impact of low-disturbing

tillage practices on ecosystem function and developing effective strategies to ensure the

sustainable development of agriculture.

Results

Soil properties and corn yield

The SOC, TN and C/N ratio substantially decreased from the soil surface to around 150

cm depths and then remained unchanged within 150-300 cm (Fig. 1). The NTSM slightly

increased SOC, TN and C/N ratio at 0-20 cm soil layers compared with the NTNS and the CT

(Fig. 1 and Table S1). The NTSM and NTNS reduced soil pH in surface and deeper layers

(Fig. 1d) and increased soil moisture at surface layers (0-60 cm) (Fig. 1e). In the CT plots,

soil NO_3^--N concentration first decreased and then increased remarkably, ranged from 4.19 to

23.32 mg kg^{-1} (Fig. 1i). However, under the NTNS and NTSM treatments, soil NO_3^--N

decreased significantly at 0-40 cm then increased to the maximum at 120-150 cm.

Interestingly, above 120-150 cm layer, NO_3^--N was significantly higher with low-disturbance

practices than conventional tillage, while the soil below 150 cm under low-disturbance

practices had much lower NO_3^--N compared to conventional tillage (Fig. 1i). The NTNS plots

contained much higher amounts of ammonium than the CT and the NTSM plots (Fig. 1h).

Soil salt-extractable organic carbon (SEOC)—a proxy for biotically-derived organic acid

declined from the surface to 40-60 cm and then increase to its peak at 60-90 cm under CT, at

90-120 cm under NTNS and at 120-150 cm under NTSM (Fig. 1b). The NTSM increased the

SEOC concentration at almost all soil layers compared with the CT and the NTNS (Fig. 1b),

in which at the surface and 120-150 cm depth the contents of SEOC with NTSM were twice

higher than CT. The relative contributions of SEOC to SOC (SEOC/SOC) in the NTSM were

also always higher than in the CT and NTNS (Fig. 1c). Based on the estimated root depths

and soil bulk density, total soil inorganic nitrogen available for the coming growing season in

the NTSM and the NTNS was approximate 427.34 and 352.34 kg ha^{-1}, respectively, while

only 179.63 kg ha^{-1} in conventional tillage.

The mean annual corn yield (2013-2016) in the NTSM is 13416.8 kg/ha, which is much

higher than the CT and NTNS (Fig. 2), particularly during the drought year of 2015, with

only 409.6 mm of rainfall during the growing season (about 100 mm lower than the mean

rainfall), while the corn yield in NTSM is 36.4% and 22.3% higher than the CT and NTNS,

respectively (Fig. 2).

Microbial diversity, composition, and structure.

The microbial richness (Chao1), observed number of species (Observed-species) and

diversity (Shannon-Index) first increased within 0-20 cm and decreased from 20 to 90 cm,

then increased hereafter (Fig. 3). The low-disturbance practices significantly increased Chao1,

Observed-species and Shannon-Index, particularly in 0-40 cm soil depths (Fig. 3). There

were 54 microbial phyla across all soil samples. The dominant phyla (relative abundance > 1%

across all soil samples) were Proteobacteria, Actinobacteria, Chloroflexi, Acidobacteria,

Nitrospirae, Gemmatimonadetes, Planctomycetes, and these phyla accounted for 60-91% of

the total microbial abundances in the whole soil profile (Fig. S1a). Bacteroidetes,

Verrucomicrobia, Latescibacteria, Parcubacteria, Firmicutes, Microgenomates and

Saccharibacteria were less dominant (relative abundance > 0.1% across all soil samples) but

were still found across all soil samples (Fig. S1a). Although no difference in the composition

of dominant phyla among treatments was found, there were more non-dominant phyla with

higher relative abundance in low disturbance practices than conventional tillage practice (Fig.

S1b).

Indicator analysis identified 16 and 51 clearly classified genera (relative abundances >

0.005%) in the NTNS and the NTSM plots, respectively, while no indicator genera were

found in the conventional tillage plots (Fig. 4 and Table S2). The indicator genera in the

NTNS plots belonged to Proteobacteria, Actinobacteria, Chloroflexi, Gemmatimonadetes and

Planctomycetes, and most of them appeared in the surface soil (0-20 cm) with only 1 genus

below 150 cm. Importantly, more extra indicator genera — including Bacteroidetes,

Acidobacteria, Deferribacteres, Firmicutes, Verrucomicrobia, Chlorobi and Spirochaetae —

existed in the NTSM plots, in which under 150 cm we observed 7 genera (Fig. 4 and Table

S2).

Microbial community structures were visualized by Non-metric multidimensional scaling

(MDS) and tested by PERMANOVA based on Bray–Curtis. The microbial communities

among treatments in the root zones were marginally different (PERMANOVA p=0.08);

however, below the root zone they differed distinctively (PERMANOVA p=0.02). The

disturbance practices influenced the vertical distribution dissimilarity in microbial

community structure (Fig. 5). Three clusters — 0-10 cm and 10-20 cm, 20-150 cm and

150-300 cm — were observed in the CT plots (PERMANOVA-F=9.57, p=0.0001) (Fig. 5). In

the NTNS plots, 0-10 cm formed an independent cluster, while other soil depths showed

some separation (e.g. 20-120 cm were separated from 150-300 cm soil depths by axis 1);

however, Bray-Curtis distances between adjacent depths were too close to be separated

(PERMANOVA-F=8.18, p=0.0001) (Fig. 5). The NTSM treatment clustered 0-10 cm and

10-20 cm together, 120-150 cm, 150-200 cm, 200-250 cm and 250-300 cm separately, and the

other depths show some separations as well (PERMANOVA-F=11.32, p=0.0001) (Fig. 5).

Predicted Ecological functions of microbial communities.

According to the results of microbial diversity, composition and structure, the metabolic

capabilities of microbial community in the whole 3-m soil profiles were predicted using

Tax4Fun (Fig. S2). Results showed that low-disturbance practices significantly increased the

abundance of predicted functions related to carbohydrate metabolism, nucleotide metabolism,

glycan biosynthesis and metabolism, lipid metabolism and metabolism related to cofactors

and vitamins (Fig. S2a). Moreover, the relative abundances of genes encoding for

assimilatory nitrate reduction in low-disturbance practices were higher than that in

conventional tillage practice (Fig. S3). The results suggested that in low disturbance practices,

microbial community prefer to convert the nitrate/nitrite to ammonia. We then further

assessed the impact of stover mulching on functional profiles (Fig. S2b). The extended error

bar plot shows that the NTNS enriched the abundance of amino acid metabolism and lipid

metabolism, while the NTSM enriched the functions associatedto energy metabolism,

carbohydrate metabolism, biosynthesis of secondary metabolites, glycan biosynthesis and

metabolism as well as metabolism of cofactors and vitamins (Fig. S2b).

Relationships between microbial communities and soil properties

Forward selection in Redundancy analysis (RDA) revealed that soil depth (pseudo-F=48, p=

0.002), SOC (pseudo-F=11.5, p= 0.002), SM (pseudo-F=3.4, p= 0.012), soil pH

(pseudo-F=2.3, p=0.018) and soil NH_4^+-N (pseudo-F=2.7, p= 0.026) significantly affected the

vertical distribution of microbial communities (Fig. S4). Furthermore, the soil properties that

regulated the distribution of soil microbes were different under different disturbance practices.

Under the CT treatment, soil microbial community was mainly affected by soil NH_4^+-N

(pseudo-F=4, p= 0.002) and soil NO_3^--N (pseudo-F=2.3, p= 0.012) that mainly came from

applied fertilizer (Fig. 6). The microbial community positively correlated to soil NH_4^+-N in

the 0-20 cm soil, to soil NO_3^--N negatively within 20-150 cm, while to soil NO_3^--N positively

after 150 cm (Fig. 6). Under the NTNS treatment, soil pH (pseudo-F=3.7, p=0.004)

constrained the distribution of the microbial community, in which strong negative

correlations occurred in 0-10 cm soil and a positive correlation in 90-150 cm (Fig. 6). Under

the NTSM treatment, soil TN (pseudo-F=11, p=0.002), SM (pseudo-F=2.6, p=0.004) and C/N

ratio (pseudo-F=1.8, p=0.016) significantly influenced the soil microbial community

separation (Fig. 6). In general, the microbes positively correlated with the soil TN and C/N

ratio in the surface soil layers (0-40 cm) and with SM in the middle layers (40-150 cm), while

they were mainly influenced by depth in the deeper soil (150-300 cm) (Fig. 6).

Root depth among tillage practices

Currently the measurement of fine roots with diameter less than 0.2 mm are still technically

difficult and their high turnover rates in-situ make the measurement even more complex[31].

Instead of detecting fine roots directly, we measured SEOC—a proxy for biotically-derived

organic acid, which is a sensitive signal of root density and could be an indicator of root

depth[6]. We also did a literature review on whether no-tillage and straw mulching extend root

depth in other crops. According to the change of SEOC with soil depth under different

disturbing practices(Fig 1b), we estimate that corn roots reached up to 60-90 cm, 90-120 cm

and 120-150 cm under the CT, the NTNS and the NTSM, respectively, which is in line with

reported corn root depths (~150 cm) [5,39].According the literature review, we can't find

root-depth studies including both no-tillage and straw mulching. However, no-tillage as a

conservative management increases root depth by 23 cm, 39 cm and 14 cm in corn, wheat,

and sunflower, respectively (Fig. S5).

Discussion

No-tillage practices promote soil nutrient, water holding capacities and corn yield.

The changes of SEOC with soil depth and the literature review of root depth among tillage indicated that no-tillage could promote root growth into deep soil, up to 150 cm in the NTSM. The root exudates with various organic acid and dead roots likely contributed to the lower soil pH in the NTSM, as shown by a significant negative relationship between SEOC and pH (r=−0.678, p<0.05). Despite function predication analyses have various uncertainty, the increased dissimilatory and assimilatory nitrate reduction genes suggested that low disturbance practices tend to convert nitrate to ammonium (Fig. S3), while lower ammonium in NTSM may be due to ammonium subsequently incorporated into biomass, as hints by the increase in genes related to biosynthesis metabolism and glycometabolism (Fig. S2). Moreover, maize stover or increased fine roots in deeper soil in NTSM provided labile carbon (Fig. 1b)to remove leaked nitrate through denitrification in deeper soil (below 1.5 m), where higher relative abundance of the denitrification bacteria (*Pseudomonas* and *Caldithrix*)[40,41] (Fig. 4 and Table. S2) were detected. However, shallower roots in the CT treatment can't provide enough labile carbon to remove extra soil NO_3^--N in deep soil, thus causing nitrite accumulation and leaching into deeper soil layers. The amount of inorganic nitrogen accumulated in the root zones under NTSM (427.34 kg ha^{-1}) likely could provide plenty of nitrogen for corn growth in the coming growing season (Fig. 2), based on the removed nitrogen in the grain (~200 kg ha^{-1}). Additionally, in line with many studies that show stover mulching reduces water evaporation and surface runoff and increase soil moisture in top soils[11,42], we found that the topsoil moisture was significantly higher in the NTSM than in the CT plots (Fig. 1e) when the precipitation during the entire dormant period was only 38.5 mm. More importantly, the topsoil (0-20 cm) SOC stock in NTSM (18.5 Mg/ha) were significantly higher than in CT (15.8 Mg/ha) due to slightly higher bulk density in NTSM. Therefore, no-tillage with stover mulching not only increased SOC stock, the holding capacities for

nutrients and water, thus reducing energy input to farm, but also tended to reduce the risk of

nitrate leaching into groundwater. And more importantly, the healthy deep soil in turn raises

corn production and promote the crop resistance to drought (Fig. 2). All these are critical to

the development of sustainable agriculture and the associated ecosystems.

No-tillage with stover mulching promotes microbial diversity, richness, and ecological

function contributing to sustainable farming.

Under the CT treatment, tillage heavily disturbed the topsoil and liberated occluded organic

materials. Microbes tended to rapidly use available nutrients in the plowed layer (e.g.

NH_4^+-N)[43], thereby causing the reduction of microbial metabolic diversity (Fig. S2a). Then,

the resistance of the soil to stress or disturbance may also decrease[44]. In deeper soil layers,

due to shallower roots, NO_3^--N could quickly move downward and accumulate in deeper soil

(Fig. 1i), which not only contaminated the underground water but also limited the activity of

non-dominant microbes with important ecological functions, as no indicator genera were

identified for each soil depth in CT treatment (Fig. 4 and Table S2). Because the microbial

communities were closely associated with inorganic nitrogen, the microbes under CT were

mainly influenced by added chemical fertilizer[45]. Although the dominant microbial

communities in CT were similar to those in the NTNS and NTSM, the loss of function

resulted from the difference of non-dominant microbes, indicating that the soil under CT had

degraded.

Under the NTNS treatment, soil pH was the major edaphic factor affecting the microbial

community and the indicator genera (Fig. 6 and Table S3). The lower soil pH possibly was

caused by deeper roots as shown by higher SEOC that is generally positively related to root

density[6]. Soil pH is often observed as a major factor determining the microbial composition

and structure in natural ecosystems[46,47], as microbes often show a narrow tolerance to soil pH.

In addition, soil pH regulates the availability of nutrient and mitigate ion toxicity[46-48]. Under

NTNS, soil pH and depth explained 35% distribution of the microbial community (Fig. 6).

We speculated that other edaphic factors (e.g. salinity and iron) directly or indirectly related

to soil pH and SEOC also influenced the changes in the microbial community.

Under NTSM treatment, TN and C/N significantly correlated with soil microbial

community due to the high C/N ratio of stover or roots (Fig. 6). Prior studies have reported

that, following maize stover mulching, more organic N, amino acid N, and amino sugar N

were observed in soil[49,50], combining with higher genes related to amino acid metabolism and

carbohydrate metabolism from function predication analyses (Fig. S2), jointly hint that

NTSM increased the retention time of nitrogen, hence meeting the nutrient requirement of

corn growth and reducing nitrate loss to underground water. The increased available nitrogen,

labile carbon and water in deep soil under NTSM can increase the resilience and resistance of

maize to disturbances with higher grain production (Fig. 2). Zhang et al.[51] also observed

litter-covered soil showed greater resistance to heating and copper addition due to the

changes in soil properties and microbial community structure. Resistance to disturbance or

stresses is the nature of a healthy soil and is essential for maintaining ecosystem functions,

such as decomposing organic matter[51,52]. Under the NTSM treatment, the microorganisms

associated with the degradation of relatively stable carbon compounds, such as

Planctomycetes and Verrucomicrobia (Table S4)[53,54] as well as the indicator *Cellulomonas*

and *Azospirillum* (Fig. 4 and Table S2) with the function of cellulose decomposition[55,56] were

increased. The predicted functional profiles related to energy metabolism (Carbon fixation

pathways in prokaryotes), carbohydrate metabolism (TCA cycle, amino sugar, nucleotide

sugar, galactose, fructose), biosynthesis of secondary metabolites (Carotenoid and Betalain)

and glycan biosynthesis were increased, suggesting a higher metabolic activity and a change

in substrate quality (Fig. S2). In addition, stover mulching also increased the ecological filter

function of soil depth for selecting microbial communities as more indicator genera of each

soil depths were identified under NTSM compared to NTNS and CT practices (Fig. 4 and

Table S2). And these indicators residing at different soil depths might enhance the

anti-disturbance ability of NTSM. For example, denitrification bacteria *Caldithrix* and

Pseudomonas[40,41] were the indicator genera of 150-200 cm and 250-300 cm, respectively (Fig.

4 and Table S2), which might explain the low nitrate in the deep soil in NTSM.

Ignavibacteria and *Spirochaeta,* the indicator genera of deep soil, have the ability to grow

under the conditions of strictly anaerobic[57] and severely limited nutrients[58], respectively.

Surface indicator genera belonging to Bacteroidetes might have the ability to degrade organic

matter that is difficult to decompose[59].

Implications for climate change and food security.

It was observed that about 179.63, 352.34 and 427.34 kg ha^{-1} inorganic N were kept in the

root-zone soil in the CT, NTNS and NTSM, respectively. Generally, corn roots reach their

maximum depth at the silking stage[60], which is also the time when the heaviest rainfall occurs

in northeastern China. We therefore expect that the available N kept in the root zone would be

utilized by crops in the coming growing season before the storm leach the nitrogen down to

ground water, which means that fertilizer N could be cut to meet crop growth in, at least,

Northeast China and also prevent reactive N losses. Since the nitrogen use efficiency (NUE)

of maize system under the conventional management is 51% in northeast China (NUE is

defined as the efficiency of fertilizer N transferring to harvested crop N)[61]. Then, we

conservatively calculate the required fertilizer N in the next year based on two assumptions: 1)

the NUE of soil available N in root zone is equal to that NUE of applied fertilizer N, both of

them are 50%; 2) the mineralized N during the coming growing season is neglected. Thus, N

supply requirement = Fertilizer N×NUE + N in root zone ×NUE + Stover-N, where Stover-N

for NTSM is 60 kg ha^{-1}. We estimated the N requirement for each disturbance practice by

multiplying grain yield by grain N concentration (1.4%)[61] plus multiplying stover yield by

stover N concentration (0.8%)[62]. For CT, NTNS and NTSM, the mean annual corn yields

were 10946.74, 12487.81 and 13416.81 kg ha^{-1}, and the stover yields were 966.67, 10083.33

and 10833.33 kg ha^{-1}, respectively. Thus, the N requirements were 230.6, 255.5 and 274.5 kg

ha^{-1} for CT, NTNS and NTSM, respectively. Therefore, the theoretically conservative

amounts of fertilizer N in the coming growing season are 281.6, 158.7 and 1.7 kg ha^{-1} for CT,

NTNS and NTSM, respectively. No fertilizer-N is needed to apply without reducing corn

yield in the NTSM plot. Compared to CT, the NTNS and NTSM could at least save

respectively about 122.9 and 281.6 kg ha^{-1} N-fertilizer. For every kilogram of fertilizer-N

produced and used on cropland, up to 87.9 MJ of energy is consumed[63] and 13.5 kg of

CO_2-equivalent (eq) (CO_2-eq) is emitted[64]. Hence, totally 24,752.6 MJ of energy

consumption could be reduced and 3,801.6 kg CO_2-eq emission could be cut per hectare

cornland in Northeast China at least by using NTSM tillage practice. If this could be applied

to all maize farmland in Northeast China (13,000,000 ha, Source: China Statistics Yearbook

2018), 0.3 EJ of energy could be saved and 49.4 Mt of CO_2-eq could be reduced. Based on

the average annual energy consumption for households of China in 2017 (15 EJ, China

Statistics Yearbook 2018) and CO_2 emissions (9,839 Mt, Global Carbon Atlas), the NTSM

practice in corn farming of Northeast China has the potential to save 2% of household energy

and to reduce 0.5% of CO_2 emissions each year in the whole country.

Our results, particularly higher SEOC content, microbial diversity, and indicator genera in the

NTSM deep soil compared with CT, clearly showed that low-disturbance practices can affect

agricultural resource in deeper soil over time. This indicates that crops, like corn in this study,

under appropriate management can ultimately explore nutrients and water from deeper soil,

thus not only increasing the volume of soil (almost double in this study) exploited without

reclaiming more natural land areas but also reducing nutrient loss into ground water.

Meanwhile, the input of labile carbon including SEOC into deeper soil in the NTSM provides

essential energy and nutrients to microbes and gradually shape highly diversified and

functional microbial communities in the deep soil over time, and hence improve the

self-sustaining ability of farmland in the face of climate change. The improvements of

microbial communities in the deep soil (1-3 m) at the end of dormant season in our study

provide evidence for the first time that a nature-based management in farmland is conducive

to deep-soil health for sustainable farm in a long run. Although many scientists have realized

the importance of deep rooting for sustainable intensification of crop production[30,31], the

deep-root studies are still rare due to technological bottleneck. According the literature review,

no-tillage extend root depth in other crops, like corn, wheat and sunflower. While some

studies also show no effects or even reduce root depth[65,66], the possible reason is the legacy

effects of tillage, for example, long-term no-tillage leads to soil compaction, and the soil

system might be still in its transition stage. Coupled with our results, low-disturbance

practices as a nature-based agricultural management likely can develop a deeper root system

to explore more resource in deep soil to sustain food production, while without threatening

environmental security and reclaiming more lands.

Methods

Site description and soil sampling.

The field experiment was established in 2007 at the Lishu Conservation Tillage Research and

Development Station of the Chinese Academy of Sciences in Jilin province, Northeast China

(43.19° N, 124.14° E). The region has a humid continental climate with a mean annual

temperature of 6.9 °C and the mean annual precipitation of 614 mm. The soils are classified

in the Mollisol order (Black Soil in Chinese Soil Classification) with a clay loam texture[67].

The site has been continuously planted with maize since 2007. We set up an experiment by a

randomized complete block design with four replicates and five treatments. Each plot area

was $261m^2$ (8.7×30m). The five treatments included conventional tillage (moldboard plowing

to a depth around 30 cm and removed the stover), no-tillage (no soil disturbance and direct

seeding), and no-tillage with three-level stover mulching (33%, 67% and 100% newly

produced maize stover were evenly spread over the soil surface each fall). For each treatment,

slow-release fertilizer was applied at one time when sowing, which was equal to 240 kg/ha N;

47 kg/ha P; 90 kg/ha K. The rainfall data were obtained from local meteorological

administration. The grain yield was estimated by manually harvesting 20 m^2 area, randomly

taken from each plot.

In this experiment, in order to reduce the damage to the plots and reduce costs, 3 plots

were randomly taken from each treatment including conventional tillage (CT), no-tillage

without stover mulching (NTNS), no-tillage with 100% stover coverage (NTSM) as three

comparative practices. In April 2017, triplicate soil cores (0-300 cm) were collected from

each plot at the end of dormant season. After removing surface stover, we took soil cores by a

stainless-steel hand auger and sliced each into ten layers: 0-10 cm, 10-20 cm, 20-40 cm,

40-60 cm, 60-90 cm, 90-120 cm, 120-150 cm, 150-200 cm, 200-250 cm, 250-300 cm. In total,

90 soil samples were collected and transported to the laboratory within 3 hours, then passed

through a 2-mm sieve. All visible roots, crop residues and stones were removed. Each soil

sample was divided into three subsamples: one subsample for DNA extraction and soil

salt-extractable organic carbon (SEOC) measurement that was immediately placed into a

polyethylene plastic bag and stored at -80 °C, one for chemical measurements including

ammonium nitrogen (NH_4^+-N) and nitrate nitrogen (NO_3^--N) (within one day), and the

remaining one was air dried for other soil physicochemical properties.

Soil properties

Soil total nitrogen (TN) content was measured by an Element analyzer Vario EL III

(Elementar Analysensysteme GmbH, Hanau, Germany). Soil organic carbon (SOC) was

converted from soil organic matter that was measured by potassium dichromate oxidation[68].

Soil pH was measured in deionized free-CO_2 water (1:2.5 w/v). Gravimetric soil moisture

was determined by oven-drying fresh soil to a constant weight at 105 °C. Soil NH_4^+-N and

NO_3^--N were extracted from fresh soil by 2 M KCl and measured by a continuous flow

analytical system (AA3, SEAI, Germany). To reflect soil soluble, exchangeable,

mineral-bound OC, SEOC was extracted from the frozen soil samples with 0.5 M K_2SO_4 (1:5

w/v)[69,70].

DNA extraction, PCR amplification and pyrosequencing

Soil DNA was extracted from the frozen soil samples (0.5 g wet weight) by using MoBio

PowerSoil DNA isolation kit (MoBio Laboratories, Carlsbad, CA, USA) following the

instructions of the manufacturer. The quality of DNA was determined by 1% agarose gel

electrophoresis. The V3–V4 region of the bacterial 16S rRNA gene was amplified by PCR

using the primers 338F and 806R with barcode for Illumina MiSeq sequencing. PCR was

performed in a total volume of 50 µl containing 30 ng DNA as a template, 20 mol of each

primer, 10mM dNTPs, 5µl   Pyrobest buffer (10×) and 0.3 U of Pyrobest polymerase (Takara

Code: DR005A). Each sample was amplified for three replicates. The PCR products from the

same sample were pooled, checked by 2% agarose gel electrophoresis and were then purified

using AxyPrepDNA agarose purification kit (AXYGEN). Finally, purified PCR products

were sequenced on an Illumina MiSeq platform PE300 sequencer (Illumina, USA).

The raw sequence data were further analyzed by the following protocol. Low-quality

sequences with an average quality score of less than 20 were filtered by employing

Trimmomatic[71]. The FLASH software was used to merge overlapping ends and treat them as

single-end reads[72]. The non-amplified region sequences, chimeras and shorter tags were also

removed using Usearch and Mothur[73]. The resulting high-quality sequences were clustered

into Operational Taxonomic Units (OTUs) at 97% sequence similarity using Usearch

(Version 8.1.1861 http://www.drive5.com/usearch/). OTUs were then classified against the

Silva (Release119 http://www.arb-silva.de) database and the taxonomic information of each

OTU representative sequence was annotated using the RDP Classifier[74]. A total of 3,255,693

high-quality reads were obtained from all soil samples, which were clustered into 9,573

unique OTUs at a 97% sequence similarity. The Good's coverage of all the samples ranged

from 0.93 to 0.98, which indicates an adequate level of sequencing to identify the majority of

diversity in the samples.

Root depth data

Plant roots, fine roots in particular, release large amounts of labile organic carbon[75] that are

essential for healthy microorganisms in soil. Generally, most studies considered that fine

roots are roots < 2mm in diameter, while roots <0.2 mm in diameter can contribute to >50 %

of the overall root length and play a major role in releasing root exudates and absorbing

nutrients and waters[31]. Currently the measurement of fine roots with diameter less than 0.2

mm are still technically difficult and their high turnover rates *in-situ* make the measurement

even more complex[31]. To minimize the root-turnover effects to the most degree, we collected

soil samples at the end of dormant season, which can likely mirror the long-term legacy

effects of our practices. Also instead of detecting fine roots directly, we measured soil

salt-extractable organic carbon (SEOC)—a proxy for biotically-derived organic acid, which is

a sensitive signal of root density and could be an indicator of root depth[6]. We also did a

literature review on whether no-tillage and straw mulching extend root depth in other crops.

We collected root depth of main crops data from literature published in 1996-2019 using the

Web of Science resource and only the data with more than two tillage managements are

retained. Finally, a total of 87 observations were selected (Table S5).

Statistical analyses

Soil properties were analyzed and plotted using Sigmaplot 12.5 software. Alpha

diversity indices were calculated in Qiime (version v.1.8) and used to reflect the diversity and

richness of the microbial community in different samples. The relative abundances of

individual phyla in different samples were computed by R packages. The indicator analysis

based on genera-specific to each soil depth was conducted using indicspecies package of R

with 9999 permutations, and the P-values were corrected for multiple testing using qvalue

package of R[76]. Functional profiles of the microbial community were predicted by Tax4fun

(an open-source package in R)[77] and further statistical analysis was conducted by STAMP

using Welch's t-test [78]. Non-metric multidimensional scaling (MDS) was performed by

"vegan" package of R to describe differences in microbial community structure among

samples. Permutational multivariate analysis of variance (PERMANOVA) was employed on

Bray-Curtis distances to test the differences in soil microbial communities among various

sample groups. The redundancy analysis (RDA, Canoco 5 software) were conducted to

identify the correlations between microbial community composition and environmental

variables. ANOVA were conducted by SPSS Version 22. Percentage data were transformed

using arcsine square root function before ANOVA test. All statistical tests were significant at

$p \leq 0.05$.

Acknowledgements

We would like to thank Dr. William H. Schlesinger at the Cary Institute of Ecosystem Studies

for his comments and Dr. Randy Neighbarger at Duke University for language editing. This

work was supported by the "National Key R&D Program" (No. 2016YFD0800103, No.

2016YFD0200307) and the National Natural Science Foundation of China (grant number,

41671297). We would like to thank Pengshuai Shao, Xuesong Ma and many individuals for

assistance with sample collection, processing and analysis.

Author Contributions

H.X., XZ and C.L. designed the experiment, F.D. did field and lab measurements, F.D., H.W.

and C.L analyzed data and wrote the manuscript, and all the authors discussed results and

commented on the manuscript.

Declaration of Interests

The authors declare no competing interests.

Sequence availability

All sequencing data that support the findings of this study have been deposited in the

National Center for Biotechnology Information (https://www.ncbi.nlm.nih.gov/), in the

Sequence Read Archive (SRA) database (BioProject number: PRJNA488172).

Supplementary information

Supporting information may be found online in the Supporting Information section at the end

of the article.

References

1 Sanderman, J., Hengl, T. & Fiske, G. J. Soil carbon debt of 12,000 years of human land use.
Proc. Natl. Acad. Sci. U.S.A. **114**, 9575-9580 (2017).
2 Congreves, K., Hayes, A., Verhallen, E. & Van Eerd, L. Long-term impact of tillage and crop
rotation on soil health at four temperate agroecosystems. *Soil Tillage Res.* **152**, 17-28 (2015).
3 Vitousek, P. M. *et al.* Nutrient imbalances in agricultural development. *Science* **324**,
1519-1520 (2009).
4 Springmann, M. *et al.* Options for keeping the food system within environmental limits.
Nature **562**, 519-525 (2018).
5 Kemper, W. D., Schneider, N. N. & Sinclair, T. R. No-till can increase earthworm populations
and rooting depths. *J. Soil Water Conserv.* **66**, 13A-17A (2011).
6 Billings, S. A. *et al.* Loss of deep roots limits biogenic agents of soil development that are
only partially restored by decades of forest regeneration. *Elem Sci Anth* **6**, 34 (2018).
7 Salem, M. A. Economics of reduced tillage technology on soil conservation and risk analysis
for Eastern Oklahoma farmers. *(Oklahoma State University, 1983)*.
8 Blanco-Canqui, H. & Lal, R. No-tillage and soil-profile carbon sequestration: An on-farm
assessment. *Soil Sci. Soc. Am. J.* **72**, 693-701 (2008).

9   Liu, C., Lu, M., Cui, J., Li, B. & Fang, C. Effects of straw carbon input on carbon dynamics
480   in agricultural soils: a meta☐analysis. *Glob. Chang. Biol.* **20**, 1366-1381 (2014).

481   10   Song, K. *et al.* Effects of tillage and straw return on water-stable aggregates, carbon
482   stabilization and crop yield in an estuarine alluvial soil. *Sci. Rep.* **9**, 4586 (2019).

483   11   Prosdocimi, M. *et al.* The immediate effectiveness of barley straw mulch in reducing soil
484   erodibility and surface runoff generation in Mediterranean vineyards. *Sci. Total Environ.* **547**,
485   323-330 (2016).

486   12   Liang, C., Schimel, J. P. & Jastrow, J. D. The importance of anabolism in microbial control
487   over soil carbon storage. *Nat. Microbiol.* **2**, 17105 (2017).

488   13   Schimel, J. & Schaeffer, S. M. Microbial control over carbon cycling in soil. *Front. Microbiol.*
489   **3**, 348 (2012).

490   14   Bardgett, R. D. & Van Der Putten, W. H. Belowground biodiversity and ecosystem
491   functioning. *Nature* **515**, 505-511 (2014).

492   15   Nielsen, M. N., Winding, A. & Binnerup, S. Microorganisms as indicators of soil health.
493   *National Environmental Research Institute, Denmark.* (2002).

494   16   Helgason, B., Walley, F. & Germida, J. Long-term no-till management affects microbial
495   biomass but not community composition in Canadian prairie agroecosytems. *Soil Biol.*
496   *Biochem.* **42**, 2192-2202 (2010).

497   17   Rincon-Florez, V. A. *et al.* Significant effects on soil microbial communities were not
498   detected after strategic tillage following 44 years of conventional or no-tillage management.
499   *Pedobiologia*, 150640 (2020).

500   18   Li, Y., Zhang, Q., Cai, Y., Yang, Q. & Chang, S. X. Minimum tillage and residue retention
501   increase soil microbial population size and diversity: Implications for conservation tillage. *Sci.*
502   *Total Environ.* **716**, 137164 (2020).

503   19   Sun, R. *et al.* Tillage changes vertical distribution of soil bacterial and fungal communities.
504   *Front. Microbiol.* **9**, 699 (2018).

505   20   Nevins, C. J., Nakatsu, C. & Armstrong, S. Characterization of microbial community
506   response to cover crop residue decomposition. *Soil Biol. Biochem.* **127**, 39-49 (2018).

507   21   Alahmad, A. *et al.* Cover crops in arable lands increase functional complementarity and
508   redundancy of bacterial communities. *J. Appl. Ecol.* **56**, 651-664 (2019).

509   22   Hartman, K. *et al.* Cropping practices manipulate abundance patterns of root and soil
510   microbiome members paving the way to smart farming. *Microbiome* **6**, 1-14 (2018).

511   23   Jobbágy, E. G. & Jackson, R. B. The vertical distribution of soil organic carbon and its
512   relation to climate and vegetation. *Ecol. Appl.* **10**, 423-436 (2000).

513   24   Eilers, K. G., Debenport, S., Anderson, S. & Fierer, N. Digging deeper to find unique
514   microbial communities: the strong effect of depth on the structure of bacterial and archaeal
515   communities in soil. *Soil Biol. Biochem.* **50**, 58-65 (2012).

516   25   Sagova-Mareckova, M. *et al.* The structure of bacterial communities along two vertical
517   profiles of a deep colluvial soil. *Soil Biol. Biochem.* **101**, 65-73 (2016).

518   26   Pries, C. E. H., Castanha, C., Porras, R. & Torn, M. The whole-soil carbon flux in response to
519   warming. *Science* **355**, 1420-1423 (2017).

520   27   Chorover, J., Kretzschmar, R., Garcia-Pichel, F. & Sparks, D. L. Soil biogeochemical
521   processes within the critical zone. *Elements* **3**, 321-326 (2007).

522   28   Stone, L. R., Goodrum, D. E., Jaafar, M. N. & Khan, A. H. Rooting front and water depletion

523   depths in grain sorghum and sunflower. *Agron. J.* **93**, 1105-1110 (2001).

524 29 Thorup-Kristensen, K. Effect of deep and shallow root systems on the dynamics of soil
525   inorganic N during 3-year crop rotations. *Plant Soil* **288**, 233-248 (2006).

526 30 Thorup-Kristensen, K. *et al.* Digging Deeper for Agricultural Resources, the Value of Deep
527   Rooting. *Trends Plant Sci.* **25**, 406-417 (2020).

528 31 Pierret, A. *et al.* Understanding deep roots and their functions in ecosystems: an advocacy for
529   more unconventional research. *Ann. Bot.* **118**, 621-635 (2016).

530 32 Nunes, M. R., Karlen, D. L., Denardin, J. E. & Cambardella, C. A. Corn root and soil health
531   indicator response to no-till production practices. *Agric., Ecosyst. Environ.* **285**, 106607
532   (2019).

533 33 Qin, R., Stamp, P. & Richner, W. Impact of tillage on root systems of winter wheat. *Agron. J.*
534   **96**, 1523-1530 (2004).

535 34 Fontaine, S. *et al.* Stability of organic carbon in deep soil layers controlled by fresh carbon
536   supply. *Nature* **450**, 277-280 (2007).

537 35 Tautges, N. E. *et al.* Deep soil inventories reveal that impacts of cover crops and compost on
538   soil carbon sequestration differ in surface and subsurface soils. *Glob. Chang. Biol.* **25**,
539   3753-3766 (2019).

540 36 Dal Ferro, N., Piccoli, I., Berti, A., Polese, R. & Morari, F. Organic carbon storage potential
541   in deep agricultural soil layers: Evidence from long-term experiments in northeast Italy.
542   *Agric., Ecosyst. Environ.* **300**, 106967 (2020).

543 37 Liu, X. *et al.* Soil degradation: a problem threatening the sustainable development of
544   agriculture in Northeast China. *Plant Soil Environ.* **56**, 87-97 (2010).

545 38 Liu, Z., Yang, X., Hubbard, K. G. & Lin, X. Maize potential yields and yield gaps in the
546   changing climate of northeast China. *Glob. Chang. Biol.* **18**, 3441-3454 (2012).

547 39 Canadell, J. *et al.* Maximum rooting depth of vegetation types at the global scale. *Oecologia*
548   **108**, 583-595, doi:10.1007/bf00329030 (1996).

549 40 Koike, I. & Hattori, A. Growth yield of a denitrifying bacterium, Pseudomonas denitrificans,
550   under aerobic and denitrifying conditions. *Microbiology* **88**, 1-10 (1975).

551 41 Miroshnichenko, M. L. *et al.* Caldithrix abyssi gen. nov., sp. nov., a nitrate-reducing,
552   thermophilic, anaerobic bacterium isolated from a Mid-Atlantic Ridge hydrothermal vent,
553   represents a novel bacterial lineage. *Int. J. Syst. Evol. Microbiol.* **53**, 323-329 (2003).

554 42 De Vita, P., Di Paolo, E., Fecondo, G., Di Fonzo, N. & Pisante, M. No-tillage and
555   conventional tillage effects on durum wheat yield, grain quality and soil moisture content in
556   southern Italy. *Soil Tillage Res.* **92**, 69-78 (2007).

557 43 Ramirez-Villanueva, D. A. *et al.* Bacterial community structure in maize residue amended soil
558   with contrasting management practices. *Appl. Soil Ecol.* **90**, 49-59 (2015).

559 44 Kremen, C. Managing ecosystem services: what do we need to know about their ecology?
560   *Ecol. Lett.* **8**, 468-479 (2005).

561 45 Wood, S. A. *et al.* Agricultural intensification and the functional capacity of soil microbes on
562   smallholder African farms. *J. Appl. Ecol.* **52**, 744-752 (2015).

563 46 Zhalnina, K. *et al.* Soil pH determines microbial diversity and composition in the park grass
564   experiment. *Microb. Ecol.* **69**, 395-406 (2015).

565 47 Lauber, C. L., Hamady, M., Knight, R. & Fierer, N. Pyrosequencing-based assessment of soil
566   pH as a predictor of soil bacterial community structure at the continental scale. *Appl. Environ.*

Microbiol. **75**, 5111-5120 (2009).

48   Rousk, J. *et al.* Soil bacterial and fungal communities across a pH gradient in an arable soil.
ISME J. **4**, 1340 (2010).

49   Lu, C. *et al.* Effects of N fertilization and maize straw on the dynamics of soil organic N and
amino acid N derived from fertilizer N as indicated by 15 N labeling. *Geoderma* **321**, 118-126
(2018).

50   Liu, X. *et al.* Linking microbial immobilization of fertilizer nitrogen to in situ turnover of soil
microbial residues in an agro-ecosystem. *Agric., Ecosyst. Environ.* **229**, 40-47 (2016).

51   Zhang, B., Wang, H., Yao, S. & Bi, L. Litter quantity confers soil functional resilience
through mediating soil biophysical habitat and microbial community structure on an eroded
bare land restored with mono Pinus massoniana. *Soil Biol. Biochem.* **57**, 556-567 (2013).

52   Kibblewhite, M., Ritz, K. & Swift, M. Soil health in agricultural systems. *Philos Trans R Soc
Lond B Biol Sci* **363**, 685-701 (2008).

53   Erbilgin, O., McDonald, K. L. & Kerfeld, C. A. Characterization of a planctomycetal
organelle: a novel bacterial microcompartment for the aerobic degradation of plant
saccharides. *Appl. Environ. Microbiol.* **80**, 2193-2205 (2014).

54   Herlemann, D. P. *et al.* Metagenomic de novo assembly of an aquatic representative of the
verrucomicrobial class Spartobacteria. *MBio* **4**, e00569-00512 (2013).

55   Halsall, D. M. & Goodchild, D. J. Nitrogen fixation associated with development and
localization of mixed populations of Cellulomonas sp. and Azospirillum brasilense grown on
cellulose or wheat straw. *Appl. Environ. Microbiol.* **51**, 849-854 (1986).

56   Pathma, J., Raman, G. & Sakthivel, N. Microbiome of Rhizospheric Soil and Vermicompost
and Their Applications in Soil Fertility, Pest and Pathogen Management for Sustainable
Agriculture. *189-210 (Springer, Singapore, 2019).*

57   Iino, T. *et al.* Ignavibacterium album gen. nov., sp. nov., a moderately thermophilic anaerobic
bacterium isolated from microbial mats at a terrestrial hot spring and proposal of
Ignavibacteria classis nov., for a novel lineage at the periphery of green sulfur bacteria. *Int. J.
Syst. Evol. Microbiol.* **60**, 1376-1382 (2010).

58   Terracciano, J. & Canale-Parola, E. Enhancement of chemotaxis in Spirochaeta aurantia
grown under conditions of nutrient limitation. *J. Bacteriol.* **159**, 173-178 (1984).

59   Thomas, F., Hehemann, J.-H., Rebuffet, E., Czjzek, M. & Michel, G. Environmental and gut
bacteroidetes: the food connection. *Front. Microbiol.* **2**, 93 (2011).

60   Archontoulis, S. & Licht, M. A. How Fast and Deep do Corn Roots Grow in Iowa? *Integrated
Crop Management News*, 2442 (2017).

61   Zhang, C., Ju, X., Powlson, D. S., Oenema, O. & Smith, P. Nitrogen surplus benchmarks for
controlling N pollution in the main cropping systems of China. *Environ. Sci. Technol.* **53**,
6678-6687 (2019).

62   Izewska, A. & Woloszyk, C. Yields of grain and straw, their content and ionic proportions of
macroelements in maize fertilized with ash from municipal sewage sludge combustion. *J
Elementology* **20**, 319-329 (2015).

63   Kennedy, S. Energy use in American agriculture. *Sustainable energy term paper* **5**, 1-26
(2000).

64   Zhang, W.-f. *et al.* New technologies reduce greenhouse gas emissions from nitrogenous
fertilizer in China. *Proc. Natl. Acad. Sci. U.S.A.* **110**, 8375-8380 (2013).

65 Dwyer, L. *et al.* Root mass distribution under conventional and conservation tillage. *Can. J.*
Soil Sci. **76**, 23-28 (1996).
66 Izumi, Y., Uchida, K. & Iijima, M. Crop production in successive wheat-soybean rotation
with no-tillage practice in relation to the root system development. *Plant production science* **7**,
329-336 (2004).
67 IUSS Working Group, W. World reference base for soil resources. *World Soil Resources*
Report **103**, FAO, Rome (2006).
68 Nelson, D. & Sommers, L. E. in *Methods of soil analysis,Part 2 (2nd).* (eds AL Page, RH
Miller, & DR Keeney)   539-579 (American Society of Agronomy and Soil Science Society
of America, 1982).
69 Jones, D. & Willett, V. Experimental evaluation of methods to quantify dissolved organic
nitrogen (DON) and dissolved organic carbon (DOC) in soil. *Soil Biol. Biochem.* **38**, 991-999
(2006).
70 Toosi, E. R., Castellano, M. J., Singer, J. W. & Mitchell, D. C. Differences in soluble organic
matter after 23 years of contrasting soil management. *Soil Sci. Soc. Am. J.* **76**, 628-637
(2012).
71 Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114-2120 (2014).
72 Derakhshani, H., Tun, H. M. & Khafipour, E. An extended single□index multiplexed 16S
rRNA sequencing for microbial community analysis on MiSeq illumina platforms. *J. Basic*
Microbiol. **56**, 321-326 (2016).
73 Mysara, M., Leys, N., Raes, J. & Monsieurs, P. IPED: a highly efficient denoising tool for
Illumina MiSeq Paired-end 16S rRNA gene amplicon sequencing data. *BMC Bioinformatics*
17, 192 (2016).
74 Wang, Q., Garrity, G. M., Tiedje, J. M. & Cole, J. R. Naive Bayesian classifier for rapid
assignment of rRNA sequences into the new bacterial taxonomy. *Appl. Environ. Microbiol.* **73**,
5261-5267 (2007).
75 Dakora, F. D. & Phillips, D. A. in *Food Security in Nutrient-Stressed Environments:*
Exploiting Plants' Genetic Capabilities   201-213 (Springer, 2002).
76 Cáceres, M. D. & Legendre, P. Associations between species and groups of sites: indices and
statistical inference. *Ecology* **90**, 3566-3574 (2009).
77 Aßhauer, K. P., Wemheuer, B., Daniel, R. & Meinicke, P. Tax4Fun: predicting functional
profiles from metagenomic 16S rRNA data. *Bioinformatics* **31**, 2882-2884 (2015).
78 Parks, D. H., Tyson, G. W., Hugenholtz, P. & Beiko, R. G. STAMP: statistical analysis of
taxonomic and functional profiles. *Bioinformatics* **30**, 3123-3124 (2014).

Figure legends

Figure 1. Soil properties (mean± SE, n = 3) along soil depth under different practices. **a**, SOC

= soil organic carbon; **b**, SEOC = salt-extractable organic carbon; **c**, SEOC/SOC = ratio of

SEOC to SOC; **d**, soil pH; **e**, SM = soil moisture; **f**, TN = total nitrogen content; **g**, C/N =

ratio of SOC to TN; **h**, NH_4^+-N = ammonium nitrogen; **i**, NO_3^--N = nitrate nitrogen; **j**,

Inorganic N = NH_4^+-N + NO_3^--N. Error bars indicate standard errors (n = 3). Different letters

indicate significant differences (P < 0.05) among disturbing practices.

Figure 2. Corn yield (line+ symbol) and annual rainfall during growing and dormant seasons

(bar) under different disturbance practices during 2013-2016. Error bars indicate standard

errors (n = 3 or 4), different letters indicate significant differences at P < 0.05. Different corn

seeds were used among years.

Figure 3. Microbial richness (Chao1), observed number of species (Observed_species) and

diversity (Shannon_Index) in the CT, NTNS and NTSM plots. Error bars indicate standard

errors (n = 3). Different letters indicate significant differences (P < 0.05) among disturbing

practices.

Figure 4. Indicator genera significantly (q < 0.1) associated with tillage practices. The size of

each circle represents the indicator value of a specific genus in the different soil depths. The

color indicates the relative abundance of each indicator genus. Taxonomic information,

indicator values, P-values, and q-values of all indicator genera are given in Table S2. Zero

indicator genera were identified in CT treatment.

Figure 5. Non-metric multidimensional scaling (MDS) ordination of soil microbial

community structures based on Bray-Curtis distances among soil depths at different

agricultural disturbance practices. Permutational multivariate analysis of variance

(PERMANOVA) revealed that the overall microbial community structures among soil depth

were significantly different at each disturbance practice. Circles, triangles and squares

represent CT, NTNS and NTSM, respectively.

Figure 6. Redundancy analysis (RDA) of the soil microbial community originating from

microbial phyla constrained by soil properties under different agricultural practices. Only soil

variables that significantly explained variability in microbial community structure in the

forward selection procedure were selected to the ordination (arrows). TN, total nitrogen

content; C/N, a ratio of carbon to nitrogen content; NH_4^+-N, ammonium nitrogen; NO_3^--N,

nitrate nitrogen；SM, soil moisture. Circles, triangles and squares represent CT, NTNS and

NTSM, respectively.

Figure 1

685   **Figure 2**

689   **Figure 3**

Figure 4

697 **Figure 5**

700 **Figure 6**

Supplementary Information

Low-disturbance Farming Regenerates Healthy Deep Soil towards Sustainable Agriculture

Fangbo Deng[1, 2#], Hongjun Wang[3#], Hongtu Xie[1#], Xuelian Bao[1], Hongbo He[1],

Xudong Zhang[1], Chao Liang[1*]

[1] Key Laboratory of Forest Ecology and Management, Institute of Applied Ecology, Chinese

Academy of Sciences, Shenyang 110016, China

[2] University of Chinese Academy of Sciences, Beijing 100049, China

[3] Duke University Wetland Center, Nicholas School of the Environment, Duke University, Durham,

NC 27708, USA

[#] These authors contributed equally to this work.

[*] Corresponding author: Chao Liang (Email: cliang823@gmail.com)

Figure S1. The relative abundance of bacterial community composition at the phylum level. **a**, Only the bacterial phyla with the relative abundance > 0.1% across all soil samples were shown; **b**, "Others" in the (**a**) panel represents the sum of bacterial phyla that individual relative abundance < 0.1% across all soil samples were shown. Abbreviations: CT (conventional tillage), NTNS (no-tillage without stover mulching) and NTSM (no-tillage with stover mulching).

Figure S2. Extended error bar plots showing significant differences of 16S rRNA gene-predicted functional profiles obtained with Tax4Fun. **a**, difference between mean proportions of conventional practice and low disturbance practices; **b**, differences between mean proportions of NTNS (no-tillage without stover mulching) and NTSM (no-tillage with 100% stover mulching).

Figure S3. The denitrification and nitrification genes that influenced by different tillage practices. Genes in red rectangles means higher abundance in low disturbing practices; Genes in blue rectangles means higher abundance in conventional tillage practice.

Figure S4. Redundancy analysis (RDA) of soil microbial community originating from microbial phyla constrained by soil properties among samples. Only soil variables that significantly explained variability in microbial community structure in the forward selection procedure were selected to the ordination (arrows). Abbreviations: SOC, soil organic carbon; NH_4^+, ammonium nitrogen; SM, soil moisture.

Figure S5. Root depths of different crops under different tillage practices. **a**, corn; **b**, wheat; **c**, soybean; **d**, sunflower; **e**, mungbean; **f**, the average root depth of crops under one tillage practice minus the average root depth under another tillage practice. n, numbers of observations; Ref, corresponding references number; Totally 87 observations were collected from following references: 1, Bunna et al., 2011; 2, Kemper et al.,2011; 3, Merrill et al.,2002; 4, Turman et al.,1995; 5, Dwyer et al.,1996; 6, Dwyer et al.,1988. See details in Table S5.

Table S1. Soil properties at different soil depths among disturbance practices.

Soil Properties	Tillage	Soil depth (cm)									
		0-10	10-20	20-40	40-60	60-90	90-120	120-150	150-200	200-250	250-300
SOC	CT	12.53±0.53a	11.18±0.40a	9.00±1.83a	7.72±3.16a	6.12±1.49a	3.06±1.10a	1.98±0.61a	1.66±0.16a	2.08±0.49a	2.23±0.10a
	NTNS	13.56±0.24a	11.12±0.47a	10.76±0.99a	8.03±2.80a	4.56±0.45a	3.01±1.27a	2.19±0.71a	2.14±0.28a	2.31±0.57a	2.82±0.40a
	NTSM	14.19±1.46a	12.72±2.88a	9.48±0.28a	6.55±0.28a	4.73±0.91a	2.80±0.17a	1.83±0.51a	1.94±0.86a	1.89±0.82a	2.31±0.78a
		A	B	C	D	E	F	F	F	F	F
SEOC	CT	91.96±3.52b	80.04±13.92a	70.49±7.75a	71.80±29.67a	105.57±53.39a	76.38±35.40a	93.72±11.15b	66.86±5.74a	59.60±10.62a	83.62±25.86a
	NTNS	127.96±33.79ab	103.30±50.78a	82.94±18.61a	60.26±37.99a	105.71±76.22a	111.66±60.32a	72.59±11.21b	85.71±64.89a	83.35±27.37a	62.26±19.63a
	NTSM	188.67±68.99a	160.78±56.73a	92.00±3.79a	86.77±40.43a	102.67±36.07a	149.09±58.04a	196.49±53.90a	97.68±31.36a	98.79±24.44a	73.03±14.14a
		A	ABC	BC	C	ABC	ABC	AB	BC	BC	C
SEOC /SOC	CT	0.0073±0.0003b	0.0072±0.0012a	0.0081±0.0022a	0.0098±0.0048a	0.0167±0.0054a	0.026±0.012a	0.0496±0.0129b	0.0406±0.006a	0.0294±0.0066a	0.0379±0.0133a
	NTNS	0.0095±0.0027ab	0.0092±0.0041a	0.0077±0.0011a	0.0072±0.0025a	0.0223±0.0141a	0.0416±0.0318a	0.036±0.0139b	0.0405±0.0311a	0.0392±0.0191a	0.0211±0.0088a
	NTSM	0.013±0.0037a	0.0126±0.0039a	0.0097±0.0002a	0.0134±0.0068a	0.0232±0.013a	0.0532±0.0209a	0.1081±0.0081a	0.0636±0.0465a	0.0587±0.0234a	0.0333±0.0076a
		A	A	A	A	AB	CD	E	D	CD	BC
pH	CT	7.65±0.10a	7.39±0.53a	7.55±0.70a	7.86±0.19a	7.81±0.22a	8.07±0.40a	8.41±0.25a	8.26±0.19a	8.06±0.22a	8.14±0.15a
	NTNS	6.99±0.06a	7.57±0.08a	7.67±0.13a	7.82±0.04a	7.82±0.05a	8.06±0.34a	8.11±0.42ab	8.10±0.49a	7.86±0.55a	7.89±0.37a
	NTSM	7.30±0.77a	7.23±0.71a	7.77±0.18a	7.85±0.06a	7.79±0.03a	7.41±0.04b	7.68±0.22b	7.75±0.36a	7.69±0.22a	7.75±0.24a
		A	AB	BC	CD	CD	CD	D	D	CD	CD
SM	CT	18.14±0.75b	20.32±3.08a	25.14±0.72a	24.03±0.52a	24.40±1.44a	22.38±0.77a	21.51±1.33a	20.60±1.82a	22.07±1.48a	23.57±0.53a
	NTNS	18.50±1.63ab	21.93±0.28a	26.58±3.00a	24.34±1.82a	25.02±1.56a	19.53±2.18a	21.71±0.39a	21.02±1.46a	22.17±0.80a	23.17±0.64a
	NTSM	20.58±0.64a	22.53±1.62a	27.90±0.27a	25.64±0.72a	23.52±1.06a	22.17±1.12a	21.50±0.78a	20.26±0.41a	22.87±0.56a	24.53±0.79a
		A	BC	E	D	D	BC	BC	B	C	D
TN	CT	1.31±0.12a	1.23±0.03a	0.99±0.20a	0.88±0.31a	0.79±0.19a	0.49±0.09a	0.47±0.05a	0.34±0.06a	0.43±0.08a	0.42±0.04a
	NTNS	1.39±0.07a	1.20±0.05a	1.21±0.22a	0.93±0.25a	0.69±0.16a	0.46±0.14a	0.42±0.08a	0.40±0.05a	0.36±0.07a	0.38±0.06a
	NTSM	1.44±0.08a	1.36±0.17a	1.04±0.09a	0.74±0.03a	0.51±0.07a	0.47±0.07a	0.39±0.09a	0.31±0.03a	0.36±0.05a	0.41±0.06a
		A	B	C	**D**	E	F	F	F	F	F

	CT	9.65±1.10a	9.09±0.14a	9.08±0.44a	8.70±0.50a	7.76±0.64a	6.16±1.23a	4.33±1.66a	4.94±1.11a	4.96±1.40a	5.34±0.33b
C/N	NTNS	9.77±0.43a	9.27±0.32a	8.99±0.86a	8.53±0.80a	6.80±1.59a	6.39±0.77a	5.14±0.71a	5.36±0.30a	6.33±0.56a	7.54±1.43a
	NTSM	9.82±0.64a	9.28±1.06a	9.14±0.48a	8.87±0.60a	9.27±1.24a	6.09±0.99a	4.71±0.65a	6.05±2.16a	5.46±2.76a	5.46±1.12ab
		A	A	A	AB	B	C	C	CD	CD	D
	CT	11.69±1.10a	7.47±1.81a	5.72±0.55b	5.11±1.33b	5.84±0.76b	6.05±1.32ab	5.77±0.80b	5.68±0.74b	6.44±0.26b	6.59±0.89a
NH_4^+-N	NTNS	8.99±1.08ab	9.18±1.84a	8.04±1.67a	8.52±1.97a	6.97±0.52a	7.39±0.64a	9.53±2.44a	8.44±0.96a	10.02±2.14a	6.94±0.99a
	NTSM	6.94±1.78b	6.13±1.59a	5.57±0.97b	5.34±0.59b	4.99±0.08b	5.3±0.68b	5.02±0.41b	5.64±0.08b	5.42±0.76b	5.93±0.63a
		A	B	BCD	BCD	D	CD	BCD	BCD	BC	BCD
	CT	12.39±3.33a	10.56±4.17a	4.80±0.93b	3.61±0.86a	6.16±4.51a	4.19±1.69b	8.30±5.65b	17.98±5.06a	20.43±8.04a	23.32±6.79a
NO_3^--N	NTNS	16.04±1.61a	10.17±1.65a	9.93±2.97a	8.03±4.36a	9.43±7.15a	13.55±4.16a	23.45±10.34a	11.63±2.74a	11.15±3.46a	12.35±4.71a
	NTSM	12.67±3.35a	10.15±5.26a	5.74±1.34b	5.71±2.04a	8.57±6.69a	16.68±4.89a	21.90±1.24ab	12.92±1.50a	17.93±7.36a	14.91±5.30a
		AB	BCD	CD	D	CD	BC	A	AB	A	A
	CT	24.08±2.24a	18.02±4.51a	10.53±1.09b	8.71±2.16b	12.01±4.05a	10.24±2.84b	14.07±5.26b	23.67±4.77a	26.87±8.30a	29.91±5.91a
Inorganic N	NTNS	25.03±2.40a	19.35±3.07a	17.97±1.37a	16.55±4.47a	16.40±6.79a	20.95±4.16a	32.97±10.59a	20.08±3.65a	21.17±4.06a	19.29±4.60a
	NTSM	19.61±4.23a	16.28±6.79a	11.31±0.65b	11.05±1.50ab	13.56±6.69a	21.99±5.33a	26.91±1.65ab	18.56±1.57a	23.34±7.40a	20.84±5.28a
		A	BC	CD	D	CD	BC	A	AB	A	A

49 Different letters indicate significant differences at $P<0.05$. Abbreviations: SOC = soil organic carbon; SEOC = salt-extractable organic carbon; SEOC/SOC = ratio of

50 SEOC to SOC; SM = soil moisture; TN = total nitrogen content; C/N = ratio of SOC to TN; NH_4^+-N = ammonium nitrogen; NO_3^--N = nitrate nitrogen ;Inorganic N =

51 NH_4^+-N + NO_3^--N.

53 Table S2. Taxonomic information, indicator values, P-values and q-values of indicator genera associated with tillage practices.

Land use	Phylum	genera	Soil Depth (cm)										p.value	qvalues
			0-10	10-20	20-40	40-60	60-90	90-120	120-150	150-200	200-250	250-300		
CT	0 indicator genera													
NTNS	Actinobacteria	Aciditerrimonas	0.712	0.067	0.283	0.067	0.189	0.267	0.115	0.249	0.067	0.231	0.0018	0.08426
NTNS	Actinobacteria	Angustibacter	0.816	0.295	0.295	0.098	0.070	0.070	0.098	0.070	0.070	0.120	0.0018	0.08426
NTNS	Actinobacteria	Aeromicrobium	0.572	0.341	0.274	0.210	0.280	0.348	0.243	0.194	0.262	0.206	0.0018	0.08426
NTNS	Actinobacteria	Amycolatopsis	0.803	0.000	0.147	0.104	0.147	0.000	0.104	0.000	0.254	0.147	0.0018	0.08426
NTNS	Chloroflexi	Nitrolancea	0.879	0.000	0.174	0.246	0.000	0.000	0.000	0.000	0.000	0.123	0.0018	0.08426
NTNS	Gemmatimonadetes	Gemmatimonas	0.712	0.315	0.363	0.178	0.190	0.231	0.190	0.152	0.253	0.135	0.0013	0.08426
NTNS	Proteobacteria	Caulobacter	0.673	0.276	0.434	0.265	0.262	0.200	0.085	0.147	0.135	0.110	0.0034	0.0975
NTNS	Proteobacteria	Phenylobacterium	0.643	0.359	0.342	0.189	0.184	0.236	0.305	0.285	0.154	0.130	0.0018	0.08426
NTNS	Proteobacteria	Rhizomicrobium	0.747	0.280	0.294	0.145	0.227	0.221	0.234	0.183	0.173	0.150	0.0018	0.08426
NTNS	Proteobacteria	Pseudolabrys	0.717	0.357	0.342	0.165	0.165	0.219	0.193	0.161	0.177	0.128	0.0018	0.08426
NTNS	Proteobacteria	Rhodovastum	0.890	0.000	0.118	0.118	0.118	0.167	0.000	0.000	0.000	0.000	0.0018	0.08426
NTNS	Proteobacteria	Castellaniella	0.903	0.094	0.000	0.094	0.000	0.094	0.094	0.094	0.187	0.000	0.0018	0.08426
NTNS	Proteobacteria	Dokdonella	0.725	0.209	0.289	0.162	0.223	0.132	0.153	0.256	0.248	0.121	0.0018	0.08426
NTNS	Proteobacteria	Mizugakiibacter	0.857	0.226	0.210	0.072	0.127	0.160	0.139	0.098	0.208	0.116	0.0018	0.08426
NTNS	Planctomycetes	AKYG587	0.306	0.709	0.301	0.200	0.193	0.237	0.178	0.285	0.125	0.223	0.0027	0.08426
NTNS	Aenigmarchaeota	Candidatus_Aenigmarchaeum	0.067	0.097	0.082	0.074	0.047	0.030	0.129	0.308	0.432	0.794	0.0026	0.08426
NTSM	Actinobacteria	Blastococcus	0.660	0.541	0.208	0.131	0.172	0.147	0.125	0.249	0.189	0.216	0.0115	0.09926
NTSM	Actinobacteria	Modestobacter	0.800	0.115	0.000	0.163	0.115	0.115	0.000	0.115	0.000	0.200	0.003	0.09403
NTSM	Actinobacteria	Angustibacter	0.699	0.176	0.000	0.125	0.125	0.125	0.088	0.249	0.088	0.431	0.0093	0.09454
NTSM	Actinobacteria	Cellulomonas	0.778	0.265	0.094	0.162	0.094	0.000	0.094	0.094	0.094	0.162	0.003	0.09403
NTSM	Actinobacteria	Aeromicrobium	0.546	0.381	0.288	0.275	0.263	0.376	0.249	0.231	0.102	0.249	0.003	0.09403
NTSM	Actinobacteria	Nocardioides	0.664	0.427	0.197	0.245	0.214	0.214	0.167	0.224	0.183	0.274	0.0063	0.09403

NTSM	Actinobacteria	Pseudonocardia	0.630	0.542	0.270	0.183	0.146	0.258	0.149	0.168	0.127	0.101	0.0063	0.09403
NTSM	Bacteroidetes	Ohtaekwangia	0.583	0.441	0.415	0.317	0.172	0.241	0.146	0.147	0.113	0.195	0.0086	0.09403
NTSM	Bacteroidetes	Flavisolibacter	0.717	0.481	0.226	0.147	0.150	0.129	0.133	0.160	0.180	0.260	0.003	0.09403
NTSM	Bacteroidetes	Flavitalea	0.618	0.552	0.202	0.295	0.128	0.140	0.128	0.128	0.214	0.268	0.0079	0.09403
NTSM	Bacteroidetes	Parafilimonas	0.687	0.467	0.294	0.175	0.029	0.101	0.041	0.133	0.101	0.377	0.0072	0.09403
NTSM	Bacteroidetes	Parasegetibacter	0.742	0.540	0.089	0.125	0.000	0.072	0.072	0.051	0.162	0.145	0.003	0.09403
NTSM	Bacteroidetes	Segetibacter	0.749	0.432	0.237	0.112	0.079	0.056	0.125	0.079	0.193	0.056	0.0074	0.09403
NTSM	Gemmatimonadetes	Gemmatimonas	0.730	0.367	0.238	0.184	0.202	0.172	0.162	0.167	0.182	0.290	0.005	0.09403
NTSM	Proteobacteria	Phenylobacterium	0.573	0.357	0.286	0.295	0.190	0.290	0.104	0.195	0.195	0.380	0.0083	0.09403
NTSM	Proteobacteria	Bradyrhizobium	0.590	0.411	0.326	0.306	0.241	0.235	0.154	0.212	0.198	0.249	0.003	0.09403
NTSM	Proteobacteria	Rhodoplanes	0.566	0.440	0.368	0.346	0.252	0.218	0.133	0.126	0.168	0.241	0.0054	0.09403
NTSM	Proteobacteria	Microvirga	0.672	0.536	0.199	0.144	0.102	0.158	0.102	0.188	0.171	0.299	0.0109	0.0984
NTSM	Proteobacteria	Labrys	0.673	0.423	0.257	0.275	0.056	0.194	0.137	0.112	0.217	0.291	0.0063	0.09403
NTSM	Proteobacteria	Pseudolabrys	0.706	0.391	0.219	0.136	0.185	0.157	0.149	0.160	0.175	0.380	0.0052	0.09403
NTSM	Proteobacteria	Variibacter	0.571	0.486	0.299	0.277	0.258	0.193	0.159	0.206	0.175	0.264	0.0059	0.09403
NTSM	Proteobacteria	Craurococcus	0.882	0.272	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.009	0.09454
NTSM	Proteobacteria	Dongia	0.740	0.355	0.237	0.084	0.084	0.068	0.076	0.244	0.153	0.360	0.0096	0.09454
NTSM	Proteobacteria	Elstera	0.894	0.183	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.183	0.0072	0.09403
NTSM	Proteobacteria	Reyranella	0.551	0.373	0.364	0.355	0.234	0.248	0.154	0.163	0.151	0.329	0.0073	0.09403
NTSM	Proteobacteria	Acidovorax	0.775	0.490	0.000	0.115	0.163	0.000	0.000	0.115	0.000	0.000	0.0063	0.09403
NTSM	Proteobacteria	Sorangium	0.491	0.417	0.341	0.209	0.287	0.231	0.231	0.333	0.251	0.251	0.0059	0.09403
NTSM	Acidobacteria	Blastocatella	0.545	0.623	0.295	0.242	0.151	0.163	0.108	0.156	0.140	0.252	0.0114	0.09926
NTSM	Actinobacteria	Luedemannella	0.372	0.608	0.386	0.399	0.175	0.101	0.101	0.238	0.189	0.101	0.0031	0.09403
NTSM	Actinobacteria	Microlunatus	0.433	0.559	0.459	0.341	0.241	0.135	0.125	0.163	0.130	0.193	0.0031	0.09403
NTSM	Planctomycetes	AKYG587	0.431	0.680	0.330	0.265	0.163	0.182	0.137	0.213	0.157	0.153	0.0031	0.09403
NTSM	Planctomycetes	SM1A02	0.449	0.568	0.318	0.406	0.173	0.223	0.189	0.183	0.169	0.179	0.0099	0.09458
NTSM	Planctomycetes	planctomycete_LX80	0.252	0.690	0.126	0.126	0.000	0.126	0.000	0.252	0.309	0.126	0.0054	0.09403

NTSM	Proteobacteria	Azospirillum	0.591	0.731	0.000	0.088	0.000	0.000	0.000	0.125	0.000	0.176	0.0079	0.09403
NTSM	Proteobacteria	Nannocystis	0.376	0.492	0.242	0.329	0.303	0.316	0.105	0.197	0.365	0.183	0.0054	0.09403
NTSM	Proteobacteria	Phaselicystis	0.340	0.555	0.248	0.328	0.203	0.304	0.175	0.263	0.226	0.248	0.0049	0.09403
NTSM	Deferribacteres	Caldithrix	0.079	0.046	0.125	0.202	0.312	0.257	0.444	0.646	0.336	0.194	0.0047	0.09403
NTSM	Firmicutes	Ruminiclostridium_1	0.302	0.220	0.302	0.191	0.078	0.246	0.220	0.505	0.246	0.405	0.0068	0.09403
NTSM	Proteobacteria	Anaeromyxobacter	0.185	0.123	0.251	0.254	0.293	0.352	0.333	0.540	0.368	0.275	0.0098	0.09454
NTSM	Actinobacteria	Iamia	0.282	0.306	0.609	0.400	0.317	0.268	0.189	0.207	0.118	0.177	0.0026	0.09403
NTSM	Actinobacteria	Frankia	0.186	0.237	0.747	0.403	0.271	0.197	0.093	0.107	0.161	0.140	0.0026	0.09403
NTSM	Bacteroidetes	Owenweeksia	0.192	0.385	0.745	0.192	0.000	0.000	0.000	0.000	0.000	0.000	0.0102	0.09559
NTSM	Chloroflexi	Anaerolinea	0.000	0.333	0.745	0.272	0.000	0.000	0.000	0.000	0.136	0.192	0.0082	0.09403
NTSM	Firmicutes	Clostridium_ sensu_stricto_13	0.108	0.324	0.675	0.296	0.216	0.153	0.350	0.076	0.076	0.076	0.0071	0.09403
NTSM	Verrucomicrobia	Candidatus_ Xiphinematobacter	0.311	0.423	0.472	0.334	0.251	0.353	0.292	0.292	0.108	0.117	0.0094	0.09454
NTSM	Chlorobi	Ignavibacterium	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	1.000	0.0035	0.09403
NTSM	Firmicutes	Thermincola	0.055	0.257	0.300	0.251	0.077	0.414	0.134	0.300	0.232	0.593	0.0068	0.09403
NTSM	Proteobacteria	Pseudomonas	0.312	0.101	0.164	0.143	0.343	0.235	0.344	0.314	0.346	0.579	0.01	0.09462
NTSM	Spirochaetae	Spirochaeta_2	0.058	0.058	0.000	0.000	0.000	0.058	0.058	0.404	0.535	0.714	0.0104	0.09654
NTSM	Actinobacteria	Haloactinopolyspora	0.203	0.224	0.498	0.601	0.423	0.204	0.162	0.163	0.115	0.116	0.0048	0.09403
NTSM	Proteobacteria	Piscinibacter	0.298	0.237	0.499	0.586	0.351	0.240	0.117	0.113	0.148	0.161	0.0114	0.09926

55 Table. S3 Spearman correlations between indicator genera and soil properties.

Land use	Phylum	genera	Depth	SOC	SEOC	SEOC/SOC	pH	SM	TN	C/N	NO$_3^-$-N	NH$_4^+$-N
CT	NA											
NTNS	Actinobacteria	Aciditerrimonas	-0.222	0.192	0.314	0.067	-0.113	-0.263	0.148	0.209	0.079	0.022
NTNS	Actinobacteria	Angustibacter	-0.554**	0.463**	0.203	-0.515**	-0.516**	-0.205	0.475**	0.461*	0.099	0.137
NTNS	Actinobacteria	Aeromicrobium	-0.536**	0.271	0.311	-0.135	-0.485**	-0.206	0.29	0.246	0.131	0.261
NTNS	Actinobacteria	Amycolatopsis	-0.241	0.242	0.109	-0.282	-0.422*	-0.171	0.226	0.29	0.297	0.12
NTNS	Chloroflexi	Nitrolancea	-0.458*	0.513**	0.04	-0.397*	-0.43*	-0.088	0.439*	0.543**	0.213	0.141
NTNS	Gemmatimonadetes	Gemmatimonas	-0.597**	0.402*	0.376*	-0.165	-0.38*	-0.317	0.39*	0.357	0.128	0.31
NTNS	Proteobacteria	Caulobacter	-0.696**	0.671**	0.231	-0.508**	-0.453*	-0.087	0.659**	0.648**	0.14	-0.018
NTNS	Proteobacteria	Phenylobacterium	-0.675**	0.4*	0.103	-0.341	-0.456*	-0.284	0.521**	0.357	0.303	0.34
NTNS	Proteobacteria	Rhizomicrobium	-0.51**	0.277	0.145	-0.25	-0.294	-0.29	0.288	0.333	0.357	0.251
NTNS	Proteobacteria	Pseudolabrys	-0.594**	0.39*	0.347	-0.287	-0.471**	-0.264	0.402*	0.409*	0.222	0.301
NTNS	Proteobacteria	Rhodovastum	-0.509**	0.456*	-0.042	-0.384*	-0.423*	-0.197	0.466**	0.384*	0.374*	-0.059
NTNS	Proteobacteria	Castellaniella	-0.249	0.103	0.209	0.011	-0.307	-0.485**	0.098	0.127	-0.046	0.387*
NTNS	Proteobacteria	Dokdonella	-0.437*	0.337	0.24	-0.206	-0.552**	-0.06	0.367*	0.304	-0.017	0.283
NTNS	Proteobacteria	Mizugakiibacter	-0.436*	0.248	0.313	-0.203	-0.465**	-0.192	0.212	0.352	0.142	0.249
NTNS	Planctomycetes	AKYG587	-0.444*	0.354	-0.128	-0.422*	-0.386*	0.099	0.344	0.378*	0.069	0.069
NTNS	Aenigmarchaeota	Candidatus_Aenigmarchaeum	0.57**	-0.388*	-0.24	0.068	-0.062	0.127	-0.483**	-0.173	-0.091	0.068
NTSM	Actinobacteria	Blastococcus	-0.383*	0.425*	0.148	-0.343	-0.144	-0.194	0.433*	0.319	-0.135	0.221
NTSM	Actinobacteria	Modestobacter	-0.343	0.334	0.193	-0.253	-0.254	-0.204	0.385*	0.251	-0.134	0.148
NTSM	Actinobacteria	Angustibacter	-0.039	0.17	0.245	0.015	0.059	-0.269	0.125	0.018	-0.045	0.418*
NTSM	Actinobacteria	Cellulomonas	-0.417*	0.477**	0.022	-0.463**	-0.014	-0.214	0.499**	0.361	-0.013	0.299
NTSM	Actinobacteria	Aeromicrobium	-0.67**	0.634**	0.459*	-0.391*	-0.447*	-0.197	0.615**	0.473**	-0.016	0.116
NTSM	Actinobacteria	Nocardioides	-0.466**	0.523**	0.171	-0.42*	-0.239	-0.111	0.539**	0.291	-0.051	0.373*
NTSM	Actinobacteria	Pseudonocardia	-0.7**	0.61**	0.272	-0.496**	-0.125	-0.18	0.631**	0.486**	-0.254	-0.001

NTSM	Bacteroidetes	Ohtaekwangia	**-0.821****	**0.786****	0.125	**-0.726****	-0.183	0.101	**0.812****	**0.601****	**-0.43***	0.03
NTSM	Bacteroidetes	Flavisolibacter	**-0.453***	**0.524****	0.18	**-0.428***	-0.276	-0.08	**0.578****	**0.416***	-0.01	0.213
NTSM	Bacteroidetes	Flavitalea	**-0.469****	**0.563****	0.023	**-0.558****	-0.163	0.02	**0.653****	**0.403***	-0.173	0.247
NTSM	Bacteroidetes	Parafilimonas	**-0.409***	**0.519****	0.122	**-0.451***	-0.218	-0.046	**0.501****	0.335	-0.146	**0.464****
NTSM	Bacteroidetes	Parasegetibacter	**-0.441***	**0.449***	0.177	**-0.363***	-0.269	-0.234	**0.478****	0.286	-0.028	0.112
NTSM	Bacteroidetes	Segetibacter	**-0.591****	**0.47****	0.324	-0.32	-0.175	-0.092	**0.535****	**0.404***	-0.216	0.288
NTSM	Gemmatimonadetes	Gemmatimonas	**-0.467****	**0.619****	-0.08	**-0.605****	-0.262	0.027	**0.658****	**0.452***	-0.032	0.291
NTSM	Proteobacteria	Phenylobacterium	**-0.427***	**0.589****	-0.163	**-0.631****	-0.246	0.158	**0.598****	**0.392***	-0.129	0.346
NTSM	Proteobacteria	Bradyrhizobium	**-0.785****	**0.809****	0.139	**-0.719****	-0.175	0.104	**0.819****	**0.669****	-0.446*	0.153
NTSM	Proteobacteria	Rhodoplanes	**-0.778****	**0.825****	0.069	**-0.774****	-0.234	0.181	**0.904****	**0.66****	**-0.427***	0.109
NTSM	Proteobacteria	Microvirga	**-0.403***	**0.493****	0.186	**-0.378***	-0.193	-0.082	**0.576****	0.263	-0.114	0.252
NTSM	Proteobacteria	Labrys	**-0.47****	**0.5****	0.182	**-0.427***	-0.254	0.022	**0.549****	0.277	-0.195	0.269
NTSM	Proteobacteria	Pseudolabrys	**-0.403***	**0.476****	0.077	**-0.418***	-0.249	-0.105	**0.578****	0.287	-0.018	0.214
NTSM	Proteobacteria	Variibacter	**-0.743****	**0.801****	0.062	**-0.737****	-0.147	0.138	**0.793****	**0.681****	**-0.439***	0.115
NTSM	Proteobacteria	Craurococcus	**-0.58****	**0.531****	0.288	**-0.401***	-0.148	-0.281	**0.53****	**0.362***	-0.118	0.308
NTSM	Proteobacteria	Dongia	-0.305	**0.365***	0.031	-0.345	-0.239	-0.265	0.35	0.28	0.006	0.335
NTSM	Proteobacteria	Elstera	**-0.406***	**0.493****	0.301	-0.317	**-0.428***	**-0.384***	**0.507****	**0.407***	0.166	**0.416***
NTSM	Proteobacteria	Reyranella	**-0.694****	**0.775****	-0.023	**-0.767****	-0.254	0.178	**0.813****	**0.614****	-0.317	0.111
NTSM	Proteobacteria	Acidovorax	**-0.639****	**0.62****	0.39*	**-0.388***	-0.311	-0.352	**0.571****	**0.521****	-0.222	0.258
NTSM	Proteobacteria	Sorangium	**-0.521****	**0.465****	0.164	**-0.372***	-0.229	-0.244	**0.499****	0.305	-0.182	0.351
NTSM	Acidobacteria	Blastocatella	**-0.67****	**0.749****	0.053	**-0.711****	-0.145	0.152	**0.793****	**0.561****	-0.41*	0.218
NTSM	Actinobacteria	Luedemannella	**-0.647****	**0.543****	0.011	**-0.559****	0.128	0.211	**0.602****	**0.431***	**-0.535****	0.028
NTSM	Actinobacteria	Microlunatus	**-0.8****	**0.838****	-0.004	**-0.828****	0.005	0.254	**0.835****	**0.728****	**-0.523****	0.057
NTSM	Planctomycetes	AKYG587	**-0.642****	**0.609****	0.088	**-0.583****	-0.033	0.111	**0.581****	**0.472****	**-0.495****	-0.014
NTSM	Planctomycetes	SM1A02	**-0.796****	**0.783****	0.165	**-0.711****	-0.151	0.176	**0.752****	**0.62****	**-0.391***	0.175
NTSM	Planctomycetes	planctomycete_LX80	-0.254	0.223	0.194	-0.087	-0.195	-0.12	0.255	0.133	-0.067	0.312
NTSM	Proteobacteria	Azospirillum	**-0.464****	**0.481****	0.236	-0.341	-0.124	-0.245	**0.488****	0.288	-0.154	0.28

NTSM	Proteobacteria	Nannocystis	**-0.466****	**0.5****	0.081	**-0.436***	-0.301	-0.171	**0.494****	**0.388***	-0.034	-0.017
NTSM	Proteobacteria	Phaselicystis	**-0.368***	**0.385***	-0.032	**-0.399***	-0.12	-0.087	0.349	0.319	0.04	0.086
NTSM	Deferribacteres	Caldithrix	**0.642****	**-0.706****	-0.057	**0.705****	0.185	-0.36	**-0.786****	**-0.525****	**0.371***	-0.112
NTSM	Firmicutes	Ruminiclostridium_1	0.315	-0.257	-0.123	0.222	0.04	-0.36	-0.305	-0.268	0.054	-0.065
NTSM	Proteobacteria	Anaeromyxobacter	0.634**	**-0.702****	-0.207	**0.604****	-0.006	-0.299	**-0.736****	**-0.541****	**0.496****	-0.15
NTSM	Actinobacteria	Iamia	**-0.656****	**0.577****	-0.124	**-0.672****	0.183	**0.527****	**0.591****	**0.503****	**-0.665****	-0.211
NTSM	Actinobacteria	Frankia	**-0.59****	**0.605****	-0.179	**-0.707****	-0.058	**0.487****	**0.589****	**0.619****	**-0.448***	-0.152
NTSM	Bacteroidetes	Owenweeksia	**-0.545****	**0.574****	-0.024	**-0.593****	0.036	0.324	**0.569****	**0.482****	**-0.399***	-0.018
NTSM	Chloroflexi	Anaerolinea	-0.247	0.321	-0.333	**-0.515****	0.208	**0.618****	0.312	0.225	**-0.407***	0.103
NTSM	Firmicutes	Clostridium_sensu_stricto_13	-0.361	0.261	0.127	-0.248	0.162	**0.421***	0.311	0.263	-0.219	-0.185
NTSM	Verrucomicrobia	Candidatus_Xiphinematobacter	**-0.674****	**0.568****	**0.384***	**-0.411***	0.021	0.165	**0.522****	**0.435***	**-0.408***	0.002
NTSM	Chlorobi	Ignavibacterium	0.521**	-0.241	**-0.401***	0.097	0.112	0.249	-0.201	-0.305	0.146	0.224
NTSM	Firmicutes	Thermincola	0.33	-0.195	-0.268	0.075	0.089	0.287	-0.168	-0.312	-0.092	-0.138
NTSM	Proteobacteria	Pseudomonas	0.621**	**-0.539****	-0.171	**0.451***	0.068	-0.169	**-0.523****	**-0.478****	**0.386***	0.007
NTSM	Spirochaetae	Spirochaeta_2	0.73**	**-0.577****	-0.306	**0.458***	0.083	-0.159	**-0.649****	**-0.512****	**0.403***	0.336
NTSM	Actinobacteria	Haloactinopolyspora	**-0.611****	**0.54****	-0.117	**-0.632****	0.181	**0.431***	**0.524****	**0.571****	**-0.575****	-0.278
NTSM	Proteobacteria	Piscinibacter	**-0.664****	**0.671****	-0.105	**-0.717****	0.069	**0.49****	**0.653****	**0.668****	**-0.616****	-0.135

56 SOC = soil organic carbon; SEOC = salt-extractable organic carbon; SEOC/SOC = ratio of SEOC to SOC; SM = soil moisture; TN = total nitrogen content; C/N =

57 ratio of SOC to TN; NH_4^+-N = ammonium nitrogen; NO_3^--N = nitrate nitrogen. * means $p < 0.05$,** means $p < 0.01$, *** means $p < 0.0001$. Red number means

58 significant negative correlation and blue number means significant positive correlation.

60   Table S4. The relative abundances of dominant phyla at different depths within three disturbance practices.

Phylum	Tillage	Soil depth(cm)									
		0-10	10-20	20-40	40-60	60-90	90-120	120-150	150-200	200-250	250-300
Acidobacteria	CT	10.97±2.42a	11.25±2.25a	10.30±3.12a	11.24±1.46a	7.94±2.63a	8.53±0.56a	10.26±1.81a	10.21±1.42a	9.94±0.62a	10.62±0.55a
	NTNS	11.25±2.39a	14.72±1.73a	13.33±1.27a	9.66±0.89a	8.90±2.61a	9.06±2.03a	6.46±2.11b	9.16±2.28a	7.02±3.10a	10.62±1.37a
	NTSM	11.22±4.10a	16.39±3.96a	13.29±1.20a	11.23±0.86a	9.58±0.83a	8.71±0.82a	9.95±0.67a	8.89±0.19a	9.45±0.92a	9.94±0.53a
		BC	D	CD	ABC	A	A	A	AB	A	ABC
Actinobacteria	CT	16.31±4.30a	25.59±10.25a	36.75±8.40a	33.29±7.31a	32.40±14.77a	39.85±4.54a	23.87±13.38a	10.73±4.61a	8.79±3.48a	14.59±6.34a
	NTNS	11.21±1.05a	14.14±4.86a	21.10±3.69a	31.95±1.76a	28.07±7.97a	26.81±10.20a	18.93±8.40a	12.45±5.72a	6.86±0.71a	10.98±4.99a
	NTSM	16.45±4.70a	15.22±0.99a	30.34±1.66a	27.92±1.27a	34.74±4.81a	34.84±3.96a	24.63±4.89a	12.89±8.00a	5.37±0.42a	7.51±0.78a
		BC	CD	E	E	E	E	B	AB	A	F
Bacteroidetes	CT	7.71±2.10a	4.59±1.00a	2.20±0.94a	3.35±1.15a	5.57±6.84a	2.36±1.63a	1.77±1.08a	1.57±1.10a	0.78±0.20a	1.58±0.32a
	NTNS	12.72±4.35a	5.61±1.35a	4.58±2.47a	3.44±1.27a	11.30±14.44a	9.12±10.07a	7.25±5.20a	1.88±0.16a	13.68±20.79a	1.55±0.55a
	NTSM	8.69±0.52a	6.62±1.02a	3.45±1.07a	2.88±0.31a	2.15±0.74a	3.51±1.46a	2.29±1.58a	3.79±2.59a	2.13±1.68a	2.73±1.29a
		C	BC	AB	AB	AB	AB	AB	AB	AB	A
Chloroflexi	CT	4.44±0.65a	8.87±1.22a	8.56±1.97a	11.52±0.99a	7.58±3.70a	12.06±2.62a	14.99±4.28ab	18.48±4.36a	15.02±1.82a	15.25±1.81a
	NTNS	5.70±0.90a	9.75±0.79a	9.99±1.50a	9.91±2.04a	8.11±3.30a	9.14±2.01a	10.80±2.25b	15.81±1.92a	12.75±7.97a	15.01±2.04a
	NTSM	5.30±0.83a	8.63±1.11a	7.82±2.38a	10.88±1.07a	9.11±1.05a	12.61±2.34a	16.89±1.40a	19.00±1.07a	16.04±2.28a	13.26±2.22a
		A	BC	BC	C	B	C	D	E	D	D
Firmicutes	CT	0.22±0.04a	0.65±0.25a	0.68±0.23a	0.41±0.15a	2.13±2.74a	0.84±0.34a	0.69±0.31a	0.39±0.31a	1.54±1.63a	1.18±0.50a
	NTNS	1.77±2.43a	0.38±0.10a	0.77±0.63a	1.14±1.43a	3.96±5.65a	3.14±3.38a	4.24±5.72a	0.62±0.42a	4.71±5.95a	0.60±0.38a
	NTSM	0.38±0.11a	0.48±0.12a	0.64±0.24a	0.38±0.06a	1.18±0.82a	1.50±0.36a	1.42±0.52a	1.23±0.59a	1.75±1.65a	1.31±0.59a
		A	A	AB	A	AB	AB	AB	AB	B	AB
Gemmatimonadetes	CT	6.72±0.07a	7.11±0.33a	6.03±0.86a	5.72±0.40a	5.63±1.79a	5.89±0.98a	6.15±0.27a	6.29±1.04a	4.42±0.97a	3.99±0.54a
	NTNS	6.65±1.75a	6.11±1.08a	5.85±0.48a	6.13±0.75a	5.34±2.21a	5.72±1.53a	6.08±1.74a	5.62±1.19a	3.79±1.23a	4.43±0.46a
	NTSM	5.68±0.12a	6.10±0.37a	5.87±0.24a	6.67±0.72a	5.58±1.05a	5.62±1.21a	6.26±0.15a	5.29±0.59a	3.76±0.50a	3.83±0.73a
		A	A	A	A	A	A	A	A	B	B
Latescibacteria	CT	0.47±0.35a	0.70±0.15b	1.50±0.66a	1.92±0.16a	1.58±1.17a	2.14±0.81a	2.62±0.99a	1.92±1.03a	1.37±0.34a	1.05±0.04a

	NTNS	0.47±0.05a	1.52±0.13a	2.00±0.78a	1.70±0.43a	1.53±0.81a	1.43±0.19a	1.25±0.72a	1.41±0.28a	0.58±0.33b	0.96±0.18a
	NTSM	0.39±0.16a	0.93±0.22b	1.73±0.31a	2.25±0.41a	2.25±0.71a	2.38±1.08a	3.32±2.31a	1.50±0.56a	0.82±0.16ab	1.03±0.39a
		A	BC	D	D	D	D	D	CD	B	BC
Microgenomates	CT	0.15±0.13a	0.34±0.27a	0.25±0.15a	0.42±0.29a	0.18±0.07a	0.50±0.41a	1.65±1.45a	2.28±1.79a	2.57±0.85a	3.15±0.33a
	NTNS	1.11±1.19a	0.56±0.27a	0.54±0.10a	0.25±0.14a	0.25±0.17a	0.47±0.34a	0.75±0.70a	2.18±0.53a	3.93±3.67a	3.77±1.33a
	NTSM	1.17±0.83a	0.85±0.21a	0.48±0.20a	0.60±0.12a	0.40±0.14a	0.60±0.32a	1.10±0.71a	2.70±0.77a	6.04±3.86a	4.01±0.49a
		AB	AB	AB	AB	A	AB	B	C	D	D
Nitrospirae	CT	1.45±0.42a	2.12±0.10b	6.56±2.39a	7.27±2.40a	6.45±1.95a	6.53±0.93a	8.37±2.48a	8.95±1.14a	10.94±2.01a	8.23±1.64a
	NTNS	1.68±0.42a	3.49±0.30a	5.44±2.79a	5.98±1.79a	5.58±2.52a	5.49±2.08a	6.30±2.51a	8.77±1.39a	6.37±3.74a	9.44±1.77a
	NTSM	1.50±0.12a	2.79±0.59ab	7.18±0.76a	7.72±0.87a	6.33±1.32a	5.99±0.69a	7.86±1.61a	9.55±1.86a	10.50±0.77a	9.66±2.27a
		A	B	C	C	C	C	D	D	D	D
Parcubacteria	CT	0.41±0.21a	0.50±0.06b	0.26±0.11b	0.40±0.13a	0.42±0.14a	0.55±0.10a	1.34±0.86a	4.58±4.11a	8.40±0.86a	7.68±2.24a
	NTNS	1.38±0.89a	0.82±0.30ab	0.52±0.09a	0.49±0.22a	0.40±0.24a	0.38±0.21a	3.23±4.78a	3.71±2.12a	5.57±4.43a	6.60±0.70a
	NTSM	1.04±0.14a	1.06±0.30a	0.43±0.15ab	0.53±0.22a	0.38±0.02a	0.56±0.16a	0.68±0.29a	5.07±3.23a	9.37±1.25a	4.75±1.92a
		AB	AB	A	AB	A	AB	B	C	D	D
Planctomycetes	CT	3.59±2.72a	2.71±0.83b	1.51±0.30b	2.31±0.92a	1.09±0.49a	1.41±0.56a	1.99±0.25a	2.01±0.34b	1.76±0.21a	2.13±0.31a
	NTNS	4.52±2.72a	7.95±2.24a	3.48±1.18a	1.93±0.52a	1.62±0.82a	1.83±1.39a	1.66±0.89a	3.09±0.39a	2.16±1.47a	2.86±0.96a
	NTSM	5.14±0.41a	6.50±1.04a	2.19±0.52ab	2.27±0.52a	1.18±0.45a	1.28±0.34a	1.74±0.45a	2.15±0.34b	2.28±0.42a	2.19±0.77a
		C	D	BC	BC	A	AB	ABC	C	ABC	C
Proteobacteria	CT	42.48±8.42a	31.01±7.20a	21.70±4.94a	17.84±2.00a	25.04±13.62a	15.23±0.20b	18.41±1.18b	22.01±2.86a	23.32±6.01a	18.93±3.75b
	NTNS	33.50±2.43a	28.43±0.73a	26.56±6.21a	22.26±4.25a	20.17±2.38a	20.42±3.04a	27.44±4.76a	28.36±11.14a	21.46±0.93a	18.58±0.86b
	NTSM	35.85±1.47a	27.79±1.87a	21.62±2.49a	20.94±0.10a	22.32±6.42a	17.21±0.65ab	16.33±0.95b	18.61±1.27a	19.48±2.98a	26.09±2.82a
		D	C	B	AB	AB	A	AB	B	AB	AB
Saccharibacteria	CT	0.70±0.33a	0.91±1.14a	0.20±0.14a	0.13±0.06a	0.23±0.14a	0.22±0.09a	0.32±0.23a	0.21±0.12a	0.20±0.12a	0.42±0.22a
	NTNS	1.45±0.58a	0.27±0.15a	0.28±0.11a	0.32±0.25a	0.25±0.12a	0.34±0.38a	0.25±0.02a	0.10±0.04a	0.45±0.26a	0.42±0.16a
	NTSM	1.59±1.39a	0.65±0.28a	0.26±0.11a	0.23±0.11a	0.25±0.09a	0.28±0.08a	0.11±0.08a	0.19±0.01a	1.67±1.82a	1.30±1.14a
		E	BCD	AB	AB	AB	ABC	AB	A	CD	D

Verrucomicrobia	CT	2.40±1.80a	1.34±0.21b	1.16±0.11b	1.56±0.64a	0.81±0.10a	1.04±0.24a	1.01±0.35a	0.84±0.08a	0.46±0.16a	1.18±0.50a
	NTNS	3.22±1.20a	2.52±0.34a	2.28±0.40a	1.55±0.39a	1.17±0.62a	1.15±0.57a	0.90±0.36a	0.93±0.29a	0.68±0.37a	1.69±0.61a
	NTSM	3.00±0.56a	2.74±0.41a	1.67±0.33ab	1.42±0.09a	0.83±0.06a	0.98±0.10a	0.74±0.31a	0.83±0.25a	0.49±0.03a	1.10±0.61a
		E	D	CD	C	B	B	AB	AB	A	BC

Different letters indicate significant differences at P<0.05.

63 Table S5 Rooting depths of crops under different tillage managements. Only crops with more than two tillage managements are retained.

ID	Reference	Crop	Tillage Practices	Straw mulching	Root depth (cm)	Location	Date
1	Bunna et al., 2011	Mungbean	Conventional	No	15.7	Takeo, Cambodia	2010, 1
		Mungbean	Conventional	Yes	17	Takeo, Cambodia	2010, 1
		Mungbean	Conventional	Yes	16	Takeo, Cambodia	2010, 1
		Mungbean	Conventional	Yes	16.8	Takeo, Cambodia	2010, 1
		Mungbean	Conventional	No	17	Kampong Thom, Cambodia	2009, 12
		Mungbean	Conventional	Yes	20	Kampong Thom, Cambodia	2009, 12
		Mungbean	Conventional	Yes	19	Kampong Thom, Cambodia	2009, 12
		Mungbean	Conventional	Yes	20	Kampong Thom, Cambodia	2009, 12
		Mungbean	Conventional	No	12.9	CARDI in Cambodia	2010, 2
		Mungbean	Conventional	Yes	13.2	CARDI in Cambodia	2010, 2
		Mungbean	Conventional	Yes	14.9	CARDI in Cambodia	2010, 2
		Mungbean	Conventional	Yes	15.7	CARDI in Cambodia	2010, 2
2	Kemper et al.,2011	Wheat	No-tillage	No	112	Mandan, ND	1988
		Wheat	Conventional	No	87	Mandan, ND	1988
		Wheat	No-tillage	No	89	Mandan, ND	1989
		Wheat	Conventional	No	69	Mandan, ND	1989
		Wheat	No-tillage	No	76	Mandan, ND	1990
		Wheat	Conventional	No	66	Mandan, ND	1990
		Sunflower	No-tillage	No	193	Mandan, ND	1992
		Sunflower	Conventional	No	143	Mandan, ND	1992
		Wheat	No-tillage	No	150	Oshkosh, WI	2008
		Wheat	Conventional	No	75	Oshkosh, WI	2008
		Corn	No-tillage	No	142	Oshkosh, WI	2009
		Corn	Conventional	No	74	Oshkosh, WI	2009
		Corn	No-tillage	No	104	Omro, WI	2009

		Corn	Conventional	No	56	Omro, WI	2009
		Corn	No-tillage	No	122	Oshkosh, WI	2009
		Corn	Conventional	No	64	Oshkosh, WI	2009
		Corn	No-tillage	No	131	Oshkosh, WI	2009
		Corn	Conventional	No	67	Oshkosh, WI	2009
		Corn	Conventional	No	58	Wisconsin	2009
		Corn	Conventional	No	69	Wisconsin	2009
		Corn	No-tillage	No	122	Wisconsin	2009
		Corn	No-tillage	No	132	Wisconsin	2009
3	Merrill et al.,2002	Sunflower	No-tillage	No	161	Morton County, ND	1995
		Sunflower	No-tillage	No	131	Morton County, ND	1996
		Sunflower	No-tillage	No	144	Morton County, ND	1997
		Wheat	No-tillage	No	126	Morton County, ND	1995
		Wheat	No-tillage	No	125	Morton County, ND	1996
		Wheat	No-tillage	No	117	Morton County, ND	1997
		Soybean	No-tillage	No	109	Morton County, ND	1995
		Soybean	No-tillage	No	103	Morton County, ND	1996
		Soybean	No-tillage	No	86	Morton County, ND	1997
4	Turman et al.,1995	Soybean	Conventional	No	80	Portageville,MO, USA	1992, 14 May
		Soybean	Conventional	No	66	Portageville,MO, USA	1992, 15 June
		Soybean	Conventional	No	93	Portageville,MO, USA	1992, 7 July
		Soybean	Conventional	No	49	Portageville,MO, USA	1993, 12 May
		Soybean	Conventional	No	41	Portageville,MO, USA	1993, 2 June
		Soybean	Conventional	No	53	Portageville,MO, USA	1993, 21 June
		Soybean	No-tillage	No	76	Portageville,MO, USA	1992, 14 May
		Soybean	No-tillage	No	87	Portageville,MO, USA	1992, 15 June
		Soybean	No-tillage	No	84	Portageville,MO, USA	1992, 7 July

		Soybean	No-tillage	No	45	Portageville,MO, USA	1993, 12 May
		Soybean	No-tillage	No	32	Portageville,MO, USA	1993, 2 June
		Soybean	No-tillage	No	40	Portageville,MO, USA	1993, 21 June
5	Dwyer et al.,1996	Corn	Conventional	No	80.6	Ottawa, Ontario	1988
		Corn	Conventional	No	101.3	Ottawa, Ontario	1989
		Corn	No-tillage	No	70.6	Ottawa, Ontario	1988
		Corn	No-tillage	No	91.3	Ottawa, Ontario	1989
		Corn	Conventional	No	90.6	Ottawa, Ontario	NA(Clay loam)
		Corn	Conventional	No	91.3	Ottawa, Ontario	NA(Sany loam)
		Corn	No-tillage	No	89.4	Ottawa, Ontario	NA(Clay loam)
		Corn	No-tillage	No	72.5	Ottawa, Ontario	NA(Sany loam)
6	Dwyer et al.,1988	Corn	Conventional	No	90	Ottawa, Ontario	1982
		Corn	Conventional	No	90	Ottawa, Ontario	1982
		Corn	Conventional	No	90	Ottawa, Ontario	1982
		Corn	Conventional	No	90	Ottawa, Ontario	1982
		Corn	Conventional	No	115	Ottawa, Ontario	1983
		Corn	Conventional	No	98	Ottawa, Ontario	1983
		Corn	Conventional	No	90	Ottawa, Ontario	1983
		Corn	Conventional	No	99	Ottawa, Ontario	1983
		Corn	Conventional	No	90	Ottawa, Ontario	1984
		Corn	Conventional	No	86	Ottawa, Ontario	1984
		Corn	Conventional	No	75	Ottawa, Ontario	1984
		Corn	Conventional	No	90	Ottawa, Ontario	1984
		Soybean	Conventional	No	75	Ottawa, Ontario	1982
		Soybean	Conventional	No	80	Ottawa, Ontario	1982
		Soybean	Conventional	No	69	Ottawa, Ontario	1982
		Soybean	Conventional	No	75	Ottawa, Ontario	1982

Soybean	Conventional	No	75	Ottawa, Ontario	1983
Soybean	Conventional	No	75	Ottawa, Ontario	1983
Soybean	Conventional	No	74	Ottawa, Ontario	1983
Soybean	Conventional	No	94	Ottawa, Ontario	1983
Soybean	Conventional	No	70	Ottawa, Ontario	1984
Soybean	Conventional	No	77	Ottawa, Ontario	1984
Soybean	Conventional	No	68	Ottawa, Ontario	1984
Soybean	Conventional	No	70	Ottawa, Ontario	1984

64
65

References

1. Bunna, S., Sinath, P., Makara, O., Mitchell, J. & Fukai, S. Effects of straw mulch on mungbean yield in rice fields with strongly compacted soils. *Field Crops Res.* **124**, 295-301 (2011).

2. Kemper, W. D., Schneider, N. N. & Sinclair, T. R. No-till can increase earthworm populations and rooting depths. *J. Soil Water Conserv.* **66**, 13A-17A (2011).

3. Merrill, S. D., Tanaka, D. L. & Hanson, J. D. Root length growth of eight crop species in Haplustoll soils. *Soil Sci. Soc. Am. J.* **66**, 913-923 (2002).

4. Turman, P., Wiebold, W., Wrather, J. & Tracy, P. Effect of planting date and tillage system on soybean root growth. *J. Plant Nutr.* **18**, 2579-2594 (1995).

5. Dwyer, L. *et al.* Root mass distribution under conventional and conservation tillage. *Can. J. Soil Sci.* **76**, 23-28 (1996).

6. Dwyer, L., Stewart, D. & Balchin, D. Rooting characteristics of corn, soybeans and barley as a function of available water and soil physical characteristics. *Can. J. Soil Sci.* **68**, 121-132 (1988).