1   **Fitness decline in hatchery-enhanced salmon populations**

2   **is manifested by global warming**

4   Running head: Hatchery-climate interaction and fitness

7   Shuichi Kitada[1] and Hirohisa Kishino[2]

[1] Tokyo University of Marine Science and Technology, Tokyo 108-8477, Japan.

[2] Graduate School of Agriculture and Life Sciences, The University of Tokyo,

Tokyo 113-8657, Japan.

Correspondence
Shuichi Kitada, Tokyo University of Marine Science and Technology, Tokyo
108-8477, Japan.
Email: kitada@kaiyodai.ac.jp

Despite historical maximum hatchery releases and biomass of chum salmon in the North Pacific, chum salmon returning to Japan, where the world largest chum salmon hatchery programme runs, have declined substantially for two decades. However, the causal mechanisms are unknown. We analysed catch/release data of chum salmon and SST in Japanese coastal waters, and synthesized LDH (lactate dehydrogenase) and PEPB (tripeptide aminopeptidase) gene frequencies of 21,911 adult chum salmon collected from 166 rivers and areas in the North Pacific Rim. We also evaluated athletic performance of Japanese chum salmon in comparison with foreign salmon populations on social network videos. We found significant negative correlation between the average SST for spring and summer and total catch of Japanese chum salmon (r=-0.77), and 59% of the variation in the decreasing catch was explained by the increasing SST. Frequencies of the thermal adaptation genes LDH-A1 and LDH-B2 were altered disregarding with directions of natural selection which eventually lead hatchery fish to adapt to colder environments, while those of the digestive gene PEPB-1 were positively correlated with latitude. Because LDH-A and LDH-B are found in skeletal and cardiac muscles, this change may have lowered the survival rates particularly for juveniles and growth of Japanese chum salmon. We argue that the long-term hatchery releases have reduced the athletic ability of hatchery fish, resulting in a continuous decline in fitness of whole populations. A genetic mechanism of fitness decline in hatchery-enhanced salmon populations is first manifested by global warming at the southern margin of the species.

Keywords: allozyme gene frequencies, hatchery, LDH, PEPB, Pacific salmon

1 | INTRODUCTION

Release of hatchery-reared animals is one of the popular tools for fishery management (Laikre, Schwartz, Waples, & Ryman, 2010) and more than 26 billion juveniles of 180 marine species are released into the wild annually (Kitada, 2018). Evidence for reduced fitness of hatchery-reared fish in the wild and altered age and size at maturation is a serious reason for caution in hatchery stocking (Araki, Cooper, & Blouin, 2007, 2009; Bolstad et al., 2017; Fleming et al., 2000; McGinnity et al., 2003; Reisenbichler & McIntyre, 1977; Reisenbichler & Rubin, 1999). However, there has been no empirical evidence for reduced fitness in wild populations caused by hatchery releases (Kitada, 2018). Decreased time duration of spawning in natal freshwater habitat (Cline, Ohlberger, & Schindler, 2019) and advanced spawning dates by 1–3 weeks (Tillotson, Barnett, Bhuthimethee, Koehler, & Quinn, 2019) were found for sockeye salmon. The results suggested that artificial selection in hatcheries might drive maladaptation to climate change and act to opposite directions of natural selection (Tillotson, Barnett, Bhuthimethee, Koehler, & Quinn, 2019). However, mechanisms for hatchery-climate interactions on fitness largely remain unknown. Understanding thermal adaptation is a challenging theme and active studies are currently performed (Chen, Farrell, Matala, & Narum, 2018). Here, we focus on declining hatchery-enhanced Japanese chum salmon (*Oncorhynchus keta*, Salmonidae) populations at the southern margin of the species range to elucidate the mechanisms for fitness decline in warming environment.

Chum salmon hatchery stocking represents one of the world's major marine stock enhancement and sea-ranching programmes (Kitada, 2018). Around 60% of chum salmon in the North Pacific are of hatchery origin (Ruggerone & Irvine, 2018). Japanese chum salmon stocking, initiated in the late 1880s, is the world's largest hatchery release

programme (Amoroso, Tillotson, & Hilborn, 2017; Kitada, 2018; Miyakoshi, Nagata, Kitada, & Kaeriyama, 2013). Releases of chum salmon juveniles from ~260 salmon hatcheries in Japan has increased remarkably since the 1970s to ~1.5 billion in 2018 (Figures S1, S2). The numbers of chum salmon returning to Japan sharply increased after the 1970s, but steeply decreased since 1996 when the catch recorded the historical maximum of 81 million fish, despite the continuous efforts of hatchery releases.

In Japanese chum salmon hatcheries, eggs and sperm are taken artificially from parent fish collected in the enhanced rivers (Miyakoshi, Nagata, Kitada, & Kaeriyama, 2013). Fertilized eggs are kept undisturbed in hatcheries, avoiding vibration, until hatching (http://salmon.fra.affrc.go.jp/zousyoku/sakemasu.html). Rearing water temperature is kept stable ~8°C on average, which mimics temperature of spring water near redds, while the mean water temperature is much lower in the natural stream for this period; it ranged 1.1–2.8°C in the tributary of Ishikari River, Hokkaido (Ando et al., 2014). Hatching begins when the accumulated temperature reaches 480 degrees (~two months after fertilisation). Hatched larvae (BL ~19mm, BW ~0.05g) emerge (BL ~38mm, BW ~0.28g) 40 days after and grow to fry (BL ~43mm, BW ~0.69g) two months after hatching (Kaeriyama, 1986). They are fed to the size at release (BL ~50mm, BW ~1g) in rearing ponds during January and May until sea water temperature exceeds 5 °C (https://www.hro.or.jp/list/fisheries/marine/o7u1kr000000dc2m.html). Rearing duration primarily depends on rearing temperature, but juveniles are reared for more than six months in hatcheries. Survival rates are very high, usually >90%.

Since the early 1980s, efforts to increase the early-run population have caused changes in run timing distribution and the late-run populations have almost disappeared in

Hokkaido (Miyakoshi, Nagata, Kitada, & Kaeriyama, 2013), the major production area

producing more than 80% of Japanese chum salmon. The past transplantation history of

Japanese chum salmon populations has altered population structures with some

populations nested within management units (Beacham, Sato, Urawa, Lei, & Wetklo,

2008; Kitada, 2014). Unintended and intended artificial selection in hatcheries can relax

natural selection and reduce fitness of hatchery-reared salmonids in the wild (Araki,

Cooper, & Blouin, 2007, 2009; Fleming et al., 2000; McGinnity et al., 2003;

Reisenbichler & McIntyre, 1977; Reisenbichler & Rubin, 1999). Genetic adaptation to

hatchery environments may occur within a single generation or early generations of

hatchery fish (Christie, Marine, French, & Blouin, 2012; Christie, Marine, Fox, French,

& Blouin, 2016; Le Luyer et al., 2017). Moreover, hatchery environments favour certain

genotypes; when hatchery broodstock is introgressed by farmed escapees, farmed

genotypes associated with domestication are amplified in the stocked population (Hagen

et al., 2019). Because all chum salmon returning to Japan could be hatchery-released

fish or wild-born hatchery descendants after long-term hatchery operations (Kitada,

2018), historical releases may have selected genes adapted to hatchery environment.

Allozyme markers were used to study the population genetics of chum salmon during

the 1970s and the 1990s. Here, we focus on LDH (lactate dehydrogenase) as candidate

gene markers by which underlying physiological adaptations were pinpointed (Chen,

Farrell, Matala, & Narum, 2018). White skeletal muscle, whose metabolism is

predominantly anaerobic expresses the Ldh-A locus, while red muscle, which possesses

significant aerobic metabolism, expresses almost exclusively Ldh-B (Somero, 2004).

The A_4-LDH gene works principally in the conversion of pyruvate to lactate (i.e.,

anaerobic glycolysis), while B_4-LDH principally converts lactate to pyruvate (i.e.,

gluconeogenesis and aerobic metabolism) (Somero, 2004, 2010; Powers, Lauerman,

Crawford, & DiMichele, 1991). We also analyse PEPB (tripeptide aminopeptidase)

because the aminopeptidase PEPB is a digestive enzyme of fish (Govoni, Boehlert, &

Watanabe, 1986), which may relate to selection in hatcheries.

2 | DECLINING JAPANESE CHUM SALMON POPULATIONS

Catch and release data were retrieved from the statistics of North Pacific Anadromous

Fish Commission (NPAFC, 2019). Simple return rates (number of returned fish after

four years /number of released juveniles) in Hokkaido and Honshu were cited from

FRA (http://salmon.fra.affrc.go.jp, accessed August, 2019). Major components of

returned fish were age 3-age 6 fish and age 4 was the largest in proportions. The simple

return rate therefore is an expectation of the true return rate throughout their lives.

Spring, summer, autumn and winter sea surface temperature (SST) data for Kushiro-oki

(Hokkaido Pacific coast), Hokkaido Sea of Japan coasts, Sanriku-oki (Honshu Pacific

coast) and Honshu Sea of Japan coasts were retrieved from Japan Meteorological

Agency (http://www.data.jma.go.jp/kaiyou/shindan/index_gw.html, accessed August,

2019). We averaged SST over the four areas for each season and for spring and

summer. Exponential curves were fitted to the simple return rates for Hokkaido and

Honshu during 1996 and 2018, respectively. The linear regression analyses between the

numbers of chum salmon returning to Japan and the SST anomaly were performed

using the lm function. We also fitted an exponential curve to the salmonid relative

reproductive success (RRS) data (Araki, Cooper, & Blouin, 2007) and estimated the

coefficients using the nls function. We calculated mean body weights of chum salmon

returning to Japan for each year between 1975 and 2018 by dividing the catch weight by

the number of returned fish, based on the catch statistics of NPAFC. All analyses were

performed and figures were created using R statistical software (https://www.r-

project.org/). All data used for the analyses were included in Supplementary Data

excluding easy-accessible NPAFC salmon statistics.

The catch of chum salmon in Japan steeply decreased, while the total catch remained

high in the North Pacific (Figure 1a). A significant exponential decline in return rates

was found for Hokkaido and Honshu (Figure 1b, $t = -5.52, p = 0.0082$; $t =$

$3.69, p = 0.0014$, respectively). The exponential rate of decline in Honshu was about

twice that in Hokkaido, suggesting that the warming effect was stronger in southern

Japan (Table 1), but there was no significant difference between them, with a combined

average of $3.1 \pm 0.9\%$ per year ($z = 0.983, p = 0.1627$). The exponential decay rates of

chum salmon were much lower than that of the red sea bream (Kitada et al, 2019), and

of RRS of captive-reared salmonids (Araki, Cooper, & Blouin, 2007) (Table 1).

Regression analysis combining Hokkaido and Honshu indicated that the average sea

surface temperature (SST) anomaly over northern Japan was negatively correlated with

the overall decrease in the Japanese catch of chum salmon (1996–2018) ($r = -0.67$)

(Table 2). Interestingly, the average SST in spring (April–June) and summer (July–

September) was highly correlated with the catch, while in autumn and winter there was

no significant correlation. The correlation between the average SST for spring and

summer and total catch was much higher ($r = -0.77, \mathrm{t} = -5.52, \mathrm{df} = 21, \mathrm{p} =$

1.76×10^{-5}), and 59% of the variation in decreasing chum salmon catch was explained

by the increasing spring and summer SST anomaly (Figure 1c).

The number of chum salmon returned and caught was the largest in the Sea of Okhotsk

and the Pacific coast of northern Japan that was affected by the cold Oyashio current

particularly in Iwate Prefecture in 1996 when the catch in Japan recorded the historical

maximum of 81 million (Figure 2), showing that chum salmon abundance was rich in

the cold waters. In contrast, they were very small on the Sea of Japan coast that is

affected by the Tsushima warm current and the southern Pacific coast. In 2006,

populations in the Pacific coast of Honshu were rapidly decreased. In contrast,

decreases in the Pacific coast of Hokkaido were not substantial and even high in the Sea

of Okhotsk. However, substantial declines occurred in all areas in 2014, which further

shrunk in 2017. On the other hand, numbers of chum salmon released were stable for

the two decades in whole Japan (Figure 3). The mean body weights of chum salmon

returning to Japan have continued to decrease since 2004 (Figure S3).

3 | THERMAL ADAPTATION AND DIGESTIVE GENE FREQUENCIES

We screened scientific literature, using the Google Scholar online database search

system to collect publications using keyword searches of "salmon", "allozyme", "LDH"

and "PEPB". Studies known to us were also manually screened. We found studies

including gene frequencies of LDH-A1, LDH-B2 and PEPB-1 in the Pacific Rim

including whole Japan (Kijima, & Fujio, 1979; Okazaki, 1982), Russia, Alaska, British

Columbia and Washington (Seeb, Crane, & Gates, 1995; Winans, Aebersold, Urawa, &

Varnavskaya, 1994). We organized frequencies of LDH-A1, LDH-B2 and PEPB-1

genes and recorded approximate longitudes and latitudes of the sampling sites based on

the names of rivers and/or areas and maps of the original studies using Google Maps.

Maps were drawn using the sf package and sampling points were located on the maps.

The allele frequency for ith sampling point (x_i) was visualized by colour gradients in

the data range based on: $\mathrm{rgb}(1 - y_{0,i}, 0, y_{0,i})$, where $y_{0,i} = (x_{0,i} - a)/(b - a)$ and

$x_{0,i} = pmin(pmax(x_i, a), b)$. We set a=0.7 and b=1.0 for LDH-A1, and a=0.6 and

b=1.0 for LDH-B2 and PEPB-1, considering the ranges of the gene frequencies. This

conversion visualizes the magnitude of the gene frequencies at the sampling point in

colours between blue and red on the map.

The gene frequencies of 21,911 adult chum salmon from 21 Russian rivers and 81

Japanese rivers (including the same rivers analysed in different studies) were organized,

and 64 rivers and areas in America (Figure 4). The Japanese samples were caught at

weirs in enhanced rivers and hatcheries (Kijima & Fujio, 1979; Okazaki, 1982) and

were, therefore, hatchery-reared fish and/or hatchery descendants, while the Russian

samples were collected by stream surveys and were expected to be wild fish because the

majority of Russian chum salmon are maintained by wild fish (Winans, Aebersold,

Urawa, & Varnavskaya, 1994). Samples collected from hatcheries were excluded in

samples from Alaska, Alaskan Peninsula, British Columbia and Washington (Seeb,

Crane, & Gates, 1995). The American samples were collected during 1986 and 1994,

when the numbers of released chum salmon were low and similar the levels in Russia

(Figure S1a). It is possible that some hatchery fish were included in the American

samples, but the numbers would have been low.

In the Russian and American samples, we found a clear geographical cline from south

to north in the LDH-A1 gene frequencies, and a negative correlation with latitude ($r =$

$-0.44, \text{t} = -4.07, \text{df} = 71, \text{p} = 0.000$) (Figure 5a, b). In the southern samples from

Washington and British Columbia, LDH-A1 gene frequencies were close to or exactly

1.0, while those from the Alaskan Peninsula were generally <0.95, and <0.8 in Alaska.

The outlier (0.63) in American samples was from the almost closed Peterson Lagoon,

Alaska (#148 in Fig. 3b). Alaska and Kamchatka had a similar range of gene

frequencies of around 0.7–0.9 (Figure 5a). Japanese samples exhibited similar variation

to the Russian and Alaska ones at higher latitudes. The gene frequency varied largely

among the Japanese samples but had a weak and positive correlation with latitude ($r =$

$0.27, t = 2.63, df = 89, p = 0.001$). Its range corresponded to the range of the gene

frequencies in Russia, Alaska and Canada (Figure 5a, b). The four outliers (#57, 97, 99,

101) were collected from the most southern coastal areas of the Sea of Japan (Figure

4d). In contrast, LDH-B2 gene frequencies were 1.0 or close to 1.0 for all samples

(Figure 5c, d), except for those from eastern Queen Charlotte Islands, Canada (#157 in

Figures. 4a, 5c) that are affected by the warm Alaska Current, and had the lowest gene

frequency (0.95). Almost all Russian samples from Kamchatka had slightly lower gene

frequencies than the others. Japanese samples had similar LDH-B2 gene frequencies to

those of the northern populations. In contrast, PEPB-1 gene frequencies were positively

correlated with latitude (Figure 5e, f).

4 | ATHLETIC PERFORMANCE OF JAPANESE CHUM SALMON: A VISUAL EVALUATION

We searched videos on the internet which could show athletic performance of Japanese

chum salmon and other salmonid species from a social network "youtube". We analysed

nine videos, of which four were on chum salmon from Hokkaido and fives were from

the Pacific Northwest, Alaska and England. In Okushibetsu River, Shiretoko, Hokkaido,

is an enhanced river, where thousands of chum salmon return every year (The Hokkaido

Shimbun Press, 2017). Almost fish were caught at the weir located ~500 m from the sea

and many were used for artificial fertilization in the hatchery. There are ~1,500 rivers in

Hokkaido and 238 rivers have lengths longer than 8 km. Hatchery releases of chum

salmon are operated in 140 major rivers and no hatchery is operated in small rivers

(Miyakoshi et al., 2013). Japanese chum salmon moved slowly. In contrast, sockeye

salmon moved vividly and went up the Hansen Creek to spawn (Taylor, 2014). Jumping

trials of returned Japanese chum salmon at a small fall, Hokkaido, Esashi Town facing

the Sea of Okhotsk (Sapporo TV House Stock Footage, 2015), resulted in no success for

64 jumps excluding two jumps which were difficult to judge. When we counted the two

jumps as succeeded ones, the rate of success was 3.0% (=2/66). At another small fall in

northern Hokkaido (Kawamura, 2008), the success was one out of 47 jumps (success

rate=2.1%). At a small weir in Hamamasu River, Hokkaido (Onishi, 2009), only one

chum salmon succeeded for 219 jumps (success rate=0.5%). These small rivers had no

hatchery, therefore they were strayed hatchery fish or natural-born fish in the rivers.

Chum salmon are very powerful swimmers even swimming across roads during floods

on the coast and along Puget Sound (Allison, 2016). The success rate of jumping was

15% (=10/66) for sockeye salmoln (*O. nerka*, Salmonidae) in Katmai National Park,

Alaska (WallnerPhotos, 2014), 33% (=2/6) for coho salmon (*O. kisutch*, Salmonidae) in

Lagunitas, California (Turtle Island Restoration Network, 2015) and 50% (=13/26) for

Atlantic salmon (*Salmo salar*, Salmonidae) in England (JettyBox Stock Video, 2015),

respectively. All species had much higher athletic performance for jumping and

upstreaming than Japanese chum salmon even in the stronger streams.

5 | DISCUSSION

A previous study found that ~30% of variation in the decreasing catches of Japanese

chum salmon was attributed to the increasing annual average SST anomaly (1996–

2016), and that 62% of the variation in Russian catches was explained by the SST

(Kitada, 2018). However, earlier tagging studies revealed that there were clear

differences in the run timing and migration routes between Russian (June–August) and

Japanese (September–November) chum salmon populations (Kondo, Hirano,

Nakayama, & Miyake, 1965; Morita, 2016). The fishing season for chum salmon begins

in late August in Hokkaido with a high fishing season in September to October. In

Honshu, the high season runs from October to November. The apparent differences in

run timing and migration routes between Japanese and Russian chum salmon suggest

that the majority of Japanese chum salmon are caught in Japanese coasts and rivers.

Therefore, the negative correlation between Japanese and Russian catches reported in a

previous study should be spurious. We found that warming SST in spring and summer

negatively correlated with declining Japan chum salmon populations. Juvenile chum

salmon (almost hatchery fish) migrate down to the sea in late winter to early summer

and remain in coastal waters until early August (Ueno & Ishida, 1996). Reduced chum

salmon returns could be caused by lower survival rates of juvenile stages in coastal

waters from spring to summer with warming SST. Ages at return of Japanese chum vary

between two and eight (http://salmon.fra.affrc.go.jp/zousyoku/sakemasu.html),

therefore the average SST value over two to eight years before the year of return could

affect the survival of juveniles. Indeed, the correlation between the eight year moving

average SST for spring and summer and total catch was significant ($r = -0.55, t =$

$-3.03, \mathrm{df} = 21, \mathrm{p} = 0.006$), suggesting that the chum salmon decline was caused by

reduced survival rates of juvenile stages in coastal waters from spring to summer with

warming SST. Natural selection on released fish that are used for broodstock may be

responsible for the lower exponential decay rate of chum salmon than that of the red sea

bream (Kitada et al., 2019) and RRS of captive-reared salmonids (Araki et al., 2007)

(Table 1). In contrast, a model-based prediction indicated that global warming

decreased suitable areas of habitat in the North Pacific Ocean and loss of migration

route to the Sea of Okhotsk, but positively affected growth and survival of Hokkaido

chum salmon populations (Kaeriyama, Seo, & Qin, 2014).

LDH-A-mediated reactions have the lowest energy requirement in most cold-adapted

species, and highest in most warm-adapted species (Somero, 2004), which agrees with

higher LDH-A gene frequencies of chum salmon in the lower latitudes. Because Japan

is located south of the other surveyed locations in Russia, Canada, and USA, a high

gene frequency close to 1.0, the value for British Columbia and Washington, is

predicted in this country. The values far below 1.0 may suggest substantial artificial

selection on the LDH-A1 gene. Population divergence for LDH measured by Q_{ST}

(=0.58) was greater than divergence measured by pairwise F_{ST} for microsatellite

markers (0-0.077) in brook trout populations (Dufresne, Barroux, DitleCadet, & Blier,

2015). The result indicated that that LDH has diversified more than neutral population

structure and significant plasticity of LDH for artificial selection (Whitlock, 2008). A

single amino acid substitution is sufficient to modify function and stability of

temperature adaptation of the LDH-A gene (Somero, 2004). An amino acid substitution

in the LDH-A gene, common in vertebrate skeletal (white) muscle, may occur, leading

to adaptation of hatchery-reared salmon to colder environments.

Furthermore, artificial selection in hatcheries slows movement of hatchery fish and

reduces agility for feeding in the wild. Indeed, mean body weights of chum salmon

returning to Japan have continued to decrease since 2004 (Figure S3). Intra-and

interspecific competition for food was very strong for chum salmon including other

salmonid species in the North Pacific (Kitada, 2018), supporting this speculation.

Rearing experiments (5°C and 9°C) for embryos and alevin of brook trout (*Salvelinus*

fontinalis, Salmonidae), cold adapted species, found that mass differences and routine

metabolic rate among fry were strongly influenced by population origins and rearing

temperatures (Cook, Wilson, & Burness, 2018). The water temperature (~8°C) for egg

hatching standardly used in Japan chum salmon hatcheries was much higher than that in

the natural stream. The artificial rearing condition with constant water temperature and

reduced vibration might alter the routine metabolic rate that needs low energy

requirement for LDH-A-mediated reactions as in most cold-adapted species.

The LDH-B2 gene frequencies in southern Hokkaido and Honshu were exactly 1.0

(Figures. 4d, 5c). The LDH-B2 gene frequencies of killifish (*Fundulus heteroclitus*,

Fundulidae) were correlated with the direction of change in mean water temperature.

They were well fitted to a sigmoid curve with values near 1.0 at latitude 44°N and 0.2–

0.8 at latitudes 36–42°N (Powers et al., 1991). LDH-B plays a key role in re-oxidizing

NADH to NAD^+ in low O_2 environment in redband trout (O. *mykiss gairdneri*,

Salmonidae) (Chen, Farrell, Matala, & Narum, 2018). Oxygen consumption in heart

ventricles was significantly greater in northern killifish populations than that in southern

populations, and the cardiac metabolism was significantly higher in populations from

cold climates than those from warm climates (Podrabsky, Javillonar, Hand, &

Crawford, 2000). These results consistently showed that LDH-B2 gene frequencies

were lower in southern populations and higher in northern populations. The LDH-B2

gene frequencies in Japan populations were 1.0, contrary to that expected in adaptation

to a warm-water environment, which might be caused by reduced heart (red muscle)

activity and athletic performance in hatchery fish. The videos were cited from

"youtube" with including only nine samples and not our observations. However, the

results agreed with our analyses, demonstrating weaker athletic performance of

Japanese chum salmon. In contrast, gene frequencies of PEPB-1 were positively

correlated with latitude. This agreed the population structure in the North Pacific based

on microsatellite markers (Beacham et al., 2009), suggesting that hatchery operations

did not compromise digestive ability.

Repeated hatchery operations substantially altered gene frequencies of thermally

adapted LDH-A1 and LDH-B2 in Japanese chum salmon populations, which could be

reflected in the reduced performance of skeletal and cardiac muscles. This may be a

causal mechanism for fitness decline in Japan chum salmon populations. Altered

distributions of the gene frequencies of LDH-A1 and LDH-B2 are likely to result from

adaptation to hatchery environments, which eventually should lead hatchery fish to

adapt to colder environments. Our results suggest that artificial and natural selection act

in opposite directions on LDH-A1 and LDH-B2 but exert no selective pressure on the

digestive enzyme PEPB-1. However, selection on LDH-A1 and LDH-B2 genes reduced

survival rates in the wild in a warming environment, particularly for juveniles. Reduced

agility for feeding could also reduce the growth of chum salmon. Reduced activity of

cardiac muscle could affect their long-distance migrations. An overwhelming proportion

of hatchery fish and hatchery descendants (Kitada, 2018) should reduce the fitness of

whole populations. The hatchery operation has continued for ~25 generations, assuming

5 years per generation. After 40 years of domestication (30–35 generations) of wild

foxes, 70%–80% became human-friendly, and all domesticated foxes that escaped for

days returned to the farm (Trut, 1999). Apparently, 'domesticated' Japan chum salmon

are now human-friendly and return to hatcheries each year (Figure S4). Fitness

reduction in a hatchery-reared population is cohort-specific and constant over time

within the cohort (Kitada et al., 2019), but it will continue as long as the current

hatchery operations continue with a variation reflecting the environmental effects.

ACKNOWLEDGEMENTS

We thank Yasuyuki Miyakoshi and Toshio Okazaki for valuable discussions. This study

was supported by the Japan Society for the Promotion of Science Grant-in-Aid for

Scientific Research 18K0578116 to S. K. and 19H04070 to HK.

AUTHOR CONTRIBUTIONS

S.K. and H.K conceived of the study, analysed the data and wrote the manuscript.

CONFLICT OF INTEREST

The authors have no conflict of interest to declare.

ORCID

Shuichi Kitada http://orcid.org/0000-0001-5838-0374

REFERENCES

Allison, T. J. (2016, Nov. 14) Skokomish River salmon cross the road/Part 2 [Video

file]. Retrieved from https://www.youtube.com/watch?v=YFZbciPzAYk,

https://oregonwild.org/wildlife/chum-salmon

Amoroso, R. O., Tillotson, M. D., & Hilborn, R. (2017). Measuring the net biological

impact of fisheries enhancement: Pink salmon hatcheries can increase yield, but

with apparent costs to wild populations. Canadian Journal of Fisheries and

Aquatic Sciences, 74, 1233–1242. https://doi.org/10.1139/cjfas-2016-0334

Ando, D., Shinriki, Y., Shimoda, K., Yasutomi, R., Sasaki, Y., Miyakoshi, Y., &

Nakajima, M. (2014). Effects of spawning time on the variation of vertebral

395 numbers in chum salmon *Oncorhynchus keta*. *Nippon Suisan Gakkaishi*, 80, 191–

396 200. [In Japanese with English abstract] https://doi.org/10.2331/suisan.80.191

397 Araki, H., Cooper, B., & Blouin, M. S. (2007). Genetic effects of captive breeding cause

398 a rapid, cumulative fitness decline in the wild. *Science*, 318, 100–103. doi:

399 10.1126/science.1145621

400 Araki, H., Cooper, B., & Blouin, M. S. (2009). Carry-over effect of captive breeding

401 reduces reproductive fitness of wild-born descendants in the wild. *Biology*

402 *Letters*, *5*(5), 621-624. https://doi.org/10.1098/rsbl.2009.0315

403 Beacham, T. D., Sato, S., Urawa, S. Lei, K. D., & Wetklo, M. (2008). Population

404 structure and stock identification of chum salmon *Oncorhynchus keta* from Japan

405 determined by microsatellite DNA variation. *Fishery Science*, 74, 983–994. doi:

406 10.1111/j.1444-2906.2008.01616.x

407 Bolstad, G. H., Hindar, K., Robertsen, G., Jonsson, B., Sægrov, H., Diserud, O. H., ... &

408 Barlaup, B. T. (2017). Gene flow from domesticated escapes alters the life history

409 of wild Atlantic salmon. Nature Ecology & Evolution, 1(5), 0124.

410 https://doi.org/10.1038/s41559-017-0124

411 Chen, Z., Farrell, A. P., Matala, A., & Narum, S. R. (2018). Mechanisms of thermal

412 adaptation and evolutionary potential of conspecific populations to changing

413 environments. *Molecular Ecology*, 27, 659–674.

414 https://doi.org/10.1111/mec.14475

415 Christie, M. R., Marine, M. L., French, R. A., & Blouin, M. S. (2012). Genetic

416 adaptation to captivity can occur in a single generation. *Proceedings of the*

417 *National Academy of Sciences*, 109, 238–242. doi: 10.1073/pnas.1111073109

Christie, M. R., Marine, M. L., Fox, S. E., French, R. A., & Blouin, M. S. (2016). A

single generation of domestication heritably alters the expression of hundreds of

genes. *Nature Communications*, *7*, 10676. https://doi.org/10.1038/ncomms10676

Cline, T. J., Ohlberger, J., & Schindler, D. E. (2019). Effects of warming climate and

competition in the ocean for life-histories of Pacific salmon. *Nature Ecology &*

Evolution, 3, 935–942. https://doi.org/10.1038/s41559-019-0901-7

Cook, C. J., Wilson, C. C., & Burness, G. (2018). Impacts of environmental matching

on the routine metabolic rate and mass of native and mixed-ancestry brook trout

(*Salvelinus fontinalis*) fry. *Conservation Physiology*, 6, coy023.

https://doi.org/10.1093/conphys/coy023

Dufresne, F., Barroux, A., DitleCadet, D., & Blier, P. U. (2015). Quantitative variation

for metabolic traits among brook trout populations inhabiting different

environments. *Journal of Zoology*, 297, 194–203.

https://doi.org/10.1111/jzo.12267

Fleming, I. A., Hindar, K., MjÖlnerÖd, I. B., Jonsson, B., Balstad, T., & Lamberg, A.

(2000). Lifetime success and interactions of farm salmon invading a native

population. *Proceedings of the Royal Society of London B*, 267, 1517–1523. doi:

10.1098/rspb.2000.1173

Govoni, J. J., Boehlert, G. W., & Watanabe, Y. (1986). The physiology of digestion in

fish larvae. *Environmental Biology of Fishes* 16, 59–77.

https://doi.org/10.1007/978-94-017-1158-6_5

Hagen, I. J., Jensen, A. J., Bolstad, G. H., Diserud, O. H., Hindar, K., Lo, H., &

Karlsson, S. (2019). Supplementary stocking selects for domesticated genotypes.

Nature Communications, 10, 199. https://doi.org/10.1038/s41467-018-08021-z

JettyBox Stock Video. (2015, Oct. 23) Salmon jumping Shrewsbury Weir [Video file].

Retrieved from https://www.youtube.com/watch?v=JjXVCZFTe70

Kaeriyama, M. (1986). Ecological study on early life of the chum salmon

Oncorhynchus keta (Walbaum). *Scientific Reports of the Hokkaido Salmon*

Hatchery, 40, 31–92.

Kaeriyama, M., Seo, H., & Qin, Y. X. (2014). Effect of global warming on the life

history and population dynamics of Japanese chum salmon. *Fisheries Science*, 80,

251–260. https://doi.org/10.1007/s12562-013-0693-7

Kawamura, S. (2008, Oct. 17) Chum salmon upstream [Video file]. Retrieved from

https://www.youtube.com/watch?v=ABD_rtZuzXo

Kijima, A., & Fujio, Y. (1979). Geographical distribution of IDH and LDH isozymes in

chum salmon population. *Nippon Suisan Gakkaishi*, 45, 287–295. [In Japanese

with English abstract] https://doi.org/10.2331/suisan.45.287

Kitada, S. (2014). Japanese chum salmon stock enhancement: current perspective and

future challenges. Fisheries Science 80, 237–249. https://doi.org/10.1007/s12562-

013-0692-8

Kitada, S. (2018). Economic, ecological and genetic impacts of marine stock

enhancement and sea ranching: A systematic review. *Fish and Fisheries*, 19, 511–

532. https://doi.org/10.1111/faf.12271

Kitada, S., Nakajima, K., Hamasaki, K., Shishidou. H., Waples, R., & Kishino, H.

(2019). Rigorous monitoring of a large-scale marine stock enhancement program

demonstrates the need for comprehensive management of fisheries and nursery

habitat. *Scientific Reports*, 9, 5290. https://doi.org/10.1038/s41598-019-39050-3

Kondo, H., Hirano, Y., Nakayama, N. & Miyake, M. (1965). Offshore distribution and

migration of Pacific salmon (genus *Oncorhynchus*) based on tagging studies

(1958-1961). *Bulletin of International North Pacific Fishery Commission,* 17, 1–213.

Laikre, L., Schwartz, M. K., Waples, R. S., Ryman, N., & GeM Working Group. (2010). Compromising genetic diversity in the wild: unmonitored large-scale release of plants and animals. Trends in Ecology & Evolution, 25, 520–529. https://doi.org/10.1016/j.tree.2010.06.013

Le Luyer, J., Laporte, M., Beacham, T. D., Kaukinen, K. H., Withler, R. E., Leong, J. S., ... & Bernatchez, L. (2017). Parallel epigenetic modifications induced by hatchery rearing in a Pacific salmon. *Proceedings of the National Academy of Sciences*, 114, 12964-12969. https://doi.org/10.1073/pnas.1711229114

McGinnity, P., Prodöhl, P., Ferguson, A., Hynes, R., ó Maoiléidigh, N., Baker, N., ... Taggart, J. (2003). Fitness reduction and potential extinction of wild populations of Atlantic salmon, *Salmo salar*, as a result of interactions with escaped farm salmon. *Proceedings of the Royal Society of London B*, 270, 2443–2450. doi: 10.1098/rspb.2003.2520

Miyakoshi, Y., Nagata, M., Kitada, S., & Kaeriyama, M. (2013). Historical and current hatchery programs and management of chum salmon in Hokkaido, northern Japan. *Reviews in Fisheries Science*, 21, 469–479. doi: org/10.1080/10641262.2013.836446

Morita, K. (2016). Trends in Russian chum salmon populations. *FRA Salmonid Research Report*, 10, 38–40 [In Japanese].

North Pacific Anadromous Fish Commission (NPAFC). *NPAFC Pacific salmonid catch statistics (updated summer 2019)*, NPAFC, Vancouver. Retrieved from http://www.npafc.org

Okazaki, T. (1982) Geographical distribution of allelic variations of enzymes in chum salmon *Oncorhynchus keta*, river populations of Japan and the effects of

transplantation. *Nippon Suisan Gakkaishi*, 48, 1525–1535.

https://doi.org/10.2331/suisan.48.1525

Onishi, Y. (2009, Sep. 25) Returning chum salmon at Hamamasu River, Hokkaido

[Video file]. Retrieved from https://www.youtube.com/watch?v=7FXF6yHoC24

Podrabsky, J. E., Javillonar, C., Hand, S. C., & Crawford, D. L. (2000). Intraspecific

variation in aerobic metabolism and glycolytic enzyme expression in heart

ventricles. *American Journal of Physiology-Regulatory, Integrative and*

Comparative Physiology, 279, R2344–R2348.

https://doi.org/10.1152/ajpregu.2000.279.6.R2344

Powers, D. A., Lauerman, T., Crawford, D., & DiMichele, L. (1991). Genetic

mechanisms for adapting to a changing environment. *Annual Review of Genetics*,

25, 629–660. https://doi.org/10.1146/annurev.ge.25.120191.003213

Reisenbichler, R. R., & McIntyre, J. D. (1977). Genetic differences in growth and

survival of juvenile hatchery and wild steelhead trout, *Salmo gairdneri. Journal of*

the Fisheries Board of Canada, 34, 123–128.

Reisenbichler, R. R., & Rubin, S. P. (1999). Genetic changes from artificial propagation

of Pacific salmon affect the productivity and viability of supplemented

populations. *ICES Journal of Marine Science*, 56, 459–466.

Ruggerone, G. T., & Irvine, J. R. (2018). Numbers and biomass of natural-and hatchery-

origin pink salmon, chum salmon, and sockeye salmon in the north Pacific Ocean,

1925–2015. *Marine and Coastal Fisheries*, 10, 152–168.

https://doi.org/10.1002/mcf2.10023

Sapporo TV House Stock Footage. (2015, June 10). Waterfall climbing of salmon

[Video file]. Retrieved from https://www.youtube.com/watch?v=QhnWlFyv7p0

Seeb, L. W., Crane, P. A. & Gates, R. B. (1995) *Progress Report of Genetic Studies of*

Pacific Rim Chum Salmon, and Preliminary Analysis of the 1993 and 1994 South

Unimak June Fisheries. Anchorage, USA: Alaska Department of Fish and Game.

Somero, G. N. (2004). Adaptation of enzymes to temperature: searching for basic

"strategies". *Comparative Biochemistry and Physiology Part B: Biochemistry and*

Molecular Biology, 139, 321–333. https://doi.org/10.1016/j.cbpc.2004.05.003

Somero, G. N. (2010). The physiology of climate change: how potentials for

acclimatization and genetic adaptation will determine 'winners' and 'losers'.

Journal of Experimental Biology, 213, 912–920. doi: 10.1242/jeb.037473

Taylor, R. (2014, July 31). Salmon swimming upstream [Video file]. Retrieved from

https://www.youtube.com/watch?v=ZjS3eSNuX_c

The Hokkaido Shimbun Press. (2017, Oct. 18). Chum salmon returning to Okushibetsu

River, Hokkaido [Video file]. Retrieved from

https://www.youtube.com/watch?v=keS0O1WLWl4

Tillotson, M. D., Barnett, H. K., Bhuthimethee, M., Koehler, M. E., & Quinn, T. P.

(2019). Artificial selection on reproductive timing in hatchery salmon drives a

phenological shift and potential maladaptation to climate change. *Evolutionary*

applications, 12, 1344–1359. https://doi.org/10.1111/eva.12730

Trut, L. N. (1999). Early Canid Domestication: The Farm-Fox Experiment: Foxes bred

for tamability in a 40-year experiment exhibit remarkable transformations that

suggest an interplay between behavioral genetics and development. *American*

Scientist, 87, 160–169. https://www.jstor.org/stable/27857815

Turtle Island Restoration Network. (2015, Dec. 22) Endangered coho salmon jump

waterfall to reach spawning grounds [Video file]. Retrieved from

https://www.youtube.com/watch?v=PCEYStqpltA

Ueno, Y. & Ishida, Y. (1996). Summer distribution and migration routes of juvenile

chum salmon (*Oncorhynchus keta*) originating from rivers in Japan. *Bulletin of*

National Research Institute of Far Seas Fisheries 33, 139–147.

WallnerPhotos. (2014, Jul. 12) Salmon jumping up waterfalls [Video file]. Retrieved

from https://www.youtube.com/watch?v=YWb8noMoM68

Whitlock, M. C. (2008). Evolutionary inference from Q_{ST}. *Molecular Ecology*, 17,

1885–1896. https://doi.org/10.1111/j.1365-294X.2008.03712.x

Winans, G. A., Aebersold, P. B., Urawa, S., & Varnavskaya, N. V. (1994). Determining

continent of origin of chum salmon (*Oncorhynchus keta*) using genetic stock

identification techniques: status of allozyme baseline in Asia. *Canadian Journal*

of Fisheries and Aquatic Sciences, 51(S1), 95–113. https://doi.org/10.1139/f94-

298

SUPPORTING INFORMATION

556   Additional supporting information may be found online in the Supporting Information

557   section at the end of the article.

TABLE 1 Exponential decay rates of fitness in the wild for captive-bred fish.

Species	Estimate	SE	t	p	Source
Chum salmon (Hokkaido) [†]	0.022	0.007	2.93	0.0082**	This study
Chum salmon (Honshu) [†]	0.040	0.011	3.69	0.0014**	This study
Red sea bream[†]	0.156	0.017	9.02	0.0000***	Kitada et al., 2019
Salmonids RRS[‡]	0.365	0.058	6.27	0.0008**	Araki et al., 2007

[†] Return rate per year. [‡] Relative reproductive success per generation in captivity.

TABLE 2 Linear regression of chum salmon catches on SST anomaly.

Season[†]	Coefficient	Estimate	*SE*	*t*-value	*p*	R^2
Average	Intercept	59.9	3.24	18.5	0.000***	0.45
	SST	-28.1	6.72	-4.2	0.000***	
Winter	Intercept	51.8	3.25	15.9	0.000***	0.07
	SST	-8.2	6.74	-1.2	0.237	
Spring	Intercept	54.4	2.93	18.6	0.000***	0.32
	SST	-15.4	4.93	-3.1	0.005*	
Summer	Intercept	56.2	3.25	17.3	0.000***	0.30
	SST	-1.0	3.32	-3.0	0.007**	
Autumn	Intercept	54.0	4.50	12.0	0.000***	0.04
	SST	-6.8	6.97	-1.0	0.339	

[†] Winter; Jan.-Mar., Spring; Apr.-June, Summer; July-Sept., Autumn; Oct.- Dec.

Figure Legends

FIGURE 1 Chum salmon in the North Pacific. (a) Changes in the total catch of chum salmon in the North Pacific and the proportion of Japan. Data from North Pacific Anadromous Fish Commission, www.npafc.org accessed August 2019. (b) Return rates in Hokkaido and Honshu. Dotted lines show exponential curves fitted to return rates (1996-2017). (c) Mean SST anomaly for spring and summer in Hokkaido and the northern Honshu vs. number of adult fish returning to Japan, with estimated regression lines. Numbers on the graph indicate years. [The colour figure can be viewed at wileyonlinelibrary.com]

FIGURE 2 Changes in number of chum salmon returned to Japan in recent two decades. (a) 1996 with the historical maximum catch. (b) 2006. (c) 2014. (d) 2017 (latest statistics). The numbers show chum salmon returned in thousand. The sizes of the circle are proportional to the numbers. Green dots show hatcheries (Figure S2).

FIGURE 3 Changes in number of chum salmon released in Japan in recent two decades. (a) 1992. (b) 2002. (c) 2010. (d) 2013. The number show chum salmon released in million. The sizes of the circles are proportional to the numbers. The years were four years before ones for catches in Figure 2. Green dots show hatcheries (Figure S2).

FIGURE 4 Locations for chum salmon allozyme gene frequencies across the Pacific Rim (*n*=166). (a) Alaska, British Columbia and Washington. (b) Alaska Peninsula. (c) Russia. (d) Japan. Numbers and names of the locations and gene frequencies were given in Supplementary Data. Data from the original study (Kijima, & Fujio, 1979; Okazaki, 1982; Winans, Aebersold, Urawa, & Varnavskaya,1994; Seeb, Crane, & Gates, 1995).

FIGURE 5 Geographical distribution of chum salmon allozyme gene frequencies in the North Pacific. (a), (b) LDH-A1. (c), (d) LDH-B2. (e), (f) PEPB-1. Colours and numbers in panels (a), (c) and (e) refer to sampling locations in Figure 3. Colours for sampling locations in panels (b), (d) and (f) visualize the magnitude of gene frequencies. The dotted circles show 95% confidence ellipses.