Consequences of hatchery stocking:

Lessons from Japan

Sort title: Japan hatchery stocking

Shuichi Kitada

[1]Tokyo University of Marine Science and Technology, Tokyo 108-8477, Japan.

E-mail: kitada@kaiyodai.ac.jp

Abstract

More than 26 billion juveniles of 180 marine species are released annually into the wild in

over 20 countries, but the usefulness remains unclear. Here, I review the effects of stocking by

Japanese marine and salmon stock-enhancement programmes, and evaluate the efficacy

through a meta-analysis, with new cases added to those previously considered. The posterior

mean recapture rates (±SD) was $8.2 \pm 4.7\%$; mean economic efficiency was 2.7 ± 4.5, with

many cases having values of 1–2, excluding consideration for personnel costs and negative

impacts on wild populations. At the macro-scale, the proportion of released seeds to total

catch was $76 \pm 20\%$ for Japanese scallop, $28 \pm 10\%$ for abalone, $20 \pm 5\%$ for swimming crab,

$13 \pm 5\%$ for kuruma prawn, $11 \pm 4\%$ for Japanese flounder and $7 \pm 2\%$ for red sea bream.

Stocking effects were generally small and population dynamics unaffected by releases but

dependent on carrying capacity of the nursery habitat. Japanese scallop in Hokkaido was the

most successful case with the highest economic efficiency (18 ± 5), and populations

maintained by releases of wild-born spat. In contrast, captive breeding reduced the fitness of

released red sea bream in the wild, while wild fish recovered due to increased seaweed

community and reduced fishing efforts. All most chum salmon returning to Japan are hatchery

fish and/or hatchery descendants, which may reduce the fitness of populations in the warming

sea temperature. Nursery-habitat recovery and fishing-effort reduction outperform hatcheries

in the long-run.

Keywords: Bayesian meta-analysis, genetic effects, nursery habitat, stocking effects

1 | Introduction

The history of stocking fish larvae dates back to the 1870s (Bell et al., 2005; Blaxter, 2000; Svåsand et al., 2000). Approximately 150 years later, seed production technology has progressed and releases of hatchery-reared animals into the wild have become a popular tool in fisheries, forestry, and wildlife management (Laikre, Schwartz, Waples, & Ryman, 2010). More than 26 billion juveniles of 180 marine species, including salmonids, are released into the wild every year (Kitada, 2018). Moreover, marine ranching is rapidly growing in China (Lee & Zhang, 2018; Wang, Yu, & Chen, 2018; Zhou, Zhao, Zhang, & Lin, 2019) and Korea (Lee & Rahman, 2018). Despite the huge number of seeds released every year, most studies have been conducted at the experimental stage, particularly for marine species, and empirical studies are still too few for determining the full effectiveness of hatchery-release efforts. The feasibility and risks of hatchery stocking cannot be tested by examining pilot-scale releases alone, but can be sufficiently tested only by considering full-scale releases (Hilborn, 2004). A global analysis of full-scale releases found that most cases were economically unprofitable as a consequence of the high costs of seed production when compared with the prevailing market prices, apart from a few successful cases or cases left unevaluated (Kitada, 2018). Further investigation is therefore needed to answer the fundamental question of whether hatchery stocking is useful or harmful (Araki & Schmid, 2010).

Japan releases the largest number of hatchery-reared species in the world and often on a large scale. Focused, in-depth analyses of full-scale releases can improve our understanding of the positive and negative effects of this practice, and would benefit future fisheries management and conservation practices. In a recent review, I summarised the status of the economic, ecological and genetic effects of marine stock enhancement and sea-ranching programmes (Kitada, 2018), but the focus was on a global scale. Several other reviews described Japanese

marine stock enhancement and sea-ranching programmes by focusing on the effects of

stocking several fishes (Kitada, 1999; Kitada & Kishino, 2006; Masuda & Tsukamoto, 1998),

kuruma prawn (Hamasaki & Kitada, 2006, 2013), abalone (Hamasaki & Kitada, 2008a), and

decapod crustaceans (Hamasaki & Kitada, 2008b; Hamasaki, Obata, Dan, & Kitada, 2011).

Attention has also been given to seed production (Fushimi, 2001; Le Vay et al., 2007;

Takeuchi, 2001), seed quality and behaviour (Tsukamoto, Kuwada, Uchida, Masuda,

Sakakura, 1999), broodstock management (Taniguchi, 2003, 2004) and the genetic effects on

wild populations (Kitada et al., 2009). For salmonids, many reviews have covered large-scale

hatchery releases (e.g., Kaeriyama, 1999; Kaeriyama, Seo, Kudo, & Nagata, 2012; Kitada,

2014; Miyakoshi, Nagata, Kitada, & Kaeriyama, 2013; Morita, 2014; Morita et al., 2006;

Nagata et al. 2012). However, as yet, there has been no integrated review that summarises

results concerning both marine and salmon stock enhancement in Japan.

Here, I first outline the history and present status of Japanese salmon and marine hatchery

programmes. Second, I evaluate the efficacy of major full-scale projects using a Bayesian

meta-analysis, with new cases added since my previous study. Third, I predict changes in the

contribution of hatchery releases to commercial catches for representative iconic species on a

macro-scale. Fourth, I summarise consequences concerning the world's largest hatchery

stocking programmes—namely those for red sea bream, Japanese scallop and chum salmon.

2 | Hatcheries, seed release, and stocking-effect surveys

Japan's marine stock-enhancement programme was initiated by the Fishery Agency in 1963.

At that time, Japan aimed to become a fully developed country under its national

industrialisation policy. Many Japanese coastlines to serve as nursery habitats for marine

species were reclaimed (Figure 1), and separate coastal industrial zones were created; many

coastal fishers moved to cities to earn higher incomes and coastal fisheries were weak. The

marine stock-enhancement programme was a mitigation policy to improve degraded habitats

and thereby enhance the coastal fisheries. Salmon hatcheries in Japan have a much longer

history, beginning in the 1880s to increase fishery production of returning chum salmon, and

have endured for over 130 years. However, chum salmon stock enhancement did not succeed

during the early 1960s as the level of spawning biomass remained as it was even before the

dramatic increase of returned fish as explained below. Therefore, Japan's marine stock-

enhancement programme was seemingly started with no successful case precedent. This

situation implies that "many believe that the future of fisheries lies with artificial propagation

to rebuild depleted fish stocks due to poor fisheries management or poor habitat management"

(Hilborn, 1992).

First, I created maps of salmon and marine hatcheries; approximate locations of salmon

hatcheries were plotted on a map of Japan using 15 regional maps of the locations of salmon

hatcheries, obtained from the website of the Japan Fisheries Research and Education Agency

(FRA, salmon.fra.affrc.go.jp). I identified 262 salmon hatcheries in 11 prefectures in northern

Japan, of which 242 were private hatcheries, managed mainly by fishers and cooperatives

(Figure 2a). The 7 prefectural and 13 national hatcheries are operated mainly for research

work. The major target salmonid species are chum salmon (*Oncorhynchus keta*, Salmonidae)

and pink salmon (*O. gorbuscha*, Salmonidae). According to statistics from the North Pacific

Anadromous Fish Commission (NPAFC), the number of released chum salmon has

remarkably increased since the 1970s, reaching a historical maximum of 2.1 billion per

annum in 1991. But since then releases of chum salmon dropped to 1.5 billion in 2018

(NPAFC, 2019) (Figure S1). A similar increasing trend since the 1970s occurred with pink

salmon; however, the number of released pink salmon was 113 million in 2018, about 7% that

of chum salmon. The number of masu salmon (*O. masou*, Salmonidae) released in 2018

amounted to 7 million. Sockeye salmon (*O. nerka*, Salmonidae) are also released, but at even

smaller numbers (NPAFC, 2019).

The locations of "saibai-gyogyo" centres in Japan (hereafter, marine hatcheries) were also

plotted, using a list of their addresses compiled by the National Association for Promotion of

Productive Seas (NAPPS). Presently, 65 prefectural marine hatcheries are operated in 43

prefectures (Figure 2b). In addition, 12 FRA research stations undertake marine stock

enhancement. These were once national marine hatcheries managed by the Japan Sea-

Farming Association (JASFA), which merged with the FRA following administrative reform

in 2003. The JASFA was disbanded after 40 years, and the promotion of hatchery releases was

inherited by the NAPPS, established in 2003. Since the FRA aimed to reduce their

involvement in hatchery releases, four national marine hatcheries were closed up to 2018.

By 2017, a total of 74 species were being released in Japan, which included 34 fishes, 10

crustaceans, 22 shellfishes and 8 other marine species (e.g., sea urchin, sea cucumber and

octopus). I organized the numbers of released seeds for the 16 major species released, whose

annual release exceeded million seeds, for the period 1983–2017, relying on the annual

statistics of seed production and releases (Fishery Agency, FRA, & NAPPS, 1985–2019).

These species are: abalone (*Haliotis* spp., Haliotidae), Japanese scallop (*Mizuhopecten*

yessoensis, Pectinidae), Manila clam (*Venerupis philippinarum*, Veneridae), green tiger prawn

(*Penaeus semisulcatus*, Penaeidae), kuruma prawn (*Marsupenaeus japonicus*, Penaeidae),

offshore greasyback prawn (*Metapenaeus ensis*, Penaeidae), swimming crab (*Portunus*

trituberculatus, Portunidae), black sea bream (*Acanthopagrus schlegeliir*, Sparidae), flatfish

(e.g. *Limanda yokohamae*, Pseuronectidae), Japanese flounder (*Paralichthys olivaceus*,

Paralichthyidae), Pacific herring (*Clupea pallasii*, Clupeidae), red sea bream (*Pagrus major*,

Sparidae), sailfin sandfish (*Arctoscopus japonicus*, Trichodontidae), tiger puffer (*Takifugu*

rubripes), sea urchins (e.g. *Strongylocentrotus intermedius*, Strongylocentroidae; *Heliocidaris*

crassispina, Echinometridae) and sea cucumber (*Apostichopus armata*, Stichopodidae). The

number of releases has increased and/or been stable for 4 species: Japanese scallop, tiger

puffer, Pacific herring and sea cucumber. Releases have been decreasing for the other 12

species, including iconic target species such as the kuruma prawn, swimming crab, abalone,

red sea bream, Japanese flounder and sea urchin (Figures S2, S3; Supplementary Data).

In earlier surveys, counting external tags reported by fishers was the main method used to

estimate recapture rates (Kitada, 1999). This type of survey was used to estimate migration,

growth and life history, including comparisons of seed quality (Kitada, Hiramatsu, & Kishino,

1994; Kitada & Hirano, 1987; Okouchi, Kitada, Morioka, Imamura, 1994; Shiota, & Kitada,

1992; Takaba, Kitada, Nakano, Morioka, 1995). However, tag-reporting rates were often very

small, and tag-shedding and tagging mortality caused biased estimates. Researchers became

aware of bias in the results when they estimated stocking effectiveness. As an alternative

approach, surveys of commercial landings at fish markets (SCFM) were applied to estimate

landings of released seeds and the hatchery contribution in the landings. For the SCFM,

researchers sampled landings by visiting fish markets to cover the range of the stocking effect.

The SCFM approach was first applied to red sea bream in Kagoshima Bay, where almost all

red sea bream landed at the Kagoshima Fish Market were checked for deformity of the

internostril epidermis to identify released fish (Kitada et al., 2019; Shishidou, 2002; Shishidou

& Kitada, 2007). In the case of Japanese flounder in Miyako Bay, all flounder at the Miyako

fish market were examined (Okouchi, Kitada, Iwamoto, & Fukunaga, 2004; Okouchi, Kitada,

Tsuzaki, Fukunaga, & Iwamoto, 1999). Such a census approach was generally difficult to

apply, therefore sample surveys were logically introduced throughout the country. The

common procedure was that all fish landed and sold at a selected market on a selected day

would be checked for released seeds versus wild individuals. This was regarded as a two-

stage sampling procedure for the sake of formulating estimators and their variance (Kitada,

Taga, & Kishino, 1992). Accordingly, this approach has been applied to abalone (Kojima,

1995), kuruma prawn (Yamaguchi, Ito, Hamasaki, & Kitada, 2006) and masu salmon

(Miyakoshi, Nagata, Sugiwaka, & Kitada, 2001). When information on the total landing days

was lacking, simple random sampling was assumed, and landings from releases were then

estimated (Obata, Yamazaki, Iwamoto, Hamasaki, & Kitada, 2008). Some other studies using

SCFM data chose to assume simple random sampling, but in most cases the precision was not

evaluated. As another approach, genetic marking was applied to mud crab to estimate

recapture rates using genetic stock identification (Obata, Imai, Kitakado, Hamasaki, & Kitada,

2006).

To enhance the effectiveness of hatchery releases, a marine-ranching project, led by the

Fishery Agency, endeavoured to shape 'natural' farms with sound habitat, such as enhanced

seaweed communities to promote nutrient-enrichment in deep-sea upwelling systems

(AFFRC, 1989). Hatchery releases and artificial reef construction were the major

technologies, and feeding systems with acoustic conditioning were the popular tool, applied

for marine fishes that could be reared and released as seeds around artificial reefs (Kayano,

Tanaka, & Hayashi, 1998; Kudo & Kimoto, 1994). However, I could not find scientific

literature that evaluated the effectiveness of Japan's marine-ranching projects, thus its

usefulness remains largely unknown, and consequently the effects of marine ranching in

Japan were excluded from my analyses.

3 | Meta-analysis of hatchery performance

3.1 | Methods

I used recapture rates, yield-per-release (YPR), and economic efficiency as indices of the performance of hatchery releases, following a preceding study. YPR is defined as weight of fish caught (g) per individual released (Hamasaki & Kitada, 2008b; Kitada & Kishino, 2006; Svåsand et al., 2000), as follows:

$$YPR = r \times w$$

where r is the recapture rate of released juveniles, and w (g) is mean body weight per recaptured individual. Economic efficiency (E) is calculated as the ratio of net income to the release costs:

$$E = YPR \times v/c$$

Here, v is fish price per gram, and c is the cost of each seed; therefore, v/c is the cost performance of a seed (Kitada, 2018).

The true mean for the recapture rate, YPR, and economic efficiency for each study (y_i) is not observed but estimated ($\hat{y}_i, i = 1, \dots, n$), and $\hat{y}_i$ may follow a normal distribution by the central limit theorem. When samples (i.e., cases within a study) are taken by random sampling, then $E(\hat{y}_i) = y_i$, and the variance within each study (σ_i^2) is also estimated from the data (s_i^2). However, the release variables, such as tags, sizes at release, release areas, and time periods of surveys, varied in each study; therefore, the summary statistics needed to take into account the variation among studies. Here, I assumed a superpopulation for the recapture studies, by applying a Bayesian approach similar as given in a previous study (Kitada, Kishino, & Hamasaki, 2011).

Studies are carried out in different areas and years, therefore each study can be regarded as

206 independent. The approximate likelihood of the sample mean $\hat{y}_i \ (i = 1, ..., n)$ can be written

207 as:

208
$$L(\hat{y}_i \ (i = 1, ..., n) | y_i \ (i = 1, ..., n)) = \prod_{i=1}^{n} \frac{1}{\sqrt{2\pi\sigma_i^2}} \exp\left(-\frac{1}{2\sigma_i^2}(\hat{y}_i - y_i)^2\right)$$

209 We assume a prior distribution for $y_i \ (i = 1, ..., n)$. Here, y_i is assumed to be normally

210 distributed around the mean μ with the variance σ^2. These are hyperparameters: namely, the

211 mean and variance of the superpopulation. The prior distribution of $y_i \ (i = 1, ..., n)$ can be

212 written as:

213
$$\pi(y_1, ..., y_n | \mu, \sigma^2) = \prod_{i=1}^{n} \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{1}{2\sigma^2}(y_i - \mu)^2\right)$$

214 By integrating the product of the prior distribution and the likelihood function in terms of y_i,

215 the marginal likelihood function is derived. In this case, it is explicitly obtained as:

216
$$\tilde{L}(\sigma^2, \mu \ | \hat{y}_i(1, ..., n)) = \prod_{i=1}^{n} \frac{1}{\sqrt{2\pi(\sigma^2 + \sigma_i^2)}} \exp\left(-\frac{1}{2(\sigma^2 + \sigma_i^2)}(\hat{y}_i - \mu)^2\right)$$

217 This shows that the sample mean $\hat{y}_i$ is distributed around the mean μ, with variances

218 calculated for between cases and within cases.

220 The log marginal likelihood function is:

221
$$\log\tilde{L}(\sigma^2, \mu \ | \hat{y}_i(1, ..., n)) = -\frac{n}{2}\log 2\pi - \frac{1}{2}\sum_{i=1}^{n}\log(\sigma^2 + \sigma_i^2) - \frac{1}{2}\sum_{i=1}^{n}\frac{1}{\sigma^2 + \sigma_i^2}(\hat{y}_i - \mu)^2$$

222 The first derivatives on μ and σ^2 are then:

223
$$\frac{\partial\log\tilde{L}}{\partial\mu} = \sum_{i=1}^{n}\frac{1}{\sigma^2 + \sigma_i^2}(\hat{y}_i - \mu)$$

224
$$\frac{\partial\log\tilde{L}}{\partial\sigma^2} = -\frac{1}{2}\sum_{i=1}^{n}\frac{1}{\sigma^2 + \sigma_i^2} + \frac{1}{2}\sum_{i=1}^{n}\frac{1}{(\sigma^2 + \sigma_i^2)^2}(\hat{y}_i - \mu)^2$$

225 From $\partial \log \tilde{L} / \partial \mu = 0$, we have the maximum likelihood estimator (MLE) of μ as a weighted

226 average of $\hat{y}_i$ ($i = 1, ..., n$):

$$\hat{\mu} = \sum_{i=1}^{n} \frac{\hat{y}_i}{\sigma^2 + \sigma_i^2} \Big/ \sum_{i=1}^{n} \frac{1}{\sigma^2 + \sigma_i^2} \qquad (1)$$

228 By substituting σ^2 with $\hat{\sigma}^2$ given in Eq. (2) and σ_i^2 with the sample variance s_i^2, we have

229 the MLE of μ (posterior mean).

231 Assuming that σ_i^2 does not depend on a case for simplicity, $\sigma_i^2 = \sigma'^2$ for all i ($1, ..., n$)

232 and from $\partial \log \tilde{L} / \partial \sigma^2 = 0$, we have the MLE of σ^2 as:

$$\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^{n} (\hat{y}_i - \mu)^2 - \sigma'^2$$

234 By estimating μ by the weighted average $\hat{\mu}$ given by Eq. (1) and σ'^2 by the mean of the

235 sample variance s_i^2, we have the approximate variance estimator as:

$$\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^{n} (\hat{y}_i - \hat{\mu})^2 - \frac{1}{n} \sum_{i=1}^{n} s_i^2 \qquad (2)$$

The MLE of μ has a different weight from the weighted mean (also the MLE) that was used

in the previous study (Kitada, 2018), while $\hat{\sigma}^2$ is the same. Eq. (1) provided a more realistic

estimate than use of the previous weighted average, particularly when very small $\hat{y}_i$ values

($\hat{y}_i < 1$) were included (i.e., when there was large variation in $\hat{y}_i$). The average and standard

deviation (SD) with a 95% confidence interval (mean $\pm$ 2 SD) were visualised for each case

using the 'forestplot' function in R. When the lower confidence limit took a negative value, it

was replaced by 0.

3.2 | Recapture rate

I summarised 21 full-scale programmes (Table S1) that reported the recapture rate of hatchery

individuals, of which 18 programmes were revisited from the previous study (Kitada, 2018).

The mean recapture rate of chum salmon for the whole of Japan was re-calculated based on

the simple return rates (number of fish returned after four years/number released) taken from

the FRA website (salmon.fra.affrc.go.jp). The recapture rates of 17 of 21 studies were

estimated using the SCFM data and/or combining use of the SCFM approach and reported

recaptures. Recapture rates for red sea bream (Kitada & Kishino, 2006), kuruma prawn

(Hamasaki & Kitada, 2006, 2008b) and abalone (Hamasaki & Kitada, 2008a) were

summarised from previous reviews for the whole country, which included black rockfish

(*Sebastes schlegelii*, Sebastidae) (Nakagawa, Okouchi, & Adachi, 2004), Japanese flounder

(Atsuchi, & Masuda, 2004; Ishino, 1999; Kitada, Taga, & Kishino, 1992; Tominaga &

Watanabe, 1998; Tomiyama, Watanabe, & Fujita, 2008), Japanese scallop (Kitada, &

Fujishima, 1997), Japanese Spanish mackerel (*Scomberomorus niphonius*, Scombridae)

(Obata, Yamazaki, Iwamoto, Hamasaki, & Kitada, 2008), mud crab (*Scylla paramamosain*,

Portunidae) (Obata, Imai, Kitakado, Hamasaki, & Kitada, 2006), short-spined sea urchin

(*Strongylocent-rotus intermedius*, Strongylocentroidae) (Sakai, Tajima, & Agatsuma, 2004),

spotted halibut (*Verasper variegatus*, Pleuronectidae) (Wada, Kamiyama, Shimamura,

Mizuno, & Nemoto, 2012), swimming crab (Okamoto, 2004) and tiger puffer (Nakajima, Kai,

Koizumi, Tanaka, & Machida, 2008). I newly added four cases of three species: barfin

flounder (*Verasper moseri*, Pleuronectidae) in Hokkaido (Murakami, 2012; Koya, 2005;

NPJSEC, 2015), red spotted grouper (*Epinephelus akaara*, Serranidae) in Osaka Bay

(Tsujimura, 2007) and tiger puffer in the Ariake Sea (Matsumura, 2005, 2006), because the

stocking effect had been recently reported for the first two species, and because the number of

tiger puffer released has increased in recent years (Figure S3). Particularly for the barfin

flounder in Hokkaido, more than one million juveniles have been released annually along the

Pacific coast since 2006.

The results for recapture rates included 14 species, consisting of eight marine fishes, one

salmonid, two crustaceans, two shellfishes, and one sea urchin (Table S1). Cases that reported

only point estimates were excluded from the following analysis. The recapture rates of 20

marine species largely varied among species and cases, and ranged from 0.9% to 34.5%

(Figure 3). The posterior mean recapture rates was 8.2 ± 4.7%, and the empirical distribution

indicated that most cases had recapture rates between 6% and 12%. Japanese scallop had the

highest recapture rate (34.5 ± 10.2%), followed by sea urchin (18.2 ± 17.5%), with relatively

large variations. Most of the marine fishes, particularly barfin flounder, black rockfish,

Japanese flounder, Japanese Spanish mackerel, spotted halibut and tiger puffer (to age 0+),

had recapture rates in the range 11–15%. Abalone likewise had a relatively high recapture rate

(12.2 ± 8.1%), also with high variation. Red sea bream had recapture rates of 7–8%, with

small variation. In contrast, the return rate of chum salmon, on Hokkaido and Honshu islands

combined, was much smaller, at 2.7 ± 0.8%, which was close to that of red spotted grouper, at

2.2 ± 1.0%. Crustaceans generally had values smaller than 5%; large variations in the

recapture rates occurred for kuruma prawn (2.8 ± 4.5%) and mud crab (0.9 ± 0.7%).

3.3 | YPR and economic efficiency

I revisited cases otherwise reported on in the previous study (Kitada, 2018), of which the

economic efficiency values of 10 cases were cited, and six cases were newly added: those of

barfin flounder in Hokkaido, Japanese flounder at Fukushima, red spotted grouper in Osaka

Bay, tiger puffer in the Ariake Sea, and swimming crab in Lake Hamana and in the Seto

Inland Sea (see references in Table S2). YPR and economic efficiency values were calculated

for kuruma prawan (Hamasaki & Kitada, 2006, 2008b), swimming crab (Hamasaki et al.,

2011) and abalone (Hamasaki & Kitada, 2008a), based on the previous reviews. For chum

salmon, YPR was recalculated based on newly estimated mean body weights from catch

statistics for 1974–2017 obtained from the NPAFC, and return rates for 1974–2017 from the

FRA (http://salmon.fra.affrc.go.jp/zousyoku/sakemasu.html). The revised YPR for chum

salmon was 86 ± 29 g. For Japanese scallop, mean body weights were recalculated based on

the work of Kurata (1999), and the revised estimate was 177 ± 30 g. YPR values were

reported for pink salmon in Hokkaido in terms of amount of money (1.5–2.2 yen/released

individual), calculated based on the estimated proportion of Hokkaido-origin hatchery fish in

the landings (Ohnuki, Morita, Tokuda, Okamoto, & Ohkuma, 2015). Thus, that case did not

give the YPR in weight, the recapture rates, nor cost performance of a seed (v/c); therefore, I

excluded the case of pink salmon in Hokkaido from the analysis.

Ultimately, YPR and economic efficiency were evaluated for 16 cases involving 12 species

(Table S2). Meta-analysis was performed on 12 cases involving 9 species for YPR, and 13

cases of 10 species for economic efficiency, where the cases with only point estimates of YPR

and/or those without information for calculating v and c were omitted from the analyses. The

YPR values varied between species and cases. The empirical distribution of the summary

statistics had a long tail to the right, with a posterior mean of 63 ± 71 g (Figure 4a). YPR was

high for both barfin flounder (182 ± 9 g) and Japanese Spanish mackerel (170 ± 8 g),

followed by that for chum salmon (86 ± 29 g) and Japanese scallop (61 ± 18 g). Red sea

bream and Japanese flounder had YPR values of 30–59 g. Other YPR values were 34 ± 10 g

for swimming crab, 26 ± 19 g for abalone, 4 ± 3 g for mud crab, and 0.9 ± 1.5 g for kuruma

prawn.

The posterior mean of economic efficiency was 2.7 ± 4.5, while that of several cases was

about 1 (the break-even point) to 2, with the lower 95% confidence limit below 0 (Figure 4b). Economic efficiency was the highest for Japanese scallop (18 ± 5), followed by chum salmon in Hokkaido and Honshu combined (7 ± 2). Red sea bream in Kagoshima Bay also had a high value (5 ± 3). The consequences of the top three species were further explained in section 5. Similar economic performance was seen for abalone in all of coastal Japan (4 ± 2) and for barfin flounder on the Pacific coast of Hokkaido (3 ± 0.1). Japanese flounder had a much smaller economic efficiency, at 0.9–1.6. Crustaceans consistently exhibited economic efficiency values around 1, and kuruma prawn showed the lowest value, at 0.7 ± 0.9. Among these species, Japanese scallop, chum salmon and barfin flounder had a lower 95% confidence limit that was >1.0. Economic efficiency is a function of YPR (recapture rate × body weight) and cost performance of a seed (v/c). A scatter plot of YPR versus economic efficiency depicted the stocking-performance characteristic of each species (Figure 4c).

It is noted that the seed cost used in the analysis did not include personnel expenses, facilities, monitoring or administration costs. I did not analyse net present value (NPV) (Moksness & Støle, 1997; Moksness, Støle, & van der Meeren, 1998; Sproul &Tominaga, 1992; Svåsand et al., 2000) because various data, such as annual costs for harvest, management and interest rates, were not available for every case. If the NPV method were used, then the economic efficiency estimates obtained here would be less (Kitada, 2018), depending on the interest rate and the time duration, although recent interest rates in Japan are low (<0.1%). The seed cost also did not account for the cost of negative effects on natural populations and ecosystems (Amoroso, Tillotson, & Hilborn, 2017; Hilborn, 1998; Waples, 1991, 1999; Winton & Hilborn, 1994). Therefore, the estimates of economic efficiency obtained here are optimistic.

4 | Contribution of released seeds in the commercial landings

I estimated the contribution of hatchery releases in the commercial catch for eight species in

the whole of Japan, based on national catch statistics (MAFF, 1964–2018). These are iconic

species of Japan's marine stock enhancement and sea-ranching programmes and ones that are

intensively released within range of Japanese waters. Among these eight species, catches

increased after releases in the cases of Japanese scallop and Japanese Spanish mackerel

(Figure 5), and catches were stable for Japanese flounder and red sea bream. In contrast,

kuruma prawn and swimming crab have shown continuous decreases in catches since the

mid-1980s. Continuous decline in abalone and sea urchin catches was also observed since the

early 1970s. For cases with an increasing catch, management agencies and hatchery advocates

often say that the practise of hatchery release should be successful. For cases displaying a

stable catch, a poplar explanation is that hatchery release might support the resource. For

instances of decreasing catch, one explanation is that this is because of decreased numbers of

releases and that the catch could increase if more seeds were released. To answer this debate,

estimates of the contribution of hatchery-released seeds in the catches could be helpful.

To evaluate the hatchery contribution for the major species in the whole of Japan, I computed

the catch expected from released seeds by multiplying the average YPR (listed in Table S2) by

the numbers of fish released for each year (Supplementary Data). Except for Japanese Spanish

mackerel the contribution was calculated for the Seto Inland Sea alone because hatchery

releases in Japan are only made at that location. This simple analysis assumed that YPR was

constant over years and that released seeds created the catch in the same year, and thus it did

not account for a time-lag. However, it allowed comparisons of the approximate contribution

of hatchery releases between species in a macro-scale (Kitada & Kishino, 2006).

371 The proportion of released seeds, at 76 ± 20%, was largest for Japanese scallop, with released

372 spat contributing stably (Figure 5, vertical bars; Figure S4) and with a wild catch created by

373 natural reproduction among released scallops. The catch decreased substantially during 2015–

374 2017, following bottom disturbance off the Okhotsk coast caused by a low-pressure bomb in

375 winter of 2014, when the estimates showed that almost all of the catch comprised released

376 spat, indicating that scallop populations were heavily damaged by the bottom disturbance in

377 2014. Although the analysis was simple, the results indicated the approach could describe the

378 population dynamics of Japanese scallop.

380 The hatchery contribution to the increased catch of Japanese Spanish mackerel was small, at 2

381 ± 2%. The catch of this fish continued to recover after releases, however the number of

382 released juveniles was reduced 10 years after the release project started. The results clearly

383 indicated that the species' population dynamics were unaffected by the releases. A previous

384 study found 35% variation in the biomass of age-0 Japanese Spanish mackerel, which could

385 be explained by the biomass of a prey fish, Japanese anchovy (Nakajima et al., 2013). Genetic

386 stock identification following releases in 2001 and 2002 found that the mixing proportion of

387 hatchery-origin fish was high, at 8–15% (Nakajima et al., 2014). Interestingly, the genetic

388 admixture of hatchery fish was much higher than the estimated contribution rates in the Seto

389 Inland Sea, as found for red sea bream in Kagoshima Bay (Kitada et al., 2019), again

390 implying a trans-generation genetic effect. The hatchery contribution for Japanese flounder

391 was 11 ± 4%, and that for red sea bream was 7 ± 2%. Over two decades, the number of

392 released juveniles was decreased by ~50% for flounder and by ~63% for red sea bream, but

393 the catches of wild fish remained stable, indicating that the hatchery releases did not boost the

394 population size of either species on a macro-scale. Natural reproduction should have

395 supported the recruitment in the carrying capacity.

For crustaceans, the hatchery contribution was $13 \pm 5\%$ for kuruma prawn. The results agreed

with the estimates of previous research (Hamasaki & Kitada, 2013), ~10% for kuruma prawn

in all of Japan for the years 1977 to 2008, wherein it was concluded that the decline in

kuruma prawn catches might be an outcome of warming ocean conditions, decreased fishing

effort because of a fewer number of fishers, and reduced hatchery releases. A continuous

decreasing trend in the catch implied a decline in natural recruitments, suggesting an

environmental effect (Figure 5). Reduced fishing efforts would likely work to increase the

population size given a sound nursery environment, as demonstrated in the case of red sea

bream (Kitada et al., 2019); therefore, a reduction in fishing effort might be excluded as a

cause of the decreasing catches of kuruma prawn. An environmental effect might be

substantial for kuruma prawn, particularly extensive loss of tidal nursery habitat (Hamasaki &

Kitada, 2006). For swimming crab, the hatchery contribution was $20 \pm 5\%$; the catch

continuously decreased, again revealing a decline in natural recruitment. Juvenile crabs are

found on tidal flats after the C4 stage (~16 mm carapace width [CW]) and/or C5 stage (~22

mm CW) (Hamasaki, Obata, Dan, & Kitada, 2011); thus, extensive loss of tidal nursery

habitat would be a logical cause of a decreasing catch. The catches of kuruma prawn and

swimming crab were positively correlated ($r = 0.52,\ t = 4.49, \mathrm{df} = 53, p = 3.9 \times 10^{-5}$),

suggesting losses of nursery habitat commonly affect these two crustaceans.

Abalone displayed a relatively high and stable hatchery contribution rate ($28 \pm 10\%$), yet the

catch continued to decrease, indicating decreases in natural recruitment. A negative

correlation was found between the Aleutian Low-Pressure Index, winter sea surface

temperatures, and catches of Ezo abalone (*Haliotis discus hannai*, Haliotidae) in northern

Japan (Hayakawa, Yamakawa, & Aoki, 2007; Nakamura, Kitada, Hamasaki, & Okouchi,

2005), plus cold winter seawater temperatures (<5 °C) affected the survival of young Ezo

abalone (Takami et al., 2008). Seaweed community richness (carrying capacity) was the key

factor for successful abalone stocking in Japan (Hamasaki, 2008). For sea urchin, I could not

calculate the hatchery contribution rates because no published information available for

calculating YPR values was found. However, the decreasing trend in the releases and catch of

sea urchin was very similar to that of abalone ($r = 0.96, t = 24.63, \mathrm{df} = 53, p =$

2.2×10^{-16}), indicating that sea urchin abundance likewise heavily depends on richness of

the seaweed community.

5 | Consequences of the world's largest programmes

5.1 | Red sea bream in Kagoshima Bay

Red sea bream in Kagoshima Bay is one of the world's largest hatchery-release programmes

for a marine fish, and it is the best-monitored one in Japan; it was started in 1974 and ~27

million hatchery-reared red sea bream (6–7 cm TL) have been released in the bay. During the

period 1989–2015, 1.6 million red sea bream were examined at fish markets to identify

hatchery fish caught in the bay. The catch of hatchery fish reached 126 tonnes by 1991, but

thereafter consistently decreased, dropping to 3 tonnes by 2016. This was due to a decline in

fitness of the hatchery-reared fish in the wild, caused by the repeated use of parent fish reared

in captivity (captive breeding) since 1974 (~9 generations). In contrast, the catch of wild fish

increased after 1991, and reached a maximum in 2016 following releases amounting to 168

tonnes. Increased seaweed communities and reduced fishing efforts were the primary factors

in recovery of Japan's wild population of red sea bream (Kitada et al., 2019).

Importantly, the example of red sea bream in Kagoshima Bay shows that the hatchery releases

substantially increased the fisheries production for the first 15 years—when this came to be

regarded as a representative, successful programme in Japan—but then the catch of hatchery

fish steadily decreased and remained very low. Ample evidence for salmonids shows that

captive breeding can reduce the fitness of hatchery salmon in the wild (Araki, Cooper, &

Blouin, 2007; Fleming et al., 2000; McGinnity et al., 2003; Reisenbichler & McIntyre, 1977;

Christie, Ford, & Blouin, 2014). The declining catch of red sea bream in Kagoshima Bay was

attributed to genetic effects through unintended domestication/selection by captive breeding

(Araki, Berejikian, Ford, & Blouin, 2008; Araki, Cooper, & Blouin, 2007; Ford, 2002).

Cumulative recapture rates until age 8+ decreased $15.6 \pm 1.7\%$ (standard error, SE) per year.

Since 1974, the broodstock of red sea bream intended for release in Kagoshima Bay have

been reared in a 100-m^3 concrete tank; about 130 non-local red seab ream broodstock have

been maintained in the concrete tank for natural spawning and were repeatedly used as

broodstock. In 1999 and 2014, a total of 395 wild, 33 farmed, and many 2-year-old hatchery

fish produced from the broodstock were added to the broodstock. The number of parent fish

used for seed production varies, between about 45 and 187, every year. The rate of fitness

reduction in hatchery-reared populations was cohort-specific and was constant over time

within the cohort, but it exponentially decreased with the duration in captivity (Kitada et al.,

2019). This suggests that, once a fitness decline arises in a hatchery population, the reduction

in fitness may continue until complete replacement of the broodstock.

Backcrossing to wild populations could work to diminish the genetic effects of captive

breeding if they were additive (Roberge, Normandeau, Einum, Guderley, & Bernatchez,

2008). The genetic effects of captive breeding could be gradually diluted if the proportion of

hatchery fish in the recipient population is not substantial. The proportion of hatchery fish in

the landings in the inner part, central part, and outside of Kagoshima Bay were highest in

1990, at 83.3%, 33.5% and 7.4%, respectively, but in 2015 the proportion was only ~1.0%

(even in the inner part). The genetic diversity indices recovered for fish in Kagoshima Bay

suggested that the genetic effects were gradually diluted. Increasing wild fish showed no

fitness decline in the recipient population due to small proportions of hatchery fish in the

population.

We should be careful when speculating about cases with very high proportions of hatchery

fish in the recipient population, particularly cases that use captive breeding. Captive-reared

parents are repeatedly used for abalone, barfin flounder, black sea bream, Japanese flounder

and tiger puffer (Table S1). In the case of barfin flounder, one million of juveniles (~8 cm TL)

were released annually on the Pacific coast of Hokkaido. The catch remarkably increased

from 0.01 tonnes in 1996 to over 100 tonnes in 2015, and almost all catches in the North

Pacific, from Ibaraki Prefecture to Hokkaido (170 tonnes), consisted of hatchery fish

(NPJSEC, 2015). In this case, ~500 parent fish have been used for seed production, for 1–3

generations in captivity, and the expected heterozygosity is very high, at 0.87 (Andoh,

Misaka, & Ichikawa, 2013). We must wait for many more generations of barfin flounder

before we can determine the consequences of this case.

5.2 | Japanese scallop

Japanese scallop account for ~27% of the landed amount of the Hokkaido fishery

(www.pref.hokkaido.lg.jp) and all landings are created by releases of wild-born spat and

naturally-reproduced spat from released individuals (Kitada & Fujishima; Nishihama, 2001;

Nishihama, 2001; Uki, 2006). The scallop-ranching programmes in Hokkaido were initiated

by fishers and are managed by cooperatives. Fishers pay most of the programme costs (Uki,

2006). The cooperatives use the following management system for the Japanese scallop

fishery: (1) mass-releases of wild-born spat (wild-born larvae collected in the sea are reared in

net cages for one year, until release); (2) removal of predators, such as starfish, before release;

(3) monitoring of the density of scallops in the fishing ground; and (4) rotation of the fishing

grounds chosen for harvesting (Goshima & Fujiwara, 1994; Kitada & Fujishima; Nishihama,

2001; Uki, 2006). The fishing grounds are generally partitioned into four areas, and 1-year-old

wild-born spat (~4.5 cm) that were reared in net cages are released into a given area after

removing starfish, and the released scallops are harvested after three years, when they reach 4

years old. In the next year, the spats are released into another area. This fishing-ground

rotation system enables complete prohibition of the fishery for three years after release.

Catches of Japanese scallop in the release areas have increased remarkably since the first

releases in the 1970s, and reached the historical maximum of 359 000 tonnes in 2014 (Figure

6a, updated from Kitada, 2018). The catch substantially dropped to 232 000 tonnes (65%) in

2015 due to a bottom disturbance on the Okhotsk coast (accidentally caused by a low-pressure

bomb on 16–17 December 2014, as reported in the previous study), but recovered to 305 000

tonnes in 2018, as expected; the catch recovery in 2018 would have been created by spat

released in 2015. The long-term trends in the release and catch statistics demonstrate that sea

ranching of scallop in Hokkaido is clearly successful with the highest economic efficiency (18

± 5).

5.3 | Chum salmon

Chum salmon hatchery-stock enhancement in Japan is one of the world's largest salmon

stocking programmes (Amoroso, Tillotson, & Hilborn, 2017). Hokkaido is Japan's major

production area for chum salmon, with the returns accounting for 21% amount of Hokkaido's

fishery (www.pref.hokkaido.lg.jp). Most of the landings are created by releases of hatchery-

born fish, and fishers pay ~7% of the landings of chum salmon for hatcheries in Hokkaido

(Kitada, 2014). The long-term trends in releases and catches indicate that the carrying

capacity in natural rivers is ~10 million fish maximum, and the great increase was created by

hatchery stocks (Figure 6b, updated from Miyakoshi, Nagata, Kitada, & Kaeriyama, 2013).

Total production in Hokkaido reached a historical maximum of 61 million returns in 2004,

then substantially decreased to 16 million in 2017. Chum salmon in Japan are at the southern

margin of the species' range; the previous study found that 30% of the variation in decreasing

catches was attributed to increasing sea surface temperature (SST) anomaly, and 62% was

explained by SST and catches by Russian after 1996 (Kitada, 2018). However, there were

clear differences between the run-timing of the chum salmon populations in Russia (Jun–Aug)

and Japan (Sep–Nov) (Kondo, Hirano, Nakayama, Miyake, 1965; Morita, 2016). Migration

routes were also different. These results suggested that the negative correlation between the

Japanese and Russian catches, reported in the previous study, were spurious. The decline in

Japan's catch might be caused by interactions between the long-term hatchery operations and

a warming climate. This hatchery–climate interaction merits further study.

There is no reported evidence of fitness decline among hatchery-born and wild-born chum

salmon in Japan. The observed genetic effect is an altered population structure, with some

populations nested over the seven and/or eight regional groups (Beacham, Sato, Urawa, Le, &

Wetklo, 2008; Kitada, 2014; Sato, Templin, Seeb, Seeb, & Urawa, 2014). This might be

caused by past translocation history (Beacham, Sato, Urawa, Le, & Wetklo, 2008; Kaeriyama

& Qin, 2014). The run-timing distribution of Hokkaido chum salmon was probably altered as

a result of preferred enhancement of early-running fish (returning in September/October)

while the late-running population almost disappeared (Miyakoshi, Nagata, Kitada, &

Kaeriyama, 2013). Lower reproductive success of the early spawners was observed in

sockeye salmon in Washington State, USA, and early-emerging juveniles had relatively low

survival in recent years, suggesting fitness costs from early spawning, which could reduce

population productivity during a warming climate (Tillotson, Barnett, Bhuthimethee, Koehler,

& Quinn, 2019).

Almost all chum salmon returning to Japan were hatchery-reared fish (Kaeriyama, 1999).

Hatchery fish are produced from returned adult fish every year, and the number of parent fish

maintained has been large (i.e., 15 000–85 000) (Kitada, 2018). Even so, natural spawning of

chum salmon was found in 31–37% of 238 non-enhanced rivers surveyed in Hokkaido

(Miyakoshi et al., 2012), and in 94% of 47 enhanced rivers and 75% of 47 non-enhanced

rivers on the northern coast of Honshu Island (Sea of Japan) (Iida, Yoshino, Katayama, 2018).

In addition, a study using otolith thermal-marking estimated the proportions of naturally-

spawned chum salmon to total production at 16–28%, among eight rivers surveyed in

Hokkaido, with large variation within rivers (range 0–50%) (Morita, Takahashi, Ohkuma, &

Nagasawa, 2013). Despite variations in the proportion of natural spawning in the rivers, gene

flow facilitates the genetic admixture of hatchery-released fish, hatchery descendants and wild

fish in the whole Japan populations.

To visualise the magnitude of gene flow between populations, I revisited the microsatellite

data of 26 chum salmon populations in Japan (14 loci, $n = 6\ 028$; Beacham, Sato, Urawa, Lei,

& Wetklo, 2008) and computed F_{ST} values between pairs of populations based on the bias-

corrected G_{ST} (Nei & Chesser, 1983), using the GstNC function in the R package

FinePop1.5.1. This G_{ST} estimator provides an unbiased estimate of F_{ST} when the number of

loci becomes large (Kitada, Nakamichi, & Kishino, 2017). I tied population pairs when the

pairwise F_{ST} value was <0.01 and superimposed this schematic with the locations of

hatcheries (Figure 7). The mean of the pairwise F_{ST} was very small (0.007 ± 0.003). The

threshold $F_{ST} = 0.01$ was equivalent to $4N_e m = 99$, where N_e is the effective population

size, and m is the migration rate. This means that 99 effective parents migrate between each

pair of populations per generation (see Waples & Gaggiotti, 2006). Figure 7 depicts

substantial gene flow between the chum salmon populations. The causal mechanisms of

population structuring are migration and genetic drift, and differentiation depends on the

number of migrants ($N_e m$) (Hauser & Calvalho, 2008; Waples & Gaggiotti, 2006). Even in a

high-gene-flow species, the Atlantic cod (*Gadus morhua*, Gadidae), temporally stable but

significantly differentiated structure was found in the populations (Hauser & Calvalho, 2008).

Constant gene flow among populations can create a stable genetic mixture in a meta-

population, as observed for Pacific herring populations in northern Japan (Kitada et al., 2017).

Hatchery practices might increase the likelihood of chum salmon to stray (Quinn, 1993). I

summarised the results of the marking studies of chum salmon juveniles in Hokkaido, where

2 028 thousand hatchery-reared salmon juveniles (3.5–4 cm BL, 0.45–5 g) were fin-clipped

and/or operculum-clipped and released without rearing between 1951 and 1955 (Sakano,

1960). The results revealed that $50 \pm 22\%$ of a total of 2 085 recoveries were recaptured in the

river of their release, and $84 \pm 12\%$ were found including recoveries in nearby coasts

(Supporting Data); hence, the spawning fidelity of hatchery-reared chum salmon was

moderate. Estimates of straying can vary largely between specific hatchery releases and

rivers, but the genetic integrity of a population can be altered by straying regardless of the

strength of the native population's spawning fidelity (Quinn, 1993). Early theoretical work

predicted that >99% (50%) of wild genes were replaced by hatchery genes in 12 (2)

generations, in the case of equal fitness between hatchery and wild fish with additive effect, at

a stocking rate of 0.5 (Matsuishi, Kishino, & Numachi, 1995; see also Kitada et al. 2019,

Figure 5). From these results together, I speculate that all chum salmon returning to Japan are

hatchery-released fish or wild-born from hatchery descendants, resulting from the large-scale,

long-term releases.

6| Conclusion

Many cases of hatchery releases were economically unprofitable if the costs of personnel and

the negative impacts on wild populations are taken into account. Stocking effects were

generally small and the population dynamics were unaffected by releases but instead

essentially depended on the carrying capacity of the nursery habitat. Captive breeding reduces

the fitness of hatchery-reared fish in the wild, and long-term, extensive hatchery releases can

replace wild genes and may cause fitness decline in the recipient population. All chum salmon

returning to Japan should be hatchery fish and/or hatchery descendants, which may reduce the

fitness of populations in the warming environment. Recovery of nursery habitats and fishing

effort reduction outperform hatcheries in the long run.

Acknowledgements

I thank Hirohisa Kishino, Yasuyuki Miyakoshi and Katsuyuki Hamasaki for useful comments

on the manuscript. I appreciate Toshio Okazaki for valuable suggestions, Tomonari Fujita and

Shigeki Dan for providing data, and Reiichiro Nakamichi for help with making the maps. This

study was supported by the Japan Society for the Promotion of Science Grant-in-Aid for

Scientific Research (KAKENHI) No. 18K0578116.

Data Availability Statement

All source data used are in the public sector and links to their online sources are specified in

the text or in Supplementary Data.

References

Agriculture, Forestry and Fisheries Research Council (AFFRC) (Ed.). (1989). *Marine Ranching*. Tokyo: Koseisha Koseikaku [in Japanese].

Amoroso, R. O., Tillotson, M. D., & Hilborn, R. (2017). Measuring the net biological impact of fisheries enhancement: Pink salmon hatcheries can increase yield, but with apparent costs to wild populations. Canadian Journal of Fisheries and Aquatic Sciences, 74, 1233–1242. https://doi.org/10.1139/cjfas-2016-0334

Andoh, T., Mikasa, T, & Ichikawa, T. (2013). Genetic monitoring and conservation programs to maintain the genetic diversity. In M. Aritaki (Ed.), *Stock Enhancement and Risk Management for Coastal Fishery Resources.-Monitoring on genetic diversity and efficiency of stock enhancement* (pp. 38–53). Tokyo: Koseisha Koseikaku. [in Japanese].

Araki, H., Cooper, B., & Blouin, M. S. (2007). Genetic effects of captive breeding cause a rapid, cumulative fitness decline in the wild. *Science*, 318, 100–103. doi: 10.1126/science.1145621

Araki, H., Berejikian, B. A., Ford, M. J., & Blouin, M. S. (2008). Fitness of hatchery‐reared salmonids in the wild. *Evolutionary Applications*, 1, 342–355. https://doi.org/10.1111/j.1752-4571.2008.00026.x

Araki, H., & Schmid, C. (2010). Is hatchery stocking a help or harm? Evidence, limitations and future directions in ecological and genetic surveys. *Aquaculture*, 308 (Sl), S2–S11. doi.org/10.1016/j.aquaculture.2010.05.036

Atsuchi, S., & Masuda, Y. (2004). Effectiveness of the releases of hatchery-produced stock of Japanese flounder *Paralichthys olivaceus* in Kagoshima Bay, southern Japan. *Nippon Suisan Gakkaishi* 70, 910–921 [In Japanese with English abstract]. https://doi.org/10.2331/suisan.70.910

Beacham, T. D., Sato, S., Urawa, S. Lei, K. D., & Wetklo, M. (2008). Population structure and

stock identification of chum salmon *Oncorhynchus keta* from Japan determined by

microsatellite DNA variation. *Fishery Science*, 74, 983–994. doi: 10.1111/j.1444-

2906.2008.01616.x

Bell, J. D., Rothlisberg, P. C., Munro, J. L., Loneragan, N. R., Nash, W. J., Ward, R. D., &

Andrew, N. L. (2005). *Restocking and stock enhancement of marine invertebrate*

fisheries. *Advances in Marine Biology*, 49, Oxford: Elsevier.

Blaxter, J. H. S. (2000). The enhancement of marine fish stocks. *Advances in Marine Biology*,

38, 1–5. doi: org/10.1016/S0065-2881(00)38002-6

Christie, M. R., Marine, M. L., French, R. A., & Blouin, M. S. (2012). Genetic adaptation to

captivity can occur in a single generation. *Proceedings of the National Academy of*

Sciences, 109, 238–242. doi: 10.1073/pnas.1111073109

Christie, M. R., Ford, M. J., & Blouin, M. S. (2014). On the reproductive success of early-

generation hatchery fish in the wild. *Evolutionary Applications*, 7, 883–896. doi:

10.1111/eva.12183

Fisheries Agency, Fisheries Research Agency (FRA) & National Association for the

Promotion of Productive Sea (NAPPS) (Eds.) (1985–2019). *Annual Statistics of Seed*

Production and Release in 1983–2017. Yokohama: FRA [in Japanese].

Fleming, I. A., Hindar, K., MjÖlnerÖd, I. B., Jonsson, B., Balstad, T., & Lamberg, A. (2000).

Lifetime success and interactions of farm salmon invading a native population.

Proceedings of the Royal Society of London B, 267, 1517–1523. doi:

10.1098/rspb.2000.1173

Ford, M. J. (2002). Selection in captivity during supportive breeding may reduce fitness in the

wild. *Conservation Biology*, 16, 815–825. https://doi.org/10.1046/j.1523-

1739.2002.00257.x

Fushimi, H. (2001). Production of juvenile marine finfish for stock enhancement in Japan.

Aquaculture, 200(1–2), 33–53. https://doi.org/10.1016/S0044-8486(01)00693-7

Goshima, S., & Fujiwara, H. (1994). Distribution and abundance of cultured scallop

Patinopecten yessoensis in extensive sea beds as assessed by underwater camera.

Marine Ecology Progress Series, 110, 151–158. https://www.jstor.org/stable/24847584

Hamasaki, K. (2008). Environmental change and resource enhancement: a case of abalone. In:

S. Kitada, M. Kaeriyama, K. Hamasaki & N. Taniguchi (Eds.), *Enhancement and*

Conservation of Fisheries Resources (pp. 86–104). Tokyo: Seizando.

Hamasaki, K., & Kitada, S. (2006). A review of kuruma prawn (*Penaeus japonicus*) stock

enhancement programs in Japan. *Fisheries Research*, 80, 80–90. doi:

org/10.1016/j.fishres.2006.03.018

Hamasaki, K, & Kitada, S. (2008a). The enhancement of abalone stocks: lessons from

Japanese case studies. *Fish and Fisheries*, 9, 243–260. doi: 10.1111/j.1467-

2979.2008.00280.x

Hamasaki, K., & Kitada, S. (2008b). Potential of stock enhancement for decapod crustaceans.

Reviews in Fisheries Science, 16, 164–174. doi: org/10.1080/10641260701681722

Hamasaki, K., & Kitada, S. (2013). Catch fluctuation of Kuruma prawn, *Penaeus japonicus* in

Japan relative to ocean climate variability and a stock enhancement program. *Reviews*

in Fisheries Science, 21(3-4), 454–468. https://doi.org/10.1080/10641262.2013.800781

Hamasaki, K., Obata, Y., Dan, S., & Kitada, S. (2011). A review of seed production and stock

enhancement for commercially important portunid crabs in Japan. *Aquaculture*

International, 19, 217–235. doi: org/10.1007/s10499-010-9387-7

Hauser, L., & Carvalho, G. R. (2008). Paradigm shifts in marine fisheries genetics: ugly

hypotheses slain by beautiful facts. *Fish and Fisheries*, 9, 333–362.

https://doi.org/10.1111/j.1467-2979.2008.00299.x

Hayakawa, J., Yamakawa, T., & Aoki, I. (2007). Long-term fluctuation in the abundance of

abalone and top shell in Japan and factors affecting those fluctuations. *Bulletin of the*

Japanese Society of Fisheries Oceanography, 71, 96–105 [In Japanese with English

abstract].

Hilborn, R. (1992). Hatcheries and the future of salmon in the Northwest. *Fisheries*, 17, 5–8.

doi: org/10.1577/1548-8446(1992)017<0005:HATFOS>2.0.CO;2

Hilborn, R. (1998). The economic performance of marine stock enhancement projects.

Bulletin of Marine Science, 62, 661–674.

Hilborn, R. (2004) Population management in stock enhancement and sea ranching. In K.

Leber, S. Kitada, H. L. Blankenship & T. Svåsand (Eds.), *Stock Enhancement and Sea*

Ranching, Developments Pitfalls and Opportunities (pp. 201–210). Oxford: Blackwell.

Iida, M., Yoshino, K., Katayama, S. (2018). Current status of natural spawning of chum

salmon *Oncorhynchus keta* in rivers with or without hatchery stocking on the Japan

Seaside of northern Honshu, Japan. *Fisheries Science* 84, 453–459.

https://doi.org/10.1007/s12562-018-1192-7

Ishino, K. (1999). Stocking effectiveness of Japanese flounder, *Paralichthys olivaceus*,

fingerlings released on the south-western coast of Hokkaido Prefecture, Japan. In: B. R.

Howel, E. Moksness & T. Svåsand (Eds.), *Stock Enhancement and Sea Ranching (*pp

557–572). Oxford: Blackwell.

Kaeriyama, M. (1999). Hatchery programmes and stock management of salmonid populations

in Japan. In: B. R. Howel, E. Moksness & T. Svåsand (Eds.), *Stock Enhancement and*

Sea Ranching (pp. 153–167). Oxford: Blackwell.

Kaeriyama, M., Seo, H., Kudo, H., & Nagata, M. (2012). Perspectives on wild and hatchery

salmon interactions at sea, potential climate effects on Japanese chum salmon, and the

need for sustainable salmon fishery management reform in Japan. *Environmental Biology of Fishes*, 94, 165–177. https://doi.org/10.1007/s10641-011-9930-z

Kaeriyama, M., Seo, H., & Qin, Y. X. (2014). Effect of global warming on the life history and population dynamics of Japanese chum salmon. *Fisheries Science*, 80, 251–260. DOI 10.1007/s12562-013-0693-7

Kaeriyama, M., & Qin, Y. (2014). Biological interactions between wild and hatchery-produced Pacific salmon. In P. T. K. Woo & D. J. Noakes (Eds.), Salmon (pp. 223–238). New York, NY: Nova Science Publishers.

Kasugai, K., Saneyoshi, H., Aoyama, T., Shinriki, Y., Iijima, A., & Miyakoshi, Y. (2016). Early marine migration of juvenile chum salmon along the Pacific coast of eastern Hokkaido. *North Pacific Anadromous Fish Commssion Bulletin*, 6, 61–72.

Kayano, Y., Tanaka, T., & Hayashi, H. (1998). Observations of artificially releasing seedlings and native fishes around the marine ranching area. *Fisheries Engineering*, 35, 303–309 [In Japanese with English abstract].

Kitada, S. (1999). Effectiveness of Japan's stock enhancement programs: current perspectives. In: B. R. Howel, E. Moksness & T. Svåsand (Eds.), *Stock Enhancement and Sea Ranching* (pp. 103–131). Oxford: Blackwell.

Kitada, S. (2014). Japanese chum salmon stock enhancement: current perspective and future challenges. Fisheries Science 80, 237–249. https://doi.org/10.1007/s12562-013-0692-8

Kitada, S. (2018). Economic, ecological and genetic impacts of marine stock enhancement and sea ranching: A systematic review. *Fish and Fisheries*, 19, 511–532. https://doi.org/10.1111/faf.12271

Kitada, S., & Fujishima, H. (1997). The stocking effectiveness of scallop in Hokkaido. *Nippon Suisan Gakkaishi*, 63, 686–693 [In Japanese with English abstract]. doi: org/10.2331/suisan.63.686

Kitada, S., Hiramatsu, K., & Kishino, H. (1994). Estimating mortality rates from tag

recoveries: incorporating over-dispersion, correlation, and change points. *ICES Journal*

of Marine Science, 51, 241–251. https://doi.org/10.1006/jmsc.1994.102

Kitada, S., & Hirano, K. (1987). Estimation of mortality coefficients from tag recovery.

Nippon Suisan Gakkaishi, 53, 1765–1770 [In Japanese with English

abstract].https://doi.org/10.2331/suisan.53.1765

Kitada, S., & Kishino, H. (2006). Lessons learned from Japanese marine finfish stock

enhancement programmes. *Fisheries Research*, 80, 101–112. doi:

org/10.1016/j.fishres.2006.03.019

Kitada, S., Kishino, H., & Hamasaki, K. (2011). Bias and significance of relative reproductive

success estimates based on steelhead trout (*Oncorhynchus mykiss*) data: a Bayesian

meta-analysis. *Canadian Journal of Fisheries and Aquatic Sciences*, 68, 1827–1835.

https://doi.org/10.1139/f2011-087

Kitada, S., Nakajima, K., Hamasaki, K., Shishidou. H., Waples, R., & Kishino, H. (2019).

Rigorous monitoring of a large-scale marine stock enhancement program demonstrates

the need for comprehensive management of fisheries and nursery habitat. *Scientific*

Reports, 9, 5290. https://doi.org/10.1038/s41598-019-39050-3

Kitada, S., Nakamichi, R., & Kishino, H. (2017). The empirical Bayes estimators of fine-scale

population structure in high gene flow species. *Molecular Ecology Resources*, 17,

1210–1222. https://doi.org/10.1111/1755-0998.12663

Kitada, S., Shishidou, H., Sugaya, T., Kitakado, T., Hamasaki, K., & Kishino, H. (2009).

Genetic effects of long-term stock enhancement programs. *Aquaculture*, 290, 69–79.

doi: org/10.1016/j.aquaculture.2009.02.011

Kitada, S., Taga, Y., & Kishino, H. (1992). Effectiveness of a stock enhancement program evaluated by a two-stage sampling survey of commercial landings. *Canadian Journal of Fisheries and Aquatic Sciences*, 49, 1573–1582. doi: org/10.1139/f92-175

Kitada, S., Yoshikai, R., Fujita, T., Hamasaki, K., Nakamichi, R., & Kishino, H. (2017). Population structure and persistence of Pacific herring following the Great Tohoku earthquake. *Conservation Genetics*, 18, 423–437. DOI https://doi.org/10.1007/s10592-016-0918-2

Kojima, H. (1995). Evaluation of abalone stock enhancement through the release of hatchery-reared seeds. *Marine and Freshwater Research*, 46, 689–695. https://doi.org/10.1071/MF9950689

Kondo, H., Hirano, Y., Nakayama, N., & Miyake, M. (1965). Offshore distribution and migration of Pacific salmon (genus *Oncorhynchus*) based on tagging studies (1958-1961). *Bulletin of International North Pacific Fisheries Commission*, 17, 1–213.

Koya, Y. (2005). Stock enhancement of barfin flounder in Hokkaido. *Technical Reports of Hokkaido Fisheries Experimental Station*, 5, 43–49 [In Japanese].

Kudo, K., & Kimoto, H. (1994). Fishery management of marine ranching in the Oita prefectural sea area. *Fisheries Engineering*, 31, 121–126 [In Japanese with English abstract].

Kurata, M. (1999). On the decline in the growth of maricultured scallop, *Patinopecten yessoensis*, in the Okhotsk coastal area of Hokkaido. *Research Reports of Hokkaido Fisheries Experimental Station*, 54, 25–32 [In Japanese].

Laikre, L., M. K. Schwartz, R. S. Waples, N. Ryman, & the GeM working group. (2010). Compromising genetic diversity in the wild: Unmonitored large-scale release of plant and animals. *Trends in Ecology and Evolution*, 25, 520–529. doi: org/10.1016/j.tree.2010.06.013

Le Vay, L., Carvalho, G. R., Quinitio, E. T., Lebata, J. H., Ut, V. N., & Fushimi, H. (2007).

Quality of hatchery-reared juveniles for marine fisheries stock enhancement.

Aquaculture, 268(1-4), 169–180. https://doi.org/10.1016/j.aquaculture.2007.04.041

Lee, S. I., & Zhang, C. I. (2018). Evaluation of the effect of marine ranching activities on the

Tongyeong marine ecosystem. *Ocean Science Journal*, 53, 557–582.

https://doi.org/10.1007/s12601-018-0045-8

Lee, S. G., & Rahman, M. A. (2018). Ecological stock enhancement programs (ESEPs) based

fisheries rebuilding plan (FRP) in Korea. *Journal of Environmental Biology*, 39, 936–

942. DOI:10.22438/jeb/39/5(SI)/31

Masuda, R., & Tsukamoto, K. (1998). Stock enhancement in Japan: review and perspective.

Bulletin of Marine Science, 62, 337–358.

Matsuishi, T., Kishino, H., & Numachi, K. (1995). A model of gene displacement by stocking

activities. *Nippon Suisan Gakkaishi*, 61, 326–330 [In Japanese with English abstract].

https://doi.org/10.2331/suisan.61.326

Matsumura, Y. (2005). Optimal release strategy of hatchery-produced ocellate puffer *Takifugu*

rubripe in Ariake Sound by mark-recapture experiments, based on the stocking

effectiveness for young-of-the-year. *Nippon Suisan Gakkaishi*, 71, 805–814 [In

Japanese with English abstract]. https://doi.org/10.2331/suisan.71.805

Matsumura, Y. (2006). Stocking effectiveness of hatchery-produced juveniles of ocellate

puffer *Takifugu rubripe* in natal spawning ground of Ariake Sound. *Nippon Suisan*

Gakkaishi, 72, 1029–1038 [In Japanese with English abstract].

https://doi.org/10.2331/suisan.72.1029

McGinnity, P., Prodöhl, P., Ferguson, A., Hynes, R., ó Maoiléidigh, N., Baker, N., ... Taggart,

J. (2003). Fitness reduction and potential extinction of wild populations of Atlantic

salmon, *Salmo salar*, as a result of interactions with escaped farm salmon. *Proceedings*

of the Royal Society of London B, 270, 2443–2450. doi: 10.1098/rspb.2003.2520

Ministry of Agriculture, Forestry and Fisheries (MAFF) (1964–2018). *Annual statistics of*

fisheries and aquaculture production in 1966–2019. Tokyo: Association of Agriculture

and Forestry Statistics [in Japanese].

Miyakoshi, Y., Nagata, M., Sugiwaka, K. I., & Kitada, S. (2001). Commercial harvest of

hatchery-reared masu salmon *Oncorhynchus masou* estimated by a coast-wide sampling

program in Hokkaido, northern Japan, and the two-stage sampling schemes of landings.

Fisheries Science, 67, 126–133. doi: 10.1046/j.1444-2906.2001.00208.x

Miyakoshi, Y., Urabe, H., Saneyoshi, H., Aoyama, T., Sakamoto, H., Ando, D., ... Nagata, M.

(2012). The occurrence and run timing of naturally spawning chum salmon in northern

Japan. *Environmental Biology of Fishes*, 94, 197–206. doi: org/10.1007/s10641-011-

9872-5

Miyakoshi, Y., Nagata, M., Kitada, S., & Kaeriyama, M. (2013). Historical and current

hatchery programs and management of chum salmon in Hokkaido, northern Japan.

Reviews in Fisheries Science, 21, 469–479. doi: org/10.1080/10641262.2013.836446

Moksness, E., & Støle, R. (1997). Larviculture of marine fish for sea ranching purposes: is it

profitable? *Aquaculture*, 155, 341–353. doi: org/10.1016/S0044-8486(97)00102-6

Moksness, E., Støle, R., & van der Meeren, G. (1998). Profitability analysis of sea ranching

with Atlantic salmon (*Salmo salar*), Arctic charr (*Salvelinus alpinus*), and European

lobster (*Homarus gammarus*) in Norway. *Bulletin of Marine Science*, 62, 689–699.

Morita, K., Saito, T., Miyakoshi, Y., Fukuwaka, M. A., Nagasawa, T., & Kaeriyama, M.

(2006). A review of Pacific salmon hatchery programmes on Hokkaido Island, Japan.

ICES Journal of Marine Science, 63, 1353–1363.

https://doi.org/10.1016/j.icesjms.2006.03.024

Morita, K., Takahashi, S., Ohkuma, K., & Nagasawa, T. (2013). Estimation of the proportion

of wild chum salmon *Oncorhynchus keta* in Japanese hatchery rivers. *Nippon Suisan*

Gakkaishi, 79, 206–213 [In Japanese with English abstract]. doi:

org/10.2331/suisan.79.206

Morita, K. (2014). Japanese wild salmon research: toward a reconciliation between hatchery

and wild salmon management. *NPAFC Newsletter*, 35, 4–14.

Morita, K. (2016). Trends in Russian chum salmon populations. *FRA Salmonid Research*

Report, 10, 38–40 [In Japanese].

Murakami, O. (2012). Effect of large-scale release of barfin flounder in the western Pacific

Hokkaido. *Reports of Hokkaido Fisheries Experimental Station*, 706 [In Japanese].

Nagata, M., Miyakoshi, Y., Urabe, H., Fujiwara, M., Sasaki, Y., Kasugai, K., ... & Kaeriyama,

M. (2012). An overview of salmon enhancement and the need to manage and monitor

natural spawning in Hokkaido, Japan. *Environmental Biology of Fishes*, 94, 311–323.

https://doi.org/10.1007/s10641-011-9882-3

Nakagawa, M., Okouchi, H., Adachi, J. (2004). Stocking effectiveness of black rockfish

Sebastes schlegeli released in Yamada Bay evaluated by a fish market census. In K.

Leber, S. Kitada, H. L. Blankenship & T. Svåsand (Eds.), *Stock Enhancement and Sea*

Ranching, Developments Pitfalls and Opportunities (pp. 501–511). Oxford: Blackwell.

Nakajima, H., Kai, M., Koizumi, K., Tanaka, T., & Machida, M. (2008). Optimal release

locations of juvenile ocellate puffer *Takifugu rubripes* identified by tag and release

experiments. *Reviews in Fisheries Science*, 16(1-3), 228–234.

https://doi.org/10.1080/10641260701684106

Nakajima, K., Kitada, S., Yamazaki, H., Takemori, H., Obata, Y., Iwamoto, A., & Hamasaki,

K. (2013). Ecological interactions between hatchery and wild fish: a case study based

on the highly piscivorous Japanese Spanish mackerel. *Aquaculture Environment Interactions*, 3, 231–243. doi: org/10.3354/aei00065

Nakajima, K., Kitada, S., Habara, Y., Sano, S., Yokoyama, E., Sugaya, T., ... & Hamasaki, K. (2014). Genetic effects of marine stock enhancement: a case study based on the highly piscivorous Japanese Spanish mackerel. *Canadian Journal of Fisheries and Aquatic Sciences*, 71, 301–314. https://doi.org/10.1139/cjfas-2013-0418

Nakamura, A., Kitada, S., Hamasaki, K., Okouchi, H. (2005). Catch fluctuation in *Haliotis* spp. abalones: annual landings of the Ezo abalone *Haliotis discus hannai* in relation to climate oscillation and stock enhancement programs. *Saibai Gyogyo Gijyutsu Kaihatsu Kenkyu*, 33, 45–54 [In Japanese with English abstract].

Nei, M., & Chesser, R. K. (1983). Estimation of fixation indices and gene diversities. *Annals of Human Genetics*, 47, 253–259. https://doi.org/10.1111/j.1469-1809.1983.tb00993.x

Nishihama, Y. (2001). *Scallop fisheries in the Okhotsk Sea*. Sapporo: Hokkaido University Press [in Japanese].

North Pacific Anadromous Fish Commission (NPAFC). *NPAFC Pacific salmonid catch and hatchery release data (updated 2019 summer)*. Vancouver, BC: NPAFC. Retrieved from http://www.npafc.org

North Pacific Japan Stock Ehancement Counci (NPJSEC) (2015). Barfin flounder enhancement plan 2015–2021. Fisheries Agency [In Japanese].

Obata, Y., Imai, H., Kitakado, T., Hamasaki, K., & Kitada, S. (2006). The contribution of stocked mud crabs *Scylla paramamosain* to commercial catches in Japan, estimated using a genetic stock identification technique. *Fisheries Research*, 80, 113–121. doi: org/10.1016/j.fishres.2006.03.016

Obata, Y., Yamazaki, H., Iwamoto, A., Hamasaki, K., & Kitada, S. (2008). Evaluation of

stocking effectiveness of the Japanese Spanish mackerel in the eastern Seto Inland Sea,

Japan. *Reviews in Fisheries Science*, 16, 35–24. doi: org/10.1080/10641260701697041

Ohnuki, T., Morita, K., Tokuda, H., Okamoto, Y., & Ohkuma, K. (2015). Numerical and

economic contributions of wild and hatchery pink salmon to commercial catches in

Japan estimated from mass otolith markings. *North American Journal of Fisheries*

Management, 35, 598–604. https://doi.org/10.1080/02755947.2015.1020078

Okamoto, K. (2004). Juvenile release and market size recapture of the swimming crab

Portunus trituberculatus (Miers) marked with coded wire tags. In K. Leber, S. Kitada,

H. L. Blankenship & T. Svåsand (Eds.), *Stock Enhancement and Sea Ranching,*

Developments Pitfalls and Opportunities (pp. 181–186). Oxford: Blackwell.

Okouchi, H., Kitada, S., Tsuzaki, T., Fukunaga, T., & Iwamoto, A. (1999). Number of returnes

and economic return rates of hatchery-released flounder *Paralichthys olivaceus* in

Miyako Bay – Evaluation by a fish market census. In: B. R. Howel, E. Moksness & T.

Svåsand (Eds.), *Stock Enhancement and Sea Ranching* (pp. 573–582).

Okouchi, H., Kitada, S., Iwamoto, A., & Fukunaga, T. (2004). The flounder stock

enhancement in in Miyako Bay. *FAO Fisheries Technical Paper*, 429, 171–202.

Okouchi, H., Kitada, S., Morioka, T., Imamura, S. (1994). A comparison of the quality of

hatchery-reared red sea bream *Pagrus major* based on mortality rate estimated from tag

recoveries. *Nippon Suisan Gakkaishi*, 60, 229–233 [in Japanese with English abstract].

https://doi.org/10.2331/suisan.60.229

Quinn, T. P. (1993). A review of homing and straying of wild and hatchery-produced salmon.

Fisheries Research, 18(1-2), 29–44. https://doi.org/10.1016/0165-7836(93)90038-9

Reisenbichler, R. R., & McIntyre, J. D. (1977). Genetic differences in growth and survival of juvenile hatchery and wild steelhead trout, *Salmo gairdneri. Journal of the Fisheries Board of Canada*, 34, 123–128. doi: org/10.1139/f77-015

Roberge, C., Normandeau, E., Einum, S., Guderley, H., & Bernatchez, L. (2008). Genetic consequences of interbreeding between farmed and wild Atlantic salmon: insights from the transcriptome. *Molecular Ecology*, 17(1), 314-324. https://doi.org/10.1111/j.1365-294X.2007.03438.x

Sakai, Y., Tajima, K., & Agatsuma, Y. (2004). Stock enhancement of the short-spined sea urchin. In K. Leber, S. Kitada, H. L. Blankenship & T. Svåsand (Eds.), *Stock Enhancement and Sea Ranching, Developments Pitfalls and Opportunities* (pp. 465–476). Oxford: Blackwell.

Sakano, E. (1960). Results from marking experiments on young chum salmon in Hokkaido, 1951-1959. *Scientific Reports of the Hokkaido Salmon Hatchery*, 15, 17–38 [in Japanese with English Abstract].

Sato, S., Templin, W. D., Seeb, L. W., Seeb, J. E., & Urawa, S. (2014). Genetic structure and diversity of Japanese chum salmon populations inferred from single-nucleotide polymorphism markers. *Transactions of the American Fisheries Society*, 143, 1231–1246. https://doi.org/10.1080/00028487.2014.901251

Shiota, K. & Kitada. S. (1992). Life history of swimming crab in Hiuchi Nada, Seto Inland Sea estimated from marking experiments. *Nippon Suisan Gakkaishi* 58, 2297–2302 [in Japanese with English abstract]. https://doi.org/10.2331/suisan.58.2297

Shishidou, H. (2002). Stocking effectiveness of red sea bream *Pagrus major* in Kagoshima Bay. *Fisheries Science*, 68 (Suppl I), 904–907 https://doi.org/10.2331/fishsci.68.sup1_904

Shishidou, H. & Kitada, S. (2007). Stocking effectiveness of red sea bream *Pagrus major* in

Kagoshima Bay, Japan. Nippon Suisan Gakkaishi 73, 270–277 [in Japanese with

English abstract].

Sproul, J. T., & Tominaga, O. (1992). An economic review of the Japanese flounder stock

enhancement project in Ishikari Bay, Hokkaido. *Bulletin of Marine Science*, 50, 75–88.

Steward, C. R., & Bjornn, T. C. (1990). Supplementation of salmon and steelhead stocks with

hatchery fish: A synthesis of published literature. *U.S. Fish and Wildlife Services,*

Technical Reports 90 (1), 202 p.

Svåsand, T., Kristiansen, T. S., Pedersen, T., Salvanes, A. V., Engelsen, R., Naevdal, G., &

Nødtvedt, M. (2000). The enhancement of cod stocks. *Fish and Fisheries*, 1, 173–205.

doi: 10.1046/j.1467-2979.2000.00017.x

Takaba, M., Kitada, S., Nakano, H., Morioka, T. (1995). Survival, feeding activity and liver

constituents of red sea bream released into the Seto Inland Sea of Japan with special

reference to fish quality. *Nippon Suisan Gakkaishi*, 61, 574–579 [in Japanese with

English abstract]. https://doi.org/10.2331/suisan.61.574

Takami, H., Saido, T., Endo, T., Noro, T., Musashi, T., & Kawamura, T. (2008). Overwinter

mortality of young-of-the-year Ezo abalone in relation to seawater temperature on the

North Pacific coast of Japan. *Marine Ecology Progress Series*, 367, 203–212.

https://doi.org/10.3354/meps07557

Takeuchi, T. (2001). A review of feed development for early life stages of marine finfish in

Japan. *Aquaculture*, 200(1–2), 203–222. https://doi.org/10.1016/S0044-8486(01)00701-

Taniguchi, N. (2003). Genetic factors in broodstock management for seed production.

Reviews in Fish Biology and Fisheries, 13, 177–185. doi:

org/10.1023/B:RFBF.0000019479.49749.fe

Taniguchi, N., (2004). Broodstock management for stock enhancement programs of marine

fish with assistance of DNA markers (a review). In K. Leber, S. Kitada, H. L.

Blankenship & T. Svåsand (Eds.), *Stock Enhancement and Sea Ranching,*

Developments Pitfalls and Opportunities (pp. 329–338). Oxford: Blackwell.

Tillotson, M. D., Barnett, H. K., Bhuthimethee, M., Koehler, M. E., & Quinn, T. P. (2019).

Artificial selection on reproductive timing in hatchery salmon drives a phenological

shift and potential maladaptation to climate change. *Evolutionary Applications*, 12,

1344–1359. https://doi.org/10.1111/eva.12730

Tominaga, O., & Watanabe, Y. (1998). Geographical dispersal and optimum release size of

hatchery-reared Japanese flounder *Paralichthys olivaceus* released in Ishikari Bay,

Hokkaido, Japan. *Journal of Sea Research*, 40(1–2), 73–81.

https://doi.org/10.1016/S1385-1101(98)00011-2

Tomiyama, T., Watanabe, M., & Fujita, T. (2008). Community-based stock enhancement and

fisheries management of the Japanese flounder in Fukushima, Japan. Reviews in

Fisheries Science, 16(1–3), 146–153. https://doi.org/10.1080/10641260701681847

Tsujimura, H. (2007). Stocking effectiveness of red spotted grouper in Osaka Bay.

www.kannousuiken-osaka.or.jp [In Japanese].

Tsukamoto, K., Kuwada, H., Uchida, K., Masuda, R., Sakakura, A. (1999). Fish quality and

stocking effectiveness: behavioural approach. In: B. R. Howel, E. Moksness & T.

Svåsand (Eds.), Stock Enhancement and Sea Ranching (pp. 203–218). Oxford:

Blackwell.

Uki, N. (2006). Stock enhancement of the Japanese scallop *Patinopecten yessoensis* in

Hokkaido. *Fisheries Research*, 80, 62–66. doi: org/10.1016/j.fishres.2006.03.013

Wada, T., Kamiyama, K., Shimamura, S., Mizuno, T., & Nemoto, Y. (2012). Effectiveness of

stock enhancement of a rare species, spotted halibut *Verasper variegatus*, in Fukushima,

Japan. *Aquaculture*, 364, 230–239. doi: org/10.1016/j.aquaculture.2012.08.030

Wang, Y., Yu, J., & Chen, P. (2018). Remote sensing assessment of ecological effects of

marine ranching in the eastern Guangdong waters, China. *Journal of Geoscience and*

Environment Protection, 6, 101–113. 10.4236/gep.2018.612007

Waples, R. S. (1991). Genetic interactions between hatchery and wild salmonids: lessons from

the Pacific Northwest. *Canadian Journal of Fisheries and Aquatic Sciences*, 48 (S1),

124–133. doi: org/10.1139/f91-311

Waples, R. S. (1999). Dispelling some myths about hatcheries. *Fisheries*, 24, 12–21. doi:

org/10.1577/1548-8446(1999)024<0012:DSMAH>2.0.CO;2

Waples, R. S., & Gaggiotti, O. (2006). What is a population? An empirical evaluation of some

genetic methods for identifying the number of gene pools and their degree of

connectivity. *Molecular Ecology*, 15, 1419–1439. https://doi.org/10.1111/j.1365-

294X.2006.02890.x

Winton, J., & Hilborn, R. (1994). Lessons from supplementation of chinook salmon in British

Columbia. *North American Journal of Fisheries Management*, 14, 1–13.

https://doi.org/10.1577/1548-8675(1994)014<0001:LFSOCS>2.3.CO;2

Zhou, X., Zhao, X., Zhang, S., & Lin, J. (2019). Marine ranching construction and

management in East China Sea: Programs for sustainable fishery and aquaculture.

Water, 11, 1237. https://doi.org/10.3390/w11061237

Yamaguchi, T., Ito, S., Hamasaki, K., & Kitada, S. (2006). Stocking effectiveness of hatchery-

released kuruma prawns estimated by two-stage sampling of commercial catch in

Ariake Sound, Japan. *Fisheries Science*, 72, 233–238. https://doi.org/10.1111/j.1444-

2906.2006.01143.x

1009

Supporting Information

1011 Additional Supporting Information may be found in the online version of this article:

1012

1013 **Supporting Data.**

1014 Catch and release data for major species in Japanese marine stock enhancement and sea-

1015 ranching programmes, and the results of marking experiments with chum salmon conducted

1016 in Hokkaido in the 1950s.

1017

Figure Legends

Figure 1 Changes in the areas of tidal flats and eel grass (*Zostera marina*, Zosteraceae) (1945–1996). From Ministry of Environment, www.env.go.jp and www.biodic.go.jp accessed July 2019

Figure 2 Maps plotting the locations of the marine and salmon hatcheries in Japan, operated by different sectors. From FRA (salmon.fra.affrc.go.jp accessed August 2019) and NAPPS (http://www.yutakanaumi.jp/, see text)

Figure 3 Forest plot of recapture rates for large-scale hatchery releases in Japan. Thin lines indicate 95% confidence intervals, and arrows indicate that the confidence intervals penetrate the scale; areas of the squares are proportional to the weight of the mean [The colour figure can be viewed at wileyonlinelibrary.com]

Figure 4 Performance of large-scale hatchery releases in Japan. Forest plots of **(a)** yield-per-release (YPR), **(b)** economic efficiency, and **(c)** YPR versus economic efficiency. Note: the seed cost used in the analysis did not include personnel expenses, facilities, monitoring, or administration costs. SIS = Seto Inland Sea [The colour figure can be viewed at wileyonlinelibrary.com]

Figure 5 Total catch and wild catches, with release statistics, for representative species in Japanese stocking programmes, for the whole of Japan. Vertical lines depict recovery (expected catch from releases), estimated by multiplying values of yield-per-release (YPR) (see Table 2) and the numbers released (Supplementary Data). For sea urchin, no YPR data were available

Figure 6 Long-term release and catch (return) statistics for **(a)** Japanese scallop (updated from Kitada, 2018) and **(b)** chum salmon in Hokkaido (updated from Miyakoshi, Nagata, Kitada, & Kaeriyama, 2013)

Figure 7 Geneflow between Japanese populations of chum salmon. Populations were tied by lines for pairwise $F_{ST} < 0.01$, estimated from the microsatellite data of 26 populations (14 loci, $n = 6\,028$; Beacham, Sato, Urawa, Lei, & Wetklo, 2008).

Tidal flat (1945-1978-1996)

Zostea marina (1960-1973-1990)

Ariake Sea

Seto Inland Sea

266 204 km²

Tokyo Bay

Mikawa Bay

Ise Bay

Yatsushiro Sea

Fig. 1

Recapture rates (%)

Species	Mean (SD)
Barfin flounder (Hokkaido)	12.1 (0.6)
Black rockfish (Yamada Bay)	11.8 (0.9)
Chum salmon (Hokkaido, Honshu)	2.6 (0.8)
Japanese flounder (Fukushima87)	15.0 (3.3)
Japanese flounder (Fukushima94_02)	12.1 (4.8)
Japanese flounder (Ishikari Bay)	5.7 (3.5)
Japanese flounder (Kagoshima Bay)	2.4 (0.7)
Japanese flounder (Miyako Bay)	14.5 (7.3)
Japanese Spanish mackerel (SIS)	15.0 (0.7)
Red sea bream (Kagoshima Bay)	8.0 (4.2)
Red sea bream (Sagami Bay)	7.1 (2.9)
Red spotted grouper (Osaka Bay)	2.2 (1.0)
Spotted halibut (Fukushima)	11.1 (11.4)
Tiger puffer (Ariake Sea, Age0)	11.6 (7.0)
Tiger puffer (Mie)	5.1 (6.5)
Kuruma prawn (Western Japan)	2.8 (4.5)
Mud crab (Urado Bay)	0.9 (0.7)
Abalone (Japan coasts)	12.2 (8.1)
Japanese scallop (Okhotsk Sea)	34.5 (10.2)
Sea urchin (Hokkaido)	18.2 (17.5)
Posterior mean	**8.2 (4.7)**

0 2.5 5 7.5 10 12.5 15 17.5 20 22.5 25 27.5 30 32.5 35 37.5 40

Economic efficiency

Species	Mean (SD)
Barfin flounder (Hokkaido)	2.7 (0.1)
Chum salmon (Hokkaido, Honshu)	6.8 (2.3)
Japanese flounder (Fukushima94_02)	0.9 (0.4)
Japanese flounder (Kagoshima Bay)	1.1 (0.1)
Japanese flounder (Miyako Bay)	1.6 (0.7)
Japanese Spanish mackerel (SIS)	1.0 (0.1)
Red sea bream (Kagoshima Bay)	5.0 (2.7)
Red sea bream (Sagami Bay)	1.4 (0.3)
Kuruma prawn (Western Japan)	0.7 (0.9)
Mud crab (Urado Bay)	1.9 (1.5)
Swimming crab (SIS)	1.0 (0.3)
Abalone (Japan coasts)	3.5 (2.4)
Japanese scallop (Okhotsk Sea)	17.9 (5.3)
Posterior mean	**2.7 (4.5)**

(a) Hokkaido scallop (1910–2018)

(b) Hokkaido chum salmon (1870–2018)