Dynamic macromolecular composition and high exudation rates in *Prochlorococcus*

Dalit Roth-Rosenberg[1], Dikla Aharonovich[1], Anne-Willem Omta[2], Michael J. Follows[2] and Daniel Sher[1]

[1] Department of Marine Biology, Leon H. Charney School of Marine Sciences, University of Haifa, Israel, [2] Department of Earth, Atmospheric and Planetary Sciences, Massachusetts Institute of Technology, USA

Abstract

Every living cell is composed of macromolecules such as proteins, DNA, RNA and pigments or cofactors. The ratio between these macromolecular pools depends on the allocation of resources within the organism to different physiological requirements, and in turn determines the elemental composition of the organism and, potentially, how it may affect biogeochemical cycles of elements such as carbon, nitrogen and phosphorus. Here, we present detailed measurements of the macromolecular composition of *Prochlorococcus* MIT9312, a representative strain of a globally abundant marine primary producer, as it grows and declines due to N starvation in laboratory batch cultures. As cells reached stationary stage and declined, protein/cell decreased by ~30% and RNA/cell and pigments/cell decreased by an order of magnitude. The decline stage was associated with the appearance of chlorotic cells which had higher forward scatter (a proxy for cell size) but lower chlorophyll autofluorescence, as well as with changes in photosynthetic pigment composition. Specifically, during culture decline divinyl-chlorophyll-like pigments emerged, which were not observed during exponential growth. These

divinyl-chlorophyll-like pigments were also observed in natural samples from the Eastern Mediterranean. Around 80-85% of the carbon fixed by *Prochlorococcus* MIT9312 (but not of a different strain, NATL2A) was released into the growth media as dissolved organic carbon under these laboratory conditions. Broadly defined, the macromolecular composition of *Prochlorococcus* MIT9312 is more similar to eukaryotic phytoplankton than to marine heterotrophic bacteria, suggesting a different set of physiological constraints determines the macromolecular composition of these two broad classes of marine microorganisms.

Keywords: Prochlorococcus, Photosynthetic pigments, Macromolecular pools, exudation, ribosomes

Introduction

Like every cell on Earth, phytoplankton and bacterioplankton are composed of several classes of biological macromolecules that function together to maintain the cells structure and activity (Geider and La Roche 2002, Finkel et al. 2016). Proteins are the most abundant macromolecule (in terms of mass), responsible for most of the metabolic functionality of the cells. DNA stores hereditary information, whereas RNA (as mRNA or other regulatory RNAs) accesses this stored data and (as rRNA and tRNAs) uses it to produce protein. Lipids form the membrane(s) that separate the cell from its environment and, in the case of photosynthetic organisms, host the molecular machinery used to harvest light for energy. This molecular machinery includes, in addition to proteins, also photosynthetic pigments and other metabolites. Finally, various storage molecules (including some lipids, carbohydrates, polyphosphate and other storage molecules) allow cells to retain energy and elements for future use, and a host of other metabolites are involved in all forms of cellular function. The relative amount of each of these classes of macromolecules changes between different phytoplankton groups (Vargas et al. 1998, Geider and La Roche 2002, Finkel et al. 2016), and may also change within a single type of organism in response to environmental conditions (Vargas et al. 1998, Liefer et al. 2019). For example, the relative amount of chlorophyll-a per cell changes in response to light intensity (e.g. (Moore et al. 1995)) and may also change in response to nutrient starvation (Lourenço et al. 1998, Liefer et al. 2019). This affects the ability of photosynthetic cells to harvest light and fix carbon, impacting cell physiology and likely the interactions of the cell with other organisms, e.g. through the release of fixed organic carbon (Dubinsky and Berman-Frank 2001). Similarly, it has been suggested that the number of ribosomes per cell is a strong determinant of cellular growth rate, and is itself affected by temperature and nutrient (particularly phosphorus, P) availability (e.g. (Elser et al. 2003, Garcia et al. 2016, Martiny et al. 2016)). Thus, to some extent, the macromolecular composition of the cell is affected by, and may shed light on, the allocation of cellular resources to different functions.

The basic building blocks of biological macromolecules such as amino acids and nucleic acids have distinct elemental composition, and the macromolecular composition of the cell is thus intimately linked with its elemental composition (Geider and La Roche 2002, Finkel et al. 2016, Garcia et al. 2016, Liefer et al. 2019). In phytoplankton, proteins generally comprise the largest pool in terms of cell biomass (dry weight) as well as nitrogen and carbon (N and C) pools (Lourenço et al. 1998, Vargas et al. 1998, Geider and La Roche 2002, Finkel et al. 2016). Other macromolecules such as Nucleic acids and pigments contribute less to total cell biomass yet contribute significantly to the pools of N and phosphorus (P) (Geider and La Roche 2002, Finkel et al. 2016, Liefer et al. 2019). Thus, the macromolecular composition of the cell will determine its elemental composition, which in turn will determine the effect of the cell on biogeochemical cycles of elements. Over the last ~85 years (Redfield 1934) significant advancements have been made in our understanding of how the elemental ratios in marine organisms change over time and space, how they are affected by cell physiology, and how they are related to surrounding nutrient element concentrations (e.g. (Martiny et al. 2013, Gruber and Deutsch 2014)). Significantly less is known about the dynamics of the macromolecular composition of marine organisms, which underlie elemental composition. Similarly, most of the models of ocean communities and their effect on biogeochemistry are formulated using elements as the model currency rather than macromolecules (Omta et al. 2009, Goebel et al. 2010, Talmy et al. 2016). In order to obtain a better understanding of the dynamics of the macromolecular composition of organisms, and to link these to elemental ratios, experimental measurements are needed, particularly in abundant marine organisms.

Here, we measure the cellular pools of major macromolecules (protein, DNA, RNA and pigments) in laboratory batch cultures of *Prochlorococcus*, a highly abundant pico-cyanobacterium which is responsible for ~8.5% of the oceanic photosynthesis (Partensky and Garczarek 2010, Flombaum et al. 2013, Biller et al. 2014). *Prochlorococcus*, as a clade, comprise multiple genotypes, each different in

its physiology, genome structure and oceanic niche (Johnson et al. 2006, Biller et al. 2014). Many

aspects of the physiology of the *Prochlorococcus* clade have bene intensively studied, ranging from its

elemental composition (Bertilsson et al. 2003, Martiny et al. 2013) through its photophysiology (Moore

et al. 1995, Moore and Chisholm 1999, Ting et al. 2002, Steglich et al. 2003, Komatsu et al. 2016) to

its system-wide transcriptomic response to changes in environmental conditions (e.g. (Martiny et al.

2006, Tolonen et al. 2006, Thompson et al. 2011, Aharonovich and Sher 2016). Yet, detailed analyses

of the macromolecular composition of the cells are lacking, even though changes in the elemental

composition of *Prochlorococcus* that are observed in response to environmental conditions are in fact

due to changes in the macromolecular composition of the cells (E.g. (Bertilsson et al. 2003, Van Mooy

et al. 2006, Martiny et al. 2013, Martiny et al. 2016)). The strain we focus on, MIT9312, represents the

high-light adapted HL-II clade which is the most abundant clade in the surface waters of large parts of

the ocean (Bouman et al. 2006, Johnson et al. 2006). We chose to follow batch cultures of

Prochlorococcus MIT9312 under conditions where entry into stationary stage occurs when the N

source in the media is depleted for several reasons. First, N stress affects *Prochlorococcus* in large

parts of the ocean (e.g.(Saito et al. 2014)), and significant information is available as to the

physiological, transcriptomic and evolutionary responses of the cells to acute N starvation (Steglich et

al. 2001, Moore et al. 2002, Tolonen et al. 2006, Gilbert and Fagan 2011, McDonagh et al. 2012, Read

et al. 2017, Berube et al. 2019, Szul et al. 2019). Second, long-term analyses of batch cultures may

identify cellular physiological processes not seen when cells are growing exponentially or exposed to

sudden or short-term nutrient starvation (e.g. (Christie-Oleza et al. 2017, Roth-Rosenberg et al. 2019)).

Indeed, it is an open question to what extent stationary-phase and culture decline in batch cultures

represent conditions found in nature, yet we show that a pigment identified in our cultures during the

culture decline stage is found in the oceans, raising the possibility that some cells in nature are indeed

found in these physiological stages. Finally, we measured, in addition to the cellular macromolecular

pools, also the concentrations of TOC in the media, extracting from these measurements estimates of

the release of DOC from the cells. This release may be due to passive exudation, active excretion and

cell lysis (reviewed by (Thornton 2014)). Some of the released macromolecules are available for

subsequent utilization by other organisms, whereas another part contributes to a huge pool of long-

lived oceanic DOM, potentially impacting global biogeochemical cycles (Hansell et al. 2009).

Materials and Methods

Growth conditions and experimental procedure

To follow the growth and macromolecular composition of *Prochlorococcus* over the various stages of

laboratory batch culture, axenic *Prochlorococcus* MIT9312 cultures were grown in 1 liter bottles of

Pro99 media (Moore et al. 2007) where the NH_4 concentration was lowered to 100µM, leading to

cessation of cell growth due to N starvation (Grossowicz et al. 2017). The experiment included 15

bottles, three of which were used for routine culture monitoring and the other 12 were used to collect

samples at four time-points, corresponding to exponential growth (t=6 days), early stationary stage

(t=10 days), late stationary stage (t=13 days) and culture decline (t=15 days). Samples were also

collected from the starter culture to represent t=0. The number and timing of the samples collected for

full analysis were selected in order to provide appropriate data for subsequent modeling (Omta et al.

2017). This experimental design was used to minimize the chance for contamination of the axenic

cultures with heterotrophic bacteria. The cell numbers of the cultures used for macromolecular analysis

were never statistically different from those of the cultures used for monitoring (Students t-test).

Prior to the beginning of the experiment, cells from a mid-exponential culture were counted by flow

cytometry (FACSCantoII, BD), and diluted in the new media to an initial cell density of 10^6/ml.

Cultures were maintained under constant light (22 µE), at 24.5 ±1°C. 1mM $NaHCO_3$ was supplemented

at three different time points (T=0, T=7, T=11 days) to verify that the cultures are not carbon-limited

(Moore et al. 2007, Grossowicz et al. 2017). Bulk culture fluorescence was measured in samples

collected aseptically from the three monitoring bottles using a Carey Eclipse spectrofluorometer (Ex

440nm/Em 680nm). Samples for cell counting were fixed in 0.25% glutaraldehyde, kept in the dark

for 10 minutes and transferred to a -80°C freezer for subsequent flow cytometry analysis.

Measurements of macromolecular pools and photosynthetic pigments.

At each of the five time-points selected, cells were collected by filtration. For protein measurements,

cells were gently filtered on 0.22µm polycarbonate filters and kept in -80°C until extraction. For protein

extraction, the filters were incubated in 500µl lysis buffer (50mM Tris pH6.8, 5mM EDTA, 2%SDS)

on a lab rotator at 37°C for 20'. Following incubation, samples were sonicated in water bath for 10',

centrifuged at 12,000g in a table-top micro-centrifuge for 5', and the supernatant liquid taken for

measurement using the Bicinchoninic acid assay (BCA, Sigma). Nucleic acids were collected similarly

on 0.22 µm Supor-200 Membrane Disc Filters (25 mm; Pall Corporation) and preserved in storage

buffer (40 mM EDTA, 50 mM Tris pH 8.3, 0.75 M sucrose) and RNA Save (Biological Industries) for

DNA and RNA measurements, respectively. Samples were kept at -80°C until extraction. DNA was

extracted from the filters according (Massana et al. 1997). Total RNA was extracted using the mirVana

miRNA kit (Ambion, Austin, TX, USA) as described in with lysozyme treatment for better lysis

(Tolonen et al. 2006), follow by removal of contaminating DNA from total RNA by Turbo DNase

(Ambion). Nucleic acids quantity measured using Qubit 2.0 (Invitrogen). For pigment analysis,

samples were collected on Glass Fiber filters (25 mm GF/F, Whatman) and kept in -80C until

extraction. Glass fiber filters were used, rather than polycarbonate filters, as the latter are not

compatible with many extraction methods for pigments. Nevertheless, preliminary experiments using

a culture of *Prochlorococcus* MED4, which is similar in size to MIT9312, showed that with the vacuum

pumps we used approximately 99.9% of the cells were retained on the GF/F membrane (Marmen et al.

2018), and thus the difference in pore size has minimal effect on the results. Cells were extracted in

100% methanol at 25°C for 2h. The cell extracts were clarified with Syringe filters (Acrodisc CR, 13

mm, 0.2 µm PTFE membrane, Pall Life Sciences), and kept at -20ºC until analysis by column

chromatography on a C8 column (1.7 µm particle size, 2.1 mm internal diameter, 50 mm column

length, ACQUITY UPLC BEH, 186002877) using an ACQUITY UPLC system equipped with a

Photodiode Array detector (Waters). The chromatography protocol was based on LOV method

(Hooker *et al.* 2005) modified for UPLC. A linear gradient was applied over 14 minutes, where solvent

A was 70:30 methanol:0.5 M ammonium acetate, and solvent B was B. 100% methanol. The flow rate

was 0.5 ml/min, and the column and injection heating were set to 50ºC and 30ºC, respectively.   To

identify and calculate pigments concentration, the following standards were used: Divinyl chlorophyll

A (DVchlA), chlorophyll B (chlB), chlorophyll C (chlC), zeaxanthin (Zea), α- carotene (α-car), Divinyl

Protochlorophyllide (MGDVP), Pheophorbide a, Pheophythin a and Chlorophyllide a. All standards

were from DHI, Denmark.

Measurements of inorganic nutrients and TOC

The GF/F filtrates (see above) were kept in polypropylene at -20ºC until analysis. For measurements,

samples were adjusted to 25ºC and diluted 4-fold in DDW to reduce the salt level. NH_4 (NH 3-N) was

measured using an HI 96715 meter (Hanna instrument), and PO_4 was measured by HI 96713 meter

(Hanna instrument). In both cases, low concentration kits were used. For Total Organic Carbon (TOC)

analysis, samples were collected into cleaned, pre combusted, acid washed (10% HCL) 40ml vials and

HCl was added to remove dissolved inorganic C before preservation at -20ºC. Before analysis, samples

were combusted at 680°C using catalytic (platinum) oxidation method in oxygen-rich environment.

TOC analysis was performed using TOC-L analyzer (Shimadzu ASI-L Autosampler, Columbia, MD).

For inorganic carbon (DIC), samples were collected in 14ml dark glass vials with screw caps and 75µl

(0.05% v/v) saturated $HgCl_2$ solution (Dickson et al. 2007) was added to the samples. For alkalinity,

60 ml samples were collected in glass vials. All samples were kept in 4°C until measurement. CO_2 was

extracted from the samples by acidification with phosphoric acid (H_3PO_4, 10%) using a custom,

automated CO_2 extractor and delivery system (AERICA by MARIANDA) using high grade N_2 as a

carrier gas, connected online with a LiCor 6252 IR CO_2 analyzer. Measurements were calibrated using

seawater Certified Reference Materials (CRMs) from Dickson's lab. In order to check and correct for

drift the CRM was run after every 4 samples.

Model of *Prochlorococcus* resolving key macromolecular pools

We formulated a mathematical model of the growth of a generalized axenic phytoplankton cell,

which resolves the macromolecular pools measured in the experiments – protein, RNA, DNA and

photosynthetic pigments, as well as N storage and dissolved organic carbon (Figure S3). The

model is formulated in units of N in each macromolecular pool, with the C in each pool determined by

the elemental ratios in each type of macromolecule (Geider and La Roche 2002). In this model, proteins

act as enzymes in the synthesis of RNA, DNA and Chl, whereas RNA acts as an enzyme for the

synthesis of proteins. The total organic carbon in the media (TOC, the sum of cellular C and DOC) is

determined by the balance between photosynthesis and respiration. Since the cells are defined in terms

of N, and the TOC is defined in units of C, the concentration of DOC is an emergent property of the

model, and enables us to test under what conditions the model can reproduce high DOC production

rates. We fit the model to the experimental data presented below, using the Metropolis-Hastings

algorithm (Metropolis et al. 1953). A more detailed description of the model, including equations, can

be found in the supplementary information, and the model itself (written in FORTRAN) is available

on https://github.com/AWO-code/DalitDaniel_Model.

Results

Dynamics of cell numbers and macromolecular composition during growth and nitrogen starvation

To determine to what extent the macromolecular composition of *Prochlorococcus* changes between the different physiological states of a batch culture, we grew strain MIT9312 in laboratory batch cultures where the N:P ratio of the media was set to 2, thus leading to cessation of growth due to N starvation (Grossowicz et al. 2017). The bulk culture fluorescence, often used to monitor phytoplankton growth in a non-invasive way, increased exponentially until day 10, after which the culture fluorescence declined rapidly, with no clearly observable stationary stage (Fig 1a). The decline of the culture fluorescence coincided with the reduction of soluble NH_4 to below detection threshold (<10uM), while PO_4 and DIC levels remained high throughout growth and decline (Fig 1c, d). Cell counts by flow cytometry also showed an increase in the cell numbers until day 10, with a growth rate (μ) of 0.353 day^{-1} (±0.0239) and doubling time of 1.968 days (±0.129). However, unlike the bulk culture fluorescence that decline rapidly after this day, the cell numbers remained relatively stable for an additional 3-4 days, before starting to decline. During this period, a clear population of cells with low chlorophyll autofluorescence (low-fl) emerged, identified by flow cytometry (Fig 1b, 67% low-fl cells on day 13 and 88% on day 15, (Roth-Rosenberg et al. 2019)). These low-fl ("chlorotic") cells were also stained much more weakly than the high-fl ones with the dye Sybr Green, which binds nucleic acids (primarily dsDNA but also ssDNA and RNA, Fig S1b). At the same time, an increase in Sybr-Green staining was observed in the high-fl cells, potentially related to the arrest of the cells in G2 stage (Sup Fig. S1, (Zinser et al. 2009)). At the same time, the median forward scatter, a proxy for cell size, increased for both the high-fl and low fl cells. (Fig S1c).

Based on the observed fluorescence curves (Fig 1a), we sampled cells during exponential stage (day 6), early stationary stage (day 10), late stationary stage (after bulk culture fluorescence had started to

decline but while cell numbers were still stable, day 13) and culture decline (day 15) (Table S1). An additional estimate of the macromolecular pools from exponentially-growing cells was obtained from the starter cultures used to start the experiment (treated here as day 0). The per-cell content of proteins and RNA were relatively stable during exponential phase (0-10 days), but was reduced during the stationary and decline phases (days 13 and 15) (Fig 1e-f). The reduction in RNA/cell (a drop of approximately 75% between days 10 and 13) was much larger than that of protein/cell (10-30%). The total photosynthetic pigment concentration per cell started declining earlier than RNA or protein, with a drop of approximately 75% between days 6 and 10, as the cells moved from exponential growth to early stationary stage (Fig 1g).

Figure 1: Dynamics of culture growth, plateau and death in batch *Prochlorococcus* MIT9312 cultures reaching stationary growth due to nitrogen starvation. (a) Cell numbers and

fluorescence curves of MIT9312 population shows a log-phase growth until day 10 than stationary and decline of the culture. (a) Flow cytometry scatter gram of specific measurements points. (c-d) Analysis of external inorganic nutrients shows that NH_4^+ drops below detection limit, while PO_4 and DIC are still available. Thus, the cultures enter stationary stage due to N starvation. Results are means and ranges of triplicate cultures. (e-g): Changes in the cell quotas of major macromolecules-DNA and RNA (e) proteins (f) and total pigments (g). While protein/cell is relatively stable, pigment and RNA pools drop as cells become starved and the culture declines. (h) Accumulation of large amounts of DOC. Triangles show measured Total Organic Carbon (TOC), squares show calculated C in cell biomass, based on flow cytometry counts and on per-cell C quotas. The error bars shown represents uncertainty due to the differences in estimates of per-cell C quotas between studies (Table 1), with the squares showing a value of 50 fg cell^{-1}.

Changes in photosynthetic pigment composition across the different growth stages

The reduction in chlorophyll autofluorescence per cell suggests changes to the photosynthetic machinery of the cells as they enter stationary stage and during culture decline. Indeed, as shown in Fig 2a, in addition to a reduction in the total pigment quota per cell (Fig 1g), the ratio of some accessory pigments to divinyl chlorophyll A (DVchlA) changed over time. Chlorophyll B and a chlorophyll-C like pigment were reduced compared to DVchlA after day 10, whereas the ratio of zeaxanthin (and also α-carotene) to DVchlA increased after this day (Fig 2b). In addition, on day 10 and later, UPLC several peaks were observed close to that of DVchlA that differed from it in their retention time. These peaks were not due to overloading of the UPLC column (were not seen when higher amounts of pigments were injected),and were reproducibly observed in other cultures from multiple

Prochlorococcus strains. One of these peaks (marked with an arrow in Fig 2c), which was especially

prominent, had an absorption spectrum similar to that of DVchlA (Fig 2d) but was not one of the known

degradation products (pheophytin-A eluted later, whereas phaeophorbide A, chlorophyllide A and

divinyl protochlorophillide A all eluted much earlier, consistent with the loss of the phytol moiety).

Assuming a similar molar absorption rate as DVchlA, this pigment (which may be a DVchlA', an

epimerization product of DVchlA, (Komatsu et al. 2016)) comprised as much as 21% of the total

DVchlA pigments on day 15, at the end of the decline stage. Importantly, pigments with the same

elution time and absorbance were repeatedly observed in samples collected from the oligotrophic

Eastern Mediterranean Sea, as demonstrated in Fig S2.

Figure 2. Pigments content and relative composition change during growth. (a) Changes in the major per-cell pigment quotas. Divinyl chlorophyll A - DVchlA, chlorophyll B - chlB, a chlorophyll C-like pigment - chlC , zeaxanthin - Zea, and α- carotene - α-car. (b) The relative abundance of zea to DVchlA increases, while that of chlB to DVchlA decreases. (c) UPLC chromatograms at different times show the appearance of additional pigments during stationary phase. The pigment marked with an arrow has an absorption spectrum similar to that of DVchlA (panel d), but differs from it in its retention time.

Reproducibly high rates of DOC exudation in *Prochlorococcus* MIT9312.

Part of the organic carbon fixed by phytoplankton is released from the cell due to exudation and cell mortality, and this organic carbon provides sustenance for co-occurring heterotrophic bacteria. To assess how much organic C is released by *Prochlorococcus*, we measured the total organic C in our cultures, and compared it to the amount of C expected to be in the particulate fraction based on the number of cells and the carbon quota of *Prochlorococcus* from previous studies (10-130 μg/cell, Table 1). During exponential stage (day 6), the measured TOC was within the range expected based on cell numbers and C quota, however, during the stationary and decline phases the amount of TOC was up to 23-fold higher than could be explained by cell biomass, suggesting a large amount of organic C was released into the growth media (Fig 1g). To check if the high release of organic carbon is common to different *Prochlorococcus* strain, we repeated this experiment using both strain MIT9312 and strain NATL2A which belongs to the Low Light I clade (Biller et al. 2014). High extracellular concentrations of organic C (up to 13-fold higher than the predicted cellular biomass) were observed again for

271 MIT9312, whereas strain NATL2A that grew under the same N starvation conditions showed TOC

272 values much lower than MIT9312, within the expected range of cell quota (Fig 3).

274 **Table 1: *Prochlorococcus* Carbon biomass estimated in different studies, expanded from**

275 (Bertilsson et al. 2003)

Strain	fg C cell^{-1}	Analytical method	Reference
Natural population, Atlantic Ocean	10-70	X ray microanalysis	(Grob et al. 2013)
PCC9511, EQPAC, SB, GP2, SARG, NATL1	17-34	X ray microanalysis	(Heldal et al. 2003)
PCC9511	17-38	POC	(Claustre et al. 2002)
MED4	17-40	POC	(Fu et al. 2007)
VOL7(MED4), VOL8(MIT9515), VOL29, VOL4 (MIT9312), VOL1(MIT9215), UH18301	18-55	POC	(Martiny et al. 2016)
MIT9301, MED4, NATL2A, MIT9313	30-33 (HL), 45 (LLI), 80 (LLIV)	Macromechanical mass sensor, assuming C is 50% of wet biomass	(Cermak et al. 2016)
MIT9301, MED4, MIT9313	44-51 (HL), ~180 (LLIV)	POC	(Becker et al. 2014)

MED4	46-61	POC	(Bertilsson et al. 2003)
CCMP1378(MED4)	49+/-9	POC	(Cailliau et al. 1996)
Generic (0.6 µm diameter cell)	53	Cell size and volume-carbon conversion (470 fg C/µm^3)	(Campbell et al. 1994)
Natural population, Atlantic Ocean	56	Cell size and volume-carbon conversion (325 fg C/mm3)	(DuRand et al. 2001)
Generic (0.8 µm diameter cell)	59	Cell size and volume-carbon conversion (220 fg C/mm3)	(Li et al. 1992)
MIT9211, MIT9215, MIT9312, MED4, SS120, MIT9302, MIT9303, MIT9313	61-94	Primary production, potential overestimation (does not account for respiration and exudation)	(Moore 1997)
MIT9312	67-123	Macromolecular pools	This study
MED4, MIT9401, SS120	78+/-27	Dry weight, assuming C is 50% of biomass	(Shaw et al. 2003)
Generic (0.8 µm diameter cell)	124	Cell size and volume-carbon conversion (294 fg C/cell)	(Veldhuis and Kraay 1990)

Figure 3: Differences in exudation rates between two axenic *Prochlorococcus* strains, MIT9312 (a) and NATL2A (b). The two strains were grown under identical conditions. Triangles show measured Total Organic Carbon (TOC), squares show calculated C in cell biomass, based on flow cytometry counts and on per-cell C quotas. The range shown represents uncertainty due to the differences in estimates of per-cell C quotas between studies (Table 1), with the square markers showing a value of 50 fg cell⁻¹.   Panel a shows an independent experiment from that shown in Fig 1h, yet the high release of DOC is consistent.

282   **Figure 3: Differences in exudation rates between *Prochlorococcus* strains.**

284   **Discussion**

285   Quantitative measurements of cellular parameters, including macromolecular composition and

286   elemental quotas, are fundamental to our understanding (and modeling) of microbial systems.

However, measurements of the per-cell C quota of *Prochlorococcus*, performed using different techniques, range over an order of magnitude (Table 1, (Bertilsson et al. 2003)). In this study, we directly measured the cell quotas of DNA, RNA and photosynthetic pigments at different physiological stages of laboratory batch culture in *Prochlorococcus* MIT9312. With some assumptions, these measurements can be used to constrain upper and lower bounds of the total cell quotas of C, N and P (Supplementary text, Supplementary Table 1). Measuring these macromolecular pools requires less biomass and can be performed at higher throughput compared to measurements of total particulate or dissolved C and N (Supplementary text). Assuming a lower bound on the C:N ratio of 6 (Bertilsson et al. 2003, Martiny et al. 2013, Martiny et al. 2016), the cellular C quota we estimate (~67-123 fg cell^{-1}, Supplementary Table 1) is at the upper range of measurements from other studies (Table 1), and is driven primarily by the high (and mostly stable) measurements of protein/cell. Like any other experimental result, these measurements are sensitive to the experimental design and the methodology used. For example, while the method we used for protein determination (BCA) is relatively insensitive to the effects of salts and detergents (if these are incorporated into the standard curve), the quantitative results can be affected by, for example, the protein standards used (e.g. Bovine Serum Albumin or Immunoglobulin). Similarly, our estimates of DNA/cell are about one-third those expected from cell number and the genome size of MIT9312 (1.7Mbp, (Kettler et al. 2007)). Experimental measurements of DNA/cell that are not consistent with genome size have been observed also in other studies (Zimmerman et al. 2014), and may be caused by differences in the DNA extraction efficiency across organisms or culture conditions. Finally, the *Prochlorococcus* population is not homogenous, containing at times two distinct sub-populations of cells, with different flow cytometry signatures (as discussed in detail below, Fig S1, see also (Coe et al. 2016, Roth-Rosenberg et al. 2019)). We did not measure the amount of the macromolecular pools in each sub-population separately, due to the difficulty in obtaining sufficient biomass from non-fixed cells sorted by fluorescence-activated cell

sorting. With these uncertainties in mind, in the following sections we use the measurements of macromolecular pools at different growth stages, as well as the estimates of cellular C, N and P, to discuss the relationship between cell physiology, macromolecular pools and elemental budgets. Doing so, we note the need for an in-depth comparison of measurements between labs, growth stages and methods, perhaps along the lines of similar studies used to calibrate pigment measurements (e.g. (Claustre et al. 2004)).

Dynamic changes in the macromolecular composition of *Prochlorococcus* in batch culture – changes in the average cell or sub-populations?

In our laboratory batch cultures, *Prochlorococcus* grow exponentially until they run out of their nitrogen source (NH_4), at which stage they stop growing, enter a short stationary stage and finally die. During this time, the cultures lose their green color, the cell population became heterogeneous, and populations of low-fl cells emerge, forming the majority of the population as the cultures decline (Fig 1a, Fig S1a). The low-fl cells also stain weaker with the dye Sybr-green (Fig S1), which binds both DNA and RNA (albeit with a higher affinity for DNA, (Martens-Habbena and Sass 2006)). We propose that a significant part of the changes in the mean per-cell concentration of chlorophyll and RNA is caused by the presence of these low-fl cells, rather than by a gradual change in the concentration of these pools per cell. In the model cyanobacteria *Synechococcus elegantus* PCC7942 and *Synechocystis* PCC6803, the loss of culture chlorophyll (chlorosis) is part of a developmental program to generate resting stages, that can survive long-term starvation (e.g. (Sauer et al. 2001, Klotz et al. 2016)). In contrast, we have recently shown that *Prochlorococcus* cultures that become chlorotic cannot revive when nutrients are added, unless co-cultured with a heterotrophic bacterium (Roth-Rosenberg et al. 2019). Nevertheless, some of the chlorotic cells are still active (photosynthesize and take up NH_4,

(Roth-Rosenberg et al. 2019)). Thus, we interpret the changes observed in the macromolecular pools

of the MIT9312 cultures, and primarily those that can be attributed to the low-fl (chlorotic) population,

as physiological responses to N starvation that nevertheless are not sufficient to enable the cells to

survive long-term nutrient stress.

In parallel to the observed loss of culture fluorescence and the appearance (and dominance) of low-fl

cells, changes were also in the per-cell concentration and composition of the photosynthetic pigments

(Fig 1h, Fig 2). Specifically, the mean relative amount of zeaxanthin increases compared to DVchlA,

whereas the relative amount of chlorophyll B decreases (Fig 2b). Similar changes were observed in

nutrient-replete batch cultures of other *Prochlorococcus* (strains MED4 and SS120) in response to

increasing light intensities (Moore et al. 1995). One possible interpretation, which we deem less likely,

is that the changes in the pigment ratio as the cells enter stationary phase and start declining are due to

increased light levels as many of the cells become chlorotic and the cultures become clear. An

alternative interpretation is that the increase in relative abundance of zeaxanthin is a general stress

response, as these pigments may act as antioxidants in cyanobacteria (Zhu et al. 2010). Indeed, an

increase in the zeaxanthin-DVchlA ratio was observed upon nitrogen starvation also in an additional

Prochlorococcus strain, SS120 (Steglich et al. 2001). The decrease in chlorophyll B can also be

interpreted independently from a potential change in light levels. The reduction in DVchlB could

simply be related to the deactivation of the photosystems, in agreement with previous studies that

showed a decrease in photosynthetic activity (e.g. Fv/Fm) and psbA protein levels upon nitrogen

starvation (Steglich et al. 2001, Tolonen et al. 2006). Alternatively, the reduction in DVchlB may be

part of a cellular mechanism to reduce the use of N-containing pigments under N starvation, as

chlorophylls contain nitrogen, whereas xanthophylls do not. This is consistent with other mechanisms

whereby *Prochlorococcus* conserve N, including changes in the transcriptional profiles (Tolonen et al.

2006), use of shorter transcripts (Read et al. 2017) and, potentially, "thrifty" N use in proteins expressed

in response to N starvation (Gilbert and Fagan 2011).

The emergence of the DVchlA-like pigment is currently unexplained, although this pigment may be

an epimerization product of DVchlA (DVchlA'). DVchlA' has been previously observed in several

Prochlororoccus strains (Komatsu et al. 2016), and in our study the DVchlA-like pigment is associated

with culture decline. Generally speaking, harvesting cells from late exponential stage or early

stationary stage is a common practice when high biomass is required, yet as discussed above the

physiology of the cells can change during stationary phase in comparison with exponentially-growing

cells. Additionally, it is currently unclear to what extent processes observed in late stages of laboratory

batch cultures, such as nutrient starvation and the subsequent changes in cell physiology, occur in

nature. Nevertheless, the observation that a pigment with the same retention time and absorption

spectrum as the DVchlA-like pigment observed at-sea suggests that at least some of the cells in nature

may be undergoing processes similar to those we observe in the laboratory batch cultures.

The high-fl cells also exhibit an increase in mean forward scatter, a proxy for cell size. This occurs

despite the overall reduction in RNA, pigments and (to a lesser extent) protein (Fig 1e, f), which are

the major N-containing macromolecular pools. Assuming the increase in forward scatter represents an

increase in cell biomass, this suggests that the cells are accumulating primarily C-rich macromolecules

such as storage carbohydrates and lipids, leading to an increased C:N ratio. A similar response to

nitrogen starvation has been predicted by mathematical models of N-starved cells (Grossowicz et al.

2017) and observed in other phytoplankton species (Hu et al. 2008, Breuer et al. 2012, Liefer et al.

2019). If these processes occur in nature, they could impact the C:N ratio of DOM released from dead

Prochlorococcus cells (e.g. (Agusti and Sanchez 2002, Llabrés et al. 2011, Ribalet et al. 2015)) as well

as POM exported by these cells (Richardson and Jackson 2007, Lomas and Moran 2012, Zhao et al.

2017).

The macromolecular composition of *Prochlorococcus* MIT9312 is more akin to eukaryotic

phytoplankton than to heterotrophic bacteria

Phytoplaknton and heterotorphic bacteria are inherently different in their life histories, and this likely

determines and constrains the relative allocation of energy and elements to different molecular

functions. Phytoplankton need to allocate resources to the photosynthetic and carbon fixation

machineries and are likely less constrained by the availability of carbon (which can be derived from

photosynthesis) compared to other elements such as nitrogen. Eukaryotic phytoplankton are also often

larger than bacteria, and thus are less constrained by the internal volume of the cell. Heterotrophic

bacteria, in contrast, do not need to allocate resources to photosynthesis but are potentially limited by

C and by cell size. As shown in Fig 4, the limited number of measurements of macromolecular pools

in marine microorganisms available in the literature supports the idea phytoplankton and bacteria

allocate resources differently.

Generally speaking, phytoplankton have a lower RNA/protein ratio than heterotrophic bacteria with

the same RNA/DNA ratio. Assuming that the ribosomes of phytoplankton and bacteria have similar

maximal rates of protein production (chain elongation), this suggests that phytoplankton ribosomes are

working at a higher relative capacity. In phytoplankton, a significant amount of protein production

needs to be invested in maintaining the photosynthetic apparatus, for example in the replacement of

the core proteins (Zavřel et al. 2019). An alternative (but non-exclusive) explanation is that

heterotrophic bacteria (primarily copiotrophic ones) maintain more ribosomes than needed, using the

"spare" production capacity to allow rapid response to changes in growth conditions. Such a strategy

has been demonstrated experimentally for *E. coli* (e.g. (Li et al. 2018). Notably, some bacteria, such as

Psychrobacter Mor119, *Halomonas Hal146*, *Vibrio Vib2d* and *Ruegeria Oce241*, are more similar in

the RNA/protein ration to phytoplankton (Fig 4). The reason for this similarity, which may represent a

lower ability to allocate ribosome resources to rapid changes in gene expression, is unclear, and may

depend on the specific experimental conditions employed in their study (Zimmerman et al. 2014).

While *Prochlorococcus* is a bacterium (prokaryote), its RNA/protein and RNA/DNA ratios at different

stages of the growth curve all fall within the range of those from eukaryotic phytoplankton. The single

measurement form *Synechococcus* is also closer to other phytoplankton (Fig 4). This could be the result

of both the need to invest in maintaining the photosystem and, potentially, that *Prochlorococcus* may

have a lower ability compared to fast-growing heterotrophs to "ramp up" their translation in order to

rapidly respond to changes in environmental conditions.

Nevertheless, *Prochlorococcus* may still maintain some unused ribosome capacity. In our study,

protein/cell remained relatively stable, with a decline of no more than ~30% as the culture declined.

RNA/cell, in contrast, dropped by more than 80% over the same period (Fig 1). The decline in

RNA/cell also started earlier. This suggests either that processes resulting in the loss of protein (e.g.

degradation, exudation or excretion) were strongly reduced in stationary and decline phase cells, or

that the actual rate of protein production per ribosome increased in declining cultures compared to

exponentially-growing ones. These two explanations - decrease in loss processes or increase in

ribosome efficiency - are not mutually exclusive, but evidence is lacking for either of them in most

organisms, including *Prochlorococcus*.

Figure 4. RNA/DNA and RNA/protein compared to those of other phytoplankton and bacteria.

Figure 4. RNA/DNA and RNA/protein compared to those of other phytoplankton and bacteria.

The ratios in phytoplankton (from (Finkel et al. 2016)), presented in cross markers , show a high

diversity of RNA/DNA, while relatively low diversity of RNA/Proteins compositions. Heterotrophs

bacteria however (squares) are highly diverse in both RNA/Proteins and RNA/DNA composition

(Churchward et al. 1982, Neidhardt et al. 1990, Zimmerman et al. 2014). *Prochlorococcus* MIT9312

at different growth phases (spheres of different sizes) are in fact similar to microalgae rather to

bacterial macromolecule composition. The most phylogenetically related cyanobacteria,

Synechococcus WH7803, marked in circle, shows similarity to MIT9312 in the RNA/DNA but not

in the RNA/Protein ratios.

DOC accumulation and its potential mechanism

One of the surprising observations of this study was the high rate of accumulation of DOC during the

exponential growth of MIT9312. By the time the cultures reached stationary stage (day 13 in Fig 1h

and day 11 in Fig 3a) the particulate organic carbon (i.e. cell biomass) was only 3-7% of the total

organic carbon (considering the two experiments shown in Fig 1h and 3h, and assuming a cell C quota

of 120 fg cell^{-1}). This suggests that 80-85% of the organic carbon fixed by *Prochlorococcus* in two

separate experiments was released as dissolved organic carbon. Pervious experimental studies where

DOC release was measured using ^{14}C as a tracer suggested that 2-24% of the primary productivity is

released (Bertilsson et al. 2005, Lopez-Sandoval et al. 2013). However, indirect assessments suggest

that a somewhat higher fraction of the fixed carbon is lost from the cells. First, a comparison of carbon

uptake through photosynthesis and growth rate in *Prochlorococcus* MED4 over a diel cycle suggested

that ~30% of the fixed carbon is released or respired (Zinser et al. 2009). Second, the theoretical growth

rate of MIT9312 based on the expected yield of photosynthesis (taking into account published values

for illumination, cross section, efficiency and chlorophyll/cell, (Moore and Chisholm 1999)) is almost

two-fold higher than measured (expected μ=0.64 day^{-1}, compared to 0.35±0.02 day^{-1} measured during

exponential stage, see Supplementary Information for more details). Third, we used a mathematical

model of a generalized phytoplankton, resolving the macromolecular pools measured in this

experiment, to test whether the exudation rates we observed are feasible. The model was able to

reproduce quite well some of the experimental results, including the accumulation of DOC (observed

as an increase in TOC, Figs S3-S5, Supplementary Table S2). This model reproduced well the cell

density, the uptake of inorganic N and the reduction after stationary stage in the cellular quotas of RNA

and chl-a, although the timing in the decline of the protein quota per cell was predicted to be earlier

than observed experimentally.   Importantly, while the values of C fixation per unit chl-a in most of

the model simulations were within the range of the maximum photosynthesis measured for

Prochlorococcus (up to 10-20mol C[g divinyl Chl-a]$^{-1}$day^{-1}, (Moore and Chisholm 1999, Bruyant et

al. 2005, Felcmanová et al. 2017)), this value had to vary over the dynamics of batch culture for the

model to reproduce well the accumulation of TOC, increasing transiently as the cells entered the

decline phase (Fig S5). This phenomenon – an increase in photosynthetic efficiency in cells entering

chlorosis (potentially caused by a reduction in shelf-shading, (Felcmanová et al. 2017)) - has not, to

the best of our knowledge, been described previously. An alternative explanation is that the respiration

rates decrease in relation to growth rate as the cells enter chlorosis (Fang et al. 2019). The current

model formulation cannot distinguish between these two possibilities, and thus experimental

measurements of photosynthesis and respiration across batch culture are required to test these two

alternative model predictions. Taken together, these considerations show that while the rates of DOC

release recorded here are 2-3 fold higher than those demonstrated experimentally (e.g. using [14]C release

assays), they are within the range of growth rates possible based on theoretical considerations.

The differences between our results, based on accumulation of DOC over time, and estimates of

exudation based on [14]C partitioning between the particulate and dissolved phases, may also be due to

loss of volatile substances during the acidification and subsequent venting of inorganic [14]C during

radioactive incorporation experiments. Indeed, *Prochloroccoccus* produce several volatile substances

(e.g. organic acids, organohalogens, methane and isoprene, (Shaw et al. 2003, Bertilsson et al. 2005,

Hughes et al. 2011, Bižić-Ionescu et al. 2019)), although to what extent these substances would be lost

from the media during the protocol for measuring [14]C incorporation is unclear. Thus, our results

showing high release of DOC are consistent with theoretical calculations and (some) previous studies,

but also highlight the need for additional studies, specifically addressing DOC release using different

analytical methods.

The DOM released by *Prochlorococcus* can include, in addition to C, also other elements such as N

and P. While we did not measure this experimentally, a rough estimate of the loss processes N and P

can be obtained by comparing the total amount of N and P in the original media and the estimated

amount of N and P in the cells at their maximum density. Since *Prochlorococcus* do not fix dinitrogen

or perform denitrification, the N "missing" from the cell biomass when reach stationary stage under

N-starved conditions should represents organic N that were lost from the cells in a format that is not

available for re-uptake (i.e. refractory to *Prochlorococcus*). At the peak of the *Prochlorococcus* growth

in our experiment, on day 10, the cellular N accounted for 79% of the total N in the media. Given that,

on that day, NH_4 in the media was depleted, this suggests that ~20% of the N had already been released

from the cells in a form that is not available for re-utilization. Thus, while C rich, the DOM released

by *Prochlorococcus* MIT9312 contains significant amounts of other elements, likely (at least in part)

in the form of proteins, amino acids, DNA, RNA and nucleotides. This is consistent with studies in

Synechococcus (Christie-Oleza et al. 2015, Christie-Oleza et al. 2017, Zhao et al. 2017) and with the

effect of *Prochlorococcus*-derived DOM on gene expression in natural communities (Sharma et al.

2014). With regard to P, a similar calculation shows that only ~3.2% of the total P in the media was

found in cellular biomass, while the PO_4 concentration in the media dropped from 50μM to 34-36μM,

a reduction of ~30%. Thus, either we have significantly underestimated total cellular P (i.e. there is an

additional, unmeasured pool besides RNA, DNA and phospholipids), or a significant amount of P has

been converted from PO_4 to extracellular DOP.

While our understanding of biogeochemistry is framed in terms of elements, cells actually release the

same macromolecules from which they are built, e.g. the proteins, DNA and RNA measured in this

study. Previous studies have suggested that *Prochlorococcus* release, in addition to the volatile

compounds described above, also TEP (Iuculano et al. 2017), and that at least part of the produced

DOM (the hydrophobic fraction that can be analyzed following solid-phase extraction) differs from

strain to strain, and even between replicate cultures (Becker et al. 2014). *Prochlorococcus* have also

been shown to release vesicles containing proteins and nucleic acids (Biller et al. 2014). Furthermore,

DOC loss from cultures can be due not only to passive leakage from the cells (exudation) or active

release (excretion) but also to lysis of dead cells (reviewed by (Thornton 2014)), with the latter process

likely releasing most or all of the cellular macromolecular content. Several studies have indicated that

up to ~20% of the cells in exponentially-growing *Prochlorococcus* cultures may be dead (Agusti and

Sanchez 2002, Hughes et al. 2011). Therefore, it is likely that all of these processes (exudation,

excretion, vesicle production and mortality) contribute to DOM formation.

Several studies have suggested that high rates of DOC release occur in oligotrophic regions of the

oceans, and that DOC, rather than POC, contributes significantly to carbon export from the photic zone

in these areas (Guyennon et al. 2015, Roshan and DeVries 2017). It is tempting to speculate that

Prochlorococcus may be at least partially responsible for this DOC accumulation, based on the high

rates of DOC production observed here by *Prochlorococcus* MIT9312, a member of the HL-II clade

that numerically dominates large areas of the oligotrophic ocean (Bouman et al. 2006, Johnson et al.

2006). In support of this speculation, DOM released by *Prochlorococcus* in the surface waters of the

Pacific Ocean (which was dominated by the HL-II clade) has been suggested to provide as much as

75% of the daily organic carbon produced (Ribalet et al. 2015), although it is unclear to what extent

mortality compared to exudation feeds this organic pulse. It has also been suggested that DOM derived

from *Prochlorococcus* contributes a significant amount to deep sea DOM (Zhao et al. 2017). We note,

however, that DOC production rates may differ based on the physiological states of the cells and the

surrounding nutrient conditions. For example, similar to our study, Bertilsson and co-workers also

noted differences between two strains of *Prochlorococcus*, MIT9312 and MED4, with the former

releasing approximately twice as much DOC (Bertilsson et al. 2005), and Becker and co-workers

documented differences in the composition of released DOM (Becker et al. 2014). Future work is

required to determine the to what extent DOM production changes between strains and across culture

conditions, what are the mechanisms of this process, and to what extent assessments of DOM release

under lab conditions can be extended to the ocean.

Acknowledgements

We thank Hila Elifantz, Ilana Berman-Frank, Eyal Geisler and Edo Bar-Zeev for the TOC analyses and Jack Silberman for the DIC analysis. This study was supported by the Gordon & Betty Moore Foundation (grant number GBMF #3778, to MJF), by the Human Frontiers Science Program (grant number grant RGP0020/2016, to DS) and by the United States-Israel Binational Science Foundation (Grant number 2010183 to DS and MJF and grant number 2016532 to DS).

Author contributions

DR, DA, AWO, MJF and DS conceived the study and designed the experiments. DR and DA performed the experiments and analyzed the samples. AWO performed the modelling. DR and DS wrote the manuscript with contributions from all authors.

Competing interests

The authors declare no competing interests

Materials and Correspondence

Please send requests for materials or other correspondence to Daniel Sher, dsher@univ.haifa.ac.il

References

Agusti, S. and M. C. Sanchez (2002). "Cell viability in natural phytoplankton communities quantified by a membrane permeability probe." Limnology and Oceanography **47**(3): 818-828.

Aharonovich, D. and D. Sher (2016). "Transcriptional response of Prochlorococcus to co-culture with a marine Alteromonas: differences between strains and the involvement of putative infochemicals." ISME J **10**(12): 2892-2906.

Becker, J. W., P. M. Berube, C. L. Follett, J. B. Waterbury, S. W. Chisholm, E. F. DeLong and D. J. Repeta (2014). "Closely related phytoplankton species produce similar suites of dissolved organic matter." Frontiers in Microbiology **5**.

Bertilsson, S., O. Berglund, D. M. Karl and S. W. Chisholm (2003). "Elemental composition of marine Prochlorococcus and Synechococcus: Implications for the ecological stoichiometry of the sea." Limnology and Oceanography **48**(5): 1721-1731.

Bertilsson, S., O. Berglund, M. J. Pullin and S. W. Chisholm (2005). "Release of dissolved organic matter by Prochlorococcus." Vie Et Milieu-Life and Environment **55**(3-4): 225-231.

Berube, P. M., A. Rasmussen, R. Braakman, R. Stepanauskas and S. W. Chisholm (2019). "Emergence of trait variability through the lens of nitrogen assimilation in Prochlorococcus." eLife **8**: e41043.

Biller, S. J., P. M. Berube, J. W. Berta-Thompson, L. Kelly, S. E. Roggensack, L. Awad, K. H. Roache-Johnson, H. Ding, S. J. Giovannoni, G. Rocap, L. R. Moore and S. W. Chisholm (2014). "Genomes of diverse isolates of the marine cyanobacterium Prochlorococcus." Scientific Data **1**.

Biller, S. J., P. M. Berube, D. Lindell and S. W. Chisholm (2014). "Prochlorococcus: the structure and function of collective diversity." Nature Reviews Microbiology **13**: 13.

Biller, S. J., F. Schubotz, S. E. Roggensack, A. W. Thompson, R. E. Summons and S. W. Chisholm (2014). "Bacterial vesicles in marine ecosystems." Science **343**(6167): 183-186.

Bižić-Ionescu, M., T. Klintzsch, D. Ionescu, M. Y. Hindiyeh, M. Günthel, A. M. Muro-Pastor, W. Eckert, F. Keppler and H.-P. Grossart (2019). "Widespread methane formation by Cyanobacteria in aquatic and terrestrial ecosystems." bioRxiv: 398958.

Bouman, H. A., O. Ulloa, D. J. Scanlan, K. Zwirglmaier, W. K. Li, T. Platt, V. Stuart, R. Barlow, O. Leth, L. Clementson, V. Lutz, M. Fukasawa, S. Watanabe and S. Sathyendranath (2006). "Oceanographic basis of the global surface distribution of Prochlorococcus ecotypes." Science **312**(5775): 918-921.

Breuer, G., P. P. Lamers, D. E. Martens, R. B. Draaisma and R. H. Wijffels (2012). "The impact of nitrogen starvation on the dynamics of triacylglycerol accumulation in nine microalgae strains." Bioresource Technology **124**: 217-226.

Bruyant, F., M. Babin, B. Genty, O. Prasil, M. J. Behrenfeld, H. Claustre, A. Bricaud, L. Garczarek, J. Holtzendorff, M. Koblizek, H. Dousova and F. Partensky (2005). "Diel variations in the photosynthetic parameters of Prochlorococcus strain PCC 9511: Combined effects of light and cell cycle." Limnology and Oceanography **50**(3): 850-863.

Cailliau, C., H. Claustre, F. Vidussi, D. Marie and D. Vaulot (1996). "Carbon biomass, and gross growth rates as estimated from ^{14}C pigment labelling, during photoacclimation in Prochlorococcus CCMP 1378." Marine Ecology Progress Series **145**(1/3): 209-221.

Campbell, L., H. A. Nolla and D. Vaulot (1994). "The importance of Prochlorococcus to community structure in
the central North Pacific Ocean." Limnology and Oceanography **39**(4): 954-961.
Cermak, N., J. W. Becker, S. M. Knudsen, S. W. Chisholm, S. R. Manalis and M. F. Polz (2016). "Direct single-
cell biomass estimates for marine bacteria via Archimedes' principle." The Isme Journal **11**: 825.
Christie-Oleza, J. A., D. J. Scanlan and J. Armengaud (2015). ""You produce while I clean up", a strategy
revealed by exoproteomics during Synechococcus–Roseobacter interactions." PROTEOMICS: n/a-n/a.
Christie-Oleza, J. A., D. Sousoni, M. Lloyd, J. Armengaud and D. J. Scanlan (2017). "Nutrient recycling
facilitates long-term stability of marine microbial phototroph–heterotroph interactions." Nature
Microbiology **2**: 17100.
Churchward, G., H. Bremer and R. Young (1982). "Macromolecular composition of bacteria." Journal of
Theoretical Biology **94**(3): 651-670.
Claustre, H., A. Bricaud, M. Babin, F. Bruyant, L. Guillou, F. Le Gall, D. Marie and F. Partensky (2002). "Diel
variations in Prochlorococcus optical properties." Limnology and Oceanography **47**(6): 1637-1647.
Claustre, H., S. B. Hooker, L. Van Heukelem, J. F. Berthon, R. Barlow, J. Ras, H. Sessions, C. Targa, C. S.
Thomas, D. van der Linde and J. C. Marty (2004). "An intercomparison of HPLC phytoplankton pigment
methods using in situ samples: application to remote sensing and database activities." Marine Chemistry
85(1-2): 41-61.
Coe, A., J. Ghizzoni, K. LeGault, S. Biller, S. E. Roggensack and S. W. Chisholm (2016). "Survival of
Prochlorococcus in extended darkness." Limnology and Oceanography **61**(4): 1375-1388.
Dickson, A. G., C. L. Sabine and J. R. e. Christian (2007). Guide to best practices for ocean CO2 measurement.
Sidney, British Columbia, North Pacific Marine Science Organization**: 191.
Dubinsky, Z. and I. Berman-Frank (2001). "Uncoupling primary production from population growth in
photosynthesizing organisms in aquatic ecosystems." Aquatic Sciences **63**(1): 4-17.
DuRand, M. D., R. J. Olson and S. W. Chisholm (2001). "Phytoplankton population dynamics at the Bermuda
Atlantic Time-series station in the Sargasso Sea." Deep Sea Research Part II: Topical Studies in Oceanography
48(8–9): 1983-2003.
Elser, J. J., K. Acharya, M. Kyle, J. Cotner, W. Makino, T. Markow, T. Watts, S. Hobbie, W. Fagan, J. Schade, J.
Hood and R. W. Sterner (2003). "Growth rate–stoichiometry couplings in diverse biota." Ecology Letters
6(10): 936-943.
Fang, X., Y. Liu, Y. Zhao, Y. Chen, R. Liu, Q.-L. Qin, G. Li, Y.-Z. Zhang, W. Chan, W. R. Hess and Q. Zeng (2019).
"Transcriptomic responses of the marine cyanobacterium Prochlorococcus to viral lysis products."
Environmental Microbiology **21**(6): 2015-2028.
Felcmanová, K., M. Lukeš, E. Kotabová, E. Lawrenz, K. H. Halsey and O. Prášil (2017). "Carbon use efficiencies
and allocation strategies in Prochlorococcus marinus strain PCC 9511 during nitrogen-limited growth."
Photosynthesis Research **134**(1): 71-82.
Finkel, Z. V., M. J. Follows, J. D. Liefer, C. M. Brown, I. Benner and A. J. Irwin (2016). "Phylogenetic Diversity in
the Macromolecular Composition of Microalgae." PLOS ONE **11**(5): e0155977.
Flombaum, P., J. L. Gallegos, R. A. Gordillo, J. Rincón, L. L. Zabala, N. Jiao, D. M. Karl, W. K. W. Li, M. W.
Lomas, D. Veneziano, C. S. Vera, J. A. Vrugt and A. C. Martiny (2013). "Present and future global distributions
of the marine Cyanobacteria Prochlorococcus and Synechococcus." Proceedings of
the National Academy of Sciences **110**(24): 9824-9829.
Fu, F.-X., M. E. Warner, Y. Zhang, Y. Feng and D. A. Hutchins (2007). "EFFECTS OF INCREASED TEMPERATURE
AND CO2 ON PHOTOSYNTHESIS, GROWTH, AND ELEMENTAL RATIOS IN MARINE SYNECHOCOCCUS AND
PROCHLOROCOCCUS (CYANOBACTERIA)1." Journal of Phycology **43**(3): 485-496.
Garcia, N. S., J. A. Bonachela and A. C. Martiny (2016). "Interactions between growth-dependent changes in
cell size, nutrient supply and cellular elemental stoichiometry of marine Synechococcus." The Isme Journal
10: 2715.
Geider, R. J. and J. La Roche (2002). "Redfield revisited: variability of C : N : P in marine microalgae and its
biochemical basis." European Journal of Phycology **37**(1): 1-17.

Gilbert, J. D. J. and W. F. Fagan (2011). "Contrasting mechanisms of proteomic nitrogen thrift in
Prochlorococcus." Molecular Ecology **20**(1): 92-104.
Goebel, N. L., C. A. Edwards, J. P. Zehr and M. J. Follows (2010). "An emergent community ecosystem model
applied to the California Current System." Journal of Marine Systems **83**(3): 221-241.
Grob, C., M. Ostrowski, R. J. Holland, M. Heldal, S. Norland, E. S. Erichsen, C. Blindauer, A. P. Martin, M. V.
Zubkov and D. J. Scanlan (2013). "Elemental composition of natural populations of key microbial groups in
Atlantic waters." Environmental Microbiology **15**(11): 3054-3064.
Grossowicz, M., D. Roth-Rosenberg, D. Aharonovich, J. Silverman, M. J. Follows and D. Sher (2017).
"Prochlorococcus in the lab and in silico: The importance of representing exudation." Limnology and
Oceanography **62**(2): 818-835.
Gruber, N. and C. A. Deutsch (2014). "Redfield's evolving legacy." Nature Geoscience **7**: 853.
Guyennon, A., M. Baklouti, F. Diaz, J. Palmieri, J. Beuvier, C. Lebaupin-Brossier, T. Arsouze, K. Béranger, J. C.
Dutay and T. Moutin (2015). "New insights into the organic carbon export in the Mediterranean Sea from 3-D
modeling." Biogeosciences **12**(23): 7025-7046.
Hansell, D. A., C. A. Carlson, D. J. Repeta and R. Schlitzer (2009). "Dissolved Organic Matter in the Ocean a
Controversy Stimulates New Insights." Oceanography **22**(4): 202-211.
Heldal, M., D. J. Scanlan, S. Norland, F. Thingstad and N. H. Mann (2003). "Elemental composition of single
cells of various strains of marine Prochlorococcus and Synechococcus using X-ray microanalysis." Limnology
and Oceanography **48**(5): 1732-1743.
Hu, Q., M. Sommerfeld, E. Jarvis, M. Ghirardi, M. Posewitz, M. Seibert and A. Darzins (2008). "Microalgal
triacylglycerols as feedstocks for biofuel production: perspectives and advances." The Plant Journal **54**(4):
621-639.
Hughes, C., D. J. Franklin and G. Malin (2011). "Iodomethane production by two important marine
cyanobacteria: Prochlorococcus marinus (CCMP 2389) and Synechococcus sp. (CCMP 2370)." Marine
Chemistry **125**(1–4): 19-25.
Iuculano, F., I. P. Mazuecos, I. Reche and S. Agustí (2017). "Prochlorococcus as a Possible Source for
Transparent Exopolymer Particles (TEP)." Frontiers in Microbiology **8**(709).
Johnson, Z. I., E. R. Zinser, A. Coe, N. P. McNulty, E. M. Woodward and S. W. Chisholm (2006). "Niche
partitioning among Prochlorococcus ecotypes along ocean-scale environmental gradients." Science
311(5768): 1737-1740.
Kettler, G. C., A. C. Martiny, K. Huang, J. Zucker, M. L. Coleman, S. Rodrigue, F. Chen, A. Lapidus, S. Ferriera, J.
Johnson, C. Steglich, G. M. Church, P. Richardson and S. W. Chisholm (2007). "Patterns and implications of
gene gain and loss in the evolution of Prochlorococcus." PLoS Genet **3**(12): e231.
Klotz, A., J. Georg, L. Bučinská, S. Watanabe, V. Reimann, W. Januszewski, R. Sobotka, D. Jendrossek,
Wolfgang R. Hess and K. Forchhammer (2016). "Awakening of a Dormant Cyanobacterium from Nitrogen
Chlorosis Reveals a Genetically Determined Program." Current Biology **26**(21): 2862-2872.
Komatsu, H., K. Wada, T. Kanjoh, H. Miyashita, M. Sato, M. Kawachi and M. Kobayashi (2016). "Unique
chlorophylls in picoplankton Prochlorococcus sp. "Physicochemical properties of divinyl chlorophylls, and the
discovery of monovinyl chlorophyll b as well as divinyl chlorophyll b in the species Prochlorococcus NIES-
2086"." Photosynth Res **130**(1-3): 445-467.
Li, S. H.-J., Z. Li, J. O. Park, C. G. King, J. D. Rabinowitz, N. S. Wingreen and Z. Gitai (2018). "Escherichia coli
translation strategies differ across carbon, nitrogen and phosphorus limitation conditions." Nature
microbiology **3**(8): 939-947.
Li, W. K. W., P. M. Dickie, B. D. Irwin and A. M. Wood (1992). "Biomass of bacteria, cyanobacteria,
prochlorophytes and photosynthetic eukaryotes in the Sargasso Sea." Deep Sea Research Part A.
Oceanographic Research Papers **39**(3): 501-519.
Liefer, J. D., A. Garg, M. H. Fyfe, A. J. Irwin, I. Benner, C. M. Brown, M. J. Follows, A. W. Omta and Z. V. Finkel
(2019). "The Macromolecular Basis of Phytoplankton C:N:P Under Nitrogen Starvation." Frontiers in
Microbiology **10**(763).

Llabrés, M., S. Agustí and G. J. Herndl (2011). "DIEL IN SITU PICOPHYTOPLANKTON CELL DEATH CYCLES
COUPLED WITH CELL DIVISION1." Journal of Phycology **47**(6): 1247-1257.
Lomas, M. W. and S. B. Moran (2012). "Evidence for aggregation and export of cyanobacteria and nano-
eukaryotes from the Sargasso Sea euphotic zone." Biogeosciences **7**(5): 7173-7206.
Lopez-Sandoval, D. C., T. Rodriguez-Ramos, P. Cermeno and E. Maranon (2013). "Exudation of organic carbon
by marine phytoplankton: dependence on taxon and cell size." Marine Ecology Progress Series **477**: 53-60.
Lourenço, S. O., E. Barbarino, U. M. L. Marquez and E. Aidar (1998). "DISTRIBUTION OF INTRACELLULAR
NITROGEN IN MARINE MICROALGAE: BASIS FOR THE CALCULATION OF SPECIFIC NITROGEN-TO-PROTEIN
CONVERSION FACTORS." Journal of Phycology **34**(5): 798-811.
Marmen, S., L. Blank, A. Al-Ashhab, A. Malik, L. Ganzert, M. Lalzar, H.-P. Grossart and D. Sher (2018). "The
role of land use types and water chemical properties in structuring the microbiome of a connected lakes
system." bioRxiv.
Martens-Habbena, W. and H. Sass (2006). "Sensitive determination of microbial growth by nucleic acid
staining in aqueous suspension." Applied and environmental microbiology **72**(1): 87-95.
Martiny, A. C., M. L. Coleman and S. W. Chisholm (2006). "Phosphate acquisition genes in Prochlorococcus
ecotypes: evidence for genome-wide adaptation." Proc Natl Acad Sci U S A **103**(33): 12552-12557.
Martiny, A. C., L. Ma, C. Mouginot, J. W. Chandler and E. R. Zinser (2016). "Interactions between Thermal
Acclimation, Growth Rate, and Phylogeny Influence Prochlorococcus Elemental Stoichiometry." PLOS ONE
11(12): e0168291.
Martiny, A. C., C. T. A. Pham, F. W. Primeau, J. A. Vrugt, J. K. Moore, S. A. Levin and M. W. Lomas (2013).
"Strong latitudinal patterns in the elemental ratios of marine plankton and organic matter." Nature
Geoscience **6**: 279.
Massana, R., A. E. Murray, C. M. Preston and E. F. DeLong (1997). "Vertical distribution and phylogenetic
characterization of marine planktonic Archaea in the Santa Barbara Channel." Applied and environmental
microbiology **63**(1): 50-56.
McDonagh, B., M. A. Domínguez-Martín, G. Gómez-Baena, A. López-Lozano, J. Diez, J. A. Bárcena and J. M.
García Fernández (2012). "Nitrogen starvation induces extensive changes in the redox proteome of
Prochlorococcus sp. strain SS120." Environmental Microbiology Reports **4**(2): 257-267.
Metropolis, N., A. W. Rosenbluth, M. N. Rosenbluth, A. H. Teller and E. Teller (1953). "Equation of State
Calculations by Fast Computing Machines." The Journal of Chemical Physics **21**(6): 1087-1092.
Moore, L. R. (1997). Physiological ecology of Prochlorococcus : a comparison of isolates from diverse
oceanographic regimes. PhD, Massachusetts Institute of Technology.
Moore, L. R. and S. W. Chisholm (1999). "Photophysiology of the marine cyanobacterium Prochlorococcus:
Ecotypic differences among cultured isolates." Limnology and Oceanography **44**(3): 628-638.
Moore, L. R., A. Coe, E. R. Zinser, M. A. Saito, M. B. Sullivan, D. Lindell, K. Frois-Moniz, J. Waterbury and S. W.
Chisholm (2007). "Culturing the marine cyanobacterium Prochlorococcus." Limnology and Oceanography-
Methods **5**: 353-362.
Moore, L. R., R. Goericke and S. W. Chisholm (1995). "Comparative physiology of Synechococcus and
Prochlorococcus: influence of light and temperature on growth, pigments, fluorescence and absorptive
properties." Marine Ecology Progress Series **116**(1/3): 259-275.
Moore, L. R., A. F. Post, G. Rocap and S. W. Chisholm (2002). "Utilization of different nitrogen sources by the
marine cyanobacteria Prochlorococcus and Synechococcus." Limnology and Oceanography **47**(4): 989-996.
Neidhardt, F. C., J. L. Ingraham and M. Schaechter (1990). Physiology of the Bacterial Cell: a Molecular
Approach, Sinauer Associates, Sunderland, Mass.
Omta, A. W., J. Bruggeman, B. Kooijman and H. Dijkstra (2009). "The organic carbon pump in the Atlantic."
Journal of Sea Research **62**(2): 179-187.
Omta, A. W., D. Talmy, D. Sher, Z. V. Finkel, A. J. Irwin and M. J. Follows (2017). "Extracting phytoplankton
physiological traits from batch and chemostat culture data." Limnology and Oceanography: Methods **15**(5):
453-466.

Partensky, F. and L. Garczarek (2010). "Prochlorococcus: Advantages and Limits of Minimalism." Annual Review of Marine Science **2**: 305-331.

Read, R. W., P. M. Berube, S. J. Biller, I. Neveux, A. Cubillos-Ruiz, S. W. Chisholm and J. J. Grzymski (2017). "Nitrogen cost minimization is promoted by structural changes in the transcriptome of N-deprived Prochlorococcus cells." The ISME journal **11**(10): 2267-2278.

Redfield, A. C. (1934). On the Proportions of Organic Derivatives in Sea Water and Their Relation to the Composition of Plankton. James Johnstone Memorial Volume, University Press of Liverpool**:** 176-192.

Ribalet, F., J. Swalwell, S. Clayton, V. Jiménez, S. Sudek, Y. Lin, Z. I. Johnson, A. Z. Worden and E. V. Armbrust (2015). "Light-driven synchrony of Prochlorococcus growth and mortality in the subtropical Pacific gyre." Proceedings of the National Academy of Sciences **112**(26): 8008-8012.

Richardson, T. L. and G. A. Jackson (2007). "Small phytoplankton and carbon export from the surface ocean." Science **315**(5813): 838-840.

Roshan, S. and T. DeVries (2017). "Efficient dissolved organic carbon production and export in the oligotrophic ocean." Nature Communications **8**(1): 2036.

Roth-Rosenberg, D., D. Aharonovich, T. Luzzatto-Knaan, A. Vogts, L. Zoccarato, F. Eigemann, N. Nago, H. P. Grossart, M. Voss and D. Sher (2019). "Prochlorococcus rely on microbial interactions rather than on chlorotic resting stages to survive long-term stress." BioRXiv **MS ID#: BIORXIV/2019/657627**.

Saito, M. A., M. R. McIlvin, D. M. Moran, T. J. Goepfert, G. R. DiTullio, A. F. Post and C. H. Lamborg (2014). "Multiple nutrient stresses at intersecting Pacific Ocean biomes detected by protein biomarkers." Science **345**(6201): 1173-1177.

Sauer, J., U. Schreiber, R. Schmid, U. Völker and K. Forchhammer (2001). "Nitrogen Starvation-Induced Chlorosis in Synechococcus PCC 7942. Low-Level Photosynthesis As a Mechanism of Long-Term Survival." Plant Physiology **126**(1): 233-243.

Sharma, A. K., J. W. Becker, E. A. Ottesen, J. A. Bryant, S. Duhamel, D. M. Karl, O. X. Cordero, D. J. Repeta and E. F. DeLong (2014). "Distinct dissolved organic matter sources induce rapid transcriptional responses in coexisting populations of Prochlorococcus, Pelagibacter and the OM60 clade." Environmental Microbiology **16**(9): 2815-2830.

Shaw, S. L., S. W. Chisholm and R. G. Prinn (2003). "Isoprene production by Prochlorococcus, a marine cyanobacterium, and other phytoplankton." Marine Chemistry **80**(4): 227-245.

Steglich, C., M. Behrenfeld, M. Koblizek, H. Claustre, S. Penno, O. Prasil, F. Partensky and W. R. Hess (2001). "Nitrogen deprivation strongly affects Photosystem II but not phycoerythrin level in the divinyl-chlorophyll b-containing cyanobacterium Prochlorococcus marinus." Biochimica et Biophysica Acta (BBA) - Bioenergetics **1503**(3): 341-349.

Steglich, C., C. W. Mullineaux, K. Teuchner, W. R. Hess and H. Lokstein (2003). "Photophysical properties of Prochlorococcus marinus SS120 divinyl chlorophylls and phycoerythrin in vitro and in vivo." FEBS Letters **553**(1–2): 79-84.

Szul, M. J., S. P. Dearth, S. R. Campagna and E. R. Zinser (2019). "Carbon Fate and Flux in Prochlorococcus under Nitrogen Limitation." mSystems **4**(1): e00254-00218.

Talmy, D., A. C. Martiny, C. Hill, A. E. Hickman and M. J. Follows (2016). "Microzooplankton regulation of surface ocean POC:PON ratios." Global Biogeochemical Cycles **30**(2): 311-332.

Thompson, A. W., K. Huang, M. A. Saito and S. W. Chisholm (2011). "Transcriptome response of high- and low-light-adapted Prochlorococcus strains to changing iron availability." Isme J **5**(10): 1580-1594.

Thornton, D. C. O. (2014). "Dissolved organic matter (DOM) release by phytoplankton in the contemporary and future ocean." European Journal of Phycology **49**(1): 20-46.

Ting, C. S., G. Rocap, J. King and S. W. Chisholm (2002). "Cyanobacterial photosynthesis in the oceans: the origins and significance of divergent light-harvesting strategies." Trends in Microbiology **10**(3): 134-142.

Tolonen, A. C., J. Aach, D. Lindell, Z. I. Johnson, T. Rector, R. Steen, G. M. Church and S. W. Chisholm (2006). "Global gene expression of Prochlorococcus ecotypes in response to changes in nitrogen availability." Mol Syst Biol **2**: 53.

Van Mooy, B. A., G. Rocap, H. F. Fredricks, C. T. Evans and A. H. Devol (2006). "Sulfolipids dramatically
decrease phosphorus demand by picocyanobacteria in oligotrophic marine environments." <u>Proc Natl Acad Sci</u>
<u>U S A</u> **103**(23): 8607-8612.
Vargas, M. A., H. Rodríguez, J. Moreno, H. Olivares, J. A. D. Campo, J. Rivas and M. G. Guerrero (1998).
"BIOCHEMICAL COMPOSITION AND FATTY ACID CONTENT OF FILAMENTOUS NITROGEN-FIXING
CYANOBACTERIA." <u>Journal of Phycology</u> **34**(5): 812-817.
Veldhuis, M. and G. W. Kraay (1990). "Vertical distribution and pigment composition of a picoplanktonic
prochlorophyte in the subtropical North Atlantic:a combined study of HPLC-analysis of pigments and flow
cytometry." <u>Marine Ecology Progress Series</u> **68**: 121-127.
Zavřel, T., M. Faizi, C. Loureiro, G. Poschmann, K. Stühler, M. Sinetova, A. Zorina, R. Steuer and J. Červený
(2019). "Quantitative insights into the cyanobacterial cell economy." <u>eLife</u> **8**: e42508.
Zhao, Z., M. Gonsior, J. Luek, S. Timko, H. Ianiri, N. Hertkorn, P. Schmitt-Kopplin, X. Fang, Q. Zeng, N. Jiao and
F. Chen (2017). "Picocyanobacteria and deep-ocean fluorescent dissolved organic matter share similar optical
properties." <u>Nature communications</u> **8**: 15284-15284.
Zhu, Y., J. E. Graham, M. Ludwig, W. Xiong, R. M. Alvey, G. Shen and D. A. Bryant (2010). "Roles of
xanthophyll carotenoids in protection against photoinhibition and oxidative stress in the cyanobacterium
Synechococcus sp. strain PCC 7002." <u>Archives of Biochemistry and Biophysics</u> **504**(1): 86-99.
Zimmerman, A. E., S. D. Allison and A. C. Martiny (2014). "Phylogenetic constraints on elemental
stoichiometry and resource allocation in heterotrophic marine bacteria." <u>Environmental Microbiology</u> **16**(5):
1398-1410.
Zinser, E. R., D. Lindell, Z. I. Johnson, M. E. Futschik, C. Steglich, M. L. Coleman, M. A. Wright, T. Rector, R.
Steen, N. McNulty, L. R. Thompson and S. W. Chisholm (2009). "Choreography of the transcriptome,
photophysiology, and cell cycle of a minimal photoautotroph, prochlorococcus." <u>PLoS ONE</u> **4**(4): e5135.

803