Title: Dietary fat and fiber impacts intestinal microbiome resilience to antibiotics and *Clostridoides difficile* infection in mice

Authors: Keith Z. Hazleton[1,2], Casey G. Martin[3], Kathleen L. Arnolds[3], Nichole M. Nusbacher[4], Nancy Moreno-Huizar[4], Michael Armstrong[5], Nichole Reisdorph[5], Catherine A. Lozupone[4]*

Affiliations:

[1] Department of Pediatrics, Section of Gastroenterology, Hepatology and Nutrition. University of Colorado, Denver Anschutz Medical Campus, Aurora, CO USA 80045.

[2] Digestive Health Institute, Children's Hospital Colorado, Aurora, CO USA 80045.

[3] Department of Immunology and Microbiology, University of Colorado, Denver Anschutz Medical Campus, Aurora, CO USA 80045.

[4] Department of Medicine, Division of Biomedical Informatics and Personalized Medicine, University of Colorado, Denver Anschutz Medical Campus, Aurora, CO USA 80045.

[5] Skaggs School of Pharmacy and Pharmaceutical Sciences, University of Colorado Anschutz Medical Campus, Aurora, CO USA 80045.

*To whom correspondence should be addressed: Catherine.lozupone@cuanschutz.edu

One Sentence Summary: High dietary fat promoted mortality in a mouse model of antibiotic-induced *C. difficile* infection and low dietary fiber caused higher microbiome disturbance upon broad-spectrum antibiotic exposure, suggesting that diets low in fat and high in fiber may protect against *C. difficile* pathogenesis.

Abstract: *Clostridoides difficile* infection (CDI) is a leading cause of hospital-acquired diarrhea and there has been a steady increase in the number of new infections, emphasizing the importance of novel prevention strategies. Use of broad-spectrum antibiotics and disruption of the intestinal microbiome is one of the most important risk factors of CDI. We used a murine model of antibiotic-induced CDI to investigate the relative contributions of high dietary fat and low dietary fiber on disease pathogenesis. We found that high fat, but not low fiber resulted in increased mortality from CDI (HR 4.95) and increased levels of *C. difficile* toxin production compared to a regular low-fat/high-fiber mouse diet even though we did not observe a significant change in *C. difficile* carriage. The high-fat diet also increased levels of primary bile acids known to be germination factors for *C. difficile* spores. Mice fed low-fat/low-fiber diets did not show increased CDI pathogenesis, but did have a larger antibiotic-induced gut microbiome disturbance compared to mice fed a high-fiber diet, characterized by a greater decrease in alpha diversity. This microbiome disturbance was associated with a loss of secondary bile acids and short chain fatty acids, which are both microbial metabolic products previously shown to protect against CDI. These data suggest that a low-fiber diet contributes to antibiotic-induced dysbiosis, while a high-fat diet promotes CDI pathogenesis. These findings indicate that dietary interventions that increase fiber and decrease fat may be an effective prevention strategy for individuals at high risk of CDI.

Introduction

Clostridoides difficile infection (CDI) is an important cause of morbidity and mortality, with 500,000 cases every year causing 30,000 deaths per year in the US alone (1). Alarmingly, there has been a steady increase in the number of new infections in spite of prevention efforts in hospitals that have focused largely on increased sanitation and antibiotic stewardship (2). These prevention approaches treat CDI as a traditional communicable infection. However, recent data suggests that CDI is often caused by the activation of strains that were already carried in the gut prior to hospital admission and prior to the onset of symptoms rather than by acquisition of a hospital-resident strain (3). Thus, determining strategies to reduce C. difficile carriage and attenuate CDI severity is critical.

A complex gut microbiome has been shown to be protective against CDI (4). Broad spectrum antibiotic usage, such as clindamycin, beta-lactams, and fluoroquinolones (5, 6) increase risk of CDI as do other illnesses associated with reduced gut microbiome diversity, such as Inflammatory Bowel Diseases (7). Individuals with recurrent CDI (rCDI) typically have microbiomes with greatly reduced complexity and altered compositions (8-12). The gut microbiome provides protection in part through conversion of conjugated primary bile acids, which are excreted by the liver into the intestine where they play a central role in fat digestion, into secondary bile acids. Taurine-conjugated primary bile acids can promote the germination and growth of C. difficile spores while secondary bile acids cause germination but arrest the growth of spores (13). Accordingly, prior studies have shown that secondary bile acid producers such as Clostridium scindens can protect against CDI in mice (14). Another class of bacterial metabolites that inhibits C. difficile are short chain fatty acids (SCFA), which are microbial

products of fiber fermentation. SCFAs can directly inhibit *C. difficile* growth *in vitro* and are

decreased in individuals with rCDI *(15, 16)*

Recent studies conducted in mice have suggested that diet modulation has the potential to

be an effective prevention strategy for CDI. Mice treated with broad-spectrum antibiotics and fed

diets devoid of microbial accessible carbohydrates (MACs; e.g. soluble fibers such as inulin)

showed increased persistence of *C. difficile* colonization consistent with a protective role of

SCFAs; inulin supplementation of diets low in MACS lead to rapid clearance of *C. difficile* in

this model *(15)*. While dietary fiber content can result in microbial metabolites that are protective

against CDI, other diet-influenced metabolites seem to promote the pathogenesis of CDI. For

instance, consistent with *C. difficile* being auxotrophic for the amino acid proline, experiments

conducted in gnotobiotic mice have supported that mice were protected from *C. difficile*

colonization with low-proline diets *(4)*.

Given that conjugated primary bile acids play a central role in fat digestion and are a

germination factor for *C. difficile* spores, we became interested in investigating a role for dietary

fat in CDI pathogenesis. In particular, since prior studies have indicated that the fraction of

taurine-conjugated colonic bile acids increase with diets high in saturated fat *(17)*, we

hypothesized that a high-fat diet could alter the intestinal bile acid composition and drive

pathology in a mouse model of CDI. We further hypothesized that the combination of high fat

and low fiber in the diet, as is common in individuals in developed countries, would together

promote CDI in mice. We found that high dietary fat content in an antibiotic-induced *C. difficile*

model induced high mortality from CDI. A low-fiber diet did not cause mortality but did

decrease the resilience of the gut microbiome to antibiotic-induced disturbance, a known risk

factor for CDI. Taken together, our results suggest that dietary interventions to increase fiber

and, perhaps more importantly, to decrease fat represent effective prevention strategies for

individuals at high risk of CDI.

Results

High dietary fat but not low dietary fiber causes increased mortality and toxin production in

murine CDI

To understand the relative effects of dietary fat and fiber on CDI, we used an established

murine model of antibiotic-induced CDI *(18)*. Specifically, conventional 6-week-old old female

C57-bl6 mice were fed 1 of 3 diets: 1) conventional mouse chow that is low-fat/high-fiber, 2) a

purified "Western" diet (WD) that had ~2x the content of fat with increased ratio of saturated-to-

unsaturated fat compared to chow and only insoluble cellulose as a source of fiber, and 3) the

same purified diet as the WD, but with a lower fat content, similar to chow (low-fat/low-fiber)

(Table 1). One week after diet switch, mice were treated with a cocktail of antibiotics in their

drinking water for 5 days (kanamycin, gentamicin, colistin, metronidazole and vancomycin)

followed by an injection of clindamycin and gavage with *C. difficile* VPI 10463 (Fig. 1A). We

conducted 2 sets of experiments. In the first set, the experiments were carried out for up to 21

days past *C. difficile* gavage, allowing us to assay effects on mortality and relate these to fecal

microbiome composition. In the second set of experiments, mice were sacrificed at day 3 past *C.*

difficile gavage (just prior to the observed onset of mortality in the first experiments), so that bile

and SCFA levels could be assessed from the cecum (Fig. 1A).

WD-fed mice showed a marked increase in mortality as compared to chow-fed mice (HR

4.95 CI 1.79-13.72) upon CDI exposure, and significantly higher levels of *C. difficile* toxin B in

the cecum after 3 days of infection (Fig. 1B, C). The low-fat/low-fiber diet-fed mice showed

survival and *C. difficile* toxin B levels comparable to the chow-fed mice (Fig. 1B, C). Using

qPCR to compare the relative abundance *C. difficile* to overall bacterial load in fecal samples

collected 2 days after *C. difficile* challenge, we did not observe any significant differences in *C.*

difficile carriage based on diet (Fig. 1D).

Because our WD and low-fat/low-fiber diet differed in sucrose content, we also tested a

fourth diet that was low in fat and fiber, but with sucrose equivalent to the WD (Table S1).

Sucrose does not appear to play a role in the increased mortality observed in the WD, as 100%

survival was observed in mice fed this fourth diet (n = 10, one cage with 5 mice in two separate

experiments) and those who received the low-fat/low-fiber/low-sucrose diet.

Table 1

	Low-fat/High fiber Chow diet	*High-fat/Low-fiber Western diet (WD)*	*Low-fat/Low-fiber diet*
Fat (% kcal)	16	34.5	17.2
(% SFA)	(N/A)	(36.2)	(19.5)
(% MUFA)		(41.3)	(41.7)
(% PUFA)		(22.5)	(38.8)
Carbohydrates (% kcal)	60	50	63.9
(Sucrose)	(0)	(23.4)	(10.6)
Protein (%kcal)	24	15.5	18.8
Fiber (g/kg)	137	50 (cellulose)	50 (cellulose)

Figure 1: Experimental design of murine model of antibiotic-induced CDI, survival curves and

C. difficile toxin production and colonization of mice on varying diets. **A)** *C. difficile* challenge

experimental design. The figure legend at the left panel indicates the samples sizes for the 2

cohorts. Grey and orange boxes indicate the timepoints at which samples were collected for the

respective cohorts **B)** Survival curves on the 3 diets. Both chow and low-fat/low-fiber (LF/LF)

diets show increased survival as compared to WD (high-fat/low-fiber). Statistical significance as

assessed by log-rank comparison is indicated. **C)** Toxin B levels measured in cecal contents of

mice by ELISA (tgcBiomics) at day 3 post infection. Statistical significance was assessed with

the Wilcoxon rank-sum test. **(D)** qPCR of *C. difficile* normalized to total 16S from fecal pellets

in fecal samples collected 2 days post *C. difficile* challenge, with significance determined by

Wilcoxon rank-sum test.

Fiber increases cecal levels of secondary bile acids and butyrate; fat increases taurine-conjugated primary bile acids

To understand the effects of diet modulation on microbial metabolites previously described to modulate *C. difficile* sporulation, growth and activity, we assessed levels of bile acids and SCFAs in aspirated cecal contents that were collected in a separate cohort of 51 mice (chow= 20, WD = 21, low-fat/low-fiber = 10) sacrificed at 3 days post *C. difficile* infection, which is just before mortality typically occurred with the WD diet (Fig. 1A). A total of 17 bile acids including conjugated and unconjugated primary bile acids found in both humans and mice and secondary bile acids, were quantified using HPLC/QTOF using the method described in *(19)* with modifications (see methods). We found that mice fed a WD showed significantly increased levels of taurine-conjugated primary bile acids compared to chow with a trend towards increased level compared to the low-fat/low-fiber diet (Fig 2A, Fig. S1C). The primary unconjugated bile acids (e.g. beta-muricholic acid, cholic acid) did not show a consistent pattern of change based on diet (Fig. S1A) and glycine conjugated bile acids were minimally detectable, which is consistent with mouse metabolism of bile acids strongly favoring taurine conjugation. Finally, the chow-fed mice demonstrated high levels of bacterially-produced secondary bile acids as compared to both WD and low-fat/low-fiber diets (Fig. 2A).

The SCFAs butyrate, acetate and propionate were also quantified in cecal contents 3 days after *C. difficile* infection using gas-chromatography/mass spectrometry (GC/MS). Consistent with SCFAs being microbial fermentation products of dietary fiber, cecal levels of butyrate were reduced in the WD and low-fat/low-fiber diets compared to chow (Fig. 2B). The SCFAs acetate and propionate showed a reduction in the low-fat/low-fiber but not the WD compared to chow Fig. 2B). Interestingly, there was a strong positive correlation between levels of the secondary

bile acid deoxycholate (DCA) and butyrate in both the chow and WD-fed mice, with a markedly

increased effect size (slope) in the chow-fed mice (Fig. 2C). For chow-fed mice DCA levels

increase by 94.4 pmol/mg wet weight for every 1 mmol/kg increase in butyrate ($p = 5.4 \times 10^{-9}$).

Diet alone was not associated with DCA levels ($p = 0.500$), but there was a significant

interaction between diet and butyrate; for WD-fed mice DCA levels increase by 18.92 pmol/mg

wet weight for every 1 mmol/kg increase in butyrate ($p = 5.1 \times 10^{-5}$). All low-fat/low-fiber

samples were omitted from the regression as none of the sample had detectable DCA.

Additionally, a single sample from the chow cohort (butyrate = 11.31 mmol/kg, DCA = 342

pmol/mg wet weight) was omitted due to its outsized influence on the regression fit (DFFITs = -

12.7).

To explore a direct relationship between microbial metabolites and *C. difficile*

pathogenicity, we determined whether any bile acids or SCFAs correlated with *C. difficile* toxin

production using linear regression. The model [Toxin B] ~ diet had an adjusted R-squared of

0.1748 and p-value of 0.0131. The only metabolite that was associated with toxin B

concentration in univariate analysis was taurocholic acid-3-sulfate (TCA-3-SO4) (Table S2).

However, from multivariate analysis it would appear that this association is driven by

collinearity of TCA-3-SO4 and Toxin B concentrations with respect to diet (Fig. S2). Addition of

other individual metabolites to the regression model with diet (e.g. Toxin B ~ metabolite + diet

or Toxin B ~ metabolite + diet + diet*metabolite) did not improve the regression model,

suggesting that dietary control of toxin production is independent of the metabolites we

interrogated.

Figure 2: Metabolite changes in cecal contents of infected mice 3 days post *C. difficile* infection.

(A) Cecal levels of taurine-conjugated primary bile acids (left column) and bacterially-produced

secondary bile acids (second column) across diets. TCA= Taurocholic acid; T-b-MCA=Tauro-b-

muricholic acid; DC=Deoxycholic acid; LCA=Lithocholic acid. **(B)** Cecal short chain fatty acid

levels across diets. P-values were determined using the Wilcoxon rank-sum statistical test. **(C)**

Multiple linear regression of DCA levels as a function of butyrate and diet. Model = DCA ~

butyrate + diet + butyrate*diet. R-squared 0.855 p <0.0001. Low-fat/low-fiber samples were

excluded from analysis since there was no detectable DCA.

Dietary fiber increases homogeneity of response, resilience and alpha-diversity of the gut

microbiome after challenge with antibiotics and CDI

We next sought to understand how the composition of the fecal microbiome was affected by diet during the course of antibiotic treatment and infection with *C. difficile* (Fig. 1A). Fecal pellets were collected upon arrival prior to diet change (Day -7), just prior to the start of oral antibiotic delivery (Day 0), after 5 days of oral antibiotics (Day 5), and daily through Day 10, which captured before and after the clindamycin injection given on day 7 and *C. difficile* gavage on Day 8 (Fig. 1A). Collected samples were subjected to 16S ribosomal RNA (rRNA) amplicon sequencing targeting the V4 region of rRNA on the MiSeq platform.

Principle coordinate analysis (PCoA) plots of a weighted UniFrac distance matrix suggested that mice fed either low-fiber diet had decreased resilience and a less homogeneous response to antibiotic challenge and CDI as compared to chow-fed mice. Mice fed either low-fiber diet showed greater divergence across PC1 upon antibiotic exposure than chow-fed mice, higher spread across mice in the same diet group, and less recovery towards their baseline after antibiotics (Fig. 3A). We quantified resilience by comparing the pairwise weighted UniFrac distances of mice across the experiment to baseline microbiota of their respective diet cohort at Day 0 (7 days post-diet change and pre-oral antibiotics; Fig. 3B). Chow-fed mice had significantly smaller weighted UniFrac distances from their baselines than the other groups at Day 5 (post 5 days antibiotic challenge) that persisted through Day 10 despite some convergence after clindamycin injection (Day 8) (Fig. 3B). By Day 9, chow-fed mice again displayed higher microbiome resilience than both low-fiber groups. We also assessed the homogeneity of response to a disturbance among mice in the same diet group. As an example, low homogeneity would occur if the mice within a diet group showed high variability in the degree to which their

gut microbiome changed upon antibiotic exposure. We quantified this as the median pairwise

weighted UniFrac distance for comparisons within samples collected at the same time point from

mice fed the same diet (Fig. 3C). Both low-fiber diets showed much lower homogeneity of gut

microbiome compositional response to antibiotic challenge, particularly to the 5 day treatment

with oral antibiotics (Day 5), compared to chow-fed mice (Fig. 3C).

Similar patterns were seen when evaluating changes in alpha-diversity across the

experiment between each diet cohort. Figure 4 shows changes in phylogenetic entropy, which is

a measure of alpha diversity that considers species richness, evenness, and distinctness *(20)*. The

phylogenetic entropy of the WD-fed mice was lower than chow-fed mice after diet change and

this difference became more pronounced upon oral antibiotics and remained so through the rest

of the experimental timeline (Fig. 4). Interestingly, the phylogenetic entropy of the low-fat/low-

fiber diet-fed mice remained equivalent to the chow-fed cohort with diet change but decreased to

the same level as the WD with antibiotic treatment (Fig. 4).

Figure 3: Beta diversity plots of fecal microbiome by diet during antibiotic treatment and

infection with *C. difficile*. Vertical red lines in panels B and C designate the day of *C. difficile*

infection. (A) Weighted UniFrac PCoA plots of all samples with each diet highlighted in

separate panels. (B) Resilience of microbiome composition assessed by within-mouse pairwise

weighted UniFrac distances between Day 0 (7 days post diet switch and prior to oral antibiotics)

and later time points and (C) Longitudinal plot of microbiome turnover homogeneity as plotted

by intra-time point pairwise Weighted UniFrac distances within diet groups. Trend lines were fit

using local polynomial regression.

Figure 4: Alpha-diversity (phylogenetic entropy) of the fecal microbiome during murine CDI

model. Data for each individual mouse is plotted as well as the fitted local polynomial regression

for each diet group. Significant differences between groups are noted as calculated with the

Wilcoxon rank-sum test. ***: p<0.001. **: p<0.01, *: p<0.05 ns= non significant.

Dietary fiber influences colonization patterns of facultative anaerobes and of secondary bile acid

and SCFA-producing bacteria

Low-diversity dysbiosis is a state of disturbance that is often characterized not only by low alpha-diversity, but also by an increased ratio of facultative to strict anaerobes *(21)*. Low-diversity dysbiosis is associated with a number of diseases including rCDI *(21)*. We sought to investigate whether high dietary fat and low dietary fiber influenced if the microbiome developed a compositional state characterized by high levels of facultative anaerobe colonization and lower levels of strict anaerobes. Since Lactobacilliales and Enterobacteriales contain many important intestinal facultative anaerobes and most members of Clostridiales are strict anaerobes and include key butyrate and secondary bile acid producers, we plotted the relative abundances of these orders over the course of the experiment (Fig. 5A). All mice had decreases in the relative abundance of Clostridiales in their fecal microbiome with oral antibiotics; however, mice fed a chow diet were able to maintain a Clostridiales population while both low-fiber diets saw near-complete elimination of these taxa (chow-WD $p<0.01$ for days 0 through 9 and $p<0.05$ on day 10, Fig. S3). Conversely, mice fed either low-fiber diet had a large bloom of Lactobacilliales after oral antibiotic treatment that was not observed in the chow-fed mice (chow-WD $p<0.001$ and chow-low-fat/low-fiber $p<0.05$). Lastly, all 3 diet groups had a large increase in Enterobacteriales in their fecal microbiome following antibiotics; however, the low-fat/low-fiber and chow groups showed earlier decrease than WD mice (chow-WD $p<0.01$ and $p<0.05$ at days 9 and 10 respectively, Fig. S3). Comparisons of the low-fat/low-fiber diet were limited due to smaller sample size ($n = 5$ vs. $n = 13$ for chow and WD).

We also used PICRUSt *(22)* to predict metagenomes using our 16S rRNA data 1) to investigate trends in the prevalence of key genes in secondary bile and butyrate production over

the course of our experimental timeline and 2) to predict which bacterial taxa were contributing

these genes. Because BaiA, BaiB, and BaiCD are not available in PICRUSt2's set of predicted

genes, we only used the genes for BaiH (KEGG ID: K15873) and BaiI (KEGG ID; K15874),

which are both genes in the Bai operon *(23)*, to assess genomic potential for secondary bile acid

metabolism. Acetoacetate co-A transferase (*but*; K01034) and Butyrate Kinase (*buk*; KEGG ID:

K00929), which are the main pathways for fermentative production of butyrate in the gut

microbiome *(24)*, were used to assess butyrate production potential. Plotting these

genes/pathways over time reveals a significant effect of fiber on their abundance and response to

antibiotics (Fig. 5C). Although all diet groups showed a marked decrease in bile acid genes with

oral antibiotics, only the chow-fed mice displayed a recovery of bile acids genes, though the

source of these genes switched from Lachnospiraceae UCG-006 to Blautia. This result is

consistent with our observation of higher cecal levels of secondary bile acids in chow-fed mice

compared to mice fed either low-fiber diets at 3 days post *C. difficile* gavage (Fig. 2A).

Fiber also influenced the butyrate coding capacity observed between diet groups. Chow-

fed mice showed minimal change in the abundance of both the *but* and *buk* genes for

fermentative butyrate production during the time course while the WD mice had a decrease of 5

orders of magnitude (Fig 5D). The low-fat/low-fiber mice showed an intermediate phenotype

with the resilience of the butyrate pathway being mostly attributed to a butyrate kinase dependent

pathway. The results for *but* and not *buk* however are consistent with our measurements of cecal

butyrate levels in these mice 3 days post *C. difficile* gavage (Fig. 2B). This is consistent with *but*

being regarded to be a more important source of butyrate in the intestine *(25)*.

Since we had observed a strong positive correlation between cecal levels of butyrate and

the secondary bile acid DCA in our mass spectrometry data (Fig. 2C), we also determined

whether there was a relationship between butyrate and secondary bile acid coding capacity. We

found a highly significant association (p = 3.6×10^{-5}), with secondary bile acid producing genes

only predicted to be present in samples that also had high predicted levels of butyrate producing

genes (Fig. 5B).

Figure 5: Changes in key taxa, and secondary bile acid and butyrate coding capacity during the

CDI protocol. The vertical red line in (A), (C) and (D) represent day of infection with *C. difficile*.

All trend lines were fit using local polynomial regression. **(A)** Relative abundance of key

bacterial orders during antibiotic treatment and infection. Clostridiales are strict anaerobes while

Enterobacteriales and Lactobacilliales are facultative anaerobes. **(B)** Violin plots of abundance of

butyrate genes from PICRUSt analysis binned by presence of secondary bile acid producing

genes (Wilcoxon p<0.001). **(C)** Time course of coding capacity of secondary bile acid genes.

The top row shows the total capacity of each sample (BaiH and BaiI) while the bottom two rows

show specific taxa contributions of key genes in the Bai operon. **(D)** Time course of coding

capacity of butyrate producing genes by diet. The top row shows the total capacity as measured

304   by *but* and *buk* genes while the bottom two rows show specific taxa contributions of *but* and *buk*

305   specifically. Taxa with mean relative abundance $< 0.01\%$ were filtered from the analysis.

Discussion

     Clostridioides difficile infection is a grave and growing health threat. Current strategies to limit its spread have focused on sanitation and antibiotic stewardship, however incidence has continued to rise *in spite* of these efforts, highlighting the need for new treatment and prevention strategies *(2)*. Because of the ubiquity of *C. difficile* spores in the environment, focusing on ways to increase the resilience of the host to colonization is one important prevention strategy. However, modulation of host factors that influence virulence of *C. difficile* already present in the gut is also a key aspect of prevention, as CDI is often caused by *C. difficile* that is already residing in the gut before the onset of symptomatic CDI rather than acquisition of a new infection *(3)*.

     A growing body of evidence points to dietary intervention as a promising new approach to prevent CDI colonization *(4, 15)*. One recent study showed that a diet poor in proline (an essential amino acid for *C. difficile* growth) prevented *C. difficile* carriage *(4)*. Another study demonstrated that mice fed a diet deficient in MACs (e.g. soluble fiber, resistant starches) had persistent *C. difficile* shedding and that there was a resolution of colonization with the reintroduction of inulin or other MACs *(15)*. Although these papers both showed a strong effect of dietary factors on *C. difficile* colonization, neither study observed the high levels of mortality that we observed here with a high-fat/low-fiber WD. This is consistent with our findings that a low-fat/low-fiber diet did not increase mortality in our antibiotic induced CDI murine model, and suggests a particularly high influence of dietary fat on CDI disease severity. This is further supported by the fact that *C. difficile* abundance 2 days post *C. difficile* challenge, as quantified by strain-specific qPCR, was not significantly influenced by diet, suggesting that in an acute infection model, *C. difficile* blooms regardless of diet, and that increased disease severity is due

to dietary fat activating toxin production and not from increased *C. difficile* carriage.

Furthermore, although new data has suggested that novel speciation of *C. difficile* may be

selecting for strains that show increased sporulation and host colonization capacity with sugar

(glucose or fructose) *(26)*, this work, conducted with a hyper-virulent *C. difficile* strain (VPI

10463), did not show differences in mortality from CDI in low-fat/low-fiber diets with different

amounts of sucrose.

Evidence to suggest that a high-fat/low-fiber western diet could have a more profound

effect on CDI was first presented over 20 years ago. In experiments designed to study the

atherogenic properties of a Western diet in Syrian hamsters, significant mortality from CDI was

observed in hamsters fed a high-fat/low-fiber pro-atherogenic diet and not a typical high-

fiber/low-fat hamster diet *(27, 28)*. Our work confirms the same effect of a WD in an antibiotic-

induced murine CDI model.

Differences in host bile acid production and microbial bile acid metabolism is one

potential mechanism of high-fat diet induced modulation of CDI severity. *In vitro* experiments

have shown that the primary bile acid taurocholate is a potent *C. difficile* germination and growth

factor *(13)*. We found that mice fed a high-fat/low-fiber WD had higher cecal levels of primary

taurine-conjugated bile acids compared to the two low-fat diets tested. This is consistent with a

prior study that found that IL10-deficient mice fed a diet high in saturated fat, had an increased

proportion of taurine-conjugated bile acids compared to standard chow and a diet high in poly-

unsaturated fats *(17)*. However, in our current study multivariate regression did not correlate any

species of bile acid with toxin B concentration. This suggests that dietary fat may directly

modulate *C. difficile* Toxin B production or act through a non-bile acid dependent pathway to

promote CDI in our model.

We also sought to explore a role for secondary bile acids in phenotypes observed in diet-

induced CDI. *In vitro* experiments have shown that the bacterially produced secondary bile acids

deoxycholate and lithocholate caused germination followed by growth arrest of *C. difficile (13)*.

In line with these effects, reduced prevalence of the secondary bile acid producer, *Clostridium*

scindens in the fecal microbiome has been associated with high incidence of CDI in both humans

and in experimental mouse models, and gavaging mice with *C. scindens* protected against CDI

and restored intestinal secondary bile acid levels *(14)*. We observed that chow-fed mice had high

levels of secondary bile acids compared to both WD and low-fat/low-fiber diets. Functional

interrogation of the microbiome using PICRUSt suggests that this might be due to a lack of

recovery of secondary bile acid producing bacteria following antibiotic disturbance in both low-

fiber diet contexts. However, mice fed the low-fat/low-fiber diet did not demonstrate the

increased mortality or high *C. difficile* toxin production observed in the WD despite a lack of

secondary bile acids, suggesting that the loss of secondary bile acid producing bacteria were not

an important mechanism for the increased mortality and *C. difficile* toxin production observed

with a WD.

While these data show that a high-fat diet increases *C. difficile* toxin production and

mortality in a mouse model of CDI, we did not explore how the composition of fat influences

these factors. Our WD composition represents a typical diet in the United States based on

population survey data and it has 34.5% of calories from fat, with a roughly equivalent

contributions of saturated (~36%), mono-unsaturdated fats (41%) and a lower contribution from

poly-unsaturated fats (~21%). In the low-fat/low-fiber diet, these contrubtions are reversed

(saturated fat ~19% and poly-unsaturated fat ~39%). Further studies to determine if total fat

intake or specific types of fat drive our observed phenotype are needed.

Our data suggests that dietary fiber is critical for the resilience and homogeneity of

response of the gut microbiome after perturbation. In both cohorts of mice fed fiber-deficient

diets, the gut microbiome was significantly more variable and slower to recover to baseline after

perturbation. By supplying the gut with a preferred fuel (fiber) for species associated with health

(e.g. strict anaerobes), the community is able to resist antibiotic induced changes and reconstitute

more quickly once the pressure of antibiotic treatment has been removed. This finding is

consistent with a recent study that showed that fiber supplementation in mice lead to a reduced

disruption of the gut microbiome to disturbance from amoxicillin, and that this was linked with

upregulation of polysaccharide utilization by *Bacteroides thetaiotaomicron*, an intestinal

commensal that individually became less susceptible to amoxicillin in the presence of dietary

fiber *(29)*.

This increased resilience of gut microbiome composition to antibiotic disturbance was

also reflected through levels of the bacterially produced metabolites that we measured. Neither

low-fiber diet was able to maintain butyrate or secondary bile acid production following

perturbation. Based on the correlation between butyrate and DCA concentrations, we speculate

that the lack of butyrate leads to increased luminal oxygen concentrations that are unsuitable for

Clostridium scindens and other secondary bile acid producers. Prior work has shown that

aerobic metabolism of butyrate by intestinal epithelial cells is a key driver of intestinal hypoxia

(30). That there may be increased luminal oxygen concentrations in the low fiber diets is

consistent with our observation of a bloom in Lactobacillales order, which is entirely composed

of facultative anaerobes, after oral antibiotic challenge in both low-fiber diets but not chow.

While our data do not suggest a role for fiber in protection against mortality from CDI in

this mouse model, it would be short-sighted to dismiss the beneficial role of fiber in maintaining

a healthy gut microbiome and resistance to CDI. Our model utilized a rather short-term diet

change and an intense antibiotic regimen that may not correlate well with human circumstances.

We also did not explore diets high in fat and high in fiber, where it is possible that increased

microbiome resilience to antibiotics due to fiber may protect from the detrimental effects of fat.

As discussed above, a fiber-deficient diet has been shown to hinder clearance of *C. difficile* after

challenge *(15)*. This is particularly relevant in a clinical context as recent studies of both

pediatric and adult oncology patients have shown asymptomatic colonization rates with *C.*

difficile of ~30% and ~10%, respectively *(3, 31)*. Further, in pediatric patients it was

demonstrated that the vast majority of "hospital-acquired" CDI may be caused by a strain of *C.*

difficile that is present at admission rather than a strain acquired during the patient stay *(3)*.

Our data along with recently published findings investigating dietary fiber *(15)* and

dietary proline *(4)* intake provides a compelling case that diet should be increasingly targeted as

a prevention and treatment modality for CDI. High-risk populations such as adult and pediatric

oncology patients may benefit from decreased *C. difficile* colonization through increased fiber

intake. For patients with active infection, limiting fat intake could decrease disease severity

while maintaining enteral nutrition.

Materials and Methods

Murine model of CDI: Mice were infected using a widely used murine CDI model *(18)*

with minor modifications. Briefly, 6-week-old female C57BL/6 mice from Taconic Bioscience

(Rensselaer, NY) arrived at University of Colorado on Day -7 of the experiment. Within 24

hours, mouse feed was changed to one of three diets: standard chow, high-fat/low-fiber (Western

Diet), or low-fat/low-fiber diet (all groups n=20). After seven days of the new diet, we placed

mice on a five-antibiotic cocktail (kanamycin (0.4 mg/ml), gentamicin (0.035 mg/ml), colistin

(850 U/ml), metronidazole (0.215 mg/ml), and vancomycin (0.045 mg/ml)) in their drinking

water. Antibiotics were removed for 48 hours, after which we administered an intraperitoneal

injection of clindamycin in normal saline (10 mg/kg body weight). Twenty-four hours after

injection, we gavaged mice with 1.75×10^5 cfu of *C. difficile* VPI 10463. We weighed mice daily

after removal of oral antibiotics and they were euthanized if they lost >15% of body weight or

were moribund. Fecal pellets were collected at arrival (Day -7), after diet change and prior to

oral antibiotics (Day 0) and then daily after removal of oral antibiotics (Day 5-10). In a separate

experiment, we performed the same experimental protocol on 51 mice (chow = 20, low-fat/low-

fiber = 10, WD = 21), but we sacrificed the mice 72 hours after infection and collected cecal

contents for short chain fatty acid, bile acid and toxin quantification. All mouse experiments

were approved by the Institutional Animal Care and Use Committee and complied with their

guidelines and NIH Guide for the Care and Use of Laboratory Animals (IACUC protocol

#00249)

DNA Extraction and Sequencing: Total genomic DNA was extracted from fecal pellets

from a subset of the mice in cohort 1 (chow = 13 mice from four separate cages over two

experiments, WD = 13 mice from four separate cages over two experiments, low-fat/low-fiber =

5 mice from two cages over two experiments) using the DNeasy PowerSoil Kit (Qiagen,

Germantown, MD). Modifications to the standard protocol included a 10 minute incubation at

65°C immediately following the addition of the lysis buffer and the use of a bead mill

homogenizer at 4.5 m/s for 1 min. The V4 variable region of the 16S rDNA gene was targeted

for sequencing (515F: GTGCCAGCMGCCGCGGTAA, 806R:

GGACTACHVGGGTWTCTAAT). The target DNA was amplified using HotMaster Mix

(Quantabio Beverly, MA). Construction of primers and amplification procedures follow the

Earth Microbiome Project guidelines (www.earthmicrobiome.org) *(32)*. Amplified DNA was

quantified in a PicoGreen (ThermoFisher Scientfic) assay and equal quantities of DNA from

each sample was pooled. The pooled DNA was sequenced using a V2 2x250 kit on the Illumina

MiSeq platform (San Diego, CA) at the University of Colorado Anschutz Medical Campus

Genomics and Microarray Core facility.

Sequence Data Analysis: Raw paired-end FASTQ files were processed with QIIME 2

version 2018.8 *(33)*. Denoising was performed with DADA2 *(34)*, a phylogenetic tree was built

using sepp *(35)* and taxonomy was assigned to amplicon sequence variants (ASVs) using the

RDP Classifier *(36)* trained on the Silva version 132 taxonomic database *(37, 38)* using QIIME

2. The data was rarefied at 5,746 sequences per sample. Alpha-diversity was measured by

phylogenetic entropy *(20)* and beta-diversity was determined by weighted UniFrac distances

(39). PCoA of weighted UniFrac plots were constructed using QIIME 2. Metagenomes were

imputed from 16S ASVs using PICRUSt2's default pipeline for stratified genome contributions

(22). Low abundance taxa (<0.01% mean relative abundance) were filtered for analysis of the

butyrogenic coding capacity. Software was installed using Anaconda *(40)*and analysis was

performed on the Fiji compute cluster at the University of Colorado Boulder BioFrontiers

Institute.

***C. difficile* and 16S Quantitative PCR:** qPCR was performed on extracted DNA from

fecal pellets from cohort 1 on day 10 of the experiment with primers targeting the V4 region of

16S rRNA (see above) and strain specific *C. difficile* VPI 10463 *(41)*. The qPCR was prepared

using Kapa SYBR Fast qPCR master mix (Roche Wilmington, MA) and completed on the

CFX96 platform (BioRad Hercules, CA).

SCFA quantification: The SCFAs butyrate, propionate, and acetate were analyzed by

stable isotope GC/MS as previously described *(42)*. Briefly, cecal samples were collected

directly into pre-weighed, sterile Eppendorf tubes and flash frozen at −80°C until processing.

Samples were then subject to an alkylation procedure in which sample and alkylating reagent

were added, vortexed for 1 min, and incubated at 60°C for 25 min. Following cooling and

addition of n-hexane to allow for separation, 170 µL of the organic phase was transferred to an

auto sampler vial and analyzed by GC/MS. Results were quantified in reference to the stable

isotope standard and normalized to sample weight.

Bile acids quantification: *Reagents:* LC/MS grade methanol, acetonitrile, and

isopropanol were obtained from Fisher Scientific (Fairlawn, New Jersey). HPLC grade water

was obtained from Burdick and Jackson (Morristown, New Jersey). Acetic acid, cholic acid,

chenodeoxycholic acid, lithocholic acid, glycolithocholic acid, glycodeoxycholic acid,

glycochenodeoxycholic acid, taurocholic acid and deoxycholic acid were obtained from Sigma

Aldrich (St. Louis, Missouri). Glycolithocholic acid, taurodeoxycholic acid,

taurochenodeoxycholic acid, taurolithocholic acid, alpha-muricholic acid and beta-muricholic

acid were obtained from Cayman Chemical (Ann Arbor, Michigan). Chenodeoxycholic acid-d4

and glycochenodeoxycholic acid-d4 were obtained from Cambridge Isotope labs (Tewksberry,

Massachusetts).

Standards preparation: An internal standard containing 21 µM of chenodeoxycholic

acid-d4 and 21 µM of glycochenodeoxycholic acid–d4 was prepared in 100% methanol. A

combined stock of all bile acid standards was prepared at 0.5mM in 100% methanol. Calibration

working standards were then prepared by diluting the combined stock over a range of 0.05 µM-

50 µM in methanol. A 20 µL aliquot of each calibration working standard was added to 120 µL

of methanol, 50 µL of water and 10 µL of internal standard (200 µL total) to create 10

calibration standards across a calibration range of 0.005 µM-5 µM.

Sample preparation: Fecal samples were prepared using the method described by

Sarafian et al *(19)* with modifications. Briefly, 15-30mg of fecal sample were weighed in a tared

microcentrifuge tube and the weight was recorded. 140 µL of methanol, 15-30 µL of water and

10 µL of internal standard were added. The sample was vortexed for 5 seconds, and then

incubated in a -20°C freezer for 20 minutes. The sample was then centrifuged at 6000RPM for

15 minutes at 4°C. 185-200 µL of the supernatant was transferred to an RSA autosampler vial

(Microsolv Technology Corporation, Leland, NC) for immediate analysis or frozen at -70°C

until analysis.

High performance liquid chromatography/quadrupole time-of-flight mass spectrometry

(HPLC/QTOF): HPLC/QTOF mass spectrometry was performed using the method described by

Sarafian et al *(19)* with modifications. Separation of bile acids was performed on a 1290 series

HPLC from Agilent (Santa Clara, CA) using an Agilent SB-C18 2.1X100mm 1.8 µm column

with a 2.1X 5mm 1.8um guard column. Buffer A consisted of 90:10 water:acetonitrile with 1mM

ammonium acetate adjusted to pH=4 with acetic acid, and buffer B consisted of 50:50

acetonitrile:isopropanol. 10 µL of the extracted sample was analyzed using the following

gradient at a flow rate of 0.6mls/min: Starting composition=10%B, linear gradient from 10-35%

B from 0.1-9.25 minutes, 35-85% B from 9.25-11.5 minutes at 0.65mls/min, 85-100% B from

11.5-11.8 minutes at 0.8mls/min, hold at 100% B from 11.8-12.4 minutes at 1.0ml/min, 100-55%

B from 12.4-12.5 minutes 0.85mls/min, followed by re-equilibration at 10%B from 12.5-15

minutes. The column temperature was held at 60°C for the entire gradient.

Mass spectrometric analysis was performed on an Agilent 6520 quadrupole time of flight

mass spectrometer in negative ionization mode. The drying gas was 300°C at a flow rate of

12mls/min. The nebulizer pressure was 30psi. The capillary voltage was 4000V. Fragmentor

voltage was 200V. Spectra were acquired in the mass range of 50-1700m/z with a scan rate of 2

spectra/sec.

Retention time and m/z for each bile acid was determined by injecting authentic

standards individually. All of the bile acids produced a prominent [M-H]$^-$ ion with negative

ionization. The observed retention time and m/z was then used to create a quantitation method.

Calibration curves for each calibrated bile acid were constructed using Masshunter Quantitative

Analysis software (Aligent Technologies). Bile acid results for feces in pmol/mg were then

quantitated using the following calculation:

Concentration in pmol/mg= $\dfrac{(X_s)(V_t)(D)}{(V_i)(W_s)}$

X_s =pmol on column

V_t =Total volume of concentrated extract (in µL)

D   =Dilution factor if sample was extracted before analysis. If no dilution D=1

V_i =Volume of extract injected (in µL)

W_s =Weight of sample extracted in mg

***C. difficile* toxin B quantification:** Toxin B concentration was determined in cecal

samples from day 3 of infection by comparison to a standard curve using ELISA (tgcBiomics,

Germany). For samples that were too small to weigh accurately, a mass of 5 mg was assigned for

concentration calculation. This mass was selected as it was the lowest weight that could be

accurately determined.

Statistics: Statistical analyses were performed in R (version 3.4.3 "Kite-Eating Tree").

Data were preprocessed using the "tidyverse" suite *(43)*. We used "survminer" and "survival"

libraries to analyze mouse survival *(44, 45)*. All other data were plotted using "ggplot2",

"ggsignif", and "cowplot" *(46-48)*.

Supplementary Materials

541
Figure S1. Cecal bile acid levels by diet. Unconjugated primary bile acids (A), secondary bile

acid (B) and taurine-conjugated bile acids (C). Glycine conjugated bile acids were omitted due to

very low concentrations. a_MCA (alpha muricholic acid); b_MCA (beta muricholic acid); CA

(cholic acid); CDCA (chenodeoxycholic acid) ;UDCA (ursodeoxycholic acid); DCA

(deoxycholic acid); LCA (lithocholic acid); T_b_MCA (tauro-beta muricholic acid); TCA

(taurocholic acid); TCA_3_SO4 (taurocholic acid 3-sulfate); TCDCA (taurochenodeoxycholic

acid); TDCA (taurodeoxycholic acid); TLCA (taurolithocholic acid).

Figure S2. Toxin B ~ TCA-3-SO4 correlations by diet. Multivariate regression was performed

allowing for an interaction between toxin B and diet.

Figure S3: Statistical summary of relative abundances of key taxa from Figure 5a. Significant

differences between groups are noted as calculated with the Wilcoxon rank-sum test. ***:

$p<0.001$. **: $p<0.01$, *: $p<0.05$, ns= non significant.

558   Table. S1. Composition of low-fat/low-fiber/high-sucrose diet

	Low-fat/Low-fiber/High-Sucrose
Fat (% kcal)	17.0
(% SFA)	(19.5)
(% MUFA)	(41.7)
(% PUFA)	(38.8)
Carbohydrates (% kcal)	64.5
(Sucrose)	(26.7)
Protein (%kcal)	18.6
Fiber (g/kg)	50 (cellulose)

561 Table S2. Correlations between *C. difficile* toxin B and metabolites

Formula = toxin_B ~ x	R-squared	p-value
~ diet	0.1748	0.0131

<div align="center">Bile acids</div>

		R-squared	p-value
TCA_3_SO4	~ metab	0.1151	0.0227
	~ metab + diet	0.1701	0.0273
	~ metab + diet + metab*diet	0.1748	0.0131
T_b_MCA	~ metab	-0.0156	0.5078
	~ metab + diet	0.1739	0.0254
	~ metab + diet + metab*diet	0.1208	0.1080
TCA	~ metab	-0.0090	0.4160
	~ metab + diet	0.1775	0.0238
	~ metab + diet + metab*diet	0.1252	0.1017
TCDCA	~ metab	-0.0042	0.3629
	~ metab + diet	0.1804	0.0226
	~ metab + diet + metab*diet	0.1287	0.0970
TLCA	~ metab	-0.0281	0.9000
	~ metab + diet	0.1612	0.0321
	~ metab + diet + metab*diet	0.1078	0.1284
TDCA	~ metab	-0.0253	0.7406
	~ metab + diet	0.1632	0.0309
	~ metab + diet + metab*diet	0.1371	0.0679
GCA	~ metab	-0.0184	0.5577
	~ metab + diet	0.1634	0.0308
	~ metab + diet + metab*diet	0.1125	0.1206
GCDCA	~ metab	0.0679	0.0653
	~ metab + diet	0.1819	0.0220
	~ metab + diet + metab*diet	0.1819	0.0220
GDCA	~ metab	0.0679	0.0653
	~ metab + diet	0.1819	0.0220
	~ metab + diet + metab*diet	0.1819	0.0220
GLCA	~ metab	-0.0212	0.6188
	~ metab + diet	0.1602	0.0327
	~ metab + diet + metab*diet	0.1867	0.0302
a_MCA	~ metab	-0.0283	0.9274
	~ metab + diet	0.1840	0.0211

		R-squared	p-value
	~ metab + diet + metab*diet	0.1332	0.0912
b_MCA	~ metab	-0.0134	0.4740
	~ metab + diet	0.1912	0.0184
	~ metab + diet + metab*diet	0.1525	0.0694
CA	~ metab	-0.0123	0.4575
	~ metab + diet	0.1677	0.0285
	~ metab + diet + metab*diet	0.1158	0.1154
CDCA	~ metab	0.0528	0.0916
	~ metab + diet	0.1602	0.0327
	~ metab + diet + metab*diet	0.1104	0.1241
UDCA	~ metab	-0.0255	0.7483
	~ metab + diet	0.1835	0.0213
	~ metab + diet + metab*diet	0.1312	0.0937
DCA	~ metab	0.0197	0.1977
	~ metab + diet	0.1628	0.0312
	~ metab + diet + metab*diet	0.1450	0.0599
LCA	~ metab	0.0259	0.1705
	~ metab + diet	0.1628	0.0312
	~ metab + diet + metab*diet	0.2009	0.0237

Short chain fatty acids

		R-squared	p-value
acetate	~ metab	-0.0250	0.6830
	~ metab + diet	0.1594	0.0389
	~ metab + diet + metab*diet	0.1020	0.1500
propinoate	~ metab	-0.0158	0.4974
	~ metab + diet	0.1654	0.0351
	~ metab + diet + metab*diet	0.1246	0.1135
butyrate	~ metab	-0.0198	0.5641
	~ metab + diet	0.1591	0.0391
	~ metab + diet + metab*diet	0.1017	0.1504

References and Notes:

1. F. C. Lessa, Y. Mu, W. M. Bamberg, Z. G. Beldavs, G. K. Dumyati, J. R. Dunn, M. M. Farley, S. M. Holzbauer, J. I. Meek, E. C. Phipps, L. E. Wilson, L. G. Winston, J. A. Cohen, B. M. Limbago, S. K. Fridkin, D. N. Gerding, L. C. McDonald, Burden of Clostridium difficile infection in the United States, *N Engl J Med* **372**, 825–834 (2015).

2. D. A. Leffler, J. T. Lamont, Clostridium difficile infection, *N Engl J Med* **372**, 1539–1548 (2015).

3. G. N. Al-Rawahi, A. Al-Najjar, R. McDonald, R. J. Deyell, G. R. Golding, R. Brant, P. Tilley, E. Thomas, S. R. Rassekh, A. O'Gorman, P. Wong, L. Turnham, S. Dobson, Pediatric oncology and stem cell transplant patients with healthcare-associated Clostridium difficileinfection were already colonized on admission, *Pediatr Blood Cancer* **353**, e27604–5 (2019).

4. E. J. Battaglioli, V. L. Hale, J. Chen, P. Jeraldo, C. Ruiz-Mojica, B. A. Schmidt, V. M. Rekdal, L. M. Till, L. Huq, S. A. Smits, W. J. Moor, Y. Jones-Hall, T. Smyrk, S. Khanna, D. S. Pardi, M. Grover, R. Patel, N. Chia, H. Nelson, J. L. Sonnenburg, G. Farrugia, P. C. Kashyap, Clostridioides difficile uses amino acids associated with gut microbial dysbiosis in a subset of patients with diarrhea, *Science Translational Medicine* **10**, eaam7019 (2018).

5. G. E. Bignardi, Risk factors for Clostridium difficile infection, *Journal of Hospital Infection* **40**, 1–15 (1998).

6. R. Fekety, L. V. McFarland, C. M. Surawicz, R. N. Greenberg, G. W. Elmer, M. E. Mulligan, Recurrent Clostridium difficile diarrhea: characteristics of and risk factors for patients enrolled in

a prospective, randomized, double-blinded trial, *Clinical Infectious Diseases* **24**, 324–333

(1997).

7. C. A. Lozupone, J. Stombaugh, A. Gonzalez, G. Ackermann, D. Wendel, Y. Vázquez-Baeza,

J. K. Jansson, J. I. Gordon, R. Knight, Meta-analyses of studies of the human microbiota,

Genome Research **23**, 1704–1714 (2013).

8. C. Staley, B. P. Vaughn, C. T. Graiziger, S. Singroy, M. J. Hamilton, D. Yao, C. Chen, A.

Khoruts, M. J. Sadowsky, Community dynamics drive punctuated engraftment of the fecal

microbiome following transplantation using freeze-dried, encapsulated fecal microbiota, *Gut*

Microbes **131**, 1–13 (2017).

9. C. Staley, C. R. Kelly, L. J. Brandt, A. Khoruts, M. J. Sadowsky, Complete Microbiota

Engraftment Is Not Essential for Recovery from Recurrent Clostridium difficileInfection

following Fecal Microbiota Transplantation, *mBio* **7**, e01965–16–9 (2016).

10. A. Weingarden, A. Gonzalez, Y. Vázquez-Baeza, S. Weiss, G. Humphry, D. Berg-Lyons, D.

Knights, T. Unno, A. Bobr, J. Kang, A. Khoruts, R. Knight, M. J. Sadowsky, Dynamic changes

in short- and long-term bacterial composition following fecal microbiota transplantation for

recurrent Clostridium difficile infection, *Microbiome* **3**, 10 (2015).

11. S. K. Dutta, M. Girotra, S. Garg, A. Dutta, E. C. von Rosenvinge, C. Maddox, Y. Song, J. G.

Bartlett, R. Vinayek, W. F. Fricke, Efficacy of Combined Jejunal and Colonic Fecal Microbiota

Transplantation for Recurrent Clostridium difficile Infection, *Clinical Gastroenterology and*

Hepatology **12**, 1–5 (2014).

12. J. R. Allegretti, S. Kearney, N. Li, E. Bogart, K. Bullock, G. K. Gerber, L. Bry, C. B. Clish,

E. Alm, J. R. Korzenik, Recurrent Clostridium difficile infection associates with distinct bile acid

and microbiome profiles, **43**, 1142–1153 (2016).

13. J. A. Sorg, A. L. Sonenshein, Bile salts and glycine as cogerminants for Clostridium difficile

spores, *J. Bacteriol.* **190**, 2505–2512 (2008).

14. C. G. Buffie, V. Bucci, R. R. Stein, P. T. McKenney, L. Ling, A. Gobourne, D. No, H. Liu,

M. Kinnebrew, A. Viale, E. Littmann, M. R. M. van den Brink, R. R. Jenq, Y. Taur, C. Sander, J.

R. Cross, N. C. Toussaint, J. B. Xavier, E. G. Pamer, Precision microbiome reconstitution

restores bile acid mediated resistance to Clostridium difficile, *Nature* **517**, 205–208 (2015).

15. A. J. Hryckowian, W. Van Treuren, S. A. Smits, N. M. Davis, J. O. Gardner, D. M. Bouley,

J. L. Sonnenburg, Microbiota-accessible carbohydrates suppress Clostridium difficile infection in

a murine model, *Nat Microbiol* **372**, 2369 (2018).

16. V. C. Antharam, E. C. Li, A. Ishmael, A. Sharma, V. Mai, K. H. Rand, G. P. Wang, Intestinal

dysbiosis and depletion of butyrogenic bacteria in Clostridium difficile infection and nosocomial

diarrhea, *Journal of Clinical Microbiology* **51**, 2884–2892 (2013).

17. S. Devkota, Y. Wang, M. W. Musch, V. Leone, H. Fehlner-Peach, A. Nadimpalli, D. A.

Antonopoulos, B. Jabri, E. B. Chang, Dietary-fat-induced taurocholic acid promotes pathobiont

expansion and colitis in Il10−/− mice, *Nature* **65**, 411–6 (2012).

18. X. Chen, K. Katchar, J. D. Goldsmith, N. Nanthakumar, A. Cheknis, D. N. Gerding, C. P.

Kelly, A Mouse Model of Clostridium difficile–Associated Disease, *Gastroenterology* **135**,

1984–1992 (2008).

19. M. H. Sarafian, M. R. Lewis, A. Pechlivanis, S. Ralphs, M. J. W. McPhail, V. C. Patel, M.-E.

Dumas, E. Holmes, J. K. Nicholson, Bile acid profiling and quantification in biofluids using

ultra-performance liquid chromatography tandem mass spectrometry, *Anal. Chem.* **87**, 9662–

9670 (2015).

20. B. Allen, M. Kon, Y. Bar-Yam, A new phylogenetic diversity measure generalizing the

shannon index and its application to phyllostomid bats, *Am. Nat.* **174**, 236–243 (2009).

21. M. Kriss, K. Z. Hazleton, N. M. Nusbacher, C. G. Martin, C. A. Lozupone, Low diversity gut

microbiota dysbiosis: drivers, functional implications and recovery, *Curr Opin Microbiol* **44**, 34–

40 (2018).

22. M. G. I. Langille, J. Zaneveld, J. G. Caporaso, D. McDonald, D. Knights, J. A. Reyes, J. C.

Clemente, D. E. Burkepile, R. L. Vega Thurber, R. Knight, R. G. Beiko, C. Huttenhower,

Predictive functional profiling of microbial communities using 16S rRNA marker gene

sequences, *Nature Biotechnology* **31**, 814–821 (2013).

23. J. M. Ridlon, D. J. Kang, P. B. Hylemon, Bile salt biotransformations by human intestinal

bacteria, (2006).

24. M. Vital, A. C. Howe, J. M. Tiedje, Revealing the bacterial butyrate synthesis pathways by

analyzing (meta)genomic data, *mBio* **5**, e00889–e00889–14 (2014).

25. M. Vital, J. Gao, M. Rizzo, T. Harrison, J. M. Tiedje, Diet is a major factor governing the

fecal butyrate-producing community structure across Mammalia, Aves and Reptilia, *ISME J* **9**,

832–843 (2015).

26. N. Kumar, H. P. Browne, E. Viciani, S. C. Forster, S. Clare, K. Harcourt, M. D. Stares, G.

Dougan, D. J. Fairley, P. Roberts, M. Pirmohamed, M. R. J. Clokie, M. B. F. Jensen, K. R.

Hargreaves, M. Ip, L. H. Wieler, C. Seyboldt, T. Norén, T. V. Riley, E. J. Kuijper, B. W. Wren,

T. D. Lawley, Adaptation of host transmission cycle during Clostridium difficile speciation,

Nature Genetics **51**, 1315–1320 (2019).

27. T. L. Blankenship-Paris, B. J. Walton, Y. O. Hayes, J. Chang, Clostridium difficile infection

in hamsters fed an atherogenic diet, *Vet. Pathol.* **32**, 269–273 (1995).

28. T. L. Blankenship-Paris, J. Chang, F. G. Dalldorf, P. H. Gilligan, In vivo and in vitro studies

of Clostridium difficile-induced disease in hamsters fed an atherogenic, high-fat diet, *Lab. Anim.*

Sci. **45**, 47–53 (1995).

29. D. J. Cabral, S. Penumutchu, E. M. Reinhart, C. Zhang, B. J. Korry, J. I. Wurster, R. Nilson,

A. Guang, W. H. Sano, A. D. Rowan-Nash, H. Li, P. Belenky, Microbial Metabolism Modulates

Antibiotic Susceptibility within the Murine Gut Microbiome, *Cell Metabolism* **30**, 800–823.e7

(2019).

30. C. J. Kelly, L. Zheng, E. L. Campbell, B. Saeedi, C. C. Scholz, A. J. Bayless, K. E. Wilson,

L. E. Glover, D. J. Kominsky, A. Magnuson, T. L. Weir, S. F. Ehrentraut, C. Pickel, K. A. Kuhn,

J. M. Lanis, V. Nguyen, C. T. Taylor, S. P. Colgan, Crosstalk between Microbiota-Derived

Short-Chain Fatty Acids and Intestinal Epithelial HIF Augments Tissue Barrier Function, *Cell*

Host Microbe **17**, 662–671 (2015).

31. C. M. Cannon, J. S. Musuuza, A. K. Barker, M. Duster, M. B. Juckett, A. E. Pop-Vicas, N.

Safdar, Risk of Clostridium difficile Infection in Hematology-Oncology Patients Colonized With

Toxigenic C. difficile, *Infect Control Hosp Epidemiol* **38**, 718–720 (2017).

32. L. R. Thompson, J. G. Sanders, D. McDonald, A. Amir, J. Ladau, K. J. Locey, R. J. Prill, A.

Tripathi, S. M. Gibbons, G. Ackermann, J. A. Navas-Molina, S. Janssen, E. Kopylova, Y.

Vázquez-Baeza, A. Gonzalez, J. T. Morton, S. Mirarab, Z. Zech Xu, L. Jiang, M. F. Haroon, J.

Kanbar, Q. Zhu, S. Jin Song, T. Kosciolek, N. A. Bokulich, J. Lefler, C. J. Brislawn, G.

Humphrey, S. M. Owens, J. Hampton-Marcell, D. Berg-Lyons, V. McKenzie, N. Fierer, J. A.

Fuhrman, A. Clauset, R. L. Stevens, A. Shade, K. S. Pollard, K. D. Goodwin, J. K. Jansson, J. A.

Gilbert, R. Knight, Earth Microbiome Project Consortium, A communal catalogue reveals

Earth's multiscale microbial diversity, *Nature* **551**, 457–463 (2017).

33. E. Bolyen, J. R. Rideout, M. R. Dillon, N. A. Bokulich, C. C. Abnet, G. A. Al-Ghalith, H.

Alexander, E. J. Alm, M. Arumugam, F. Asnicar, Y. Bai, J. E. Bisanz, K. Bittinger, A. Brejnrod,

C. J. Brislawn, C. T. Brown, B. J. Callahan, A. M. Caraballo-Rodríguez, J. Chase, E. K. Cope, R.

Da Silva, C. Diener, P. C. Dorrestein, G. M. Douglas, D. M. Durall, C. Duvallet, C. F.

Edwardson, M. Ernst, M. Estaki, J. Fouquier, J. M. Gauglitz, S. M. Gibbons, D. L. Gibson, A.

Gonzalez, K. Gorlick, J. Guo, B. Hillmann, S. Holmes, H. Holste, C. Huttenhower, G. A.

Huttley, S. Janssen, A. K. Jarmusch, L. Jiang, B. D. Kaehler, K. B. Kang, C. R. Keefe, P. Keim,

S. T. Kelley, D. Knights, I. Koester, T. Kosciolek, J. Kreps, M. G. I. Langille, J. Lee, R. Ley, Y.-

X. Liu, E. Loftfield, C. Lozupone, M. Maher, C. Marotz, B. D. Martin, D. McDonald, L. J.

McIver, A. V. Melnik, J. L. Metcalf, S. C. Morgan, J. T. Morton, A. T. Naimey, J. A. Navas-

Molina, L. F. Nothias, S. B. Orchanian, T. Pearson, S. L. Peoples, D. Petras, M. L. Preuss, E.

Pruesse, L. B. Rasmussen, A. Rivers, M. S. Robeson, P. Rosenthal, N. Segata, M. Shaffer, A.

Shiffer, R. Sinha, S. J. Song, J. R. Spear, A. D. Swafford, L. R. Thompson, P. J. Torres, P. Trinh,
A. Tripathi, P. J. Turnbaugh, S. Ul-Hasan, J. J. J. van der Hooft, F. Vargas, Y. Vázquez-Baeza,
E. Vogtmann, M. von Hippel, W. Walters, Y. Wan, M. Wang, J. Warren, K. C. Weber, C. H. D.
Williamson, A. D. Willis, Z. Z. Xu, J. R. Zaneveld, Y. Zhang, Q. Zhu, R. Knight, J. G. Caporaso,
Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2,
Nature Biotechnology **37**, 852–857 (2019).

34. B. J. Callahan, P. J. McMurdie, M. J. Rosen, A. W. Han, A. J. A. Johnson, S. P. Holmes,
DADA2: High-resolution sample inference from Illumina amplicon data, *Nat Meth* **13**, 581–583
(2016).

35. S. Janssen, D. McDonald, A. Gonzalez, J. A. Navas-Molina, L. Jiang, Z. Z. Xu, K. Winker,
D. M. Kado, E. Orwoll, M. Manary, S. Mirarab, R. Knight, N. Chia, Ed. Phylogenetic Placement
of Exact Amplicon Sequences Improves Associations with Clinical Information, *mSystems* **3**,
581 (2018).

36. Q. Wang, G. M. Garrity, J. M. Tiedje, J. R. Cole, Naive Bayesian classifier for rapid
assignment of rRNA sequences into the new bacterial taxonomy, *Appl. Environ. Microbiol.* **73**,
5261–5267 (2007).

37. C. Quast, E. Pruesse, P. Yilmaz, J. Gerken, T. Schweer, P. Yarza, J. Peplies, F. O. Glöckner,
The SILVA ribosomal RNA gene database project: improved data processing and web-based
tools, *Nucleic Acids Res* **41**, D590–6 (2013).

38. P. Yilmaz, L. W. Parfrey, P. Yarza, J. Gerken, E. Pruesse, C. Quast, T. Schweer, J. Peplies,

W. Ludwig, F. O. Glöckner, The SILVA and "All-species Living Tree Project (LTP)" taxonomic

frameworks, *Nucleic Acids Res* **42**, D643–8 (2014).

39. C. A. Lozupone, M. Hamady, S. T. Kelley, R. Knight, Quantitative and Qualitative

Diversity Measures Lead to Different Insights into Factors That Structure Microbial

Communities, *Appl. Environ. Microbiol.* **73**, 1576–1585 (2007).

40. Anaconda Software Distribution (available at https://anaconda.com).

41. L. Etienne-Mesmin, B. Chassaing, O. Adekunle, L. M. Mattei, F. D. Bushman, A. T.

Gewirtz, Toxin-positive Clostridium difficile latently infect mouse colonies and protect against

highly pathogenic C. difficile, *Gut* **67**, 860–871 (2018).

42. C. J. Kelly, E. E. Alexeev, L. Farb, T. W. Vickery, L. Zheng, C. Eric L, D. A. Kitzenberg, K.

D. Battista, D. J. Kominsky, C. E. Robertson, D. N. Frank, S. P. Stabler, S. P. Colgan, Oral

vitamin B12 supplement is delivered to the distal gut, altering the corrinoid profile and

selectively depleting Bacteroides in C57BL/6 mice, *Gut Microbes* **87**, 1–9 (2019).

43. H. Wickham, Tidyverse (available at https://CRAN.R-project.org/package=tidyverse).

44. A. Kassambara, M. Kosinski, P. Biecek, S. Fabian, survminer: Drawing Survival Curves

using "ggplot2" (available at http://www.sthda.com/english/rpkgs/survminer/).

45. T. M. Therneau, T. Lumley, survival: Survival Analysis (available at

https://github.com/therneau/survival).

46. H. Wickham, *ggplot2* (Springer, 2010).

47. C. Ahlmann-Eltze, ggsignif: Significance Brackets for "ggplot2" (available at

https://github.com/const-ae/ggsignif).

48. C. O. Wilke, cowplot: Streamlined Plot Theme and Plot Annotations for "ggplot2" (available

at https://wilkelab.org/cowplot).

49. A. Gonzalez, J. A. Navas-Molina, T. Kosciolek, D. McDonald, Y. Vázquez-Baeza, G.

Ackermann, J. DeReus, S. Janssen, A. D. Swafford, S. B. Orchanian, J. G. Sanders, J.

Shorenstein, H. Holste, S. Petrus, A. Robbins-Pianka, C. J. Brislawn, M. Wang, J. R. Rideout, E.

Bolyen, M. Dillon, J. G. Caporaso, P. C. Dorrestein, R. Knight, Qiita: rapid, web-enabled

microbiome meta-analysis, *Nature* **15**, 796–798 (2018).

Acknowledgments: We would like to thank Jordi Lanis and Sean Colgan for advice on the

employed CDI mouse model and Sally Stabler for her assistance in measuring SCFAs. **Funding:**

We would like to thank the University of Colorado Department of Medicine's Outstanding Early

Career Science Award for supporting this work as well as support to Keith Hazleton from the

Institutional Training Grant for Pediatric Gastroenterology from NIDDK (5T32-DK067009-12)

and Clinical Fellow Awards from the Cystic Fibrosis Foundation (HAZLET18DO and

HAZLET19DO). Kathleen Arnolds was supported by T32-AI007405 Training Program in

Immunology. High performance computing was supported by a cluster at the University of

Colorado Boulder funded by National Institutes of Health 1S10OD012300. **Author**

contributions: KH conceived of and conducted experiments, analyzed data and wrote the paper;

CM analyzed sequence data and made figures; KA generated and analyzed data from qPCR

experiments and toxin ELISA; NN generated 16S rRNA sequence and qPCR data and aided in

analysis and results interpretation; NMH aided in mouse experiments, 16S rRNA sequencing and

generated data from toxin ELISA; NR and MAworked with KH to develop a novel bile acid

panel and aided in analysis and interpretation of results, CL directed and contributed to all

aspects of the project. All authors contributed to the manuscript. **Competing interests:** The

authors declare no competing financial interests. **Data and materials availability:** The 16S

rRNA has been deposited in QIITA *(49)*(Qiita Study ID: 12849) and at EBI (pending)