Dietary fat promotes antibiotic-induced *Clostridioides difficile* mortality in mice

Authors: Keith Z. Hazleton[1,2,#], Casey G. Martin[3], David J Orlicky[4], Kathleen L. Arnolds[3], Nichole M. Nusbacher[5], Nancy Moreno-Huizar[5], Michael Armstrong[6], Nichole Reisdorph[6], Catherine A. Lozupone[5]*

Affiliations:

[1] Department of Pediatrics, Section of Gastroenterology, Hepatology and Nutrition. University of Colorado, Denver Anschutz Medical Campus, Aurora, CO USA 80045.

[2] Digestive Health Institute, Children's Hospital Colorado, Aurora, CO USA 80045.

[#] Current address, Department of Pediatrics, Division of Gastroenterology, Hepatology and Nutrition, University of Arizona, Tucson, AZ 85719

[3] Department of Immunology and Microbiology, University of Colorado, Denver Anschutz Medical Campus, Aurora, CO USA 80045.

[4] Department of Pathology, University of Colorado, Denver Anschutz Medical Campus, Aurora, CO USA 80045.

[5] Department of Medicine, Division of Biomedical Informatics and Personalized Medicine, University of Colorado, Denver Anschutz Medical Campus, Aurora, CO USA 80045.

[6] Skaggs School of Pharmacy and Pharmaceutical Sciences, University of Colorado Anschutz Medical Campus, Aurora, CO USA 80045.

*To whom correspondence should be addressed: Catherine.lozupone@cuanschutz.edu

One Sentence Summary: A high-fat/low-fiber Western type diet promoted mortality in a mouse model of antibiotic-induced *C. difficile* infection compared to a low-fat/high-fiber diet and chow diet, suggesting that lower dietary fat may be an effective strategy for preventing C. difficile pathology.

Abstract: *Clostridioides difficile* infection (CDI), is the leading cause of hospital-acquired

diarrhea and emerging evidence has linked dietary components with CDI pathogenesis,

suggesting that dietary modulation may be an effective strategy for prevention. Here, we show

that mice fed a high-fat/low-fiber "Western type" diet (WD) had dramatically increased mortality

in a murine model of antibiotic-induced CDI compared to a low-fat/low-fiber (LF/LF) diet and

standard mouse chow controls. We found that the WD had a pro- *C. difficile* bile acid

composition that was driven in part by higher levels of primary bile acids that are produced to

digest fat, and a lower level of secondary bile acids that are produced by the gut microbiome.

This lack of secondary bile acids was associated with a greater disturbance to the gut

microbiome with antibiotics in both the WD and LF/LF diet compared to mouse chow. Mice fed

the WD also had the highest level of toxin TcdA just prior to the onset of mortality, but not of

TcdB or increased inflammation. These findings indicate that dietary intervention to decrease fat

may complement previously proposed dietary intervention strategies to prevent CDI in high-risk

individuals.

Introduction

Clostridioides difficile infection (CDI) is an important cause of morbidity and mortality, with 500,000 cases every year causing 30,000 deaths per year in the US alone[1]. Alarmingly, there has been a steady increase in the number of new infections in spite of prevention efforts in hospitals that have focused largely on increased sanitation and antibiotic stewardship[2]. C. difficile induced pathology has been linked to the production of two different toxins, TcdA and TcdB, which can directly induce intestinal damage and inflammation [3].

A complex gut microbiome is protective against CDI[4]. Illnesses associated with reduced gut microbiome diversity, such as Inflammatory Bowel Disease[5] increase risk of CDI, as does broad spectrum antibiotic usage, such as clindamycin, beta-lactams, and fluoroquinolones [6,7]. Antibiotics have been shown to predispose mice to CDI via modified metabolic activity of the altered gut microbiome [8] . Individuals with recurrent CDI (rCDI) typically have microbiomes with greatly reduced complexity and altered composition [9–11]. The gut microbiome provides protection from CDI in part through metabolism of primary bile acids, which are excreted by the liver into the intestine where they play a central role in fat digestion[12]. The primary bile acids taurocholic acid (TCA) and cholic acid (CA) can promote the germination of C. difficile spores. However a healthy gut microbiome can metabolize TCA and CA into the secondary bile acid deoxycholate (DCA), a derivative that can arrest the growth of vegetative C. difficile [13]. Accordingly, prior studies have shown that secondary bile acid producers such as Clostridium scindens can protect against CDI in mice[14]. Short chain fatty acids (SCFA), which are commensal microbial products of fermentation of dietary microbial accessible carbohydrates (MACs; e.g. soluble fibers such as inulin), have also been shown to directly suppress C. difficile growth in vitro [15]and are decreased in individuals with rCDI [16].

Recent studies conducted in mouse models of antibiotic-induced CDI have suggested that

diet modulation has the potential to be an effective prevention strategy for antibiotic-induced

CDI. Diets high in MACs[15] and low in proline [4] reduced *C. difficile* colonization and persistence.

Excess dietary zinc reduces the threshold of antibiotics needed to confer susceptibility to CDI

and increased cecal inflammation and toxin activity [17]. High-fat/high-protein diets [18] and high-fat

induced obesity [19] resulted in more severe disease and/or increased mortality. Furthermore, mice

fed a protein deficient defined diet, had increased survival, decreased weight loss, and decreased

overall disease severity [20].

Given that primary bile acids play a central role in fat digestion, increase with diets high

in saturated fat[21] and are a germination factor for *C. difficile* spores, we became interested in

investigating a role for dietary fat in antibiotic-induced CDI pathogenesis. We hypothesized that

a high-fat diet coupled with low-fiber in the context of antibiotic treatment would provide a

"double hit" for shifting towards a pro-*C. difficile* bile acid pool – with dietary fat increasing

excretion of pro- *C. difficile* primary bile acids into the gut and increased antibiotic-induced gut

microbiome disturbance decreasing their conversion into protective secondary bile acids. We

found that high dietary fat content in the context of a low fiber diet (a high-fat/low-fiber Western

Diet; WD) induced high mortality from CDI in an antibiotic-induced *C. difficile* model. This

higher mortality was linked with higher levels of *C. difficile* toxin TcdA, but not with higher

levels of TcdB or increased intestinal inflammation just prior to the onset of mortality. Bile acid

pools were strongly influenced by diet to a pro-*C. difficile* composition but more work needs to

be done to determine the degree to which these differences were driving the higher mortality

observed with the WD. Our work suggests that dietary interventions to decrease fat intake may

complement previously proposed strategies that target fiber and protein to prevent CDI in high-

risk individuals.

Results

High dietary fat in the context of low dietary fiber causes increased mortality in murine

antibiotic-induced CDI

To understand the effects of dietary fat on CDI, we used an established murine model of

antibiotic-induced CDI[22]. Specifically, conventional 6-week-old old female C57-bl6 mice were

fed 1 of 3 diets: 1) conventional mouse chow that is low-fat/high-fiber, 2) a purified "Western"

diet (WD) that had ~2x the content of fat with increased ratio of saturated-to-unsaturated fat

compared to chow and only insoluble cellulose as a source of fiber, and 3) a similar purified diet

as the WD, but with a lower fat content, similar to chow (low-fat/low-fiber; LF/LF) (Table 1;

Table S2). This WD composition represents a typical diet in the United States based on

population survey data with 34.5% of calories from fat, with a roughly equivalent contributions

of saturated (~36%), mono-unsaturated fats (41%) and a lower contribution from poly-

unsaturated fats (~21%). In the LF/LF diet, these contributions are reversed (saturated fat ~19%

and poly-unsaturated fat ~39%). One week after diet switch, mice were treated with a cocktail of

antibiotics in their drinking water for 5 days (kanamycin, gentamicin, colistin, metronidazole and

vancomycin) followed by an injection of clindamycin and gavage with *C. difficile* VPI 10463

(Fig. 1A). The experiments were carried out for up to 21 days past *C. difficile* gavage, allowing

us to assay effects on mortality and relate these to fecal microbiome composition. Experiments

were conducted on 2 separate cohorts – 2 cages of five mice per diet, with technical replicates of

mice with a total sample size of 20 per diet (Figure 1, Table S2).

The WD-fed mice showed a marked increase in mortality as compared to both the LF/LF

(HR 7.403 p = 0.0041) and chow-fed mice (HR 4.95 p = 0.00208) upon *C. difficile* exposure.

Mortality onset began at Day 4 in the WD and chow-fed mice and then continued to Day 8 in the

WD and to Day 6 in the chow-fed mice before stabilizing with the remaining mice appearing to

recover. The LF/LF diet-fed mice showed survival levels comparable to the chow-fed mice with

a slightly delayed onset of mortality (Fig. 1B). Qualitatively, WD-fed mice had more purulent

and liquid stools, and poorer grooming than LF/LF and chow-fed mice starting 2 days after

infection. Because our WD and LF/LF diet differed in sucrose content, we also tested a fourth

diet that was low in fat and fiber, but with sucrose equivalent to the WD (Table S1). Sucrose did

not appear to play a role in the increased mortality observed in the WD, as 100% survival was

observed in mice fed the low-fat/low-fiber/low-sucrose diet. (n = 10, one cage with 5 mice in two

separate experiments).

Table 1

	Chow diet	**WD**	**LF/LF diet**
Fat (% kcal) *(% SFA)* *(% MUFA)* *(% PUFA)*	16 *(N/A)*	34.5 *(36.2)* *(41.3)* *(22.5)*	17.2 *(19.5)* *(41.7)* *(38.8)*
Carbohydrates (% kcal) *(Sucrose)*	60 *(0)*	50 *(23.4)*	63.9 *(10.6)*
Protein (%kcal)	24	15.5	18.8
Fiber (g/kg)	137	50 (cellulose)	50 (cellulose)

Figure 1: *Experimental design of murine model of antibiotic-induced CDI and survival curves.*

A) *C. difficile* challenge experimental design. The figure legend at the left panel indicates the samples sizes for 2 cohorts: more information on batching and n's per assay is given in Table S2). Cohort 1 was followed for 13 days post *C. difficile* gavage to monitor survival and gut microbiome composition over time. Cohort 2 was sacrificed at 3 days post *C. difficile* gavage to collect cecal contents for measurement of metabolites and toxin and colon and cecal mucosa for histopathology (some assays were only conducted on a subset of Cohort 2; but all in at least 2 independent experiments; see Table S2 for details). Grey and orange boxes indicate the timepoints at which samples were collected for the respective cohorts. **B)** Survival curves on the 3 diets. Statistical significance as assessed by log-rank comparison is indicated.

WD associated with increased C. difficile toxin TcdA but not TcdB or intestinal inflammation

To further explore the mechanisms of increased mortality in the WD-fed mice by

assessing factors that required the collection of host tissues, we conducted a second set of

experiments in which mice were sacrificed at day 3 post *C. difficile* gavage (cohort 2; Figure 1).

We chose 3 days post *C. difficile* gavage because this was just prior to the observed onset of

mortality in the first cohort across all 3 diets (Fig. 1B), and because we felt it was important to

compare all mice at a standard time point.

We measured cecal levels of *C. difficile* Toxins A (TcdA) and B (TcdB) by ELISA (see

Table S2 for cohort size and batch information). Interestingly, TcdA and not TcdB showed

differences with diet consistent with mortality patterns, with TcdA being much higher in the WD

compared to the LF/LF diet. The LF/LF diet also had slightly lower levels of TcdA than the

chow diet (Figure 2A), which is consistent with a delayed onset of mortality in the LF/LF diet

compared to chow (Figure 1). To understand whether intestinal inflammation was related to

toxin levels and differed between diets, we evaluated the transverse colon and cecum by

histology (Fig. 2B, Fig. S1).   Cecal and transverse colon tissues from mice sacrificed three days

post-infection with *C. difficile* were fixed and stained with hematoxylin and eosin and were

scored by the Barthel and Dieleman scoring systems respectively by a trained histologist blinded

to the treatments and grouping of individuals [23,24]. The cecum and distal colon samples showed

mild to moderate inflammation, but the level of inflammation did not differ across diet groups (p

= 0.1375, Fig 2B; representative histology Figure S1). Cecal levels of TcdB and not TcdA

strongly correlated with cecal inflammation (Figure 2; Table S3). Taken together, our results

suggest that although TcdB levels are associated with higher intestinal inflammation in these

mice just prior to onset of mortality, the differences cannot solely explain the increased mortality

observed in the WD compared to low fat diets. Our data support a potential role for TcdA in the

increased mortality with the WD, but via a mechanism independent of intestinal inflammation.

Figure 2: *Toxin and histopathology scores by diet.* **(A)** TcdA and tcdB production across diets as

determined by ELISA. **(B)** Histologic inflammation scores across diets as determined by a

blinded histologist. Significant differences calculated with a Kruskal-Wallis and Dunn's post hoc

test. Kruskal-Wallis p-values were corrected for multiple comparisons with the FDR algorithm

of Benjamini and Hochberg [25]. Median and Interquartile Range (IQR) indicated. **(C)** Linear

regression of cecal histopathology against *C. difficile* tcdB burden. (p = 0.001 with model

cecum_infl ~ tcdB and FDR correction).

Cecal levels of bile acids and their relationship to diet, cecal levels of C. difficile toxins, and

inflammation

To further explore mechanism, we used targeted LC/MS to measure the levels of a pool

of 13 different bile acids in the aspirated cecal contents of a separate cohort of mice that were

sacrificed at 3 days post infection (Table S2). Bile acids have a complex relationship with *C.*

difficile germination and growth[13,26–29]. The primary bile acids TCA and CA can promote the

germination of *C. difficile* spores *in vitro*[13] and primary bile acids including CA are elevated in

individuals with first time or rCDI compared to controls [30, 31]. The primary bile acid

chenodeoxycholic acid (CDCA) can block TCA-induced spore germination[27,28] and another

primary bile acid, Ursodeoxycholic acid (UDCA), can inhibit both *C. difficile* spore germination

and *C. difficile* growth [29]. Furthermore, the murine primary bile acids alpha muricholic acid

(a_MCA) and beta muricolic acid (b_MCA) can inhibit *C. difficile* spore germination and growth

[32]. Of particular interest in this study are also the secondary bile acids DCA and lithocholate

(LCA); these molecules are produced by the metabolic transformation of primary bile acids by

intestinal microbes [14], can arrest the growth of vegetative *C. difficile* [13], and are lower in

individuals with CDI [30,31]. We measured the levels of these bile acids with known effects on *C.*

difficile as well as 5 other taurine-conjugated bile acids (Figure S2). To consider known effects

of bile acids on *C. difficile* growth and germination in our analyses, we binned the bile acids that

were inhibitors of *C. difficile* germination and/or growth (*CDCA, UDCA, a_MCA, b_MCA, LCA,*

DCA), and *C. difficile* germination promoters (*TCA* and *CA*). We also evaluated the ratio of *C.*

difficile promoters to inhibitors, as has been done previously [19].

When comparing across diets, we were most interested in bile acid measures that showed

differential levels between the WD and both the LF/LF and chow diets, since the WD had high

CDI mortality compared to both the LF/LF and chow diets. *C. difficile* inhibitors were

significantly lower in the WD compared to the chow diet, but there was not a difference between

the WD and the LF/LF diet (Figure 3A). *C. difficile* promoters had significantly higher levels in

the WD compared to chow but not compared to the LF/LF diet (Figure 3A). Interestingly, the

ratio of promoters:inhibitors was significantly higher in the WD compared to both the LF/LF diet

and chow, consistent with mortality differences. We also analyzed whether each of the 13 bile

acids individually differed across diets, and diet significantly affected the levels of most (Figure

S2). However, none were individually significantly different in the WD compared to both the

LF/LF and chow diet. While regressing cecal and colon inflammation scores (Figure 3B) and

toxins tcdA and tcdB (data not shown) against bile acid summary measures, the only significant

relationship observed was a negative correlation between *C. difficile* inhibitors and colon

inflammation, however, as noted above, inflammation at 3 days post infection did not differ

across diets in this model. Taken together, these data support that bile acid pools were strongly

influenced by diet, with the WD having the most pro-*C. difficile* bile acid composition, but more

work needs to be done to determine the degree to which these differences were driving the higher

mortality observed with the WD.

Figure 3: *Bile acid pools in cecal contents of infected mice 3 days post C. difficile infection.* **(A)**

Cecal levels of *C. difficile* inhibitors (*CDCA, UDCA, a_MCA, b_MCA, LCA, DCA*), *C. difficile*

promotors (*TCA, CA*), and ratios of promoters:inhibitors across diets (chow n = 20, LF n = 20,

WD n = 25; Table S2). Significant differences calculated with a Kruskal-Wallis and Dunn's post

hoc test. Kruskal-Wallis p-values were corrected for multiple comparisons with the FDR

algorithm of Benjamini and Hochberg. Median and IQR indicated. **(B)** Linear regressions of

cecal or colonic histology against CD inhibitors, CD promotors, and the promotor:inhibitor

ratios. Only colon inflammation versus *C. difficile* inhibitors was significant.

Cecal levels of SCFAs and their relationship diet, secondary bile acids, cecal levels of C.

difficile toxins, and inflammation

To further explore mechanism, we also used targeted GC/MS to measure the levels of the

SCFAs butyrate, propionate, and acetate in the aspirated cecal contents in mice that were

sacrificed at 3 days post infection (Table S2). SCFAs are of interest because they are microbial

products of fermentation of dietary fiber, have been previously implicated in the positive effects

of a diet rich in MACs, [15] and are decreased in individuals with rCDI [16]. Although butyrate can

directly suppress *C. difficile* growth *in vitro* [15], butyrate also enhances *C. difficile* toxin

production *in vitro* [15 33]. We also directly evaluated the secondary bile acid DCA, since it can

arrest the growth of vegetative *C. difficile* [13] and prior studies have shown that secondary bile

acid producers such as *Clostridium scindens* can protect against CDI in mice[14].

Butyrate, acetate, and DCA were all significantly higher in the chow diet compared to

both the LF/LF diet and WD (Figure 4A). There was also a significant correlation between levels

of DCA and butyrate in a multivariate regression that accounted for differences across diets

(Figure 4B). This is consistent with both DCA and butyrate having been linked with the presence

of a healthy protective gut microbiome composition and low levels of both have been observed

in individuals with rCDI[30,34]. Surprisingly, butyrate positively correlated with TcdB (Figure 4C)

and cecal and colonic inflammation (Figure 4D), but linear regression indicated that this

relationship was dependent on diet, being driven by a positive association in the WD and LF/LF

diet contexts only (Figure 4C, Table S3). DCA also correlated with tcdB levels and cecal and

colonic inflammation in a diet dependent manner, with a positive relationship in LF/LF and WD

and the expected negative (protective) relationship only in chow(Figure 4D, Table S3).

Figure 4: *Relationships between microbial metabolites (SCFAs and the secondary bile acid*

DCA) and diet, toxin, and inflammation. **(A)** Cecal levels of the SCFAs acetate, butyrate, and

propionate, and the secondary bile acid DCA across diets. p-values were determined using an

Kruskal-Wallis with Dunn's post hoc test. Median and IQR indicated. **(B)** Multiple linear

regression of DCA levels as a function of butyrate and diet. Model = ln(DCA) ~ ln(butyrate) +

diet + ln(butyrate)*diet. R-squared 0.855 p <0.0001. **(C)** Multiple linear regressions of *C.*

difficile toxin tcdA and tcdB concentrations against butyrate and DCA while controlling for

dietary interactions. **(D)** Multiple linear regressions of cecal or colonic histology against

butyrate and DCA while controlling for dietary interactions.

 A conventional chow diet increases homogeneity of response, resilience and alpha-diversity of

the gut microbiome after challenge with antibiotics and CDI compared to both purified diets

We next sought to understand how the composition of the fecal microbiome was affected

by diet during the course of antibiotic treatment and infection with *C. difficile* (Fig. 1A). Fecal

pellets were collected during experiment 1 upon arrival prior to diet change (Day -7), just prior

to the start of oral antibiotic delivery (Day 0), after 5 days of oral antibiotics (Day 5), and daily

through Day 10, which captured before and after the clindamycin injection given on day 7 and *C.*

difficile gavage on Day 8 (Fig. 1A). Collected samples were subjected to 16S ribosomal RNA

(rRNA) amplicon sequencing targeting the V4 region of rRNA on the MiSeq platform.

Principle coordinate analysis (PCoA) plots of a weighted UniFrac [35] distance matrix

suggested that mice fed either the WD or LF/LF diet had decreased resilience and a less

homogeneous response to antibiotic challenge and CDI as compared to chow-fed mice (Figure

5). Mice fed either the WD or LF/LF diet showed greater divergence across PC1 upon antibiotic

exposure than chow-fed mice, higher spread across mice in the same diet group, and less

recovery towards their baseline after antibiotics (Fig. 5A). We quantified resilience by

comparing the pairwise weighted UniFrac distances of mice across the experiment to baseline

microbiota of their respective diet cohort at Day 0 (7 days post-diet change and pre-oral

antibiotics; Fig. 5B). Chow-fed mice had significantly smaller weighted UniFrac distances from

their baselines than the other groups at Day 5 (post 5 days antibiotic challenge) that persisted

through Day 10 despite some convergence after clindamycin injection (Day 8) (Fig. 5B). By Day

9, chow-fed mice again displayed higher microbiome resilience than both the WD and LF/LF

diet groups. We also assessed the homogeneity of response to a disturbance among mice in the

same diet group. As an example, low homogeneity would occur if the mice within a diet group

showed high variability in the degree to which their gut microbiome changed upon antibiotic

exposure. We quantified this as the median pairwise weighted UniFrac distance for comparisons

within samples collected at the same time point from mice fed the same diet (Fig. 5C). Both the

WD and LF/LF diet showed much lower homogeneity of gut microbiome compositional

response to antibiotic challenge, particularly to the 5-day treatment with oral antibiotics (Day 5),

compared to chow-fed mice (Fig. 5C).

Similar patterns were seen when evaluating changes in alpha-diversity across the

experiment between each diet cohort. Figure 6 shows changes in phylogenetic entropy, which is

a measure of alpha diversity that considers species richness, evenness, and distinctness[36]. The

phylogenetic entropy of the WD-fed mice was lower than chow-fed mice after diet change and

this difference became more pronounced upon oral antibiotics and remained so through the rest

of the experimental timeline (Fig. 6). Interestingly, the phylogenetic entropy of the LF/LF diet-

fed mice remained equivalent to the chow-fed cohort with diet change but decreased to the same

level as the WD with antibiotic treatment (Fig. 6).

Figure 5: Beta diversity plots of fecal microbiome by diet during antibiotic treatment and

infection with *C. difficile*. Vertical red lines in panels B and C designate the day of *C. difficile*

infection (chow n = 13, LF n = 5, WD n = 13). (A) Weighted UniFrac PCoA plots of all samples

with each diet highlighted in separate panels. (B) Resilience of microbiome composition

assessed by within-mouse pairwise weighted UniFrac distances between Day 0 (7 days post diet

switch and prior to oral antibiotics) and later time points and (C) Longitudinal plot of

microbiome turnover homogeneity as plotted by intra-time point pairwise Weighted UniFrac

distances within diet groups. Significant differences between diet groups were calculated by

Kruskal Wallis followed by Dunn's post hoc test. Trend lines were fit using local polynomial

regression. ***: p<0.001. **: p<0.01, *: p<0.05 ns= non-significant.

Figure 6: Alpha-diversity (phylogenetic entropy) of the fecal microbiome during murine CDI

model. Data for each individual mouse is plotted as well as the fitted local polynomial regression

for each diet group. Significant differences between diet groups were calculated by Kruskal

Wallis followed by Dunn's post hoc test.. ***: p<0.001. **: p<0.01, *: p<0.05 ns= non-

significant.

The WD and LF/LF diets had increased facultative anaerobe colonization and decreased

secondary bile acid and SCFA-producing bacteria compared to conventional chow diet

Low-diversity dysbiosis is a state of disturbance that is often characterized not only by

low alpha-diversity, but also by an increased ratio of facultative to strict anaerobes [37]. Low-

diversity dysbiosis is associated with a number of diseases including rCDI [37]. We sought to

investigate whether the different diets tested influenced if the microbiome developed a

compositional state characterized by high levels of facultative anaerobe colonization and lower

levels of strict anaerobes. Since Lactobacilliales and Enterobacteriales contain many important

intestinal facultative anaerobes and most members of Clostridiales are strict anaerobes and

include key butyrate and secondary bile acid producers, we plotted the relative abundances of

these orders over the course of the experiment (Fig. 7A). All mice had decreases in the relative

abundance of Clostridiales in their fecal microbiome with oral antibiotics; however, mice fed a

chow diet were able to maintain a Clostridiales population while both the WD and LF/LF diets

saw near-complete elimination of these taxa (chow-WD p<0.01 for days 0 through 9 and p<0.05

on day 10, Fig. S3). Conversely, mice fed either the WD or LF/LF diet had a large bloom of

Lactobacillales after oral antibiotic treatment that was not observed in the chow-fed mice (chow-

WD p <0.001 and chow-low-fat/low-fiber p<0.05). Lastly, all 3 diet groups had a large increase

in Enterobacteriales in their fecal microbiome following antibiotics; however, the LF/LF and

chow groups showed earlier decrease than WD mice (chow-WD p <0.01 and p<0.05 at days 9

and 10 respectively, Fig. S3). Comparisons of the LF/LF diet were limited due to smaller sample

size (n = 5 vs. n = 13 for chow and WD).

We also used PICRUSt[38] to predict metagenomes using our 16S rRNA data to 1)

investigate trends in the prevalence of key genes in secondary bile and butyrate production over

the course of our experimental timeline and 2) predict which bacterial taxa were contributing

these genes. Because *baiA*, *baiB*, and *baiCD* are not available in PICRUSt2's set of predicted

genes, we only used the genes for *baiH* (KEGG ID: K15873) and *baiI* (KEGG ID; K15874),

which are both genes in the *bai* operon[39], to assess genomic potential for secondary bile acid

metabolism. Acetoacetate co-A transferase (*but*; K01034) and Butyrate Kinase (*buk*; KEGG ID:

K00929), which are the main pathways for fermentative production of butyrate in the gut

microbiome[40], were used to assess butyrate production potential. Plotting these genes/pathways

over time reveals a significant effect of diet on their abundance and response to antibiotics (Fig.

7C). Although all diet groups showed a marked decrease in bile acid genes with oral antibiotics,

only the chow-fed mice displayed a recovery of secondary bile acid genes, though the source of

these genes switched from Lachnospiraceae UCG-006 to Blautia. This result is consistent with

our observation of higher cecal levels of secondary bile acids in chow-fed mice compared to

mice fed either the WD or LF/LF diets at 3 days post *C. difficile* gavage (Fig. 4A).

319   Butyrate coding capacity also differed between diet groups. Chow-fed mice showed

minimal change in the abundance of both the *but* and *buk* genes for fermentative butyrate

production during the time course while the WD mice had a decrease of 5 orders of magnitude

(Fig. 7D). The LF/LF diet-fed mice showed an intermediate phenotype with the resilience of the

butyrate pathway being mostly attributed to a butyrate kinase dependent pathway. The results for

but and not *buk* however are consistent with our measurements of cecal butyrate levels in these

mice 3 days post *C. difficile* gavage (Fig. 4A). This is consistent with *but* being regarded to be a

more important source of butyrate in the intestine [41].

327   Since we had observed a strong positive correlation between cecal levels of butyrate and

328   the secondary bile acid DCA in our mass spectrometry data (Fig. 4B), we also determined

329   whether there was a relationship between butyrate and secondary bile acid coding capacity. We

330   found a highly significant association (p = 3.6×10^{-5}), with secondary bile acid producing genes

331   only predicted to be present in samples that also had high predicted levels of butyrate producing

332   genes (Fig. 7B).

Figure 7: Changes in key taxa, and secondary bile acid and butyrate coding capacity during the
CDI protocol. The vertical red line in (A), (C) and (D) indicates the day of *C. difficile* infection.
All trend lines were fit using local polynomial regression. **(A)** Relative abundance of key
bacterial orders during antibiotic treatment and infection. A summary of significant differences
of these taxa across diets are in Fig. S3. **(B)** Violin plots of abundance of butyrate genes from
PICRUSt analysis binned by presence of secondary bile acid producing genes (Wilcoxon
p<0.001). **(C)** Time course of coding capacity of secondary bile acid genes. The top row shows
the total capacity of each sample (*baiH* and *baiI*) while the bottom two rows show specific taxa
contributions of key genes in the Bai operon. **(D)** Time course of coding capacity of butyrate
producing genes by diet. The top row shows the total capacity as measured by *but* and *buk* genes
while the bottom two rows show specific taxa contributions of *but* and *buk* specifically. Taxa
with mean relative abundance < 0.01% were filtered from the analysis.

Discussion

C. difficile infection is a grave and growing health threat. Current strategies to limit its

spread have focused on sanitation and antibiotic stewardship, however incidence has continued

to rise despite these efforts, highlighting the need for new treatment and prevention strategies[2].

Because of the ubiquity of *C. difficile* spores in the environment and high levels of colonization,

focusing on ways to increase the resilience of the host to *C. difficile* disease is one important

prevention strategy [42].

Our results augment a growing body of evidence from studies conducted in mouse

models of antibiotic induced *C. difficile* infection that points to dietary intervention as a

promising approach to prevent antibiotic-induced CDI[4,15,18,19]. Prior studies have suggested the

importance of a variety of macronutrients, including MACs, protein, and fat. Specifically, for

MACs, one study that used an antibiotic-induced murine model demonstrated that mice fed a diet

deficient in MACs (e.g. soluble fiber, resistant starches) had persistent *C. difficile* shedding and

that there was a resolution of colonization with the reintroduction of inulin or other MACs[15]. A

protective effect of dietary MACS was also demonstrated in a human clinical trial in which a

decrease in *C. difficile* recurrence from 34.3% to 8.3% was observed with prebiotic

supplementation[43]. Our results are consistent with these studies in that both the WD and LF/LF

diets were low in MACs compared to the chow diet and had a greater antibiotic-induced

disturbance to the gut microbiome and loss of CDI protective microbial metabolites such as

DCA. However, it is important to note that the differences between conventional chow, WD and

LF/LF diets extend well beyond fiber, and these other dietary components could also have

influenced our observation. Also, our LF/LF diet had greatly reduced mortality compared to the

WD, even though the LF/LF diet-fed mice was low in MACs and had a comparable level of

microbiome disturbance and loss of protective metabolites with antibiotic treatment.

Influence of dietary protein has also been noted in a few studies. Specifically, one study

found a low-protein diet to be protective in an antibiotic-induced CDI murine model, with mice

fed a 2% protein diet having increased survival, decreased weight loss, and decreased overall

disease severity compared to mice fed a 20% protein defined diet [20]. Another study showed that a

diet poor in proline (an essential amino acid for *C. difficile* growth) prevented *C. difficile*

carriage[4]. Furthermore, in a recent study that evaluated both a high-fat/high-protein Atkins-type

diet and a high-fat/low-protein diet in a mouse model of antibiotic-induced CDI, the high-

fat/high-protein diet promoted severe CDI and 100% mortality, while the high-fat/low-protein

diet had variable disease severity and survival, showing a strong effect of dietary protein but

indicating that the effects of fats were uncertain [18].

Another had found that a diet that was high in refined carbohydrates and low in fiber had

improved CDI severity compared to mice fed a standard chow diet [18]. New data has suggested

that novel speciation of *C. difficile* may be selecting for strains that show increased sporulation

and host colonization capacity with sugar availability (glucose or fructose) [47.] This work,

conducted with *C. difficile* strain (VPI 10463), did not show differences in mortality from CDI in

low-fat/low-fiber diets with different amounts of sucrose[18,19].

Our results show that high dietary fat in the context of low dietary fiber had a strong

effect on CDI-induced mortality, with mechanisms distinct from a loss of beneficial microbial

metabolites. Evidence to suggest that a high-fat/low-fiber western-type diet could have a

profound effect on CDI was first presented over 20 years ago in experiments designed to study

the atherogenic properties of a Western diet in Syrian hamsters[44,45]. Significant mortality from

CDI was observed in hamsters fed a high-fat/low-fiber pro-atherogenic diet and not a typical

high-fiber/low-fat hamster diet, even in the absence of an antibiotic disturbance [44,45]. Another

recent study that conducted a study of antibiotic-induced CDI in a high-fat-diet (HFD) induced

obesity model found protracted disease in the HFD compared to a chow diet[19], but not the severe

mortality that we observed with a high-fat/low-fiber diet. We posit that high dietary fat may have

a more profound influence on CDI than low dietary fiber since a prior study of MAC deficient

diets found that low fiber was associated with higher *C. difficile* carriage but did not describe the

severe disease/mortality that was observed here while using a similar mouse model[15]. However,

since we did not test a high-fat/high-fiber diet, it is unclear whether the high mortality that we

observed was due to a combination of high-fat and low-fiber in the diet, or just dietary fat.

Although these studies taken together support a potential synergy of high-fat and low-

fiber leading to severe disease, it is important to note that these papers differ in many

experimental parameters including the source of the mice (which has been shown to influence

response to antibiotic perturbation and *C. difficile* clearance in mice[46]), types of antibiotics used,

strain of *C. difficile*, and whether *C. difficile* was used as active growing bacteria (as done in our

study) or as spores.

The Role of Toxin Production and Inflammation

In order to further explore potential causes of death, we looked at both inflammation by

histology and levels of the toxins TcdA and TcdB by ELISA in cecal contents collected 3 days

post *C. difficile* infection, which was just prior to the onset of mortality in our longitudinal

cohort. Both TcdA and TcdB can disrupt cytoskeletal structure and tight junctions of target cells

[48] and induce inflammation [49, 3]. We did not observe any differences in TcdB or cecal or colon

inflammation scores across diets. However, cecal levels of TcdB did correlate with cecal

inflammation, consistent with known effects of TcdB [48] [50]. This supports that levels of TcdB

produced by *C. difficile* may indeed be causing pathology in these mice, but higher levels of

TcdB at Day 3 post CDI cannot alone explain the higher mortality that we began to observe at

Day 4 post CDI infection in the WD-fed mice. Interestingly, TcdA and TcdB levels did not

correlate with each other, and TcdA levels did show a pattern at Day 3 post-CDI consistent with

mortality, being significantly higher in the WD mice compared to the LF/LF-fed mice.

Consistent with the LF/LF diet mice having delayed mortality compared to chow-fed mice, there

were also lower levels of TcdA in the LF/LF fed mice compared to those fed a conventional

chow diet at Day 3 post-CDI. However, unlike TcdB, TcdA levels did not correlate with

inflammation. The lack of correlation of TcdA and TcdB with each other is surprising since they

are often co-expressed, although their transcription is regulated by unique promoter regions[51],

and other post-transcriptional factors at the RNA or protein degradation level may also be at

play. Studies of the activity of TcdA versus TcdB in various animal models have more strongly

supported the importance of TcdB in CDI pathogenesis, and studies investigating TcdA have had

mixed results[48], although none of these studies were conducted in the context of a high-fat/low-

fiber diet. Our results support a potential importance of TcdA and not TcdB in diet-associated

differences in CDI pathogenesis, but further studies that sample the toxin levels at more time

points over disease progression might prove illuminating. Indeed, another prior study that

showed higher CDI pathology in HFD-induced obesity model versus a regular chow diet did not

observe higher toxin levels (while binning TcdA and TcdB ELISA data) at day 3 post infection

(acute phase), but did find higher toxin levels and intestinal inflammation between diets at day

10 post infection, due to recovery occurring in the chow fed but not HFD-obese mice [19].

Although it is possible that differences in TcdB and inflammation across diets in our study may

have emerged over time, it was not possible to evaluate this since we had much higher mortality

in our model, and most of our WD fed mice would have died by day 10. Further studies that use

complementary methods to measure toxin besides just ELISA, which can lack specificity for

TcdB in particular[52,53], or with strains of *C. difficile* that produce TcdA or TcdB only would be

required for further validation [52]. Also, in these studies we measured toxin levels but were unable

to produce quality data regarding levels of *C. difficile* bacteria in the cecal materials. We thus

cannot evaluate whether these differences in toxin levels are driven by more bacteria or increased

toxin production by similar loads of bacteria.

Effects of bile acids

Differences in host bile acid production and microbial bile acid metabolism is one

potential mechanism of high-fat diet induced modulation of CDI severity. In the clinical setting,

studies have shown that patients with CDI have increased TCA and decreased concentrations of

the secondary bile acids DCA and LCA in their feces [30,34] as well as other complex alterations to

bile acid pools[31]. These derangements are corrected with fecal microbiota transplant for

treatment of *C. difficile* (FMT) [34]. *In vitro* experiments have shown that the primary bile acids

TCA and CA are potent *C. difficile* germination factors [13] while UDCA and CDCA have been

shown to inhibit germination and growth of *C. difficile in vitro* [27] [28] [29]. The microbially-produced

secondary bile acids DCA and LCA have also been shown to affect *C. difficile in vitro*: DCA

promotes germination of *C. difficile* spores [13] while LCA inhibits germination [29] and both inhibit

growth of vegetative *C. difficile* [13] [29]. In line with these effects, reduced prevalence of the

secondary bile acid producer, *Clostridium scindens* in the fecal microbiome has been associated

with high incidence of CDI in both humans and in experimental mouse models, and gavaging

mice with *C. scindens* protected against CDI and restored intestinal secondary bile acid levels [14].

Despite this strong evidence of a role of a protective effect of microbially produced secondary

bile metabolites in protection from CDI, this mechanism did not appear to be a sole driving

factor of the mortality that we observed in mice fed a WD, since the levels of these metabolites

were lowest in the mice fed the LF/LF diet even though the LF/LF mice did not experience

increased mortality. Levels of *C. difficile* inhibitors, which included the secondary bile acids

DCA and LCA, did negatively correlate with colonic inflammation, suggesting some degree of

protection in these mice. Functional interrogation of the microbiome using PICRUSt suggests

that the lack of secondary bile acids in the WD and LF/LF diet fed mice might be due to a lack of

recovery of secondary bile acid producing bacteria following antibiotic disturbance in both the

WD and LF/LF diet contexts.

We did find that the ratio of *C. difficile* promoters:inhibitors was significantly higher in

the WD compared to both the LF/LF and chow diets, consistent with mortality differences. Our

results support that a high-fat diet coupled with low-fiber and antibiotic treatment may provide a

"double hit" for shifting towards a pro-*C. difficile* bile acid pool – with dietary fat increasing

excretion of pro- *C. difficile* primary bile acids into the gut and antibiotic-induced gut

microbiome disturbance decreasing their conversion into protective secondary bile acids. *In vitro*

assays have demonstrated that variable mixtures of primary and secondary bile assays have

different impacts on *C. difficile* germination and growth[54]. However, more work needs to be

done to determine the degree to which these differences were driving the higher mortality

observed with the WD. A more convincing result would be if the *C. difficile* promoter:inhibitor

ratio also predicted *C. difficile* toxin production while controlling for diet, but this was not the

case (Fig. 3B).

We also found that diet had a significant effect on 4 of the 5 taurine conjugated bile acids

that we assayed, with TCA, T_b_MCA, TDCA, and TCDCA all showing a pattern of increased

levels in the WD compared to both the chow and LF/LF diets, but only comparisons of chow

versus WD reaching statistical significance (Fig. S2). It is probable that the further decrease in

levels of these taurine conjugated bile acids in the chow compared to the LF/LF diet is because

the primary bile acids that are produced by the host are converted by microbes to secondary bile

acids in only the chow diet. Our finding of increased TCA in the WD compared to chow is

consistent with a prior study that found that IL10-deficient mice fed a diet high in saturated fat,

had an increased proportion of taurine-conjugated bile acids compared to standard chow and a

diet high in poly-unsaturated fats [21]. One prior study demonstrated that both TDCA and TCDCA

have pro-germinative effects on *C. difficile*, though in our study, their cecal concentrations were

orders of magnitude lower than TCA which is also a much stronger germinant[55].

One weakness of our study is that we cannot differentiate between the complex changes

of the bile acid pools and the effects of the dietary components themselves – such as known

effects of high-fat diet on inflammation [56,57]. Controlled studies that directly alter bile acid pools

without also altering diet are valuable. In one study of CDI in HFD-induced obesity, inhibiting

primary bile acid synthesis with the FXR antagonist obeticholic acid ameliorated CDI disease

during later phases of infection but not in acute CDI [19]. Another factor that may have influenced

our result is that like other related studies [15], our infection procedure used a sample cultured for

~24 hours without enumerating or enriching the sporulated fraction of the inoculum. As spores

are the likely infective form of *C. difficile* in clinical settings, and many bile acids influence *C.*

difficile pathogenesis by promoting or inhibiting germination, it is of interest to determine how

the variability in vegetative composition influences the relationship between bile acid pools and

CDI pathogenesis.

Effects of the microbiome and their metabolites

Our data suggests that a complex diet is critical for the resilience and homogeneity of

response of the gut microbiome after perturbation. In both cohorts of mice fed a purified diet that

was deficient in fiber, the gut microbiome was significantly more variable and slower to recover

to baseline after perturbation. We hypothesize that by supplying the gut with a preferred fuel

(fiber) for species associated with health (e.g. strict anaerobes), the community is able to resist

antibiotic induced changes and reconstitute more quickly once the pressure of antibiotic

treatment has been removed. Since the chow diet differed from the purified diets in many

components besides the levels of fiber, we cannot conclude from our study alone that increased

resilience to microbiome disturbance with antibiotics in chow is driven by differences in fiber.

However, our results are consistent with previous murine studies that have shown that low fiber

diets can increase antibiotic-induced microbiome disturbance and delay recovery from treatment

with ciprofloxacin [58] and that fiber supplementation can lead to a reduced disruption of the gut

microbiome to disturbance from amoxicillin [59].

The increased resilience of gut microbiome composition to antibiotic disturbance was

also reflected through levels of the bacterially produced metabolites that we measured. Neither

the WD or LF/LF diets were able to maintain butyrate or secondary bile acid production

following antibiotic perturbation. Based on the correlation between butyrate and DCA

concentrations, we speculate that the lack of butyrate leads to increased luminal oxygen

concentrations that are unsuitable for *Clostridium scindens* and other secondary bile acid

producers. Prior work has shown that aerobic metabolism of butyrate by intestinal epithelial cells

is a key driver of intestinal hypoxia [60]. That there may be increased luminal oxygen

concentrations in the LF/LF and WD is consistent with our observation of a bloom in

Lactobacillales order, which is entirely composed of facultative anaerobes, after oral antibiotic

challenge in the WD and LF/LF diets but not chow.

While our data do not suggest a role for fiber in protection against mortality from CDI in

this mouse model since the LF/LF diet fed mice were protected without fiber in the diet, it would

be short-sighted to dismiss the beneficial role of fiber in maintaining a healthy gut microbiome

and resistance to CDI. Our model utilized a rather short-term diet change and an intense

antibiotic regimen. We also did not explore diets high in fat and high in fiber, where it is possible

that increased microbiome resilience to antibiotics due to fiber may protect from the detrimental

effects of fat. As discussed above, a fiber-deficient diet has been shown to hinder clearance of *C.*

difficile after challenge [15].

One surprising finding of our work, however, given these protective effects of dietary

Microbiota Accessible Carbohydrates (MACs) from other studies, was that butyrate, a major

fermentation product of MACs, positively correlated with TcdB and cecal and colonic

inflammation, driven by an association in the WD and LF/LF diet contexts and not chow. DCA

also correlated with butyrate and with TcdB levels and cecal and colonic inflammation in a diet

dependent manner, with a positive relationship in LF/LF and WD and the expected negative

(protective) relationship only in chow. Butyrate is typically associated with beneficial effects on

gut health, including supporting intestinal barrier function [61 62 63], suppressing inflammation

through induction of T regulatory cells [64], and directly suppressing *C. difficile* growth *in vitro* in

a dose dependent manner[15]. Also, lower butyrate and DCA has been observed clinically in

individuals with CDI [30 34]. However, a positive correlation between butyrate and TcdB is

consistent with prior studies showing that butyrate enhances *C. difficile* toxin production *in vitro*

[15,33], leading some to suggest that butyrate may signal to *C. difficile* a competitive gut

environment [15]. A similar diet-dependent detriment of SCFAs was observed in a study that

showed that soluble fiber-supplementation drove hepatocellular carcinoma in mice in a manner

dependent on microbial fermentation to SCFAs, but this effect occurred when soluble fiber was

added to a compositionally defined diet and not to a conventional chow diet [65]. Our results

suggest that supplementation with soluble fibers such as inulin to prevent *C. difficile* may not

produce the desired result in individuals who are otherwise consuming highly refined diets.

Limitations of our study:

We have demonstrated a striking difference in diet-mediated mortality in an antibiotic-

induced murine CDI model, but our study does have limitations. We did not explore how the

composition of fat influences these factors. Our WD composition represents a typical diet in the

United States based on population survey data. Further studies to determine if total fat intake or

specific types of fat drive our observed phenotype are needed. Furthermore, we only evaluated

the effects of fat in a low-fiber context. Evaluating a high-fat/high-fiber diet would elucidate

whether the expected beneficial effects of fiber on the microbiome would temper the negative

effects of high-fat. Comparisons between chow-fed mice and those receiving a purified diet are

limited due to the marked differences in the composition of macronutrients [66]. Since this is an

antibiotic-induced CDI model, our results only reflect effects of diet in the context of antibiotic

disturbance. Finally, we note that we induced CDI infection using a standardized amount of live

C. difficile, which is a commonly used in murine studies of *C. difficile* [22]. However, we note that

since different bile acids influence spore germination as well as growth, results may vary in

challenge models that instead use spores. Future experiments to compare these results to a spore-

infection model would elucidate the effects of germination versus growth on diet effects on CDI

pathogenicity.

CONCLUSIONS

This study along with recently published findings investigating dietary fiber [15], dietary

proline [4], protein [18], and fat [18] [19] intake provides a compelling case that diet should be

increasingly targeted as a prevention and treatment modality for CDI. High-risk populations such

as elderly hospitalized individuals subjected to antibiotics and adult and pediatric oncology

patients may benefit from decreased *C. difficile* colonization through diets with decreased fat and

increased fiber. For patients with active infection, limiting fat intake could decrease disease

severity while maintaining enteric nutrition.

Methods

Mouse diets: Diets were all obtained from Envigo (Indiana): Standard chow - Teklad

global soy protein-free extruded (item 2920X - https://www.envigo.com/resources/data-

sheets/2020x-datasheet-0915.pdf), Western Diet – New Total Western Diet (item TD.110919),

Low-fat/low-fiber – variant of AIN93G (item TD.180811). See Table S4 for detailed

composition of purified diets.

Murine model of CDI: Mice were infected using a widely used murine CDI model [22]

with minor modifications. Briefly, 6-week-old female C57BL/6 mice from Taconic Bioscience

(Rensselaer, NY) arrived at University of Colorado on Day -7 of the experiment. During

experiments, mice were cohoused in groups of 4-5 mice per cage. Survival experiments were

conducted in four independent experiments at four separate starting dates. For cecal metabolite

and toxin analysis, 2-7 independent experiments were conducted with separate starting dates

(Table S2). Within 24 hours, mouse feed was changed to one of three diets: standard chow, high-

fat/low-fiber (WD), or LF/LF diet (all groups n=20 over 4 batches; Table S2). After seven days

of the new diet, we placed mice on a five-antibiotic cocktail (kanamycin (0.4 mg/ml), gentamicin

(0.035 mg/ml), colistin (850 U/ml), metronidazole (0.215 mg/ml), and vancomycin (0.045

mg/ml)) in their drinking water. Antibiotics were removed for 48 hours, after which we

administered an intraperitoneal injection of clindamycin in normal saline (10 mg/kg body

weight). Twenty-four hours after injection, we gavaged mice with 1.75×10^5 cfu of *C. difficile*

VPI 10463. We weighed mice daily after removal of oral antibiotics and they were euthanized if

they lost >15% of body weight or were moribund. Fecal pellets were collected at arrival (Day -

7), after diet change and prior to oral antibiotics (Day 0) and then daily after removal of oral

antibiotics (Day 5-10). In a separate set of experiments, we performed the same experimental

protocol on 66 mice (chow = 20, low-fat/low-fiber = 20, WD = 26) over 7 different batches (see

Table S2), but we sacrificed the mice 72 hours after infection and collected cecal contents for

SCFA, bile acid and toxin quantification and cecum and intestines for histopathology. Mice for

the second experiments were also obtained from Taconic Bioscience. All mouse experiments

were approved by the Institutional Animal Care and Use Committee and complied with their

guidelines and NIH Guide for the Care and Use of Laboratory Animals (IACUC protocol

#00249).

DNA Extraction and Sequencing: Total genomic DNA was extracted from fecal pellets

from a subset of the mice in cohort 1 (chow = 13 mice from four separate cages over two

experiments, WD = 13 mice from four separate cages over two experiments, LF/LF = 5 mice

from two cages over two experiments) using the DNeasy PowerSoil Kit (Qiagen, Germantown,

MD). Modifications to the standard protocol included a 10-minute incubation at 65°C

immediately following the addition of the lysis buffer and the use of a bead mill homogenizer at

4.5 m/s for 1 min. The V4 variable region of the 16S rDNA gene was targeted for sequencing

(515F: GTGCCAGCMGCCGCGGTAA, 806R: GGACTACHVGGGTWTCTAAT). The target

DNA was amplified using HotMaster Mix (Quantabio Beverly, MA). Construction of primers

and amplification procedures follow the Earth Microbiome Project guidelines

(www.earthmicrobiome.org) [67]. Amplified DNA was quantified in a PicoGreen (ThermoFisher

Scientific) assay and equal quantities of DNA from each sample was pooled. The pooled DNA

was sequenced using a V2 2x250 kit on the Illumina MiSeq platform (San Diego, CA) at the

University of Colorado Anschutz Medical Campus Genomics and Microarray Core facility.

Sequence Data Analysis: Raw paired-end FASTQ files were processed with QIIME 2

version 2018.8 [68]. Denoising was performed with DADA2 [69], a phylogenetic tree was built using

sepp [70] and taxonomy was assigned to amplicon sequence variants (ASVs) using the RDP

Classifier [71] trained on the Silva version 132 taxonomic database [72,73] using QIIME 2 [68]. The data

was rarefied at 5,746 sequences per sample. Alpha-diversity was measured by phylogenetic

entropy [36] and beta-diversity was determined by weighted UniFrac distances [35]. PCoA of

weighted UniFrac plots were constructed using QIIME 2. Metagenomes were imputed from 16S

ASVs using PICRUSt2's default pipeline for stratified genome contributions [38]. Low abundance

taxa (<0.01% mean relative abundance) were filtered for analysis of the butyrogenic coding

capacity. Software was installed using Anaconda [74] and analysis was performed on the Fiji

compute cluster at the University of Colorado Boulder BioFrontiers Institute.

SCFA quantification: The SCFAs butyrate, propionate, and acetate were analyzed by

stable isotope GC/MS as previously described [75]. Briefly, cecal samples were collected directly

into pre-weighed, sterile Eppendorf tubes and flash frozen at −80°C until processing. Samples

were then subject to an alkylation procedure in which sample and alkylating reagent were added,

vortexed for 1 min, and incubated at 60°C for 25 min. Following cooling and addition of n-

hexane to allow for separation, 170 μL of the organic phase was transferred to an auto sampler

vial and analyzed by GC/MS. Results were quantified in reference to the stable isotope standard

and normalized to sample weight.

Bile acids quantification: *Reagents:* LC/MS grade methanol, acetonitrile, and

isopropanol were obtained from Fisher Scientific (Fairlawn, New Jersey). HPLC grade water

was obtained from Burdick and Jackson (Morristown, New Jersey). Acetic acid, cholic acid

(CA), chenodeoxycholic acid (CDCA), lithocholic acid (LCA), taurocholic acid (TCA) and

deoxycholic acid (DCA) were obtained from Sigma Aldrich (St. Louis, Missouri).

Taurodeoxycholic acid (TCDCA), taurochenodeoxycholic acid (TCDCA), taurolithocholic acid

(TLCA), alpha-muricholic acid (a_MCA) and beta-muricholic acid (b_MCA) were obtained

from Cayman Chemical (Ann Arbor, Michigan). Chenodeoxycholic acid-d4 (CDCA) and

glycochenodeoxycholic acid-d4 were obtained from Cambridge Isotope labs (Tewksberry,

Massachusetts).

Standards preparation: An internal standard containing 21 μM of chenodeoxycholic

acid-d4 and 21 μM of glycochenodeoxycholic acid–d4 was prepared in 100% methanol. A

combined stock of all bile acid standards was prepared at 0.5mM in 100% methanol. Calibration

working standards were then prepared by diluting the combined stock over a range of 0.05 μM-

50 μM in methanol. A 20 μL aliquot of each calibration working standard was added to 120 μL

of methanol, 50 μL of water and 10 μL of internal standard (200 μL total) to create 10

calibration standards across a calibration range of 0.005 μM-5 μM.

Sample preparation: Fecal samples were prepared using the method described by

Sarafian et al[76] with modifications. Briefly, 15-30mg of fecal sample were weighed in a tared

microcentrifuge tube and the weight was recorded. 140 µL of methanol, 15-30 µL of water and

10 µL of internal standard were added. The sample was vortexed for 5 seconds, and then

incubated in a -20°C freezer for 20 minutes. The sample was then centrifuged at 6000RPM for

15 minutes at 4°C. 185-200 µL of the supernatant was transferred to an RSA autosampler vial

(Microsolv Technology Corporation, Leland, NC) for immediate analysis or frozen at -70°C

until analysis.

High performance liquid chromatography/quadrupole time-of-flight mass spectrometry

(HPLC/QTOF): HPLC/QTOF mass spectrometry was performed using the method described by

Sarafian et al [76] with modifications. Separation of bile acids was performed on a 1290 series

HPLC from Agilent (Santa Clara, CA) using an Agilent SB-C18 2.1X100mm 1.8 µm column

with a 2.1X 5mm 1.8um guard column. Buffer A consisted of 90:10 water:acetonitrile with 1mM

ammonium acetate adjusted to pH=4 with acetic acid, and buffer B consisted of 50:50

acetonitrile:isopropanol. 10 µL of the extracted sample was analyzed using the following

gradient at a flow rate of 0.6mls/min: Starting composition=10%B, linear gradient from 10-35%

B from 0.1-9.25 minutes, 35-85% B from 9.25-11.5 minutes at 0.65mls/min, 85-100% B from

11.5-11.8 minutes at 0.8mls/min, hold at 100% B from 11.8-12.4 minutes at 1.0ml/min, 100-55%

B from 12.4-12.5 minutes 0.85mls/min, followed by re-equilibration at 10%B from 12.5-15

minutes. The column temperature was held at 60°C for the entire gradient.

Mass spectrometric analysis was performed on an Agilent 6520 quadrupole time of flight

mass spectrometer in negative ionization mode. The drying gas was 300°C at a flow rate of

12mls/min. The nebulizer pressure was 30psi. The capillary voltage was 4000V. Fragmentor

voltage was 200V. Spectra were acquired in the mass range of 50-1700m/z with a scan rate of 2

spectra/sec.

Retention time and m/z for each bile acid was determined by injecting authentic

standards individually. All of the bile acids produced a prominent [M-H]$^-$ ion with negative

ionization. The observed retention time and m/z was then used to create a quantitation method.

Calibration curves for each calibrated bile acid were constructed using Masshunter Quantitative

Analysis software (Aligent Technologies). Bile acid results for feces in pmol/mg were then

quantitated using the following calculation:

Concentration in pmol/mg= $\dfrac{(X_s)(V_t)(D)}{(V_i)(W_s)}$

X_s =pmol on column

V_t =Total volume of concentrated extract (in µL)

D   =Dilution factor if sample was extracted before analysis. If no dilution D=1

V_i =Volume of extract injected (in µL)

W_s =Weight of sample extracted in mg

***C. difficile* toxin TcdA and TcdB quantification:** TcdA and TcdB concentrations were

determined in cecal samples from day 3 of infection by comparison to a standard curve using

ELISA (tgcBiomics, Germany). For samples that were too small to weigh accurately, a mass of 5

mg was assigned for concentration calculation. This mass was selected as it was the lowest

weight that could be accurately determined.

Histologic evaluation of large intestinal tissue: Cecum and transverse colon were

harvested from mice three days after infection with *C. difficile* from mice fed either a chow diet

(n=9), LF/LF (n=10) or WD (n=14) total in 3 separate experiments (Table S2). Tissue was fixed

in 10% formalin in PBS, paraffin embedded and sections cut before hematoxylin and eosin

staining by the University of Colorado Histopathology Core. Inflammation was assessed in the

cecum using the Barthel scoring system [24] and in the colon using the Dieleman scoring system [23]

by a trained histologist.

Statistics: Statistical analyses were performed in R (version 3.4.3 "Kite-Eating Tree").

Data were preprocessed using the "tidyverse" suite [77]. We used "survminer" and "survival"

libraries to analyze mouse survival [78,79]. All other data were plotted using "ggplot2", "ggsignif",

and "cowplot" [80,81,82]. All statistical tests were two-tailed with measurements from distinct

samples.

Data availability

The 16S rRNA has been deposited in QIITA [83] (Qiita Study ID: 12849) and at EBI

(pending)

References:

727   1.   Lessa, F. C. et al. Burden of Clostridium difficile infection in the United States. *N Engl J*

728   *Med* **372**, 825-834 (2015).

729   2.   Leffler, D. A. & Lamont, J. T. Clostridium difficile infection. *N Engl J Med* **372**, 1539-

730   1548 (2015).

731   3.   Ng, J. et al. Clostridium difficile toxin-induced inflammation and intestinal injury are

732   mediated by the inflammasome. *Gastroenterology* **139**, 542-52, 552.e1 (2010).

733   4.   Battaglioli, E. J. et al. *Clostridioides difficile* uses amino acids associated with gut

734   microbial dysbiosis in a subset of patients with diarrhea. *Sci Transl Med* **10**, eaam7019

735   (2018).

736   5.   Lozupone, C. A. et al. Meta-analyses of studies of the human microbiota. *Genome Res* **23**,

737   1704-1714 (2013).

738   6.   Bignardi, G. E. Risk factors for Clostridium difficile infection. *J Hosp Infect* **40**, 1-15

739   (1998).

740   7.   Fekety, R. et al. Recurrent Clostridium difficile diarrhea: characteristics of and risk factors

741   for patients enrolled in a prospective, randomized, double-blinded trial. *Clin Infect Dis* **24**,

742   324-333 (1997).

743   8.   Theriot, C. M. et al. Antibiotic-induced shifts in the mouse gut microbiome and

744   metabolome increase susceptibility to Clostridium difficile infection. *Nat Commun* **5**, 3114

745   (2014).

746   9.   Chang, J. Y. et al. Decreased diversity of the fecal Microbiome in recurrent Clostridium

747   difficile-associated diarrhea. *J Infect Dis* **197**, 435-438 (2008).

748   10.   Seekatz, A. M., Rao, K., Santhosh, K. & Young, V. B. Dynamics of the fecal microbiome

749   in patients with recurrent and nonrecurrent Clostridium difficile infection. *Genome Med* **8**,

750   47 (2016).

751   11.   Khanna, S. et al. Gut microbiome predictors of treatment response and recurrence in

752   primary Clostridium difficile infection. *Aliment Pharmacol Ther* **44**, 715-727 (2016).

753   12.   Martinez-Augustin, O. & Sanchez de Medina, F. Intestinal bile acid physiology and

754   pathophysiology. *World J Gastroenterol* **14**, 5630-5640 (2008).

755   13.   Sorg, J. A. & Sonenshein, A. L. Bile salts and glycine as cogerminants for Clostridium

756   difficile spores. *J Bacteriol* **190**, 2505-2512 (2008).

757   14.   Buffie, C. G. et al. Precision microbiome reconstitution restores bile acid mediated

758   resistance to Clostridium difficile. *Nature* **517**, 205-208 (2015).

759   15.   Hryckowian, A. J. et al. Microbiota-accessible carbohydrates suppress Clostridium difficile

760   infection in a murine model. *Nat Microbiol* **3**, 662-669 (2018).

761   16.   Antharam, V. C. et al. Intestinal dysbiosis and depletion of butyrogenic bacteria in

762   Clostridium difficile infection and nosocomial diarrhea. *J Clin Microbiol* **51**, 2884-2892

763   (2013).

764   17.   Zackular, J. P. et al. Dietary zinc alters the microbiota and decreases resistance to

765   Clostridium difficile infection. *Nat Med* **22**, 1330-1334 (2016).

766   18.   Mefferd, C. C. et al. A High-Fat/High-Protein, Atkins-Type Diet Exacerbates

767   Clostridioides (Clostridium) difficile Infection in Mice, whereas a High-Carbohydrate Diet

768   Protects. *mSystems* **5**, e00765-19 (2020).

769   19.   Jose, S. et al. Obeticholic acid ameliorates severity of Clostridioides difficile infection in

770   high fat diet-induced obese mice. *Mucosal Immunology* **14**, 500-510 (2021).

20. Moore, J. H. et al. Defined Nutrient Diets Alter Susceptibility to Clostridium difficile

Associated Disease in a Murine Model. *PLoS One* **10**, e0131829 (2015).

21. Devkota, S. et al. Dietary-fat-induced taurocholic acid promotes pathobiont expansion and

colitis in Il10-/- mice. *Nature* **487**, 104-108 (2012).

22. Chen, X. et al. A mouse model of Clostridium difficile-associated disease.

Gastroenterology **135**, 1984-1992 (2008).

23. Dieleman, L. A. et al. Chronic experimental colitis induced by dextran sulphate sodium

(DSS) is characterized by Th1 and Th2 cytokines. *Clin Exp Immunol* **114**, 385-391 (1998).

24. Barthel, M. et al. Pretreatment of mice with streptomycin provides a Salmonella enterica

serovar Typhimurium colitis model that allows analysis of both pathogen and host. *Infect*

Immun **71**, 2839-2858 (2003).

25. Benjamini, Y. & Hochberg, Y. Multiple Hypotheses Testing with Weights. *Scandinavian*

Journal of Statistics **24**, 407-418 (1997).

26. Francis, M. B., Allen, C. A., Shrestha, R. & Sorg, J. A. Bile acid recognition by the

Clostridium difficile germinant receptor, CspC, is important for establishing infection.

PLoS Pathog **9**, e1003356 (2013).

27. Sorg, J. A. & Sonenshein, A. L. Chenodeoxycholate is an inhibitor of Clostridium difficile

spore germination. *J Bacteriol* **191**, 1115-1117 (2009).

28. Sorg, J. A. & Sonenshein, A. L. Inhibiting the initiation of Clostridium difficile spore

germination using analogs of chenodeoxycholic acid, a bile acid. *J Bacteriol* **192**, 4983-

4990 (2010).

29.   Weingarden, A. R. et al. Ursodeoxycholic Acid Inhibits Clostridium difficile Spore

793   Germination and Vegetative Growth, and Prevents the Recurrence of Ileal Pouchitis

794   Associated With the Infection. *J Clin Gastroenterol* **50**, 624-630 (2016).

795   30.   Allegretti, J. R. et al. Recurrent Clostridium difficile infection associates with distinct bile

796   acid and microbiome profiles. *Aliment Pharmacol Ther* **43**, 1142-1153 (2016).

797   31.   Robinson, J. I. et al. Metabolomic networks connect host-microbiome processes to human

798   Clostridioides difficile infections. *J Clin Invest* **129**, 3792-3806 (2019).

799   32.   Francis, M. B., Allen, C. A. & Sorg, J. A. Muricholic acids inhibit Clostridium difficile

800   spore germination and growth. *PLoS One* **8**, e73653 (2013).

801   33.   Karlsson, S., Lindberg, A., Norin, E., Burman, L. G. & Akerlund, T. Toxins, butyric acid,

802   and other short-chain fatty acids are coordinately expressed and down-regulated by

803   cysteine in Clostridium difficile. *Infect Immun* **68**, 5881-5888 (2000).

804   34.   Seekatz, A. M. et al. Restoration of short chain fatty acid and bile acid metabolism

805   following fecal microbiota transplantation in patients with recurrent Clostridium difficile

806   infection. *Anaerobe* **53**, 64-73 (2018).

807   35.   Lozupone, C. A., Hamady, M., Kelley, S. T. & Knight, R. Quantitative and qualitative beta

808   diversity measures lead to different insights into factors that structure microbial

809   communities. *Appl Environ Microbiol* **73**, 1576-1585 (2007).

810   36.   Allen, B., Kon, M. & Bar-Yam, Y. A new phylogenetic diversity measure generalizing the

811   shannon index and its application to phyllostomid bats. *Am Nat* **174**, 236-243 (2009).

812   37.   Kriss, M., Hazleton, K. Z., Nusbacher, N. M., Martin, C. G. & Lozupone, C. A. Low

813   diversity gut microbiota dysbiosis: drivers, functional implications and recovery. *Curr*

814   *Opin Microbiol* **44**, 34-40 (2018).

38.   Langille, M. G. et al. Predictive functional profiling of microbial communities using 16S

rRNA marker gene sequences. *Nat Biotechnol* **31**, 814-821 (2013).

39.   Ridlon, J. M., Kang, D. J. & Hylemon, P. B. Bile salt biotransformations by human

intestinal bacteria. *The Journal of Lipid Research* **47**, 241-259 (2006).

40.   Vital, M., Howe, A. C. & Tiedje, J. M. Revealing the bacterial butyrate synthesis pathways

by analyzing (meta)genomic data. *mBio* **5**, e00889 (2014).

41.   Vital, M., Gao, J., Rizzo, M., Harrison, T. & Tiedje, J. M. Diet is a major factor governing

the fecal butyrate-producing community structure across Mammalia, Aves and Reptilia.

ISME J **9**, 832-843 (2015).

42.   Zacharioudakis, I. M., Zervou, F. N., Pliakos, E. E., Ziakas, P. D. & Mylonakis, E.

Colonization with toxinogenic C. difficile upon hospital admission, and risk of infection: a

systematic review and meta-analysis. *Am J Gastroenterol* **110**, 381-90; quiz 391 (2015).

43.   Lewis, S., Burmeister, S. & Brazier, J. Effect of the prebiotic oligofructose on relapse of

Clostridium difficile-associated diarrhea: a randomized, controlled study. *Clin*

Gastroenterol Hepatol **3**, 442-448 (2005).

44.   Blankenship-Paris, T. L., Chang, J., Dalldorf, F. G. & Gilligan, P. H. In vivo and in vitro

studies of Clostridium difficile-induced disease in hamsters fed an atherogenic, high-fat

diet. *Lab Anim Sci* **45**, 47-53 (1995).

45.   Blankenship-Paris, T. L., Walton, B. J., Hayes, Y. O. & Chang, J. Clostridium difficile

infection in hamsters fed an atherogenic diet. *Vet Pathol* **32**, 269-273 (1995).

46.   Tomkovich, S., Stough, J. M. A., Bishop, L. & Schloss, P. D. The Initial Gut Microbiota

and Response to Antibiotic Perturbation Influence Clostridioides difficile Clearance in

Mice. *mSphere* **5**, e00869-20 (2020).

47. Kumar, N. et al. Adaptation of host transmission cycle during Clostridium difficile

speciation. *Nat Genet* **51**, 1315-1320 (2019).

48. Di Bella, S., Ascenzi, P., Siarakas, S., Petrosillo, N. & di Masi, A. Clostridium difficile

Toxins A and B: Insights into Pathogenic Properties and Extraintestinal Effects. *Toxins*

(Basel) **8**, 134-125 (2016).

49. Carter, G. P. et al. Defining the Roles of TcdA and TcdB in Localized Gastrointestinal

Disease, Systemic Organ Damage, and the Host Response during Clostridium difficile

Infections. *mBio* **6**, e00551 (2015).

50. Savidge, T. C. et al. Clostridium difficile toxin B is an inflammatory enterotoxin in human

intestine. *Gastroenterology* **125**, 413-420 (2003).

51. Dupuy, B. & Sonenshein, A. L. Regulated transcription of Clostridium difficile toxin

genes. *Mol Microbiol* **27**, 107-120 (1998).

52. Litvin, M. et al. Identification of a pseudo-outbreak of Clostridium difficile infection (CDI)

and the effect of repeated testing, sensitivity, and specificity on perceived prevalence of

CDI. *Infect Control Hosp Epidemiol* **30**, 1166-1171 (2009).

53. Walker, R. C. et al. Comparison of culture, cytotoxicity assays, and enzyme-linked

immunosorbent assay for toxin A and toxin B in the diagnosis of Clostridium difficile-

related enteric disease. *Diagn Microbiol Infect Dis* **5**, 61-69 (1986).

54. Theriot, C. M., Bowman, A. A. & Young, V. B. Antibiotic-Induced Alterations of the Gut

Microbiota Alter Secondary Bile Acid Production and Allow for Clostridium difficile

Spore Germination and Outgrowth in the Large Intestine. *mSphere* **1**, e00045-15 (2016).

55. Howerton, A., Ramirez, N. & Abel-Santos, E. Mapping interactions between germinants

and Clostridium difficile spores. *J Bacteriol* **193**, 274-282 (2011).

56.   Ding, S. et al. High-fat diet: bacteria interactions promote intestinal inflammation which

precedes and correlates with obesity and insulin resistance in mouse. *PLoS One* **5**, e12191

(2010).

57.   Doerner, S. K. et al. High-Fat Diet-Induced Complement Activation Mediates Intestinal

Inflammation and Neoplasia, Independent of Obesity. *Mol Cancer Res* **14**, 953-965 (2016).

58.   Ng, K. M. et al. Recovery of the Gut Microbiota after Antibiotics Depends on Host Diet,

Community Context, and Environmental Reservoirs. *Cell Host Microbe* **26**, 650-665.e4

(2019).

59.   Cabral, D. J. et al. Microbial Metabolism Modulates Antibiotic Susceptibility within the

Murine Gut Microbiome. *Cell Metab* **30**, 800-823.e7 (2019).

60.   Kelly, C. J. et al. Crosstalk between Microbiota-Derived Short-Chain Fatty Acids and

Intestinal Epithelial HIF Augments Tissue Barrier Function. *Cell Host Microbe* **17**, 662-

671 (2015).

61.   Kanauchi, O. et al. Butyrate from bacterial fermentation of germinated barley foodstuff

preserves intestinal barrier function in experimental colitis in the rat model. *J Gastroenterol*

Hepatol **14**, 880-888 (1999).

62.   Wang, H. B., Wang, P. Y., Wang, X., Wan, Y. L. & Liu, Y. C. Butyrate enhances intestinal

epithelial barrier function via up-regulation of tight junction protein Claudin-1

transcription. *Dig Dis Sci* **57**, 3126-3135 (2012).

63.   Peng, L., He, Z., Chen, W., Holzman, I. R. & Lin, J. Effects of butyrate on intestinal barrier

function in a Caco-2 cell monolayer model of intestinal barrier. *Pediatr Res* **61**, 37-41

(2007).

64.   Furusawa, Y. et al. Commensal microbe-derived butyrate induces the differentiation of

884   colonic regulatory T cells. *Nature* **504**, 446-450 (2013).

885   65.   Singh, V. et al. Dysregulated Microbial Fermentation of Soluble Fiber Induces Cholestatic

886   Liver Cancer. *Cell* **175**, 679-694.e22 (2018).

887   66.   Klurfeld, D. M. et al. Considerations for best practices in studies of fiber or other dietary

888   components and the intestinal microbiome. *Am J Physiol Endocrinol Metab* **315**, E1087-

889   E1097 (2018).

890   67.   Thompson, L. R. et al. A communal catalogue reveals Earth's multiscale microbial

891   diversity. *Nature* **551**, 457-463 (2017).

892   68.   Bolyen, E. et al. Reproducible, interactive, scalable and extensible microbiome data science

893   using QIIME 2. *Nat Biotechnol* **37**, 852-857 (2019).

894   69.   Callahan, B. J. et al. DADA2: High-resolution sample inference from Illumina amplicon

895   data. *Nat Methods* **13**, 581-583 (2016).

896   70.   Janssen, S. et al. Phylogenetic Placement of Exact Amplicon Sequences Improves

897   Associations with Clinical Information. *mSystems* **3**, 581 (2018).

898   71.   Wang, Q., Garrity, G. M., Tiedje, J. M. & Cole, J. R. Naive Bayesian classifier for rapid

899   assignment of rRNA sequences into the new bacterial taxonomy. *Appl Environ Microbiol*

900   **73**, 5261-5267 (2007).

901   72.   Quast, C. et al. The SILVA ribosomal RNA gene database project: improved data

902   processing and web-based tools. *Nucleic Acids Res* **41**, D590-6 (2013).

903   73.   Yilmaz, P. et al. The SILVA and "All-species Living Tree Project (LTP)" taxonomic

904   frameworks. *Nucleic Acids Res* **42**, D643-8 (2014).

905   74.   Anaconda Software Distribution.

906   75.   Kelly, C. J. et al. Oral vitamin B_{12} supplement is delivered to the distal gut, altering the

907   corrinoid profile and selectively depleting Bacteroides in C57BL/6 mice. *Gut Microbes* **10**,

908   654-662 (2019).

909   76.   Sarafian, M. H. et al. Bile acid profiling and quantification in biofluids using ultra-

910   performance liquid chromatography tandem mass spectrometry. *Anal Chem* **87**, 9662-9670

911   (2015).

912   77.   Wickham, H. Tidyverse.

913   78.   Kassambara, A., Kosinski, M., Biecek, P. & Fabian, S. survminer: Drawing Survival

914   Curves using 'ggplot2'

915   79.   Therneau, T. M. & Lumley, T. survival: Survival Analysis.

916   80.   Wickham, H. *ggplot2* (Springer, 2010).

917   81.   Ahlmann-Eltze, C. ggsignif: Significance Brackets for 'ggplot2'.

918   82.   Wilke, C. O. cowplot: Streamlined Plot Theme and Plot Annotations for

919   'ggplot2'

920   83.   Gonzalez, A. et al. Qiita: rapid, web-enabled microbiome meta-analysis. *Nat Methods* **15**,

921   796-798 (2018).

Supplementary Materials

Figure S1: Example sections of cecal (left) and colon (right) tissues with low or medium

inflammation. No samples had levels of inflammation considered to be high. The cecum was

scored for injury according to the system of Barthel et al, 2003 [24]. Scoring of inflammation

using the Barthel scoring system is restricted to neutrophils in the mucosa portion of the cecum.

In low cecal inflammation, no neutrophils are observed in the mucosa (red bracket) while they

are observed with medium inflammation. Medium cecal inflammation also displayed

submucosal edema (blue bracket) that is thought to occur at least to some degree due to the

neutrophils present in the submucosa. The colon (right panels) was scored for injury according to

the system of Dieleman et al, 1998 [23]. This system takes into account the relative quantity of

inflammatory cells as well as whether they are found only in the mucosa layer (red bracket), are

also in the submucosa (green arrow), or are found all the way through the muscularis (blue

bracket) and into the peritoneal cavity.

Figure S2. Cecal bile acid concentrations (pmol/kg) by diet. Unconjugated primary bile acids

(A), secondary bile acid (B) and taurine-conjugated bile acids (C). a_MCA (alpha muricholic

acid); b_MCA (beta muricholic acid); CA (cholic acid); CDCA (chenodeoxycholic acid); UDCA

(ursodeoxycholic acid); DCA (deoxycholic acid); LCA (lithocholic acid); T_b_MCA (tauro-beta

muricholic acid); TCA (taurocholic acid); TCA_3_SO4 (taurocholic acid 3-sulfate); TCDCA

(taurochenodeoxycholic acid); TDCA (taurodeoxycholic acid); TLCA (taurolithocholic acid). P-

values for the Kruskal-Wallis test with Dunn's post hoc are noted. Median and IQR are

indicated. Plots are colored based on the previously described influence of each bile acid on *C.*

difficile.

Figure S3: Statistical summary of relative abundances of key taxa from Figure 7a. Significant

differences between groups are noted as calculated with the Wilcoxon rank-sum test. ***:

p<0.001. **: p<0.01, *: p<0.05, ns= non significant.

Table S1: Composition of low-fat/low-fiber/high-sucrose diet

	Low-fat/Low-fiber/High-Sucrose

Fat (% kcal)
  (% SFA)
  (% MUFA)
  (% PUFA)
	17.0
	(19.5)
	(41.7)
	(38.8)

Carbohydrates (% kcal) *(Sucrose)*	64.5 *(26.7)*
Protein (%kcal)	18.6
Fiber (g/kg)	50 (cellulose)

961 **Table S2:** Description of samples sizes across different assays in cohorts 1 and 2.
962

		Batches	Chow	LF/LF	WD
Cohort 1 (stool collection)		Total	20	20	20
		Batch1	10	0	10
		Batch2	10	10	0
		Batch3	0	5	5
		Batch4	0	5	5
	16S rRNA	Batches 1,2,3	13	5	13
Cohort 2 (tissue collection)		Total	20	20	26
		Batch1	0	5	0
		Batch2	0	5	0
		Batch3	6	0	6
		Batch4	5	5	5
		Batch5	0	5	5
		Batch6	5	0	5
		Batch7	4	0	5
	histology	Batches 4,5,7	9	10	14
	TcdA	4,5	5	10	9
	TcdB	1,2,4,5,7	8	20	15
	Bile acids	All batches	20	20	25
	SCFAs	1,2,4-7	14	18	19

963
Cohort 1 (longitudinal mortality analysis with serial fecal collection and microbiome
sequencing) was conducted in 4 different batches of mice with per batch counts across diets
indicated. Cohort 2 (tissue collection – intestines for histopathology and cecal aspirates for bile
acids, SCFAs, and toxins) was conducted in 7 different batches of mice with per batch counts
across diets indicated. 16S rRNA, histology, toxins TcdA and TcdB, bile acids and SCFAs were
all measured for particular sets of batches as indicated.

Table S3

Complete results of linear modeling statistical analyses performed. **[Table in separate Excel**

spreadsheet file]

973
974
975   **Table S4**
976

Ingredient (g/KG)	WD	LF/LF
Anhydrous Milkfat	36.3	50
Beef Tallow	24.8	0
Casein	190	0
Cellulose	30	35
Cholesterol	0.4	0
Choline Bitartrate	2.1	0.01
Corn Oil	16.5	6.5
Corn Starch	230	392.2
L-Cystine	2.85	3
Lard	28	0
Maltodextrin	70	100
Mineral Mix, nTWD (110422)	35	15
Olive Oil	28	16.5
Sodium Chloride	4	0
Soybean Oil	31.4	23
Sucrose	255.6	24
TBHQ, antioxidant	0.028	0.014
Thiamin (81%)	0.015	0.002
Vitamin K1, phylloquinone	0.003	0.014
Vitamin Mix*	15	2.75

977
978   * WD used the Vitamin Mix nTWD (110423) and LF/LF diet used AIN-93-VX (94047)
979

Acknowledgments: We would like to thank Jordi Lanis and Sean Colgan for advice on the

employed CDI mouse model and Sally Stabler and Whitney Phinney for their assistance in

measuring SCFAs. We also appreciate the contribution to this research made by E. Erin Smith,

TL(ASCP)CMQIHC, Jenna Van Der Volgen, HT(ASCP)CM, Allison Quador, HTL

(ASCP)CM, and Jessica Arnold HTL(ASCP)CM of the University of Colorado

Funding: This work was supported by NIH U01 AI150589. Additional support was provided by

the University of Colorado Department of Medicine's Outstanding Early Career Science Award

program as well as support to Keith Hazleton from the Institutional Training Grant for Pediatric

Gastroenterology from NIDDK (5T32-DK067009-12), Clinical Fellow Awards from the Cystic

Fibrosis Foundation (HAZLET18DO and HAZLET19DO) and The Judith Sondheimer Pediatric

GI Fellow Research Fund. Kathleen Arnolds was supported by T32-AI007405 Training Program

in Immunology. High performance computing was supported by a cluster at the University of

Colorado Boulder funded by National Institutes of Health 1S10OD012300. The Denver

Histology Shared Resource is supported in part by the Cancer Center Support Grant

(P30CA046934).

Author contributions: KH conceived of and conducted experiments, analyzed data and co-

wrote the paper; CM analyzed data, made figures, and contributed to data interpretation and

writing; KA generated and analyzed data from toxin ELISA; NN generated 16S rRNA sequence

and qPCR data and aided in analysis and results interpretation; NMH aided in mouse

experiments, 16S rRNA sequencing and generation of toxin ELISA data; NR and MA worked

with KH to develop a novel bile acid panel and aided in analysis and interpretation of results; CL

directed and contributed to all aspects of the project. All authors contributed to the manuscript.

1004

1005　**Competing interests:** The authors declare that there are no competing interests.

1006