Stability of neocortical synapses across sleep and wake

Brian A. Cary[1], Gina G. Turrigiano[1,*]

1: Dept of Biology, Brandeis University, Waltham, MA 02453, USA

*: Corresponding author (email: turrigiano@brandeis.edu)

ABSTRACT

Sleep is important for brain plasticity, but its exact function remains mysterious. An influential but controversial idea is that a crucial function of sleep is to drive widespread downscaling of excitatory synaptic strengths. Here we used real-time sleep classification, *ex vivo* measurements of postsynaptic strength, and *in vivo* optogenetic monitoring of thalamocortical synaptic efficacy to ask whether sleep and wake states can constitutively drive changes in synaptic strength within the neocortex of juvenile rats. We found that miniature EPSC amplitudes onto L4 and L2/3 pyramidal neurons were stable across sleep and wake dense epochs in both primary visual (V1) and prefrontal cortex (PFC). Further, chronic monitoring of thalamocortical synaptic efficacy in V1 of freely behaving animals revealed stable responses across even prolonged periods of natural sleep and wake. Together these data demonstrate that neocortical synaptic strengths are remarkably stable across sleep and wake states, and provide strong evidence against the view that sleep drives widespread synaptic downscaling at neocortical synapses.

INTRODUCTION

Sleep is a widely expressed behavior, present in animals as evolutionarily distant as Cassiopea jellyfish and the Australian dragon lizard (Nath et al., 2017; Idelson et al., 2016). Despite its ubiquity and long history of scientific study, sleep – and more broadly the function of brain states – remains deeply mysterious. Sleep disruption can perturb learning and memory consolidation (Walker and Stickgold, 2004; Krause et al., 2017), presumably through the modulation of synaptic plasticity (Frank and Cantera, 2014). However, there is little agreement on the nature of this regulation. Researchers have variously proposed that sleep stabilizes, strengthens, weakens, or even prunes synapses (Kavanau, 1996; Datta et al., 2008; Chauvette et al., 2012; de Vivo et al., 2017; Li et al., 2017). Further, there is disagreement on whether sleep primarily enables correlation-based plasticity mechanisms such as long-term potentiation (LTP),

or homeostatic forms of plasticity that serve the function of regulating overall synaptic strength

to stabilize neuron and circuit function (Frank and Cantera, 2014, Aton et al., 2014; Hengen et

al., 2016). Finally, it is unclear whether sleep is merely permissive for some forms of synaptic

plasticity (Torrado Pacheco et al., 2020), or whether being asleep is by itself sufficient to induce

synaptic changes without a preceding salient learning event (Tononi and Cirelli, 2014). Here we

use a combination of real-time sleep classification, *ex vivo* measurements of postsynaptic

strength in two neocortical areas and two different cell types, and *in vivo* optogenetic monitoring

of evoked thalamocortical transmission in primary visual cortex (V1) to ask whether sleep and

wake states constitutively drive widespread synaptic plasticity within neocortical circuits.

One influential hypothesis, the synaptic homeostasis hypothesis (SHY), has motivated much

work on the role of sleep in brain plasticity (Tononi and Cirelli, 2014). SHY proposes that

memories are formed during wake when animals actively sample their environment, primarily

through the induction of Hebbian LTP-like mechanisms that causes a net potentiation of

synapses. This process would saturate synapses if left unopposed (Miller and MacKay, 1994;

Abbott and Nelson, 2000; Turrigiano and Nelson, 2004), and sleep is proposed to be an offline

state that allows neurons to renormalize synaptic weights by downscaling synaptic strengths

(Tonoi and Cirelli, 2014). This renormalization is postulated to be a global process that affects

all or most synapses in many brain regions (Tononi and Cirelli, 2014). Critically, SHY predicts

that excitatory synapses should on average be stronger after a period of wake, and weaker after a

period of sleep. A feature of NREM sleep is slow wave activity (SWA), in which the

electroencephalogram (EEG) expresses large, low frequency waves (<4.5 Hz, Dijk, 2009). It is

well established that SWA is high during NREM immediately after prolonged wake and

progressively decreases as sleep pressure diminishes (Dijk, 2009). SHY additionally proposes

that the size of slow waves is correlated with cortical synaptic strengths, and that the oscillation

in SWA with sleep and wake is driven by sleep-dependent synaptic downscaling that then

diminishes SWA (Tononi and Cirelli, 2014; Tononi, 2009).

Studies supporting SHY have found that the expression of proteins associated with synaptic

potentiation are higher, axon-spine interfaces (a corollary of synaptic strength) are larger, and

evoked transcallosal cortical responses increase in slope after a period of wake (Vyazovksiy et

al., 2008; de Vivo et al., 2017; de Vivo et al., 2019; Diering et al., 2017). Importantly, these

changes were observed without introducing any salient learning experiences for the animal,

suggesting that simply being asleep or awake reduces or increases net synaptic weights,

respectively. In contrast other recent studies have found that synaptic transmission is potentiated

by sleep (Chauvette et al., 2012) or is unaffected by sleep deprivation (Matsumoto et al., 2020).

If excitatory synapses indeed oscillate in strength across sleep and wake states, then the firing

rate of individual neurons would be expected to oscillate as well; while some studies have found

such oscillations (but with variable effects between brain regions and neuronal populations;

Vyazovskiy et al., 2009; Miyawaki & Diba, 2016; Miyawaki et al., 2019; Watson et al., 2016),

others have observed stable firing across periods of sleep and wake (Hengen et al., 2016;

Torrado-Pacheco et al., 2020).

Together, these studies paint a complex picture of the possible roles of sleep and wake states in

modulating synaptic plasticity. Some discrepancies between studies are likely due to

methodological differences; for instance, not all studies clearly differentiate between effects of

sleep and circadian cycle, carefully classify sleep states and prior sleep history, or directly

measured synaptic strengths. Here we set out to carefully test the hypothesis that neocortical

synaptic weights weaken during sleep and strengthen during wake, using both *ex vivo* slice

physiology and *in vivo* monitoring of synaptic strengths and evoked spiking. We used *in vivo*

behavioral state classification in real time to track the accumulation of sleep and wake, which

allowed us to detect natural sleep or wake-dense epochs that occurred within a defined 5 hour

circadian window (ZT 3-8). We then immediately cut slices and measured miniature excitatory

postsynaptic currents (mEPSCs) from pyramidal neurons in V1 or prefrontal cortex (PFC;

specifically, prelimbic and infralimbic cortex). We found that mEPSC amplitudes were stable

across sleep and wake dense epochs in both brain regions. To extend this finding to evoked

transmission, we used *in vivo* optogenetics to monitor thalamocortical synaptic drive to visual

cortex across natural sleep-wake epochs. Again, we found that thalamocortical synaptic drive

and evoked spiking were stable across even prolonged periods of sleep and wake. Together our

data show that neocortical synaptic strengths are remarkably stable across naturally occurring

periods of sleep and wake and provide strong evidence against the view that a universal function

of sleep is to drive widespread constitutive weakening of excitatory synaptic strengths.

RESULTS

Sleep/wake behavior of Juvenile Long Evans rats

We wished to compare several functional measures of neocortical synaptic strength after natural periods of prolonged waking or sleeping. Here and in subsequent experiments we used juvenile rats (postnatal days 25 to 31) to allow for high-quality endpoint slice recordings, and to ensure that our experiments were performed within a highly plastic developmental window when Hebbian and homeostatic plasticity are known to be pronounced (Lambo et al., 2013; Hengen et al., 2016; Smith et al., 2008; Espinosa and Stryker, 2012). To this end we first analyzed behavioral data from animals in this age range, to understand their natural rhythms of sleep and wake across Zeitgeber (ZT) time, and determine when they are likely to experience prolonged periods of sleep or wake (Fig. 1A-D). We monitored electromyograms (EMGs), EEG, and video, and used standard approaches to classify sleep states into wake and rapid eye movement (REM) or non-rapid eye movement slow wave (NREM) sleep, as described previously (Hengen et al., 2016; Torrado Pacheco et al., 2019, 2020; see Methods). In some experiments we also differentiated active and quiet wake, as noted (Fig. 5-8, Fig 2 - supplement 1). As expected, animals on average slept more during the light phase (ZT 0-12, ~60% asleep) than during the dark phase (ZT 12-24, ~40% asleep, Fig. 1B, D; Frank and Heller, 1997; Frank et al., 2017); however, there was considerable variability from day to day and animal to animal in when they experienced periods of relatively consolidated sleep or wake (defined as a 4 hr period with >65% of time in that state, Fig. 1A, compare animals 1 and 2). Another prominent feature of sleep is the modulation of slow wave amplitude by prior sleep history (Dijk, 2009; Tononi and Cirelli, 2014). As expected, the size of slow waves in NREM (measured as average delta power, Dijk, 2009; Vyazovksiy et al., 2008; Fig. 1C), rose as net time spent awake accumulated in the dark, and dropped during the day when animals typically spent more time sleeping (Fig 1B,C). This oscillation in delta power is similar to that seen in adult rodents (Leemburg et al., 2010). Thus, these animals exhibit homeostasis in slow wave amplitude across sleep and wake cycles, which SHY suggests is causally linked to synaptic downscaling (Tononi and Cirelli, 2014; Tononi, 2009).

Real-time sleep classification

To study the impact of sleep and wake on synaptic efficacy it is necessary to disentangle sleep

from circadian effects. Since prolonged natural periods of wake and sleep tend to occur during

distinct circadian periods, many studies have accomplished this by sacrificing animals at the

same ZT (typically several hours after lights on) and comparing animals that have slept naturally

to those that were continuously kept awake via sleep deprivation (Liu et al., 2010; Diering et al.,

2017; de Vivo et al., 2017, 2019). However, the animal-to-animal and cycle-to-cycle variability

in the timing of consolidated sleep means that even when animals are sacrificed at the exact same

ZT, there are likely to be considerable differences in the amount of consolidated sleep/wake (for

example, compare animals 1 and 2 several hours after lights on, red arrows in Fig. 1A). This

prompted us to develop an approach to classify sleep and wake in real time, so we could reliably

detect consolidated periods of sleep or wake within a defined circadian window for each animal,

and then sacrifice them to probe synaptic strengths *ex vivo*.

To classify behavioral state in real time, we collected EEG, EMG from the nuchal (neck)

muscles, and video as they freely behaved in an enriched environment. The recording chamber

included a littermate separated by a thin, transparent plexiglass wall with nose-poke holes to

allow for social interactions, toys for stimulation and play, and food and water available ad

libitum. The data were acquired, analyzed, and plotted all within a custom program, which uses

canonical markers to classify behavioral state into NREM, REM, or wake (Fig. 2A). Briefly, the

program computes a delta/beta and a theta/delta ratio from the corresponding frequency bands in

the EEG (delta: 0.5-4 Hz; theta: 5-8 Hz; beta ~ 20-35 Hz). Large deflections in the EMG are z-

scored and animal pixel movement is extracted from the video recording. The classifier reads in

these variables and applies a simple semi-automated decision tree, which uses 3 manually

adjustable thresholds (delta, theta, and movement thresholds). Given what is known about the

probability of sleep transitions, the classifier then adjusts these thresholds based on recent state

history (additional details in Methods). With this technique, we could determine the sleep/wake

state of the animal with an accuracy of 93.3% when benchmarked against manual coding (Fig.

2D). All classifications made in real time were manually verified post hoc. We could specify the

length and required sleep or wake density for an epoch to pass threshold; for this set of

experiments we chose threshold values that were within the range used by others to test SHY

(Liu et al., 2010; Diering et al., 2017; Vyazovskiy et al., 2009; de Vivo et al., 2019): 4 hr of

>65% within a state, with the last hour >70% within the state. The 4 hr requirement was chosen

because it is the longest span of time the rats naturally spent in a sleep dense epoch in a typical

day, and the last hr threshold was to ensure that synaptic differences were not rapidly reversed by

recent behavioral state changes. The average density of sleep epochs that exceeded threshold and

were included in the mEPSC analysis was 69%; the average wake density during the light period

was 76% and during the dark period was 71%.

We wished to compare three conditions: sleep dense and wake dense epochs that ended during a

similar circadian period (ZT 3-8, Fig. 3A, B), and wake dense epochs that ended during the

opposite circadian period (dark period, ZT 13-16, Fig. 3C); this 3-way comparison allowed us to

look for differences driven by sleep/wake (sleep and wake at the same ZT) and differences

driven by circadian time (wake at opposite ZTs). Spontaneous wake dense epochs were

infrequent in the light period, so when we detected a long natural wake epoch during the early

light phase we added new toys to the chamber (example: red arrow in Fig. 3B, top panel) to

encourage additional waking. This was sufficient to extend natural waking epochs to reach

threshold for consolidated wake within the specified circadian window (Fig 3B, Fig 3 –

supplement 2).

Postsynaptic strengths are stable across extended periods of sleep or wake

In order to assess whether sleep induces a global downscaling of postsynaptic strengths, we

tracked sleep/wake history in real time, cut acute slices once an animal finished a sleep or wake

dense epoch (Fig. 3), and performed whole-cell patch electrophysiology to record miniature

excitatory postsynaptic currents (mEPSCs) onto pyramidal neurons in two brain regions (visual

cortex and prefrontal cortex, PFC) and two layers (L4 and L2/3, see below for rationale for these

targets). Ex *vivo* slice physiology is an established method to probe experience-dependent

changes in synaptic strengths (Khurana and Li, 2013; Lambo and Turrigiano, 2013; Nataraj et

al., 2010; Heynan et al., 2003; Nate Miska et al., 2018), and mEPSC amplitudes are a standard

readout of synaptic up- or downscaling and are a direct correlate of postsynaptic strength

(Turrigiano, 2008). The sleep history (expressed as hypnograms that show the time in each

sleep/wake state) of each animal included in our mEPSC analysis are shown in Fig. 3A-C, top

traces, and in Fig. 3 – supplement 2. Another way of visualizing an individual animal's

sleep/wake history is to generate a moving average of the percent of time spent awake in the last

4 hours (Fig 3A-C, bottom traces; Fig. 3 – supplement 2); this illustrates when in ZT each animal

passed the threshold for our endpoint mEPSC analysis. Using this approach we were able to

measure mEPSC amplitudes at similar circadian time points from animals that had spent the

majority of the proceeding 4 hours either awake or asleep, with minimal perturbation (Fig 3A,

B); and from animals with similar wake experience at the opposite circadian time (Fig. 3C).

We first recorded from layer 2/3 pyramidal neurons in primary visual cortex (Fig. 3D). We chose

visual cortex as it is highly plastic during this developmental window, which is within the classic

visual system critical period (Espinosa and Stryker, 2012). Further, homeostatic mechanisms are

present and are able to up- and downscale mEPSC amplitudes onto L2/3 pyramidal neurons

during this developmental window (Kaneko et al., 2008; Lambo et al., 2013; Hengen et al., 2013,

Torrado Pacheco et al., 2020). When we computed the mean mEPSC amplitude by cell and

compared this across conditions, we found no differences in amplitude after sleep dense or wake

dense epochs in the light phase (Sleep and L.Wake), or wake dense epochs in the dark period (D.

Wake; Fig. 3D, E). Here and below, cellular properties such as input resistance and resting

potential were not different between conditions; mEPSC kinetics (rise times, decay tau, and

scaled waveforms) were also not different between conditions. To examine more closely the

mEPSC amplitude distribution in the different conditions we randomly selected the same number

of events from each neuron and plotted individual mEPSC amplitudes as a cumulative

distribution (Fig. 3E, right). There were no significant differences in the amplitude distribution

between conditions. Thus neither prior sleep history nor circadian time significantly modulated

mEPSC amplitude onto visual cortical L2/3 pyramidal neurons.

We next examined mEPSCs onto pyramidal neurons in layer 4 of primary visual cortex. L4

pyramidal neurons receive extensive monosynaptic input from thalamic relay neurons, which

vary their firing patterns dramatically during sleep (Steriade, 2001; Steriade and Timofeev,

2003), and these thalamocortical synapses are also highly plastic at this time (Cooke and Bear,

2010; Miska et al., 2018). Similar to our observations in L2/3, mEPSC amplitudes were

indistinguishable after wake dense or sleep dense epochs, and there was no difference in the

cumulative amplitude distribution between conditions (Fig. 3F).

To determine whether this stability in postsynaptic strength was evident outside of primary

sensory cortex, we next examined mEPSCs onto L2/3 pyramidal neurons in PFC. The PFC

shows a strong modulation of slow-wave activity (Leemburg et al., 2010), has been suggested to

be especially impacted by extended wake (Muzur et al., 2002; Jones and Harrison, 2001), and a

previous study reported weaker synapses within prefrontal cortex after sleep (Liu et al., 2010).

However, consistent with our findings in primary visual cortex, mean mEPSC amplitude was

also stable across sleep and wake dense epochs in PFC, and was not significantly different after

extended wake at opposite circadian periods (Fig. 3G, left). The cumulative amplitude

distributions following extended sleep or wake in the light phase were indistinguishable (Fig.

3G, right), but interestingly there was a small but significant rightward shift at the high end of

this distribution in the D. Wake (SD - D. Wake, p=0.015). This suggests that there is a small

circadian modulation of mEPSC amplitude onto L2/3 pyramidal neurons in PFC.

We next examined the frequency of mEPSC events. MEPSC frequency can be influenced by

many factors (Han and Stevens, 2009; Wierenga et al., 2006), and is generally more variable

between neurons than amplitude (Lambo et al., 2013; Hengen et al., 2013; Liu et al., 2010; Fig.

4). We found no significant differences in mean mEPSC frequency across any of the conditions

or brain regions examined here (Fig. 4). Inter-event intervals were not different between

conditions in visual cortex L2/3 or L4. In PFC inter-event intervals were shorter in L. Wake

compared to both D. Wake and L. Sleep; thus there may be some interaction between time of day

and sleep/wake state on mEPSC frequency in PFC, that does not generalize to visual cortex.

Because we have the full behavioral history of each animal prior to preparing brain slices, we

could examine potential influences on mEPSCs beyond the duration of sleep or wake. There was

a reasonable range of wake densities (from 65% to 90%) within the wake dense condition, which

allowed us to evaluate possible correlations between wake density and amplitude or frequency of

mEPSCs; neither of these were significantly correlated (Fig. 3 – supplement 1). Further, we

could test an important proposed relationship between changes in slow wave amplitude and

excitatory synapses: according to SHY a larger reduction in slow wave amplitude (i.e. decrease

in delta power) should correlate with weaker synapses (Tononi and Cirelli, 2014; Tononi, 2009).

However, when we plotted the decrease in delta power during prior sleep against the endpoint

mEPSC amplitude or frequency, we again found no relationship (Fig. 3 – supplement 1A, B).

Together our data show that postsynaptic strength in a range of excitatory neocortical cell types

is remarkably stable across prolonged periods of sleep or wake.

Thalamocortical evoked field EPSPs are stable over the course of sleep and wake

MEPSC amplitude is a useful measure of postsynaptic strength, but does not take into account

potential changes in presynaptic function, and cannot be used to follow synaptic strengths

prospectively across sleep and wake epochs. To complement the mEPSC measurements we

therefore designed a paradigm to measure functional synaptic strength continuously *in vivo*

across sleep and wake epochs in freely behaving animals. In order to evoke synaptic responses in

visual cortex from a defined population of inputs we took advantage of our ability to express

ChR2 in thalamocortical terminals within V1 (Miska et al., 2018) by virally expressing ChR2

(AAV-ChR2-mCherry) in dorsal lateral geniculate nucleus (dLGN) of the thalamus, which

projects extensively to layer 4 of V1. After viral injection into dLGN, animals were given 1.5-2

weeks to allow for expression and transport of ChR2 to thalamocortical axon terminals in V1

(Fig. 5A), and we then implanted linear 16 channel silicon probes into V1 with adhered optic

fibers that rested on the cortical surface. With this configuration we could reliably evoke

thalamocortical responses with 1 ms pulses of blue light in freely behaving animals (Fig. 5B-E).

The linear design of the probe with exact electrode spacing, combined with current-source

density analysis (Fig. 5B) and post hoc histology, allowed us to estimate the placement of

electrical sites with high confidence. These evoked responses were similar to those evoked by

electrical stimulation of LGN, as well as visually evoked potentials (Cooke and Bear, 2010; Niell

and Stryker, 2008). The amplitude of the first negative trough of this evoked response has the

shortest latency in L4 (Fig. 5B-D), where it arises primarily from monosynaptic thalamic

excitatory synapses onto L4 neurons (Khibnik et al., 2010; Cooke and Bear, 2010), and thus

represents an evoked field excitatory postsynaptic potential (fEPSP). We monitored and analyzed

the amplitude and slope of this short-latency fEPSP (Fig. 5B-G, Fig. 6 – supplement 1) as a read-

out of functional thalamocortical synaptic efficacy in L4.

To chronically monitor the amplitude of thalamocortical fEPSPs, we used light intensities below

the peak saturation of responses (~50% of the peak response), and very low stimulation

frequencies (1/20 - 1/40 Hz) to avoid phototoxicity or the induction of plasticity from the

stimulation itself. With these parameters we were able to measure stable fEPSPs continuously for

several days (Fig. 5E), while simultaneously monitoring the sleep/wake state of the animal. An

interesting feature of these recordings is that fEPSP amplitude was rapidly modulated by

behavioral states (Fig. 5E, F), as previously reported (Vyazovskiy et al., 2008; Matsumoto et al.,

2020). To more fully capture behavioral state modulation of the fEPSP we further classified

wake into Quiet Wake and Active Wake using LFP and video data (Fig. 2 – supplement 1). We

found that evoked fEPSPs were largest on average during NREM, of similar size during quiet

wake and REM, and smallest during active wake (Fig. 5F). These changes in fEPSP amplitude

manifested rapidly upon transitions between states (Fig. 5E), consistent with a rapid

neuromodulatory effect (Lee and Dan, 2012; Brown et al., 2012; Matsumoto et al., 2020). A

second feature of these fEPSPs is that they were of consistently higher amplitude in all

behavioral states in the light than in the dark; there were no obvious circadian oscillations

beyond a rapid change in amplitude at the transitions between light and dark periods (Fig. 5G).

SHY predicts that there should be a gradual decrease in the size of the fEPSPs during sleep,

while wake should produce a gradual increase (Vyazovskiy et al., 2008). To test this we evoked

fEPSPs continuously at low frequency, identified sleep or wake dense epochs (using either a

>65% or >75% time-in-state threshold, Fig. 6A), and separately measured the fEPSP amplitude

during each of the 4 behavioral states (REM, NREM, Active Wake, and Quiet Wake; Fig. 6). We

then plotted fEPSP amplitude across sleep dense epochs measured either in NREM or REM (Fig.

6B, C) and across wake dense epochs measured in Active or Quiet Wake (Fig. 6D, E). Using a

65% threshold (Fig. 6B-E, left column) we were able to isolate natural sleep dense epochs of up

to 4.5 hours and wake dense epochs of up to 12.5 hours in duration; with a more stringent >75%

threshold (Fig. 6B-E, right column) these durations were somewhat shorter; up to 2.5 hr for

Sleep Dense, and 9 hr for wake dense.

Strikingly, we saw no progressive changes in fEPSP amplitude during prolonged sleep (Fig. 6B,

C) or wake (Fig. 6D, E) using either density threshold, even for the longest duration epochs. This

stability of the responses was evident regardless of which state we measured fEPSPs in (Fig. 6).

As fEPSP amplitude estimates can be contaminated by polysynaptic or spiking activity occurring

after the evoked monosynaptic thalamocortical synaptic potentials, we repeated this analysis

using a second standard measure of fEPSP magnitude, the slope of the initial downward

deflection. Consistent with the amplitude measurements, the slopes did not change across sleep

and wake states (Fig. 6 – supplement 1).

As a second means of probing for sleep or wake-induced changes in the fEPSP, we designed a

"sandwich" method for detecting changes across extended periods of sleep, by measuring the

amplitude in flanking periods of wake separated by an extended period of sleep, and vice versa

for extended periods of wake (Fig. 7A); all measurement were made in NREM for flanking

periods of sleep, and Quiet Wake for flanking periods of wake, and the results were identical

when we instead used REM and Active Wake, respectively. For this analysis we defined dense

epochs as >30 minute periods when the animal was >75% asleep/awake, and identified all such

epochs for each animal. Using this sandwich approach we again found that fEPSP amplitudes

were largely stable across periods of sleep or wake (Fig. 7B, C; < 2% change), indicating that the

intervening periods of sleep and wake produce no persistent change in synaptic strength. When

we plotted the change in fEPSP amplitude against the duration of the intervening sleep or wake

epoch, we find no significant correlation between epoch duration and change in fEPSP amplitude

(Fig.7D, E). Taken together with our mEPSC data, these experiments show that synaptic efficacy

onto L4 excitatory neurons is stable across extended periods of sleep and wake.

The ability of thalamocortical inputs to evoke spikes is stable over the course of sleep and

wake

Although thalamocortical fEPSP amplitudes were not modulated by time spent awake or asleep,

it is possible that the ability of thalamocortical synapses to evoke spiking in postsynaptic V1

neurons – a readout of functional connection strength – could be modified by prolonged periods

of sleep or wake. In addition to recording fEPSPs, our optrodes allowed us to sample spikes in

V1 evoked by stimulation of thalamocortical inputs (Fig. 8A). Using spike sorting as described

(Hengen et al., 2016; Torrado Pacheco et al., 2020), we were able to follow spiking in individual

regular-spiking (putative pyramidal) neurons over the course of sleep and wake epochs.

Although the probability of evoking a spike for any given stimulus was low, when averaged

across many stimuli we found that most well-isolated units had a detectable response to the

thalamocortical stimulation (Fig. 8A, B). Consistent with the rapid brain-state modulation of

fEPSP amplitude, there were more evoked spikes on average during NREM, an intermediate

number during REM, and fewer during wake (Fig. 8B). To determine whether evoked firing

changed as a result of time spent asleep or awake, we used the same "sandwich" approach we

used for the fEPSP data to compare the magnitude of evoked firing in flanking periods of wake

separated by an extended period of sleep, and vice versa for extended periods of wake (Fig. 8 C-

E). We averaged the response for each isolated unit across all sleep or wake dense epochs

(defined as >30 minute periods when the animal was >75% asleep/awake), and found no

significant differences in evoked spiking (Fig. 8E).

Next, we took advantage of the variation in epoch durations to look for a correlation between the

length of an epoch and the change in number of evoked spikes. We plotted the change in evoked

spikes for each unit for each epoch, and found no significant correlation between duration of a

sleep/wake epoch and ability of the thalamocortical stimulation to elicit spikes. Taken in concert

with the fEPSP results, this shows that time spent asleep or awake does not cause progressive

changes in the efficacy of thalamocortical synaptic transmission in vivo.

DISCUSSION

It has been postulated that patterns of brain activity during sleep induce a widespread

downscaling of excitatory synaptic strengths throughout CNS circuits, but the evidence for this

has been contradictory. Here we used three complementary measures of functional synaptic

strengths, coupled with real-time sleep classification in freely behaving animals, to ask whether

simply being asleep is able to constitutively drive widespread synaptic weakening. We found that

mEPSCs onto L4 or L2/3 pyramidal neurons were stable across sleep and wake dense epochs in

both V1 and PFC. Further, chronic monitoring of thalamocortical synaptic efficacy in V1 of

freely behaving animals revealed remarkable stability of thalamocortical transmission across

prolonged natural sleep and wake epochs. Together these data provide strong evidence that sleep

does not drive widespread constitutive weakening of excitatory synaptic strengths.

Several previous studies have reported that molecular or morphological correlates of synaptic
strength are lower after a period of sleep than a period of wake (Vyazovksiy et al., 2008; de Vivo
et al., 2017; Diering et al., 2017). There are several methodological differences between our
approach and previous studies that might explain some of these differences. First, we carefully
classified the sleep history of every animal used for our *ex vivo* recordings, which allowed us to
control for the considerable variability across individual animals in sleep habits. In the young
animals studied here (4-5 weeks old), the average chance of having experienced a sleep dense
epoch at ZT 4 or 6 (time points used in many studies) is < 50%; our approach allowed us to wait
until a sleep or wake dense epochs was detected within a specified circadian window for each
animal to ensure consistency in sleep history. Second, we carefully controlled for circadian time,
which has often been used as a proxy for sleep and wake (e.g. Lui et al., 2010, Diering et al.
2017), despite the potential impact of circadian time itself on synaptic strengths (discussed in
Frank and Cantera, 2014; Bridi et al., 2020). Third, we used direct electrophysiological
measurements of synaptic strength across largely unperturbed natural sleep and wake cycles,
whereas many previous studies used indirect measures of synaptic function as proxies for
synaptic strength (Diering et al., 2017; De Vivo et al., 2017), and/or relied on extended sleep
deprivation paradigms to manipulate the amount of sleep (Diering et al., 2017; De Vivo et al.,
2017, Vyazovksiy et al., 2008; Liu et al., 2010, Khlghatyan et al., 2020). Finally, there is wide
variation in brain areas and cell types examined across studies. Correlates of synaptic weakening
after sleep have been reported in motor cortex, somatosensory cortex, and PFC (Vyazovksiy et
al., 2008; Lui et al., 2010; de Vivo et al., 2017; Diering et al., 2017), while here we examined
synaptic strengths in visual cortex and PFC. It is possible that the impact of sleep on synaptic
function is distinct in different brain areas and cell types (discussed in Frank and Cantera, 2014).
Regardless, our data show that synaptic downscaling is not a universal function of natural sleep
across cortical cell types and brain regions.
397
Using a direct measure of postsynaptic strength, we found that mEPSC amplitude was stable
across sleep and wake epochs in three cell types from two very different cortical areas, V1 and
PFC. These findings are inconsistent with a previous study that reported that mEPSCs in frontal
cortex (including the PFC region we recorded from in this study) were larger and more frequent
after a period of sleep deprivation (Lui et al., 2010); a third recent study in PFC found increased

mEPSC amplitude but no change in frequency following sleep deprivation (Khlghatyan 2020).

While there are a number of possible explanations for these discrepancies between studies, one

major difference is that here we more carefully controlled for sleep history in individual animals,

and our study design allowed us to examine mEPSCs after relatively unmanipulated periods of

natural sleep and wake that ended in the same circadian window. A second major difference is

that we recorded from defined cell types and layers within well-defined brain regions, whereas

previous studies combined recordings from several distinct regions and did not report cell type

(Lui et al., 2010), or combined data from pyramidal neurons in several layers (Khlghatyan et al.,

2020); as mEPSC frequency and amplitude can vary considerably between cell types, these

measures could be strongly affected by sampling differences between conditions. By virtue of

the EEG recordings performed in our experiments (generally absent in these previous studies for

mEPSC datasets), we were also able to look for correlations in delta power drops and mEPSC

amplitude, and were unable to see a significant relationship. An important question is whether

mEPSC measurements are sensitive enough to detect changes in synaptic strength induced by 4

hours of consolidated sleep or wake. We have used a similar approach to successfully quantify

changes in mEPSC amplitude induced by visual deprivation and eye reopening, and can reliably

detect both increases and decreases of <10% with a similar sample size to the present study

(Lambo and Turrigiano, 2013; Hengen et al., 2016; Torrado Pacheco et al., 2020). Previously

reported changes in synaptic parameters following sleep or sleep deprivation are larger than that

(De Vivo et al., 2017; Liu et al., 2010; Vyazovskiy et al., 2008), suggesting that if they were

occurring at these synapses we would have been able to detect them. Taken together, our data

show that natural periods of sleep and wake do not by themselves globally modulate postsynaptic

strengths.

Whereas mEPSC amplitude strongly correlates with synaptic glutamate receptor accumulation

and postsynaptic strength (e.g. Ibata et al., 2008; Wierenga et al., 2005), mEPSC frequency is

influenced by many parameters and does not have a straightforward correlate with presynaptic

function. The frequency of mEPSCs can be impacted by changes in the number of functional

synaptic sites, but while some studies have found learning-dependent changes in spine turnover

during sleep (Li et al., 2017), sleep does not constitutively drive changes in spine density (de

Vivo et al., 2017). MEPSC frequency can also be modulated by factors that affect presynaptic

excitability, including presence of neuromodulators (Choy et al., 2018; Sharma &

Vijayaraghavan, 2003), and by vesicle refilling and pool size (Zhou et al., 2000; Liu & Tsien,

1995). Previous studies from PFC found inconsistent effects of sleep on mEPSC frequency; one

study found increased mEPSC frequency after sleep deprivation (Lui et al., 2010) while another

found no effect (Khlghatyan et al., 2020). While we found that mean mEPSC frequency was

unaffected by sleep or wake in any condition, there were small shifts in inter-event interval

distributions in some conditions. It is not clear if these effects are tied to circadian or sleep/wake

differences, and they were not consistent across brain regions. Taken together with previous

studies, our data show that sleep and wake do not drive widespread, consistent changes in

mEPSC frequency.

To complement our *ex vivo* mEPSC recordings, we also devised an approach that let us follow

synaptic strength at a defined set of synaptic inputs in real time during naturally occurring

periods of extended sleep or wake. We evoked thalamocortical fEPSPs at low frequency in Layer

4 of V1 while monitoring vigilance state, and found that – like mEPSC amplitude – these evoked

synaptic events were remarkably stable even across very long periods of sleep and wake. As for

changes in mEPSCs, previous studies have reported a wide range of effects of sleep on evoked

synaptic transmission. Sleep and/or slow wave activity has been variously found to potentiate

evoked responses (Chauvette et al., 2012) or bias evoked responses towards depression

(González-Rueda et al., 2018; Vyazovskiy et al., 2008), while a recent study using

optogenetically-evoked responses in mouse motor cortex found no impact of sleep deprivation

on evoked transmission (Matsumoto et al., 2020). Our approach (similar to Mastumoto et al.,

2020) has the advantage of sampling thalamocortical synaptic strengths continuously over many

iterations of nature sleep and wake cycles, and revealed no impact of time spent asleep or awake

on evoked synaptic strengths in V1. One concern with this technique is that repeated optogenetic

measurements from the same synapses might itself affect the synaptic responses, as high levels

of blue light can affect neuronal health and RNA expression (Tyssowski and Gray, 2019).

However, we used low light intensities and very low stimulation frequencies (1/20 to 1/40 Hz),

so that total exposure to blue light was ~8000 fold less per hour than levels shown to impact

function (Tyssowski and Gray, 2019); further, we were able to evoke consistent and stable

responses over several days, suggesting that our stimulation paradigm was not inducing

significant phototoxicity or otherwise compromising evoked transmission.

While time spent awake or asleep had no impact on evoked thalamocortical fEPSP responses,

these responses were rapidly modulated by transitions between brain states, as has been noted

previously (Matsumoto et al., 2020; Vyazovskiy et al., 2008; Reinhold et al., 2015; Hall and

Borbely, 1970). This is likely due to rapid changes in neuromodulatory tone (Lee and Dan,

2012), and/or changes in the network activity in the thalamus or V1 (Steriade, 2001; Steriade and

Timofeev, 2003) between sleep and wake states. The changes we observed in fEPSP amplitude

correlated with changes in the ability of thalamocortical activation to evoke spikes in V1

neurons. Interestingly, thalamocortical transmission was most effective at evoking spikes during

NREM sleep, with implications for information transfer during sleep states. We also found rapid

modulation of thalamocortical efficacy at transitions between the light and dark phase of the

12/12 L/D cycle; as there were no other obvious circadian oscillations in efficacy, this is likely a

rapid effect of changes in visual input. These data make clear that we can readily detect changes

in thalamocortical synaptic efficacy driven by neuromodulatory or sensory input, indicating that

the stability we observe during prolonged periods of sleep and wake is not due to a lack of

sensitivity of our measurements.

Another method researchers have used to probe for sleep/wake driven changes in excitability is

to measure spontaneous firing of neurons *in vivo* across sleep and wake states. While some

chronic long term recordings have revealed an oscillation in frontal cortex firing rates consistent

with changes in net excitatory drive (Vyazovskiy et al., 2009), other studies have either found

variable effects between regions and across neuronal populations (Miyawaki & Diba, 2016;

Miyawaki et al., 2019; Watson et al., 2016) or have observed stable firing rates absent a

plasticity induction paradigm (Hengen et al., 2016; Torrado Pacheco et al., 2020), suggesting that

the impact of sleep and wake on spontaneous firing is complex and brain-area dependent.

Spontaneous firing arises from many sources, so here we instead measured the ability of

thalamocortical inputs to V1 to evoke spikes. Consistent with our findings that mEPSC

amplitudes and thalamocortical fEPSPs are stable across extended periods of sleep and wake, we

found that the functional connection strength between thalamus and V1 was also stable (Fig. 8).

Thus, in this study we used three complementary approaches to measure synaptic efficacy, and

all three measures paint the same picture: that synaptic efficacy in V1 (as well as mEPSC

amplitude in PFC) is not constitutively modulated by time spent awake or asleep. We note that

this does not rule out a role for sleep in increasing or decreasing specific connections within

specific brain circuits during specific developmental windows (e.g. Chauvette et al., 2012,

Vyazovskiy et al., 2008), but taken together our data show that synaptic downscaling is not a

universal constitutive function of sleep.

While sleep is not sufficient to induce widespread changes in cortical synaptic strengths, there is

a large body of work showing that sleep and wake states can profoundly affect the induction of

various forms of plasticity when they are initiated by robust learning or sensory deprivation

paradigms. For example, sleep facilitates and is required for several forms of visual system

plasticity, and roles for NREM and REM sleep have been found in promoting growth,

maintenance, and loss of specific synaptic connections during learning (Frank et al., 2001; Aton

et al., 2014; Yang et al., 2014; Li et al., 2017; Durkin et al., 2017). Further, when firing in V1 is

perturbed using visual deprivation/eye reopening paradigms to induce homeostatic

compensation, upward firing rate homeostasis is confined to periods of active wake (Hengen et

al., 2016), while downward firing rate homeostasis is confined to periods of sleep (Torrado

Pacheco et al., 2020). Upward and downward firing rate homeostasis are driven in part through

homeostatic changes in synaptic strengths (Hengen et al., 2013; Torrado Pacheco et al., 2020),

indicating that while natural wake and sleep episodes do not drive constitutive changes in V1

excitatory synaptic strengths, they are able to gate the induction of upward and downward

homeostatic synaptic plasticity, respectively. Combined with this larger literature, our findings

are most consistent with the view that sleep does not have a single unified effect on synaptic

strengths, but rather that sleep and wake states are able to gate various forms of synaptic

plasticity when they are induced by salient experiences.

METHODS

Overview

All experimental measurements were performed on Long-Evans rats of both sexes between postnatal day P25 and 31. Litters were housed on a 12:12 light cycle with the dam and with free access to food and water. All animals were housed, cared for, surgerized, and sacrificed in accordance with Brandeis IBC and IACAUC protocols. Rat pups were weaned at ~P21. All animals received two days of post-operative care including daily injection of Meloxicam and Penicillin. During the few days before recording, animals were handled twice daily by the experimenter. The day before recording started animals were transferred to a clear plexiglass cage of dimensions 12"x12"x16" (length, depth, height) and separated into two arenas by a clear plastic divider with 1" holes, which safely allows for tactile and olfactory interaction with a littermate that was present for all recordings. The recording chamber contained toys and had walls with black/white edged patterns for enrichment. Animals were kept on a 12h/12h light/dark cycle in a temperature controlled room (21ºC, 25-55% humidity). Animals in the dark period wake dense condition (D. Wake) were on an inverted L-D cycle, in order to keep all other experimental variables (time of slicing and recording) the same for the experimenter. To do this, the whole littler with dam were transferred to a L-D chamber with inverted light schedule (lights off 7:30am-7:30pm) at least 2 weeks prior to surgery, and maintained on an inverted schedule while recording EEG/EMG.

Virus injections into dLGN

Viral injections of AAV-CAG-ChR2-mCherry were performed between P12-P16 using stereotaxic surgery under isoflurane anesthesia at 1.5-2.5%. Dorsal LGN was targeted bilaterally using stereotaxic coordinates after adjusting for the lambda-bregma distance for age. A glass micropipette pulled to a fine point delivered ~400 nL of virus-containing solution at the targeted depth. Targeting and expression was verified via posthoc histology.

Electrode implantation surgeries

Animals underwent electrode implantation at ages P22-P26. For animals to undergo acute slice physiology, we implanted 3 EEG screws (PlasticsOne, Roanoke, VA): frontal lobe (B: +1-2, L: 1.5-2.5 mm), parietal lobe (B: -3-4, L: 1.5-2.5 mm), and cerebellum (for reference). In addition,

two stainless steel spring electrodes were placed in the nuchal (neck) muscles of the rats for

EMG recording. Screw electrodes were fed into a pedestal (PlasticsOne) and everything was

secured with dental cement. For *in vivo* recording and stimulation experiments, a 16 channel

silicon probe (Neuronexus, Ann Arbor, MI) with adhered optic fiber (200-400 μm diameter;

Doric Lenses, Quebec City, Quebec; ThorLabs, Newton, New Jersey) were stereotaxically

inserted into V1 in dLGN virus-injected animals such that the fiber tip rested on the surface of

the brain. In the majority of these surgeries light stimulation and recording were performed

during the operation to verify correct placement of the optrode. The exposed craniotomy was

surrounded with a silicone elastomer (Kwik-Cast, World Precision Instruments, Sarasota, FL).

The array was secured using dental cement, then grounded to two screws (cerebellum and frontal

lobe) using steel wire and soldering paste. Screws were not used for EEG data collection in the

optrode animals as LFP was sufficient for behavioral classification. Animals were given 2-3 days

of recovery before data collection, at which point sleep/wake behavior had stabilized and was

comparable to unsurgerized animals.

Real time behavioral recording and classification

Animals with EEG/EMG head caps were connected to a flexible cable and low resistance

commutator (PlasticsOne). EEG signal was recorded as the difference between the parietal and

frontal screw, grounded to the cerebellum screw. The signal was amplified and filtered with EEG

and EMG amplifiers at x5000 and x500, respectively (BioPac, Goleta, CA). This analog signal

was then digitized with a NIDAQ board (National Instruments, Austin, TX). The data were

acquired, analyzed, and plotted all within a custom MATLAB program, which uses canonical

markers to classify behavioral state as either: NREM, REM, or wake. Briefly, the program

computes a delta/beta and a theta/delta ratio from the corresponding frequency bands in the EEG

(delta: 0.5-4 hz; theta: 5-8 hz; beta ~ 20-35 hz). Ratios were chosen as features because they

normalize the power band measurements between animals. Because delta is high during NREM,

and beta typically low (Uchida et al., 1992), taking the delta/beta ratio was logical and proved to

be a stable and informative feature for classifying NREM. Likewise, theta is high during REM,

and delta low, and therefore was indicative of REM sleep (Ratios used in Rempe et al., 2015).

EMG data is normalized to standard deviation and large deflections that cross a threshold (3-4

STDs) were extracted and considered as significant muscle activity. Animal pixel movement is

extracted from the video recording (infrared light allowed video monitoring in darkness) using

open-source software written in C++ (Video Blob Tracking, Open Source Instruments,

Watertown, MA, USA; Github: https://github.com/OSI-INC/VBT) and modified in-house to suit

our needs (adding video cropping, tracking in both light and dark conditions). The program uses

3 manually adjustable thresholds (delta, theta, and movement thresholds) for classification,

which are additionally adjusted given recent behavioral states. This moving threshold was

chosen because the transition probabilities between different states are far from uniform (e.g. 10

s of NREM is more likely to follow prior NREM than wake; Perez-Atencio et al., 2018). Using

known transition probabilities makes a scalar threshold more accurate. The classifier uses the

EEG ratios and movement measures and applies a decision tree which predicts states in 10

second epochs given the thresholds. The recent behavioral data is plotted and added to a graph

that contains the full behavioral history of the animal. This cycle repeats every 4 mins. Animals

were randomly assigned to either the sleep or wake dense conditions.

Post hoc semi-automated behavioral state scoring

In order to perform the final classification of behavior after experiments an in-house

MATLAB/Python GUI was used. The GUI initially uses the threshold-based algorithm used in

the real time classifier for the first several hours, which is manually reviewed for accuracy.

These data from each animal are then used as training input for the machine learning algorithm, a

200-300 tree random forest machine learning algorithm that uses the features described above

and additional features found to be useful for the decision tree. These include: the proportional

contribution of delta, theta, and gamma powers; min/max EEG; and a past value of many of

these features up to three 10 s bins ago. The random forest classifier was then applied and

manually reviewed for the rest of the recording. While the automated classification step occurred

in 10 s blocks, the subsequent manual review had a resolution of 1 s, ensuring our ability to

capture particularly short microarousals. Random forest accuracy was roughly 97%

benchmarked against manual coding.

In order to classify active vs. quiet wake, the same procedure was used except the training data

included a manual separation of these states (Fig. 2 – supplement 1). Manual differentiation of

active vs. quiet wake relied primarily on the video data (extracted movement measures and

manual review of video segments to observe animal movement by eye), but also included use of

gamma power. Animals were considered to be in active wake when LFP signals were especially

desynchronized (high gamma power, low delta), and were physically moving around the

cage/exploring. In contrast, quiet wake periods contained much less full-body displacement and

were mostly composed of sitting or grooming/ingesting.

NREM delta analysis

Delta power (0.5 Hz < Freq. < 4 Hz) was extracted from the slice behavioral data using FFT with

10 second windows. As delta power varies greatly between animals/recordings, we normalized it

to the value at the end of the light period where it tended to be lowest (ZT 7-11). To calculate the

change in delta power for each animal (Fig 3. – supplement 1), we found the percentage drop

from sleep epochs that occurred in the window ZT23.5-1.5 to the last hour of sleep right before

sacrificing animals.

Ex vivo **acute brain slice preparation**

For brain slice preparation, animals between P25 and P31 were briefly anesthetized with

isoflurane (usually under 60 seconds), and coronal brain slices (300 μm) containing V1 or

prefrontal cortex were obtained from both hemispheres of each animal. After slicing in

carbogenated (95% O_2, 5% CO_2) standard ACSF (in mM: 126 NaCl, 25 $NaHCO_3$, 3 KCl, 2

$CaCl_2$, 2 $MgSO_4$, 1 NaH_2PO_4, 0.5 Na-Ascorbate, osmolarity adjusted to 315 mOsm with

dextrose, pH 7.35), slices were immediately transferred to a warm (34°C) chamber filled with a

continuously carbogenated "protective recovery" (Ting et al., 2014) choline-based solution (in

mM: 110 Choline-Cl, 25 $NaHCO_3$, 11.6 Na-Ascorbate, 7 $MgCl_2$, 3.1 Na-Pyruvate, 2.5 KCl, 1.25

NaH_2PO_4, and 0.5 $CaCl_2$, osmolarity 315 mOsm, pH 7.35) for 10 min, then transferred back to

warm (34°C) carbogenated standard ACSF and incubated another 20-30 min. Slices were used

for electrophysiology between 1 – 6 hr. post-slicing.

Whole cell recordings

V1 and prefrontal cortex were identified in acute slices using the shape and morphology of the

white matter as a reference. Pyramidal neurons were visually targeted and identified by the

presence of an apical dendrite and teardrop shaped soma, and morphology was confirmed by

post hoc reconstruction of biocytin fills. Borosilicate glass recording pipettes were pulled using a

Sutter P-97 micropipette puller, with acceptable tip resistances ranging from 3 to 5 MΩ. Cs^+

Methanesulfonate-based internal recording solution was modified from Xue et al., 2014, and

contained (in mM) 115 Cs-Methanesulfonate, 10 HEPES, 10 BAPTA•4Cs, 5.37 Biocytin, 2 QX-

314 Cl, 1.5 $MgCl_2$, 1 EGTA, 10 Na_2-Phosphocreatine, 4 ATP-Mg, and 0.3 GTP-Na, with sucrose

added to bring osmolarity to 295 mOsm, and CsOH added to bring pH to 7.35. Inclusion criteria

included V_m, R_{in}, and R_s cut-offs as appropriate for experiment type and internal solution; values

are listed below.

All recordings were performed on submerged slices, continuously perfused with carbogenated

34°C recording solution. Neurons were visualized on an Olympus upright epifluorescence

microscope using a 10x air (0.13 numerical aperture) and 40x water-immersion objective (0.8

numerical aperture) with infrared-differential interference contrast optics and an infrared CCD

camera. Data were low-pass filtered at 5 kHz and acquired at 10 kHz with Axopatch 200B

amplifiers and CV-7B headstages (Molecular Devices, Sunnyvale, CA). Data were acquired

using an in-house program written either in Igor Pro (Wavemetrics, Lake Oswego, OR) or

MATLAB (Mathworks, Natick, MA), and all post-hoc data analysis was performed using in-

house scripts written in MATLAB.

mEPSC recordings

For spontaneous mEPSC recordings, pyramidal neurons were voltage clamped to −70 mV in

standard ACSF containing a drug cocktail of TTX (0.2 μM), APV (50 μM), picrotoxin (25 μM).

10 s traces were obtained and amplified (10-20x). Event inclusion criteria included amplitudes >

5 pA and rise times < 3 ms. Neurons were excluded from analysis if R_s > 25 MΩ or V_m > −50

mV.

mEPSC analysis

To reliably detect mEPSC events above noise and limit bias in selection, we used an in-house

program written in MATLAB that employs a semi-automated template-based detection method

contained in a GUI (Torrado Pacheco et al., 2020). In brief, the program first filters the raw

current traces and then applies a canonical mEPSC event shaped template to detect regions of

best fit. Multiple tunable parameters for template threshold and event kinetics that aid in

detection were optimized and then chosen to stay constant for all analyses. Briefly, sample traces

were chosen, which had all of the mEPSC events manually annotated. This training dataset was

used while we sampled much of the parameter space for template-matching parameters. These

parameters include the mEPSC shape used for template, how the error around the template was

calculated, the threshold cutoff, and a coefficient that balances detection of larger mini events

versus smaller ones to correct for larger error in larger events. After identification, putative

events were then examined for hallmark features of mEPSC (rise times, decay kinetics,

conforming to amplitude cutoff, not saturated) and some detected events (~4%) were excluded if

they did not conform to these features. Finally, the automatic detection missed ~6% of events

and these were manually added. Results did not differ substantially when the data were not

manually corrected.

Array recording and stimulation

The arrays were connected to an Intan RHD2216 amplifier board, which in turn was connected

to a RHD2000 SPI interface cable (Intan, Los Angeles, California). The cable connected to a

custom designed Hall effect-based active commutator (NeuroTek Innovative Technology

Incorporated, Toronto, Ontario, Canada). The commutator fed this digitized data into the

RHD2000 USB board (Intan). Data were recorded at 25 kHz continuously for up to 100 hours

using the RHD2000 Interface Software (Intan). Spike extraction, clustering, and sorting were

done using custom Matlab and Python code (see below). Light for stimulation was produced

using a fiber coupled 470 nm LED system (M470F3/LEDD1B, Thorlabs). Light irradiance was

adjusted for each animal and never exceeded 18 mW/cm^2. Stimulation was triggered using

voltage pulses of varying magnitude from a NIDAQ board (controlled by custom Matlab

program) to the LED driver, set to modulation mode.

Semi-automated spike extraction, clustering, and sorting

Spike extraction, clustering, and sorting was performed as previously described (Hengen et al.,

2016; Torrado Pacheco et al., 2020). Spikes were detected in the raw signal as threshold

crossings (-4 standard deviations from mean signal) and re-sampled at 3x the original rate.

Principal component analysis (PCA) was done on all spikes from each channel, and the first four

principal components were used for clustering (Harris et al., 2000). A random forest classifier

implemented in python was used to classify spikes according to a model built on a dataset of

1200 clusters manually scored by expert observers. A set of 19 features, including ISI

contamination (% of ISIs < 3 ms), similarity to regular spiking unit (RSU) and fast-spiking unit

(FS) waveform templates, amount of 60 Hz noise contamination and kinetics of the mean

waveform. Cluster quality was also ensured by thresholding of L-ratio and isolation distance

(Schmitzer-Torbert et al., 2005). Clusters were classified as noise, multi-unit or single-unit. Only

single-unit clusters with a clear refractory period were used for FR analysis. We classified units

as RSU or FS based on established criteria (mean waveform trough-to-peak and tail slope,

Hengen et al., 2016). Only RSUs (putative excitatory neurons) were used for analysis. Some

units were lost during the recording, presumably due to electrode drift or gliosis. To establish

periods of time when units could be clearly detected and isolated, we used ISI contamination:

when hourly % of ISIs < 2 msec was above 4%, the unit was considered to be poorly isolated;

neurons were included for analysis for each behavioral epoch in which unit isolation quality met

criterion. Using these criteria, 93% of cells were well isolated for >1/3 of the full experiment

time, and 80% of cells were well isolated for >1/2 of full experiment time of 2-3 days. Results of

the automated scoring were manually reviewed for final inclusion of neurons.

FEPSP analysis

For fEPSP analysis, the data were bandpass filtered (high-pass = 5 Hz, low-pass = 450, filter

order = 2). For each optogenetic stimulation, a section of data was extracted (100 ms prior to

stimulation and 80 ms after) for each channel in the array. While all the channels were used for

estimation of cortical layer placement, only channels in L4 to upper L5 where the response was

largest and shortest-latency, and channels where signal-to-noise was stable, were chosen for

fEPSP analysis. Evoked events were rejected if the standard deviation was greater than 200 µV

in the last 5 ms before stimulation or if an evoked trough of >30 µV magnitude could not be

found. Slopes were calculated for the rise of the waveform between 20 to 80% of the maximum

amplitude. For the sandwich analysis (Fig. 7), flanking periods of 1 hour before and after a given

epoch were compared. Normalized change in amplitude was calculated as: post/pre – 1. For time

course analysis (Fig. 6), fEPSP amplitudes were normalized to the first hour of a given epoch.

For analysis of the rare epochs that spanned L-D transitions we normalized the amplitude just

after the transition to the amplitude just before the transition to account for L-D mediated

changes in amplitude; results were not different if instead these epochs were excluded or were

used without this normalization.

Evoked firing rate analysis

For the evoked spike analysis (Fig. 8), spike responses were determined for each RSU for each

stimulation. Spike responses were measured as the number of spikes that occurred in the 20 ms

interval post stimulation above what was expected given the previous baseline firing rate. A

baseline firing rate was determined in the 60 seconds prior to each stimulation for each cell, and

the expected number of spikes given the baseline firing rate were subtracted from the actual

spike number that occurred in the 20 ms post interval: Response = Spike Count - (baseline FR *

0.02 seconds). To determine whether evoked spikes were above chance levels for a given cell,

we performed a bootstrap analysis by randomly sampling times where no stimulations occurred,

and performed the same analysis. This produced a null distribution of "responses" to no

stimulation. Cells that had responses <95th percentile of randomly sampled responses were

considered unresponsive to stimulation and excluded from analysis.

Statistical analysis

All data analysis was performed using in-house scripts written in MATLAB. For each

experiment, means ± SEM derived from individual cell measurements are provided within the

results section of the text, and n's (number of cells and animals), P values, and statistical tests are

provided within figure captions. Generally, a Kruskal-Wallis test was performed on 2-3

condition experiments in Figure 3 as the distributions did not pass normality tests. Two-sample

Kolmogorov-Smirnov test with a Bonferroni correction for multiple comparisons was used for

comparisons between distributions. Wilcoxon sign rank test was used to compare distributions

that were not normally distributed in Fig 7 and 8. Fig 8E's data was log normal and so a 2-

sample t-test after log transformation was used.

For slice experiments where expected means and variance were known, target sample sizes were

determined beforehand using power analyses, using a detection threshold of ~10%. For our *in*

vivo experiments this information was not known so an *a priori* power analysis was not possible.

A post-hoc calculation based on sample size and variance shows that changes of >10% would

have been detectable.

Code and data Availability

776 All analysis code is available here https://github.com/BrianAndCary/papers.

ACKNOWLEDGEMENTS

779 We thank Alejandro Torrado Pacheco for his invaluable help with the surgery and

780 implementation of *in vivo* recording and expertise in handling extracellular data and the

781 programming involved in behavioral analysis. We thank Raul Ramos for his technical expertise

782 and aid in imaging of brain slices. Supported by NIH EY025613 (GGT).

COMPETING INTERESTS

785 No competing interests declared.

REFERENCES

Abbott, L. F., & Nelson, S. B. (2000). Synaptic plasticity: taming the beast. *Nature Neuroscience*, *3*(november), 1178–1183. https://doi.org/10.1038/81453

Aton, S. J., Suresh, A., Broussard, C., & Frank, M. G. (2014). Sleep promotes cortical response potentiation following visual experience. *Sleep*, *37*(7), 1163–1170. https://doi.org/10.5665/sleep.3830

Brown, R. E., Basheer, R., McKenna, J. T., Strecker, R. E., & McCarley, R. W. (2012). Control of Sleep and Wakefulness. *Physiological Reviews*, *92*(3). https://doi.org/10.1152/physrev.00032.2011

Chauvette, S., Seigneur, J., & Timofeev, I. (2012). Sleep Oscillations in the Thalamocortical System Induce Long-Term Neuronal Plasticity. *Neuron*, *75*(6), 1105–1113. https://doi.org/10.1016/j.neuron.2012.08.034

Cooke, S. F., & Bear, M. F. (2010). Visual experience induces long-term potentiation in the primary visual cortex. *J Neurosci*, *30*(48), 16304–16313. https://doi.org/10.1523/JNEUROSCI.4333-10.2010

Datta, S., Li, G., & Auerbach, S. (2008). Activation of phasic pontine-wave generator in the rat: A mechanism for expression of plasticity-related genes and proteins in the dorsal hippocampus and amygdala. *European Journal of Neuroscience*, *27*(7), 1876–1892. https://doi.org/10.1111/j.1460-9568.2008.06166.x

de Vivo, L., Bellesi, M., Marshall, W., Bushong, E. A., Ellisman, M. H., Tononi, G., & Cirelli, C. (2017). Ultrastructural evidence for synaptic scaling across the wake/sleep cycle. *Science*, *355*(6324), 507–510. https://doi.org/10.1126/science.aah5982

Diering, G. H., Nirujogi, R. S., Roth, R. H., Worley, P. F., Pandey, A., & Huganir, R. L. (2017). Homer1a drives homeostatic scaling-down of excitatory synapses during sleep. *Science*, *515*(February), 511–515.

Dijk, D. J. (2009). Regulation and functional correlates of slow wave sleep. *Journal of Clinical Sleep Medicine*, *5*(2 SUPPL.), 6. https://doi.org/10.5664/jcsm.5.2s.s6

Durkin, J., Suresh, A. K., Colbath, J., Broussard, C., Wu, J., Zochowski, M., & Aton, S. J. (2017). Cortically coordinated NREM thalamocortical oscillations play an essential, instructive role in visual system plasticity. *Proceedings of the National Academy of Sciences of the United States of America*, *114*(39), 10485–10490. https://doi.org/10.1073/pnas.1710613114

Espinosa, J. S., & Stryker, M. P. (2012, July 26). Development and Plasticity of the Primary Visual Cortex. *Neuron*, Vol. 75, pp. 230–249. https://doi.org/10.1016/j.neuron.2012.06.009

Frank, M. G., & Cantera, R. (2014). Sleep, clocks, and synaptic plasticity. *Trends in Neurosciences*, *37*(9), 491–501. https://doi.org/10.1016/j.tins.2014.06.005

Frank, M. G., Issa, N. P., & Stryker, M. P. (2001). Sleep enhances plasticity in the developing visual cortex. *Neuron*, *30*(1), 275–287. https://doi.org/10.1016/S0896-6273(01)00279-3

Frank, M. G., Ruby, N. F., Heller, H. C., & Franken, P. (2017). Development of circadian sleep regulation in the rat: A longitudinal study under constant conditions. *Sleep*, *40*(3). https://doi.org/10.1093/sleep/zsw077

Hall, R. D., & Borbely, A. A. (1970). Acoustically evoked potentials in the rat during sleep and waking. *Experimental Brain Research*, *11*(1), 93–110. https://doi.org/10.1007/BF00234203

Han, E. B., & Stevens, C. F. (2009). Development regulates a switch between postand presynaptic strengthening in response to activity deprivation. *Proceedings of the National Academy of Sciences of the United States of America*, *106*(26), 10817–10822. https://doi.org/10.1073/pnas.0903603106

Hengen, K. B., Lambo, M. E., VanHooser, S. D., Katz, D. B., & Turrigiano, G. G. (2013). Firing rate homeostasis in visual cortex of freely behaving rodents. *Neuron*, *80*(2), 335–342. https://doi.org/10.1016/j.neuron.2013.08.038

Hengen, K. B., Torrado Pacheco, A., McGregor, J. N., Van Hooser, S. D., & Turrigiano, G. G. (2016). Neuronal Firing Rate Homeostasis Is Inhibited by Sleep and Promoted by Wake. *Cell*, *165*(1), 1–12. https://doi.org/10.1016/j.cell.2016.01.046

Heynen, A. J., Yoon, B. J., Liu, C. H., Chung, H. J., Huganir, R. L., & Bear, M. F. (2003). Molecular mechanism for loss of visual cortical responsiveness following brief monocular deprivation. *Nature Neuroscience*, *6*(8), 854–862. https://doi.org/10.1038/nn1100

Jones, K., & Harrison, Y. (2001, December 1). Frontal lobe function, sleep loss and fragmented sleep. *Sleep Medicine Reviews*, Vol. 5, pp. 463–475. https://doi.org/10.1053/smrv.2001.0203

Kavanau, J. L. (1996). Memory, sleep, and dynamic stabilization of neural circuitry: Evolutionary perspectives. *Neuroscience and Biobehavioral Reviews*, *20*(2), 289–311. https://doi.org/10.1016/0149-7634(95)00019-4

Khurana, S., & Li, W. (2013). Baptisms of fire or death knells for acute-slice physiology in the age of "omics" and light? *Rev. Neurosci.*, *24*(5).

Krause, A. J., Simon, E. Ben, Mander, B. A., Greer, S. M., Saletin, J. M., Goldstein-Piekarski, A. N., & Walker, M. P. (2017, July 1). The sleep-deprived human brain. *Nature Reviews Neuroscience*, Vol. 18, pp. 404–418. https://doi.org/10.1038/nrn.2017.55

Lambo, M. E., & Turrigiano, G. G. (2013). Synaptic and intrinsic homeostatic mechanisms cooperate to increase L2/3 pyramidal neuron excitability during a late phase of critical period plasticity. *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience*, *33*(20), 8810–8819. https://doi.org/10.1523/JNEUROSCI.4502-12.2013

Lee, S. H., & Dan, Y. (2012). Neuromodulation of Brain States. *Neuron*, *76*(1), 109–222. https://doi.org/10.1016/j.neuron.2012.09.012

Leemburg, S., Vyazovskiy, V. V., Olcese, U., Bassetti, C. L., Tononi, G., & Cirelli, C. (2010). Sleep homeostasis in the rat is preserved during chronic sleep restriction. *Proceedings of the National Academy of Sciences of the United States of America*, *107*(36), 15939–15944. https://doi.org/10.1073/pnas.1002570107

Li, W., Ma, L., Yang, G., & Gan, W.-B. (2017). REM sleep selectively prunes and maintains new synapses in development and learning. *Nature Neuroscience*, *20*(3). https://doi.org/10.1038/nn.4479

Liu, Z.-W., Faraguna, U., Cirelli, C., Tononi, G., & Gao, X.-B. (2010). Direct Evidence for Wake-Related Increases and Sleep-Related Decreases in Synaptic Strength in Rodent Cortex. *Journal of Neuroscience*, *30*(25), 8671–8675. https://doi.org/10.1523/JNEUROSCI.1409-10.2010

Matsumoto, S., Ohyama, K., Díaz, J., Yanagisawa, M., Greene, R. W., & Vogt, K. E. (2020). Enhanced cortical responsiveness during natural sleep in freely behaving mice. *Scientific Reports*, *10*

Miller, K. D., & MacKay, D. J. C. (1994). The Role of Constraints in Hebbian Learning. *Neural Computation*, *6*(1), 100–126. https://doi.org/10.1162/neco.1994.6.1.100

Miska, N. J., Richter, L. M. A., Cary, B. A., Gjorgjieva, J., & Turrigiano, G. G. (2018). Sensory experience inversely regulates feedforward and feedback excitation-inhibition ratio in rodent visual cortex. *ELife*, *7*. https://doi.org/10.7554/eLife.38846

Miyawaki, H., & Diba, K. (2016). Regulation of Hippocampal Firing by Network Oscillations during Sleep. *Current Biology*, *26*(7), 893–902. https://doi.org/10.1016/j.cub.2016.02.024

Miyawaki, H., Watson, B. O., & Diba, K. (2019). Neuronal firing rates diverge during REM and homogenize during non-REM. *Scientific Reports*, *9*(1). https://doi.org/10.1038/s41598-018-36710-8

Muzur, A., Pace-Schott, E. F., & Hobson, J. A. (2002, November 1). The prefrontal cortex in sleep. *Trends in Cognitive Sciences*, Vol. 6, pp. 475–481. https://doi.org/10.1016/S1364-6613(02)01992-7

Nataraj, K., Le Roux, N., Nahmani, M., Lefort, S., & Turrigiano, G. (2010). Visual deprivation suppresses L5 Pyramidal Neuron Excitability by Preventing the Induction of Intrinsic Plasticity. *Neuron*, *68*(4), 750–762. https://doi.org/10.1016/j.neuron.2010.09.033

Nath, R. D., Bedbrook, C. N., Abrams, M. J., Basinger, T., Bois, J. S., Prober, D. A., … Goentoro, L. (2017). The Jellyfish Cassiopea Exhibits a Sleep-like State. *Current Biology*, *27*(19), 2984-2990.e3. https://doi.org/10.1016/j.cub.2017.08.014

Niell, C. M., & Stryker, M. P. (2008). Highly selective receptive fields in mouse visual cortex. *Journal of Neuroscience*, *28*(30), 7520–7536. https://doi.org/10.1523/JNEUROSCI.0623-08.2008

Perez-Atencio, L., Garcia-Aracil, N., Fernandez, E., Barrio, L. C., & Barios, J. A. (2018). A four-state Markov model of sleep-wakefulness dynamics along light/dark cycle in mice. *PLOS ONE*, *13*(1), e0189931. https://doi.org/10.1371/journal.pone.0189931

Reinhold, K., Lien, A. D., & Scanziani, M. (2015). Distinct recurrent versus afferent dynamics in cortical visual processing. *Nature Neuroscience*, *18*(12), 1789–1797. https://doi.org/10.1038/nn.4153

Rempe, M. J., Clegern, W. C., & Wisor, J. P. (2015). An automated sleep-state classification algorithm for quantifying sleep timing and sleep-dependent dynamics of electroencephalographic and cerebral metabolic parameters. *Nature and Science of Sleep*, *7*, 85–99. https://doi.org/10.2147/NSS.S84548

Shein-Idelson, M., Ondracek, J. M., Liaw, H.-P., Reiter, S., & Laurent, G. (2016). Slow waves, sharp waves, ripples, and REM in sleeping dragons. *Science*, *352*(6285), 590–595. https://doi.org/10.1126/science.aaf3621

Smith, G. B., & Bear, M. F. (2010). Bidirectional ocular dominance plasticity of inhibitory networks: Recent advances and unresolved questions. *Frontiers in Cellular Neuroscience*, *4*(JUN). https://doi.org/10.3389/fncel.2010.00021

Steriade, M. (2001). Impact of network activities on neuronal properties in corticothalamic systems. *Journal of Neurophysiology*, *86*(1), 1–39.

Steriade, M., & Timofeev, I. (2003). Neuronal plasticity in thalamocortical networks during sleep and waking oscillations. *Neuron*, *37*, 563–576.

Ting, J. T., Daigle, T. L., Chen, Q., & Feng, G. (2014). Acute brain slice methods for adult and aging animals: application of targeted patch clamp analysis and optogenetics. *Methods in Molecular Biology (Clifton, N.J.)*, *1183*, 221. https://doi.org/10.1007/978-1-4939-1096-0_14

Tononi, G. (2009). Slow wave homeostasis and synaptic plasticity. *Journal of Clinical Sleep Medicine*, *5*(2 SUPPL.), S16. https://doi.org/10.5664/jcsm.5.2s.s16

Tononi, G., & Cirelli, C. (2014). Sleep and the Price of Plasticity: From Synaptic and Cellular Homeostasis to Memory Consolidation and Integration. *Neuron*, *81*(1), 12–34. https://doi.org/10.1016/j.neuron.2013.12.025

Torrado Pacheco, A., Bottorff, J., & Turrigiano, G. G. (2019). Sleep promotes downward firing rate homeostasis. *BioRxiv*, 827832. https://doi.org/10.1101/827832

Torrado Pacheco, A., Tilden, E. I., Grutzner, S. M., Lane, B. J., Wu, Y., Hengen, K. B., … Turrigiano, G. G. (2019). Rapid and active stabilization of visual cortical firing rates across light-dark transitions. *Proceedings of the National Academy of Sciences of the United States of America*, *116*(36), 18068–18077. https://doi.org/10.1073/pnas.1906595116

Turrigiano, G. G. (2008, October 31). The Self-Tuning Neuron: Synaptic Scaling of Excitatory Synapses. *Cell*, Vol. 135, pp. 422–435. https://doi.org/10.1016/j.cell.2008.10.008

Turrigiano, G. G., & Nelson, S. B. (2004). Homeostatic plasticity in the developing nervous system. *Nature Reviews. Neuroscience*, *5*(2), 97–107. https://doi.org/10.1038/nrn1327

Tyssowski, K. M., & Gray, J. M. (2019). Blue light induces neuronal-activity-regulated gene expression in the absence of optogenetic proteins. *BioRxiv*, 572370. https://doi.org/10.1101/572370

Uchida, S., Maloney, T., & Feinberg, I. (1992). Beta (20-28 Hz) and Delta (0.3-3 Hz) EEGs Oscillate Reciprocally Across l~REM and REM Sleep. In *Sleep* (Vol. 15). Retrieved from https://academic.oup.com/sleep/article/15/4/352/2749299

Vyazovskiy, V. V., Olcese, U., Lazimy, Y. M., Faraguna, U., Esser, S. K., Williams, J. C., … Tononi, G. (2009). Cortical Firing and Sleep Homeostasis. *Neuron*, *63*(6), 865–878. https://doi.org/10.1016/j.neuron.2009.08.024

Vyazovskiy, V. V, Cirelli, C., Pfister-Genskow, M., Faraguna, U., & Tononi, G. (2008). Molecular and electrophysiological evidence for net synaptic potentiation in wake and depression in sleep. *Nature Neuroscience*, *11*(2), 200–208. https://doi.org/10.1038/nn2035

Walker, M. P., & Stickgold, R. (2004, September 30). Sleep-dependent learning and memory consolidation. *Neuron*, Vol. 44, pp. 121–133. https://doi.org/10.1016/j.neuron.2004.08.031

Watson, B. O., Levenstein, D., Greene, J. P., Gelinas, J. N., & Buzsáki, G. (2016). Network Homeostasis and State Dynamics of Neocortical Sleep. *Neuron*, *90*(4), 839–852. https://doi.org/10.1016/j.neuron.2016.03.036

Wierenga, C. J., Walsh, M. F., & Turrigiano, G. G. (2006). Temporal regulation of the expression locus of homeostatic plasticity. *Journal of Neurophysiology*, *96*(4), 2127–2133. https://doi.org/10.1152/jn.00107.2006

Xue, M., Atallah, B. V., & Scanziani, M. (2014). Equalizing excitation–inhibition ratios across visual cortical neurons. *Nature*, *511*(7511), 596–600. https://doi.org/10.1038/nature13321

Yang, G., Sau, C., Lai, W., Cichon, J., Ma, L., Li, W., & Gan, W. (2014). Sleep promotes branch-specific formation of dendritic spines after learning. *Science*, *344*(6188).

FIGURES

Fig 1: Characterization of sleep/wake behavior in juvenile Long-Evans rats. (**A**) Two example sleep/wake histories from two different animals. Y axis is a moving mean showing percent of time spent awake in the previous 4 hours; x axis is zeitgeber time (ZT) in hours. Periods where the animal is in a sleep dense epoch (>65% time asleep in previous 4 hrs) are colored blue, while >65% wake in previous 4 hrs are colored green. Red arrows indicate points showing differing sleep/wake dense experiences between animals at the same ZT. (**B**) Average percentage time spent awake in the previous 4 hours as a function of ZT time. Light green shading indicates the standard deviation between animals; n = 30 animals. (**C**) Normalized delta power from the EEG of animals in B as a percent of baseline (hours 7-11 in the light period). Shading represents SEM between animals. (**D**) Average time spent in each behavioral state shown as a proportion for each animal broken down by light period (left) and dark period (right); each point represents one animal. Mean and SEM shown, n = 30 animals.

Fig 2: Real time behavioral classification. (**A**) 1 hour of behavioral data illustrating variables used for classification of sleep/wake state, described in descending order. Top shows expanded raw EEG traces from the indicated periods; scale bar next to NREM example, 1 V, 0.5 s. Below expanded EEG is the sleep/wake classification expressed as a hypnogram, where colored bars indicate periods of wake (green), NREM (blue), or REM (magenta). Beneath the hypnogram is the full raw EEG trace; scale is 2 V. Spectral Frequency (Spect. Freq.) plots the full spectrogram of the EEG from 0-20 Hz, colored by power (normalized to maximum). Extracted features of the Spect. Freq. are plotted below: delta(0.5-4 Hz)/beta(20-35 Hz) power ratio (shown on log scale) is high during periods of NREM, while the theta(5-8 Hz)/delta(0.5-4 Hz) power ratio is high during periods of REM. Absolute EMG values were normalized and expressed as standard deviation. Finally, animal movement in pixels is plotted. The bottom 4 plots of vigilance state features each have an adjustable threshold for state classification, shown as dashed line; points above the threshold are shown as colored dots. (**B**) Schematic of real time classifier rig. (**C**) 3D plot of vigilance state features: delta ratio, theta ratio, and movement measures (zero movement assigned lowest observed value for log axis) colored by brain state. Data from example animal. (**D**) Accuracy of real time classifier compared to manual scoring by state: Overall (% time matching between all 3 states) = 93.3%; Wake = 94.2%; NREM = 93.2%; REM = 90.5%; Wake Dense (matching wake dense on and off times) = 96.0%; Sleep Dense = 96.0%.

Fig 2 – figure supplement 1: **Discrimination of active and quiet wake.** (**A**) Top, hypnogram indicating periods of active (dark green) or quiet (light green) wake. Some states labeled with specific activities the animal was performing, as determined by video recording. Top line plot is the gamma power of the LFP; which is higher during more active wake behaviors. Bottom line plot is the animal's movement in pixels; higher movement is associated with active wake. (**B**) Cumulative plot of the animals' movement (same animal from A) for active and quiet wake states.

Fig 3: Stability of mEPSC amplitude after prolonged periods of sleep or wake. The left column represents example sleep histories for the VC L2/3 experiments while right column shows mEPSC characteristics for the indicated cell types and brain regions. (**A**) Top figure, hypnograms for three animals used for sleep dense mEPSC recordings. Green bars indicate instances of wake, magenta indicates REM, and blue NREM. Bottom, this same behavioral state data represented as a moving average of % time spent awake in the preceding 4 hr; the dashed lines represent the thresholds for achieving the required density in a given state, and the dots represent endpoints for slice recordings. (**B**) and (**C**) are same as for (A), but for wake dense recordings performed during the light and dark periods, respectively. Red arrow indicates wake encouragement (introduction of toy). Black circle around end point dot indicates which trace is represented by example hypnogram. (**D**) Traces on the left, example mEPSC recordings from L2/3 of V1; scale bar, 10 pA, 1 s. On the right, cell average mEPSC events; scale bar, 5 pA, 5 ms. (**E**) On left, cell average mEPSC amplitudes from L2/3 V1; error bars = 95% CI of mean (p=0.17; Kruskal-Wallis). (**E**) On right, cumulative histogram of sampled mEPSC amplitudes (SD – L. WD p=0.075, L. WD – D. WD p>0.9, SD – D. WD p=0.77; KS test with Bonferroni correction). Inset for (E) – (G) is a peak scaled average mEPSC waveform; scale bar, 50% peak, 5 ms. Sleep dense n = 17 cells, 3 animals; L. wake dense n = 25 cells, 4 animals; D. wake dense n = 18 cells, 5 animals. (**F**) On left, cell average mEPSC from L4 V1 (p=0.52; Kruskal-Wallis). (**F**) On right, cumulative mEPSC amp. from L4 V1 (p=0.99; KS test). Sleep dense n = 47 cells, 6 animals; L. wake dense n = 32 cells, 4 animals. (**G**) On left, cell average mEPSC amplitudes from L2/3 PFC (p=0.13; Kruskal-Wallis). (**G**) On right, cumulative histogram of sampled mEPSC amplitudes (SD – L. WD p>0.9, L. WD – D. WD p=0.096, SD – D. WD p=0.015; KS test with Bonferroni correction). Sleep dense n = 31 cells, 6 animals; L. wake dense n = 28 cells, 5 animals; D. wake dense n = 18 cells, 6 animals.

Fig 3 – figure supplement 1: Drop in delta power and time spent awake do not correlate with mEPSC amplitudes or frequency. (**A**) Schematic showing how drop in delta power is calculated. Delta drop was calculated as percentage drop from sleep epochs that occurred in the window ZT23.5-1.5 to the last hour of sleep right before sacrificing animals. Peak delta power after extended wake at t1 is normalized to the baseline recovery period of t2. (**B**) mEPSC amplitude after a sleep dense epoch, plotted against normalized drop in delta power during that epoch. Black line = linear fit; R^2 value and p value included on plot. Cell averages from different regions are indicated in legend. (**C**) Same as B) but for mEPSC frequency. (**D**) Schematic illustrating time spent awake in preceding 4 hours of a wake dense epoch. (**E**) mEPSC amplitude after a wake dense epoch, plotted against proportion of time spent awake in previous 4 hours. Cell averages from different regions are indicated in legend; light period recordings in green, dark period recordings in Red. (**F**) Same as (**E**) but for mEPSC frequency.

Fig 3 – figure supplement 2: Sleep/wake histories for additional *ex vivo* slice recordings. Each experiment includes an example hypnogram from one animal above summary data of wake history. (**A**) Animals from visual cortex L4 recordings. Top figure, example hypnogram from a single animal. Green bars indicate instances of wake, magenta indicates REM, and blue NREM. Below, this same behavioral state data is represented as a moving average of % time spent awake in the preceding 4 hr but for each animal included in the experiment; the dashed lines represent the thresholds for achieving the required density in a given state, and the dots represent endpoints for slice recordings. Black circle around end point dot indicates which trace is represented by example hypnogram. (**B**) Same as (A) but for wake dense recordings (with green dots). (**C-D**) Same as (A-B) but for prefrontal cortex L2/3 recordings. (**E**) Same as (D) but for the D. wake dense recordings (i.e. sacrifice occurred in the dark period).

Fig 4: Impact of prolonged sleep/wake on mEPSC frequency. MEPSC frequency is not consistently modulated by sleep/wake history across brain regions. (**A**) Data from visual cortex L2/3. Left plot, cell average mEPSC frequencies for the indicated conditions; error bars = 95% CI of mean (p=0.75, Kruskal Wallace). Right plot, cumulative plot of inter event interval (SD - L. WD p=0.35, SD - D. WD p=0.15, L. WD - D. WD p>0.9; KS test with Bonferroni correction). Sleep dense n = 17 cells, 3 animals; L. wake dense n = 25 cells, 4 animals; D. wake dense n = 18 cells, 5 animals. (**B**) Same as in (A) but data from visual cortex L4; cell average on left (p>0.9, Kruskal Wallace); cumulative plot on right (p=0.45, KS test). Sleep dense n = 47 cells, 6 animals; L. wake dense n = 32 cells, 4 animals. (**C**) Same as in (A) but data from prefrontal cortex L2/3. Left, cell average values for the indicated conditions (p=0.056, Kruskal Wallace). Cumulative plot of inter-event interval (SD - L. WD p<1e-6, SD - D. WD p=0.74, L. WD - D. WD p=0.007; KS test with Bonferroni correction). Sleep dense n = 31 cells, 6 animals; L. wake dense n = 28 cells, 5 animals; D. wake dense n = 18 cells, 6 animals.

Fig. 5: Thalamocortical evoked synaptic responses in freely behaving animals. (**A**) AAV-Chr2-mCherry viral expression targeted to dLGN projects to V1. Left, coronal slice showing viral expression in magenta. Right, visual cortex showing neuronal nuclei marked with NeuN and ChR2-mCherry in magenta. (**B**) Evoked fEPSPs recorded with a linear silicon array show clear layer-specific waveforms consistently with TC afferent location. (**C**) Current source density heat plot identifies layers; cool colors indicate current sink. (**D**) FEPSP latency to first negative peak identifies layers. Red circles represent latency of individual events; black line represents mean for each channel. (**E**) FEPSPs recorded over 3 days in a single freely behaving animal. Amplitude of the response plotted and colored by brain state. Dark periods (lights off) are represented by grey rectangles. (**F**) fEPSP waveforms by brain state, where each waveform is from a different animal; normalized to QW peak (indicated by dashed line) for each animal. Scale bar, 50% of peak amplitude, 10 ms. (**G**) FEPSP amplitudes across circadian time, normalized to quiet wake in the light period. N = 4 animals.

Fig 6: Stability of thalamocortical fEPSP amplitude during prolonged periods of sleep or wake. Time series data of thalamocortical evoked fEPSPs over the course of either sleep or wake dense epochs. (**A**) Example hypnograms showing wake (left) and sleep (right) dense epochs. Grey rectangles indicate times in which stimulations were analyzed for this example. (**B**) FEPSP amplitudes during sleep dense epochs, measured in NREM (as illustrated by box around hypnogram on left). Epochs passing the >65% threshold are shown on left, and those passing the >75% threshold are shown on right. FEPSP amplitude was normalized to the beginning of each epoch. Each dot is data from one epoch, averaged over 10 minute bins; n = 4 animals. Blue line = mean, shading = SEM, black dotted line = 1 (i.e. no change). (**C**) Same as (B) but for REM events. (**D**) Same as (B) but normalized fEPSP amplitude during wake, measured during active wake. (**E**) Same as (D) but for quiet wake events.

Fig 6 – figure supplement 1: Stability of fEPSP slope during prolonged periods of sleep or wake. Time series data of thalamocortical evoked fEPSP slopes over the course of either sleep or wake dense epochs. (**A**) Hypnograms showing an example of a wake and sleep dense epoch. Black rectangles indicate example times in which stimulations were analyzed for this example. (**B**) FEPSP slopes during sleep dense epochs, measured in NREM (as illustrated by box around hypnogram on left). Epochs passing the >65% threshold are shown on left, and those passing the >75% threshold are shown on right. FEPSP slope was normalized to the beginning of each epoch. Each dot is data from one epoch, averaged over 10 minute bins; n = 4 animals. Blue line = mean, shading = SEM, black dotted line = 1 (i.e. no change). (**C**) Same as (B) but for REM events. (**D**) Same as (B) but normalized fEPSP slope during wake, measured during active wake. (**E**) Same as (D) but for quiet wake events.

Fig 7: **Thalamocortical fEPSP amplitude measured before and after prolonged periods of sleep or wake.** (**A**) Hypnograms showing an example of a wake (left) and sleep (right) dense epoch, with flanking periods of the opposite state. Grey rectangles indicate when measurements were taken for comparison. (**B**) Example fEPSP waveforms before and after a wake (left) or sleep (right) dense epoch. (**C**) Normalized change in fEPSP amplitude after wake or sleep (wake change, p>0.9; sleep change, p>0.9; Wilcoxon signed-rank test). (**D**) Correlative plot of duration of wake dense epoch against normalized change in fEPSP amplitude. Black line = linear fit; R^2 value and p value included on plot. Inset in top right is an expansion of plot between 0 and 3.5 hours. (**E**) Same as (D) but for sleep dense. N = 39 wake dense epochs, 55 sleep dense epochs, 4 animals.

Fig 8: Characterization of thalamocortical evoked spiking in visual cortex between behavioral states and its stability across sleep and wake. (**A**) Heat plot response of the firing rate of each cell isolated during stimulation. Data are grouped by layer, and sorted by the magnitude of response within each layer. N = 45 cells, 4 animals. (**B**) Evoked spikes per stimulation for each of the 4 vigilance states. The y axis represents the average number of evoked spikes for a single stimulation, calculated as spikes in a 15 ms post stimulus window, minus the expected number of baseline spikes in that window; each dot represents a cell. Evoked spikes varied significantly by vigilance state (AW-QW, p=3.4e-4; AW-NREM, p=6.56e-4; AW-REM, p=4.70e-4; QW-NREM, p=0.0022; QW-REM, p=0.0122; NREM-REM, p=0.0380). (**C**) To determine whether evoked spike rates were affected by time spent awake or asleep, we compared evoked spiking before and after a sleep or wake dense epoch, using a >75% time in state threshold, as for fEPSPs in Figure 7. Hypnograms show examples of wake or sleep dense epochs with flanking periods of the opposite state. Gray rectangles indicate the stimulation times chosen for the analysis. (**D**) example raster plots and post-stimulus time histograms of firing rate for an example cell pre-post sleep. (**E**) Shows the number of evoked spikes before and after wake (left) or sleep (right) dense epochs; each dot is one cell, averaged across all of the wake or sleep dense epochs (wake dense n = 32 cells, 4 animals; sleep dense n = 24 cells, 4 animals). (**F**) Correlative plot of duration of wake dense epoch against change in evoked spikes; each dot represents the response of a single cell to a single epoch (post – pre). Black line = linear fit; R^2 value and p value included on plot. Inset is an expansion of plot between 0.25-3.5 hours. (**G**) Same as (F) but for sleep dense epochs. N = 35 wake dense epochs, 50 sleep dense epochs, 4 animals.