Using genetic variation to disentangle the complex relationship between food intake and

health outcomes.

Authors

Nicola Pirastu[1][†], Ciara McDonnell[1,2][*],Eryk J. Grzeszkowiak[1][*], Ninon Mounier[3, 4], Fumiaki

Imamura[5], Jordi Merino[6,7,8], Felix R. Day[5], Jie Zheng[9], Nele Taba[10,11], Maria Pina Concas[11],

Linda Repetto[1], Katherine A. Kentistou[1,2], Antonietta Robino[12], Tõnu Esko[11,8],Peter K.

Joshi[1], Krista Fischer[11], Ken K. Ong[5],Tom R. Gaunt[9], Zoltan Kutalik[3,4], John R. B. Perry[5],

James F. Wilson[1,13].

Affiliations

[1] Centre for Global Health Research, Usher Institute, University of Edinburgh, Teviot Place,

Edinburgh, EH8 9AG, Scotland.

[2] Centre for Cardiovascular Sciences, Queen's Medical Research Institute, University of

Edinburgh, Royal Infirmary of Edinburgh, Little France Crescent, Edinburgh EH16 4TJ,

Scotland

[3] Center for Primary Care and Public Health, University of Lausanne, Lausanne, Switzerland

[4] Swiss Institute of Bioinformatics, Lausanne, Switzerland

[5] MRC Epidemiology Unit, Institute of Metabolic Science, Cambridge Biomedical Campus,

University of Cambridge School of Clinical Medicine, Box 285, Cambridge, CB2 0QQ, UK

[6] Diabetes Unit and Center for Genomic Medicine, Massachusetts General Hospital, Boston,

MA, USA

[7] Program in Medical and Population Genetics, Broad Institute, Cambridge, MA, USA

[8] Department of Medicine, Harvard Medical School, Boston, MA, USA

[9] MRC Integrative Epidemiology Unit, Bristol Medical School, Bristol, UK

[10] Estonian Genome Center, Institute of Genomics, University of Tartu, Tartu, Riia 23b,

51010, Estonia

[11] Institute for Maternal and Child Health - IRCCS "Burlo Garofolo", Trieste, Italy

[12] Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Nobels väg

12A, SE-171 77 Stockholm, Sweden

[13] MRC Human Genetics Unit, Institute of Genetic and Molecular Medicine, University of

Edinburgh, Western General Hospital, Crewe Road, Edinburgh, EH4 2XU, Scotland

[14] Institute of Molecular and Cell Biology, University of Tartu, Tartu, Riia 23, 51010, Estonia

*Authors contributed equally to this work.

†Correspondence should be addressed to Nicola Pirastu, nicola.pirastu@ed.ac.uk

Abstract:

Despite food choices being one of the most important factors influencing health, efforts

to identify individual food groups and dietary patterns that cause disease have been

challenging, with traditional nutritional epidemiological approaches plagued by biases

and confounding. After identifying 302 individual genetic determinants of dietary

intake in 445,779 individuals in the UK Biobank study, we develop a statistical genetics

framework that enables us, to directly assess the impact of food choices on health

outcomes. We show that the biases which affect observational studies extend also to

GWAS, genetic correlations and causal inference through genetics, which can be

corrected by applying our methods. Finally, by applying Mendelian Randomization

approaches to the corrected results we identify some of the first robust causal

associations between eating patterns and cancer, heart disease, obesity, and several

other health related risk factors, distinguishing between the effects of specific foods or

dietary patterns.

Introduction

Given their profound impact on human well-being, diet is one of the most studied human behaviours. Quality, quantity, and patterns of consumed foods are associated with a wide range of medical conditions such as metabolic, inflammatory, or mental health diseases[1]. However, despite the growing number of studies reporting associations between diet and health outcomes, it has been challenging to establish causal relationships due methodological limitations such as measurement error, confounding, and reverse causation. To date, several methods have been devised to try to account for intrinsic limitations in nutritional studies such as calibration of food records[2] or the implementation of domiciled feeding studies (ie. the PREDICT study[3]) in which participants are instructed to eat only the food provided by the study. Although these methods have helped in addressing some the limitations related to food consumption measurement, problems still remain especially when it comes to measure the effects of food on health over a long period of time.

In this context genetics may represent an alternative approach through the use of Mendellian Randomization. Mendelian Randomization (MR) is a methodological approach in which genetic variants associated with a phenotype of interest are used as instrumental variables to measure the "life-long effect of an exposure" to an outcome.[4] To date, several MR studies have been designed to investigate the associations between the consumption of single food groups, such as alcoholic beverages[5] , coffee[6], milk[7–9] and specific health outcomes, but a systematic study investigating the overall role of diet is missing. In addition, previous MR studies have not accounted for the fact that genetic variants associated with reported dietary intake may be primarily associated with other risk factors or social determinants of health which may confound the causal estimates if used. In addition, previous studies on single food groups have not accounted for inter-relationships between different foods thus limiting the interpretability of the findings.

Given the complex number of factors that are driving the association between diet and health

outcomes, the present study was designed to initially identify the genetic variants associated

with reported food consumption, and then to leverage a causal inference statistical framework

to systematically investigate the causal effects of dietary factors on health outcomes, while

accounting for the effects that health determinants have on habitual dietary intake reporting.

Methods

Study population and genome-wide association for dietary intake

The UK Biobank[10] is a large population-based cohort including 500 000 adults aged between

40 and 69 years at baseline across 22 assessments centers in the United Kingdom. Data were

collected based on clinical examinations, assays of biological samples, detailed information

on self-reported health characteristics, and genome-wide genotyping. Dietary intake in UK

Biobank was assessed using a food frequency questionnaire which included questions about

the frequency of consumption specific foods and beverages over the past year. The number of

samples used for each trait can be found in table S1 while a detailed description of the

phenotypes, can be found in the in the supplementary methods 1.2 and table S2.

We used the BOLT-LMM software[11] to assess the association between the genetic variants

across the human genome and 29 food phenotypes. Analyses were conducted on genetic data

release version 3 imputed to the HRC panel[12], as provided by the UK Biobank

(http://www.ukbiobank.ac.uk/wp-

content/uploads/2014/04/UKBiobank_genotyping_QC_documentation-web.pdf). Population

stratification was assessed using LD-score regression as implemented in LD Hub[13,14] using

the LD scores provided with the software. Table S15 reports for each food trait the LD

regression intercept and heritability estimation using ldsc. Cluster analysis conducted on the

foods identified 5 main groups of traits (see additional online methods paragraph 1.8 and 2.2

for details of group definition) and we thus set the genome-wide significance threshold at

1×10^{-8}. Work within was conducted under UKB application 19655. Participants enrolled in

UK Biobank have signed consent forms. Replication analyses for identified signals

associated with food phenotypes were conducted independently by using genetic and dietary

data from the EPIC-Norfolk Study[15] and the Fenland Study[16]. Details additional online

methods 1.4.

Investigating the effect of health outcomes on reported food intake using MR.

Univariable MR analyses were initially conducted to measure the causal effect of health

outcomes on food consumption using the TwoSampleMR[17] R package. Exposures of interest

were selected amongst those for which nutritional advice is given and included body mass

index (BMI), low density lipoprotein cholesterol (LDLc), high density lipoprotein cholesterol

(HDLc), Total cholesterol, Triglycerides, Diastolic and Systolic blood pressure, Type 2

diabetes, and coronary artery disease. In addition, we included educational attainment as a

proxy of socio-economic status which is likely to affect food consumption. The full list of

studies from which the summary statistics were derived is detailed in Table S6. For each

exposure we selected all SNPs with $p < 5 \times 10^{-8}$ and $r^2 < 0.001$ to be used as instruments in the

MR analysis. After performing stepwise heterogeneity pruning we performed MR analysis

using the inverse variance method[18]. We then tested if the intercept from the MR-Egger[19]

regression was different from zero ($p < 0.05$). If this was the case, MR-Egger was used for the

analysis instead.

Measuring the direct effects of food types on health outcomes and identifying genetic

variants with predominantly direct-effects

One of the most important assumptions in MR is that the effect of the instrument on the

outcome must be mediated only through the exposure of interest (sometimes referred as

exclusion restriction criteria)[20]. In this light the instruments whose effect on food is mediated

through the health outcomes or through educational attainment may violate this assumption

acting as confounders in the relationship between the exposure and the outcome. Moreover if

the mediating trait is acting on the reporting of food consumption and not food consumption

itself it would mean that the genetic variant is not truly associated to food consumption and it

would thus not be a valid instrument. It is thus important to estimate the direct effect(i.e., the

effect that acts directly on food intake rather than is mediated through other factors see Figure

1) the SNPs are exerting on actual food consumption in order to properly select the genetic

variants to be used as instrumental variables.

To this end we use a modified version of the method implemented in bGWAS[21]. This method

consists of a first step were the phenotype of interest (i.e., food consumption) is used as

outcome in multivariable MR. Next, exposures of interest are selected using a forward step

wise regression selection algorithm where each exposure is added until their p-value is less

than 0.05. The method provides a corrected estimate for each genetic variant of its effect on

the outcome trait once all mediated effects are removed. Further details can be found in

supplementary methods 1.6. In order to identify genetic variants with only a direct effect on

the phenotype of interest we defined the corrected to uncorrected ratio (CUR) as the ratio

between the corrected and the uncorrected effects (see additional methods 1.7 for a detailed

explanation).

Fig. 1 Direct and indirect SNP effects. *The plot shows the causal path of exemplar genes identified for cheese consumption.*
In the multivariable MR model cheese consumption is causally influenced by educational attainment (EDU), low density
lipoprotein cholesterol levels (LDL) and systolic blood pressure (SBP). The effect of PDCH17 and is mediated through
educational attainment, while SIX3 has a direct effect on cheese consumption. The mediated effects cannot be used reliably
as MR instruments as they could be affecting either consumption or its reporting. Moreover, they could act as confounders

in the MR analysis and thus they need to be identified.

The threshold to define genetic variants with non-mediated effects (CUR=1±0.05) is based on

simulations provided in the supplementary note 2.1 and on the genetic variants with known

biological function (ie. bitter receptors). We defined as "non-mediated" those SNPs whose

CUR fell within the defined ranges while "uncertain" the others. We applied bGWAS to all

29 food phenotypes. As potential mediators, we used the same cardiometabolic phenotypes as

before except total cholesterol to avoid collinearity issues with LDL and HDL cholesterol,

and we added summary statistics from Crohn's disease and ulcerative colitis as they are

likely to affect dietary patterns. A Detailed discussion of this approach can be found in

supplementary methods 1.6.

Genome-wide genetic correlations between corrected dietary intake and health

outcomes.

We used LD-score regression implemented in LD Hub[13,14] to estimate genome-wide genetic

correlations between dietary intake phenotypes and 844 health outcomes and intermediary

phenotypes. Genetic correlations were estimated both with the corrected and uncorrected

GWAS summary statistics using the bivariate LD-score regression model. Stratified LD-

score regression[22] analyses were implemented using ldsc and the annotation files available on

the ldsc website.

Definition of food group variables

In order to define measures of dietary patterns we first performed cluster analysis of the 29

food items applying iCLUST[23] to the corrected genetic correlation matrix between the

different foods. iCLUST clusters items in different groups based on a hierarchical structure

(Details additional methods 1.8). Figure 2 shows the resulting dendrogram and its

comparison with the genetic correlation matrix.

Fig2 Clustering of the food traits and definition of measures of dietary patterns. *The plot reports the genetic correlation*
plot amongst the food traits after applying the correction. The stars report the Bonferroni-corrected significant correlations.
The dendrogram and the boxes represent the clustering according to the ICLUST algorithm. The labels on the dendrogram
branches show the traits used to define each measure of dietary pattern. The dashed line represents the traits excluded from
the estimation of the dietary patterns traits. The "Vegetarian" trait was excluded from the "Meat PC" trait but was included
in the overall dietary pattern measure (All PC).

We then defined based on the resulting structure several measures of dietary pattern at

different levels of the dendrogram as shown in Figure 2. For each measure we performed

principal component analysis of the items which participated to each group. The rotation

matrix was derived from the eigen decomposition of the correlation matrix of the foods in the

PC trait of interest. For example for the Coffee PC measure we performed principal

component analysis of "Ground Coffee", "Instant Coffee" and "Decaf Coffee". Once the

rotation matrix was estimated for each SNP its effect on the new measure was estimated as

the linear combination of the effect on each food trait using as weights the loadings on each

PC. A correlation plot of the loadings of each item onto the PC traits can be found in figure

S3.

MR analyses to assess causal relationships between food intake and health outcomes

MR analyses were conducted to estimate the effects of the food phenotypes on 79 health

related phenotypes (see table S17 for details) available in MR-base.[17] Genetic instruments for

each exposure of interest included independent genetic variants ($p<5x10^{-8}$ and pruning for

LD ($r^2<0.001$)). For dietary patterns exposures SNPs were selected as outlined in additional

methods 1.12. For the main analysis we restricted the genetic instruments to those that only

had evidence of a direct effect (i.e., not affecting the main exposure through a different

pathway; CUR 1±0.05). Discussion of the relationship with other methods can be found in

supplementary note 2.7. Weights for the genetic instruments were based on the uncorrected

effects. To verify the effects of using only direct effect only SNPs on MR, all the analyses

were also conducted without applying the CUR filtering.

After selecting the genetic instruments, exposure and outcome data were harmonised. The

MR estimates were tested for heterogeneity and outliers were removed using the MR-Radial

method.[24] MR analyses were based on the inverse variance weighted method, which

estimates the causal effect of an exposure on an outcome by combining ratio estimates using

each variant. A random effect model was used if significant heterogeneity between the

different estimates was detected. We then tested for the presence of directional pleiotropy

using the intercept from the MR-Egger regression. MR median and MR-Raps were used as

sensitivity analyses. All results have been made available through an online app (

https://npirastu.shinyapps.io/Food_MR/) and can be found in additional table S18.

Patient and public involvement

This research did not involve patients or the public as it uses data from the UK Biobank study

that were previously obtained from a cohort of people who had already been recruited. As

such, no patients or member of the public were involved in the design or implementation of

this study or the research questions addressed.

Results

Genetic variants associated with food intake

In a GWAS of 29 food phenotypes we identified 414 genetic associations in 260 independent

loci (Fig 3 and additional table S4) at Bonferroni corrected level of significance (P< 1×10^{-8}).

Fig. 3 302 independent genomic loci associate with food choices. *Results for both univariate (260 loci) and multivariate*
(additional 42 loci see paragraph S2.3) analyses are included. For each SNP the lowest p-value for all traits was plotted.
The upper panel represents the unadjusted GWAS associations while the lower panel represents the association with food
choices, after adjustment for mediating traits, such as health status.

Replication was sought in two additional UK-based cohorts including up to 32,779

participants. Despite relatively limited power in replication cohorts, concordant direction of

effect was observed for 82% of the signals (p=7.82×10^{-35}, Binomial test; Table S5), and

nominal significance was achieved by 32% of the signals (p=9.47×10^{-54}). Gene prioritization

is described in supplementary methods 1.10 while biological annotation, network analysis

and tissue enrichment analysis are discussed in additional paragraphs 1.11, 2.4 and 2.5.

Several of the identified loci have been previously associated with BMI. However, contrary

to our expectations, the BMI-raising allele was consistently associated with lower reported

consumption of energy-dense foods such as meat or fat, and higher reported intake of low-

calorie foods.

Genetic variants associated with food intake are strongly influenced by other

phenotypes

In univariable MR we identified 81 instances in which health-related traits significantly

influencing food intake (Fig. 4 additional table S7). In particular BMI and Educational

attainment influenced more than 50% of the food traits. Similar effects extend to a broad

range of traits, for example LDL and triglycerides influenced 15 and 18 traits respectively.

Higher genetically-determined CAD associates with higher consumption of fish and red

wine, and lower consumption of whole milk, salt and lamb. These findings suggest that some

of the signals identified in GWAS for reported food phenotypes are not directly associated

with food intake but are mediated through a wide range of potential confounders.

Fig 4. Health status influences reported food choices. *The plot reports only the univariable MR results which were*
significant at FDR<0.05. For each food outcome the effect estimate (β) is reported in standard deviations of the exposure
trait, together with 95% confidence intervals. Each colour represents a different exposure. BMI, body mass index; CHD,
coronary heart disease; DBP, diastolic blood pressure; HDL, high density lipoprotein cholesterol; LDL, low density
lipoprotein cholesterol; TotalC, total cholesterol. Champ/Wh wine, champagne, white wine. Temp, temperature.

The Multivariable MR confirmed the univariable MR results (Supplementary Fig S4 panel A

and Supplementary Table S8). The percentage of genetic variance for the reported food

phenotypes explained by health determinants ranged from 42% for cheese to ~0% for

fortified wine and white wine/champagne (Supplementary Fig S3 panel B and Supplementary

Table S16). We systematically compared the estimated effect sizes of each genetic variants

influencing food consumption before and after correcting for the effect of health determinants

and showed that in many loci the variant initially identified for food phenotypes changed

dramatically after taking into account the effect of health factors (Fig. 3, see Supplementary

file 1 for trait-specific plots). For example, the effect size of the lead *FTO* variant

(rs55872725, *p*=2x10^{-29}) on milk fat percentage chosen decreased three-fold after accounting

for the mediated effects. To further explore the magnitude of this indirect effect on food

intake phenotypes, we compared the correlation patterns between the 29 food phenotypes and

832 phenotypes present in the LD hub[14] database identifying great differences. For example,

low fat milk intake was correlated with a beneficial effect on body fat percentage (r_G = -0.43)

but this association diminished to near zero (r_G = -0.04) after accounting for indirect effects

(Supplementary Data 2.2 and additional table S10). The effects of the correction procedure

on the genetic correlation amongst the traits and with the 844 health traits are discussed in

supplementary note 2.2 while full results can be found at in table S9 and browsed at

https://npirastu.shinyapps.io/rg_plotter_2/. These findings highlight the relevance of biases

and confounding in genetic correlation studies, and provide the framework to study complex

physiological relationships.

Causal inference analyses for diet phenotypes and health outcomes

A total of 230 out of 414 genetic variants initially associated with food phenotypes

(corresponding to 169/260 loci) were categorized as "non-mediated" associations (Table S3).

The balance of uncertain to non-mediated genetic associations varied by food group, ranging

from none uncertain for tea, spirits and processed meat, to all uncertain for percentage fat in

milk and adding spread to bread (Table S3).

In two-sample MR analyses we found 141 significant associations between food phenotypes

and health outcomes after multiple test correction (pFDR < 0.05, Table S18).

Of these 89 showed no sign of heterogeneity amongst the estimates (heterogeneity test p

>0.05). Figure 5 reports full results for all significant food exposure trait outcome pairs.

Fig 5. Significant effects of food choice on disease related traits. *The heatmap reports the results for all significant food*
trait exposure trait outcome. Only dietary pattern exposures summarising the overall group consumption (PC1) have been
reported. All exposures have been aligned to have a positive loading onto the "overall unhealthy diet" measure. Significant
*food/trait association are indicated with * if they show no sign of heterogeneity while @ if they show significant*
heterogeneity. To facilitate meaningful visualisation and maximise the appearance of signal rather than noise, we applied a
shrinkage method - imposing a bayesian prior assumption on the distribution of beta (mean 0, SD 0.1), and conjugating that
with the likelihood of our results and then taking mean beta from the resulting distribution, thus shrinking estimates with
larger SEs more towards 0. Abbreviations: BMI Body Mass Index, WHR Waist to Hip Ratio, TRY trygliceries, TC total
cholesterol, HDL HDL cholesterol, LDL LDL cholesterol, Hb% Haemoglobin percentage, MCV Mean Corpuscolar Volume,
MPV Mean Platelet Volume, PLT Platelet count, Edu Educational attainment, CHD Coronary Heart Disease, MI
Myocardial Infarction, CD Chron's Disease, IBD Inflammatory Bowel Disease, Serous ovarian cancer:LG/LP low grade
low potential. H/L serous ovarian cancer High and Low grade serous ovarian cancer, RA Rheumatoid Arthritis.

Overall we found evidence supporting the beneficial effect of a healthy diet on health

outcomes. For example, for obesity/adiposity outcomes, genetically-determined unhealthy

diet leads to very similar effects across, increasing obesity measurements. For lipid-related

outcomes, the overall unhealthy diet is associated with higher levels of LDLc with no

significant heterogeneity, but no association with any of the other dietary traits. The overall

unhealthy diet was also strongly associated with Lung adenocarcinoma (OR 1.4xSD CI 1.2-

1.9) which seemed to be driven mostly by alcoholic beverages.

We identified 51 instances in which we would have not detected a significant result without

filtering out the non-direct effect instruments such as the effect of increased fruit

consumption on triglycerides levels (estimated uncorrected effect= -0.03 (SE=0.05) vs.

estimated corrected effect = -0.17 (SE=0.05)) or the effect of increased beef consumption on

height (uncorrected effect = -0.02 (-0.17, 0.13) vs corrected effect = -0.52 (0.29, 0.74). In

addition, we found 124 food/trait relationships which were not significant after applying

CUR filtering, showing that either confounding effects or reduced power explain the lack of

association (see additional note 2.6). For example, red wine consumption was initially

associated with increased BMI (uncorrected effect =0.22 (SE 0.05)) and waist circumference

(uncorrected beta= 0.26 (SE 0.07), but after correcting for CAD liability, both effects

disappeared (corrected effect for BMI 0.05 (SE 0.06), corrected effect for WC 0.005 (0.08)).

On the flip side, we showed that the effect of red wine on mean corpuscular volume remains

substantially unchanged when applying the filtering approach (beta 0.07 (SE 0.02)

uncorrected and 0.065 (SE 0.02) corrected), suggesting that our approach could precisely

identify relevant biological relationships.

A full description of our findings are found in table S18 and have been made available

through an online app (https://npirastu.shinyapps.io/Food_MR/).

Discussion

In this study we have provided quantitative data about the complex interplay between diet and health outcomes showing that the causal path from food intake to adverse health outcomes is not unidirectional and may be influenced by reverse causation and confounding even when MR is used. We showed that genetic correlations and causal inference can be improved by leveraging statistical approaches that take into account this mediated effects and identify genetic variants that have a only non-mediated effects on the exposure of interest. This information allowed us to perform causal inference analyses that helped identifying more reliable potential causal effects of food on health outcomes.

Results in context

Previous MR studies have mainly focused on specific food groups such as coffee, alcohol and milk consumption while none has comprehensively investigated the role of different food groups on health outcomes. Our results support previous observations such as the effect of alcohol consumption on coronary artery diseases reported in previous MR studies. In addition, we were able to confirm similar previous results detecting no evidence of an effect on IBD and CD[25] , ovarian cancer[26] or rheumatoid arthritis[27].

Findings from this study also suggest that the same biases that affect measures of food consumption such as reporting bias, confounding and reverse causation are reflected also in studies focusing on genetic associations. We have shown that these issues extend beyond obesity and socio-economic status including a broader range of intermediate factors. For example blood LDL and triglycerides concentration influence a wide variety of food traits thus being important factors to be considered as potential sources of bias, yet to our knowledge this is the first time this has been reported. For our analyses we have used UK biobank in which participants were aged between 40 and 60 at the time of the questionnaire,

it is likely that a younger cohort will suffer less from some of these (ie. LDL cholesterol or

blood pressure) as it is unlikely that they will display pathological level of these traits.

Our results are in contradiction to some previous studies in which no evidence of reverse

causation influencing genetic susceptibility for dietary patterns was reported.[28,29] We believe

that this difference is due to our novel approach, which is not based on using the potential

mediators as covariates, but rather exploits MR, which should be able to distinguish the

forward and reverse effects when the causal relationship is bidirectional. We have thus shown

that it is possible, through the use of available data and methods, to disentangle these

different colliding effects and to select the instrumental variables which show a non-mediated

effect, thus enabling the use of MR for the assessment of causal relationships between food

and health.

Many studies have looked at the relationship between nutritional composition and health

outcomes. One of the most salient examples is the relationship between saturated fat intake

and cardiovascular disease and all-cause mortality, in which recent studies suggest that food

sources of saturated fatty acids are more important than saturated fat content per se[Citation error].

Our study provide a new angle on the importance of food sources by providing evidence that

foods with similar nutrient profile, for example cheese and meat, which are both relatively

high in saturated fat and protein, have opposite effects on some metabolic risk factors such as

BMI (Figure S24 A) but there is no difference in other phenotypes such as blood lipids. A

similar conclusion can be drawn if we look at the foods which have the greatest effect on

triglycerides, fruit, vegetables and fish; all with very similar lowering effects (Figure S24 B),

which have relatively different macronutrient compositions. While the findings require

further investigations in mechanisms and related behaviours, our genetic evidence lends

support for the importance of studying foods in their complexity and not as a mere mixture of

nutrients. This approach, in fact, does not consider that the sources of the nutrients are not

equal due to the food matrix, the different preparations and that foods are seldom consumed

by themselves but in patterns which are likely to modify the effects on health.

Our findings illustrate that the effect of diet on health outcomes is complex, and components

of specific food groups have a differential association with health. In this case, although fish

and fruit and vegetables have a very different macronutrient composition it was impossible to

separate their effect on triglyceride concentrations. This suggests that at least in this case the

macronutrient composition is not as important as the an overall tendency to eat certain foods

and it highlights the importance of always including the assessment of dietary patterns before

claiming health effects of single foods or nutrients.

Some of the effects we have identified are more complex to explain and will need different

sources of evidence to be understood. For example we have found that the overall unhealthy

diet is associated to a higher risk of both lung andenocarcinoma and lung cancer. When

looking more closely to which of food explain this association the most we can see that

Alcohol seems to be driving the overall effect. One possibility is that this relationship is

confounded by smoking through a common tendency to addictive behaviours. However a

recent GWAS on cigarette smoking in Japan Biobank[30] reported a strong association between

the ALDH2 gene and number of cigarettes per day smoked which has also been associated to

differences in alcohol consumption[31], suggesting a causal effect of increased alcohol

consumption on increased smoking thus predisposing to lung cancer. Regardless of the

interpretation this example shows how complex the interpretation of MR results are when

behavioural traits are involved as they influence each other constantly creating a complex net

of interrelationships. This also points to the need of extreme care when claiming beneficial

health effects of food and multiple sources of evidence and approaches should always be

used before translating these findings into public policies.

Our study has several potential limitations. First, the number of items available in the dietary

questionnaire in the UK BioBank is limited, and therefore it limited our ability to capture

overall diet or specific food groups not detailed. The inclusion of white and relatively healthy

and educated participants from UK Biobank may have limited the generalisability of our

findings. Estimated effect sizes could be inflated because of the underestimation of the SNP

effects on the actual food trait consumption, rather than its self-report, if so, this will have

inflated our estimates of the effects of food on health, due to the noise in the questionnaire

responses, and warrants further statistical investigations. Even so, our method should not

have falsely identified a causal effect or reversed its direction, but further studies are needed

to assess the precise effect sizes.

In conclusion, our findings show that overall what is generally considered a healthy diet leads

to many favourable health outcomes and to reducing a wide range of risk factors broadly

agreeing with current guidelines aimed at reducing meat and alcohol consumption while

increasing fruit vegetables and fish. We also show that some of these effects are mostly

reconductable to specific food or group of foods which however are not characterized by

common nutrient composition thus adding granularity to our knowledge on the effect of diet

on health. This information can be useful to inform the design and implementation of future

studies to reduce the burden of diet-related diseases.

Author Contributions

NP,JFW,JRBP,ZK,EJG,FRD,KKO contributed to the study design.

JFW,TE,JRBP,AR,TG,FI,KKO,FRD contributed data.

NP,CMD,EJG,NM,FI,JZ,NT,KAK,MPC, performed the statistical analyses. NP, JFW,ZK,

JRBP, JM,TE, NT,KF,CMD,LR,EJG,FI,KKO,FRD contributed to the interpretation of the

results. All authors contributed to writing and editing of the text.

Acknowledgements

J.F.W. acknowledges support from the MRC Human Genetics Unit quinquennial programme grant "QTL in Health and Disease" (MC_UU_00007/10). This research has been conducted using the UK Biobank Resource under Application Number 19655. We would like to thank Professor George Davey Smith for the precious feedback, Erin MacDonald-Dunlop, and Pascale Lubbe for help with statistical analyses and Dr. Nana Matoba for providing the results from the smoking GWAS.

EGCUT was funded by Estonian Research Council Grant IUT20-60, PUT1660 (T.E.), PUT1665 (K.F.), the European Union through the European Regional Development Fund grant no. 2014-2020.4.01.15-0012 GENTRANSMED and 2014-2020.4.2.2, and Estonian and European Research Roadmap grant no.2014-2020.4.01.16-0125. The EPIC-Norfolk study (DOI 10.22025/2019.10.105.00004) has received funding from the Medical Research Council (MR/N003284/1, MC-PC_13048, and MC-UU_12015/1). The Fenland study (DOI: 10.1186/ISRCTN72077169) was funded by the Medical Research Council and the Wellcome Trust (Ref: 074548). J.P., K.O., F.I., and F.R.D. were funded by the UK Medical Research Council Epidemiology Unit core grant (MC-UU_12015/2, MC_UU_12015/5). T.R.G. receives funding from the UK Medical Research Council (MC_UU_00011/4). Z.K. received funding from the Swiss National Science Foundation (31003A_169929). We are grateful to all the participants who have been part of the project and to the many members of the study teams at the University of Cambridge who have enabled this research. J.M. was partially supported by a fellowship funded by the European Commission Horizon 2020 program (H2020-MSCA-IF-2015-703787) and by the National Institutes of Health grant P30 DK40561

Data Availability

All GWAS results will be made available through GWAS catalog at the time of publication.

All results from the MR analyses have been shared in the additional tables.

References

1. Misra, A. & Khurana, L. Obesity and the Metabolic Syndrome in Developing Countries. *The Journal of Clinical Endocrinology & Metabolism* vol. 93 s9–s30 (2008).

2. Naska, A., Lagiou, A. & Lagiou, P. Dietary assessment methods in epidemiological research: current state of the art and future prospects. *F1000Res.* **6**, 926 (2017).

3. Berry, S. E. *et al.* Human postprandial responses to food and potential for precision nutrition. *Nat. Med.* **26**, 964–973 (2020).

4. Zheng, J. *et al.* Recent Developments in Mendelian Randomization Studies. *Curr Epidemiol Rep* **4**, 330–345 (2017).

5. Millwood, I. Y. *et al.* Conventional and genetic evidence on alcohol and vascular disease aetiology: a prospective study of 500 000 men and women in China. *Lancet* **393**, 1831–1842 (2019).

6. Cornelis, M. C. & Munafo, M. R. Mendelian Randomization Studies of Coffee and Caffeine Consumption. *Nutrients* **10**, (2018).

7. Bergholdt, H. K. M., Nordestgaard, B. G., Varbo, A. & Ellervik, C. Milk intake is not associated with ischaemic heart disease in observational or Mendelian randomization analyses in 98,529 Danish adults. *Int. J. Epidemiol.* **44**, 587–603 (2015).

8. Vissers, L. E. T. *et al.* Dairy Product Intake and Risk of Type 2 Diabetes in EPIC-InterAct: A Mendelian Randomization Study. *Diabetes Care* **42**, 568–575 (2019).

9. Hartwig, F. P., Horta, B. L., Smith, G. D., de Mola, C. L. & Victora, C. G. Association

of lactase persistence genotype with milk consumption, obesity and blood pressure: a

Mendelian randomization study in the 1982 Pelotas (Brazil) Birth Cohort, with a

systematic review and meta-analysis. *Int. J. Epidemiol.* **45**, 1573–1587 (2016).

10. Sudlow, C. *et al.* UK biobank: an open access resource for identifying the causes of a

wide range of complex diseases of middle and old age. *PLoS Med.* **12**, e1001779 (2015).

11. Loh, P.-R. *et al.* Efficient Bayesian mixed-model analysis increases association power in

large cohorts. *Nat. Genet.* **47**, 284–290 (2015).

12. McCarthy, S. *et al.* A reference panel of 64,976 haplotypes for genotype imputation. *Nat.*

Genet. **48**, 1279–1283 (2016).

13. Bulik-Sullivan, B. K. *et al.* LD Score regression distinguishes confounding from

polygenicity in genome-wide association studies. *Nat. Genet.* **47**, 291–295 (2015).

14. Zheng, J. *et al.* LD Hub: a centralized database and web interface to perform LD score

regression that maximizes the potential of summary level GWAS data for SNP

heritability and genetic correlation analysis. *Bioinformatics* vol. 33 272–279 (2017).

15. Day, N. *et al.* EPIC-Norfolk: study design and characteristics of the cohort. European

Prospective Investigation of Cancer. *Br. J. Cancer* **80 Suppl 1**, 95–103 (1999).

16. Lotta, L. A. *et al.* Integrative genomic analysis implicates limited peripheral adipose

storage capacity in the pathogenesis of human insulin resistance. *Nat. Genet.* **49**, 17–26

(2017).

17. Hemani, G. *et al.* The MR-Base platform supports systematic causal inference across the

human phenome. *Elife* **7**, (2018).

18. Burgess, S., Butterworth, A. & Thompson, S. G. Mendelian randomization analysis with

multiple genetic variants using summarized data. *Genet. Epidemiol.* **37**, 658–665 (2013).

19. Bowden, J., Davey Smith, G. & Burgess, S. Mendelian randomization with invalid

instruments: effect estimation and bias detection through Egger regression. *Int. J.*

Epidemiol. **44**, 512–525 (2015).

20. Davies, N. M., Holmes, M. V. & Davey Smith, G. Reading Mendelian randomisation

studies: a guide, glossary, and checklist for clinicians. *BMJ* **362**, k601 (2018).

21. Mounier, N. & Kutalik, Z. bGWAS: an R package to perform Bayesian Genome Wide

Association Studies. *Bioinformatics* (2020) doi:10.1093/bioinformatics/btaa549.

22. Finucane, H. K. *et al.* Partitioning heritability by functional category using GWAS

summary statistics. doi:10.1101/014241.

23. Revelle, W. Hierarchical cluster analysis and the internal structure of tests. *Multivariate*

Behav. Res. **14**, 57–74 (1979).

24. Bowden, J. *et al.* Improving the visualization, interpretation and analysis of two-sample

summary data Mendelian randomization via the Radial plot and Radial regression. *Int. J.*

Epidemiol. **47**, 2100 (2018).

25. Georgiou, A. N., Ntritsos, G., Papadimitriou, N., Dimou, N. & Evangelou, E. Cigarette

smoking, coffee consumption, alcohol intake, and risk of crohn's disease and ulcerative

colitis: A Mendelian randomization study. *Inflamm. Bowel Dis.* (2020)

doi:10.1093/ibd/izaa152.

26. Zhu, J., Jiang, X. & Niu, Z. Alcohol consumption and risk of breast and ovarian cancer:

A Mendelian randomization study. *Cancer Genet.* **245**, 35–41 (2020).

27. Jiang, X. & Alfredsson, L. Modifiable environmental exposure and risk of rheumatoid

arthritis-current evidence from genetic studies. *Arthritis Res. Ther.* **22**, 154 (2020).

28. Meddens, S. F. W., de Vlaming, R., Bowers, P. & Burik, C. A. P. Genomic analysis of
diet composition finds novel loci and associations with health and lifestyle. *bioRxiv*
(2018).

29. Cole, J. B., Florez, J. C. & Hirschhorn, J. N. Comprehensive genomic analysis of dietary
habits in UK Biobank identifies hundreds of genetic loci and establishes causal
relationships between educational attainment and healthy eating. doi:10.1101/662239.

30. Matoba, N. *et al.* GWAS of smoking behaviour in 165,436 Japanese people reveals
seven new loci and shared genetic architecture. *Nat. Hum. Behav.* **3**, 471–477 (2019).

31. Matoba, N. *et al.* GWAS of 165,084 Japanese individuals identified nine loci associated
with dietary habits. *Nat. Hum. Behav.* **4**, 308–316 (2020).