The immunocytokine L19-TNF eradicates Sarcomas in combination with Chemotherapy agents or with immune check-point inhibitors

Riccardo Corbellari[1,2] [*], Lisa Nadal[1,2] [*], Alessandra Villa[2], Dario Neri[3] and Roberto De Luca[2] [#]

[1] University of Trento, Italy, CiBIO (Department of Cellular, Computational and Integrative Biology), Via Sommarive 9, 38123 Povo (Trento)
[2] Philochem AG, Otelfingen, Switzerland
[3] Department of Chemistry and Applied Biosciences, Swiss Federal Institute of Technology (ETH Zürich), Zurich, Switzerland

* The two authors have contributed equally to this work

Corresponding author
Roberto De Luca
Philochem AG, Philogen Group, Libernstrasse 3, CH-8112 Otelfingen (Switzerland)
Telephone: +41-43-5448811
Fax: +41-43-5448809
e-mail: roberto.deluca@philogen.com

Posted on non-commercial preprint servers
The data that support the findings of this study are openly available in bioRxiv.

Abstract

Antibody-cytokine fusion proteins (also called "immunocytokines") represent an emerging class of biopharmaceutical products, which are being considered for cancer immunotherapy. When used as single agents, pro-inflammatory immunocytokines are rarely capable of induce complete and durable cancer regression in mouse models and in patients. However, the combination treatment with conventional chemotherapy or with other immune-stimulatory agents typically increases the therapeutic efficacy of immunocytokines.

In this article, we describe combination treatments of a tumor-targeting antibody-cytokine fusion protein based on the L19 antibody (specific to a splice isoform of fibronectin) fused to murine tumor necrosis factor (TNF) with standard chemotherapy (dacarbazine, trabectedin or melphalan) or with an immune check-point inhibitor (anti-PD-1) in a BALB/c derived immunocompetent murine model of sarcoma (WEHI-164).

All combination treatments led to improved tumor remission compared to single agent treatments, suggesting that these combination partners may be suitable for further clinical development in sarcoma patients.

Introduction

Sarcomas represent an heterogenous group of malignancies that can originate from soft tissue, cartilage, bone and other connective tissues. In particular, soft tissue sarcomas are tumors, which stem from non-epithelial extra skeletal tissue (e.g. muscle, fat, blood vessels and others). These cancer types are rare and they mainly affect young patients, however they represent 15% of all pediatric cancers [1–3].

For small and localized tumors (i.e. in the limbs) surgical resection (sometimes combined with radiation) represents a curative therapy, however postsurgical recurrence is observed in 50% of patients. This treatment option is often not applicable for patients with metastatic disease or with tumors with a different histology (e.g. head, neck or abdomen) and in these cases, chemotherapy is administered as first-line treatment [4]. Doxorubicin represents the standard-of-care for sarcomas with a response rate of 12-30% [5]. Doxorubicin may be combined with dacarbazine or other agents (e.g. trabectedin), however these combination treatments typically do not increase the survival benefit compared to the monotherapy [6]. Moreover, they are usually associated with substantial toxicities.

Already in the late 19th century, William B Coley observed a case of a sarcoma patient who experienced a spontaneous tumor regression as a consequence of an erysipelas infection. This finding was the basis for the development of the "Coley's toxin", a mixture of heat-inactivated bacteria, which was found to increase the overall survival of sarcoma patients, in spite of the development of high fever and chill [7,8]. It was later demonstrated that Coley's toxin treatment induced a dramatic increase of TNF levels [9].

TNF is a homo-trimeric pro-inflammatory cytokine, mainly produced by macrophages, NK and T cells [10,11]. TNF is involved in inflammatory processes and as indicated by its name, it may display a potent anticancer activity against some malignancies, by promoting hemorrhagic necrosis and by inducing direct apoptosis of tumor cells [12–17]. Moreover, TNF may activate the immune system as a result of the increased inflammation [18–20].
The systemic administration of recombinant human TNF (Beromun®) in the clinic is limited by substantial side effects and the maximal tolerated dose (MTD) was found to be 300µg/m^2 [21]. Beromun® obtained marketing authorization only for locoregional treatments (i.e., for isolated

limb perfusion procedure) in patients with soft-tissue sarcomas. The product is frequently administered in combination with melphalan [22].

The use of recombinant cytokines for cancer therapy may cause substantial toxicities already at low doses, therefore limiting their clinical use and preventing escalation to therapeutically active dose regimens [23–26]. To increase the therapeutic efficacy of cytokine payloads, the generation of fusion proteins with tumor-homing antibodies has been proposed. Antibody-cytokine fusion proteins (also termed "immunocytokines") are capable of increasing the therapeutic index of the corresponding cytokine payload, as results of a selective localization at the site of disease [24–28].

We have previously described the immunocytokine L19-mTNF, consisting of the fully human antibody fragment L19 (specific to the alternatively spliced EDB domain of fibronectin, a marker of angiogenesis) [29,30] fused to murine TNF. The fusion protein was able to induce tumor regression in different murine cancer types either alone or in combination with chemotherapy or other immunocytokines [31–35].

The fully human L19-TNF (featuring human TNF) has been investigated in cancer patients (phase I and II completed) either as systemic administration, intralesional injection or in isolated limb perfusion settings [36–38]. Moreover, L19-TNF is currently being investigated in Phase III clinical trials for the treatment of Stage III B,C melanoma (EudraCT number: 2015-002549-72 and NCT03567889) in combination with L19-IL2 and of metastatic soft-tissue sarcoma (EudraCT number: 2016-003239-38 and NCT03420014) in combination with doxorubicin.

As previoulsy mentioned, melphalan, trabectedin and dacarbazine have been already considered for the treatment of sarcoma patients. For this reason, in this work we evaluated the combination of these chemotherepeutic agents with the immunocytokine L19-mTNF.

In the last decade, immune checkpoint inhibitors (e.g. anti CTLA-4, anti PD-1 and anti PD-L1) have shown encouraging clinical efficacy in the treatment of different malignancies [39–42]. However, not all cancer patients respond equally well to these drugs and combination strategies are currently being investigated in clinical trials aiming to improve therapeutic efficacy [43–45]. Moreover, the combination of immune checkpoint inhibitors with immunocytokines

showed interesting results in mouse models of cancers [15,46,47]. Encouraged by these results, we now investigated the combination of an anti-PD-1 antibody with L19-mTNF for the treatment of sarcoma lesions.

L19-mTNF and anti PD-1 antibodies explicates their function in activating the immune system. For this reason, immunocompetent BALB/c mice were chosen as animal model. Mice models have become indispensable in pharmaceutical and biomedical research to verify the effectiveness of drugs and treatment regimens. These analysis may ensure safe and valuable data for a future clinical trial application [48].

Aim of the work

The aim of this work was to investigate the combination of the immunocytokine L19-mTNF with chemotherapy agents (melphalan, trabectedin and dacarbazine) and with immune checkpoint inhibitors (anti PD-1) in a mouse model of sarcoma. The results of this work may provide a rationale and facilitate a future clinical investigation of these agents in sarcoma patients.

Materials and Methods

Cell lines and reagents

CHO cells and WEHI-164 were obtained from the ATCC. Cell lines were received between 2017 and 2019, expanded, and stored as cryopreserved aliquots in liquid nitrogen. Cells were grown according the supplier's protocol and kept in culture for no longer than 14 passages. Authentication of the cell lines also including check of postfreeze viability, growth properties, and morphology, test for mycoplasma contamination, isoenzyme assay, and sterility test were performed by the cell bank before shipment.

The production and purification of L19-mTNF was performed as described before [31]. Commercially available dacarbazine (Dacin®, Lipomed AG), trabectedin (Yondelis®, PharmaMar AG), melphalan (Alkeran®, Aspen Pharma Schweiz GmbH) and the anti-PD-1 antibody (clone 29F.1A12, BioXCell, Catalog # BE0273) were purchased.

Ethic statement

All animal experiments and maintenance were performed under a project license (license number 04/2018) granted by the Veterinäramt des Kantons Zürich, Switzerland, in compliance with the Swiss Animal Protection Act (TSchG) and the Swiss Animal Protection Ordinance (TSchV).

Study design

Two different therapy experiments were performed with 24 and 30 BALB/c mice, respectively. One control group (mice injected only with saline) was used for each experiment, 5 groups were used as experimental. Mice were randomized into treatment groups according to their tumor volume (**Supplementary table 1 and 2**); measurements were taken by the same experimenter to minimize any subjective bias. Mice were divided in six groups of 4 and 5 animals (4 mice per group for the first reported therapy, 5 mice per group for the second therapy) per cage before the first drug injection. Blinded experiments were not performed.

Experimental animals

A total of 54 female immunocompetent BALB/c mice, aged 8 weeks with an average weight of 20 g were used in this study. Mice were purchased from Janvier (Route du Genest, 53940 Le Genest-Saint-Isle, France).

Mice were raised in a pathogen-free environment with a relative humidity of 40-60% at a temperature between 18 and 26°C with daily cycles of 12 hours light/darkness according to guidelines (GV-SOLAS; FELASA). Mice were kept in an OHB Animal Facility in cages of maximum 5 mice, left for one-week acclimatization upon arrival and subsequently handled under sterile BL2 workbenches. Specialized personnel were responsible for their feeding; food and water were provided ad libitum. Mice were monitored daily (in the morning) in weight, tumor load, appearance (coat, posture, eyes and mouth moisture) and behavior (movements, attentiveness and social behavior).

Euthanasia criteria adopted were body weight loss > 15% and/or ulceration of the s.c. tumor and/or tumor diameter > 1500 mm and/or mice pain and discomfort. Mice were euthanized in CO_2 chambers.

Experimental procedure

Tumor cells were implanted subcutaneously in the flank of BALB/c mice using 5×10^6 cells (WEHI-164) with 0.5 ml 29G insulin syringes (MicroFine™+, BD medical). Tumor volume was measured with a caliper and volume was calculated using the formula: Tumor Volume = (Length[mm]*Width [mm]*Width [mm])/2. When tumors reached a suitable volume (approx. 70-100 mm^3), pharmacological treatment was performed.

L19-mTNF was dissolved in phosphate buffered saline (PBS), also used as negative control, sterile-filtered and administered at 0.1µg/g into the lateral tail vein three times every 48h. The intravenously route was chosen to achieve systemic administration of the bioactive payload. Dacarbazine and trabectedin were dissolved at 0.15 µg/g and administered intraperitoneally once, four hours after the first L19-mTNF injection. The intraperitoneal route was chosen based on previously reported experiments [49,50].

The anti-PD-1 antibody was administered at 10 µg/g into the lateral tail vein three times every 48h alone or in combination with L19-mTNF. Melphalan was administered once intravenously into the lateral vein once at 4.5 µg/g. Also in this case the intravenously route was chosen to achieve systemic administration of the bioactive payload [28,33,47].

All injections were performed in the morning with 0.5 ml 30G insulin syringes (Omnican® 50, Braun) in home cage for all mice groups. No anesthesia or analgesia were used for the injections.

Experimental outcomes

The primary experimental outcome was the reduction in tumor volume of treated mice, indication of the effect of the administrated bioactive agents and drugs.

The survival end point was reached when the tumor diameter exceeded 1500 mm, body weight loss > 15% and/or ulceration of the s.c. tumor.

Statistical method

Data were analyzed using Prism 7.0 (GraphPad Software, Inc.). Differences in tumor volume between therapeutic groups (until day 17, when n = 4 for the first therapy and until day 13, when n = 5 for the second therapy) were evaluated with the two-way ANOVA followed by Tukey post hoc test. P < 0.05 was considered statistically significant (*P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001).

Results

In a first study, we compared the therapeutic activity of L19-mTNF, dacarbazine and trabectedin, either alone or in combination, in immunocompetent BALB/c mice bearing WEHI-164 sarcomas [**Figure 1**], according to the experimental scheme [**Figure 1A**].

As expected, the treatment with saline was not able to induce cancer regression. When used as single agents, L19-mTNF and trabectedin exhibited a mild tumor growth retardation compared to the saline group. In addition, 1 mouse out of 4 treated with L19-mTNF showed a complete response. However, the combination therapy with these two agents led to an enhanced antitumor effect (3 out of 4 complete response). Dacarbazine administered as a single agent was capable of eradicate tumors in 3 out of 4 mice. The therapeutic efficacy was improved when administered in combination with L19-mTNF with (4 out of 4 complete response) [**Figure 1B, D**]. All treatments were well tolerated as evidenced by the comparison of body weight profiles. Only a transient toxicity was observed upon the first pharmacological treatment with the two combination treatments (L19-mTNF with either dacarbazine or trabectedin). However, all treated mice recovered the body weight loss in few days [**Figure 1C**].

In a second experiment, we compared the therapeutic activity of L19-mTNF in combination with melphalan or with an immune check-point inhibitor (anti-PD-1) in the same murine sarcoma model (WEHI-164) [**Figure 2**], according to the experimental scheme [**Figure 2A**]. The treatment with saline and melphalan alone was not able to induce cancer regression, by contrast, when used as single agent, L19mTNF was able to cure 1 mouse out of 5. Treatment with anti-PD1 alone induced cancer eradication in 2 out of 5 mice. The therapeutic activity of L19-mTNF was potentiated when used in combination with both melphalan (3 out of 5 complete response) or with an anti-PD1 antibody (4 out of 5 complete response) [**Figure 2B, D**]. All the treatments were well tolerated as evidenced by the comparison of body weight profiles [**Figure 2C**].

Discussion

We have now reported a pre-clinical evaluation of combination therapies with the fusion protein L19-mTNF.

BALC/c mice bearing WEHI-164 lesions were used for this purpose. In this study, we decided to use 4 and 5 mice per group, for the first and second therapy respectively, to have a robust statistic and reliable data applying the 3Rs principle. Mice health and comfort were constantly checked; the treatments were administrated trying to minimize any pain or distress of the animal. Mice cage were provided with housing, bedding and tissue; any cause of distress or suffering can indeed alter animals' physiology and immunology.

Dacarbazine is already an established treatment for sarcoma patients and in this study, we confirmed its potency as single agent. Moreover, the therapeutic efficacy was furhter improved when used in combination with L19-mTNF. A synergistic effect was also observed when L19-mTNF was used in combination with trabectedin and with melphalan.

Immune checkpoint inhibitors are rapidly gaining acceptance as broad-spectrum agents for many cancer indications and in this study, we demonstrated that L19mTNF efficiently synergized with PD-1 blockade, thus providing evidence for a further clinical investigation in sarcoma patients.

Conclusions

In this work, we presented a pre-clinical evaluation of combination modalities for the treatment of sarcoma lesions. The immunocytokine L19-mTNF was able to synergize with chemotherapeutic agents, which are already being considered for the treatment of sarcoma patients. Moreover, L19-mTNF was able to potentiate the anti-cancer activity of an anti-PD-1 antibody. The fully-human analog of L19-mTNF (L19-TNF), which is currently being investigated in phase I clinical trials in combination with doxorubicin, may also be considered in combination with other chemotherapeutic agents or with immune check-point inhibitors for the treatment of sarcoma patients. This study may also provide pre-clinical evidences for the upcoming phase I clinical trial of L19-TNF in pretreated Soft Tissue Sarcoma in combination with dacarbazine foreseen for 2020.

Acknowledgments

We would like to thank Dr. Teresa Hemmerle for helpful discussions and we gratefully acknowledge Alessandro Pedrioli for his help with experimental procedures.

Funding

We gratefully acknowledge funding from ETH Zürich and the Swiss National Science Foundation (Grant Nr. 310030_182003/1). This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (grant agreement 670603).

Conflict of interest

Dario Neri is co-founder, shareholder and member of the board of Philogen, a company working on antibody therapeutics. The authors declare no additional conflict of interest.

References

[1]   S. Singer, G.D. Demetri, E.H. Baldini, C.D.M. Fletcher, Management of soft-tissue sarcomas: an overview and update, Lancet Oncol. 1 (2000) 75–85. doi:10.1016/S1470-2045(00)00016-4.

[2]   M.A. Clark, C. Fisher, I. Judson, J.M. Thomas, Soft-Tissue Sarcomas in Adults, N. Engl. J. Med. 353 (2005) 701–711. doi:10.1056/NEJMra041866.

[3]   S. Sinha, A.H.S. Peach, Diagnosis and management of soft tissue sarcoma, BMJ. 341 (2010). http://www.bmj.com/content/341/bmj.c7170.abstract.

[4]   R.R. F., Systemic therapy for advanced soft tissue sarcomas , Cancer. 118 (2011) 1474–1485. doi:10.1002/cncr.26415.

[5]   G.K. In, J.S. Hu, W.W. Tseng, Treatment of advanced, metastatic soft tissue sarcoma: latest evidence and clinical considerations, Ther. Adv. Med. Oncol. 9 (2017) 533–550. doi:10.1177/1758834017712963.

[6]   R. Ratan, S.R. Patel, Chemotherapy for soft tissue sarcoma, Cancer. 122 (2016) 2952–2960. doi:10.1002/cncr.30191.

[7]   A.H. Shah, I. Jusué-Torres, M.E. Ivan, R.J. Komotar, N. Kasahara, Pathogens and glioma: A history of unexpected discoveries ushering in novel therapy, J. Neurosurg. 128 (2018) 1139–1146. doi:10.3171/2016.12.JNS162123.

[8]   C.O. Starnes, Coley's toxins in perspective, Nature. (1992). doi:10.1038/357011a0.

[9]   E.A. Carswell, L.J. Old, R.L. Kassel, S. Green, N. Fiore, B. Williamson, An endotoxin induced serum factor that cuases necrosis of tumors, Proc. Natl. Acad. Sci. U. S. A. (1975). doi:10.1073/pnas.72.9.3666.

[10]   E.Y. Jones; D.I. Stuart; N.P.C. Walker, Structure of tumour necrosis factor, (1989) 338:225–228. doi:338:225-228.

[11]   B.B. Aggarwal, S.C. Gupta, J.H. Kim, Historical perspectives on tumor necrosis factor and its superfamily: 25 years later, a golden journey, Blood. (2012). doi:10.1182/blood-2011-04-325225.

[12]   T. Hemmerle, P. Probst, L. Giovannoni, A.J. Green, T. Meyer, D. Neri, The antibody-based targeted delivery of TNF in combination with doxorubicin eradicates sarcomas in mice and confers protective immunity, Br. J. Cancer. (2013). doi:10.1038/bjc.2013.421.

[13]   P. Probst, J. Kopp, A. Oxenius, M.P. Colombo, D. Ritz, T. Fugmann, D. Neri, Sarcoma eradication by doxorubicin and targeted TNF relies upon CD8 + T-cell recognition of a

retroviral antigen, Cancer Res. (2017). doi:10.1158/0008-5472.CAN-16-2946.

[14] R. De Luca, A. Soltermann, F. Pretto, C. Pemberton-Ross, G. Pellegrini, S. Wulhfard, D. Neri, Potency-matched Dual Cytokine–Antibody Fusion Proteins for Cancer Therapy, Mol. Cancer Ther. (2017). doi:10.1158/1535-7163.mct-17-0211.

[15] R. De Luca, D. Neri, Potentiation of PD-L1 blockade with a potency-matched dual cytokine–antibody fusion protein leads to cancer eradication in BALB/c-derived tumors but not in other mouse strains, Cancer Immunol. Immunother. (2018). doi:10.1007/s00262-018-2194-0.

[16] B. Ziffels, M. Stringhini, P. Probst, T. Fugmann, T. Sturm, D. Neri, Antibody-based delivery of cytokine payloads to carbonic anhydrase IX leads to cancer cures in immunocompetent tumor-bearing mice, Mol. Cancer Ther. (2019) molcanther.1301.2018. doi:10.1158/1535-7163.MCT-18-1301.

[17] P. Murer, J.D. Kiefer, L. Plüss, M. Matasci, S.L. Blümich, M. Stringhini, D. Neri, Targeted Delivery of TNF Potentiates the Antibody-Dependent Cell-Mediated Cytotoxicity of an Anti-Melanoma Immunoglobulin, J. Invest. Dermatol. (2019). doi:10.1016/j.jid.2018.11.028.

[18] L.M. Sedger, M.F. McDermott, TNF and TNF-receptors: From mediators of cell death and inflammation to therapeutic giants - past, present and future, Cytokine Growth Factor Rev. (2014). doi:10.1016/j.cytogfr.2014.07.016.

[19] R. van Horssen, TNF- in Cancer Treatment: Molecular Insights, Antitumor Effects, and Clinical Utility, Oncologist. (2006). doi:10.1634/theoncologist.11-4-397.

[20] A.H. Van Der Veen, J.H.W. De Wilt, A.M.M. Eggermont, S.T. Van Tiel, A.L.B. Seynhaeve, T.L.M. Ten Hagen, TNF-α augments intratumoural concentrations of doxorubicin in TNF-α-based isolated limb perfusion in rat sarcoma models and enhances anti-tumour effects, Br. J. Cancer. (2000). doi:10.1054/bjoc.1999.1027.

[21] N.J. Roberts, S. Zhou, L.A. Diaz, M. Holdhoff, Systemic use of tumor necrosis factor alpha as an anticancer agent, Oncotarget. 2 (2011) 739–751. doi:10.18632/oncotarget.344.

[22] D.J. Grunhagen, J.H.W. De Wilt, W.J. Graveland, C. Verhoef, A.N. Van Geel, A.M.M. Eggermont, Outcome and prognostic factor analysis of 217 consecutive isolated limb perfusions with tumor necrosis factor-α and melphalan for limb-threatening soft tissue sarcoma, Cancer. (2006). doi:10.1002/cncr.21802.

[23] D. Schrama, R.A. Reisfeld, J.C. Becker, Antibody targeted drugs as cancer therapeutics, Nat. Rev. Drug Discov. (2006). doi:10.1038/nrd1957.

[24] D. Neri, Antibody–Cytokine Fusions: Versatile Products for the Modulation of Anticancer Immunity, Cancer Immunol. Res. (2019). doi:10.1158/2326-6066.CIR-18-0622.

[25] N. Pasche, D. Neri, Immunocytokines: A novel class of potent armed antibodies, Drug Discov. Today. (2012). doi:10.1016/j.drudis.2012.01.007.

[26] F. Bootz, D. Neri, Immunocytokines: A novel class of products for the treatment of chronic inflammation and autoimmune conditions, Drug Discov. Today. (2016). doi:10.1016/j.drudis.2015.10.012.

[27] P. Murer, D. Neri, Antibody-cytokine fusion proteins: A novel class of biopharmaceuticals for the therapy of cancer and of chronic inflammation, N. Biotechnol. (2019). doi:10.1016/j.nbt.2019.04.002.

[28] C. Hutmacher, D. Neri, Antibody-cytokine fusion proteins: Biopharmaceuticals with immunomodulatory properties for cancer therapy, Adv. Drug Deliv. Rev. (2018). doi:10.1016/j.addr.2018.09.002.

[29] L. Tarli, E. Balza, F. Viti, L. Borsi, P. Castellani, D. Berndorff, L. Dinkelborg, D. Neri, L. Zardi, A high-affinity human antibody that targets tumoral blood vessels, Blood. (1999).

[30] L. Borsi, E. Balza, M. Bestagno, P. Castellani, B. Carnemolla, A. Biro, A. Leprini, J. Sepulveda, O. Burrone, D. Neri, L. Zardi, Selective targeting of tumoral vasculature: Comparison of different formats of an antibody (l19) to the ED-B domain of fibronectin, Int. J. Cancer. (2002). doi:10.1002/ijc.10662.

[31] L. Borsi, E. Balza, B. Carnemolla, F. Sassi, P. Castellani, A. Berndt, H. Kosmehl, A. Birò, A. Siri, P. Orecchia, J. Grassi, D. Neri, L. Zardi, Selective targeted delivery of TNFα to tumor blood vessels, Blood. (2003). doi:10.1182/blood-2003-04-1039.

[32] E. Balza, B. Carnemolla, L. Mortara, P. Castellani, D. Soncini, R.S. Accolla, L. Borsi, Therapy-induced antitumor vaccination in neuroblastomas by the combined targeting of IL-2 and TNFα, Int. J. Cancer. (2010). doi:10.1002/ijc.25018.

[33] E. Balza, L. Mortara, F. Sassi, S. Monteghirfo, B. Carnemolla, P. Castellani, D. Neri, R.S. Accolla, L. Zardi, L. Borsi, Targeted delivery of tumor necrosis factor-α to tumor vessels induces a therapeutic T cell-mediated immune response that protects the host against syngeneic tumors of different histologic origin, Clin. Cancer Res. (2006). doi:10.1158/1078-0432.CCR-05-2448.

[34] C. Halin, V. Gafner, M.E. Villani, L. Borsi, A. Berndt, H. Kosmehl, L. Zardi, D. Neri, Synergistic therapeutic effects of a tumor targeting antibody fragment, fused to

interleukin 12 and to Tumor Necrosis Factor α, Cancer Res. (2003).

[35] E. Balza, S. Zanellato, A. Poggi, D. Reverberi, A. Rubartelli, L. Mortara, The therapeutic T-cell response induced by tumor delivery of TNF and melphalan is dependent on early triggering of natural killer and dendritic cells, Eur. J. Immunol. (2017). doi:10.1002/eji.201646544.

[36] G. Spitaleri, R. Berardi, C. Pierantoni, T. De Pas, C. Noberasco, C. Libbra, R. González-Iglesias, L. Giovannoni, A. Tasciotti, D. Neri, H.D. Menssen, F. De Braud, Phase I/II study of the tumour-targeting human monoclonal antibody-cytokine fusion protein L19-TNF in patients with advanced solid tumours, J. Cancer Res. Clin. Oncol. (2013). doi:10.1007/s00432-012-1327-7.

[37] F. Papadia, V. Basso, R. Patuzzo, A. Maurichi, A. Di Florio, L. Zardi, E. Ventura, R. González-Iglesias, V. Lovato, L. Giovannoni, A. Tasciotti, D. Neri, M. Santinami, H.D. Menssen, F. De Cian, Isolated limb perfusion with the tumor-targeting human monoclonal antibody-cytokine fusion protein L19-TNF plus melphalan and mild hyperthermia in patients with locally advanced extremity melanoma, J. Surg. Oncol. (2013). doi:10.1002/jso.23168.

[38] R. Danielli, R. Patuzzo, A.M. Di Giacomo, G. Gallino, A. Maurichi, A. Di Florio, O. Cutaia, A. Lazzeri, C. Fazio, C. Miracco, L. Giovannoni, G. Elia, D. Neri, M. Maio, M. Santinami, Intralesional administration of L19-IL2/L19-TNF in stage III or stage IVM1a melanoma patients: Results of a phase II study, Cancer Immunol. Immunother. (2015). doi:10.1007/s00262-015-1704-6.

[39] P. Sharma, J.P. Allison, Immune checkpoint targeting in cancer therapy: Toward combination strategies with curative potential, Cell. (2015). doi:10.1016/j.cell.2015.03.030.

[40] S.C. Wei, J.H. Levine, A.P. Cogdill, Y. Zhao, N.A.A.S. Anang, M.C. Andrews, P. Sharma, J. Wang, J.A. Wargo, D. Pe'er, J.P. Allison, Distinct Cellular Mechanisms Underlie Anti-CTLA-4 and Anti-PD-1 Checkpoint Blockade, Cell. (2017). doi:10.1016/j.cell.2017.07.024.

[41] H.L. Kaufman, J. Russell, O. Hamid, S. Bhatia, P. Terheyden, S.P. D'Angelo, K.C. Shih, C. Lebbé, G.P. Linette, M. Milella, I. Brownell, K.D. Lewis, J.H. Lorch, K. Chin, L. Mahnke, A. von Heydebreck, J.M. Cuillerot, P. Nghiem, Avelumab in patients with chemotherapy-refractory metastatic Merkel cell carcinoma: a multicentre, single-group, open-label, phase 2 trial, Lancet Oncol. (2016). doi:10.1016/S1470-2045(16)30364-3.

[42] L. Fehrenbacher, A. Spira, M. Ballinger, M. Kowanetz, J. Vansteenkiste, J. Mazieres,

K. Park, D. Smith, A. Artal-Cortes, C. Lewanski, F. Braiteh, D. Waterkamp, P. He, W. Zou, D.S. Chen, J. Yi, A. Sandler, A. Rittmeyer, Atezolizumab versus docetaxel for patients with previously treated non-small-cell lung cancer (POPLAR): A multicentre, open-label, phase 2 randomised controlled trial, Lancet. (2016). doi:10.1016/S0140-6736(16)00587-0.

[43] C. Robert, J. Schachter, G. V Long, A. Arance, J.J. Grob, L. Mortier, A. Daud, M.S. Carlino, C. McNeil, M. Lotem, J. Larkin, P. Lorigan, B. Neyns, C.U. Blank, O. Hamid, C. Mateus, R. Shapira-Frommer, M. Kosh, H. Zhou, N. Ibrahim, S. Ebbinghaus, A. Ribas, KEYNOTE-006 investigators, Pembrolizumab versus Ipilimumab in Advanced Melanoma., N. Engl. J. Med. (2015). doi:10.1056/NEJMoa1503093.

[44] J. Larkin, V. Chiarion-Sileni, R. Gonzalez, J.J. Grob, C.L. Cowey, C.D. Lao, D. Schadendorf, R. Dummer, M. Smylie, P. Rutkowski, P.F. Ferrucci, A. Hill, J. Wagstaff, M.S. Carlino, J.B. Haanen, M. Maio, I. Marquez-Rodas, G.A. McArthur, P.A. Ascierto, G. V Long, M.K. Callahan, M.A. Postow, K. Grossmann, M. Sznol, B. Dreno, L. Bastholt, A. Yang, L.M. Rollin, C. Horak, F.S. Hodi, J.D. Wolchok, Combined Nivolumab and Ipilimumab or Monotherapy in Untreated Melanoma., N. Engl. J. Med. (2015). doi:10.1056/NEJMoa1504030.

[45] C. Robert, L. Thomas, I. Bondarenko, S. O'Day, J. Weber, C. Garbe, C. Lebbe, J.-F. Baurain, A. Testori, J.-J. Grob, N. Davidson, J. Richards, M. Maio, A. Hauschild, W.H. Miller, P. Gascon, M. Lotem, K. Harmankaya, R. Ibrahim, S. Francis, T.-T. Chen, R. Humphrey, A. Hoos, J.D. Wolchok, Ipilimumab plus dacarbazine for previously untreated metastatic melanoma., N. Engl. J. Med. (2011). doi:10.1056/NEJMoa1104621.

[46] C. Hutmacher, N. Gonzalo Núñez, A.R. Liuzzi, B. Becher, D. Neri, Targeted Delivery of IL2 to the Tumor Stroma Potentiates the Action of Immune Checkpoint Inhibitors by Preferential Activation of NK and CD8 + T Cells , Cancer Immunol. Res. (2019). doi:10.1158/2326-6066.cir-18-0566.

[47] E. Puca, P. Probst, M. Stringhini, P. Murer, G. Pellegrini, S. Cazzamalli, C. Hutmacher, B. Gouyou, S. Wulhfard, M. Matasci, A. Villa, D. Neri, The antibody-based delivery of interleukin-12 to solid tumors boosts NK and CD8+ T cell activity and synergizes with immune checkpoint inhibitors, Int. J. Cancer. (2019). doi:10.1002/ijc.32603.

[48] T. Denayer, T. Stöhrn, M. Van Roy, Animal models in translational medicine: Validation and prediction, New Horizons Transl. Med. (2014).

doi:10.1016/j.nhtm.2014.08.001.

[49]   F. Pretto, G. Elia, N. Castioni, D. Neri, Preclinical evaluation of IL2-based immunocytokines supports their use in combination with dacarbazine, paclitaxel and TNF-based immunotherapy, Cancer Immunol. Immunother. (2014). doi:10.1007/s00262-014-1562-7.

[50]   C. Ratti, L. Botti, V. Cancila, S. Galvan, I. Torselli, C. Garofalo, M.C. Manara, L. Bongiovanni, C.F. Valenti, A. Burocchi, M. Parenza, B. Cappetti, S. Sangaletti, C. Tripodo, K. Scotlandi, M.P. Colombo, C. Chiodoni, Trabectedin overrides osteosarcoma differentiative block and reprograms the tumor immune environment enabling effective combination with immune checkpoint inhibitors, Clin. Cancer Res. (2017). doi:10.1158/1078-0432.CCR-16-3186.

Figure 1

A

B

C

D

Figure 2

Figure captions

Figure 1 Therapeutic activity of L19-TNF in combination with chemotherapeutics (Dacarbazine and Trabectedin) in BALB/c mice bearing WEHI-164 tumors. (**A**) Experimental scheme. (**B**) Treatment started when tumors reached a volume between 70-100 mm^3, mice were injected three times intravenously (black arrows) every 48 hours with either PBS or L19-TNF. Dacarbazine and Trabectedin were administered intraperitoneally once (gray arrow), for hours after the first L19-TNF injection (combination group). Data represent mean tumor volume ± SEM, n = 4 mice per group, CR = complete response. (**C**) Body weight monitoring. Weight change during the therapy of subcutaneous WEHI-164 tumors. Data represent mean % weight (± SEM), n = 4 mice per group. (**D**) Survival curves.

Figure 2 Therapeutic activity of L19-TNF in combination with immune-checkpoint inhibitor (anti-PD1) or chemotherapeutics (Melphalan) in BALB/c mice bearing WEHI-164 tumors. (**A**) Experimental scheme. (**B**) Treatment started when tumors reached a volume between 70-100 mm^3, mice were injected three times intravenously (black arrows) every 48 hours with either PBS, L19-TNF, anti-PD-1 or the combination of L19-TNF plus anti-PD1. Melphalan were also administered intravenously once (gray arrow), both as a monotherapy and in combination with L19-TNF. Data represent mean tumor volume ± SEM, n = 5 mice per group, CR = complete response. (**C**) Body weight monitoring. Weight change during the therapy of subcutaneous WEHI-164 tumors. Data represent mean % weight (± SEM), n = 5 mice per group. (**D**) Survival curves.