Running head: Turanose induced WOX5 restores symbiosis in *cre1*

*Author for correspondence:**

Maitrayee DasGupta

Address: Department of Biochemistry, University of Calcutta,

35, Ballygunge Circular Road, Kolkata-700019, West Bengal, India

Phone No. +91-33-2475-4680; Fax: +91-33-2476-4419

Email ID: maitrayee_d@hotmail.com

Journal Research area:

Signaling and response

17
18

19 **Turanose induced WOX5 restores symbiosis in the *Medicago truncatula* cytokinin**
20 **perception mutant *cre1***
21

Anindya Kundu[# 1], Firoz Molla[#] and Maitrayee DasGupta[*]

Department of Biochemistry, University of Calcutta, Kolkata 700019, India

# Equal contribution by both authors

One sentence summary:

Activation of sugar-WOX5 signaling axis restores root nodule symbiosis in cytokinin perception

mutant *cre1*

Keywords:

Root Nodule Symbiosis, Cytokinin signaling, sugar, WOX5, *MtIAR33*

Footnotes:

*Address correspondence to *maitrayee_d@hotmail.com*

Author contribution:

A.K. M.D. conceived the idea; A.K. and F.M. performed experiments and analyzed the data. All authors wrote and approved the final manuscript.

Financial support:

This work was funded by Grants from Govt. of India: DST (DST EMR/2015/001006). Fellowship to A. K (Council of Scientific and Industrial Research, CSIR-09/028[0756]/2009– EMR–I) and fellowship to F.M (University Grant Commission, No.F.16-6(DEC.2016)/2017 (NET). UGC Ref no- 771

[1]**Present address:** NIAB, Cambridge, UK

The authors declare no conflict of interest.

ABSTRACT

Rhizobia-legume interaction recruits cytokinin-signaling that causes local auxin accumulation for the induction of nodule primordia in the cortex. Since sugar signaling can trigger auxin responses and regulate developmental processes, we explored whether sugar treatments could rescue cre1. Here we demonstrate that turanose, a non-metabolizable sucrose analogue can recover functional symbiosis in cytokinin perception mutant *cre1*. Additionally, turanose significantly upregulated the expression of WUSCHEL-related homeobox 5 (*MtWOX5*) which prompted us to check if ectopic expression of WOX5 could rescue *cre1*. Overexpression of WOX5 from *Arachis hypogaea* (*AhWOX5*), but not the intrinsic *MtWOX5* could completely restore functional symbiosis in *cre1* though both WOX5 (Mt and Ah) were functionally equivalent in inducing the expression of cytokinin inducible transcription factor Nodule Inception (*NIN*). Among the tested markers for cytokinin and auxin responses, significant differences were noted in the expression of IAA-Ala Resistant3 (*MtIAR33*), an auxin conjugate hydrolase. Turanose and *AhWOX5* overexpression resulted in upregulation of *MtIAR33* that further increased significantly in presence of rhizobia. On the other hand, *MtIAR33* expression was unaffected in *MtWOX5* overexpressed roots suggesting deconjugation driven auxin pool to be critical for rescuing symbiosis in *cre1*. We hypothesize a working model for sugar and WOX5 mediated rescue of symbiosis in *cre1*.

INTRODUCTION

In root nodule symbiosis (RNS) rhizobia-legume interaction activates the SYM pathway which in turn activates cytokinin response for the induction of a *de novo* meristem in the root cortical cells (Frugier et al., 2008). Cytokinin-signaling causes local auxin accumulation at the site of

incipient nodule primordia by modulating the expression of auxin transporters (Plet et al., 2011;

Ng et al., 2015). These phytohormonal signals and the SYM pathway together reprogram the

cortical cells and regulate their division, ultimately building a nodule primordium for the

endocytic accommodation of the symbionts (Mathesius et al., 2000; Suzaki et al., 2012).

Several evidences projected the hierarchical organization of the Nod factor-dependent activation

of SYM-Pathway and cytokinin signaling pathway (Frugier et al., 2008). For example, Nod

factor induced accumulation of cytokinin by upregulation of several cytokinin biosynthesis genes

indicates its precedence to cytokinin signaling(Held et al., 2014; van Zeijl et al., 2015). This

explains why the majority of Nod factor-induced transcriptional changes were absent in

cytokinin perception mutant *cre1* in *M. truncatula*(van Zeijl et al., 2015). Cytokinin signaling

through CRE1 (CYTOKININ RESPONSE 1) is required for the induction of NIN, a key

transcription factor required for nodule organogenesis as well as for infection (Gonzalez-Rizzo et

al., 2006; Plet et al., 2011; Vernié et al., 2015; Liu et al., 2019). Cortical activation of cytokinin

signaling that is prerequisite for nodule organogenesis is also dependent on NIN (Vernié et al.,

2015). Genetic evidences further confirm this hierarchy, when constitutive activation of CCaMK

(*snf1*) fails to trigger spontaneous nodulation in absence of cytokinin signaling in *cre1* whereas

constitutive activation of cytokinin receptor LHK1 (*snf2*) triggered nodulation in *dmi2* (Madsen

et al., 2010). Cytokinin oxidase/dehydrogenase expressed during nodule development restricts

the level of active cytokinin for balancing its positive role in nodule organogenesis with its

negative effect on rhizobial infection at the root epidermis (Reid et al., 2016). Therefore, apart

from being the key endogenous signal for nodule development and differentiation (Plet et al.,

2011; Kundu and DasGupta, 2018), cytokinin also maintains the homeostasis of the symbiotic

interaction indicating that the *SYM* pathway and cytokinin signaling are connected through

feedback loops (Miri et al., 2016).

Hierarchically auxin is projected to act next to cytokinin during nodule development (Plet et al.,

2011; Suzaki et al., 2012). Several evidences indicated directional transport and local

accumulation of auxin to be important for the development of nodule primordia and accordingly

application of auxin transport inhibitors is sufficient to induce nodule-like cell division

(Rightmyer and Long, 2011; Ng et al., 2015). Accumulation of auxin could be either due to

regulation of its transport or biosynthesis and cytokinin is known to regulate both these processes

(van Noorden et al., 2006; Pernisová et al., 2009; Jones et al., 2010). The response regulators
(RRs), functioning downstream to cytokinin receptors modulate auxin signaling via SHY2
mediated downregulation of auxin transporters (Ioio et al., 2008; Moubayidin et al., 2010).
Recent evidence revealed that cytokinin controls flavonoid concentration in roots and local
application of flavonoid could recover nodulation in the cytokinin perception mutant *cre1* of *M.*
truncatula (Ng et al., 2015). Though auxin accumulation by its biosynthesis depends on
cytokinin inducible NIN expression (Schiessl et al., 2019), the auxin transport inhibitors can
cause pseudonodule formation in mutants like *nsp2* and *nin* suggesting auxin accumulation to be
driven by diverse upstream cues (Rightmyer and Long, 2011). Overall auxin accumulation and
response appear to be the final call for initiation of cortical cell division during RNS (Ng et al.,
2015).

Sugars play a role in plant development and several evidences indicate a crosstalk between sugar
signaling and auxin responses. For example, (i) there are mutants that are commonly resistant to
both sugar and auxin responses (Ohto et al., 2006), (ii) auxin biosynthetic genes are modulated
by sugar status (LeClere et al., 2010; Sairanen et al., 2012) and (iii) there is a connection
between PAT and sugar metabolism (Stokes et al., 2013). Moreover, sugar signaling through
auxin-based signal transduction is also responsible for specific developmental responses like
trehalose mediated inhibition of root elongation and glucose affecting root architecture (Wingler
et al., 2000; Mishra et al., 2009). Sugar signaling is also connected with auxin response of the
conserved WUS (WUSCHEL)/WOX (WUSCHEL-related homeobox gene)-CLV regulatory
system that are involved in meristem maintenance. For example, WOX9 is typically required for
auxin response in Shoot Apical meristem (SAM) and sucrose can compensate for loss of WOX9
(stip), that arrests growth soon after germination (Wu et al., 2005). Also, the expression of
WOX5, which maintains localized auxin maxima in the root apical meristem (RAM) is induced
by auxin and turanose, a non-metabolizable sucrose analogue (Gonzali et al., 2005).

CRE1, an orthologue of *AtAHK4*, is a membrane-bound cytokinin receptor necessary for
nodulation in *M. truncatula* (Gonzalez-Rizzo et al., 2006; Plet et al., 2011). Earlier reports
indicated that the *M. truncatula cre1* mutant is defective in auxin accumulation in the root cortex
following *S. meliloti* inoculation (Ng et al., 2015). A plausible hypothesis could be that sugar
signaling would rescue cytokinin perception mutant *cre1* by inducing auxin responses. Sucrose is

an important metabolite transported to nodules and sucrose transporters are upregulated with
rhizobial infection but their specific role in signaling during nodule development is still oblivious
(Lalonde et al., 2004; Ayre, 2011). The dual role of sucrose as an energy source and a signaling
molecule is evidenced by many experiments and turanose, a non-metabolizable analogue of
sucrose provides an opportunity to study its role as a signaling molecule (Sinha et al., 2002).
Herein we show that turanose treatment upregulates both cytokinin and auxin responses and
successfully rescues the nodulation efficiency in *cre1*. Turanose treatment induced the
expression of WOX5, and ectopic expression of *WOX5* can also restore symbiosis in *cre1*.

RESULTS

Turanose restores root nodule symbiosis in *M. truncatula* cytokinin perception mutant *cre1*

Based on several evidences that highlight the crosstalk between sugar signaling and auxin
responses we wanted to explore whether sugar treatments could rescue rhizobial symbiosis in
cytokinin perception mutant *cre1* (LeClere et al., 2010; Ljung et al., 2015). Since sugars can act
as energy source, as well as signaling molecules, we disentangled these functions by using
sucrose as well as turanose, a non-metabolizable sucrose which is previously reported to trigger
auxin response in *Arabidopsis* (Gonzali et al., 2005). Both *A17* and *cre1* roots were pre-treated
with sucrose and turanose (10^{-4}M to 10^{-2}M) for 1 week before being infected with *Sinorhizobium*
meliloti Sm2011-pBHR-mRFP. Under these conditions, there was no significant change in root or
shoot length or the lateral root number in either *A17* or *cre1* plants in presence of turanose
whereas sucrose strongly affected both root and shoot growth (Supplementary Fig.1). On the
other hand, while 10^{-3}M turanose treatment could completely recover symbiosis in *cre1*,
treatment with sucrose at the same concentration was unable to restore symbiosis (Fig.1;
Supplementary Table 1). These findings suggested that sucrose act as a signaling molecule rather
than as a metabolizable energy source for the activation of RNS in *cre1*. At 3 weeks after
infection (WAI), there were ~1.4 ± 0.9 nodules in untreated *cre1* roots whereas turanose
treatment resulted in formation of ~12 ± 3.3 nodules per plant (Fig.1A). In untreated A17 there

were ~21 ± 2.9 nodules at 3WAI which increased to ~28 ± 8.7 nodules upon turanose treatment

(Fig.1A). Almost 42% of the turanose rescued nodules in *cre1* were pink nodules (Fig.1B and E),

with proper colonization and differentiation of bacteroids (Fig.1I). For the rest of the nodules that

were white and merged (Fig.1B and F), rhizobia were entrapped in cortical infection threads

(Fig.1J). The increase in nodule number in turanose treated A17 was due to significant increase

in merged white nodules (Fig.1B and H). In these nonfunctional white nodules rhizobia were

entrapped in infection threads in the infection zone as opposed to the proper colonization and

differentiation of bacteroids in the pink functional nodules (Fig.1K-L). It may be noted that

formation of merged nodules was a common feature in turanose treated *cre1* and A17 roots

indicating that turanose induced loss of spatial distribution is independent of CRE1 mediated

cytokinin signaling.

Turanose mediated distribution of cytokinin and auxin response maxima in transgenic

roots of composite *M. truncatula cre1* mutants

To understand the probable mechanism behind turanose mediated rescue of symbiosis in *cre1* we

analyzed the cytokinin and auxin responses at early stages of rhizobial infection. As a readout for

cytokinin response we used *pTCS:GUS,* the cytokinin responsive two-component signaling

(TCS) reporter that uses the target binding sequence of ARR-Bs (Müller and Sheen, 2008;

Zürcher et al., 2013). The *pTCS:GUS* transformed hairy roots of both *A17* and *cre1* were pre-

treated with 10^{-3}M turanose for 1 week before being infected with *Sinorhizobium meliloti* for 24

hours. Histochemical analysis was done for spatial localization of GUS activity where staining

was strictly done for 15 minutes to highlight the noted contrast in presence of turanose. In

uninfected mock treated A17, *pTCS:GUS* was only detectable in meristematic zone (MZ) which

in infected roots spread into the epidermal layer of the both MZ and root hair zone (RHZ)

(Fig.2A-B). Earlier report have also shown a similar activation of *TCSn:GUS* during early stages

of rhizobial invasion, with maximal expression in epidermal and outer cortical cells in the

nodulation-competent RHZ (Jardinaud et al., 2016). Interestingly in turanose treated A17 roots

even in absence of rhizobia, the expression of *pTCS:GUS* was noted in the epidermal layers of

both MZ and RHZ which at 24hpi further extended in the cortex and along the root stele (Fig.2C-

D). Such expansion of *pTCS:GUS* in inner cortical cells is usually noted at later states of
rhizobial infection (Vernié et al., 2015). Under identical conditions *pTCS:GUS* was not
detectable in mock treated *cre1,* but upon infection, it was noted near the outer layers of MZ
(Fig.2E-F). On the other hand, turanose treated *cre1* roots have a significantly enhanced
pTCS:GUS expression in the MZ and in the epidermal and cortical layers of RHZ. Thus
turanose appears to have changed the cytokinin signaling landscape in *M. truncatula* roots and
the increased expression of *pTCS:GUS* in *cre1* roots indicate CRE1 to spatially restrict the level
of cytokinin responses (Fig.2G-H).

As a readout for auxin response we used *pDR5:GUS,* that uses multiple tandem repeats of a
highly active synthetic auxin response element (Ulmasov et al., 1997; Sabatini et al., 1999).
Under identical conditions, in wild type A17 roots *pDR5:GUS* was noted along the stele near the
MZ but not in the RHZ (Fig.2I). This expression increased along the root stele in the RHZ in
infected A17 roots but activity in the root cortex was very low (Fig.2J). Earlier observations also
reported auxin response to be restricted to vascular bundles and pericycle in *M. truncatula* roots
(van Noorden et al., 2007; Demina et al., 2019). Interestingly in turanose treated roots this
increased auxin response along the root stele in RHZ was noted in absence of rhizobia and this
pattern remained unchanged in turanose treated infected roots (Fig.2K-L). In uninfected *cre1*
roots the pattern of auxin response was similar to A17 except that it was also detected in the
RHZ along the stele and the pattern remained similar in the infected roots as well (Fig.2M-N).
However in turanose treated *cre1*, *pDR5:GUS* expression was significantly higher in RHZ with
GUS staining in the epidermal and cortical layers (Fig. 2O). Concomitantly, 1µM IAA treatment
of *M. truncatula* was previously reported to cause a similar auxin response that spread along the
whole root system (van Noorden et al., 2007). Thus, our observations suggest a similar surge of
auxin inside the root upon turanose treatment. Such patches of *pDR5:GUS* expression in *cre1*
were also noted in the cortex of turanose treated infected roots but in a lesser frequency (Fig.2P).
It is well documented that cytokinin signaling through CRE1 is required for the accumulation of
auxin in cortical cells to initiate nodule organogenesis (Ng et al., 2015). Since auxin response is
significantly high in turanose treated *cre1* roots, our results additionally demonstrate that CRE1
has a significant role in restricting the level and spatial spread of the auxin response. In
summary, turanose treatment induced significant cytokinin and auxin responses in the RHZ of
the *cre1* roots and thus indicating a probable cause behind the rescue of nodulation phenotype.

Turanose treatment elevated the expression of *MtWOX5* in *cre1*

The crosstalk of WOX family of homeobox transcription factors with the sugar and phytohormonal signaling networks play a key role in meristem development and maintenance (Wu et al., 2005; Wahl et al., 2010; Kong et al., 2016). Earlier observations indicated *MtWOX5* expression to coincide with the site of nodule primordium formation where the first divisions are initiated in the pericycle, endodermis, and inner cortical cells (Osipova et al., 2011). We were curious to check how the changes in phytohormonal responses in turanose treated roots affected the expression of *MtWOX5* in the early stages of rhizobial infection (Fig.2A-P). Experimental conditions were therefore kept unchanged and *MtWOX5* expression was followed using *pWOX5:GUS* in turanose treated roots in presence (24hpi) and in absence of rhizobia. qRT-PCR estimation revealed *MtWOX5* expression to be elevated in presence of turanose with the effect being more pronounced in *cre1* as compared to A17. This increased expression of *MtWOX5* in *cre1* indicates cytokinin signaling through MtCRE1 to have a role in restricting *MtWOX5* expression in A17 (Supplementary fig.2).

We then analyzed the spatial expression pattern of *pWOX5:GUS* and compared it with cytokinin (*pTCS:GUS*) and auxin (*pDR5:GUS*) responses (Fig.2A-P). In absence of turanose in uninfected roots of both A17 and *cre1*, expression of *pWOX5:GUS* was restricted to root apex (Fig.2Q and U). Upon infection *pWOX5: GUS* spread in the inner layers of the MZ in A17, whereas in *cre1* roots it spread in both outer and inner layers. This indicates that CRE1 has a role in restricting the expression of *MtWOX5* (Fig.2R and V). In turanose treated A17 roots, both in presence or in absence of rhizobia, *pWOX5:GUS* expression expanded to the epidermal layers of RHZ. This highlights an overlap of *pWOX5:GUS* with cytokinin response pattern and suggests a role of cytokinin signaling in turanose mediated transcriptional activation of *MtWOX5* (Fig.2C, S and T). In uninfected roots of turanose treated *cre1*, *pWOX5:GUS* expression was also noted in epidermal layers of RHZ indicating that CRE1 may not have a role in turanose mediated *WOX5* expression (Fig.2W). In turanose treated infected *cre1* roots *pWOX5:GUS* expression was primarily centered towards stele showing an overlap between auxin response, but was also present throughout the cortex that shows overlap with both cytokinin and auxin responses (Fig.2H, P and X). Thus, both cytokinin and auxin responses show significant overlap with

261 *MtWOX5* expression and could be involved together in determining the level and distribution of
262 *MtWOX5* in turanose treated infected roots of *cre1*. Intriguingly, previous reports also indicated
263 *MtWOX5* expression to be a function of both cytokinin and auxin response during the onset of
264 nodule development (Osipova et al., 2012).

Ectopic expression of WOX5 rescues nodulation in *cre1*

Earlier observations have demonstrated that *STIP (WOX9)* overexpression (OE) restores SAM in
cytokinin sensing mutants(Wu et al., 2005). The significant increase in *MtWOX5* expression in
turanose treated *cre1* roots raises the question whether OE-WOX5 would rescue the symbiotic
phenotypes in *cre1*. Phylogenetically WOX5 from determinate nodule forming legumes like
Arachis hypogaea, *Glycine max*, *Vigna angularis*, *Cajanus cajan* and *Phaseolus vulgaris* and
indeterminate nodulators like *Cicer arietinum*, *Pisum sativum*, *Medicago truncatula*and
Trifolium subterraneum distinctly clusters in a distance tree and therefore could be amenable to
distinct regulations (Supplementary fig.3). While acropetal auxin transport inhibition is essential
for indeterminate nodule development, it is dispensable for determinate nodulation (Ng and
Mathesius, 2018). We therefore attempted to rescue *cre1* by OE-*MtWOX5* as well as OE-
AhWOX5 from a determinate legume like *Arachis hypogaea*. *Mt*WOX5 is 86% similar to
*Ah*WOX5 and the homeobox domain is highly conserved with 96% similarity (Supplementary
fig.3).

Full length *MtWOX5* and *AhWOX5* was amplified from cDNA prepared from respective plant
roots. *Mt*WOX5 codes for a protein of 184aa (XP_003616581.1) and *Ah*WOX5 codes for a
protein of 215aa (KT820790). Transformed roots of *cre1* overexpressing *MtWOX5* and *AhWOX5*
were inoculated with *S. meliloti Sm2011-pBHR-mRFP* or *Sm1021-pXLGD4-lacZ*. Infection
threads (ITs) were detected within 2WAI in both empty vector transformed and OE-*MtWOX5*
cre1 roots (Fig.3A, Supplementary Fig.4). But in both the cases IT development was abnormal
(A) where ~75% ITs were stalled in epidermal cortical barrier and ~20% were stalled in
microcolonies in root hair . In both cases, cortical cell divisions were noted but these primordia-
like structures remain uninfected (Supplementary Fig.4C and F). These problems in progress of
infection were previously reported for *cre1* (Plet et al., 2011), which indicates that OE-*MtWOX5*
could not overcome the infection related abnormalities. On the other hand, within 2WAI ~75%

of the ITs in OE-*AhWOX5* roots were normal (N) (Fig.3A). The normal ITs progressed toward

the subtending nodule primordia generated in the cortex to ensure proper rhizobial

accommodation (Supplementary fig.4G-I). It is worth noting that with increased incidences of

normal ITs in OE-*AhWOX5* roots, there was a significant reduction in the total number of ITs

observed in comparison to empty vector or OE-*MtWOX5* roots (Fig.3B). In accordance with the

abnormal progression of ITs there was no improvement of nodulation efficiency in OE-*MtWOX5*

cre1 roots even at 6WAI (n=10) (Fig.3C; Supplementary Table 2). But overexpression of

AhWOX5 led to significant nodulation in *cre1* roots by 4WAI where the total number of nodules

was comparable to wild type A17 roots (Supplementary fig.4; Supplementary Table 2). Almost

75% of these nodules were pink (Fig.3D) and could reduce acetylene (Fig. 3G) with

ultrastructure analysis revealing properly differentiated bacteroids (Fig.3M). Rest of the nodules

that were whitish, resembled the nodules formed in *cre1* or OE-*MtWOX5 cre1* roots, where

bacteria were trapped within a network of infection threads in the infection zone (Fig.3H-M).

The level of expression of *MtWOX5* and *AhWOX5* were comparable in respective OE-*cre1* roots

indicating an additional layer of regulation required for MtWOX5 to become functional in *cre1*

(Fig.3E and F). Since overexpression of neither *AhWOX5* nor *MtWOX5* had any effect in nodule

number or their organization in A17, it is clear that their signaling output is under the

homeostatic control of a large network of fate governing factors (Supplementary Table 2).

MtWOX5 and AhWOX5 overexpression has contrasting effects on expression of auxin and

cytokinin response markers

To understand how OE-*AhWOX5* roots have altered cytokinin and auxin responses at early

stages of infection (24hpi), we checked the expression of several markers as a readout of these

phytohormonal responses in OE-*AhWOX5* roots of both A17 and *cre1*. The same markers were

also evaluated in OE-*MtWOX5* roots and turanose treated roots under otherwise identical

conditions to have possible clues towards understanding the similarities and differences in their

ability to restore symbiosis in *cre1*.

For cytokinin response the expression of *MtCRE1* and its paralogous *MtHK2* and *MtHK3* and

several type-A RRs such as *MtRR4*, *MtRR8* and *MtRR9* that are induced during nodule

organogenesis were monitored (Gonzalez-Rizzo et al., 2006; Tirichine et al., 2007; den Camp et
al., 2011). It was noted earlier that the late and inefficient nodule development in *cre1* is due to
the redundant involvement of receptors *MtHK2* and *MtHK3* (Held et al., 2014; Boivin et al.,
2016). In both A17 and *cre1* at early stages of infection, OE-*WOX5* (*Mt* or *Ah*), had little or no
effect in the level of expression of *MtCRE1*, *MtHK2* and *MtHK3* (Fig.4A-C). Turanose treatment
however had interesting consequences. In A17 roots, turanose downregulated *MtCRE1* but not
MtHK2 and *MtHK3*, suggesting a CRE1 independent cytokinin perception by these paralogues
must be functional in presence of turanose (Fig.4A-C). In *cre1* roots turanose doubled *MtHK2*
expression suggesting *MtHK2* expression could be important in turanose mediated recovery of
symbiosis in *cre1* (Fig. 4B). There were notable differences in the expression of RRs between
OE-*MtWOX5* and OE-*AhWOX5*. Both OE-WOX5 (*Mt* and *Ah*) induced the expression of *MtRR4*,
MtRR8 and *MtRR9* in A17 and *cre1* indicating MtCRE1 independent regulation of their
expression in presence of WOX5 (Fig.4D-F). Upon infection, expression of all 3 RRs
significantly increased in A17 but in *cre1* their expression was differentially affected. For
example, the level of *MtRR8* further increased upon infection but *MtRR9* was unaffected
(Fig.4E-F). These effects were however similar in both OE-*MtWOX5* and OE-*AhWOX5*. On the
other hand, *MtRR4* expression in infected *cre1* roots was downregulated in OE-*MtWOX5* but
significantly upregulated in OE-*AhWOX5* (Fig.4D). This contrast in regulation of *MtRR4*
expression in OE-*MtWOX5* and OE-*AhWOX5* might be crucial in determining the difference in
their ability to rescue *cre1*. Additionally, similar to OE-*AhWOX5* the turanose treated roots also
had increased expression of *MtRR4* in both A17 and *cre1* whereas expression of *MtRR8* and
MtRR9 was unaffected (Fig.4D-F). Thus, for rescue of symbiosis turanose could use MtHK2 as a
proxy for MtCRE1, whereas WOX5 might directly target the downstream RRs. This possibility
is corroborated by an earlier report which shows that WOX proteins can directly regulate the
expression of Type-A RR genes (Zhao et al., 2009).

In the early stages of rhizobial infection cytokinin-inducible transcription factor NIN activates
LBD16 to promote local accumulation of auxin by triggering its biosynthesis (Schiessl et al.,
2019). RNAseq analysis in *cre1* shows that *MtLBD16* and *MtYUCCA2/8* are very feebly
expressed in the mutant indicating a role of CRE1 in inducing auxin biosynthesis. We therefore
monitored the expression of *MtNIN*, along with *MtLBD16* and *MtYUCCA*2/8 as a readout of the
auxin response (Fig.5). In both A17 and *cre1*, OE-*MtWOX5* and OE-*AhWOX5* upregulated the

expression of *MtNIN* (~5-10 fold) which further increased with infection (~15-20 fold) (Fig.5A).

Also, in both A17 and *cre1*, turanose treatment significantly upregulated *MtNIN* expression in

infected roots (~10-25 fold). Thus, till the expression of *MtNIN* the response of *cre1* for OE-

MtWOX5 and OE-*AhWOX5* was identical. As noted earlier, *MtLBD16* expression was very low

in *cre1* mutants and turanose has no effect on *MtLBD16* expression in either A17 or *cre1* roots

making it unlikely for *MtLBD16* to participate in the Turanose-WOX5 signaling axis (Fig.5B).

On the other hand, both OE-*MtWOX5* and OE-*AhWOX5* led to increased expression of *MtLBD16*

(~4-5 fold) in *cre1* but not in A17, indicating WOX5 mediated induction of *MtLBD16* to be

CRE1 independent and also subject to CRE1 mediated inhibition. Upon infection, only in OE-

AhWOX5 roots there was significant increase in expression of *MtLBD16*. This increase was again

significantly more in *cre1* as compared to A17 confirming *AhWOX5* to activate *MtLBD16* in a

CRE1 independent pathway (Fig.5B). Similar to *MtLBD16*, turanose treatment had no effect on

expression of either *MtYUCCA2* or *MtYUCCA8* indicating that turanose signaling does not

directly recruit auxin biosynthesis gene to induce the auxin response (Fig.5C-D). *MtYUCCA2*

expression was unaffected by OE-*MtWOX5*, but *MtYUCCA8* expression was upregulated

similarly in both OE-*MtWOX5* and OE-*AhWOX5*, and in both A17 and *cre1* indicating a CRE1

independent pathway to activate WOX5 dependent *MtYUCCA8* expression (Fig.5C-D). Local

auxin accumulation can also happen by regulating free auxin availability through auxin

deconjugation or by affecting auxin efflux transporters (Plet et al., 2011; Deinum et al., 2012).

For deconjugation, we chose to monitor the expression of *MtIAR33* as it is evidenced to be

highly expressed during nodule organogenesis (Campanella et al., 2008). Unlike in OE-

MtWOX5, OE-*AhWOX5* resulted in ~10-fold increase in *MtIAR33* expression under uninoculated

condition. The *MtIAR33* expression further spiked to ~50-fold post infection in OE-*AhWOX5* in

both A17 and *cre1* indicating that its induction under this condition is independent of CRE1

(Fig.5E). Similarly, turanose treatment also resulted in ~150-fold increase in *MtIAR3* expression

in infected roots of *cre1* as compared to A17 (~50-fold) suggesting CRE1 might be playing a

role in restricting turanose dependent *MtIAR3* expression. Genetic evidence supports the fact that

acropetal auxin transport mediated by MtPIN4 plays a very significant role in increasing local

auxin concentration during nodule organogenesis (Plet et al., 2011; Ng et al., 2015). However,

neither *WOX5* overexpression nor turanose treatment had any effect on *MtPIN4* expression

suggesting auxin transport may not be critical in turanose or WOX5 mediated effects (Fig.5F). It

is worth mentioning that *MtPIN* expression is not always in concert with the actual auxin
transport capacity during nodule development (Plet et al., 2011; Ng et al., 2015).

DISCUSSION

The symbiotic inefficiency in *cre1* is due to its inability to induce the synthesis of flavonoids that
regulate auxin transport for reactivation of cortical cells (Ng et al., 2015). In accordance,
treatment with flavonoids allowed auxin transport control and rescued nodulation efficiency in
cre1. Herein we demonstrate that (i) turanose treatment restores symbiosis in *cre1* (Fig.1). While
sugar signaling is known to control several distinct aspects of plant development our results for
the first time show its role in RNS. (ii) Turanose induces the expression of *MtWOX5* and ectopic
expression of *AhWOX5*, but not *MtWOX5* could restore symbiosis in *cre1* (Fig.2-3). (iii) Both
turanose and OE-*AhWOX5* mediated rescue of symbiosis in *cre1* is associated with significant
increase in the expression of an auxin conjugate hydrolase *MtIAR33* at early stages of rhizobial
infection. As hydrolysis of auxin conjugates are known to be critical in determining free auxin
availability (Rampey et al., 2004), we propose this increase in expression of *MtIAR33*
downstream to turanose-WOX5 signaling pathway to be responsible for the restoration of
symbiosis in cre1 (Fig. 5). Based on our observations we propose a working model for turanose
and WOX5 mediated rescue of symbiosis in *cre1* (Fig.6).

Several evidences indicate a connection between sucrose signaling, the conserved WOX-CLV
regulatory system, and the phytohormonal network that are involved in meristem maintenance
(Francis and Halford, 2006). STIMPY (STIP; WOX9), a homeodomain transcription factor
promotes WUS expression in the shoot meristem (Wu et al., 2005). Cytokinin signaling is
required for *AtWOX9* expression (Skylar et al., 2010) and sucrose treatment could rescue *stip*
(AtWOX9) mutants in *Arabidopsis* which has defects in SAM illustrating a cytokinin-WOX-
sucrose module working in concert to control complex developmental processes (Wu et al.,
2005). Accordingly in cytokinin receptor mutants, ectopic expression of *AtWOX9* could also
restore establishment of SAM (Skylar et al., 2010). Similar example is restoration of shoot
regenerative capacity in the type-B response regulator (*arr1* and *arr12)* mutants by ectopic
expression of WUS (Meng et al., 2017). Overexpression of Fantastic 4 (*AtFAF4*) that controls

meristem size by repressing WUS expression causes root growth arrest after germination and this

aberration can also be rescued by sucrose treatment (Wahl et al., 2010). Our results further

exemplify such sugar-WOX-phytohormone signaling axis by demonstrating turanose and WOX5

dependent recovery of RNS in a cytokinin perception mutant *cre1*. These examples point to a

deep conservation of signaling for developing and sustenance of meristems with a direct link

between WOX, sugar and phytohormonal networks (Gonzali et al., 2005; Wu et al., 2005; Wahl

et al., 2010).

The homeobox transcription factors of WOX family are common regulators of cell proliferation

and differentiation (Richards et al., 2015). The common mode of action for WOX5 proteins is

prevention of premature differentiation by transcriptional repression and thereby maintaining a

stem cell niche in meristems(Sarkar et al., 2007). The complete rescue of symbiosis in *cre1* by

AhWOX5 but not the intrinsic *MtWOX5* was intriguing because the reverse was actually expected

(Fig.3; Supplementary table 2). This difference is possibly due to variation in post transcriptional

or post translational regulations because (i) the transcript levels of both *WOX5* was similar in

respective OE roots (Fig. 3E-F) and (ii) there was a significant overlap in the downstream

responses induced by both the WOX5. Overexpression of WOX5 (*Mt* or *Ah*) had no effect on

nodulation efficiency in wildtype background (A17) which further confirms WOX5 to be tightly

controlled at post transcriptional level (Fig.3C; Supplementary Table 2). The homeobox domain

is highly conserved between *Mt*WOX5 and *Ah*WOX5 with 96% similarity but there are notable

features in AhWOX5 that distinguishes it from MtWOX5 for example (i) AhWOX5 has a 23

amino acid leader sequence in the N-terminus which is not present in MtWOX5 (Supplementary

Fig. 3). This is important because intercellular mobility of WOX5 is important for stem-cell

maintenance activity in roots (Pi et al., 2015) (ii) The WUS box motif is essential for WOX5 to

act as transcriptional activator or repressor depending on developmental context and is absolutely

essential for stem cell maintenance and interaction with TPL/TPR proteins (Leibfried et al.,

2005; Ikeda et al., 2009; Dolzblasz et al., 2016). This motif is different between AhWOX5

(TL**Q**LFP) and MtWOX5 (TL**E**LFP). Earlier reports indicate substitutions in WUS box can

completely alter the function of WOX proteins (Lin et al., 2013) (iii) AhWOX5 and MtWOX5

could be differentially regulated by the non-conserved regions in WOX proteins, as its essential

role in determining the mobility and stability was previously noted in *Arabidopsis* (Daum et al.,

2014). At present it is not clear which of these features of AhWOX5 are actually critical in the

recovery of symbiosis in *cre1*, but the difference between MtWOX5 and AhWOX5 could

emerge to be a model for understanding WOX5 action during nodule organogenesis in *M.*

truncatula.

Nodule organogenesis in response to rhizobia is dependent on the cytokinin receptor CRE1 and

the cytokinin-inducible transcription factor NIN (Gonzalez-Rizzo et al., 2006; Marsh et al., 2007;

Plet et al., 2011; Schiessl et al., 2019). Interestingly, though ectopic expression of *MtWOX5*

failed to restore symbiosis in *cre1*, it was functionally equivalent to *AhWOX5* in inducing *MtNIN*

in both uninfected and infected roots of A17 and *cre1* (Fig.5A). This MtCRE1 and rhizobia

independent mode of induction of *MtNIN* has not been noted earlier and highlights another layer

to NIN regulation apart from the classical cytokinin-CRE1-NIN module that governs rhizobial

symbiosis (Fig.6) (Laffont et al., 2020). Involvement of the paralogs *MtHK2/HK3* may be an

unlikely option since WOX5 (*Mt* and *Ah*) expression has no effect on their expression (Fig.4B-

C). Rather WOX5 might be directly targeting the response regulators downstream to CRE1 as

previously observed in *Arabidopsis* SAM (Skylar and Wu, 2011). *MtNIN* expression also

increased significantly by turanose treatment in both A17 and *cre1*, but only in infected roots

(Fig.5A). Thus, turanose dependent *MtWOX5* is unable to trigger *MtNIN* expression in absence

of rhizobia, which is unlike OE-WOX5 (*Mt* and *Ah*) roots where *MtNIN* expression increased in

uninfected roots as well. Interestingly, we noted a significant inhibition of *MtCRE1* expression in

turanose treated A17 roots and therefore turanose mediated *MtNIN* expression was MtCRE1

independent but absolutely dependent on upstream activation of SYM Pathway. These complex

responses of *MtNIN* expression are in tune to distinct *cis*-regulatory sequences in NIN promoter

that guide its complex spatio-temporal expression during symbiosis (Liu et al., 2019). Finally,

just as turanose inhibited *MtCRE1* expression, the *pTCS:GUS* expression was always higher in

turanose treated *cre1* highlighting a reciprocity in sugar and *MtCRE1* signaling (Fig.2G-H). Our

results therefore suggest an alternate MtCRE1 independent module for *MtNIN* expression that is

mediated through sugar and WOX5 and that reciprocally regulate the classical cytokinin-CRE1-

NIN module (Fig. 6). Similar reciprocity between cytokinin and sugar signaling is noted earlier

where plants with impaired cytokinin receptors display increased sugar sensitivity (Moore et al.,

2003) and plants with decreased sensitivity to sugar was accompanied with increased cytokinin

response (Franco-Zorrilla et al., 2005), suggesting a conserved relationship between sugar and

cytokinin across plant species.

Both cytokinin and auxin responses are dependent on NIN action during symbiosis (Heckmann

et al., 2011; Plet et al., 2011; Ng et al., 2015; Schiessl et al., 2019). Interestingly, though OE-

MtWOX5 and OE-*AhWOX5* was functionally equivalent in inducing *MtNIN*, there were distinct

differences in the cytokinin and auxin responses that explains the difference in their ability to

restore symbiosis in *cre1*. For example, the contrast in expression of *MtRR4*, a type-A response

regulator could be critical. During symbiosis *MtRR4* expression depends on NIN and is

associated with cortical cell division during nodule meristem formation (Vernié et al., 2015). In

both OE-*AhWOX5* and OE-*MtWOX5*, expression of *MtRR4* increased in uninfected roots of A17

and *cre1* indicating a CRE1 independent pathway to induce the expression. In infected OE-

AhWOX5 roots expression of *MtRR4* further increased (Fig.4D). But in infected roots of OE-

MtWOX5, the expression of *MtRR4* significantly decreased specifically in *cre1* roots but not in

A17 indicating MtCRE1 signaling to be essential for *MtRR4* expression in infected roots. Earlier

reports have shown WOX proteins to directly regulate the expression of Type-A RRs genes

(Zhao et al., 2009) and the distinct differences between AhWOX5 and MtWOX5 could be the

reason behind the differences in *MtRR4* expression.

Auxin accumulation (*pDR5:GUS* expression) depends on NIN, as it does not occur in a *nin* null

mutant (Suzaki et al., 2012). Downstream to NIN, the genes that regulate auxin availability was

differentially affected by WOX5 (*Ah* and *Mt*) and turanose (Fig.5). The most distinctive was the

expression of an auxin conjugate hydrolase *MtIAR33* that significantly increased in OE-*AhWOX5*

and turanose treated roots indicating a WOX5 dependent pathway to upregulate *MtIAR33*.

Expression of *MtIAR33* in OE-*AhWOX5* roots or turanose treated roots was equivalent or more in

cre1 as compared to A17 indicating a CRE1 independent pathway to induce *MtIAR33* expression

(Fig.5E). Again auxin response (*pDR5:GUS*) in turanose treated *cre1* roots was significantly

higher than auxin response in A17 roots indicating CRE1 restricts sugar mediated auxin response

(Fig.2K and O). Taken together these results prompt us to hypothesize that the turanose-WOX5

signaling axis primarily functioned through *MtNIN* and *MtIAR33*, and hydrolysis of auxin

conjugates was preferentially used by this axis in determining the bioactive pool of auxin for

sugar mediated restoration of symbiosis in *cre1*. There was no change in expression of *MtIAR33*

in OE-*MtWOX5* indicating the importance of deconjugation of auxin in restoration of symbiosis

in *cre1*. Thus, apart from auxin transport (Rightmyer and Long, 2011; Ng et al., 2015), and

biosynthesis (Schiessl et al., 2019), that functions in an MtCRE1 dependent manner, our results

suggest a CRE1 independent auxin deconjugation to be a potential pathway for auxin
accumulation.

Several earlier evidences indicate auxin responsive WOX5 expression (Ding and Friml, 2010;
Osipova et al., 2012) whereas in our results WOX5 dependent IAR33 expression suggest auxin
accumulation to happen downstream to WOX5. This is similar to a previous report where
turanose treatment inhibited auxin conjugation to increase free auxin content but failed to do so
in turanose insensitive (*tin*) loss of function *wox5* mutants thereby indicating WOX5 to mediate
the accumulation of bioactive auxin pool (Gonzali et al., 2005). It is possible that WOX5
mediates a positive feedback loop for rapid auxin response (Fig.6). We envisage this positive
loop to involve NIN because a WOX-NIN crosstalk would mean a cross talk between a sugar
sensing and a nitrogen sensing network being connected during inception of symbiosis. It may
be noted that *MtIAR33* was significantly upregulated during early stages of rhizobial infection in
M. truncatula (Campanella et al., 2008). However, there was no upregulation of *MtIAR33* in our
experimental conditions in absence of turanose and the difference is most likely due to presence
of β-aminobutyric acid (BABA) in their nodulation media that is known for having priming
effects on plants against stress responses (Rejeb et al., 2018). In analogy, it is likely that turanose
treatment and *AhWOX5* expression actually primed the *cre1* roots with increased expression of
MtIAR33 that developed a CRE1 independent robust auxin response and thereby restored
symbiosis in *cre1*. Incidence of merged nodules which is an indication of higher auxin
concentration was only observed in presence of turanose which could be due to increased
expression of *MtIAR33* in presence of turanose or a better homeostatic control over auxin
availability in presence of *AhWOX5* (Penmetsa et al., 2003; van Noorden et al., 2007; Suzaki et
al., 2012). For example, OE-*AhWOX5*, but not the turanose treated roots have increased
expression of *MtLBD16* and *MtYUCCA8* and might therefore have better control over auxin
response that led to higher efficacy of restoration of symbiosis in *cre1* than turanose treated
plants that recruited only auxin deconjugation (*MtIAR33*).

In a working model we propose how turanose-wox5 signaling might have restored symbiosis
in *cre1* (black arrow) in the backdrop of the present understanding about cytokinin signaling
(grey arrows) during nodule initiation in *M. truncatula* (Fig.6). We propose a turanose
dependent WOX5 expression to turn on feed forward positive loop(s) to increase cytokinin and

auxin response and NIN expression. This turanose-WOX5 signaling axis is CRE1 independent
and reciprocally regulates the classical cytokinin-CRE1-NIN module. Future work is needed to
understand the basis of differential action of MtWOX5 and AhWOX5 and the role of NIN in
deconjugation mediated auxin accumulation. We also need to better understand the sugar
perception and signaling network to understand its role in nodule meristem development.

MATERIALS AND METHODS

Plant and rhizobial strains

Medicago cre1 seeds (Plet et al., 2011), *Agrobacterium rhizogenes* strain MSU440 and
Sinorhizobium meliloti Sm2011-pBHR-mRFP and *Sm1021-pXLGD4-lacZ* (Boivin et al., 1990)
strains were used.

Medicago growth conditions:

Medicago was grown as described in (Garcia et al., 2006). In brief, seeds were scarified for 10min in
H_2SO_4, rinsed with sterile water, followed by surface sterilization with 6% sodium hypochlorite
for 3 min containing a few drops of Tween-20 and then again rinsed with sterile water. Surface-
sterilized seeds were imbibed with sterile water with gentle rotation for 5 to 7h at ambient
temperature on a rotating shaker and then stored in water at 4^0 C overnight. Seeds were then placed
on inverted plates containing moist filter papers and then left in dark at 20^0Cfor 36 h for
germination. Germinated seedlings were placed in Fahraeus (Catoira et al., 2000) agar (0.8%)
medium in square petri dishes (12×12 cm) and maintained in a growth chamberat 20ºC on a 16-
h-light and 8-h-dark schedule under 100-110 μmol/m^2/s.

Turanose and sucrose treatment

Plants are germinated as previously described by (Saha et al., 2014) followed by their transfer
into Fahraeus medium containing different concentration of turanose and sucrose:10^{-4}M, 10^{-3}M
and 10^{-2}M for 1 week. For promoter assay plants are maintained for 1 week in Turanose plates
before infected with *Sm2011-pBHR-mRFP* for 24 hours then plants are harvested and stained. For
nodulation assay plants are inoculated in the plates with *Sm2011-pBHR-mRFP/ Sm1021-*
pXLGD4-lacZ grown overnight at 28ºC in YM to OD600=1.0 and then diluted 1:50 in half-

strength B&D (Broughton and Dilworth, 1971). From this diluted bacterial culture 2ml/plate was

used for inoculation. Nodulation is scored 3WAI.

Constructs

Full length MtWOX5 and AhWOX5 are amplified from cDNA prepared from nodulated roots of

Medicago truncatula and *Arachis hypogaea* (Kundu and DasGupta, 2018) with primers 5'-

CACCATGGAAGAGAGCATGTCAGG-3' and 5'-ACTTACGGTTGAGTTTTGTGTAA-3',

and (*MtWOX5*) 5'-CACCATGCAGACGGTCCGAGATCTGTC-3' and 5'-

CCTTCGCTTAAGTTTCATGTAA-3' (*AhWOX5*) and cloned into pENTR-dTOPO

(Invitrogen). The entry clones of *AhWOX5 & MtWOX5* recombined through LR clonase

Gateway technology (Invitrogen) into *pK7WGF2* to generate *p35S::eGFP-AhWOX5*

p35S::eGFP-MtWOX5 respectively. *pDR5:GUS*, *pWOX5:GUS* and *pTCS:GUS* were obtained

from (Franssen et al., 2015).

Induction of hairy roots and generation of composite *M. truncatula* plants and scoring nodulation

Seedlings were transformed with indicated constructs in *Agrobacterium rhizogenes* MSU440 by

following a standard procedure (Boisson-Dernier et al., 2001). Briefly, when seedlings had a radicle

length of approximately 10 mm, the radicle was sectioned approximately 3 mm from the root tip

with a sterile scalpel. Sectioned radicles were inoculated by coating the freshly cutsurface with *A.*

rhizogenes grown on TY solid medium. Thereafter, the inoculated sectioned seedlings were

placed on slanted agar (Sigma) in square petri dishes containing Fahraeus medium with

kanamycin at 25 mg/liter and maintained at 20ºC. Within 3-4 weeks, transgenic hairy roots were

screened for GFP fluorescence by Leica stereo fluorescence microscope M205FA equipped with a

Leica DFC310FX digital camera (Leica Microsystems). The selected composite transgenic plants

were transplanted in vermiculite pots and the plants were maintained in a growth chamber at

20ºC on a 16-h-lightand 8-h-dark schedule under 100-110 µmols/m^2/s. For nodulation assay

plants are inoculated with *Sm2011-pBHR-mRFP*/ *Sm1021-pXLGD4-lacZ* grown overnight at

28ºC in YM to OD600=1.0 and then diluted 1:50 in half-strength B&D (Broughton and

Dilworth, 1971). From this diluted bacterial culture 2ml/pot was used for inoculation. Nodulation

was scored 4 and 6WAI with *S. meliloti* expressing mRFP (Smit et al., 2005) . For observing ITs,

plants were harvested 2WAI.

Acetylene reduction assay

The nitrogen-fixing activity of the nodules was examined indirectly by Acetylene Reduction

Assay as described by (Koch and Evans, 1966) using gas chromatography, the bacterial

nitrogenase activity of the nodules at 6WAI with *Sm2011-pBHR-mRFP* was measured. The roots

were placed into 10mL glass vials sealed with rubber septa, and 200μL of acetylene was injected

into each vial, incubated for 90 min at room temperature, and ethylene production was measured

using Clarus 460 PerkinElmer Gas Chromatograph. The total nitrogenase activity was calculated

as ARA units (nanomoles of ethylene. Hour^{-1} mg nodule^{-1}). Three independent biological

replicates were performed, with n>4 plants analyzed per replicate. Nodules from individual plant

root systems were then pulled out and weighted to determine specific nitrogenase activity (ARA

units per mg of nodule).

Phenotypic analysis, histochemical staining and confocal microscopy

Images of whole mount nodulated roots were captured using a Leica stereo fluorescence

microscope M205FA equipped with a Leica DFC310FX digital camera (Leica Microsystems).

Histological assay for checking GUS expression was performed according to Saha et. al.,2014 with a

short staining period. Briefly, roots were washed with 0.1 M potassium phosphate buffer (pH 7.0)

and immersed and incubated in the dark in staining solution 1 mM 5-bromo-4-chloro-3-indolyl-β-

D-glucuronicacid (X-Gluc, Thermo Scientific), 50mM sodium phosphate buffer, 0.5mM

potassium ferrocyanide, 0.5mM potassium ferricyanide, 20% methanol and 0.1%Triton X-100

for 15 minutes at 37ºC. After rinsing in phosphate buffer and cleaning in 70% ethanol, the roots

were analyzed and imaged using Leica stereo fluorescence microscope M205FA (Leica

Microsystems) and Olympus fluorescence microscope (IX71, Japan). Sectioning of Gus stained

roots was done using cryo microtome (Leica Microsystem) and sliced into 40-μm thick sections and

imaged using Olympus fluorescence microscope (IX71, Japan). Detached nodules were embedded

in 5% low melting point agarose and sliced into 30-μm thick sections with a rotary microtome

RM2235 (Leica Microsystems). Roots were also stained with X-Gal (Affymetrix) for lacZ-

activity, asdescribed in (Arrighi et al., 2008) and imaged under an Olympus IX71 microscope

(Japan). For confocal microscopy, sample preparation was done according to Haynes and
associates (Haynes et al., 2004). Images were acquired with a Leica TCS SP5 II AOBS confocal
laser scanning microscope (Leica Microsystems). All digital micrographs wereprocessed using
Adobe Photoshop CS6.

Real time PCR:

Superscript III RT (Life Technologies) and anchored dT17 primers as described in the
manufacturer's protocol. 1μl of cDNA was used to set up real time qRT-PCR in a 20μl reaction
system with 2X SYBR Green PCR master mix (Life Technologies). Experimental setup and
execution were conducted using an ABI 7500 Fast Real-Time PCR system. PCR program: 1cycle
at 50°C, 1cycle at 95°C for 5 min, 40 cycles at 95°C for 30s and 52°C for 30s. Expression data were
obtained from three independent biological repetitions. Calculations were done using the cycle
threshold method using *MtACTIN2* as the endogenous control. All primers including the genes
used for normalization are given in (Supplementary Table 3). Statistical significance was
determined based on Mann-Whitney test.

Accession number

Nucleotide sequence data for the genes used in thisarticle can be found in the GenBank data
library under the following accession numbers: MtCRE1, XM_024773926.1;*Mt*HK2,
XP_003617960.1; *Mt*HK3, XP_003601762.1; MtRR4, XM_003613375.3; MtRR8,
003608889.3; MtRR9, XM_024778303.1;*Mt*NIN, FJ719774; MtLBD16, XM_013594406.2;
MtYUCCA2, XM_013597647.2; MtYUCCA8, XM_003625408; MtIAR33, DQ489994;
MtPIN4, AY115839.1;*Mt*ACTIN2, JQ02873.

ACKNOWLEDGEMENT

We thank Florian Frugier for *cre1* seeds; Henk J. Franssen for providing us with *pDR5:GUS*,
pTCS:GUS and *pWOX5:GUS* constructs; Ton Bisseling & Erik Limpens for *S. meliloti* harboring
pBHR-mRFP; Douglas R. Cook for *Agrobacterium* strain MSU440.

SUPPLEMENTAL DATA

Supplementary Table 1: Nodule number in A17 and *cre1* upon turanose and sucrose treatment.

Supplementary Table 2: Nodule number in A17and *cre1* by overexpression of *AhWOX5* and
MtWOX5.

Supplementary Table 3: List of primers used for cloning and qRT-PCR.

Supplementary figure 1: Effect of Turanose and Sucrose treatment on Nodulation in A17 and
cre1.

Supplementary figure 2: *MtWOX5* expression in response to turanose treatment.

Supplementary figure 3: Sequence alignment and phylogenetic tree of WOX5.

Supplementary figure 4: IT morphology and nodule number under WOX5 overexpression.

Supplementary figure 5: Relative expression of *MtWOX5* and *AhWOX5* in A17.

FIGURE LEGENDS

Figure 1: Turanose mediated recovery of symbiosis in *cre1* mutants. **(A-B)** Box plot
represents (A) total nodule number per plant and (B) total white (white box) and pink nodules
(red box) under control (mock) and turanose treatment (10^{-3}M) in A17 and *cre1* after infection
with *Sinorhizobiummeliloti Sm2011-pBHR-mRFP*. Number above each box in (A) indicates the
number of root systems with nodules per total number of scored roots. Mann-Whitney test were
used to assess significant differences, where ***, ** and * indicates P<0.0005, <0.002 and <0.02
respectively. **(C-H)** Morphology of infection event or nodules in *cre1* and A17 3WAI under (C-

D) control condition and (E-H) after turanose treatment where (E and G) solitary pink nodules

and (F and H) merged white nodules. Representative micrograph of control and turanose treated

nodulated roots indicated as bright field and bright field + mRFP merged. **(I-L)** Ultrastructure

analysis of nodules in *cre1* and A17 after turanose treatment, where the perforated box represents

the infection zone with its corresponding enlarged view below the respective images. Total

number of each type of nodules observed out of the total observed nodules are indicated for each

panel. (upper panel) Brightfield + mRFP merged and (lower panel) mRFP. Scale bar: C-H=

500µm, I-L (upper panel) = 100µm and (lower panel) = 10µm.

Figure 2: Effect of turanose treatment on *pTCS:GUS*, *pDR5:GUS* and *pWOX5:GUS*

expression pattern in A17 and *cre1* roots. (A-H) *pTCS:GUS* expression in (A-D) A17 and (E-

H) *cre1* roots,(I-P) *pDR5:GUS* expression in (I-L) A17 and (M-P) *cre1* roots and (Q-X)

pWOX5:GUS expression in (Q-T) A17 and (U-X) *cre1* roots under following conditions. **(A-H)**

mock treated uninfected (A, E), infected (B, F), turanose treated uninfected (C, G) and infected

(D, H). **(I-P)** mock treated uninfected (I, M), infected (J, N), turanose treated uninfected (K, O)

and infected (L, P). **(Q-X)** mock treated uninfected (Q, U), infected(R, V), turanose treated

uninfected (S, W) and infected (T, X). All plants were mock or Turanose (10^{-3}M) treated for 7

days followed by infection with *Sinorhizobium meliloti Sm2011-pBHR-mRFP*for 24 hours before

being harvested for GUS staining. Every root image has its corresponding longitudinal section of

the root tip and the root hair zone (box). At least 5 independent roots were analyzed for each

condition from three independent experiments. Scale bars = 300 µm (left panel single root),

Scale bar = 50µm (right panel upper and lower longitudinal section).

Figure 3: Rescue of symbiosis in *cre1* by ectopic expression of *AhWOX5*. (A) Percentile

projection of epidermal infection thread (IT) observed 2WAI with *Sm1021-pXLGD4-lacZ* in

cre1 roots transformed with empty vector, *p35S::eGFP-MtWOX5* and *p35S::eGFP-AhWOX5*.

Each of the infection events has been roughly classified into 4 categories: (i) successful cortical

invasion and intracellular colonization (ii) Stalled in nodule apex, (iii) stalled in epidermal

cortical barrier and (iv) stalled in microcolonies in root hair. Category (i) is considered normal

whereas Category (ii-iv) is considered as abnormality.**(B)** Box plot representing cumulative
number of normal and abnormal ITs. **(C-D)** Box plot represents (C) total nodule number per
plant and (D) total white (white box) and pink nodules (red box) per plant in empty vector,
p35S::eGFP-MtWOX5 and *p35S::eGFP-AhWOX5* transformed hairy-root systems in A17 and
cre1 respectively, harvested 6WAI with *Sinorhizobiummeliloti Sm2011-pBHR-mRFP*. n
indicates the number of root systems with nodules per total number of scored roots. Mann-
Whitney test was used to assess significant differences, where ***, ** and * indicates P<0.0005,
0.002 and <0.02 respectively.**(E-F)** qRT-PCR analysis of (E) *MtWOX5* and (F) *AhWOX5* relative
to empty vector transformed (*cre1*) roots normalized against *MtActin*. **(G)** Acetylene reduction
assay (nmole C_2H_4/h/mg nodule) in indicated root system. (E-G) Histogram represent an average
of three biological replicates each having n>4 plants and error bar represents SD. Mann-Whitney
test was used to assess significant differences, where ***, ** and * indicates P<0.0005, 0.002
and <0.02 respectively.**(H-M)** Morphology of nodules in transgenic hairy-roots of *cre1* mutant
4WAI with *Sm2011-pBHR-mRFP* (H, J and L). Representative micrograph of transgenic roots
indicated as bright field + mRFP merged and perforated box is represented as eGFP + mRFP
merged and bright field. Scale bar = 1mm and box = 500µm. (I, K and M) Ultrastructure of
nodules developed in *cre1* mutant transformed with different constructs indicated as bright field+
mRFP merged where the perforated box in the infection zone is enlarged separately. Scale bar: I,
K and M = 100µm and box = 10µm. Arrow indicate infection thread.

Figure 4: Expression analysis of cytokinin responsive genes under turanose treatment and
WOX5 overexpression in A17 and *cre1*. Quantitative reverse transcriptase-PCR analysis of
indicated genes in A17 and *cre1* roots **(A, B, C, D, E, F)** treated with turanose (10^{-3}M) and water
(mock) followed by infection with *Sinorhizobiummeliloti Sm2011-pBHR-mRFP*relative to mock
treated uninoculated control or transformed with empty vector, *p35S::AhWOX5* and
p35S::MtWOX5 followed by infection with *Sm2011-pBHR-mRFP* relative to empty vector
transformed uninoculated control. *MtActin* was used as a reference gene. Histogram represents
an average of three biological replicates each having n>3 plants and error bar represent standard
deviation (SD). Mann–Whitney test was used to access significant differences where ***, ** and
* represents P < 0.001, <0.01 and <0.05 respectively.

Figure 5: Expression analysis of auxin responsive genes under turanose treatment and WOX5 overexpression in A17 and *cre1*. Quantitative reverse transcriptase-PCR analysis of indicated genes inA17 and *cre1* roots**(A, B, C, D, E, F)** treated with turanose (10^{-3}M) and water (mock) followed by infection with *Sinorhizobiummeliloti Sm2011-pBHR-mRFP* relative to mock treated uninoculated control or transformed with empty vector, *p35S::AhWOX5* and *p35S::MtWOX5* followed by infection with *Sm2011-pBHR-mRFP* relative to empty vector transformed uninoculated control. *MtActin* was used as a reference gene. Histogram represents an average of three biological replicates each having n>3 plants and error bar represent standard deviation (SD). Mann-Whitney test was used to access significant differences where ***, ** and * represents $P < 0.001$, <0.01 and <0.05 respectively.

Figure 6: Proposed Model for the role of sugar signaling and *WOX5* expression in restoration of symbiotic efficiency in *M*. truncatula cytokinin perception mutant *cre1*. Open arrows indicate SYM pathway dependent *CRE1* activation that leads to auxin accumulation by auxin biosynthesis and/or auxin transport inhibition. The local auxin accumulation induces *WOX5* to generate the nodule primordia. On the basis of our observations we propose the black arrows indicating sugar induced cytokinin response, *NIN* expression, deconjugation mediated auxin response and *WOX5* expression. We propose feed back loops to indicate *WOX5* induced cytokinin response, *NIN* expression and deconjugation mediated auxin response. The blunt arrows indicate Sugar and *CRE1* mediated cytokinin response to be mutually antagonistic. Dashed arrow indicates differential activity of *AhWOX5*.

REFERENCES

750
751
752

Arrighi J-F, Godfroy O, de Billy F, Saurat O, Jauneau A, Gough C (2008) The RPG gene of Medicago truncatula controls Rhizobium-directed polar growth during infection. Proceedings of the National Academy of Sciences **105:** 9817-9822

Ayre BG (2011) Membrane-transport systems for sucrose in relation to whole-plant carbon partitioning. Molecular plant **4:** 377-394

Boisson-Dernier A, Chabaud M, Garcia F, Bécard G, Rosenberg C, Barker DG (2001) Agrobacterium rhizogenes-transformed roots of Medicago truncatula for the study of nitrogen-fixing and endomycorrhizal symbiotic associations. Molecular Plant-Microbe Interactions **14:** 695-700

Boivin C, Camut S, Malpica CA, Truchet G, Rosenberg C (1990) Rhizobium meliloti genes encoding catabolism of trigonelline are induced under symbiotic conditions. The Plant Cell **2:** 1157-1170

Boivin S, Kazmierczak T, Brault M, Wen J, Gamas P, Mysore KS, Frugier F (2016) Different cytokinin histidine kinase receptors regulate nodule initiation as well as later nodule developmental stages in Medicago truncatula. Plant, cell & environment **39:** 2198-2209

Broughton W, Dilworth M (1971) Control of leghaemoglobin synthesis in snake beans. Biochemical journal **125:** 1075-1080

Campanella JJ, Smith SM, Leibu D, Wexler S, Ludwig-Müller J (2008) The auxin conjugate hydrolase family of Medicago truncatula and their expression during the interaction with two symbionts. Journal of Plant Growth Regulation **27:** 26-38

Catoira R, Galera C, de Billy F, Penmetsa RV, Journet E-P, Maillet F, Rosenberg C, Cook D, Gough C, Dénarié J (2000) Four genes of Medicago truncatula controlling components of a Nod factor transduction pathway. The Plant Cell **12:** 1647-1665

Daum G, Medzihradszky A, Suzaki T, Lohmann JU (2014) A mechanistic framework for noncell autonomous stem cell induction in Arabidopsis. Proceedings of the National Academy of Sciences **111:** 14619-14624

Deinum EE, Geurts R, Bisseling T, Mulder BM (2012) Modeling a cortical auxin maximum for nodulation: different signatures of potential strategies. Frontiers in plant science **3:** 96

Demina IV, Maity PJ, Nagchowdhury A, Ng JL, van der Graaff E, Demchenko KN, Roitsch T, Mathesius U, Pawlowski K (2019) Accumulation of and response to auxins in roots and nodules of the actinorhizal plant Datisca glomerata compared to the model legume Medicago truncatula. Frontiers in plant science **10:** 1085

den Camp RHO, De Mita S, Lillo A, Cao Q, Limpens E, Bisseling T, Geurts R (2011) A phylogenetic strategy based on a legume-specific whole genome duplication yields symbiotic cytokinin type-A response regulators. Plant physiology **157:** 2013-2022

Ding Z, Friml J (2010) Auxin regulates distal stem cell differentiation in Arabidopsis roots. Proceedings of the National Academy of Sciences **107:** 12046-12051

Dolzblasz A, Nardmann J, Clerici E, Causier B, van der Graaff E, Chen J, Davies B, Werr W, Laux T (2016) Stem cell regulation by Arabidopsis WOX genes. Molecular plant **9:** 1028-1039

Francis D, Halford NG (2006) Nutrient sensing in plant meristems. Plant Molecular Biology **60:** 981-993

Franco-Zorrilla JM, Martín AC, Leyva A, Paz-Ares J (2005) Interaction between phosphate-starvation, sugar, and cytokinin signaling in Arabidopsis and the roles of cytokinin receptors CRE1/AHK4 and AHK3. Plant physiology **138:** 847-857

Franssen HJ, Xiao TT, Kulikova O, Wan X, Bisseling T, Scheres B, Heidstra R (2015) Root developmental programs shape the Medicago truncatula nodule meristem. Development **142:** 2941-2950

Frugier F, Kosuta S, Murray JD, Crespi M, Szczyglowski K (2008) Cytokinin: secret agent of symbiosis. Trends in plant science **13:** 115-120

Garcia J, Barker DG, Journet E-P (2006) Seed storage and germination. The Medicago
truncatula Handbook. The Samuel Roberts Noble Foundation, Ardmore, OK, http://www.
noble. org/MedicagoHandbook
Gonzalez-Rizzo S, Crespi M, Frugier F (2006) The Medicago truncatula CRE1 cytokinin
receptor regulates lateral root development and early symbiotic interaction with
Sinorhizobium meliloti. The Plant Cell **18**: 2680-2693
Gonzali S, Novi G, Loreti E, Paolicchi F, Poggi A, Alpi A, Perata P (2005) A
turanose-insensitive mutant suggests a role for WOX5 in auxin homeostasis in
Arabidopsis thaliana. The Plant Journal **44**: 633-645
Haynes JG, Czymmek KJ, Carlson CA, Veereshlingam H, Dickstein R, Sherrier DJ (2004)
Rapid analysis of legume root nodule development using confocal microscopy. New
Phytologist **163**: 661-668
Heckmann AB, Sandal N, Bek AS, Madsen LH, Jurkiewicz A, Nielsen MW, Tirichine L,
Stougaard J (2011) Cytokinin induction of root nodule primordia in Lotus japonicus is
regulated by a mechanism operating in the root cortex. Molecular plant-microbe
interactions **24**: 1385-1395
Held M, Hou H, Miri M, Huynh C, Ross L, Hossain MS, Sato S, Tabata S, Perry J, Wang TL
(2014) Lotus japonicus cytokinin receptors work partially redundantly to mediate nodule
formation. The Plant Cell **26**: 678-694
Ikeda M, Mitsuda N, Ohme-Takagi M (2009) Arabidopsis WUSCHEL is a bifunctional
transcription factor that acts as a repressor in stem cell regulation and as an activator in
floral patterning. The Plant Cell **21**: 3493-3505
Ioio RD, Nakamura K, Moubayidin L, Perilli S, Taniguchi M, Morita MT, Aoyama T,
Costantino P, Sabatini S (2008) A genetic framework for the control of cell division and
differentiation in the root meristem. Science **322**: 1380-1384
Jardinaud M-F, Boivin S, Rodde N, Catrice O, Kisiala A, Lepage A, Moreau S, Roux B,
Cottret L, Sallet E (2016) A laser dissection-RNAseq analysis highlights the activation
of cytokinin pathways by Nod factors in the Medicago truncatula root epidermis. Plant
physiology **171**: 2256-2276
Jones B, Gunnerås SA, Petersson SV, Tarkowski P, Graham N, May S, Dolezal K,
Sandberg G, Ljung K (2010) Cytokinin regulation of auxin synthesis in Arabidopsis
involves a homeostatic feedback loop regulated via auxin and cytokinin signal
transduction. The Plant Cell **22**: 2956-2969
Koch B, Evans HJ (1966) Reduction of acetylene to ethylene by soybean root nodules. Plant
Physiology **41**: 1748
Kong D, Hao Y, Cui H (2016) The WUSCHEL related homeobox protein WOX7 regulates the
sugar response of lateral root development in Arabidopsis thaliana. Molecular plant **9**:
261-270
Kundu A, DasGupta M (2018) Silencing of putative cytokinin receptor histidine kinase1 inhibits
both inception and differentiation of root nodules in Arachis hypogaea. Molecular plant-
microbe interactions **31**: 187-199
Laffont C, Ivanovici A, Gautrat P, Brault M, Djordjevic MA, Frugier F (2020) The NIN
transcription factor coordinates CEP and CLE signaling peptides that regulate nodulation
antagonistically. Nature communications **11**: 1-13
Lalonde S, Wipf D, Frommer WB (2004) Transport mechanisms for organic forms of carbon
and nitrogen between source and sink. Annu. Rev. Plant Biol. **55**: 341-372
LeClere S, Schmelz EA, Chourey PS (2010) Sugar levels regulate tryptophan-dependent
auxin biosynthesis in developing maize kernels. Plant Physiology **153**: 306-318
Leibfried A, To JP, Busch W, Stehling S, Kehle A, Demar M, Kieber JJ, Lohmann JU
(2005) WUSCHEL controls meristem function by direct regulation of cytokinin-inducible
response regulators. Nature **438**: 1172-1175

Lin H, Niu L, McHale NA, Ohme-Takagi M, Mysore KS, Tadege M (2013) Evolutionarily
conserved repressive activity of WOX proteins mediates leaf blade outgrowth and floral
organ development in plants. Proceedings of the National Academy of Sciences **110:**
366-371
Liu J, Rutten L, Limpens E, Van Der Molen T, Van Velzen R, Chen R, Chen Y, Geurts R,
Kohlen W, Kulikova O (2019) A remote cis-regulatory region is required for NIN
expression in the pericycle to initiate nodule primordium formation in Medicago
truncatula. The Plant Cell **31:** 68-83
Ljung K, Nemhauser JL, Perata P (2015) New mechanistic links between sugar and hormone
signalling networks. Current Opinion in Plant Biology **25:** 130-137
Madsen LH, Tirichine L, Jurkiewicz A, Sullivan JT, Heckmann AB, Bek AS, Ronson CW,
James EK, Stougaard J (2010) The molecular network governing nodule
organogenesis and infection in the model legume Lotus japonicus. Nature
communications **1:** 1-12
Marsh JF, Rakocevic A, Mitra RM, Brocard L, Sun J, Eschstruth A, Long SR, Schultze M,
Ratet P, Oldroyd GE (2007) Medicago truncatula NIN is essential for rhizobial-
independent nodule organogenesis induced by autoactive calcium/calmodulin-
dependent protein kinase. Plant physiology **144:** 324-335
Mathesius U, Weinman JJ, Rolfe BG, Djordjevic MA (2000) Rhizobia can induce nodules in
white clover by "hijacking" mature cortical cells activated during lateral root development.
Molecular Plant-Microbe Interactions **13:** 170-182
Meng WJ, Cheng ZJ, Sang YL, Zhang MM, Rong XF, Wang ZW, Tang YY, Zhang XS (2017)
Type-B ARABIDOPSIS RESPONSE REGULATORs specify the shoot stem cell niche by
dual regulation of WUSCHEL. The Plant Cell **29:** 1357-1372
Miri M, Janakirama P, Held M, Ross L, Szczyglowski K (2016) Into the root: how cytokinin
controls rhizobial infection. Trends in plant science **21:** 178-186
Mishra BS, Singh M, Aggrawal P, Laxmi A (2009) Glucose and auxin signaling interaction in
controlling Arabidopsis thaliana seedlings root growth and development. PLOS one **4:**
e4502
Moore B, Zhou L, Rolland F, Hall Q, Cheng W-H, Liu Y-X, Hwang I, Jones T, Sheen J
(2003) Role of the Arabidopsis glucose sensor HXK1 in nutrient, light, and hormonal
signaling. Science **300:** 332-336
Moubayidin L, Perilli S, Ioio RD, Di Mambro R, Costantino P, Sabatini S (2010) The rate of
cell differentiation controls the Arabidopsis root meristem growth phase. Current Biology
20: 1138-1143
Müller B, Sheen J (2008) Cytokinin and auxin interaction in root stem-cell specification during
early embryogenesis. Nature **453:** 1094-1097
Ng JL, Mathesius U (2018) Acropetal auxin transport inhibition is involved in indeterminate but
not determinate nodule formation. Frontiers in plant science **9:** 169
Ng JLP, Hassan S, Truong TT, Hocart CH, Laffont C, Frugier F, Mathesius U (2015)
Flavonoids and auxin transport inhibitors rescue symbiotic nodulation in the Medicago
truncatula cytokinin perception mutant cre1. The Plant Cell **27:** 2210-2226
Ohto M-a, Hayashi S, Sawa S, Hashimoto-Ohta A, Nakamura K (2006) Involvement of HLS1
in sugar and auxin signaling in Arabidopsis leaves. Plant and cell physiology **47:** 1603-
1611
Osipova M, Dolgikh E, Lutova L (2011) Peculiarities of meristem-specific WOX5 gene
expression during nodule organogenesis in legumes. Russian Journal of Developmental
Biology **42:** 226
Osipova MA, Mortier V, Demchenko KN, Tsyganov VE, Tikhonovich IA, Lutova LA,
Dolgikh EA, Goormachtig S (2012) WUSCHEL-RELATED HOMEOBOX5 gene
expression and interaction of CLE peptides with components of the systemic control add

two pieces to the puzzle of autoregulation of nodulation. Plant physiology **158:** 1329-
1341

Penmetsa RV, Frugoli JA, Smith LS, Long SR, Cook DR (2003) Dual genetic pathways
controlling nodule number inMedicago truncatula. Plant Physiology **131:** 998-1008

Pernisová M, Klíma P, Horák J, Válková M, Malbeck J, Souček P, Reichman P, Hoyerová
K, Dubová J, Friml J (2009) Cytokinins modulate auxin-induced organogenesis in
plants via regulation of the auxin efflux. Proceedings of the National Academy of
Sciences **106:** 3609-3614

Pi L, Aichinger E, van der Graaff E, Llavata-Peris CI, Weijers D, Hennig L, Groot E, Laux T
(2015) Organizer-derived WOX5 signal maintains root columella stem cells through
chromatin-mediated repression of CDF4 expression. Developmental cell **33:** 576-588

Plet J, Wasson A, Ariel F, Le Signor C, Baker D, Mathesius U, Crespi M, Frugier F (2011)
MtCRE1-dependent cytokinin signaling integrates bacterial and plant cues to coordinate
symbiotic nodule organogenesis in Medicago truncatula. The Plant Journal **65:** 622-633

Rampey RA, LeClere S, Kowalczyk M, Ljung K, Sandberg G, Bartel B (2004) A family of
auxin-conjugate hydrolases that contributes to free indole-3-acetic acid levels during
Arabidopsis germination. Plant physiology **135:** 978-988

Reid DE, Heckmann AB, Novák O, Kelly S, Stougaard J (2016) CYTOKININ
OXIDASE/DEHYDROGENASE3 maintains cytokinin homeostasis during root and
nodule development in Lotus japonicus. Plant Physiology **170:** 1060-1074

Rejeb IB, Pastor V, Gravel V, Mauch-Mani B (2018) Impact of β-aminobutyric acid on induced
resistance in tomato plants exposed to a combination of abiotic and biotic stress. J Agric
Sci **2:** 12

Richards S, Wink RH, Simon R (2015) Mathematical modelling of WOX5-and CLE40-
mediated columella stem cell homeostasis in Arabidopsis. Journal of experimental
botany **66:** 5375-5384

Rightmyer AP, Long SR (2011) Pseudonodule formation by wild-type and symbiotic mutant
Medicago truncatula in response to auxin transport inhibitors. Molecular plant-microbe
interactions **24:** 1372-1384

Sabatini S, Beis D, Wolkenfelt H, Murfett J, Guilfoyle T, Malamy J, Benfey P, Leyser O,
Bechtold N, Weisbeek P (1999) An auxin-dependent distal organizer of pattern and
polarity in the Arabidopsis root. Cell **99:** 463-472

Saha S, Dutta A, Bhattacharya A, DasGupta M (2014) Intracellular catalytic domain of
symbiosis receptor kinase hyperactivates spontaneous nodulation in absence of
rhizobia. Plant physiology **166:** 1699-1708

Sairanen I, Novák O, Pěnčík A, Ikeda Y, Jones B, Sandberg G, Ljung K (2012) Soluble
carbohydrates regulate auxin biosynthesis via PIF proteins in Arabidopsis. The Plant
Cell **24:** 4907-4916

Sarkar AK, Luijten M, Miyashima S, Lenhard M, Hashimoto T, Nakajima K, Scheres B,
Heidstra R, Laux T (2007) Conserved factors regulate signalling in Arabidopsis thaliana
shoot and root stem cell organizers. Nature **446:** 811-814

Schiessl K, Lilley JL, Lee T, Tamvakis I, Kohlen W, Bailey PC, Thomas A, Luptak J,
Ramakrishnan K, Carpenter MD (2019) NODULE INCEPTION recruits the lateral root
developmental program for symbiotic nodule organogenesis in Medicago truncatula.
Current Biology **29:** 3657-3668. e3655

Sinha AK, Hofmann MG, Römer U, Köckenberger W, Elling L, Roitsch T (2002)
Metabolizable and non-metabolizable sugars activate different signal transduction
pathways in tomato. Plant physiology **128:** 1480-1489

Skylar A, Hong F, Chory J, Weigel D, Wu X (2010) STIMPY mediates cytokinin signaling
during shoot meristem establishment in Arabidopsis seedlings. Development **137:** 541-
549

Skylar A, Wu X (2011) Regulation of Meristem Size by Cytokinin Signaling F. Journal of Integrative Plant Biology **53:** 446-454

Smit P, Raedts J, Portyanko V, Debellé F, Gough C, Bisseling T, Geurts R (2005) NSP1 of the GRAS protein family is essential for rhizobial Nod factor-induced transcription. Science **308:** 1789-1791

Stokes ME, Chattopadhyay A, Wilkins O, Nambara E, Campbell MM (2013) Interplay between sucrose and folate modulates auxin signaling in Arabidopsis. Plant Physiology **162:** 1552-1565

Suzaki T, Yano K, Ito M, Umehara Y, Suganuma N, Kawaguchi M (2012) Positive and negative regulation of cortical cell division during root nodule development in Lotus japonicus is accompanied by auxin response. Development **139:** 3997-4006

Tirichine L, Sandal N, Madsen LH, Radutoiu S, Albrektsen AS, Sato S, Asamizu E, Tabata S, Stougaard J (2007) A gain-of-function mutation in a cytokinin receptor triggers spontaneous root nodule organogenesis. Science **315:** 104-107

Ulmasov T, Murfett J, Hagen G, Guilfoyle TJ (1997) Aux/IAA proteins repress expression of reporter genes containing natural and highly active synthetic auxin response elements. The Plant Cell **9:** 1963-1971

van Noorden GE, Kerim T, Goffard N, Wiblin R, Pellerone FI, Rolfe BG, Mathesius U (2007) Overlap of proteome changes in Medicago truncatula in response to auxin and Sinorhizobium meliloti. Plant Physiology **144:** 1115-1131

van Noorden GE, Ross JJ, Reid JB, Rolfe BG, Mathesius U (2006) Defective long-distance auxin transport regulation in the Medicago truncatula super numeric nodules mutant. Plant Physiology **140:** 1494-1506

van Zeijl A, den Camp RHO, Deinum EE, Charnikhova T, Franssen H, den Camp HJO, Bouwmeester H, Kohlen W, Bisseling T, Geurts R (2015) Rhizobium lipo-chitooligosaccharide signaling triggers accumulation of cytokinins in Medicago truncatula roots. Molecular plant **8:** 1213-1226

Vernié T, Kim J, Frances L, Ding Y, Sun J, Guan D, Niebel A, Gifford ML, de Carvalho-Niebel F, Oldroyd GE (2015) The NIN transcription factor coordinates diverse nodulation programs in different tissues of the Medicago truncatula root. The Plant Cell **27:** 3410-3424

Wahl V, Brand LH, Guo Y-L, Schmid M (2010) The FANTASTIC FOUR proteins influence shoot meristem size in Arabidopsis thaliana. BMC Plant Biology **10:** 1-12

Wingler A, Fritzius T, Wiemken A, Boller T, Aeschbacher RA (2000) Trehalose induces the ADP-glucose pyrophosphorylase gene, ApL3, and starch synthesis in Arabidopsis. Plant Physiology **124:** 105-114

Wu X, Dabi T, Weigel D (2005) Requirement of homeobox gene STIMPY/WOX9 for Arabidopsis meristem growth and maintenance. Current Biology **15:** 436-440

Zhao Y, Hu Y, Dai M, Huang L, Zhou D-X (2009) The WUSCHEL-related homeobox gene WOX11 is required to activate shoot-borne crown root development in rice. The Plant Cell **21:** 736-748

Zürcher E, Tavor-Deslex D, Lituiev D, Enkerli K, Tarr PT, Müller B (2013) A robust and sensitive synthetic sensor to monitor the transcriptional output of the cytokinin signaling network in planta. Plant physiology **161:** 1066-1075

Figure 1: Turanose mediated recovery of symbiosis in *cre1* mutants. (A-B) Box plot represents (A) total nodule number per plant and (B) total white (white box) and pink nodules (red box) under control (mock) and Turanose treatment (10^{-3}M) in A17 and *cre1* after infection with *Sinorhizobium meliloti Sm2011-pBHR-mRFP*. Number above each box in (A) indicates the number of root systems with nodules per total number of scored roots. Mann and Whitney test was used to assess significant differences, where ***, ** and * indicates P<0.0005, <0.002 and <0.02 respectively.

(C-H) Morphology of infection event or nodules in *cre1* and A17 3WAI under (C-D) control condition and (E-H) after turanose treatment where (E and G) solitary pink nodules and (F and H) merged white nodules. Representative micrograph of control and turanose treated nodulated roots indicated as bright field and bright field+ mRFP merged.

(I-L) Ultrastructure analysis of nodules in *cre1* and A17 after turanose treatment, where the perforated box represents the infection zone with its corresponding enlarged view below the respective images. Total number of each type of nodules observed out of the total observed nodules are indicated for each panel. (upper panel) Brightfield + mRFP merged and (lower panel) mRFP. Scale bar: C-H = 500µm, I-L (upper panel) = 100µm and (lower panel) = 10µm.

Figure 2: Effect of turanose treatment on *pTCS:GUS*, *pDR5:GUS* and *pWOX5:GUS* expression pattern in A17 and *cre1* roots.
(A-H) *pTCS:GUS* expression in (A-D) A17 and (E-H) *cre1* roots, (I-P) *pDR5:GUS* expression in (I-L) A17 and (M-P) *cre1* roots, and (Q-X) *pWOX5:GUS* expression in (Q-T) A17 and (U-X) *cre1* roots under following conditions. **(A-H)** mock treated uninfected (A, E), infected (B, F), turanose treated uninfected (C, G) and infected (D, H). **(I-P)** mock treated uninfected (I, M), infected (J, N), turanose treated uninfected (K, O) and infected (L, P). **(Q-X)** mock treated uninfected (Q, U), infected (R, V), turanose treated uninfected (S, W) and infected (T, X). All plants were mock or Turanose (10^{-3}M) treated for 7 days followed by infection with *Sinorhizobium meliloti Sm2011-pBHR-mRFP* for 24 hours before being harvested for GUS staining. Every root image has its corresponding longitudinal section of the root tip and the root hair zone (box). At least 5 independent roots were analyzed for each condition from three independent experiments. Scale bars = 300 µm (left panel single root), Scale bar = 50µm (right panel upper and lower longitudinal section).

Figure 3: Rescue of symbiosis in *cre1* by ectopic expression of *AhWOX5*. (A) Percentile projection of epidermal infection thread (IT) observed 2WAI with *Sm1021-pXLGD4-lacZ* in *cre1* roots transformed with empty vector, *p35S::eGFP-MtWOX5* and *p35S::eGFP-AhWOX5*. Each of the infection events has been roughly classified into 4 categories: (i) successful cortical invasion and intracellular colonization (ii) Stalled in nodule apex, (iii) stalled in epidermal cortical barrier and (iv) stalled in microcolonies in root hair. Category (i) is considered normal whereas Category (ii-iv) is considered as abnormality. **(B)** Box plot representing cumulative number of normal and abnormal ITs. **(C-D)** Box plot represents(C) total nodule number per plant and **(D)** total white (white box) and pink nodules (red box) per plant in empty vector, *p35S::eGFP-MtWOX5* and *p35S::eGFP-AhWOX5* transformed hairy-root systems in A17 and *cre1* respectively, harvested 6WAI with *Sinorhizobium meliloti Sm2011-pBHR-mRFP*. n indicates the number of root systems with nodules per total number of scored roots. Mann-Whitney test was used to assess significant differences, where ***, ** and * indicates P<0.0005, 0.002 and <0.02 respectively. **(E-F)** qRT-PCR analysis of (E) *MtWOX5* and (F) *AhWOX5* relative to empty vector transformed (*cre1*) roots normalized against *MtActin*. **(G)** Acetylene reduction assay (nmole C_2H_4/h/mg nodule) in indicated root system. (E-G) Histogram represent an average of three biological replicates each having n>4 plants and error bar represents SD. Mann-Whitney test was used to assess significant differences, where ***, ** and * indicates P<0.0005, 0.002 and <0.02 respectively. **(H-M)** Morphology of nodules in transgenic hairy-roots of *cre1* mutant 4WAI with *Sm2011-pBHR-mRFP* (H, J and L). Representative micrograph of transgenic roots indicated as bright field + mRFP merged and perforated box is represented as eGFP + mRFP merged and bright field. Scale bar = 1mm and box = 500μm. (I, K and M) Ultrastructure of nodules developed in *cre1* mutant transformed with different constructs indicated as bright field+ mRFP merged where the perforated box in the infection zone is enlarged separately. Scale bar: I, K and M = 100μm and box = 10μm. Arrow indicate infection thread.

Figure 4: Expression analysis of cytokinin responsive genes under turanose treatment and WOX5 overexpression in A17 and *cre1*. Quantitative reverse transcriptase-PCR analysis of indicated genes in A17 and *cre1* roots (**A, B, C, D, E, F**) treated with turanose (10^{-3}M) and water (mock) followed by infection with *Sinorhizobium meliloti* Sm2011-pBHR-mRFP relative to mock treated uninoculated control or transformed with empty vector, *p35S::AhWOX5* and *p35S::MtWOX5* followed by infection with *Sm2011-pBHR-mRFP* relative to empty vector transformed uninoculated control. *MtActin* was used as a reference gene. Histogram represents an average of three biological replicates each having n>3 plants and error bar represent standard deviation (SD). Mann–Whitney test was used to access significant differences where ***, ** and * represents P < 0.001, <0.01 and <0.05 respectively.

Figure 5: Expression analysis of auxin responsive genes under turanose treatment and WOX5 overexpression in A17 and *cre1*. Quantitative reverse transcriptase-PCR analysis of indicated genes in A17 and *cre1* roots **(A, B, C, D, E, F)** treated with turanose (10^{-3}M) and water (mock) followed by infection with *Sinorhizobium meliloti Sm2011-pBHR-mRFP* relative to mock treated uninoculated control or transformed with empty vector, *p35S:AhWOX5* and *p35S:MtWOX5* followed by infection with *Sm2011-pBHR-mRFP* relative to empty vector transformed uninoculated control. *MtActin* was used as a reference gene. Histogram represents an average of three biological replicates each having n>3 plants and error bar represent standard deviation (SD). Mann–Whitney test was used to access significant differences where ***, ** and * represents P < 0.001, <0.01 and <0.05 respectively.

Figure 6: Proposed Model for the role of sugar signaling and *WOX5* expression in restoration of symbiotic efficiency in *M. truncatula* cytokinin perception mutant *cre1*. Open arrows indicate SYM pathway dependent *CRE1* activation that leads to auxin accumulation by auxin biosynthesis and/or auxin transport inhibition. The local auxin accumulation induces *WOX5* to generate the nodule primordia. On the basis of our observations we propose the black arrows indicating sugar induced cytokinin response, *NIN* expression, deconjugation mediated auxin response and *WOX5* expression. We propose feedback loops to indicate *WOX5* induced cytokinin response, *NIN* expression and deconjugation mediated auxin response. The blunt arrows indicate Sugar and *CRE1* mediated cytokinin response to be mutually antagonistic. Dashed arrow indicates differential activity of *AhWOX5*.