TITLE

Immune checkpoint proteins are conserved across 160 million years of evolution and are expressed

on transmissible cancers

RUNNING TITLE

Transmissible cancers express evolutionarily conserved immune checkpoint molecules

AUTHORS

Andrew S. Flies[1]*, Jocelyn M. Darby[1], Patrick R. Lennard[1,2], Peter R. Murphy[1,3], Chrissie E. B.

Ong[1], Terry L. Pinfold[4], A. Bruce Lyons[4], Gregory M. Woods[1], Amanda L. Patchett[1]

AFFILIATIONS

[1]Menzies Institute for Medical Research, College of Health and Medicine, University of Tasmania,

Hobart, TAS 7000, Australia

[2]The Roslin Institute and Royal School of Veterinary Studies, University of Edinburgh, Easter

Bush Campus, Midlothian, EH25 9RG, UK

[3]University of Queensland Diamantina Institute, The University of Queensland, Translational

Research Institute, Woolloongabba, Queensland, Australia

[4]School of Medicine, College of Health and Medicine, University of Tasmania, Hobart, TAS 7000,

Australia

CORRESPONDING AUTHOR CONTACT INFORMATION

Andrew S. Flies, PhD

Menzies Institute for Medical Research, College of Health and Medicine

University of Tasmania

Private Bag 23, Hobart TAS 7000

phone: +61 3 6226 4614; email: Andy.Flies@utas.edu.au

GRAPHICAL ABSTRACT

SIGNIFICANCE

Immune checkpoint immunotherapy has revolutionized medicine, but translational success for

new treatments remains low. Around 40% of humans and Tasmanian devils (*Sarcophilus harrisii*)

develop cancer in their lifetime, compared to less than 10% for most species. Additionally, devils

are affected by two of the three known transmissible cancers in mammals. Unfortunately, little is

known about the role of immune checkpoints in devils and other non-model species, largely due

to a lack of species-specific reagents. We developed a simple cut-and-paste reagent development

method applicable to any vertebrate species and show that immune checkpoint interactions are

conserved across 160 million years of evolution. The inhibitory checkpoint molecule CD200 is

highly expressed on devil facial tumor cells. We are the first to demonstrate that co-expression of

CD200R1 can block CD200 expression. The evolutionarily conserved pathways suggest that

naturally occurring cancers in devils and other species can serve as models for understanding

cancer and immunological tolerance.

ABSTRACT

Around 40% of humans and captive Tasmanian devils (*Sarcophilus harrisii*) develop cancer, compared to less than 10% for most species. Tasmanian devils also suffer from two of the three known naturally-occurring transmissible cancers detected in vertebrate species. Transmissible cancers are a unique form of cancer in which tumor cells act as an infectious pathogen and an allograft. The two different transmissible devil facial tumors (DFT1 and DFT2) overcome immunological barriers (e.g. major histocompatibility complex) and have killed thousands of devils. Immune checkpoint immunotherapy has revolutionized human oncology in recent years. However, immune checkpoints in transmissible and non-transmissible cancers remains largely unexplored in most species due to a lack of species-specific reagents. To overcome this, we developed a "cut-and-paste" Fluorescent Adaptable Simple Theranostic (FAST) protein system, adaptable for any vertebrate species. This method facilitated rapid confirmation of seven receptor-ligand interactions between twelve immune checkpoint proteins in Tasmanian devils, thus filling a 160 million year gap in our understanding of the evolution of the mammalian immune system. We used this system to investigate the checkpoint molecule CD200, which can inhibit natural killer cell responses to cancer and facilitate graft tolerance in humans and mice. CD200 was highly expressed on DFT cells and can be used to identify DFT cells in devil blood. Understanding how transmissible tumor cells graft onto new hosts and evade immune defenses will help to identify evolutionarily conserved immunological principles relevant to transplant immunology, cancer, and infectious disease for human and veterinary medicine.

KEYWORDS

INTRODUCTION

Metastatic cancer affects most mammals, but the cancer incidence can vary widely across phylogenetic groups and species (**Figure 1, Table S1**) [1–9]. In humans, the lifetime risk of developing cancer is around 40% [10]. This is in stark contrast to a general cancer incidence of 3% for mammals, 2% for birds, and 2% for reptiles reported by the San Diego Zoo (n=10,317) [7,11]. A more recent study at the Taipei Zoo reported cancer incidence of 8%, 4%, and 1% for mammals, birds, and reptiles, respectively (n=2,657) [4]. Cancer incidence in domestic animals is generally less than 10% (n=202,277) [9]. However, two studies performed 40 years apart reported that greater than 40% of Tasmanian devils develop spontaneous, often severe neoplasia in their lifetime [11,12]. Devils are also unique because they are affected by two of the three known naturally-occurring transmissible cancers in vertebrate species [13,14]. Transmissible cancers are a distinct form of cancer in which the tumor cells function as an infectious pathogen and an allograft. Dogs (*Canis lupus familiaris*) are the only other vertebrate species affected by a transmissible cancer [15], and interestingly some breeds of dogs also have high cancer incidence [9,16].

The devil facial tumor (DFT) disease was first detected in Northwest Tasmania and has been a primary driver of an 80% decline in the wild Tasmanian devil population [13,17]. The clonal devil facial tumor (DFT1) cells have been continually transmitted among devils and is estimated to have killed at least 10,000 individuals since at least 1996. In 2014 a second independent transmissible Tasmanian devil facial tumor (DFT2) was discovered in wild devils [14] and 23 cases have been reported to date [18]. Genetic mismatches, particularly in the major histocompatibility complex (MHC) genes should lead to rejection of these transmissible tumors. Consequently, the role of devil MHC has been a focus of numerous studies (**Figure 1, Table S1**) to understand the lack of rejection of the transmissible tumors. These studies have revealed that the DFT1 cells

downregulate MHC class I (MHC-I) expression [19], a phenomenon observed in many human cancers [20]. In contrast to DFT1 cells, the DFT2 cells do express MHC-I [21]. DFT1 and DFT2 cells also have 2,884 and 3,591 single nucleotide variants, respectively, that are not present in 46 normal devil genomes [22]. The continual transmission of DFT1 and DFT2, despite MHC-I expression by DFT2 cells and genetic mismatches between host and tumor, suggests that additional pathways are likely involved in immune evasion.

Human cancer treatment has been revolutionized in the past decade by manipulating interactions among immune checkpoint molecules [23,24]. These have proven broadly effective in part because they function across many different MHC types and tumor mutational patterns. However, these pathways have received little attention in transmissible cancers and other naturally occurring cancers in non-model species (**Figure 1, Table S1**) [25–27]. We have previously shown that the inhibitory immune checkpoint molecule programmed death ligand 1 (PDL1) is expressed in the DFT microenvironment and is upregulated by interferon-gamma (IFNγ) *in vitro* [25]. This finding led us to question which other immune checkpoint molecules play a role in immune evasion by the transmissible cancers and the devil's high spontaneous cancer incidence. Understanding this immune evasion in a natural environment has the potential to help protect this endangered species and identify protein interactions that are conserved across divergent species to improve translational success of animal models [27]. Unfortunately, a persistent limitation for immunology in non-traditional study species is a lack of species-specific reagents. Wildlife biologists and veterinarians are at the front lines of emerging infectious disease outbreaks, but they often lack species-specific reagents to fulfil the World Health Organization's call for "cross-cutting R&D preparedness" and perform mechanistic immunological investigations [28].

To solve the paucity of reagents available for Tasmanian devils and address ongoing

limitations for non-traditional study species, we developed a Fluorescent Adaptable Simple

Theranostic (FAST) protein system that builds upon the diverse uses of fluorescent proteins

previously reported [29–33]. This simple system can be used for rapid development of diagnostic and

therapeutic (i.e. theranostic) immunological toolkits for any animal species (**Figure 2**). We

demonstrate the versatility and impact of the FAST system by using it to confirm seven receptor-

ligand interactions among twelve checkpoint proteins in devils.

In humans, these checkpoint proteins have been targets of immunotherapy in clinical trials,

but the functional role and binding patterns of these proteins are unknown for most other species.

We have used the FAST system to show that the inhibitory checkpoint protein CD200 is highly

expressed on DFT cells, opening the door to single-cell phenotyping of circulating tumor cells

(CTCs) in devil blood. Furthermore, we are the first to report that co-expression of CD200R1 can

block surface expression of CD200 in any species. Understanding how clonal tumor cells graft

onto new hosts, evade immune defenses and metastasize within a host will identify evolutionarily

conserved immunological mechanisms to help improve cancer, infectious disease, and transplant

outcomes for human and veterinary medicine.

RESULTS

Fluorescent fusion proteins can be secreted from mammalian cells

Initially, we developed FAST proteins to determine whether monomeric fluorescent

proteins could be fused to devil proteins and secreted from mammalian cells (**Figure 2A** and **Table**

S2). We used 41BB (TNFRSF9) for proof-of-concept studies by fusing the extracellular domain

of devil 41BB checkpoint molecule to monomeric fluorescent proteins (**Figure 2A-B** and **Figure**

S1). We used wild-type Chinese hamster ovary (CHO) cells and CHO cells transfected with

41BBL (TNFSF9) to confirm specificity of the 41BB FAST proteins and demonstrate that the

fluorescent proteins (mAzurite, mCerulean3, mCherry, mCitrine, mOrange, Neptune2, and mTag-

BFP (aka mBFP)) remained fluorescent when secreted from mammalian cells (**Figure 2C**).

We chose mCherry, mCitrine, mOrange, and mBFP for ongoing FAST protein

development. Initial batches of FAST proteins were purified using the 6xHis-tag and eluted with

imidazole. The gradient of FAST protein in the collection tubes was apparent when excited with

blue light and visualized with an amber filter unit (**Figure 2D**), allowing immediate confirmation

that the fluorescent protein DNA coding sequences were in-frame and the proteins were properly

folded. mCherry was visible without excitation or filters (**Figure 2E**). After combining,

concentrating, and sterile filtering the eluted fractions, 100 µL was aliquoted and visualized again

using blue light to confirm fluorescent signal (**Figure 2F**). A full step-by-step protocol and set of

experimental templates for creating and testing FAST proteins for any species is available online

with the supplementary material.

Receptor-ligand binding confirmed in single-step staining assays

We chose additional immune checkpoint molecules for FAST protein development

(**Figure 3A**) based on targets of clinical trials and sequence analysis of devil genes [27,34,35]. We

transfected the FAST protein expression vectors (**Table S2**) into CHO cells and tested the

supernatant against CHO cell lines expressing full-length receptors. 41BB FAST proteins in

supernatant exhibited strong binding to 41BBL cell lines, but the fluorescent signals from most

other FAST proteins were too weak to confirm binding to the expected receptors (**Figure S2**). As

FAST proteins do not require secondary reagents, we next incubated target cells with purified

FAST proteins and added chloroquine to block the lysosomal protein degradation pathway [36]. This

allowed us to take advantage of receptor-mediated endocytosis, which can allow accumulation of captured fluorescent signals inside the target cells [37]. This protocol adjustment allowed confirmation that CD47-mCherry, CD200-mBFP, CD200-mOrange, CD200R1-mBFP, and CD200R1-mOrange, and PD1-mCitrine bound to their expected receptors (**Figure 3B**). We also demonstrated the flexibility of the FAST proteins by showing that alternative fusion conformations (**Figure S1C-D**), such as type II proteins (e.g. mCherry-41BBL) and a devil Fc-tag (e.g. CD80-Fc-mCherry) bound to their expected ligands (**Figure 3B**). The stability of the fusion proteins was demonstrated using supernatants that were stored at 4 °C for two months prior to use in a one-hour live-culture assay with chloroquine (**Figure S3**).

Cell lines secreting FAST proteins confirm protein interactions in live coculture assays

To further streamline the reagent development process, we next took advantage of the single-step nature of FAST proteins (i.e. no secondary antibodies or labels needed) in live-cell coculture assays (**Figure 4A**). Cell lines secreting 41BB-mCherry, 41BBL-mCherry, or CD80-Fc-mCherry FAST proteins were mixed with cell lines expressing full-length 41BB, 41BBL, or CTLA4-mCitrine and cocultured at a 1:1 ratio overnight with chloroquine. Singlet cells were gated (**Figure 4B**) and binding of mCherry FAST proteins to CFSE or mCitrine-labelled target cells was analyzed (**Figure 4C**). The strongest fluorescent signal from 41BB-mCherry, 41BBL-mCherry and CD80-Fc-mCherry were detected when cocultured with their predicted receptors, 41BBL, 41BB and CTLA4, respectively.

Optimization of the FAST-Fc construct

The fluorescent binding signal of CD80-Fc-mCherry was lower than expected, so we next re-examined our Fc-tag construct. In humans and all other mammals examined to date the IgG heavy chain has glycine-lysine (Gly-Lys) residues at the C-terminus [38]; the initial devil IgG

constant region sequence available to us had an incomplete C-terminus, and thus our initial CD80-Fc-mCherry vector did not have the C-terminal Gly-Lys. We subsequently made a new FAST-Fc construct with CTLA4-Fc-mCherry, which exhibited strong binding to both CD80 and CD86 transfected DFT cells (**Figure 4D**).

CD200 mRNA and protein are highly expressed in DFT cells

Analysis of previously published devil and DFT cell transcriptomes suggested that CD200 mRNA is highly expressed in DFT2 cells and peripheral nerves, moderately expressed in DFT1 cells, and lower in other healthy devil tissues (**Figure 5A**) [34,35,39]. As CD200 is an inhibitory molecule expressed on most human neuroendocrine neoplasms [40], and both DFT1 and DFT2 originated from Schwann cells [35,41], we sought to investigate CD200 expression on DFT cells at the protein level. Staining of wild-type DFT1 and DFT2 cells with CD200R1-mOrange FAST protein showed minimal fluorescent signal (**Figure 5B**). However, overexpression of CD200 using a human EF1α promoter yielded a detectable signal with CD200R1-mOrange binding to CD200 on DFT1 cells. A weak signal from CD200-mOrange was detected on DFT1 cells overexpressing CD200R1 (**Figure 5B**). To confirm naturally-expressed CD200 on DFT cells we digested CD200 and 41BB FAST proteins using TEV protease to remove the linker and fluorescent reporter. The digested proteins were then used to immunize mice for polyclonal sera production. We stained target CHO cell lines with pre-immune (PI) or immune (I) mouse sera collected after 3X immunizations. Only the immune sera showed strong binding to the respective CD200 and 41BB target cell lines (**Figure 5C**). After the final immunization (4X), we collected another batch of sera and tested it on DFT1 and DFT2 cells (**Figure 5D**). In agreement with the transcriptomic data for DFT cells [34], the polyclonal sera revealed high levels of CD200 on DFT cells, but low levels of 41BB.

Overexpression of CD200R1 blocks surface expression of CD200

In humans, overexpression of some checkpoint proteins can block surface expression of heterophilic binding partners in *cis* (e.g. CD80 and PDL1) [42,43]. As a potential route for disrupting the inhibitory effects of CD200 on anti-tumor immunity, we tested if overexpression of CD200R1 on DFT cells could reduce CD200 surface expression. We stained a DFT1 strain C5065, and DFT1 C5065 cells transfected to overexpress CD200 or CD200R1 with polyclonal anti-CD200 sera and secondary anti-mouse IgG AF647. We detected no surface protein expression of CD200 DFT1 cells overexpressing CD200R1 (**Figure 5E**).

Identification of DFT cells in whole blood using anti-CD200

In addition to high expression of CD200 on neuroendocrine neoplasms [40,44], CD200 is used as a diagnostic marker for several human blood cancers [45–47]. DFT cells metastasize in the majority of cases [48] and our transcriptome results (**Figure 5A**) suggest that CD200 mRNA is more highly expressed in DFT cells than in peripheral blood mononuclear cells (PBMCs) [34,35]. As a result, we tested if CD200 could be used to identify DFT cells in blood. We stained PBMCs and DFT2 cells separately with polyclonal anti-CD200 sera and anti-mouse AlexaFluor 647 and then analyzed CD200 expression by flow cytometry (**Figure S4A**). We then mixed the stained PBMCs and DFT2 cells at ratios of 1:10 (**Figure S4A**) and 1:5 (**Figure S4B**) and analyzed the mixed populations. PBMCs showed minimal CD200 expression and background staining (**Figure S4**), whereas CD200 was highly expressed on DFT2 cells. CD200+ DFT2 cells were readily distinguishable from PBMCs.

As our RNAseq results only included mononuclear cells, we next performed a pilot test to determine if DFT cells could be spiked into whole devil blood and identified via flow cytometry using CD200 staining. DFT1 and DFT2 cells were labeled with CellTrace violet (CTV) and 10,000

cells were diluted directly into 100 µL of whole blood from a healthy devil (n=1/treatment; n = 1

devil). The cells were then stained with purified polyclonal anti-CD200 with and without

secondary anti-mouse IgG AF647 prior to red blood cell (RBC) lysis. Initial results showed that

DFT2 cells expressed CD200 above the leukocyte background, but that DFT1 cells could not be

distinguished from leukocytes (**Figure S5**). To eliminate the secondary antibody step from the

whole blood staining protocol we next labelled the polyclonal anti-CD200 and normal mouse

serum (NMS) with a no-wash Zenon mouse IgG AF647 labeling reagent (n = 1/treatment; n = 2

devils). This system again showed that CD200 expression could be used to identify DFT2 cells in

blood (**Figure 6**), suggesting that CD200 is a candidate marker for identification of metastasizing

DFT2 cells.

DISCUSSION

Naturally occurring cancers provide a unique opportunity to study immune evasion and the

metastatic process across diverse hosts and environments. The exceptionally high cancer rate in

Tasmanian devils coupled with the two transmissible tumors currently circulating in the wild

warrants a thorough investigation of the devil immune system. However, taking advantage of such

natural disease models has been out of reach for most species due to a lack of reagents. The FAST

protein system we developed here is well-suited to discovering additional DFT markers, and more

generally, filling the reagent gap for non-traditional species. For proteins like 41BB that have high

affinity for 41BBL, FAST proteins can be used as detection reagents directly from supernatant.

For other molecules with lower receptor-ligand affinity, the FAST proteins can be purified,

digested with a protease to remove the non-target proteins and used for production of polyclonal

or monoclonal antibodies.

The simple cut-and-paste methods for vector assembly lend the FAST protein system to

entry level immunology and molecular biology skill sets. Additionally, the ability of FAST

proteins to be used in live coculture assays and with elimination of secondary reagents, will

increase efficiency and reduce experimental error for advanced human and mouse cancer

immunology studies. For example, previous high-throughput studies have used a two-step staining

process (i.e. recombinant protein + secondary antibody) to screen more than 2000 protein

interactions [49,50]; this type of assay can be streamlined using FAST proteins to eliminate the need

for secondary antibodies. Fc-tags or other homodimerization domains can be incorporated into

FAST proteins to increase binding for low-affinity interactions.

Production of recombinant proteins in cell lines that closely resemble the physiological

conditions of the native cell type (i.e. mammalian proteins produced in mammalian cell lines) are

more likely to yield correct protein folding, glycosylation, and function than proteins produced

using evolutionarily distant cell lines [51]. The fluorescent fusion proteins developed here that take

advantage of natural receptor expression and cycling processes (e.g. CTLA4 transendocytosis) in

eukaryotic target cells; bacterial protein production methods are not amenable to coculture with

eukaryotic target cells in immunological assays [52]. Our demonstration of the FAST protein system

in CHO cells, which are used to produce approximately 70% of recombinant pharmaceutical

proteins [53], suggest that this method can be efficiently integrated into existing research and

development pipelines for humans and other vertebrate species.

A primary question in transmissible tumor research is why genetically mismatched cells

are not rejected by the host. Successful infection of devils with DFT cells relies on the ability of

the tumor allograft to evade and manipulate host defences. The "missing-self" hypothesis suggests

that the lack of constitutive MHC-I expression on DFT1 cells should lead to natural killer (NK)

cell-mediated killing of the allograft tumor cells. Here we used the FAST protein system to develop

a tool set to address this question and show that DFT1 and DFT2 cells express CD200 at higher

levels than most other devil tissues examined to date. CD200 has been shown to directly inhibit

NK cells in other species [54–56], so overexpression of CD200 is a potential mechanism of immune

evasion of NK responses by DFT cells.

We hypothesize that CD200 could be particularly important in DFT transmission as the

CD200-CD200R pathway is critical to the initial stages of establishing transplant and allograft

tolerance in other species [57–59]. In line with this hypothesis, a recent study reported that

overexpressing several checkpoint molecules, including CD200, PDL1, and CD47, in mouse

embryonic stem cells could be used to generate teratomas that could establish long-term allograft

tolerance in fully immunocompetent hosts [60]. We have previously reported that PDL1 mRNA and

protein are upregulated on DFT2 cells in response to IFNγ [25], and our transcriptome results show

that CD47 is expressed at moderate to high levels in DFT cells. Here we show that overexpression

of CD200R1 on DFT1 eliminates binding of our polyclonal anti-CD200 antibodies, suggesting

that DFT cells overexpressing CD200R1 could be used to test the role of CD200 in allograft

tolerance. Alternatively, genetic ablation of CD200 in DFT cells could be used as a complementary

approach to examine the role of immune checkpoint molecules in DFT allograft tolerance. The

CD200-CD200R1 pathway has been implicated in reducing IFNγ production by dendritic cells [59]

and decreasing the responsiveness of myeloid cells to IFNγ stimulation [61]. Low MHC-I expression

is a primary means of immune evasion by DFT1 cells, and disrupting the CD200-CD200R1

pathway could facilitate improved recognition of DFT1 cells by CD8 T cells by enhancing IFNγ-

mediated MHC-I upregulation. Recent work in mice has identified immunosuppressive natural

regulatory plasma cells that express CD200, LAG3, PDL1, and PDL2; we have previously

identified PDL1+ cells with plasma cell morphology near or within the DFT microenvironment [25].

Previous DFT vaccine efforts have used killed DFT cells with adjuvants [62,63]. A similar

approach to treat gliomas in dogs reported that tumor-lysate with CD200 peptides nearly doubled

progression-free survival compared to tumor lysate alone [64]. Like devils, several breeds of dog are

prone to cancer and these genetically-outbred large animal models provide a fertile ground for

testing cancer therapies. Interestingly, the CD200 peptides are reported to provide agonistic

function through CD200-like activation receptors (CD200R4) rather than by blocking CD200R1

[44,64]. The functional role of CD200-CD200R pathway in devils remains to be elucidated, but the

CD200R1$_{NPLY}$ inhibitory motif and key tyrosine residues are conserved in devil CD200R [27,65,66],

demonstrating this motif is conserved over 160 million years of evolutionary history [67]. In addition

to agonistic peptides, several other options for countering CD200-CD200R immune inhibition are

possible. Human chronic lymphocytic leukemia cells often express high levels of CD200, which

can be downregulated in response to imiquimod [68]. Likewise, we have previously shown that DFT1

cells downregulate expression of CD200 mRNA *in vitro* in response to imiquimod treatment [34].

In mice, chronic salmonella and schistosoma infections upregulated both CD200 and

CD200R [69]. Several viruses, including cytomegalovirus [70] and herpesvirus [71] manipulated the

CD200-CD200R pathway as a means of immune evasion. Interestingly, in one of the longest

running and most in-depth studies of host-pathogen coevolution, CD200R was shown to be under

selection in rabbits in response to myxoma virus biocontrol agent [72]. As DFT1 and DFT2 have

been circulating in devils for more than 20 years and 5 years, respectively, it will be important to

monitor CD200/R expression and the potential evolution of paired activating and inhibitory

receptors in these natural disease models [73].

Immunophenotyping and single-cell RNAseq of circulating tumor cells (CTCs) has

potential to identify key gene expression patterns associated with metastasis and tissue invasion.

Periaxin (PRX) is the most sensitive and specific marker for DFT1 cells in immunohistochemistry

assays [74]. Unfortunately, PRX is expressed primarily in the cytoplasm, which eliminates the

possibility of using PRX as a marker to sort live cells via flow cytometry for single-cell RNAseq.

However, CD200 is a potential marker for the identification of circulating tumor cells (CTCs) from

devil blood. As proof of concept, DFT2 cells could be identified in devil blood spiked with DFT2

cells. As CTCs are likely to be rare in the blood of most infected devils, CD200 alone would be

insufficient for identifying DFT1 cells. Additional surface DFT markers would be required to

purify CTCs for metastases and tissue invasion analyses. The FAST protein system provides a

simple procedure to facilitate the production of a panel of DFT-markers to help identify key

proteins in the metastatic process.

In summary, the simple "cut-and-paste" production of the vectors and single-step testing

pipeline of the FAST system provided multiple benefits. The FAST system allowed us to

characterize receptor-ligand interactions, and to identify evolutionarily conserved immune evasion

pathways in naturally occurring transmissible cancers. Our initial implementation of the system

confirmed numerous predicted protein interactions for the first time in a marsupial species and

documented high expression of the inhibitory molecule CD200 on DFT cells. The high expression

of CD200 in devil nervous tissues and neuroendocrine tumors, downregulation of CD200 in

response to imiquimod, and binding of CD200 to CD200R1, is consistent with results from human

and mouse studies. Consequently, the CD200/R pathway provides a promising immunotherapy

and vaccine target for DFTs. Beyond this study, FAST proteins meet the key attributes needed for

reagent development, such as being straightforward to make, stable, versatile, renewable, cheap,

and amenable to high-throughput testing [75]. The direct fusion of the reporter protein to the protein-

of-interest allows for immediate feedback during transfection, supernatant testing, and protein

purification; proteins with frameshifts, introduced stop codons, or folded improperly will not

fluoresce and can be discarded after a simple visualization, rather than only after extensive

downstream testing. Efficient mapping of immune checkpoint interactions across species can

identify evolutionarily conserved immune evasion pathways and appropriate large animal models

with naturally occurring cancer. This knowledge could inform veterinary and human medicine in

the fields of immunological tolerance to tissue transplants, infectious disease, and cancer.

MATERIALS AND METHODS

Study design

The objectives of this study were to fill a major gap in our understanding of the mammalian

immune system and to understand how genetically mismatched transmissible tumors evade host

immunity. To achieve this goal, we developed a recombinant protein system that directly fuses

proteins-of-interest to a fluorescent reporter protein. The first phase was to determine if the

fluorescent protein remained fluorescent after secretion from mammalian cells and to confirm that

proteins bound to their predicted receptors (i.e. ligands). Initial testing was performed in CHO

cells and follow-up assays used devil cells. To further demonstrate the functionality of this system

for antibody development, mice were immunized with either 41BB or CD200 proteins. Pre- and

post-immunization polyclonal sera was used to confirm that the proteins used for immunization

induced antibodies that specifically bound to surface-expressed recombinant proteins and native

proteins on devil facial tumor cells.

Target transcript amplification

Target gene DNA sequences for vector construction were retrieved from Genbank,

Ensembl or de novo transcriptome assemblies (**Table S2**) [76]. Target DNA was amplified from a

cDNA template or existing plasmids using primers and PCR conditions shown in **Tables S2-S4**

using Q5 High-Fidelity 2X Master Mix (New England Biolabs # M0494L). Primers were ordered

with 5' base extensions that overlapped expression vectors on either side of the restriction sites.

The amplified products were identified by gel electrophoresis and purified using Nucleospin PCR

and Gel Clean Up Kit (Macherey-Nagel # 740609.5). Alternatively, DNA sequences were

purchased as double stranded DNA gblocks (**Table S5**) (Integrated DNA Technologies) for direct

assembly into expression vectors.

Construction of all-in-one Sleeping Beauty transposon vectors

All new plasmids were assembled using NEBuilder kit (NEB # E5520S) following the

manufacturer's recommendations unless otherwise noted. DNA inserts, digested plasmids, and

NEBuilder master mix were incubated for 60 minutes at 50 °C and then transformed into DH5α

included with the NEBuilder kit. Plasmid digestions were performed following manufacturer

recommendations and generally subjected to Antarctic phosphatase (New England Biolabs #

M0289S) treatment to prevent potential re-annealing. Sleeping Beauty transposon vectors pSBbi-

Hyg (Addgene # 60524), pSBbi-BH (Addgene # 60515), pSBtet-Hyg (Addgene # 60508), pSBtet-

RH (Addgene # 60500) were gifts to Addgene from Eric Kowarz [77]. The pCMV(CAT)T7-SB100

containing the CMV promoter and SB100X transposase was a gift to Addgene from Zsuzsanna

Izsvak (Addgene # 34879) [78]. We first constructed an all-in-one Sleeping Beauty vector by

inserting a CMV promoter and SB100X transposase from pCMV(CAT)T7-SB100 [78] into pSBi-

BH [77] (**Tables S3-S4**). This was accomplished by using pAF111-vec.FOR and pAF111.1.REV

primers to amplify an overlap region from pSBbi-BH (insert 1) and pAF111-2.FOR and pAF111-

2.REV to amplify the CMV-SB100X region from pCMV(CAT)T7-SB100 (insert 2). The purified

amplicons were then used for NEBuilder assembly of pAF111. The final all-in-one vectors

pAF112 (hygromycin resistance and luciferase) and pAF123 (hygromycin resistance) were

assembled from the pAF111 components. pAF112 was assembled by amplifying the Luc2

luciferase gene (insert 1) from pSBtet-Hyg and the P2A-hygromycin resistance gene (insert 2)

from pSBbi-BH and inserting into the pAF111 Bsu36I digest using NEBuilder. pSBbi-Hyg was

Bsu36I-digested to obtain the hygromycin resistance gene, and this fragment was inserted into

Bsu36I-digested pAF111 using T4 ligase cloning to replace the BFP-P2A-hygromycin segment in

pAF111.

Construction of full-length protein vectors

All full-length gene coding sequences except CTLA4 were cloned into the a pAF112 SfiI

digest (**Table S2**). All full-length vectors also contain luciferase with T2A peptide linked to the

hygromycin resistance protein; luciferase was included for use in downstream functional testing

that was not part of this study. Tasmanian devil CTLA4 was cloned into a NotI-HF and XmaI

digest of pAF100 that was used in a different study but is derived from vectors pAF112 and

pAF138. Additionally, we also used devil PDL1 (CHO.pAF48) and 41BBL (CHO.pAF56) cell

lines developed using a vector system described previously [25].

Construction of FAST protein vectors

401   Plasmids containing fluorescent protein coding sequences mCerulean3-N1 (Addgene #

402 54730), mAzurite-N1 (Addgene # 54617), mOrange-N1 (Addgene # 54499), mNeptune2-N1

403 (Addgene # 54837) were gifts to Addgene from Michael Davidson. mTag-BFP was amplified from

404 pSBbi-BH, mCitrine was amplified from pAF71, and mCherry was amplified from pTRE-Dual2

405 (Clontech # PT5038-5). pAF137 was constructed by amplifying the devil 41BB extracellular

domain with primers pAF137-1.FOR and pAF137-1.REV and amplifying mCherry with pAF137-

2a.FOR and pAF137-2.REV (**Table S3-4**). 5' extensions on pAF137-1.FOR and pAF137-2.REV

were used to create overlaps for NEBuilder assembly of pAF137 from a pAF123 SfiI-digested

base vector. 3' extensions on pAF137-1.REV and pAF137-2a.FOR were used to create the linker

that included an XmaI/SmaI restriction site, TEV cleavage tag, GSAGSAAGSGEF linker peptide,

and 6x-His tag between the gene-of-interest and fluorescent reporter. The GSAGSAAGSGEF was

chosen due to the low number of large hydrophobic residues and less repeated nucleic acids than

are needed with other flexible linkers such as $(GGGS)_4$ [79]. The pAF137 primer extensions also

created 5' NotI and 3' NheI sites in the FAST vector to facilitate downstream swapping of

functional genes and to create a Kozak sequence [80] (GCCGCCACC) upstream of the FAST protein

open-reading frame. Following confirmation of correct assembly via DNA sequencing, the FAST

41BB-mCherry (pAF137) was digested and used as the base vector (**Figure 2B** and **Figure S1A-**

B) for development of FAST vectors with alternative fluorescent proteins. This was accomplished

by digestion of pAF137 with SalI and NheI and then inserting PCR-amplified coding sequences

for other fluorescent proteins using NEBuilder (**Tables S3-S4**).

Type I FAST (extracellular N-terminus, cytoplasmic C-terminus) protein vectors were

constructed by digestion of 41BB FAST vectors with NotI and either XmaI or SmaI (**Figure 2B**

and **Figure S1A-B**), and then inserting genes-of-interest (**Tables S2-4**). To create an Fc-tagged

FAST protein we fused the extracellular domain of devil CD80 to the Fc region of the devil IgG

(**Figure S1C**). The Fc region was amplified from a devil IgG plasmid provided by Lynn Corcoran

(Walter and Eliza Hall Institute of Medical Research). All secreted FAST proteins in this study

used their native signal peptides, except for 41BBL. 41BBL is a type II transmembrane protein in

which the signal peptide directly precedes the cytoplasmic and transmembrane domains of the

protein (cytoplasmic N-terminus, extracellular C-terminus). As type I FAST vectors cannot

accommodate this domain architecture, we developed an alternative base vector for type II

transmembrane FAST proteins (**Figure S1D**). To increase the probability of efficient secretion of

type II FAST proteins from CHO cells, we used the hamster IL-2 signal peptide (accession #

NM_001281629.1) at the N-terminus of the protein, followed by a SalI restriction site, mCherry,

an NheI restriction site, 6x-His tag, GSAGSAAGSGEF linker, TEV cleavage site, XmaI/SmaI

restriction site, the gene-of-interest, and a PmeI restriction site following the stop codon.

General plasmid assembly, transformation, and sequencing

Following transformation of assembled plasmids, colony PCR was performed as an initial

test of the candidate plasmids. Single colonies were inoculated directly into a OneTaq Hot Start

Quick-Load 2X Master Mix (NEB # M0488) with primers pSB_EF1a_seq.FOR

(atcttggttcattctcaagcctcag) and pSB_bGH_seq.REV (aggcacagtcgaggctgat). PCR was performed

with 60 °C annealing temperature for 25-35 cycles. Colonies yielding appropriate band sizes were

used to inoculate Luria broth with 100 μg/mL ampicillin for bacterial outgrowth overnight at 37

°C and 200 RPM. The plasmids were purified using standard plasmid kits and prepared for Sanger

sequencing using BigDye Terminator v3.1 Cycle Sequencing Kit (ThermoFisher # 4337455) with

pSB_EF1a_seq.FOR and pSB_bGH_seq.REV primers. The BigDye® Terminator was removed

using Agencourt CleanSEQ® (Beckman Coulter # A29151) before loading samples to an Applied

Biosystems® 3500XL Genetic Analyzer (Applied Biosystems) for sequencing by fluorescence-

based capillary electrophoresis.

General cell culture conditions

DFT1 cell line C5065 and DFT2 cell line JV were cultured at 35 °C with 5% CO_2 in cRF10

(10% complete RPMI (Gibco # 11875-093) with 2 mM L-glutamine, supplemented with 10% heat-

inactivated fetal bovine serum (FBS), and 1% antibiotic-antimycotic (ThermoFisher # 15240062).

RPMI without phenol red (Sigma # R7509) was used to culture FAST protein cell lines when

supernatants were collected for downstream flow cytometry assays. Devil peripheral blood cells

were cultured in cRF10 at 35 °C with 5% CO_2. CHO cells were cultured at 37 °C in cRF10 during

transfections and drug selection but were otherwise cultured at 35 °C in cRF5 (5% complete

RPMI). For production of purified recombinant proteins, stably transfected CHO cells were

cultured in suspension in spinner flasks in chemically defined, serum-free CHO Ex-Cell (Sigma #

14361C) media supplemented with 8 mM L-glutamine, 10 mM HEPES, 50 μM 2-ME, 1% (v/v)

antibiotic-antimycotic, and 1 mM sodium pyruvate and without hygromycin.

Transfection and Generation of Recombinant Cell Lines

Stable transfections of CHO and DFT cells were accomplished by adding 3 x 10^5 cells to

each well in 6-well plates in cRF10 and allowing the cells to adhere overnight. The next day, 2 μg

of plasmid DNA was added to 100 μL of PBS in microfuge tubes. Polyethylenimine (PEI) (linear,

MW 25,000; Polysciences # 23966-2) was diluted to 60 μg/mL in PBS and incubated for at least

two minutes. 100 μL of the PEI solution was added to the 100 μL of plasmid DNA in each tube to

achieve a 3:1 ratio of PEI:DNA. The solution was mixed by gentle pipetting and incubated at room

temperature for 15 minutes. Whilst the solution was incubating, the media on the CHO cells were

replaced with fresh cRF10. All 200 μL from each DNA:PEI mix was then added dropwise to the

CHO cells and gently rocked side-to-side and front-to-back to evenly spread the solution

throughout the well. The plates were then incubated overnight at 37 °C with 5% CO_2. The next

day the plates were inspected for fluorescence and then the media was removed and replaced with

cRF10 containing 1 mg/mL hygromycin (Sigma # H0654). The media was replaced with fresh

cRF10 1 mg/ml hygromycin every 2-3 days for the next seven days until selection was complete.

The cells were then maintained in 0.2 mg/mL hygromycin in cRF5 at 35 °C with 5% CO_2.

Supernatant was collected 2-3 weeks post-transfection and stored at 4 °C for two months to assess

stability of secreted FAST proteins.

Protein production and purification

16 days post-transfection the first batch of FAST protein cell lines were adapted to a 1:1

mix of cRF5 and chemically defined, serum free CHO Ex-cell media for 1-2 days to facilitate

adaptation of the adherent CHO cells to suspension culture in serum-free media. At least $5x10^7$

cells were then transferred to Proculture spinner flasks (Sigma # CLS45001L, CLS4500250) and

stirred at 75 RPM at 35 °C in 5% CO_2 on magnetic stirring platforms (Integra Bioscience #

183001). Cells were maintained at a density ranging from $5x10^5$ to $2x10^6$ cells/ml for 8-14 days.

Supernatant was collected every 2-3 days, centrifuged at 3200 RCF for 10 minutes, stored at 4 °C,

and then purified using the ÄKTA start protein purification system (GE Life Sciences #

29022094). The supernatant was diluted 1:1 with 20 mM sodium phosphate pH 7.4 and then

purified using HisTrap Excel columns (GE Life Sciences # 17-3712-05) according to the

manufacturer's instructions. Samples were passed through the columns using a flow rate of 2

mL/minute at 4 °C; all wash and elution steps were done at 1 mL/minute. Elution from HisTrap

columns (GE Life Sciences # 17-3712-05) was accomplished using 0.5 M imidazole and

fractionated into 1 ml aliquots using the Frac30 fraction collector (GE Life Sciences # 29023051).

Fluorescence of FAST proteins was checked via brief excitation (**Figure 2D**) on a blue light

transilluminator with an amber filter unit. In the case of mCherry chromogenic color was visible

(**Figure 2E**) without excitation. Fractions containing target proteins were combined and diluted to

15 mL with cold PBS, dialyzed (Sigma # PURX60005) in PBS at 4 °C, 0.22 μm sterile-filtered

(Millipore # SLGV033RS) and concentrated using Amicon Ultra centrifugal filter units (Sigma #

Z706345). The protein concentration was quantified using the 280 nm absorbance on a Nanodrop

spectrophotometer. Extinction coefficients using for each protein were calculated using the

ProtParam algorithm [81]. The proteins were then aliquoted into microfuge tubes and frozen at -80

°C until further use. The CTLA4-Fc-mCherry protein was designed, assembled, and tested

separately from the other FAST proteins and was tested directly in supernatant without

purification.

Preparation of CHO cells expressing full-length proteins for flow cytometry

CHO cells expressing full-length proteins were thawed in cRF10 and then maintained in

cRF5 with 0.2 mg/mL hygromycin. The adherent CHO cells were washed with PBS and incubated

with trypsin for 5 minutes at 37 °C to remove cells from the culture flask. Trypsin was diluted 5X

with cRF5 and centrifuged at 200 RCF for 5 minutes. Cells were resuspended in cRF5, counted

(viability > 95% in all cases), and resuspended and aliquoted for assays as described below.

Initial staining of CHO cells with 41BB FAST protein supernatants (without chloroquine)

Supernatants (cRF5) were collected from CHO cells expressing devil 41BB-extracellular

domain fused to either mCherry (pAF137), mCitrine (pAF138), mOrange (pAF164), mBFP

(pAF139), mAzurite (pAF160), mCerulean3 (pAF161), or mNeptune2 (pAF163) (**Tables S2-4**).

The supernatant was spun for 10 minutes at 3200 RCF to remove cells and cellular debris, and

then stored at 4 °C until further use. CHO cells expressing devil 41BBL (CHO.pAF56) and

untransfected CHO cells were prepared as described above. Flow cytometry tubes were loaded

with 5 x 10^4 target CHO cells/well in cRF5, centrifuged 500 RCF for 3 minutes, and then

resuspended in 200 μL of supernatant from the 41BB FAST cell lines (n=1/treatment). The tubes

were then incubated for 15 minutes at 4 °C, centrifuged at 500 RCF for 3 minutes, resuspended in

400 μL of cold FACS buffer, and stored on ice until the data were acquired on a Beckman-Coulter

521 Astrios flow cytometer (**Figure 2C**). All flow cytometry data was analyzed using FCS Express 6

522 Flow Cytometry Software version 6 (Denovo Software).

Staining CHO cells with FAST protein supernatants (without chloroquine)

U-bottom 96-well plates were loaded with 1×10^5 target CHO cells/well in cRF5,

centrifuged 500 RCF for 3 minutes, and then resuspended in 175 µL of cRF5 supernatant from

FAST cell lines collected 11 days after transfection (n=1/treatment). The plates were then

incubated for 30 minutes at room temperature, centrifuged at 500 RCF for 3 minutes, resuspended

in 200 µL of cold FACS buffer, centrifuged again and fixed with FACS fix buffer (PBS, 0.02%

NaN3, 0.4% formalin, 10g/L glucose). The cells were transferred to tubes, diluted with FACS

buffer and analyzed on a Beckman-Coulter Astrios flow cytometer (**Figure S2**).

Staining CHO cells with purified FAST proteins (with chloroquine)

Purified FAST proteins were diluted to 20 µg/mL in cRF5, aliquoted into V-bottom 96-

well transfer plates, and then stored at 37 °C until target cells were ready for staining. Target cells

were resuspended in cRF5 with 100 µM chloroquine and 100,000 cells/well were aliquoted into

U-bottom 96-well plates. 100 µL of the diluted FAST proteins (n=1/treatment, 2

timepoints/treatment) were then transferred from the V-bottom plates into the U-bottom 96-well

plates containing target cells. The final volumes and concentrations were 200 µL/well in cRF5

with 50 µM chloroquine and 2 µg/well of FAST proteins. One set of plates was incubated at 37 °C

for 30 minutes and another set of plates was incubated at 37 °C overnight. The cells were then

centrifuged 500 RCF for 3 minutes, the media decanted, and incubated for 5 minutes with 100 µL

of trypsin to dislodge adherent cells. The cells were then washed with 200 µL of cold FACS buffer,

fixed, resuspended in cold FACS buffer, and transferred to tubes for analysis on the Astrios flow

cytometer (**Figure 3B**).

Staining CHO cells with FAST supernatants (with chloroquine)

The protocol for using FAST protein supernatants was the same above as the preceding experiment except for the modifications described here. Supernatants were collected 2-3 weeks post-transfection, centrifuged at 3200 RCF for 10 minutes, and stored at 4 °C for 2 months. Prior to staining for flow cytometry, the supernatant was 0.22 μm filtered. Supernatant was then loaded into V-bottom 96-well plates to facilitate rapid transfer to staining plates and stored at 37 °C until target cells were ready for staining. Target cells were prepared as described above except for being diluted in cRF5 with 100 μM chloroquine. 2×10^5 cells/well (100 μL) were then loaded into U-bottom 96-well plates. 100 μL of FAST protein supernatant (n=1/treatment) was then transferred from the V-bottom plates to achieve 50 μM chloroquine and the cells were then incubated at 37 °C for 60 minutes. The plates were then washed, fixed, and analyzed on the Astrios flow cytometer (**Figure S3**). A similar procedure was used for staining stably-transfected DFT cells with CTLA4-Fc-mCherry, except that the supernatant was used fresh (**Figure 4D**).

Coculture assay with full-length target and FAST protein CHO cell lines (with chloroquine)

CHO cells expressing full-length CTLA4 with a C-terminal mCitrine, and CHO cells expressing full-length 41BB or 41BBL were labelled with 5 μM CFSE; CFSE and mCitrine were analyzed using the same excitation laser (488 nm) and emission filters (513/26 nm). 1×10^5 FAST protein-secreting cells were mixed with 1×10^5 target cells in cRF5 with 50 μM chloroquine and incubated overnight at 37 °C in 96-well U-bottom plates (**Figure 4A**). The next day the cells were rinsed with PBS, trypsinized, washed, fixed, and resuspended in FACS buffer prior to running flow cytometry. Cells were gated on forward and side scatter (FSC x SSC) and for singlets (FSC-H x FSC-A). (**Figure 4B**). Data shown in **Figure 4C** is representative of n=3 technical

566 replicates/treatment. Data was collected using a Beckman Coulter MoFlo Astrios and analyzed

567 using FCS Express.

Analysis of checkpoint molecule expression in DFT cells and Tasmanian devil tissues

RNAseq data was generated during previous experiments, aligned against the reference

Tasmanian devil genome Devil_ref v7.0 (GCA_000189315.1) and summarised into normalized

read counts as previously described [34,35]. RPKM-normalized read counts were produced in R using

edgeR[82]. Genes were ranked from highest RPKM-normalized count to lowest RPKM-normalized

count, and a heat map was produced for the genes of interest using the heatmap.2 function of

gplots. Heatmap color represents gene ranking among 18,788 predicted protein-coding genes in

the reference genome.

Staining DFTs cell with CD200/R FAST proteins

577 50,000 DFT cells/well were aliquoted into u-bottom 96-well plates, washed with 150 μL

578 of cRF10, and resuspended in 100 μL of warm cRF10 containing 100 μM chloroquine. 5 μg of

579 FAST protein/well was then added and mixed by pipetting. The plates were then incubated at 37

580 °C for 30 minutes. The cells were then transferred to microfuge tubes without washing, stored on

581 ice, and analyzed on a Beckman Coulter MoFlo Astrios (n=2/treatment).

Polyclonal antibody development

CD200 and 41BB FAST proteins were digested overnight with TEV protease (Sigma #

T4455) at 4 °C in PBS. The cleaved linker and 6x-His tag were then removed using a His SpinTrap

kit (GE Healthcare # 28-9321-71). Digested proteins in PBS were diluted 1:1 in Squalvax (Oz

Biosciences # SQ0010) to a final concentration of 0.1 μg/μL and was mixed using interlocked

syringes to form an emulsion. Immunization of BALB/c mice for antibody production was

approved by the University of Tasmania Animal Ethics Committee (# A0014680). Pre-immune

sera were collected prior to subcutaneous immunization with at least 50 µL of the emulsion. On

day 14 post-immunization the mice were boosted using a similar procedure. On day 50 the mice

received a booster with proteins in IFAVax (Oz Biosciences # IFA0050); mice immunized with

CD200 again received subcutaneous injections, whereas 41BB mice received subcutaneous and

intraperitoneal injections. Pre-immune and sera collected after 3X immunizations were then tested

by flow cytometry against CHO cells expressing either 41BB or CD200. CHO cells were prepared

as described above and $2x10^5$ cells were incubated with mouse serum diluted 1:200 in PBS for 30

minutes at 4 °C. The cells were then washed 2X and stained with 50 µL of anti-mouse IgG

AlexaFluor 647 diluted 1:1000 in FACS buffer. The cells were then washed 2X, stained with DAPI

to identify live cells, and analyzed on a Cyan ADP flow cytometer (**Figure 5C**). CD200 and 41BB

expression on DFT cells was tested using a procedure similar to the CHO cell staining, except the

sera used was collected after 4X immunizations and was diluted 1:500 and analyzed on the BD

FACSCanto II (**Figure 5D**).

Purification of antibodies from normal mouse serum (NMS) and anti-CD200 serum

Approximately 200 µL of normal mouse serum or anti-CD200 serum day 157 (after 4X

immunizations) were purified using a protein G SpinTrap (GE Healthcare # 28-4083-47) according

to the manufacturer's instrutions. Serum was diluted 1:1 with 20 mM sodium phosphate, pH 7.0

binding buffer, eluted with 0.1 M glycine-HCl, pH 2.7, and the pH was neutralized with 0.1 M

glycine-HCl, pH 2.7. The eluted antibodies were then concentrated using an Amicon Ultra 0.5

centrifugal until (Merck # UFC500308) by centrifuging at 14,000 RCF for 30 minutes at 4 °C and

then washing the antibodies with 400 µL of PBS twice. The protein concentration was then

quantified on a Nanodrop spectrophotometer at 280 nm using the extinction coefficients for IgG.

Testing CD200 expression on DFT cells that overexpress CD200R1

50,000 DFT cells/well were aliquoted into u-bottom 96-well plates and washed with 200

µL of cold FACS buffer. Purified polyclonal anti-CD200 was diluted to 2.5 µg/mL in cold FACS

buffer and the cells in appropriate wells were resuspend in 100 µL/well (0.25 µg/well) diluted

antibody; wells that did not receive antibody were resuspended in 100 µL of FACS buffer. The

cells were incubated on ice for 20 minutes and then washed with 200 µL of FACS buffer. Whilst

incubating, anti-mouse IgG-AF647 was diluted to 1 µg/mL in cold FACS buffer and then used to

resuspend cells in the appropriate wells. The plates were incubated on ice for 20 minutes, then

washed with 100 µL of cold FACS buffer. The cells were then resuspended in 200 µL of FACS

fix and incubated on a rocking platform at room temp for 15 minutes. The cells were then

centrifuged 500 RCF for 3 minutes at 4 °C, resuspended in 200 µL FACS buffer and stored at 4

°C until they were analyzed on a FACSCanto II (n=2/treatment).

Isolation of devil peripheral blood mononuclear cells

Blood collection from Tasmanian devils was approved by the University of Tasmania

Animal Ethics Committee (permit # A0014599) and the Tasmanian Department of Primary

Industries, Parks, Water and Environment (DPIPWE). Blood was collected from the jugular vein

and stored in EDTA tubes for transport to the lab. Blood was processed within three hours by

diluting 1:1 with serum-free RPMI and then layering onto Histopaque (Sigma # 10771) before

centrifuging at 400 RCF for 30 minutes. The interface containing the peripheral blood

mononuclear cells was then collected using a transfer pipette, diluted with 50 mL of serum-free

RPMI and centrifuged for 5 minutes at 500 RCF. Cells were washed with again with cRF10 and

then either used fresh or stored at -80 °C until further use.

Detecting DFT2 cells in PBMC using CD200

Frozen devil PBMC were thawed and cultured in cRF10 at 35 °C with 5% CO_2 for 2 hours,

cells were then washed in FACS buffer, counted and $3x10^5$ PBMC cells used per sample. DFT2.JV

cells were removed from culture flasks, counted, and $2x10^5$ cells used per sample. Samples were

incubated with 50 µL normal goat serum (Thermo Cat # 01-6201) diluted 1:200 in FACS buffer

for 15 minutes at 4 °C, 50 µL of anti-CD200 serum diluted 1:100 was added (1:200 final) for 30

minutes at 4 °C. Cells were then washed 2X and stained with 50 µL of anti-mouse IgG AlexaFluor

647 diluted to 1 µg/mL in FACS buffer for 30 minutes at 4 °C. The cells were then washed 2X,

stained with DAPI (Sigma Cat # D9542) to identify live cells, and analyzed on the BD FACSCanto

II. PBMC and DFT cells were run separately then PBMC and DFT2 mixed at a ratio of 10:1 by

volume for the combined samples (n=1/treatment) (**Figure S4A**). The experiment was repeated

(n=1/treatment), except that PBMCs and DFT cells were mixed at a 5:1 ratio (**Figure S4B**).

Staining of DFT cells in devil whole blood

DFT1.C5065 and DFT2.JV cells were labelled with 5 µM CellTrace violet (CTV) and

cultured for three days at 37 °C. On the day of the assays peripheral blood from one devil was

collected and stored at ambient temp for less than three hours. 100 µL of whole blood was aliquoted

into 15 mL tubes and stored at ambient temperature whilst DFT cells were prepared. The media

on CTV-labeled DFT cells were decanted and the cells were detached from the flask by incubating

in 2.5 mL of TrypLE Select for 5 minutes at 37 °C. The cells were washed with cRF10,

resuspended in cRF10, and counted. DFT cells were then diluted to $1x10^4$ cells/mL in cRF10 and

100 µL were aliquoted into appropriate 15 mL tubes containing 100 µL of whole blood. 1 µL of

purified anti-CD200 (0.5 µg/tube) was diluted into the appropriate tubes and incubated for 15

minutes at ambient temperature. Next, 0.5 µg/tube of anti-mouse IgG AF647 was added to each

tube. Note: 0.5 µL (0.5 µg) of concentrated secondary antibody was accidentally added directly to

the tube for the data shown in the top row and middle column of **Figure S5A;** for all other tubes

the secondary antibody was diluted 1:20 in PBS and 10 μL was added to each tube. The cells were

then incubated for 15 minutes at ambient temperature. The cells were then diluted in 1 mL

ammonium chloride red blood cell (RBC) lysis buffer (150 mM NH_4Cl, 10 mM $KHCO_3$, 0.1 mM

EDTA disodium (Na_2-$2H_2O$)) and mixed immediately gently pipetting five times. The cells were

incubated at ambient temperature for 10 minutes and then diluted with 5 mL of PBS and

centrifuged 500 RCF for 3 minutes. Some tubes contained residual RBCs, so the pellet was

vigorously resuspended in 5 mL of RBC lysis buffer, incubated for 5 minutes, diluted with 5 mL

of cold FACS buffer, and centrifuged 500 RCF for 3 minutes. The cells were then resuspended in

250 μL of FACS buffer and stored on ice until analysis on a Beckman Coulter MoFlo Astrios

(n=1/treatment). Data were analyzed in FCS Express version 6 (**Figure S5**).

The experiment above was repeated with the following modifications. DFT cells were

labelled with 5 μM of CFSE and incubated for two days at 37 °C. On the day of the assays fresh

blood was collected from two devils. Purified anti-CD200 and NMS were labeled with Zenon

mouse IgG AF647 (ThermoFisher # Z25008) and blocked with the Zenon blocking agent. $1x10^4$

CFSE-labeled DFT cells were diluted directly into 100 μL of whole blood in 15 mL tubes and 12

μL (2 μL antibody, 5 μL labeling agent, 5 μL blocking agent) of Zenon AF647-labeled purified

NMS or anti-CD200 were added directly to the cells. The cells were incubated for 30 minutes at

ambient temperature. The cells were then gently resuspended in 2.5 mL of RBC lysis buffer and

incubated for 10 minutes at ambient temperature. The cells were diluted with 10 mL of PBS and

centrifuged 500 RCF for 3 minutes. The cells were resuspended in 1.5 mL of RBC lysis buffer and

incubated for another 10 minutes to lyse residual RBCs. The tubes were then resuspended in 9 mL

of cRF10 and centrifuged 500 RCF for 3 minutes. The cells were resuspended in 350 μL of cold

FACS buffer containing 200 ng/mL of DAPI and stored on ice until analysis on a Beckman Coulter

MoFlo Astrios (n=1/treatment for n = 2 devils) (**Figure 6**).

ACKNOWLEDGMENTS

We wish to thank Ginny Ralph for her ongoing care of Tasmanian devils, and the Bonorong

Wildlife Sanctuary for providing access to Tasmanian devils, and Ruth Pye for providing care for

devils and collecting blood samples. We would like to thank John Hayball and Georgina

Kalodimos for advice and assistance, Mahalia Kingsley and Nirdesh Poudel for plasmid

construction, Nick Blackburn for bioinformatics assistance, Lynn Corcoran for the devil IgG

plasmid, James Murphy for a devil IL-15 plasmid, Emily Flies for editing the manuscript, and

Kay Holekamp for her comments on the manuscript. **Funding:** ARC DECRA grant #

DE180100484, ARC Linkage grant # LP0989727, ARC Discovery grant # DP130100715, Morris

Animal Foundation Grant-in-Aid # D14ZO-410, University of Tasmania Foundation Dr Eric

Guiler Tasmanian Devil Research Grant through funds raised by the Save the Tasmanian Devil

Appeal (2013, 2015, 2017, 2018), and Entrepreneurs' Programme - Research Connections grant

with Nexvet Australia Pty. Ltd. # RC50680.

AUTHOR CONTRIBUTIONS

ASF designed the study; ALP, ASF, CEBO, PRL, and PRM developed the technology; ASF,

CEBO, PRL, PRM, JMD, and TLP performed the experiments; ALP performed bioinformatic

analyses; ALP, ASF, JMD and PRL created the figures; ALP, ASF, PRL, JMD, TLP, and GMW

analyzed the data; ASF wrote the manuscript and all authors edited the manuscript.

AVAILABILITY OF DATA AND MATERIALS

All data are included with the paper. The FAST base vectors (pAF92.3 pAF112.7, pAF123.1,

pAF137.4c1, pAF138.7, pAF139.2, pAF160.1, pAF161.3, pAF163.1, pAF164.3) will be available

through Addgene (deposit # 77504).

CONFLICT OF INTEREST

The authors received funding from Nexvet Australia Pty. Ltd for related studies.

REFERENCES

Albuquerque TAF, Drummond do Val L, Doherty A, de Magalhães JP. From humans to hydra: patterns of cancer across the tree of life. *Biol Rev* 2018; **93**: 1715–1734.

Abu-Helil B, Van Der Weyden L. Metastasis in the wild: investigating metastasis in non-laboratory animals. *Clin Exp Metastasis* 2019. doi:10.1007/s10585-019-09956-3.

Olds JE, Burrough ER, Fales-Williams AJ, Lehmkuhl A, Madson D, Patterson AJ *et al.* Retrospective Evaluation of Cases of Neoplasia in a Captive Population of Egyptian Fruit Bats (Rousettus Aegyptiacus). *J Zoo Wildl Med* 2015; **46**: 325–332.

Chu PY, Zhuo YX, Wang FI, Jeng CR, Pang VF, Chang PH *et al.* Spontaneous neoplasms in zoo mammals, birds, and reptiles in Taiwan - A 10-year survey. *Anim Biol* 2012; **62**: 95–110.

Deus A De, Alves F, Siqueira DB De, Rameh-de-albuquerque LC. Breast Carcinoma with Pulmonary Metastasis in Armadillo (Eupharactus sexcinctus). *Acta Sci Vet* 2018; **46**: 1–5.

Kruse TN, Garner MM, Bonar CJ. A Retrospective Study of Pathologic Findings in Captive Rock Hyrax (Procavia Capensis) in the United States. *J Zoo Wildl Med* 2015; **46**: 798–805.

Effron M, Griner L, Benirschke K. Nature and rate of neoplasia found in captive wild mammals, birds, and reptiles at necropsy. *J Natl Cancer Inst* 1977; **59**: 185–198.

Ratcliffe HL. Incidence and nature of tumors in captive wild mammals and birds. *Am J Cancer* 1933; **17**: 116–135.

Priester WA, Mantel N. Occurrence of tumors in domestic animals. Data from 12 United States and Canadian colleges of veterinary medicine. *J Natl Cancer Inst* 1971; **47**: 1333–1344.

10   Howlader N, Noone A, Krapcho M, Miller D, Brest A, Yu M *et al.* SEER Cancer

Statistics Review, 1975-2016. Bethesda, MD, United States, 2016.

11   Griner LA. Neoplasms in Tasmanian devils (Sarcophilus harrisii). *J Natl Cancer Inst*

1979; **62**: 589–595.

12   Peck SJ, Michael SA, Knowles G, Davis A, Pemberton D. Causes of mortality and severe

morbidity requiring euthanasia in captive Tasmanian devils (Sarcophilus harrisii) in

Tasmania. *Aust Vet J* 2019; **97**: 89–92.

13   Pearse A-M, Swift K. Allograft theory: transmission of devil facial-tumour disease.

Nature 2006; **439**: 549.

14   Pye RJ, Pemberton D, Tovar C, Tubio JMC, Dun KA, Fox S *et al.* A second transmissible

cancer in Tasmanian devils. *Proc Natl Acad Sci* 2016; **113**: 374–379.

15   Murgia C, Pritchard JK, Kim SY, Fassati A, Weiss RA. Clonal Origin and Evolution of a

Transmissible Cancer. *Cell* 2006; **126**: 477–487.

16   Fleming JM, Creevy KE, Promislow DEL. Mortality in North American Dogs from 1984

to 2004: An Investigation into Age-, Size-, and Breed-Related Causes of Death. *J Vet*

Intern Med 2011; **25**: 187–198.

17   Lazenby BT, Tobler MW, Brown WE, Hawkins CE, Hocking GJ, Hume F *et al.* Density

trends and demographic signals uncover the long-term impact of transmissible cancer in

Tasmanian devils. *J Appl Ecol* 2018; **55**: 1368–1379.

18   James S, Jennings G, Kwon YM, Stammnitz M, Fraik A, Storfer A *et al.* Tracing the rise

of malignant cell lines: distribution, epidemiology and evolutionary interactions of two

transmissible cancers in Tasmanian devils. *Evol Appl* 2019; : eva.12831.

19   Siddle H V., Kreiss A, Tovar C, Yuen CK, Cheng Y, Belov K *et al.* Reversible epigenetic

down-regulation of MHC molecules by devil facial tumour disease illustrates immune

escape by a contagious cancer. *Proc Natl Acad Sci* 2013; **110**: 5103–8.

20 Yoshihama S, Roszik J, Downs I, Meissner TB, Vijayan S, Chapuy B *et al.* NLRC5/MHC

class I transactivator is a target for immune evasion in cancer. *Proc Natl Acad Sci* 2016;

113: 5999–6004.

21 Caldwell A, Coleby R, Tovar C, Stammnitz MR, Mi Kwon Y, Owen RS *et al.* The newly-

arisen devil facial tumour disease 2 (DFT2) reveals a mechanism for the emergence of a

contagious cancer. *Elife* 2018; **7**. doi:10.7554/eLife.35314.

22 Stammnitz MR, Coorens THH, Gori KC, Hayes D, Fu B, Wang J *et al.* The Origins and

Vulnerabilities of Two Transmissible Cancers in Tasmanian Devils. *Cancer Cell* 2018;

33: 607-619.e15.

23 Topalian SL, Hodi FS, Brahmer JR, Gettinger SN, Smith DC, McDermott DF *et al.*

Safety, Activity, and Immune Correlates of Anti–PD-1 Antibody in Cancer. *N Engl J Med*

2012; **366**: 2443–2454.

24 Larkin J, Chiarion-Sileni V, Gonzalez R, Grob JJ, Cowey CL, Lao CD *et al.* Combined

Nivolumab and Ipilimumab or Monotherapy in Untreated Melanoma. *N Engl J Med* 2015;

373: 23–34.

25 Flies AS, Bruce Lyons A, Corcoran LM, Papenfuss AT, Murphy JM, Knowles GW *et al.*

PD-L1 is not constitutively expressed on tasmanian devil facial tumor cells but is strongly

upregulated in response to IFN-γ and can be expressed in the tumor microenvironment.

Front Immunol 2016; **7**: 581.

26 Ong CEB, Lyons AB, Woods GM, Flies AS. Inducible IFN-γ Expression for MHC-I

Upregulation in Devil Facial Tumor Cells. *Front Immunol* 2018; **9**: 3117.

27   Flies AS, Blackburn NB, Lyons AB, Hayball JD, Woods GM. Comparative analysis of

781   immune checkpoint molecules and their potential role in the transmissible tasmanian devil

782   facial tumor disease. *Front Immunol* 2017; **8**: 513.

783   28   World Health Organization (WHO). WHO R&D Blueprint for action to prevent

784   epidemics. World Health Organization, 2016http://www.who.int/blueprint/en/ (accessed 1

785   May2018).

786   29   Shaner NC, Campbell RE, Steinbach PA, Giepmans BNG, Palmer AE, Tsien RY.

787   Improved monomeric red, orange and yellow fluorescent proteins derived from

788   Discosoma sp. red fluorescent protein. *Nat Biotechnol* 2004; **22**: 1567–72.

789   30   Cabantous S, Terwilliger TC, Waldo GS. Protein tagging and detection with engineered

790   self-assembling fragments of green fluorescent protein. *Nat Biotechnol* 2005; **23**: 102–

791   107.

792   31   Cha HJ, Dalal NN, Bentley WE. Secretion of human interleukin-2 fused with green

793   fluorescent protein in recombinant Pichia pastoris. *Appl Biochem Biotechnol* 2005; **126**:

794   1–11.

795   32   Duellman T, Burnett J, Yang J. Quantitation of secreted proteins using mCherry fusion

796   constructs and a fluorescent microplate reader. *Anal Biochem* 2015; **473**: 34–40.

797   33   Kammertoens T, Friese C, Arina A, Idel C, Briesemeister D, Rothe M *et al.* Tumour

798   ischaemia by interferon-γ resembles physiological blood vessel regression. *Nature* 2017;

799   **545**: 98–102.

800   34   Patchett AL, Wilson R, Charlesworth JC, Corcoran LM, Papenfuss AT, Lyons AB *et al.*

801   Transcriptome and proteome profiling reveals stress-induced expression signatures of

802   imiquimod-treated Tasmanian devil facial tumor disease (DFTD) cells. *Oncotarget* 2018;

9: 15895–15914.

35 Patchett AL, Coorens THH, Darby J, Wilson R, McKay MJ, Kamath KS *et al.* Two of a

kind: transmissible Schwann cell cancers in the endangered Tasmanian devil (Sarcophilus

harrisii). *Cell Mol Life Sci* 2019; : 1–12.

36 Seglen PO, Grinde B, Solheim AE. Inhibition of the Lysosomal Pathway of Protein

Degradation in Isolated Rat Hepatocytes by Ammonia, Methylamine, Chloroquine and

Leupeptin. *Eur J Biochem* 1979; **95**: 215–225.

37 Qureshi OS, Zheng Y, Nakamura K, Attridge K, Manzotti C, Schmidt EM *et al.* Trans-

endocytosis of CD80 and CD86: a molecular basis for the cell extrinsic function of

CTLA-4. *Science* 2011; **332**: 600–603.

38 van den Bremer ET, Beurskens FJ, Voorhorst M, Engelberts PJ, de Jong RN, van der

Boom BG *et al.* Human IgG is produced in a pro-form that requires clipping of C-terminal

lysines for maximal complement activation. *MAbs* 2015; **7**: 672–680.

39 Patchett AL, Latham R, Brettingham-Moore KH, Tovar C, Lyons AB, Woods GM. Toll-

like receptor signaling is functional in immune cells of the endangered Tasmanian devil.

Dev Comp Immunol 2015; **53**: 123–133.

40 Love JE, Thompson K, Kilgore MR, Westerhoff M, Murphy CE, Papanicolau-Sengos A

et al. CD200 Expression in Neuroendocrine Neoplasms. *Am J Clin Pathol* 2017; **148**:

236–242.

41 Murchison EP, Tovar C, Hsu A, Bender HS, Kheradpour P, Rebbeck CA *et al.* The

Tasmanian devil transcriptome reveals schwann cell origins of a clonally transmissible

cancer. *Science* 2010; **327**: 84–87.

42 Haile ST, Bosch JJ, Agu NI, Zeender AM, Somasundaram P, Srivastava MK *et al.* Tumor

cell programmed death ligand 1-mediated T cell suppression is overcome by coexpression

of CD80. *J Immunol* 2011; **186**: 6822–9.

43   Sugiura D, Maruhashi T, Okazaki I, Shimizu K, Maeda TK, Takemoto T *et al.* Restriction

of PD-1 function by *cis* -PD-L1/CD80 interactions is required for optimal T cell

responses. *Science* 2019; : eaav7062.

44   Moertel CL, Xia J, LaRue R, Waldron NN, Andersen BM, Prins RM *et al.* CD200 in CNS

tumor-induced immunosuppression: The role for CD200 pathway blockade in targeted

immunotherapy. *J Immunother Cancer* 2014; **2**: 46.

45   Challagundla P, Medeiros LJ, Kanagal-Shamanna R, Miranda RN, Jorgensen JL.

Differential Expression of CD200 in B-Cell Neoplasms by Flow Cytometry Can Assist in

Diagnosis, Subclassification, and Bone Marrow Staging. *AJCP / Orig Artic Am J Clin*

Pathol 2014; **142**: 837–844.

46   Spacek M, Karban J, Radek M, Babunkova E, Kvasnicka J, Jaksa R *et al.* CD200

Expression Improves Differential Diagnosis Between Chronic Lymphocytic Leukemia

and Mantle Cell Lymphoma. *Blood* 2014; **124**.

47   Saksena A, Yin CC, Xu J, Li J, Zhou J, Wang SA *et al.* CD23 expression in mantle cell

lymphoma is associated with CD200 expression, leukemic non-nodal form, and a better

prognosis. *Hum Pathol* 2019; **89**: 71–80.

48   Loh R, Hayes D, Mahjoor A, O'Hara A, Pyecroft S, Raidal S. The immunohistochemical

characterization of devil facial tumor disease (DFTD) in the Tasmanian Devil

(Sarcophilus harrisii). *Vet Pathol* 2006; **43**: 896–903.

49   Yao S, Zhu Y, Zhu G, Augustine M, Zheng L, Goode DJ *et al.* B7-h2 is a costimulatory

ligand for CD28 in human. *Immunity* 2011; **34**: 729–740.

50   Wang J, Sanmamed MF, Datar I, Su TT, Ji L, Sun J *et al.* Fibrinogen-like Protein 1 Is a

Major Immune Inhibitory Ligand of LAG-3. *Cell* 2018; **0**.

doi:10.1016/J.CELL.2018.11.010.

51   Aricescu AR, Lu W, Jones EY. A time- and cost-efficient system for high-level protein

production in mammalian cells. *Acta Crystallogr Sect D Biol Crystallogr* 2006; **62**: 1243–

1250.

52   Consortium SG, Biologiques A et F des M, Center BSG, Consortium CSG, Innovation IC

for S and F, Center ISP *et al.* Protein production and purification. *Nat Methods* 2008; **5**:

135.

53   Jayapal KP, Wlaschin KF, Hu WS, Yap MGS. Recombinant protein therapeutics from

CHO cells - 20 years and counting. *Chem Eng Prog* 2007; **103**: 40–47.

54   Atfy M. CD200 Suppresses the Natural Killer Cells and Decreased its Activity in Acute

Myeloid Leukemia Patients. *J Leuk* 2015; **3**.https://www.omicsonline.org/open-

access/cd200-suppresses-the-natural-killer-cells-and-decreased-its-activity-in-

acutemyeloid-leukemia-patients-2329-6917-1000190.pdf (accessed 18 Apr2019).

55   Coles SJ, Man S, Hills R, Wang EC, Burnett A, Darley RL *et al.* Over-Expression of

CD200 In Acute Myeloid Leukemia Mediates the Expansion of Regulatory T-

Lymphocytes and Directly Inhibits Natural Killer Cell Tumor Immunity. *Blood* 2015; **116**:

491.

56   Coles SJ, Wang ECY, Man S, Hills RK, Burnett AK, Tonks A *et al.* CD200 expression

suppresses natural killer cell function and directly inhibits patient anti-tumor response in

acute myeloid leukemia. *Leukemia* 2011; **25**: 792–9.

57   Gorczynski RM, Chen Z, Khatri I, Yu K. Graft-infiltrating cells expressing a CD200

872  transgene prolong allogeneic skin graft survival in association with local increases in

873  Foxp3 +Treg and mast cells. *Transpl Immunol* 2011; **25**: 187–193.

874 58 Gorczynski RM, Chen Z, He W, Khatri I, Sun Y, Yu K *et al.* Expression of a CD200

875  transgene is necessary for induction but not maintenance of tolerance to cardiac and skin

876  allografts. *J Immunol* 2009; **183**: 1560–1568.

877 59 Gorczynski L, Chen Z, Hu J, Kai Y, Lei J, Ramakrishna V *et al.* Evidence That an OX-2-

878  Positive Cell Can Inhibit the Stimulation of Type 1 Cytokine Production by Bone

879  Marrow-Derived B7-1 (and B7-2)-Positive Dendritic Cells. *J Immunol* 1999; **162**: 774–

880  781.

881 60 Harding J, Vintersten-Nagy K, Shutova M, Yang H, Tang JK, Massumi M *et al.* Induction

882  of long-term allogeneic cell acceptance and formation of immune privileged tissue in

883  immunocompetent hosts. Cold Spring Harbor Laboratory, 2019 doi:10.1101/716571.

884 61 Jenmalm MC, Cherwinski H, Bowman EP, Phillips JH, Sedgwick JD. Regulation of

885  Myeloid Cell Function through the CD200 Receptor. *J Immunol* 2014; **176**: 191–199.

886 62 Tovar C, Pye RJ, Kreiss A, Cheng Y, Brown GK, Darby J *et al.* Regression of devil facial

887  tumour disease following immunotherapy in immunised Tasmanian devils. *Sci Rep* 2017;

888  **7**: 43827.

889 63 Pye R, Patchett A, McLennan E, Thomson R, Carver S, Fox S *et al.* Immunization

890  strategies producing a humoral IgG immune response against devil facial tumor disease in

891  the majority of Tasmanian devils destined for wild release. *Front Immunol* 2018; **9**: 259.

892 64 Olin MR, Ampudia-Mesias E, Pennell CA, Sarver A, Chen CC, Moertel CL *et al.*

893  Treatment combining CD200 immune checkpoint inhibitor and tumor-lysate vaccination

894  after surgery for pet dogs with high-grade glioma. *Cancers (Basel)* 2019; **11**: 137.

895   65   Zhang S, Phillips JH. Identification of tyrosine residues crucial for CD200R-mediated

896   inhibition of mast cell activation. *J Leukoc Biol* 2005; **79**: 363–368.

897   66   Hatherley D, Lea SM, Johnson S, Barclay AN. Structures of CD200/CD200 receptor

898   family and implications for topology, regulation, and evolution. *Structure* 2013; **21**: 820–

899   832.

900   67   Luo ZX, Yuan CX, Meng QJ, Ji Q. A Jurassic eutherian mammal and divergence of

901   marsupials and placentals. *Nature* 2011; **476**: 442–445.

902   68   Wong KK, Khatri I, Shaha S, Spaner DE, Gorczynski RM. The role of CD200 in

903   immunity to B cell lymphoma. *J Leukoc Biol* 2010; **88**: 361–372.

904   69   Caserta S, Nausch N, Sawtell A, Drummond R, Barr T, MacDonald AS *et al.* Chronic

905   infection drives expression of the inhibitory receptor CD200R, and its ligand CD200, by

906   mouse and human CD4 T cells. *PLoS One* 2012; **7**: e35466.

907   70   Foster-Cuevas M, Westerholt T, Ahmed M, Brown MH, Barclay AN, Voigt S.

908   Cytomegalovirus e127 protein interacts with the inhibitory CD200 receptor. *J Virol* 2011;

909   **85**: 6055–9.

910   71   Foster-Cuevas M, Wright GJ, Puklavec MJ, Brown MH, Barclay AN. Human herpesvirus

911   8 K14 protein mimics CD200 in down-regulating macrophage activation through CD200

912   receptor. *J Virol* 2004; **78**: 7667–76.

913   72   Alves JM, Carneiro M, Cheng JY, Matos AL de, Rahman MM, Loog L *et al.* Parallel

914   adaptation of rabbit populations to myxoma virus. *Science* 2019; : eaau7285.

915   73   Barclay AN, Hatherley D. The Counterbalance Theory for Evolution and Function of

916   Paired Receptors. *Immunity* 2008; **29**: 675–678.

74   Tovar C, Obendorf D, Murchison EP, Papenfuss AT, Kreiss A, Woods GM. Tumor-

specific diagnostic marker for transmissible facial tumors of tasmanian devils:

Immunohistochemistry studies. *Vet Pathol* 2011; **48**: 1195–1203.

75   Marx V. Calling the next generation of affinity reagents. *Nat Methods* 2013; **10**: 829–833.

76   Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I *et al.* Full-length

transcriptome assembly from RNA-Seq data without a reference genome. *Nat Biotechnol*

2011; **29**: 644–52.

77   Kowarz E, Löscher D, Marschalek R. Optimized Sleeping Beauty transposons rapidly

generate stable transgenic cell lines. *Biotechnol J* 2015; **10**: 647–653.

78   Mátés L, Chuah MKL, Belay E, Jerchow B, Manoj N, Acosta-Sanchez A *et al.* Molecular

evolution of a novel hyperactive Sleeping Beauty transposase enables robust stable gene

transfer in vertebrates. *Nat Genet* 2009; **41**: 753–761.

79   Waldo GS, Standish BM, Berendzen J, Terwilliger TC. Rapid protein-folding assay using

green fluorescent protein. *Nat Biotechnol* 1999; **17**: 691–5.

80   Kozak M. An analysis of 5'-noncoding sequences from 699 vertebrate messenger RNAs.

Nucleic Acids Res 1987; **15(20)**: 8125–8148.

81   Gasteiger E, Hoogland C, Gattiker A, Duvaud S, Wilkins MR, Appel RD *et al.* Protein

identification and analysis tools in the ExPASy server. In: Walker JM (ed). *The*

Proteomics Protocols Handbook. Humana Press, 2005, pp 571–607.

82   Robinson M, McCarthy D, Smyth G. edgeR: a Bioconductor package for differential

expression analysis of digital gene expression data. *Bioinformatics* 2010; **26**: 139–140.

FIGURES

Figure 1. Phylogenetic tree of immune system related studies in mammal orders 2009-2019.

Metastatic cancer has been reported in nearly all mammalian orders and major histocompatibility

complexes (MHC) have been the most intensely studied molecules in most orders. In the past

decade, studies of immune checkpoint molecules (PD1, PDL1, CTLA4) have become a primary

focus in humans and rodents. However, immune checkpoint studies in other species are limited,

particularly at the protein level, due to the lack of species-specific reagents. This creates a vast gap

in our understanding of the evolution of the mammalian immune system. The numbers in the

columns represent the number studies matching Web of Science search results between 2009-2019.

See Table S1 for search terms.

Figure 2. FAST protein schematic and initial testing. (A) Schematic diagram of FAST protein

therapeutic and diagnostic (i.e. theranostic) features and (B) vector map showing restriction sites

for swapping the target gene (i.e. gene-of-interest) and color genes (i.e. fluorescent protein). (C)

Results of flow cytometry binding assay with devil 41BB FAST proteins. The colored lines in

the histograms show binding of devil 41BB fused to mCherry, mCitrine, mOrange, mTagBFP,

mAzurite, mCerulean3, or mNeptune2 to CHO cells transfected with devil 41BBL, and the black

lines show binding to untransfected CHO cells. (D-E) Images showing the gradient of FAST

proteins eluted from HisTrap columns. (D) mCitrine excited with blue light and (E) chromogenic

visualization of mCherry without excitation. (F) Image of 100 μL of FAST protein excited with

blue light (NOTE: mBFP appears clear with blue excitation and amber filter unit).

Figure 3. Map and testing of soluble proteins FAST proteins. (A) Diagram of soluble FAST proteins and full-length proteins used for testing of FAST proteins. 41BBL is a type II transmembrane protein; all other proteins are type I. CD80 and CTLA4 soluble FAST proteins included a devil IgG Fc-tag. Arrows indicate interactions confirmed in this study. (B) Histograms showing binding of FAST proteins to CHO cells expressing full-length devil proteins. Target CHO cells were cultured with chloroquine to block lysosomal degradation of FAST proteins and maintain fluorescent signal during live-culture binding assays with 2 μg/well of purified FAST proteins for 30 minutes or 18 hours to assess receptor-ligand binding (n=1/time point).

Figure 4. Live-cell coculture assays with FAST proteins. (A) Schematic of coculture assays to assess checkpoint molecule interactions (Absent, Weak, Strong). Cells were mixed and cultured overnight with chloroquine. Protein binding and/or transfer were assessed using flow cytometry. (B) Gating strategy for coculture assays. (C) CHO cells that secrete 41BBL-mCherry, 41BB-mCherry, or CD80-Fc-mCherry were cocultured overnight with target CHO cells that express full-length 41BB, 41BBL, or CTLA4. 41BB and 41BB-L were labeled with CFSE, whereas full-length CTLA4 was directly fused to mCitrine. Cells that secrete mCherry FAST proteins appear in the upper left quadrant. Cells expressing full-length proteins and labeled with CFSE or mCitrine appear in the lower right quadrant. Cells in the upper right quadrant represent binding of mCherry FAST proteins to full-length proteins on CFSE or mCitrine labeled cells. Results shown are representative of n=3/treatment. (D) CTLA4-Fc-mCherry FAST protein binding to DFT cells. DFT1 C5065 cells transfected with control vector (black), 41BB (gray), CD80 (red), or CD86 (blue) were stained with CTLA4-Fc-mCherry supernatant with chloroquine. Results are representative of n=2 replicates/treatment.

Figure 5. Elevated CD200 expression on DFT cells. (A) Heatmap showing within sample transcript ranking (1 = highest expression) according to RPKM-normalized mRNA sequencing counts of 18,788 annotated coding genes (devil_refv7.0; GCA_000189315.1). Genes-of-interest for this study are plotted as heatmap with dark blue indicating the most highly expressed genes. Technical replicates (n=2) were used for all tissues, except peripheral nerve (PN) (n=1). (B) Wild type DFT1.C5065, DFT2.JV, DFT2.SN, and DFT1.C5065 transfected to overexpress CD200 or CD200R1 were stained with 5 µg of either CD200R1-mOrange or CD200-mOrange FAST protein. Histograms filled with blue or red highlight the cells overexpressing CD200 or CD200R1 and their expected binding interactions with CD200R1 and CD200, respectively. Target cells were cultured

with FAST proteins in chloroquine, incubated 30 minutes, and run without washing. Results are

representative of n=2 replicates/treatment. (C) Mice were immunized with TEV digested 41BB or

CD200 FAST proteins. Black = pre-immune (PI) and gray = immune (I) sera from a mouse

immunized with 41BB; red = pre-immune (PI) and blue = immune (I) sera from a mouse

immunized with CD200. CHO cells transfected with either full-length 41BB or CD200 were

stained with sera and then anti-mouse AlexaFluor-647. Results are representative of

n=2/treatment. (D) Sera was used to screen two strains of DFT1 and two strains of DFT2 cells for

41BB and CD200 expression. PI sera was negative in all cases, whereas all DFT1 and DFT2 cells

expressed CD200. Results are representative of n=3/treatment. (E) DFT1 C5065 transfected with

either vector control, CD200, or CD200R1 were stained with purified polyclonal anti-CD200 and

anti-mouse IgG AlexaFluor 647 (black = no antibodies, red = secondary antibody only, blue =

primary + secondary antibody). Results are representative of n=2/treatment.

1015

1016

1017

Figure 6. CD200 identifies DFT cells in whole blood. Color dot plots showing DFT cells in green

(CFSE), PBMCs in black, DFT Alexa Fluor 647+ (AF647) cells in red, and PBMC AF647+ in

blue. (A) Forward- and side-scatter plot of DFT.JV cells and (B) DFT.JV cells mixed with PBMCs.

(C) Color dot plot showing dead cells stained with DAPI (right quadrants) and CFSE-labeled DFT

cells (upper quadrants). (D) The top row shows unmixed PBMCs. The middle row and bottom row

show DFT1.C5065 and DFT2.JV cells, respectively, mixed with PBMCs. Alexa Fluor 647+ DFT

(red) and PBMC (blue) are in the right quadrants. (E) Histogram overlays to highlight AF647+

(right quadrants) from DFT1-PBMC and DFT2-PBMC mixtures. Cells were analyzed on the

Beckman-Coulter MoFlo Astrios.

TABLES (See supplementary Materials)

Group		Cancer	Checkpoint	MHC
Human		47,927	6,511	5,493
Primatomorpha		860	95	324
Lagomorpha		9	0	4
Rodentia		14,337	1,665	1,807
Cetartiodactyla		11	1	22
Artiodactyla		431	29	234
Perissodactyla		211	8	72
Carnivora		494	34	85
Pholidoea		0	0	0
Chiroptera		5	0	7
Eulipotyphla		6	0	6
Xenarthra		4	0	1
Hyracoidea		0	0	0
Proboscidea		9	0	3
Sirenia		4	0	2
Marsupial		78	3	52
Monotreme		5	0	2