1

Full title: Loss of pseudouridine synthases in the RluA family causes hypersensitive

nociception in *Drosophila*

Short title: Pseudouridine synthase RluA in nociception

Wan Song[1,2] & W. Daniel Tracey[1,2,*]

[1]Gill Center for Biomolecular Research, Indiana University, Bloomington, Indiana, USA

[2]Department of Biology, Indiana University, Bloomington, Indiana, USA

[*] Corresponding author

Email: dtracey@indiana.edu

Abstract

Nociceptive neurons of *Drosophila* larvae are characterized by highly branched dendritic processes whose proper morphogenesis relies on a large number of RNA-binding proteins. Post-transcriptional regulation of RNA in these dendrites has been found to play an important role in their function. Here, we investigate the neuronal functions of two putative RNA modification genes, *RluA-1* and *RluA-2*, which are predicted to encode pseudouridine synthases. *RluA-1* is specifically expressed in larval sensory neurons while *RluA-2* expression is ubiquitous. Nociceptor-specific RNAi knockdown of *RluA-1* caused hypersensitive nociception phenotypes, which were recapitulated with genetic null alleles. These were rescued with genomic duplication and nociceptor-specific expression of *UAS-RluA-1-cDNA*. As with *RluA-1*, *RluA-2* loss of function mutants also displayed hyperalgesia. Interestingly, nociceptor neuron dendrites showed a hyperbranched morphology in the *RluA-1* mutants. The latter may be a cause or a consequence of heightened sensitivity in mutant nociception behaviors.

Author Summary

Pseudouridine (Psi) is a C5-glycoside isomer of uridine and it is the most common posttranscriptional modification of RNAs, including noncoding tRNAs, rRNAs, snRNAs as well as mRNAs. Although first discovered in the 1950s, the biological functions of Psi in multicellular organisms are not well understood. Interestingly, a marker for sensory neurons in *Drosophila* encodes for a putative pseudouridine synthase of unknown function called RluA-1. Here, we report our characterization of nociception phenotypes for larvae with RluA-1 loss of function along with that of a related gene RluA-2. Disrupting either or both RluA-1 and RluA-2 resulted in hypersensitive nociception. In addition, RluA-1 mutants have more highly branched nociceptor neurites that innervate the epidermis. Our studies suggest an important role for the

RluA family in nociception. This may be through its action on RNAs that regulate neuronal

excitability and/or dendrite morphogenesis.

Introduction

Pain serves an indispensable, protective role but when pain becomes pathological it can

have a debilitating impact on human life. The total annual cost of pain to society in the United

States was estimated by the Institute of Medicine to be up to $635 billion, which is greater than

that of heart disease, cancer, and diabetes combined (1). It is therefore of urgent importance to

uncover the basic molecular and cellular mechanisms involved in pain in order to better treat it.

Laboratory animal models of pain and nociception have played an essential role in identifying

such mechanisms. *Drosophila* larvae respond to noxious thermal and mechanical stimuli

through stereotyped rolling escape locomotion (in which the larva rotates around its long body

axis), which is robust and easily distinguishable from other forms of locomotion (2). Rolling is

used as a behavioral defense against attacks by parasitoid wasps which lay their eggs within the

fly larvae using a sharp ovipositor (3). Successful wasp infection is lethal to the larvae and is a

strong selective pressure in natural populations of *Drosophila*. Combined with the unparalleled

genetic tools available for *Drosophila melanogaster*, this behavioral readout provides an

excellent system to study the genetics of nociception and pain (2, 4-8). Previous studies have

demonstrated a specific subset of dendritic arborization (da) sensory neurons in the peripheral

nervous system, Class IV multidendritic da (cIVda) neurons, are of critical importance for

thermal, mechanical and high intensity light nociception (3, 9, 10). Further evidence suggests a

lesser but significant contribution of Class II (cIIda) and Class III da (cIIIda) neurons in mechanical

nociception (3, 11). In addition, great progress has been made in identifying the circuits in the

larval abdominal ganglion that are involved in rolling escape locomotion (12-15).

Genes that are specifically expressed in multidendritic (md) neurons have been found to

play a role in nociception. For instance, *painless* (2) is required for mechanical and thermal

nociception and it is expressed in all four classes of md neurons. Mechanical nociception genes

such as *pickpocket* (16), *ppk26/balboa* (17, 18) and the polymodal nociception gene *dTRPA1-C/D*

(19) each show very specific expression in the cIVda neurons. Forward genetic screens have also

identified a set of genes with enriched expression in the cIVda neurons which either inhibit or

activate nociceptive pathways (20).

The historically first known genetic marker with specific expression pattern in the da

neurons was the *lacZ* enhancer trap *E7-2-36* (21). Later studies reported that this enhancer trap

gene was inserted upstream of the *RluA-1* gene and DNA sequences from upstream of *RluA-1*

caused expression of GAL4 in multidendritic neurons (22). The *RluA1* gene encodes an enzyme

that is predicted to have pseudouridine synthase activity, but how this RNA modifying protein

is involved in the function of nociceptive multidendritic neurons remains unknown.

A widespread importance of RNA-binding proteins in cIVda neuron dendrite

morphogenesis and function was found in a recent large-scale RNAi screen that identified 88

genes encoding RNA-binding proteins whose knockdown caused aberrant dendrite

morphogenesis (23). The elaborate dendrite arbors of cIVda neurons project long distances

from the neuronal cell body and mRNA granules are trafficked to these distant sites where they

may undergo local translation. Indeed, RNA granules containing Nanos (Nos), Pumilio (Pum),

Oskar (Osk), Fragile X Mental Retardation (FMRP) and other proteins have been shown to

regulate the formation of higher order dendrites in these cells (24-28). Interestingly, the *nos*

mRNA itself is transported into dendrites via a pathway that relies on Oskar and

Rumplestiltskin, two proteins that also play a role regulating *nos* in the early embryo (28).

Mechanical nociception defects are also observed in animals with disruption in these pathways

(28).

Pseudouridylation is the most common post-transcriptional RNA modification.

Pseudouridine (Ψ), the C5-glycoside isomer of uridine, was initially found in many positions in

rRNA, tRNA and snRNA in all organisms that have been investigated (29). RNA-seq based

global pseudouridine profiling has shown the presence of Ψ in many mRNAs and a large

number of those sites were found to be dynamically regulated in yeast and human cells (30-33).

Dysfunctional pseudouridylation has been linked to several human diseases (34-36). Since many

sites of pseudouridylation in different organisms are evolutionarily conserved, *Drosophila*

melanogaster may provide an excellent and genetically tractable metazoan system to elucidate

some of these functions.

The isomerization of uridine to pseudouridine is catalyzed by six families of

pseudouridine synthases. They function either as guide RNA directed ribonucleoprotein

complexes or as stand-alone proteins (37, 38). In the *Drosophila* genome, 9 proteins have been

identified with annotated pseudouridine synthase domains. Minifly (mfl), the RNA-dependent

pseudouridine synthase homolog of human dyskerin (mouse NAP57 and yeast Cbf5), is required

for somatic stem cell homeostasis and is essential for *Drosophila* viability and fertility (39-41).

Knockout of *Drosophila* Pus7, the human and yeast Pus7 homolog, results in increased

aggressiveness in adult flies (36). The function and specificity of other predicted pseudouridine

synthases are largely unknown. Among the six families, the RluA family, which does not rely on

guide RNAs, appears to be the most complex based on divergent substrate specificities in

bacteria and yeast (42). RluA family members in bacteria are involved in ribosomal assembly

and growth (43, 44) but their function in multicellular organisms has not been studied. Although

pseudouridine synthases appear to function ubiquitously, as noted above, *Drosophila RluA-1*, a

member in RluA family, has been reported to be specifically expressed in md neurons (22). Thus,

we have investigated the role for *RluA-1* and its paralog *RluA-2*, in nociception pathways that

are known to depend on md neurons. Our results indicate an important function for *RluA-1* and

RluA-2 in the regulation of nociception.

Results

RluA-1 is expressed in the multidendritic neurons of the peripheral nervous system

and in the cells of brain

Previously described reporter genes for *RluA-1* showed specific expression in larval

multidendritic neurons (22). We replicated this finding by generating an *RluA-1*^{GAL4} driver at the

endogenous gene locus through recombination mediated cassette exchange (RMCE) of *RluA-*

1^{MI06897} (an intronic MiMIC) and a Trojan exon cassette (45). An mCD8GFP reporter driven by

RluA-1^{GAL4} was expressed in peripheral sensory neurons in each segment of the larval body wall

(**Fig 1A**). In the dorsal cluster, GFP-positive signals were clearly detected in all four classes of

md-da sensory neurons, dorsal multiple dendrite neuron (dmd1), external sensory (ES) and

dorsal bipolar dendritic (dbd) neurons (**Fig 1B**). In the larval ventral nerve cord, the GFP-positive

signals were seen in the axonal projections of the sensory neurons (**Fig 1C**). GFP signal was also

observed in the unidentified clusters of neurons in the larval brain (**Fig 1C**). In the adult brain,

significant signals were detected in the optic lobes and other small cell clusters of the central

brain (**S1A Fig**).

To determine the localization of the RluA-1 proteins, we generated a GFSTF exon trap

(46) that expresses an in frame GFP fusion with *RluA-1* at the endogenous genomic locus (with

RMCE of the *RluA-1*^{MI06897} MiMIC element) (46). Although live imaging did not detect the EGFP

tagged RluA-1 protein, immunostaining with anti-GFP labelled nuclei of larval multidendritic

neurons, ES and dbd neurons also expressing a membrane-localized RFP (*Gal4109(2)80 > UAS-*

CD8-RFP, **Fig 1D**). The tagged RluA-1 protein was also detected in the cell bodies of neurons in

optic lobes and other yet-to-be identified cells in the adult brain (**S1B Fig**).

Reducing or removing *RluA-1* results in a hypersensitive thermal nociception

phenotype

Given our confirmation of the expression of the *RluA1* gene and protein in the

multidendritic neurons we tested for potential roles of RluA-1 in the regulation of nociception.

To do so, we first performed tissue-specific knockdown using GAL4/UAS based RNA

interference (RNAi). A cIVda specific driver (*ppk1.9-GAL4 UAS-dicer2*) (47) was employed to

drive an *RluA-1* RNAi construct in the larval cIVda nociceptors. We assessed potential

insensitive phenotypes (with a probe temperature of 46°C) and potential hypersensitive

phenotypes (with a temperature of 42°C) as previously described (20). When stimulated with

the higher temperature 46°C probe the *RluA-1-RNAi* larvae responded significantly faster than

the *ppk-GAL4* driver alone controls. This genotype also responded faster than the UAS-RNAi

controls but this difference was not statistically significant (**Fig 2A**). The results suggested that

reducing *RluA-1* in classIV neurons may have made the larvae more sensitive to noxious heat.

Indeed, a clearly hypersensitive phenotype for the *RluA1*-RNAi animals was seen when testing

with 42°C probe (**Fig 2A**). The average latency to roll in the *RluA-1* knock-down animals was

significantly faster than the driver alone animals or the *UAS-RNAi* control animals. These data

combined suggest that reducing the activity of *RluA-1* in the noxious heat-responsive cIVda

cells caused thermal hyperalgesia.

To further test the function of *RluA-1* we next generated a precise genetic deletion

mutant of *RluA-1* in which 11.14 kbp including the entire *RluA-1* genomic region was removed

by CRISPR-guided homologous recombination-directed repair (HDR) (48). Homology arms of

~1kb immediately flanking the CRIPSPR cleavage sites were used to direct the HDR (**S2A Fig**).

The resultant deletion mutant ($RluA-1^{del-HDR}$) was confirmed by PCR amplification and

sequencing of PCR products from the targeted $RluA-1$ locus (**S2B Fig**). To facilitate the

behavioral comparisons, the $RluA-1^{del-HDR}$ mutant was backcrossed six times to our most

commonly used genetic background strains Canton-S (CS), w^{1118}, and isogenized w^{1118} (isow^{1118}).

As with class IV md neuron-specific RNAi knockdown, $RluA-1^{del-HDR}$ larvae showed significantly

faster responses to noxious heat stimulation of 42°C compared to the corresponding control

animals, regardless of genetic background (**Fig 2B**). Note that for unknown reasons, these

genetic backgrounds vary in their baseline responses. This genetic background effect has a

significant impact on the latency of larval nociception responses to noxious heat stimuli, with

the most striking differences between homozygous $RluA-1^{-/-}$ and the relatively insensitive

isow^{1118} background, followed by that of w^{1118} background, and then the CS background (**Fig**

2B).

Given the sensitivity to genetic background, we performed an additional genetic test

for the importance of $RluA-1$, by generating an independent mutant allele

($RluA-1^{del-FRT}$) using FLP recombinase and FRT-bearing insertions (49) ($PBac\{WH\}^{f02750}$(+) and

$p\{XP\}^{d2586}$(-), **S2A Fig**). Transheterozygous $RluA-1^{del-FRT}/RluA-1^{del-HDR}$ mutant larvae displayed

hypersensitivity to a 42°C stimulus (**Fig 2C**)) indicating that $RluA-1^{del-FRT}$ failed to complement

$RluA-1^{del-HDR}$. Failure of complementation of independently generated alleles created in distinct

genetic backgrounds provides strong evidence that the hypersensitive nociceptive phenotypes

observed are a consequence of the mutation of $RluA-1$ and not due to unlinked mutations

present in one background or another.

For the remainder of our behavioral studies, we relied on the isogenized w^{1118}

background. This had the advantage of greater genetic uniformity relative to w^{1118} and CS, as

well as showing the strongest hypersensitive mutant phenotype for $RluA-1$.

Genetic rescue of *RluA-1* mutant restores thermal nociception response

To test that the mutation in *RluA-1* was the underlying cause of the hypersensitive nociception phenotype, we introduced a genomic rescue construct into the *RluA-1*$^{del\text{-}HDR}$ mutant background (a Bacterial Artificial Chromosome (BAC) (P6-D7)) (50) covering the *RluA-1* gene region). With the 42°C stimulus, larvae with two copies of the duplication (Dp) in the background of *RluA-1*$^{del\text{-}HDR}$ showed rescue of the hypersensitivity (**Fig 3A**). This rescue with the genomic duplication was dosage dependent. Larvae with only one copy of the Dp in the *RluA-1*$^{del\text{-}HDR}$ background (*RluA-1*$^{-/-}$; *Dp*$^{+/-}$) responded more slowly than the mutant this difference was not statistically significant (**Fig 3A**). Combined, the results of genomic rescue experiments support the hypothesis that mutation of *RluA-1* is indeed the cause of the hypersensitive thermal nociception phenotype.

A caveat remained in that the genomic rescue construct included other genes in addition to *RluA-1*. Thus, genomic rescue did not rule out the possibility that a mutation tightly linked to *RluA-1*, but not in *RluA-1* itself, was responsible for the mutant phenotype. Thus, as a final test, we generated transgenic lines to express an *RluA-1* cDNA under the control of the GAL4/UAS system (*UAS-RluA-1*). Using the *UAS-RluA-1* we specifically restored *RluA-1* to cIVda neurons in the *RluA-1* null mutant background. When stimulated with the 42°C probe the animals with both the *ppkGal4* driver and the *UAS-RluA-1-cDNA* transgene in the *RluA-1*$^{del\text{-}HDR}$ showed a complete rescue from the hypersensitivity seen in the null mutant (**Fig 3B**). Neither the *ppkGal4* driver alone nor the *UAS-RluA-1* had an effect on the hypersensitive thermal nociception phenotype in the *RluA-1* null mutant background, excluding the possibility of non-specific effects of these transgenes (**Fig 3B**). In addition, overexpression of *UAS-RluA1* with a md neuron driver (MD-Gal4) had no effect on nociception behavior at 42°C stimulus ruling out the possibility that the increased latency seen in the rescue effect was a non-specific

consequence of over-expression (**S3 Fig**). Combined, these nociceptor-specific rescue

experiments provide genetic confirmation that loss-of-function mutation in *RluA-1* causes

hypersensitive thermal nociception and localizes the site of action for RluA-1 in this process to

the nociceptors.

RluA-1 requirement for mechanosensory thresholds

The cIVda neurons are not only required for detection of noxious heat, they also

contribute to sensing harsh mechanical stimulation (3, 16, 17). Thus, we investigated the *RluA-1*

mutant responses to noxious mechanical stimuli. When stimulated with a 30mN/720kPa Von

Frey fiber, significantly more *RluA-1* $^{del\text{-}HDR}$ null mutant larvae performed the typical nociceptive

rolling behavior compared to the isow^{1118} controls (**Fig 4A**). Even when the probe

strength/pressure was reduced to 15mN/360kPa, the majority of *RluA-1*$^{-/-}$ larvae still rolled while

less than half of control larvae rolled (**Fig 4A**), indicating the defect in *RluA-1* also caused

hypersensitive mechanical nociception. Loss of *RluA-1* did not have any impact on behavioral

responses to gentle touch (**Fig 4B**) suggesting a more specific involvement in nociception than

for sensory processing in general.

Expression pattern and nociception functions for RluA-2

The *RluA-2* locus is adjacent to *RluA-1* on the second chromosome of *Drosophila*

melanogaster. *RluA-2* has significant sequence similarity with *RluA-1* within the evolutionarily

conserved pseudouridine synthase domain. This amino acid similarity of *RluA1* and *RluA2*

suggested possible functional overlap for the encoded proteins. To investigate the expression

of RluA-2 we generated a GFSTF line with RMCE of the *RluA-2*^{MI12981} mimic element (**S4A Fig**)

(46). Immunostaining with anti-GFP labelled nuclei in all of the cell types that we observed in

third instar larvae (**Fig 5A**), including md neurons (**Fig 5B**).

To test the function of *RluA-2* we generated a deletion (*RluA-2*$^{del-HDR}$) to remove its

pseudouridine synthase domain via CRISPR/Cas9 HDR (**S4A Fig**). Also, since RluA-1 and RluA-2

are both expressed in the md neurons (**Fig 1D** and **Fig 5B**), we generated a double mutant

(*RluA-1*$^{del-HDR}$*RluA-2*$^{del-HDR}$), by injecting the *RluA-2*$^{del-HDR}$ constructs in the *RluA-1*$^{del-HDR}$ null

mutant background (**S4 Fig**). The double mutant allowed us to completely remove RluA

pseudouridine synthase gene activity from the flies. Prior to functional assessment, the single

mutant and the double mutant were backcrossed six times to the genetic background of

iso*w*^{1118}.

We tested each single mutant (*RluA-1*$^{del-HDR}$ and *RluA-2*$^{del-HDR}$) together with the double

mutant *RluA-1*$^{del-HDR}$ *RluA-2*$^{del-HDR}$ side by side in thermal nociception assays with the 42°C

thermal stimulus. *RluA-1*$^{del-HDR}$ larvae (*RluA-1*$^{-/-}$) again responded significantly faster than the

genetic background control (**Fig 6**). The *RluA-2*$^{del-HDR}$ single mutant larvae also displayed a faster

response to the stimulus (**Fig 6**) and similar hypersensitivity was also seen in an independent

allele for *RluA-2* that we generated by FLP/FRT mediated recombination (*RluA-2*$^{del-FRT}$) allele (**S5**

Fig). Finally, the double mutant *RluA-1*$^{del-HDR}$ *RluA-2*$^{del-HDR}$ larvae showed a faster response than

the control larvae (**Fig 6**) to the same extent as the single mutant of *RluA-2*$^{del-HDR}$. These results

indicated that *RluA-2*, like *RluA-1*, negatively regulates nociception. The finding that the double

mutant did not show a more severe phenotype than either single mutant suggests that RluA-2

and RluA-1 have non-redundant functional roles, and that they may function in the same

molecular pathway. When this pathway is disrupted, hypersensitive nociception results.

RluA-1 regulates neuronal dendrite morphology of nociceptors

A nociceptor-specific RNAi screen with thermal nociception assay discovered dozens of

genes whose reduction caused either insensitive or hypersensitive thermal nociception (20).

Interestingly, some of those genes targeted with RNAi showed a reduced or increased

branching of Class IV neuron dendrites. Reduced dendrite branching was often seen with

nociceptive insensitivity while increased branching was found in some hypersensitive

genotypes. Thus, regulation of Class IV neuron dendrite morphology is a commonly affected

developmental pathway that is related to nociception phenotypes. Given this, we investigated

the dendrite morphology of the cIVda neuron dendrites in the $RluA-1^{del-HDR}$ mutant. In mutant

ddaC neurons visualized with $ppk-CD4-tdTom$ato, we observed a modest but significant

increase in the number of dendrite branches (normalized by neuron size) and shorter average

branch length in comparison to control animals (**Figs 7A and 7B**). We also found that dendritic

branches in the $RluA-1^{del-HDR}$ ddaC neurons had higher frequency of isoneuronal cross-over

events compared to the control (**Figs 7C and 7D**). This latter phenotype is suggestive of an

isoneuronal tiling defect. Increased isoneuronal crossovers are also seen in mutants that affect

dendrite attachment to the basal lamina (51-54). Whether or not these dendrite abnormalities

play a causal role in the hypersensitive nociception phenotypes of the $RluA-1$ mutant will be an

interesting subject for future investigation.

Discussion

Given the well-established nociceptive role of md-neurons, we have investigated the

historically first known molecular marker for md-neurons in nociception pathways. This gene

encodes the RluA-1 protein in the RluA family of pseudouridine synthases. Our studies clearly

demonstrate that loss of function for either $RluA-1$ or $RluA-2$ produce hyperalgesia in third

instar $Drosophila$ larvae. Tissue-specific RNAi, genetic null mutant, and cDNA rescue

experiments all indicate that loss of the $RluA-1$ gene from whole animals, or specifically from

nociceptors, results in hyperalgesia. A newly generated $RluA-1^{GAL4}$ driver showed specific

expression in larval multidendritic and ES neurons. As well, a GFP exon trap for RluA-1 protein

localized to the nuclei of these neurons. A small number of unidentified neurons in the larval

brain were also revealed by *RluA-1*^{GAL4} and we observed expression of *RluA-1*^{GAL4} driven

mCD8GFP and GFP tagged RluA-1 in cells of the adult brain, which included the optic lobe.

Although loss of *RluA-2* also caused hyperalgesia, its expression pattern was ubiquitous

and included multidendritic neurons. Like the RluA-1 GFP exon trap, the RluA-2 GFP exon trap

labelled nuclei. The nuclear localization for both RluA-1 and RluA-2 may indicate that these

proteins act on RNA targets prior to export of the nucleus, or that they predominantly act upon

nuclear localized RNAs. A caveat to this interpretation is that we have yet to demonstrate that

these molecules function as genuine pseudouridine synthases, and this function remains a

hypothesis that is based on the known function of evolutionary homologues. This hypothetical

function requires future biochemical verification.

We also observed that *RluA-1* mutants showed an increase in the number of dendrite

branches relative to control genotypes as well as an increase in isoneuronal crossovers.

Transcription factors, cytoskeletal regulators, motor proteins, secretory pathways and cell

adhesion molecules all function in concert to develop and maintain optimum dendrite

morphology (55). RluA-1 may modify RNAs for those genes that regulate dendritic morphology,

or changes in the dendrite morphology could be an indirect consequence of neuronal sensitivity

that is regulated by RluA-1. It is noteworthy that prior studies have noted a potential link

between the degree of dendrite branching and the sensitivity of nociception behaviors (20). In

other cases, axonal factors such as the ion channel SK have been found to be important in

regulating the cIVda neuron excitability (56, 57). Whether the dendrite branching phenotype

that we observe in *RluA-1* mutants is a cause or a consequence of hypersensitivity will be an

interesting question for future studies.

A large body of literature indicates that RNA trafficking and local translation is

important in dendrites and axons of neurons (58, 59). Relative to uridine the pseudourine base

is believed to have enhanced rotational freedom which may alter conformation of RNA

secondary structures. As well, an additional hydrogen bond donor present in pseudouridine

may favor alternative base-pairing interactions in RNA. These properties may consequently

alter RNA localization, stability and/or efficiency of translation (60-62). Pseudouridines can also

influence decoding during translation as pseudouridylation of nonsense codons has been

shown to suppress translation termination both *in vitro* and *in vivo* (63). Thus, another possible

function for pseudouridylation in nociceptive neurons could be to favor read-through of

pseudouridylated stop codons to generate novel sequences at protein carboxy termini.

The precise mechanism explaining the involvement of RluA proteins in nociception can

only be elucidated by identifying the RNA targets of these enzymes. Several groups have

developed methods using next generation sequencing methods to identify the pseudouridine

sites in transcriptomes (30-33, 64). Future investigations applying these methods to wild type

and *RluA* mutants in *Drosophila* will help us to identify the RluA targets and to further define the

underlying mechanisms.

Materials and methods

Fly strains and husbandry

The following fly strains were obtained from Bloomington Stock Center: (w^{1118};

PBac{vas-Cas9}VK00027), Mi{MIC}RluA-1$^{[MI06897]}$, Mi{MIC}RluA-2$^{[MI12981]}$, Df(2L)Exel7048/CyO, (yw;

Sp/CyO; pC-(lox-attB2-SA-T2A-Gal4-Hsp70)3), (yw hs-cre, vasΦC31; Sp/CyO; Sb/TM3Ser),

(P{ry=hsFLP}, yw M{vas-int.B}ZH-2A; Sp/CyO; P{FRT-attB-{GFSTF}-attB(w+)-FRT}), ppk-CD4-

tdTom, md-Gal4, UAS-mCD8::RFP, 40XUAS-mCD8::GFP. The following fly strains were obtained

from Exelixis collection at Harvard Medical School: PBac{WH}^{f02750}, P{XP}d2586,

PBac{WH}Grip75$^{[f05483]}$, PBac{WH}RluA-2$^{[f07702]}$. The RNAi lines targeting *RluA-1* (31719-R1, 31719-

R2) were obtained from Kyoto Stock Center. The genetic duplication line (BAC ID: CH321-

49P21) covering *RluA-1* (starting at 24,819,420 and ending at 24,910,132 on 2R) was obtained

from Genetivision. The nos-Cas9 line used for generating *RluA-2*$^{del-HDR}$ and double mutant *RluA-*

1$^{del-HDR}$*RluA-2*$^{del-HDR}$ [*y sc v; {nos-Cas9} attP2* (TH00787.N)] was kindly provided by the Norbert

Perrimon lab. All larvae used in experiments were reared on the Bloomington *Drosophila*

medium in an incubator with controlled temperature (25°C) and humidity (70%) on a 12h

light/12h dark cycle. Strains were otherwise maintained at room temperature.

CRIPSR targeting of *RluA-1* and *RluA-2*

HDR for mutagenesis used the CRISPR/Cas9 (Gratz et al. 2013) to generate *RluA-1*$^{del-HDR}$,

RluA-2$^{del-HDR}$ and the double mutant *RluA-1*$^{del-HDR}$*RluA-2*$^{del-HDR}$. Target sites were selected using

the flyCRISPR Optimal Target Finder. For the precise deletion of *RluA-1*, the gRNA target sites

are 5′ end 5′-CCACTG|TGCAGCGGAAAATTCAC-3′ (where "|" represents the Cas9 cut site,

which is 71bp upstream of the *RluA-1* transcriptional start) and 3′ end: 5′-

ACATATATTCAAAAGCT|CTTTGG-3′ (cut site at 60bp downstream of *RluA-1* 3′UTR). We used

the following primer pairs to amplify the 885bp *RluA-1* homology ARM1: *RluA-1*-ARM1F: 5′-

AATACACCTGCATTATCGCTGGTCCCTGTGGCTTTGCAC-3′,

RluA-1-ARM1R: 5′-AATACACCTGCAATTCTACCAGTGGGGCAAACCGCATTT-3′. We used *RluA-*

1-ARM2F: 5′-GACTGCTCTTCGTATCTTTGGATGGTAAGTGCTTAAAC-3′ and *RluA-1*-ARM2R: 5′-

ATTAGCTCTTCTGACCTTACAACTCCTTCAAGTCA-3′ to amplify 969 bp of *RluA-1* homology

ARM2. For *RluA-2*, we used a target site at the 5′ end: 5′-GCAATCTATAGGTCTGC|GGAAGC-3′

(cut site at bp 5 of exon 3) and at the 3′ end: 5′AAGATAAACTACAGAGA|CCCCGG-3′ (cutting

site in the middle of exon 8) so that the conserved pseudouridine synthase domain (located in

exon 5) would be deleted. Primer pairs *RluA-2*-ARM1F:

5′CTAACACCTGCATATTCGCTCGAAACCCATTGTTAGCTG3′ and *RluA-2* ARM1R: 5′

CATTCACCTGCATTACTACGCAGACCTATAGATTGCAAT3′ were used to amplify the 979bp

RluA-2 homology ARM1. *RluA-2-ARM2F*: 5'CAGTGCTCTTCGTATCCCCGGCACAAAGGATCTCA-

3', and *RluA-2-ARM2R*: 5' GACTGCTCTTCCGACATTTCAATGCCCTTGGCCAA-3' were used to

amplify the RluA-2 ARM2 (998bp). Rapid dsDNA donor cloning was carried out with the *pHD-*

DsRed-attP vector (65) and the guide RNA (gRNAs) were cloned into *pU6-BsbI-chiRNA* vector

(66). Embryos of *vas-Cas9* on chromosome III (w^{1118}; *PBac{y[+mDint2]=vas-Cas9}VK00027,*

"injection line 1") were injected with *RluA-1* dsDNA donor and gRNAs to generate G_0 founders

for *RluA-1$^{del-HDR}$*. Embryos with *nos-Cas9* on chromosome III [*y sc v; {nos-Cas9}attP2* (TH00787.N),

"injection line 2"] were injected with *RluA-2* dsDNA donor and gRNAs to create the G_0 founders

for *RluA-2$^{del-HDR}$*. For generating the double mutant *RluA-1$^{del-HDR}$RluA-2$^{del-HDR}$*, *nos-Cas9* was first

introduced to the *RluA-1$^{del-HDR}$* mutant background and the DsRed marker in *RluA-1$^{del-HDR}$*

removed with CRE recombinase and the resultant homozygous strain [*yw; RluA-1$^{del-HDR}$ΔDsRed;*

{nos-Cas9}attP2 (TH00787.N), "injection line 3"] was injected with the *RluA-2* dsDNA donor and

gRNAs. All embryo injections of the dsDNA donor (at a concentration of 500ng/ul) and gRNAs

(100ng/ul for each) were performed by the Model system Injections

(modelsysteminjections@flymsi.com).

To identify the desired HDR mutants, G_0 flies were crossed to w^{1118} and single F_1

founders were identified with DsRed fluorescence in the eyes (from *3XP3-DsRed reporter*) and

mated with a second chromosome balancer strain to establish independent lines. Molecular

testing for the desired events was performed by PCR on gDNA extracted from candidates

placed over a deficiency (*Df(2L)Exelixis7048*) covering the *RluA-1* and *RluA-2* region (primer

pairs marked in **S2A** and **S4A Figs**). PCR products sequenced across the *attp-loxP-3XP3-DsRed-*

SV40-loxP fragment (**S2B** and **S4B Figs, Fig 4C**) confirmed accurate targeting of the loci. For

behavioral analysis, the original deletion mutants were backcrossed to *CS*, w^{1118} or isow^{1118} for

six times. For each generation, five heterozygous females were selected for six successive

backcrosses in vials and about 10 heterozygous females were used to cross to a second
chromosome balancer to produce balanced mutant males in bottles and finally heterozygous
mutant virgins and balanced males were crossed "en masse" in bottles to establish the
balanced and homozygous mutant lines. In all crosses the dsRed fluorescence marker was used
to follow the presence of the mutant.

Generation of deletion line in *RluA-1* and *RluA-2* using FRT-mediated deficiency

FRT bearing insertions *{WH+}^{fo2750}* and *{XP-}^{d2586}* and in *RluA-1* (locations marked in **S2A**
Fig) were used for generating a deficiency allele *RluA-1$^{del-FRT}$*. Insertions of *{WH-}RluA-2^{F07702}* and
{WH-}Grip75^{fo5483} (locations marked in **S4A Fig**) were used for generating the deficiency allele
RluA-2$^{del-FRT}$. Crossing and heat-shock schemes followed (49). Hybrid PCR with corresponding
primers (WH5′ plus / XP5′ minus left and right primers) was used to screen for candidate lines
with *w-* deletion in *RluA-1$^{del-FRT}$* and two-sided PCR with left and right primers for WH3′ minus/
WH5′ minus was used for screening for candidate lines with *w+* deletion in *RluA-2$^{del-FRT}$* (49).
Molecular testing for the deletion was performed by PCR on gDNA extracted from positive
candidates placed over a deficiency (*Df(2L)Exelixis7048*) covering the *RluA-1* and *RluA-2* region.
The primers used for PCR and sequencing verification in *RluA-1$^{del-FRT}$* are *RluA-1-d2586*-up-F: 5′-
AAAAATGCGGTTTGCCCC-3′, located upstream of the *{XP-}d2586* insertion site and *RluA-1-*
fo2750-down-R: 5′-AAGGGGTAACAAAAGTCCAG-3′, downstream of *{WH+}fo2750 insertion site.*

Generation of *RluA-1^{GAL4}* using "Trojan-exon"

A triplet Trojan exon donor line on the 3rd chromosome *(yw; Sp/CyO; pC-(lox-attB2-SA-*
T2A-Gal4-Hsp70)3) and an *RluA-1* insertion line containing an intronic MiMIC element
(*Mi{MIC}RluA-1$^{[MI06897]}$*) were used to generate the *RluA-1^{Gal4}* driver using a crossing scheme as
described by (45). Candidate males who have lost the *y^{+}* selection marker associated with
MiMIC were crossed to *40XUAS-mCD8::GFP* line and animals expressing GFP were selected to

establish a stable line. The line with the correct linker (phase 0) was confirmed with sequencing

of PCR products amplifying the left side with primers *RluA-1*-5494F: 5'-

TGATGTTGCCCCATAACG-3' and *T2A-Gal4*-Seq-1R: 5' CGCTATCGATGCTCACGGTC-3' and the

right side with the primers *T2A-Gal4*-4F: 5'-ACACCGTGCTGATGCTGC-3' and *RluA-1*-5907R: 5'-

GAAAACATCGCACATCTGG-3' of the *RluA-1* genome-*T2A-Gal4* insertion bordering region.

Generation of GFSTF insertions in RluA-1 and RluA-2 by recombination mediated

cassette exchange (RMCE)

Crossing, heat shocking and screening for EGFP tagged MiMIC lines in *RluA-1* and *RluA-2* were

essentially carried out as described (46). Males carrying a MiMIC insertion in a coding intron of

RluA-1 (*RluA-1^{MI06897}*) or *RluA-2* (*RluA-2^{MI12981}*) were crossed to females carrying the *hs-FLP* and

vasa-phiC31 integrase on the X chromosome and a frame-specific ("phase 0") *FRT* flanked

multiple tag (*GFSTF*) cassette on chromosome III. Candidate males with mosaic *w-* eyes and *y-*

bodies were individually crossed to *w; Sco/CyO; Sb/TM3 Ser* balancers to establish stocks. The

presence and direction of the insertion were tested by PCR assays described (67). Since the

original MiMIC insertion in *RluA-1* or *RluA-2* and the respective gene are in the same orientation,

positive PCR reaction 1 (with primers MiLF and TagR) and 4 (with primers MiLR and TagF) as

described (67) indicated a successful RMCE event and resulted in expression.

Generation of *UAS-RluA-1*

Drosophila RluA-1 full length cDNA clone for *RluA-1* transcript A (FI04540) was obtained

from the Drosophila Genome Resource Center (DGRC). The *UAS-RluA-1* expression constructs

were generated with the ENTR/gateway system following the instructions of the manufacturer

(Invitrogen). The *RluA-1*-cDNA-F (5'-CACCATGCAGAATTCTCCGGCT-3', and *RluA-1*-

cDNA+STOP-R (5'-TCATGCCGAGTCTAAGTG-3' primers were used to amplify the open reading

frame and cloned into *pENTR-D-TOPO* vector. The sequence was verified and then cloned to

the P-element based destination vector *pTW*. Model System Injections performed injections

into w^{1118} embryos. F_0 flies were crossed to w^{1118} and single F_1 founders were identified based on

the w^+ marker. Individual lines were mapped and balanced to establish stable stocks. A *UAS-*

RluA-1 line with relatively weak expression evaluated by Q-PCR in *RluA-1*-cDNA lines driven by

md-GAL4 was used for the cIV-neuron specific *RluA-1*-cDNA rescue experiment (**Fig 3B**).

Larval behavioral analyses

Wandering 3^{rd} instar larvae were washed out from vials and acclimated for 5 min in petri

dishes before testing. Larval thermal and mechanical nociception response assays were

conducted as previously described (4, 17, 56). Behavioral recording and scoring were performed

with the observer blinded to the genotype.

For gentle touch assays, early L3 larvae were scooped out from the top layer of the fly

food in the vials and 5-10 larvae were briefly rinsed with PBS and allowed to acclimate on 1%

agarose in a plate for 5 min before testing with an eyelash fixed to the end of a paintbrush. Each

larva was brushed with the eyelash on segments T1-A3 for 4 times and the responses were

recorded and summarized using a gentle touch scale (68).

Immunohistochemistry and microscopy

The following primary antibodies were used for immunofluorescence: rabbit anti-GFP (ab6556,

Abcam, 1:500), mouse anti-GFP (ab38689, Abcam, 1:500), Rabbit anti-HRP (1:100), mouse anti-

nc82 (DSHB, supernatant, 1:30). Alexa Fluor 488, 633 were used at 1:1000 as secondary

antibodies. Detailed immunostaining protocol is available on request. Images for

immunostained tissues were taken on a Zeiss LSM 5 LIVE confocal microscope using a 40X

objective except for **Fig 1D** and **S1B Fig,** which were taken on a Zeiss LSM880 using the 63X

objective.

CIV dendrite imaging, tracing and analysis

To image class IV dendrites, the *DsRed* marker in *RluA-1$^{del-HDR}$* was first removed with CRE

recombinase (*w^{1118}; RluA-1$^{del-HDRΔDsRed}$*) and *ppk1.9-CD4::tdTom* was introduced to the mutant

background and a stable homozygous stock was established. For CIV dendrite analysis, six

virgins and three males were crossed in each vial for the mutant (*w^{1118}; RluA-1$^{del-HDRΔDsRed}$;*

ppk1.9-CD4::tdTom) and control *(ppk1.9-CD4::-td-Tom)*. Wandering larvae were anesthetized

with diethyl ether in a sealed glass chamber for 15min before being arranged on a slide and

covered with 50 mm glass coverslip. Neurons expressing the fluorescently tagged markers were

visualized on a Zeiss LSM 5 Live confocal microscope with a 40X oil objective (Plan Apo M27,

NA 0.8). Images were collected as 5x3 tiles scans of z-stacks with 512x512 resolution. A MatLab

build was used for initial automatic tracing of the ddaC neuron dendrites from the confocal z-

stack series TIFF images (69). The generated SWC files were overlaid onto the maximum

intensity projected image of the neuron in neuTube (70) and manually curated to eliminate

tracing errors made by MatLab. The corrected images were then analyzed with MatLab to

extract neuron features of interest including number of branches, average branch length, and

neuron size (the estimated size of the neuron, defined as the area of the minimum bounding

circle) (69). Iso-neuronal crossover events were quantified manually from the traced dendrites

for each genotype (n=6 neurons).

Statistical analysis

Statistics were performed using GraphPad Prism 4. Thermal and gentle touch

behavioral data were compared with an unpaired non-parametric Mann-Whitney test when

comparing two groups and Kruskal-Wallis test when comparing three or more groups.

Mechanical nociception behavioral data were compared with Fisher's exact test. Dendrite

morphology data were compared with the Student's t test. Error bars represent S.D. in all the

figures unless otherwise specified.

Acknowledgements

The authors thank Drosophila Genomic Resource Center (supported by NIH grant 2P40OD010949) for clones. Stocks obtained from the Bloomington Drosophila Stock Center (NIH P40OD018537) were used in this study. We also thank the KYOTO Stock Center Kyoto in Kyoto Institute of Technology and Harvard Medical School for providing fly stocks used in this study. Members of Tracey lab provided helpful feedback on the manuscript and Jayce Brown Culbertson assisted with dendrite tracing.

References

1. Gaskin DJ, Richard P. The economic costs of pain in the United States. J Pain. 2012;13(8):715-24.

2. Tracey WD, Wilson RI, Laurent G, Benzer S. painless, a Drosophila gene essential for nociception. Cell. 2003;113(2):261-73.

3. Hwang RY, Zhong L, Xu Y, Johnson T, Zhang F, Deisseroth K, et al. Nociceptive neurons protect Drosophila larvae from parasitoid wasps. Curr Biol. 2007;17(24):2105-16.

4. Caldwell JC, Tracey WD. Alternatives to Mammalian Pain Models 2: Using Drosophila to Identify Novel Genes Involved in Nociception. Methods Mol Biol. 2010;617:19-29.

5. Milinkeviciute G, Gentile C, Neely GG. Drosophila as a tool for studying the conserved genetics of pain. Clin Genet. 2012;82(4):359-66.

6. Im SH, Galko MJ. Pokes, sunburn, and hot sauce: Drosophila as an emerging model for the biology of nociception. Dev Dynam. 2012;241(1):16-26.

7. Tracey WD, Jr. Nociception. Curr Biol. 2017;27(4):R129-R33.

8. Khuong TM, Wang QP, Manion J, Oyston LJ, Lau MT, Towler H, et al. Nerve injury drives a heightened state of vigilance and neuropathic sensitization in Drosophila. Sci Adv. 2019;5(7):eaaw4099.

9.   Xiang Y, Yuan Q, Vogt N, Looger LL, Jan LY, Jan YN. Light-avoidance-mediating

photoreceptors tile the Drosophila larval body wall. Nature. 2010;468(7326):921-6.

10.   Grueber WB, Jan LY, Jan YN. Tiling of the Drosophila epidermis by multidendritic

sensory neurons. Development. 2002;129(12):2867-78.

11.   Hu C, Petersen M, Hoyer N, Spitzweck B, Tenedini F, Wang D, et al. Sensory integration

and neuromodulatory feedback facilitate Drosophila mechanonociceptive behavior. Nat

Neurosci. 2017;20(8):1085-95.

12.   Ohyama T, Schneider-Mizell CM, Fetter RD, Aleman JV, Franconville R, Rivera-Alba M,

et al. A multilevel multimodal circuit enhances action selection in Drosophila. Nature.

2015;520(7549):633-9.

13.   Chin MR, Tracey WD, Jr. Nociceptive Circuits: Can't Escape Detection. Curr Biol.

2017;27(16):R796-R8.

14.   Burgos A, Honjo K, Ohyama T, Qian CS, Shin GJ, Gohl DM, et al. Nociceptive

interneurons control modular motor pathways to promote escape behavior in Drosophila. Elife.

2018;7.

15.   Ohyama T, Jovanic T, Denisov G, Dang TC, Hoffmann D, Kerr RA, et al. High-

throughput analysis of stimulus-evoked behaviors in Drosophila larva reveals multiple

modality-specific escape strategies. Plos One. 2013;8(8):e71706.

16.   Zhong L, Hwang RY, Tracey WD. Pickpocket is a DEG/ENaC protein required for

mechanical nociception in Drosophila larvae. Curr Biol. 2010;20(5):429-34.

17.   Mauthner SE, Hwang RY, Lewis AH, Xiao Q, Tsubouchi A, Wang Y, et al. Balboa Binds

to Pickpocket In Vivo and Is Required for Mechanical Nociception in Drosophila Larvae. Curr

Biol. 2014;24(24).

18.   Gorczyca DA, Younger S, Meltzer S, Kim SE, Cheng L, Song W, et al. Identification of

Ppk26, a DEG/ENaC Channel Functioning with Ppk1 in a Mutually Dependent Manner to Guide

Locomotion Behavior in Drosophila. Cell Rep. 2014;9(4):1446-58.

19.   Zhong L, Bellemer A, Yan H, Ken H, Jessica R, Hwang RY, et al. Thermosensory and

nonthermosensory isoforms of Drosophila melanogaster TRPA1 reveal heat-sensor domains of

a thermoTRP Channel. Cell Rep. 2012;1(1):43-55.

20.   Honjo K, Mauthner SE, Wang Y, Skene JH, Tracey WD, Jr. Nociceptor-Enriched Genes

Required for Normal Thermal Nociception. Cell Rep. 2016;16(2):295-303.

21.   Brewster R, Bodmer R. Origin and specification of type II sensory neurons in Drosophila.

Development. 1995;121(9):2923-36.

22.   Wang CC, Lo JC, Chien CT, Huang ML. Spatially controlled expression of the Drosophila

pseudouridine synthase RluA-1. Int J Dev Biol. 2011;55(2):223-7.

23.   Olesnicky EC, Killian DJ, Garcia E, Morton MC, Rathjen AR, Sola IE, et al. Extensive use

of RNA-binding proteins in Drosophila sensory neuron dendrite morphogenesis. G3 (Bethesda).

2014;4(2):297-306.

24.   Ye B, Petritsch C, Clark IE, Gavis ER, Jan LY, Jan YN. Nanos and Pumilio are essential for

dendrite morphogenesis in Drosophila peripheral neurons. Curr Biol. 2004;14(4):314-21.

25.   Pan L, Zhang YQ, Woodruff E, Broadie K. The Drosophila fragile X gene negatively

regulates neuronal elaboration and synaptic differentiation. Curr Biol. 2004;14(20):1863-70.

26.   Brechbiel JL, Gavis ER. Spatial regulation of nanos is required for its function in dendrite

morphogenesis. Curr Biol. 2008;18(10):745-50.

27.   Bianco A, Dienstbier M, Salter HK, Gatto G, Bullock SL. Bicaudal-D regulates fragile X

mental retardation protein levels, motility, and function during neuronal morphogenesis. Curr

Biol. 2010;20(16):1487-92.

28.   Xu X, Brechbiel JL, Gavis ER. Dynein-dependent transport of nanos RNA in Drosophila

sensory neurons requires Rumpelstiltskin and the germ plasm organizer Oskar. J Neurosci.

2013;33(37):14791-800.

29.   Ge J, Yu YT. RNA pseudouridylation: new insights into an old modification. Trends

Biochem Sci. 2013;38(4):210-8.

30.   Carlile TM, Rojas-Duran MF, Zinshteyn B, Shin H, Bartoli KM, Gilbert WV.

Pseudouridine profiling reveals regulated mRNA pseudouridylation in yeast and human cells.

Nature. 2014;515(7525):143-+.

31.   Lovejoy AF, Riordan DP, Brown PO. Transcriptome-Wide Mapping of Pseudouridines:

Pseudouridine Synthases Modify Specific mRNAs in S. cerevisiae. Plos One. 2014;9(10).

32.   Schwartz S, Bernstein DA, Mumbach MR, Jovanovic M, Herbst RH, Leon-Ricardo BX, et

al. Transcriptome-wide mapping reveals widespread dynamic-regulated pseudouridylation of

ncRNA and mRNA. Cell. 2014;159(1):148-62.

33.   Khoddami V, Yerra A, Mosbruger TL, Fleming AM, Burrows CJ, Cairns BR.

Transcriptome-wide profiling of multiple RNA modifications simultaneously at single-base

resolution. Proc Natl Acad Sci U S A. 2019;116(14):6784-9.

34.   Knight SW, Heiss NS, Vulliamy TJ, Greschner S, Stavrides G, Pai GS, et al. X-linked

dyskeratosis congenita is predominantly caused by missense mutations in the DKC1 gene. Am J

Hum Genet. 1999;65(1):50-8.

35.   Fujiwara T, Harigae H. Pathophysiology and genetic mutations in congenital

sideroblastic anemia. Pediatr Int. 2013;55(6):675-9.

36.   de Brouwer APM, Abou Jamra R, Kortel N, Soyris C, Polla DL, Safra M, et al. Variants in

PUS7 Cause Intellectual Disability with Speech Delay, Microcephaly, Short Stature, and

Aggressive Behavior. Am J Hum Genet. 2018;103(6):1045-52.

37.   Hammal T, Ferre-D'Amare AR. Pseudouridine synthases. Chem Biol. 2006;13(11):1125-

35.

38.   Koonin EV. Pseudouridine synthases: four families of enzymes containing a putative

uridine-binding motif also conserved in dUTPases and dCTP deaminases. Nucleic Acids Res.

1996;24(12):2411-5.

39.   Phillips B, Billin AN, Cadwell C, Buchholz R, Erickson C, Merriam JR, et al. The Nop60B

gene of Drosophila encodes an essential nucleolar protein that functions in yeast. Mol Gen

Genet. 1998;260(1):20-9.

40.   Giordano E, Peluso I, Senger S, Furia M. minifly, a Drosophila gene required for

ribosome biogenesis. J Cell Biol. 1999;144(6):1123-33.

41.   Vicidomini R, Petrizzo A, di Giovanni A, Cassese L, Lombardi AA, Pragliola C, et al.

Drosophila dyskerin is required for somatic stem cell homeostasis. Sci Rep. 2017;7(1):347.

42.   Hoang C, Chen JJ, Vizthum CA, Kandel JM, Hamilton CS, Mueller EG, et al. Crystal

structure of pseudouridine synthase RluA: Indirect sequence readout through protein-induced

RNA structure. Mol Cell. 2006;24(4):535-45.

43.   Gutgsell NS, Deutscher MP, Ofengand J. The pseudouridine synthase RluD is required

for normal ribosome assembly and function in Escherichia coli. Rna. 2005;11(7):1141-52.

44.   Raychaudhuri S, Niu L, Conrad J, Lane BG, Ofengand J. Functional effect of deletion

and mutation of the Escherichia coli ribosomal RNA and tRNA pseudouridine synthase RluA. J

Biol Chem. 1999;274(27):18880-6.

45.   Diao F, Ironfield H, Luan H, Diao F, Shropshire WC, Ewer J, et al. Plug-and-play genetic

access to drosophila cell types using exchangeable exon cassettes. Cell Rep. 2015;10(8):1410-21.

46.   Nagarkar-Jaiswal S, DeLuca SZ, Lee P-T, Lin W-W, Pan H, Zuo Z, et al. A genetic toolkit

for tagging intronic MiMIC containing genes. Elife. 2015;4.

47.   Ainsley JA, Pettus JM, Bosenko D, Gerstein CE, Zinkevich N, Anderson MG, et al.

Enhanced locomotion caused by loss of the Drosophila DEG/ENaC protein Pickpocket1. Curr

Biol. 2003;13(17):1557-63.

48.   Ran FA, Hsu PD, Wright J, Agarwala V, Scott DA, Zhang F. Genome engineering using

the CRISPR-Cas9 system. Nat Protoc. 2013;8(11):2281-308.

49.   Parks AL, Cook KR, Belvin M, Dompe NA, Fawcett R, Huppert K, et al. Systematic

generation of high-resolution deletion coverage of the Drosophila melanogaster genome. Nat

Genet. 2004;36(3):288-92.

50.   Venken KJ, Carlson JW, Schulze KL, Pan H, He Y, Spokony R, et al. Versatile P[acman]

BAC libraries for transgenesis studies in Drosophila melanogaster. Nat Methods. 2009;6(6):431-

4.

51.   Han C, Wang D, Soba P, Zhu S, Lin X, Jan LY, et al. Integrins regulate repulsion-

mediated dendritic patterning of drosophila sensory neurons by restricting dendrites in a 2D

space. Neuron. 2012;73(1):64-78.

52.   Kim SE, Coste B, Chadha A, Cook B, Patapoutian A. The role of Drosophila Piezo in

mechanical nociception. Nature. 2012;483(7388):209-12.

53.   Tenenbaum CM, Misra M, Alizzi RA, Gavis ER. Enclosure of Dendrites by Epidermal

Cells Restricts Branching and Permits Coordinated Development of Spatially Overlapping

Sensory Neurons. Cell Rep. 2017;20(13):3043-56.

54.   Meltzer S, Yadav S, Lee J, Soba P, Younger SH, Jin P, et al. Epidermis-Derived

Semaphorin Promotes Dendrite Self-Avoidance by Regulating Dendrite-Substrate Adhesion in

Drosophila Sensory Neurons. Neuron. 2016;89(4):741-55.

55.   Corty MM, Matthews BJ, Grueber WB. Molecules and mechanisms of dendrite

development in Drosophila. Development. 2009;136(7):1049-61.

56.   Walcott KCE, Mauthner SE, Tsubouchi A, Robertson J, Tracey WD. The Drosophila

Small Conductance Calcium-Activated Potassium Channel Negatively Regulates Nociception.

Cell Rep. 2018;24(12):3125-32 e3.

57.   Onodera K, Baba S, Murakami A, Uemura T, Usui T. Small conductance Ca(2+)-

activated K(+) channels induce the firing pause periods during the activation of Drosophila

nociceptive neurons. Elife. 2017;6.

58.   Rangaraju V, Tom Dieck S, Schuman EM. Local translation in neuronal compartments:

how local is local? EMBO Rep. 2017;18(5):693-711.

59.   Glock C, Heumuller M, Schuman EM. mRNA transport & local translation in neurons.

Curr Opin Neurobiol. 2017;45:169-77.

60.   Arnez JG, Steitz TA. Crystal structure of unmodified tRNA(Gln) complexed with

glutaminyl-tRNA synthetase and ATP suggests a possible role for pseudo-uridines in

stabilization of RNA structure. Biochemistry-Us. 1994;33(24):7560-7.

61.   Newby MI, Greenbaum NL. Sculpting of the spliceosomal branch site recognition motif

by a conserved pseudouridine. Nat Struct Biol. 2002;9(12):958-65.

62.   Kariko K, Muramatsu H, Welsh FA, Ludwig J, Kato H, Akira S, et al. Incorporation of

pseudouridine into mRNA yields superior nonimmunogenic vector with increased translational

capacity and biological stability. Mol Ther. 2008;16(11):1833-40.

63.   Karijolich J, Yu YT. Converting nonsense codons into sense codons by targeted

pseudouridylation. Nature. 2011;474(7351):395-+.

64.   Li XY, Zhu P, Ma SQ, Song JH, Bai JY, Sun FF, et al. Chemical pulldown reveals dynamic

pseudouridylation of the mammalian transcriptome. Nat Chem Biol. 2015;11(8):592-U93.

65.   Beumer KJ, Carroll D. Targeted genome engineering techniques in Drosophila.

Methods. 2014;68(1):29-37.

66.   Gratz SJ, Cummings AM, Nguyen JN, Hamm DC, Donohue LK, Harrison MM, et al.

Genome engineering of Drosophila with the CRISPR RNA-guided Cas9 nuclease. Genetics.

2013;194(4):1029-35.

67.   Venken KJT, Schulze KL, Haelterman NA, Pan H, He Y, Evans-Holm M, et al. MiMIC: a

highly versatile transposon insertion resource for engineering Drosophila melanogaster genes.

Nat Methods. 2011;8(9):737-43.

68.   Tsubouchi A, Caldwell JC, Tracey WD. Dendritic filopodia, Ripped Pocket, NOMPC, and

NMDARs contribute to the sense of touch in Drosophila larvae. Curr Biol. 2012;22(22):2124-34.

69.   Gulyanon S, Sharifai, N., Kim, M. D., Chiba, A., Tsechpenakis, G. CRF formulation of

active contour population for efficient three-dimensional neurite tracing. 2016 IEEE 13th

International Symposium on Biomedical Imaging2016. p. 593-7.

70.   Feng L, Zhao T, Kim J. neuTube 1.0: A New Design for Efficient Neuron Reconstruction

Software Based on the SWC Format. eNeuro. 2015;2(1).

Figure Legends

Fig 1. *RluA-1* **gene and protein expression in** *Drosophila melanogaster.* (**A**) A low

magnification confocal micrograph of *RluA-1* gene expression pattern in the larval PNS (third

instar w^{1118}; *RluA-1^{Gal4}/40xUAS-mCD8::GFP*). Note that GFP positive signals are detected in

peripheral sensory neurons in each segment along the larval body wall. Scale bar=500μm. (**B**)

Higher magnification of an abdominal dorsal PNS cluster. *RluA-1^{Gal4}* is expressed in the cell body

and dendrites of all classes of multidendritic neurons, external sensory (ES), dorsal multiple

dendrite neuron (dmd1) and dorsal bipolar dendritic (dbd) neurons (ddaD and ddaE (Class I),

ddaB (Class II), ddaA and ddaF (Class III), and ddaC (ClassIV), ES, dmd1 and dbd neurons are

labeled). Scale bar = 50μm. (**C**). *RluA-1^{Gal4}/+* driving expression of *40xUAS-mCD8::GFP/+* in the

larval CNS. Labelling is observed in axonal projections of sensory neurons in the larval ventral

nerve cord and unidentified clusters of neurons in the larval brain. Scale bar =5 0μm. (**D[1-3]**) anti-

GFP immunohistochemistry of a third instar larval fillet preparation of *RluA-1-GFSTF/*

Gal4109(2)80>UAS-mCD8-RFP, immunoreactive signals are detected in the nuclei of multi-

dendritic neurons (D[1] green, GFP) surrounded by the membrane-localized RFP signal (D[2]

magenta, RFP) and merged image in D[3]. Scale bar = 50μm.

Fig 2. Thermal nociception in animals with *RluA-1* loss of function. (A). Class IV specific

knock-down in *RluA-1* results in significant hypersensitive thermal nociception in larvae

compared to the *ppk-GAL4* driver alone animals at 46▯ (average latency of 1.33 ± 1.10 sec for

ppk-Gal4 > UAS-RluA-1-RNAi (31719-R1), n=38 vs. 2.04 ± 1.89 sec for *ppk-GAL4* driver alone,

n=37. P<0.05), albeit not statistically significant compared to the UAS-RNAi alone (1.96 ± 1.32

sec for UAS-RNAi alone, n=27). These differences are more pronounced at 42▯ (average latency

of 1.92 ± 1.16 sec for *ppk-Gal4 > UAS-RluA-1-RNAi*, n=35 vs. 6.25 ± 2.51 sec for *ppk-GAL4* driver

alone controls, n=42 or 5.20 ± 3.28 sec for UAS-RNAi alone controls, n=49, P<0.0001). Kruskal-

Wallis test with Dunn's multiple comparison tests, only significant comparisons were labeled.

(B). Homozygous null mutant *RluA-1[del-HDR]* larvae showed significantly faster response to

noxious heat stimulation of 42°C compared to the corresponding control animals of *Canton-S*

(average latency of 1.29 ± 0.78 sec in *RluA-1[-/-]*, n=54 vs. 2.22 ± 1.32 sec in *CS*, n=39), w^{1118} (2.23 ±

1.57 sec in *RluA-1[-/-]*, n=51 vs. 4.16 ± 2.31 sec in w^{1118}, n=36) and isow^{1118} (4.82 ± 2.53 sec in *RluA-1[-/-]*,

n=30 vs. 11.99 ± 3.98 sec in isow^{1118}, n=17). Significance of comparisons are marked as ****

(p<0.0001). Data were analyzed using Mann-Whitney non-parametric test. **(C).**

Transheterozygote *RluA-1[del-HDR]/RluA-1[del-FRT]* showed hypersensitive thermal nociception

responses compared to the controls (average latency of 3.07 ± 1.72 sec in *RluA-1[del-HDR]/RluA-1[del-*

FRT], n=195 vs. 6.17 ± 3.95 sec in +/+, n=125). The genetic background is w^{1118} for *RluA-1[del-HDR]*, and

isow^{1118} for *RluA-1*$^{del\text{-}FRT}$. For the *RluA-1*$^{del\text{-}HDR}$/*RluA-1*$^{del\text{-}FRT}$ transheterozygotes, data were pooled

from the progeny from reciprocal crosses of *RluA-1*$^{del\text{-}HDR}$ to *RluA-1*$^{del\text{-}FRT}$. To generate the control

larvae, reciprocal crosses were made between the genetic background of w^{1118} and isow^{1118} and

the data from the progeny of these crosses were pooled. Significance of the comparison is

marked as **** (exact p<0.0001). Data were analyzed using Mann-Whitney U test.

Fig 3: ***RluA-1* nociception phenotypes with (A) chromosome duplication or (B) Class IV-**

specific expression of *UAS-RluA-1 cDNA*. (A). The hypersensitive nociception phenotype with

42▯ thermal stimulus in *RluA-1*$^{del\text{-}HDR}$ null mutant was rescued by introducing a duplication (Dp)

on the third chromosome which covers the *RluA-1* gene region (average latency of 8.02 ± 3.61

sec in *RluA-1*$^{-/-}$; Dp/Dp, n=38, which is similar to the isow^{1118} genetic background (+/+, average

latency of 10.16±3.87 sec, n=22) , but significantly slower than the latency of 5.31 ± 2.16 sec in

RluA-1$^{-/-}$, n=34 (p=0.02)). Larvae with one copy of duplication showed slower response (average

latency of 6.90 ± 4.81 sec in *RluA-1*$^{-/-}$; *Dp/+*, n=45) without statistical significance compared to

RluA-1$^{del\text{-}HDR}$. Data were analyzed using Kruskal-Wallis test with Dunn's multiple comparison

tests. **(B)**. The hypersensitive thermal nociception phenotype in *RluA-1*$^{del\text{-}HDR}$ larvae was

completely reversed to that of the heterozygous *RluA-1* by Class IV specific expression of full

length *RluA-1-cDNA* (*RluA-1*$^{-/-}$; *ppk-Gal4*$^{+/-}$; *UAS-RluA-1*$^{+/-}$, average latency of 12.37 ±6.48 sec,

n=30, is similar to *RluA-1*$^{+/-}$; *ppk1.9-Gal4*$^{+/-}$, average latency of 11.04 ± 5.44 sec, n=20 but

significantly slower than either the driver alone (*RluA-1*$^{-/-}$; *Ppk1.9-Gal4*$^{+/-}$, average latency of

3.78±2.11 sec, n=25 or transgene alone controls (*RluA-1*$^{-/-}$; *UAS-RluA-1*$^{+/-}$, average latency of 2.86

± 1.39 sec, n=33) at 42▯ thermal stimulus. Significance of the comparisons are marked as ****

(p<0.0001). Data were analyzed using Kruskal-Wallis test with Dunn's multiple comparison tests.

Fig 4: ***RluA-1* mechanical nociception and gentle touch responses. (A)**. In response to the

noxious mechanical stimulus of 30mN/720kPa, all (100%) of the *RluA-1*$^{del\text{-}HDR}$ null mutant larvae

(*RluA-1*, n=28) rolled compared to the 78.57% of control larvae (iso*w*^{1118} n=28) rolling. At the

reduced stimulus of 15mN/360kPa, 95.56% of *RluA-1*$^{-/-}$ (n=45) rolled while only 48.53% of

control (n=68) rolled. Significance of the comparisons are marked as *(p<0.05) and ****

(p<0.0001). Data were analyzed using Fisher's exact test and presented as percentages ± 95%

confidence intervals. (**B**). *RluA-1*$^{del-HDR}$ larvae (*RluA-1*$^{-/-}$) had a gentle touch response score (6.70

± 2.48, n=36) similar to the control larvae (iso*w*^{1118}, 7.83 ± 2.69, n=30). ns, not significant

(p>0.05). Data were analyzed with Student's t test.

Fig 5: **RluA-2 expression in *Drosophila* larvae. (A)**. In third instar larvae of *RluA-2-GFSTF* line,

anti-GFP (green) immunoreactive signals are detected in the nuclei of all types of cells (neurons

are labelled with anti-HRP (red)). Scale bar = 100μm. **(B).** In the larval abdominal dorsal PNS

cluster of *RluA-2-GFSTF* line, GFP signal (green) is also detected in the nuclei of the md neurons,

whose membranes are marked with anti-HRP (magenta). Arrow points to Class IV ddaC neuron.

Scale bar = 50μm.

Fig 6. **Thermal nociception responses of *RluA-1* and *RluA-2* single mutant, and *RluA-1 RluA-***

2 double mutant. Larvae of single mutant *RluA-1*$^{del-HDR}$ (*RluA-1*$^{-/-}$), *RluA-2*$^{del-HDR}$ (*RluA-2*$^{-/-}$),

double mutant *RluA-1*$^{del-HDR}$ *RluA-2*$^{del-HDR}$ (*RluA-1*$^{-/-}$ *RluA-2*$^{-/-}$) and the genetic background iso*w*^{1118}

(+/+) were stimulated with noxious heat probe of 42⬚. *RluA-1*$^{-/-}$, *RluA-2*$^{-/-}$, and *RluA-1*$^{-/-}$ *RluA-2*$^{-/-}$ all

displayed faster responses to the stimulus compared to controls (*RluA-1*$^{-/-}$ average latency of

4.32 ± 2.80 sec, n= 79; *RluA-2*$^{-/-}$, 6.63 ± 3.31 sec, n=68; (*RluA-1*$^{-/-}$ *RluA-2*$^{-/-}$), 6.26 ± 3.46 sec, n=57;

+/+, 13.72 ± 6.03 sec, n=46). Significance of comparisons are marked as ** (p<0.01), ***

(p<0.001) or **** (p<0.0001). Data were analyzed using Kruskal-Wallis test with Dunn's

multiple comparison tests.

Fig 7. ***RluA-1* ddaC neuron dendrite morphology. (A-C)**. Quantification of the branch number

(**A**), average branch length (**B**), and isoneuronal cross-over points (**C**). (**A**). Homozygous *RluA-*

$1^{del\text{-}HDR}$ larvae showed a higher normalized branch number (total number of branches / (neuron

size x $10^{-4}\mu m^{2}$), 32.56 ± 7.83 in $RluA\text{-}1^{-/-}$; $ppk\text{-}tdTom$, n=6 vs 23.10 ± 4.96 in $ppk\text{-}tdTom$ control,

n=6, p<0.05). (**B**). The mean ddaC dendritic branch length in homozygous $RluA\text{-}1^{del\text{-}HDR}$ was

shorter than that of the control (average branch length is 14.01 ± 1.29 μm in $RluA\text{-}1^{-/-}$; $ppk\text{-}$

$tdTom$, n=6 vs 16.30 ± 1.07 μm in $ppk\text{-}tdTom$ control, n=6, p<0.01). (**C**). The number of ddaC

dendritic isoneuronal crossovers in homozygous $RluA\text{-}1^{del\text{-}HDR}$ was higher compared to that of

the control (normalized crossover points = crossover points / (neuron size x $10^{-7}\mu m^{2}$), $25.03 \pm$

8.24 in $RluA\text{-}1^{-/-}$; $ppk\text{-}tdTom$, n=6 vs 13.48 ± 4.08 in $ppk\text{-}tdTom$ control, n=6, p<0.05). Significant

differences are marked as * (p<0.05) and ** (p<0.01). Data were analyzed with Student's t test.

(**D**). Representative traced dendritic structure of a class IV ddaC neuron of an L3 larva

expressing $ppk\text{-}GAL4 > UAS\text{-}td\text{-}Tom$ in homozygous $RluA\text{-}1^{del\text{-}HDR}$ ($RluA\text{-}1^{-/-}$, **left**) in comparison

with that of the $ppk\text{-}tdTom$ control (+/+, **right**). The position of the cell body is marked with blue

circle and axon with red circles.

A

B
ES
ddaD   ddaE
ddaF
ddaC
                ES
ddaB   dmd1
ddaA
dbd

C

D¹   D²   D³
anti-GFP   RFP   merge

Figure 1

Figure 2

Figure 3

A

B

Figure 4

Figure 5

Figure 6

+/+   RluA-1⁻/⁻

Figure 7