The Chloroplast RNA binding protein CP31A shows cold-dependent RNA association and mediates cold-resistance via its acidic domain

Ayako Okuzaki[#$1], Marie-Kristin Lehniger[1$], Jose M Muino[2], Benjamin Lenzen[1], Thilo Rühe[3], Dario Leister[3], Uwe Ohler[2], Christian Schmitz-Linneweber[1*]

[1] Molecular Genetics, Humboldt-University Berlin, Philippstr.13, 10115 Berlin, Germany
[2] Computational Regulatory Genomics, Humboldt-University Berlin / Max Delbrück Centre for Molecular Medicine, Philippstr.13, 10115 Berlin, Germany
[3] Plant Molecular Biology, Department of Biology, Ludwig Maximilian University of Munich, Munich, Germany.

[#] current address: College of Agriculture, Tamagawa University, Tamagawagakuen 6-1-1, Machida-shi, Tokyo, Japan

[$] equally contributed

*corresponding author:

Christian Schmitz-Linneweber
phone: +49 30 2093-49700
fax: +49 30 2093-49701
Email: smitzlic@rz.hu-berlin.de

Running title: Cold resistance by a chloroplast RNA binding protein

Key words: chloroplast, RNA processing, RNA binding, acidic domain, RNA editing, RNA stability, organelle, *Arabidopsis thaliana*

Materials distribution footnote:

Abstract

Chloroplast gene expression is characterized by a multitude of post-transcriptional events as well as long-lived mRNAs. Given that the chloroplast is responsible for expressing core components of the photosynthetic apparatus, it is not surprising that the accumulation and processing of chloroplast RNA is heavily affected by various external and internal cues, including light and temperature changes. A multitude of nuclear-encoded RNA-binding proteins (RBPs) are known to be required for chloroplast RNA metabolism, but we do not yet know how chloroplast RBPs convert abiotic signals into gene expression changes. Previous studies showed that chloroplast ribonucleoprotein 31A (CP31A) is required for the stabilization of multiple chloroplast mRNAs in the cold, and that the phosphorylation of CP31A at various residues within its N-terminal acidic domain (AD) can alter its affinity for RNA *in vitro*. Here, we demonstrate that CP31A shows increased affinity for a large number of chloroplast RNAs *in vivo*. Loss of CP31A leads to cold sensitivity, and cold-dependent RNA binding and cold resistance both depend on the AD of CP31A. Together, our analysis establishes the AD of CP31A as a key mediator of cold acclimation of the chloroplast transcriptome.

Introduction

Chloroplasts contain genetic information that is essential for photosynthesis. The expression of this information is realized by a unique mixture of ancestral bacterial and derived eukaryotic features (Barkan, 2011). Chloroplast gene expression adapts to various environmental changes, including light and temperature (e.g. Klein, 1991; Mentzen and Wurtele, 2008; Cho et al., 2009; Castandet et al., 2016). Contributions to such acclimation processes have been described on the transcriptional level (Pfannschmidt, 2003; Tsunoyama et al., 2004), but post-transcriptional processes are likely to dominate (Deng and Gruissem, 1987; Eberhard et al., 2002; Udy et al., 2012). One key change in posttranscriptional processes between chloroplast and their bacterial ancestors are the vastly increased RNA half lives in the organelle. In bacteria, transcription and translation are usually directly coupled and mRNAs have short half-lives (in the range of minutes; Selinger et al., 2003). In chloroplasts, in contrast, the half-lives of mRNAs are long (in the range of hours) and untranslated RNAs accumulate in large amounts (Klaff and Gruissem, 1991; Germain et al., 2012). The turnover rates of chloroplast RNAs change in response to developmental and environmental cues, which is suggesting that RNA stability is regulated in chloroplasts (Deng et al., 1989; Klaff and Gruissem, 1991; Biehl et al., 2005; Bollenbach et al., 2007; Germain et al., 2013; Manavski et al., 2018), but the underlying regulatory factors are largely unknown. Possible candidates for regulators of RNA stability are pentatricopeptide repeat (PPR) proteins and chloroplast ribonucleoproteins (cpRNPs). PPR proteins specifically associate with one or few RNAs (Barkan and Small, 2014), while cpRNPs are generalists that bind to a large number of mRNAs (Kupsch et al., 2012; Teubner et al., 2017). Two cpRNPs

named CP29A and CP31A were previously shown to be required for the accumulation of chloroplast RNAs

in the cold, which makes them interesting candidates to be mediators of global RNA stability during the

acclimation to changing temperature conditions (Kupsch et al., 2012).

The cpRNP protein family consists of ten members in *Arabidopsis thaliana* (Ruwe et al., 2011). All cpRNPs

are targeted to the chloroplast post-translationally, and dedicated import receptors appear to be responsible

for their transport across the chloroplast envelope (Li and Sugiura, 1990; Grimmer et al., 2014). cpRNPs

are highly regulated proteins that react to various external and internal signals, particularly light, which

controls both their expression and their protein modification state (summarized in Ruwe et al., 2011).

Several cpRNPs have been identified as phosphoproteins (Reiland et al., 2009) and an N-terminally

acetylated isoform of CP29A was shown to respond rapidly to changes in light and developmental stages

(Wang et al., 2006). The phosphorylation of cpRNPs can alter their RNA-binding characteristics *in vitro*

(Lisitsky and Schuster, 1995; Loza-Tavera et al., 2006), but evidence for condition-dependent RNA binding

of cpRNPs or any other chloroplast RBPs *in vivo* are lacking.

Several molecular functions have been suggested for cpRNPs. *In vitro*, tobacco CP31 has been shown to

support RNA editing of multiple sites (Hirose and Sugiura, 2001). Other cpRNPs have been reported to be

required for the 3'-end processing of several mRNAs (Schuster and Gruissem, 1991; Hayes et al., 1996;

Schuster et al., 1999). They also support the ribozymatic maturation of a viroid RNA genome (Daros and

Flores, 2002). The most notable and general function of cpRNPs is however their role in stabilizing mRNAs,

which has been shown *in vitro*, where they protect the mRNA encoding the D1 subunit of photosystem II

against degradation (Nakamura et al., 2001), as well as *in vivo*, where they are required for the accumulation

of a multitude of mRNAs (Kupsch et al., 2012; Teubner et al., 2017). Co-immunoprecipitation analyses

have demonstrated that cpRNPs are associated with multiple chloroplast RNAs, and that they prefer

unprocessed (unspliced) RNAs over mature forms (Nakamura et al., 1999). Together, the data obtained from

these functional analyses show that cpRNPs have a broad target range and contribute to various RNA-

processing events and to RNA stabilization.

All cpRNPs share a similar design, with two RNA recognition motifs (RRMs) that are preceded by a domain

rich in acidic residues (acidic domain; AD). While the RRMs are well characterized RNA binding domains

(Li and Sugiura, 1991; Ye and Sugiura, 1992; Lisitsky et al., 1995), the role of the AD remains to be

determined. The cpRNP CP31A stands out from the family in having a particularly long acidic domain,

which contains two phosphorylated serine residues within a short repeat element of five amino acids

(Reiland et al., 2009). CP31A's AD was suggested to be important for RNA editing based on *in vitro* assays

(Hirose and Sugiura, 2001). The AD of the spinach homolog of CP31A is supportive of RNA binding, but

does not bind RNA itself (Lisitsky and Schuster, 1995). Recently, plastid casein kinase II (pCKII) was

demonstrated to phosphorylate the acidic domain of CP31A *in vitro* (Schonberg et al., 2014). Genetic

analyses have demonstrated that Arabidopsis CP31A supports RNA editing at multiple sites and modulates

the stability of multiple mRNAs (Tillich et al., 2009; Kupsch et al., 2012). An RNA strongly affected in

cp31a mutants was the *ndhF* mRNA, which encodes a subunit of the NDH complex. CP31A binds to the

3'-UTR of *ndhF* and is required for the generation of the *ndhF* 3'-terminus (Kupsch et al., 2012). Mutants

of CP31A display cold sensitivity: the germination rate of null mutants is reduced and their newly emerging

leaf tissue bleaches at 8°C (Kupsch et al., 2012). All analyzed proteins of the photosynthetic apparatus are

reduced in this defective tissue, which is at least in part due to multiple defects in RNA processing (e.g.,

RNA splicing, RNA editing, and intercistronic processing), but is likely primarily caused by the strong

reduction of multiple chloroplast mRNAs (Kupsch et al., 2012). However, it remains unclear how CP31A

mechanistically confers cold resistance and whether it directly perceives cold as a signal. We hypothesized

that the AD could act as a regulatory domain for RNA binding and cold-responsiveness. We here therefore

analyzed the ability of CP31A to associate with RNA in response to cold and assessed the role of the AD

for CP31A's ability to bind and process chloroplast RNA. Our results demonstrate that the AD of CP31A

is essential for plant cold acclimation, at least in part because of its supportive role for cold-dependent RNA

binding.

Results

The RRM domains of CP31A are sufficient for RNA editing of CP31A-dependent sites

The binding of RBPs to their RNA targets can be altered by protein modifications. For several members of

the cpRNP family, phosphorylation has been demonstrated to alter RNA binding *in vitro* (Lisitsky and

Schuster, 1995; Loza-Tavera et al., 2006; Reiland et al., 2009). As phosphoproteomic analyses identified

several phosphorylation sites within the acidic N-terminal domain of CP31A (Reiland et al., 2009;

Schonberg et al., 2014), we decided to test the function of this acidic phosphodomain by deletion

mutagenesis. We designed three T-DNA based constructs to express CP31A variants in Arabidopsis (**Fig.**

1A). The first two expressed variants lack the AD. One was driven by the 35S promoter, while the other

was driven by the endogenous genomic region upstream of the transcriptional start site, presumably

encompassing the unknown *CP31A* promoter (12769646 to 12767953 of the *Arabidopsis thaliana*

chromosome 4 sequence; nucleotides –1 to –1694 relative to the *CP31A* start codon). These constructs were

introduced into a *cp31a*-deficient background (homozygous *cp31a*-1 null allele) by Agrobacterium-

mediated gene transfer. In addition, the full-length protein driven by its native promoter was used as a

positive control. All plant lines retrieved showed normal development and were indistinguishable from wt

plants under normal growth conditions (**Fig.** 4A).

We assessed CP31A protein accumulation in our transgenic lines by using an antibody that reacts to two peptides: one situated in the linker between the two RRM domains and another at the C-terminus of the protein (Kupsch et al. 2012). The antibody thus detects both the full-length as well as the AD-deletion proteins. We observed accumulation of proteins that were of the appropriate size for all transgenic lines (**Fig.** 1B). These data demonstrate that the full-length and AD-deficient proteins were expressed and specifically detected by the antibody. We quantified signals from three independent transgenic lines for each construct (**Fig.** 1B, C). As expected, there is variability in expression between individual lines, which can be attributed to position effects of the transgene insertion site. In sum, these analyses demonstrate that all three constructs were successfully expressed in a CP31A-deficient background.

Figure 1: Construction of AD deletion mutants.
A) Schematic overview of constructs used to complement *cp31a* null mutants. bar = selectable marker; black arrows indicate promoters. TP = transit peptide. NosT = the terminator of the nos gene. LB/RB = borders of the Agrobacterium T-DNA.
B) Western analysis of transgenic lines. Blots were probed with a HSP70 antibody to control for loading and with an antiserum recognizing CP31A. The position of the full-length protein and the truncated, AD-less version are indicated by filled and open arrowheads, respectively.
C) Quantification of CP31A signals normalized to HSP70 signals (from (B)).

Since CP31A has been shown to support RNA editing at 13 specific sites (Tillich et al., 2009), we investigated these and additional sites in our complementation lines by performing next-generation sequencing of cloned amplified cDNA (Bentolila et al., 2013), testing a total of 13 sites on 10 amplicons. This included sites that were previously shown not to be affected in *cp31a* mutants. Two wt and two *cp31a-1* null mutant plants served as controls. The lowest read coverage observed for any individual site was above 500 reads per site and was found for site *ndhD* 116281 (number refers to the position in the Arabidopsis chloroplast genome), while the average coverage was 3132 reads per site for all sites and all genotypes analyzed.

Of the 13 sites previously described to be dependent on CP31A, we confirmed that RNA editing of 11 sites was reduced in *cp31a*-1 mutants (**Fig.** 2A). Additional defects were found for two previously unreported

sites in *ndhB* (94999, 96579), whereas no defect was observed for the previously reported sites, *petL* (65716)

and *ndhB* (952225). We speculate that these differences reflect differences in growth conditions: our plants

were grown under long-day conditions (16-h light/8-h dark), whereas the previous study used short-day

conditions (8-h light/16-h dark; Tillich et al., 2009). Importantly, all our complementation lines, irrespective

of the utilized promoter or the presence of the AD showed wt-like editing states at most analyzed sites (**Fig.**

2A). We confirmed this finding by independent amplification and Sanger sequencing of selected RNA-

editing sites of the *ndhD* and *rps14* mRNAs. The amplicons we sequenced harbored both, CP31A-dependent

and CP31A-independent editing sites (**Fig.** 2B). Our sequencing confirmed that in *cp31a* mutants, sites

ndhD-3, 4 and 5, *rps14*-2 show reduced editing efficiency when compared to wt. This defect is

complemented in the AD-less CP31A protein (**Fig.** 2C). The successful complementation demonstrates that

the RRM domains are sufficient for the RNA-editing function of CP31A.

Figure 2: Analysis of RNA editing sites in cp31a complementation lines.

A) Summary of results from amplicon sequencing of RNA editing sites in complementation lines. Editing sites were
amplified by PCR and sequenced on an Illumina platform using barcoded adapters that allowed to eliminate duplicate
amplifications. Each individual cDNA sequence is scored for its editing status and the frequency of edited versus non-
editied cDNAs is shown. Error bars represent the standard deviation calculated from three replicate WT and *cp31a*
mutant experiments, respectively. Editing sites marked with asterisks are dependent on CP31A. Numbers refer to the
position of the editing site in the *Arabidopsis thaliana* chloroplast genome.
B) Schematic overview of two genes with their editing sites that were assessed by Sanger sequencing of amplified
cDNAs. Arrows indicate positions of primers used for cDNA amplification.

C) Excerpts from electropherograms showing fluorescencec signals for base triplets with the editing site always at the
center. Bases marked by red arrows are target of CP31A. Green traces refer to T signals, blue to C signals and red to
A signals. The expected edited (on cDNA) and unedited (on DNA) triplets are shown above. CR = control sites not
affected by CP31A.
187

The RRM domains of CP31A are sufficient to
stabilize the *ndhF* mRNA

We next analyzed the RNA-stabilizing function of
CP31A by focusing on *ndhF*, which is its major
target under normal growth conditions (Tillich et
al., 2009). Using RNA gel-blot hybridization, we
quantified the amount of *ndhF* mRNA in the
transgenic lines. The accumulation of this transcript
is at least partially restored in all complementing
lines relative to the *cp31a* mutant (**Fig.** 3A, B). Even
in a line with very little CP31A accumulation and
lacking the AD, a signal for *ndhF* is visible (#3-3-3.
Fig. 1A, B). This indicates that the RRM domains
were sufficient to stabilize the *ndhF* mRNA.
However, we observed marked differences in the
efficiency of *ndhF* stabilization between the full-
length and AD-deficient proteins. Even lines
accumulating large amounts of the AD-less CP31A
protein had less *ndhF* mRNA than lines expressing
much less full-length CP31A protein (compare for
example line #3-1-3 with #4-17-1 in Fig. 1B and
Fig. 3A). This was visualized by calculating the
ratio of *ndhF* mRNA levels to CP31A protein levels
(i.e., the stabilization efficiencies; **Fig.** 3C), which

Figure 3: Analysis of the accumulation of the *ndhF* mRNA in cp31a complementation lines.
A) RNA gel blot analysis of the *ndhF* mRNA in various *cp31a* complementation lines. 4 µg total cell RNA were separated by denaturing agarose gel electrophoresis, transferred to a nylon membrane and probed with a radiolabelled DNA probe specific for *ndhF*. The transcript labelled with an arrowhead represents the full-length *ndhF* mRNA, whereas smaller signals correspond to degradation products (Kupsch et al. 2012). The nylon membrane was labeled with methylene blue to control for equal loading (excerpt with 23S rRNA shown below blot).
B) Quantification of the full-length *ndhF* transcript labeled with an arrowhead in (A).
C) Ratio of *ndhF* mRNA levels over CP31A protein levels (from Fig. 1C).

clearly showed that the full-length proteins outperformed the AD-less proteins in stabilizing the *ndhF*
mRNA. Together, these results show that the AD supports but is not essential for stabilizing the *ndhF* mRNA
and that the RRM domains are capable of doing the job by themselves.

The AD is required for cold-stress tolerance

To test the impact of the AD on CP31A-mediated cold tolerance, we challenged 2-week-old plants from the
transgenic lines with low temperature (8°C) for 5 weeks. Under this treatment, the *cp31a*-null lines are
known to display bleaching of the freshly emerging tissue at the center of the leaf rosette (**Fig.** 4A). Similar

losses of pigment were also found in the lines expressing the AD-less CP31A, but not in wt plants or in mutant plants complemented with the full-length protein (**Fig.** 4A). The pigment deficiency has been shown to be paralleled by a global reduction of the chloroplast-encoded proteins required for photosynthesis in *cp31a* mutants (Kupsch et al., 2012). This reduction of photosynthetic proteins was also observed in plants expressing the AD-less CP31A proteins, while plants expressing the full-length CP31A protein accumulated protein amounts similar to those seen in wt plants (Suppl. **Fig.** 1). Next, we tested the accumulation of CP31A and *ndhF* in these lines under cold exposure, harvesting separately full leaves and the bleached center parts of rosettes, respectively. Quantification of protein and RNA signals demonstrated that the levels of CP31A and *ndhF* were well correlated in both tissues (**Fig.** 4B/C), similar to our findings when these plants were grown at standard temperature. In line with the findings at normal growth temperatures, the *ndhF*/CP31A ratio was consistently higher for plants expressing full-length CP31A versus plants expressing the AD-less protein also in the cold (**Fig.** 4B/C and Suppl. **Fig.** 2). These experiments show that the AD of CP31A supports the accumulation of chloroplast RNA and is essential for conferring cold resistance to emerging leaf tissue in Arabidopsis.

Figure 4: Analysis of *cp31a* complementation lines during cold acclimation.
A) Phenotypes of complementation lines grown at 23°C or challenged with 8°C for three weeks. Note the bleached tissue in the center of the rosette in *cp31a* deficient plants and in plants lacking the AD.
B) Analysis of CP31A and HSP70 protein accumulation and *ndhF* mRNA accumulation in *cp31a* complementation lines. Here, the entire rosette leaves were used for protein and RNA extraction. Panels from top to bottom: immunoblot analysis of HSP70 proteins as loading control; immunoblot analysis of CP31A proteins; RNA gel blot analysis of *ndhF*; Methylene

blue stain of cytosolic rRNA as loading control; Ratio of quantified *ndhF* (transcript 'a') and CP31A signals. Filled
and open arrowheads indicate full-length and AD-less CP31A, respectively.
C) Same analysis as in (B), but with tissue only form the centre of the rosette (bleached area in *cp31a* null mutants).

CP31A associates with transcripts encoding subunits of the NDH complex and is required for their accumulation

RBPs of various origins and organisms show differential expression under temperature changes, and genetic studies indicate that they are important for temperature-dependent RNA processing (Lin et al., 2010; Malay et al., 2011; Gotic et al., 2016). That said, few studies have examined whether the binding of RBPs to their RNA targets is modified in response to external and internal cues *in vivo*. Since CP31A is required for cold resistance, we herein tested whether its ability to bind RNA is modified at low temperature and also assayed the importance of the AD in this response. Previous studies using low-resolution techniques could only identify whole transcripts as targets of CP31A (Kupsch et al., 2012). We here used a RIP-Seq approach to obtain better resolution (Suppl. **Fig.** 3A). For this, wt *Arabidopsis* seedlings and plants expressing the AD-less CP31A protein in a *cp31a* background were grown for 14 days under normal growth conditions and then subjected to formaldehyde cross-linking. For cold tests, seedlings were grown for 13 days at the standard growth temperature (21°C), transferred to 4°C for 24 hours, and then subjected to cross-linking. Immunoprecipitation (IP) was performed using a previously described antibody that targets the linker domain between the two RRM motifs of CP31A (Kupsch et al., 2012). The efficiency of precipitation was comparable between samples grown at normal and low temperatures (Suppl. **Fig.** 3b). RIP-seq libraries were prepared in duplicate from input and pellet samples (Suppl. **Fig.** 3C). We tested the reproducibility of the RIP-seq experiments by calculating pairwise correlation coefficients across all samples (Suppl. **Fig.** 3D). We found strong correlations among the input samples (average Pearson coefficient (R): 0.95), and the pellet samples from the IP (average Pearson coefficient (R): 0.97). As expected, input and pellet samples formed separate clusters (Suppl. **Fig.** 3D). This analysis demonstrates high reproducibility between biological replicates.

For the analysis of RIP-Seq reads, we followed established procedures originally developed for ChIP-Seq (Bardet et al., 2011; Muino et al., 2011). We started by identifying transcript regions with an enrichment of RNA in pellet samples over input samples. In total, 75 different binding regions were identified as significant in at least one of the four experiment, which are henceforth called binding sites (BSs). The 75 BSs of CP31A are located in 44 transcripts of diverse functionality. Of these 44 transcripts, 31 were also identified by previous RIP-Chip experiments (Suppl. **Tab**. 1, Suppl. **Fig.** 4). The difference observed is likely due at least in part to methodological dissimilarities. Purified chloroplast stroma was used for the previous RIP-Chip experiments without a cross-link, while here, we used cross-linked frozen total leaf tissue in the RIP-seq. Given these technical differences, it was gratifying to see a similar set of transcripts enriched in the two types of RIP experiments. We next focused on those BSs that can be clearly assigned to a specific

gene since they are located within an open reading frame. Such BSs are are found in genes for the
photosynthetic complexes as well as for the gene expression apparatus (**Tab. 1)**. We compared the actual
distribution of BSs with the calculated numbers of BSs expected for each functional category (if random
binding in the transcriptome is assumed; **Tab. 1**). This demonstrated that the NDH complex is the only
functional category overrepresented among RIP-Seq targets (12 found versus 3 expected; **Tab. 1**). We also
found an unexpected large number of binding sites antisense to coding regions of known genes (**Tab. 1**).
Although antisense RNAs are known to exist in chloroplasts, their levels do not (with a few exceptions)
reach those of sense RNAs and their functions (if any) remain unclear (Nakamura et al., 2003; Hotto et al.,
2010; Hotto et al., 2011; Sharwood et al., 2011). Consequently, the functional significance of CP31A'a
association with antisense RNAs remains unclear as well. It is however interesting that at least for the main
target category, the *ndh* genes, a preferences for sense sites is evident (**Tab. 1**), suggesting that CP31A is
relevant for *ndh* mRNA expression. We therefore analyzed chloroplast RNA levels in *cp31a* versus wt
samples using an oligonucleotide tiling array that represents the entire chloroplast genome of *Arabidopsis*
thaliana. We scored only probes whose RNA levels were at least one third lower in the mutant compared
to wt. Of all exon probes in the array, 16% contained *ndh* sequences, but 75% of all exon probes whose
signals were decreased by at least 0.6-fold contained *ndh* sequences (**Fig**. 5A,B). No other functional
category showed enrichment in this analysis.

To confirm the results of our microarray analysis, we performed RNA gel-blot hybridization experiments
for four *ndh* genes that represented the four *ndh* operons in the chloroplast genome (**Fig**. 5C). We analyzed
cp31a mutant RNAs alongside RNAs from seedlings and control lines with impairment in the expression of
SIGMA FACTOR 4 (*SIG4*), which is specifically required for the transcription of the *ndhF* mRNA (Favory
et al., 2005), and *CHLORORESPIRATORY REDUCTION 2* (*CRR2*), which is required for the accumulation
of monocistronic *ndhB* transcripts (Hashimoto et al., 2003). As expected, the control mutants showed
specific defects for their known target RNAs (**Fig**. 5C). By contrast, the *ndhK* transcripts accumulated to
normal levels in these two control lines, and the *ndhD* mRNA was only slightly decreased (**Fig**. 5C). In
contrast and consistent with our microarray results, the *cp31a* mutants displayed reductions in all four *ndh*

transcripts analyzed. The decrease was most pronounced for *ndhF*, strong for *ndhK* and *ndhB*, and somewhat weaker for *ndhD* (**Fig**. 5C). Collectively, these findings indicate that CP31A stabilizes mRNAs from all four *ndh* operon-encoding genomic loci, suggesting that CP31A regulates *ndh* mRNAs as a group.

Arabidopsis **CP31A prefers U-rich sequences** *in vivo*

Prior *in vitro* experiments showed that a CP31A homolog in tobacco prefers homopolymers of G and U (Li and Sugiura, 1991). We asked whether a sequence preference can be identified in our dataset, i.e. *in vivo,* as well. We used the SSMART motif finder (Munteanu et al., 2018) to query BSs in our RIP-Seq data for common sequences. This uncovered consensus motifs with a clear preference for Us (**Fig.** 6A; Suppl. **Fig.** 5). We next compared the frequencies of all possible trinucleotides within significant CP31A BSs against those in all chloroplast coding sequences. Indeed, we

Figure 5: Analysis of RNA accumulation in *cp31a* mutants.
A) Summary of microarray analyses of wt and *cp31a* mutants. Relative abundance of probes showing at least 1.5 fold stronger signals in wt versus *cp31a* mutants. Two replicate microarray hybridizations were considered and assigned to different gene categories.
B) Relative distribution of all probes on the microarray to different gene categories.
C) RNA gel blot analysis of 14 days old Arabidopsis seedlings. 4 µg RNA from from wt, *cp31a* mutants and two control mutants (*crr2* and *sig4*) together with dilutions of wt probes (1/2 and 1/4) were probed with radiolabeled DNA probes against four different *ndh* genes. The resulting autoradiographs are always shown with the corresponding methylene blue stains of the membranes (below). The 2kb marker band is shown as a reference.

found that UUU was the most strongly enriched

trinucleotide in this comparison, followed by a number of other U-rich trinucleotides (**Fig.** 6B). We therefore

conclude that CP31A preferentially binds U-rich sequences. As U-rich trinucleotides are the most common

340 trinucleotides in the AT-rich chloroplast genome, our findings suggest that CP31A has evolved to allow a
341 broad RNA target range in the chloroplast transcriptome.

Figure 6: Analysis of sequence elements in CP31A BSs
A) Top scoring consensus motifs identified in CP31A BSs. The SSMART RNA motif finder was run using all significant BSs detected in WT plants grown in cold conditions, as under these conditions, the number of BSs was the highest. The input sequences of the respective BSs made a contribution to the consensus motifs according to their adjusted p-value - the more significant the BS, the stronger its impact on the consensus.
B) Overrepresentation of the TTT trinucleotides in CP31A BSs. The scatterplot shows the frequency of all possible trinucleotide sequences in the CP31A BSs regions (y-axis) versus the trinucleotide frequency in all chloroplast coding sequences. Trinucleotides with significant enrichment in RIP-seq BS regions are marked in red ($p<0.01$) or blue ($p<0.05$).

CP31A shows an increased association with RNA after cold exposure

We hypothesized that CP31A might impact cold resistance in Arabidopsis plants through temperature-
dependent changes in its affinity for some or all of its target RNAs, which could affect their stability,
processing, and/or translation. We therefore compared BSs according to how reliable their enrichment in a
given dataset is, based on its FDR value. Using the FDR values, we will first focus on the effect of cold on
RNA binding of full-length CP31A before discussing the effect of the loss of the AD on binding in the next
chapter. A majority of BSs were found with high significance in CP31A precipitates in the cold, but were
not significantly enriched at higher temperature. This was already visible for some sites in the coverage
graphs, as shown in **Fig.** 7A for binding sites within the *ndhJ* and *rpl2* mRNAs, or for a site in the intergenic
region between the *rps14* and *psbZ* genes. Many BSs show at least some read coverage at both temperatures,
but the FDR analysis separates significant from non-significant BSs. The FDR level is displayed for all 75
binding sites as a heat map in Suppl. **Fig.** 5 and is summarized in **Fig**. 7B. Most importantly, 30 BS show

significant enrichment with full-length CP31A at 8°C, but not at 23°C (**Fig**. 7B). Only 9 sites show the
opposite behavior, that is, significant enrichment only at 23°C, but not at 8°C. Thus many more sites showed
increased versus decreased association with CP31A after cold exposure. Three of the BSs gained in the cold
were located in the *ndhF* mRNA, which is known to be a prime target of CP31A (Kupsch et al., 2012). Four
more CP31A BSs in *ndh* transcripts were cold-dependent as well (Suppl. **Fig.** 6). In general, *ndh* mRNAs
appeared to be important targets of CP31A under both temperature conditions (Suppl. **Fig.** 6). Taken
together, the results of our temperature dependent RIP-Seq analysis demonstrate that RNA binding of full-
length CP31A is higher under cold exposure than at normal temperature, and that *ndh* mRNAs are prime
targets under both normal and low temperatures.

The AD of CP31A supports cold-dependent RNA binding

The RRM domain is a highly versatile protein domain that functions in the context of adjacent protein
regions. Often, linkers between RRMs or protein domains outside the RRM domains support RNA
interactions and can make direct contact with RNAs in addition to the canonical RRM contacts (Lunde et
al., 2007). Given this, we used RIP-Seq to test whether the RNA-binding behavior of the AD-less CP31A
protein differed from that of the full-length protein. Using the same strategy described for our binding-site
analysis with the full-length protein, we scored significant BSs in plants expressing the AD-less variant
grown under both normal and cold conditions. We found that the AD-less version of CP31A bound far fewer
RNAs than its full-length counterpart (**Fig.** 7B). Under normal growth temperatures, we identified 41
significant BSs (FDR < 0.05) for the full-length protein and only 10 for the AD-less version (**Fig.** 7B).
Interestingly, the AD-less protein is still responsive to cold: The AD-less protein associated with 18 sites in
the cold that are not found to co-precipitate significantly with the protein at normal temperatures (**Fig.** 7B).
On the other hand, AD-less CP31A bound 39 fewer sites than the full-length protein after cold exposure
(**Fig.** 7B). This demonstrates that the AD supports RNA binding under both conditions. Together, our results
show that (i) the AD substantially supports but is not essential for RNA binding; and that (ii) the remainder
of the protein, which almost exclusively consists of the two RRM domains, can itself react to lowered
temperatures.

Figure 7: RIP-Seq uncovers cold-induced RNA binding of CP31A.
A) Visualization of three examples for temperature-dependent CP31A binding sites. Top track shows the normalized read coverage of RIP-Seq experiments from pellet fractions of WT plants grown at 8°C or 23°C (for two biological replicates at each growth temperature). The track below represents in grey boxes the identified binding sites. The bottom track shows the gene models in black and an intron in *rpl2* as a hatched line. The numbers on the thin black line refer to positions on the Arabidopsis chloroplast genome. The red rectangles indicate temperature-dependent CP31A BS.
B) Venn diagram summarizing the occurrence of binding sites in the two genotypes and the two temperature regimes used for growing the plants. For each genotype-temperature combination, only the BS with an FDR < 0.05 were counted.

Discussion

CP31A co-regulates *ndh* genes

During the evolution of chloroplasts from cyanobacterial ancestors, operon structures were disrupted and operons were shuffled. Many chloroplast operons therefore include genes with different functions. The *ndh* genes, for example, are separated into four transcriptional units in Arabidopsis and are mixed with genes from other functional categories. This lack of conservation of operon structures suggests that transcriptional units are less important than other processes for *ndh* gene regulation in the chloroplast context, giving way to post-transcriptional processes. Translation plays an important role for regulation (Zoschke and Bock, 2018), but it remains unclear, how post-transcriptional co-regulation is achieved prior to translation. We herein demonstrate that CP31A associates with many *ndh* mRNAs under both normal and low temperatures. All analyzed *ndh* mRNAs were reduced in *cp31a* mutants under normal growth conditions, indicating that the CP31A-*ndh* interaction is functionally important. Since the loss of CP31A does not impede transcription (Kupsch et al., 2012), we conclude that the protein is required for the stability of *ndh* mRNAs. An alternative explanation is that the loss of one RNA of the NDH complex has an indirect, hierarchical effect on the other *ndh* RNAs that yields the observed reduction in all *ndh* mRNAs. This phenomenon has been well described for hierarchical protein synthesis cascades in chloroplasts (Choquet et al., 2001), but it has not been previously shown to impact mRNA synthesis or stability. Moreover, the likelihood of this scenario is

weakened by our observation that the losses of *ndhF* or *ndhB* in the *sig4* or *crr2* mutants, respectively, were

not followed by the loss of all *ndh* mRNAs. Taken together, our data indicate that CP31A combines ndh

mRNAs from different genomic loci into a post-transcriptional operon in the chloroplast. Similar post-

transcriptional operons, or "RNA regulons", have been well-described in fruit flies, budding yeast, and

mammalian cells (Keene, 2007). In most cases, these RNA operons function in the combined translation

and/or stabilization of the participating RNAs (Gerber et al., 2004; Lykke-Andersen and Wagner, 2005;

Townley-Tilson et al., 2006). Given the RNA-stabilizing roles of cpRNPs in general and of CP31A in

particular, we propose that CP31A adjusts the RNA stability of the *ndh* genes.

The AD contributes to the ability of CP31A to recognize RNA

CP31A has two canonical RRM motifs, which contain the expected aromatic amino acid residues in the

RRM's RNP1 and RNP2 motifs known to be crucial for RNA binding (Ruwe et al., 2011). Here, we

show that these RRMs are sufficient to edit chloroplast RNAs and stabilize the *ndhF* mRNA but. Still,

compared to the wt protein, more AD-less protein is needed to stabilize a similar level of *ndhF* mRNA.

Moreover, loss of the AD decreases the ability of CP31A to bind RNA. Thus, the AD domain supports but

is not essential for the RNA-binding and RNA-stabilizing functions of CP31A. We do not yet know how

the AD achieves its effects at the molecular level. Two non-exclusive possible modes of action are (i) a

direct involvement in protein-RNA interactions, and (ii) a role in protein-protein interactions, both already

described in non-plant systems for acidic domains (Lunde et al., 2007). Such functional diversity of an AD

has been described for two structural relatives of CP31A, the non-plant RNA binding proteins hnRNP Q

and hnRNP R, that also have an N-terminal AD and C-terminal RRM motifs (Geuens et al., 2016). The AD

of hnRNP R was shown to support RNA binding and is essential for the role of hnRNP R in the re-initiation

of transcription (Fukuda et al., 2013). Similarly, the AD of hnRNP Q forms part of the RNA interaction

surface (Hobor et al., 2018) and in addition interacts with the editing factor Apobec (Quaresma et al., 2006).

These examples show that it is conceivable that the AD of CP31A contributes to protein-protein interactions

(e.g., with other RNA-binding proteins) and also contributes directly to RNA binding. This could serve to

increase the affinity for RNA and mediate RNA stabilization and RNA-processing events. A future

structural analysis of CP31A bound to RNA would greatly help to understand the mechanistic function of

CP31A's AD.

Cold-dependent RNA association of CP31A

Many RBPs show differential association with RNA depending on physico-chemical conditions *in vitro*,

but relatively little is known about how external and internal cues modulate the target spectrum and target

affinity of an RBP *in vivo*. Changed binding to RNA due to external signals has been shown for mammalian

RNA binding proteins (Rousseau et al., 2002; Cloutier et al., 2018), but even here, rarely on a genome-wide

scale (Benegiamo et al., 2018) and comparable data are missing for plants. Our present results on CP31A
show that plant RNA-binding proteins can change their RNA target profiles in response to an external signal,
here low temperatures, but it remains unclear how this differential RNA binding is induced. Cold is expected
to change RNA structures and CP31A can bind dsDNA *in vitro*, albeit with less affinity than ssDNA and
ssRNA (Li and Sugiura, 1991). It is also notable that related spinach cpRNPs have been implicated in the
establishment of the structured 3'-ends of several mRNAs *in vitro* (Schuster and Gruissem, 1991; Lisitsky
et al., 1995; Hayes et al., 1996). These RNA termini require the formation of stem-loop RNA structures,
and cpRNP-like proteins have been found to associate with such stem-loops in UV cross-linking
experiments (Stern et al., 1989; Chen and Stern, 1991; Danon and Mayfield, 1991; Nickelsen and Link,
1993). Thus, the binding of CP31A to RNAs under cold exposure could reflect the emergence of RNA
structures that become more stable at lower temperatures. An *in vivo* analysis of the structure of CP31A
target sites under different temperatures could help address the question of why CP31A associates more
extensively with RNA in the cold.

What is the functional consequence of the increased RNA association of CP31A under cold exposure? Given
that chloroplast RNAs are lost in the bleached tissue of *cp31a* mutants, we propose that an increased
interaction with target transcripts could impact the resilience of these transcripts against RNases. An
interesting open question is, which of the many CP31A – transcript interactions in the cold is key to avoid
bleaching. Although it is likely that many of these interactions play a role, we can make some guesses about
major contributors. Loss of *ndh* mRNAs, although a prime target of CP31A, is unlikely to be responsible
for the bleaching phenotype since *ndh* genes are downregulated in the cold in Arabidopsis (Ivanov et al.,
2012), and mutants devoid of the NDH complex do not show bleaching in the cold (Li et al., 2004; Yamori
et al., 2011). By contrast, there is a number of other CP31A target RNAs that are known to be essential for
chloroplast development and cold-dependent loss of their processing and stabilization could well explain
bleaching. This includes mRNAs for subunits of the ATP synthase (*atpA, atpB*), of the cytochrome b_6f
complex (e.g. *petB, petD*), of the photosystems (e.g. *psaA, psbB*), as well as mRNAs for subunits of the
plastid-encoded RNA polymerase (*rpoB, rpoC2*). Loss of the expression of any of these genes leads to
defective chloroplasts including pale phenotypes (Hajdukiewicz et al., 1997; Rogalski et al., 2006; Stoppel
et al., 2011; Manavski et al., 2015; Chen et al., 2016). Furthermore, many mutants affected in chloroplast
translation are more sensitive to cold than wt (Rogalski et al., 2008; Fleischmann et al., 2011; Gu et al.,
2014; Wang et al., 2016; Zhang et al., 2016; Paieri et al., 2018; Pulido et al., 2018). This includes mutants
of chloroplast RNA binding proteins that affect translation and that show cold-dependent bleaching of the
center of the Arabidopsis rosette (Wang et al., 2016), which is strikingly similar to the phenotype of *cp31a*
mutants in the cold. CP31A binds to several mRNAs coding for ribosomal proteins (e.g. *rps18*), thus being
potentially important for the production of the translational machinery in the cold. Taken together, we

hypothesize that CP31A becomes important for the expression of a combination of its target mRNAs in the

cold; their compromised expression in cold-treated *cp31a* mutants leads to loss of chloroplast development

and thus bleaching. Complementation analyses and time-resolved analysis of the accumulation of mRNAs

in *cp31a* mutants during cold treatment could reveal, which mRNAs are key targets for CP31A during cold

acclimation.

Methods

Plant growth

Arabidopsis thaliana Columbia-0, *cp31a-1* T-DNA insertion mutants (Tillich et al., 2009) and AD-deletion

mutants were grown on soil with a 16-h-light/8-h-dark cycle at 23°C in a CLF growth cabinet at 120

$\mu mol \cdot m^{-2} \cdot s^{-1}$. For cold stress application, temperatures were lowered to 8°C when plants reached an age of

two weeks. Cold exposure was then continued for another five weeks to allow the emergence of sufficient

new tissue prior to harvest. For RIP-seq experiments plants were grown on a soil/vermiculite 4:1 mixture at

21°C for 14 days (normal conditions). Cold stress was applied after 13 days at 21°C with a 24 h exposure

to 4°C.

Vector construction and production of transgenic plants

A full length Arabidopsis CP31A cDNA clone was prepared from total Arabidopsis RNA. The genomic

region encoding the acidic domain (+241 to +447 bp from the start codon of the open reading frame) was

deleted as follows: The 5' region of *CP31A* (1-240 bp) and the 3' region of CP31A (448-990 bp) were

amplified with the primers 31A_gateway_for and A31-minusADrev and A31-minusAD_for and

31A_gateway_rev, respectively. These 5' and 3' regions were joined by restriction-based ligation into a

pBluescript vector. The *p35SP:31AΔAD* binary vector for constitutive expression of CP31A was constructed

as follows: The *31AΔAD* fragment was amplified with the primer set (SmaI-31AF01; 31A-SmaIR01) and

ligated into *pJET1.2* cloning vector (Thermo Scientific), and the *Sma*I digested product was integrated into

a *Sma*I site between the *35S* promoter and *Nos* terminator sequences of the binary vector *pGL1* (provided

by Dr. Boris Hedtke, HU Berlin), which also contains the bar expression cassette. For producing the

p31AP:31A vector, the *CP31A* 5' UTR and promoter region (1,694 bp of the 5' region of *CP31A*; 12769646

to 12767953 of the *Arabidopsis thaliana* chromosome 4 sequence; Acc. No. CP002687.1) was cloned with

the specific primer set (CP31A5'up-1694; CP31A-CDS5'_rev). Then, the *CP31A* cDNA was combined with

the promotor by Gibson cloning using the primers (XhoI-31Apro F; Cp31A-CDS5'_rev; CP31A-

ADtest_for; 31A-SmaIR01) and cloned into *pJET1.2*, yielding *pXhoI-31AP:31A-SmaI*. The 35S promoter

was removed and replaced with multi cloning site including *Xho*I and *Sma*I sites, yielding vector *pGL1-*

MCS. XhoI/SmaI digestion products from *pXhoI-31AP:31A-SmaI* were inserted into *Xho*I/*Sma*I sites of

pGL1-MCS, yielding *p31AP:31A* vector. *p31AP:31AΔAD* was constructed in an analogous way. These three

vectors were introduced into the Agrobacterium GV3103 strain independently, and integrated into the

knock-out mutant *cp31a-1* (SALK_109613; Tillich et al., 2009) by the floral-dip method. The T_0 generation

was grown on 15 cm diameter plastic plates filled with soil (mixture of horticulture soil and vermiculite).

1/2000 diluted BASTA was sprayed on the plants several times in order to select for BASTA resistant plants.

BASTA resistant plants were transferred to plastic pots filled with soil and cultivated at 23°C, 8-h-dark/16-

h-light at a light intensity of 120-130 µmol·m^{-2}·s^{-1}.

Immunoblot analysis

506 Total protein (15 µg) were extracted from fully developed leaves grown at normal conditions and

507 immunological analysis were carried out and antibodies were used as previously reported (Kupsch et al.,

508 2012). Immunoblot analysis for the RIP-seq experiments was performed in the same way with protein

509 samples taken from the input, supernatant and pellet fraction of the coimmunoprecipitation.

RIP-Seq analysis

For RIP-seq experiments *Arabidopsis thaliana* Columbia-0 and AD-deletion mutants under the native

promoter (#5-7-1, #5-13-2) were harvested after 14 days and flash-frozen in liquid nitrogen. The flash-

frozen plant material was grinded under liquid nitrogen to a homogeneous powder. Between 250 to 350 mg

plant material was suspended in 3 ml RIP-seq lysis buffer containing formaldehyde (50 mM HEPES-KOH

pH 8.0, 200 mM KCl, 5 mM MgCl$_2$, 5 mM CaCl$_2$, 0.5% Nonidet P-40, 0.5% Sodium deoxycholate, 1x

cOmplete™, EDTA-free Protease Inhibitor Cocktail (Roche), 100 U RiboLock RNase Inhibitor

(ThermoFisher Scientific), 1% formaldehyde) per 1 g plant material. Crosslinking was performed for 10

min under rotation at room temperature and stopped by the addition of 125 mM glycine and a 5 min

incubation. The plant extract was centrifuged for 10 min at 20,000xg and 4°C to remove unsoluble plant

material and flash-frozen in liquid nitrogen.

For the coimmunoprecipitation (CoIP) 8 µl affinity-purified anti-CP31A antibody (Kupsch et al., 2012) was

bound to 50 µl Dynabeads ProteinG (Invitrogen) under rotation (15 rpm). The plant extract was thawed and

centrifuged for 10 min at 20,000xg and 4°C. 350 µl of the supernatant were diluted 1:1 with CoIP buffer

(150 mM NaCl, 20 mM Tris-HCl pH 7.5, 2 mM MgCl$_2$, 0.5% Nonidet P-40, 5µg/ml Aprotinin) and

incubated with the antibody-coated magnetic beads for 75 min at 15 rpm and 4°C. An aliquot of the

antibody-bead solution was taken to serve as the input control. The beads were washed four times in CoIP

buffer and resuspended in Proteinase K buffer (100 mM NaCl, 10 mM Tris-HCl pH 7.0, 1 mM EDTA, 0.5%

SDS).

The crosslink was reversed with 0.1 mg/ml Proteinase K (ThermoFisher Scientific) at 50°C for 1 h and RNA
was extracted from input and pellet fractions using TRIzol and RNA Clean and Concentrator Columns
(Zymo Research) according to the manufacturer's instructions.

Library preparation was performed with the NEBNext® Multiplex Small RNA Library Prep Set for Illumina
(New England BioLabs) according to the manufacturer's instructions with few deviations. Library
preparation was performed for half the volume. Additionally, a 5' adaptor including unique molecular
identifiers (UMI) was used (5´-
rGrUrUrCrArGrArGrUrUrCrUrArCrArGrUrCrCrGrArCrGrArUrCGATCNNNNNNNN-3´). PCR
amplification was performed using the KAPA HiFi HotStart ReadyMix with the cycling protocol for library
amplification for Illumina platforms and an annealing temperature of 62°C. The PCR amplified cDNA
construct was purified using the GeneJET PCR Purification Kit (ThermoFisher Scientific) according to the
manufacturer's instructions and then separated on 6% polyacrylamide gel. Library fragments between
160bp and 190bp were extracted from the gel according to the NEBNext® protocol and subjected to Illumina
sequencing.

For the bioinformatic identification of the CP31A binding events the following steps were performed:

1) Mapping of the Illumina reads. First, reads were split depending on its barcode using
fastx_barcode_splitter (version 11sep2008) to identify to which sample they correspond, next the UMI
barcode was extracted for each read using *umi_tools,* later, reads were trimmed using Trimmomatic version
0.36 with the next parameters: *LEADING:3 TRAILING:3 SLIDINGWINDOW:4:15 MINLEN:10* in order to
eliminate Illumina adapters and bad quality sequence regions. The obtained trimmed reads were aligned to
the *Arabidopsis thaliana* genomes (Araport11) using STAR version 2.5.2a with parameter: *--
outFilterMultimapNmax 2 --outSAMtype BAM SortedByCoordinate --alignIntronMax 3000 --
outFilterIntronMotifs RemoveNoncanonical --outSAMstrandField intronMotif --outSAMattrIHstart 0*. At
this point, duplicated reads were removed from the BAM file using *umi_tools*. Only reads that mapped to
the chloroplast genome with a length bigger than 35 bp were used for downstream analysis.

2) Identification of candidate BSs in each condition studied. Reads mapped in rRNA were eliminated.
Remaining reads were used to identify BSs using CSAR (version 1.31.0; parameters: *b=10, w=70,*
nper=100, test="Ratio") for each strand and sample/replicate independently. Only those regions that
showed a significant enrichment in mapped reads when comparing IP versus corresponding input samples
at a false discovery rate (FDR) < 0.05 were kept (Muino et al., 2011), which led to the discovery of 371
candidate BSs across all samples. In order to obtain a common set of candidate BSs for all samples, the list
of BSs for each sample/replicate were merged using the software *mergeBed* from the package *bedtools*
(v2.26.0). This resulted in 75 reference BSs.

3) Quantification of BSs. The program *featurecounts* (v1.6.0; parameters: *-s 1 -M*) was used to count the
number of reads strand-specifically mapping to the common set of candidate BSs obtained from the previous
step. DESeq2 (v1.14.1) was used with defaults parameters except *fitType="local"* to obtain normalized
number of mapped read in each sample, next it was used to calculate enrichment and significance of number
of the candidate BSs comparing IP vs control (2 biological replicates). The normalized number of reads
were used to calculate Pearson correlation coefficients.

SSMART analysis of CP31A RIP-seq data

SSMART analysis was carried out as described (Munteanu et al., 2018). We used all significant BSs
identified in wild type plants grown in cold conditions as input, as in this sequence set most binding events
were detected. The input sequences of the BSs were scored using their adjusted p-value according to the
actual binding site analysis of the RIP-seq data.

RNA extraction and editing analysis

Total RNA was extracted from fully developed leafs (0.1 g) powdered in liquid nitrogen using Trizol
(Thermo Fisher) according to the manufacturer's protocol. DNA was removed from RNA samples by three
consecutive DNase I treatments and Phenol/Chloroform/Isoamyl alcohol extractions. DNA removal was
checked by PCR with the chloroplast-specific *rps14* primer set. cDNA synthesis was performed with 2 µg
RNA using SuperScript III first strand synthesis system (Invitrogen) according to the manufacturer's
protocol. A one-to-ten dilution of cDNA was used as a template for amplifying ten cDNAs encompassing
16 editing sites (**Fig.** 6A) on nine plastid genes using primer sets previously reported (Tillich et al., 2009).
RT-PCR products were purified with MinElute PCR Purification Kit (Qiagen). Bulk RT-PCR products were
cloned into the barcoded cloning vector and sequenced using the Illumina MiSeq machine. Sequenced data
were sorted according to barcoding of vectors and analyzed with the CLC Genomics Workbench (CLC bio).
Selected RT-PCR products were directly analyzed by Sanger Sequencing and analyzed using the Geneious
software.

RNA gel blot analysis

Total RNA (4 µg) was fractioned on 1.2% agarose gels containing 1.2% formaldehyde, blotted and
hybridized with radiolabeled RNA probes produced by T7 in vitro transcription from PCR products
generated with primer combinations described in Suppl. Tab. 2.

Author contributions and Acknowledgement

We wish to acknowledge expert technical support by Irina Passow. This work was supported by
grants of the Deutsche Forschungsgemeinschaft to CSL (A02 of TRR175), DL (C05 of TRR175)

and TR (FOR 2092). Support of MKL by the IRI Life Sciences and the "Frauenförderung des
Instituts für Biologie" of the Humboldt University is gratefully acknowledged. AO was supported
by JSPS Overseas Research Fellowships. We thank Deserah Strand and Hannes Ruwe for critical
discussion of the data.

AO, MKL, and TR performed the research and analyzed the data. JMM performed the
computational analysis of the RNA seq data. BL performed the analysis of the CP31A BS
consensus. DL, UO and CSL designed the research and analyzed the data. CSL wrote the paper
with contributions of all authors to the final version.

Table 1: Summary of target sites of CP31A across all samples analyzed in RIP-Seq experiments

	Unique CP31A binding sites[1]		Unique CP31A target genes[2]		Length of potential target genes (bp)[3]		Expected unique binding sites[4]	
	sense	antisense	Sense	antisense	sense	antisense	sense	antisense
NDH complex	12	0	6	0	13002	13002	3	3
Photosynthesis (Except *ndh* genes)	6	3	5	3	23644	23644	6	6
Other metabolism	4	0	2	0	4580	4580	1	1
Ribosome	0	5	0	5	22778	22778	6	6
RNA polymerase	3	1	3	1	11170	11170	3	3
tRNA	2	1	2	1	10196	10196	2	2
Other	2	9	2	4	24177	24177	6	6
All BS	29	19	20	14			27	27
Total BSs in intergenic regions	27		NA		44931		21	
Total BSs	75		34				75	

602
603 [1] this includes binding sites identified in all samples; rRNAs were excluded from calling binding sites
604 [2] gene annotations according to Araport11
605 [3] sum of the length of all genes within a category
606 [4] number of target sites expected in a gene class if binding is random and abundance of transcripts is
607 considered equal between all genes (i.e. depends only on the combined length of all genes within a class)
608

609 **References**
610

Bardet, A.F., He, Q., Zeitlinger, J., and Stark, A. (2011). A computational pipeline for comparative ChIP-seq
analyses. Nat Protoc **7,** 45-61.
Barkan, A. (2011). Expression of plastid genes: organelle-specific elaborations on a prokaryotic scaffold. Plant
Physiol **155,** 1520-1532.

Barkan, A., and Small, I. (2014). Pentatricopeptide repeat proteins in plants. Annu Rev Plant Biol **65,** 415-442.
Benegiamo, G., Mure, L.S., Erikson, G., Le, H.D., Moriggi, E., Brown, S.A., and Panda, S. (2018). The RNA-
Binding Protein NONO Coordinates Hepatic Adaptation to Feeding. Cell Metab **27,** 404-418 e407.
Bentolila, S., Oh, J., Hanson, M.R., and Bukowski, R. (2013). Comprehensive high-resolution analysis of the role
of an Arabidopsis gene family in RNA editing. PLoS Genet **9,** e1003584.
Biehl, A., Richly, E., Noutsos, C., Salamini, F., and Leister, D. (2005). Analysis of 101 nuclear transcriptomes
reveals 23 distinct regulons and their relationship to metabolism, chromosomal gene distribution and co-
ordination of nuclear and plastid gene expression. Gene **344,** 33-41.
Bollenbach, T.J., Schuster, G., Portnoy, V., and Stern, D. (2007). Processing, degradation, and polyadenylation of
chloroplast transcripts. In Cell and Molecular Biology of Plastids, R. Bock, ed (Berlin, Heidelberg:
Springer), pp. 175-211.
Castandet, B., Hotto, A.M., Strickler, S.R., and Stern, D.B. (2016). ChloroSeq, an Optimized Chloroplast RNA-
Seq Bioinformatic Pipeline, Reveals Remodeling of the Organellar Transcriptome Under Heat Stress. G3
(Bethesda) **6,** 2817-2827.
Chen, F., Dong, G., Wu, L., Wang, F., Yang, X., Ma, X., Wang, H., Wu, J., Zhang, Y., Wang, H., Qian, Q., and
Yu, Y. (2016). A Nucleus-Encoded Chloroplast Protein YL1 Is Involved in Chloroplast Development and
Efficient Biogenesis of Chloroplast ATP Synthase in Rice. Sci Rep **6,** 32295.
Chen, H.C., and Stern, D.B. (1991). Specific ribonuclease activities in spinach chloroplasts promote mRNA
maturation and degradation. J Biol Chem **266,** 24205-24211.
Cho, W.K., Geimer, S., and Meurer, J. (2009). Cluster analysis and comparison of various chloroplast
transcriptomes and genes in Arabidopsis thaliana. DNA Res **16,** 31-44.
Choquet, Y., Wostrikoff, K., Rimbault, B., Zito, F., Girard-Bascou, J., Drapier, D., and Wollman, F.A. (2001).
Assembly-controlled regulation of chloroplast gene translation. Biochem Soc Trans **29,** 421-426.
Cloutier, A., Shkreta, L., Toutant, J., Durand, M., Thibault, P., and Chabot, B. (2018). hnRNP A1/A2 and
Sam68 collaborate with SRSF10 to control the alternative splicing response to oxaliplatin-mediated DNA
damage. Sci Rep **8,** 2206.
Danon, A., and Mayfield, A.P.Y. (1991). Light regulated translational activators: identification of chloroplast gene
specific mRNA binding proteins. EMBO J. **10,** 3993-4001.
Daros, J.A., and Flores, R. (2002). A chloroplast protein binds a viroid RNA in vivo and facilitates its
hammerhead-mediated self-cleavage. Embo J **21,** 749-759.
Deng, X.-W., and Gruissem, W. (1987). Control of plastid gene expression during development: the limited role of
transcriptional regulation. Cell **49,** 379-387.
Deng, X.W., Tonkyn, J.C., Peter, G.F., Thornber, J.P., and Gruissem, W. (1989). Post-transcriptional control of
plastid mRNA accumulation during adaptation of chloroplasts to different light quality environments. Plant
Cell **1,** 645-654.
Eberhard, S., Drapier, D., and Wollman, F. (2002). Searching limiting steps in the expression of chloroplast-
encoded proteins: relations between gene copy number, transcription, transcript abundance and translation
rate in the chloroplast of *Chlamydomonas reinhardtii*. Plant J. **31,** 149-160.
Favory, J.J., Kobayshi, M., Tanaka, K., Peltier, G., Kreis, M., Valay, J.G., and Lerbs-Mache, S. (2005).
Specific function of a plastid sigma factor for ndhF gene transcription. Nucleic Acids Res **33,** 5991-5999.
Fleischmann, T.T., Scharff, L.B., Alkatib, S., Hasdorf, S., Schottler, M.A., and Bock, R. (2011). Nonessential
plastid-encoded ribosomal proteins in tobacco: a developmental role for plastid translation and implications
for reductive genome evolution. Plant Cell **23,** 3137-3155.
Fukuda, A., Shimada, M., Nakadai, T., Nishimura, K., and Hisatake, K. (2013). Heterogeneous nuclear
ribonucleoprotein R cooperates with mediator to facilitate transcription reinitiation on the c-Fos gene. PLoS
One **8,** e72496.
Gerber, A.P., Herschlag, D., and Brown, P.O. (2004). Extensive association of functionally and cytotopically
related mRNAs with Puf family RNA-binding proteins in yeast. PLoS Biol **2,** E79.
Germain, A., Kim, S.H., Gutierrez, R., and Stern, D.B. (2012). Ribonuclease II preserves chloroplast RNA
homeostasis by increasing mRNA decay rates, and cooperates with polynucleotide phosphorylase in 3' end
maturation. Plant J **72,** 960-971.
Germain, A., Hotto, A.M., Barkan, A., and Stern, D.B. (2013). RNA processing and decay in plastids. Wiley
Interdiscip Rev RNA **4,** 295-316.
Geuens, T., Bouhy, D., and Timmerman, V. (2016). The hnRNP family: insights into their role in health and
disease. Hum Genet **135,** 851-867.
Gotic, I., Omidi, S., Fleury-Olela, F., Molina, N., Naef, F., and Schibler, U. (2016). Temperature regulates
splicing efficiency of the cold-inducible RNA-binding protein gene Cirbp. Genes Dev **30,** 2005-2017.

Grimmer, J., Rodiger, A., Hoehenwarter, W., Helm, S., and Baginsky, S. (2014). The RNA-binding protein
RNP29 is an unusual Toc159 transport substrate. Front Plant Sci **5,** 258.
Gu, L., Xu, T., Lee, K., Lee, K., and Kang, H. (2014). A chloroplast-localized DEAD-box RNA helicase AtRH3 is
essential for intron splicing and plays an important role in the growth and stress response in Arabidopsis
thaliana. Plant Physiol Biochem **82,** 309-318.
Hajdukiewicz, P.T., Allison, L.A., and Maliga, P. (1997). The two RNA polymerases encoded by the nuclear and
the plastid compartments transcribe distinct groups of genes in tobacco plastids. Embo J **16,** 4041-4048.
Hashimoto, M., Endo, T., Peltier, G., Tasaka, M., and Shikanai, T. (2003). A nucleus-encoded factor, CRR2, is
essential for the expression of chloroplast ndhB in Arabidopsis. Plant J **36,** 541-549.
Hayes, R., Kudla, J., Schuster, G., Gabay, L., Maliga, P., and Gruissem, W. (1996). Chloroplast mRNA 3'-end
processing by a high molecular weight protein complex is regulated by nuclear encoded RNA binding
proteins. Embo J **15,** 1132-1141.
Hirose, T., and Sugiura, M. (2001). Involvement of a site-specific *trans*-acting factor and a common RNA-binding
protein in the editing of chloroplast mRNAs: development of a chloroplast in vitro RNA editing system.
Embo J **20,** 1144-1152.
Hobor, F., Dallmann, A., Ball, N.J., Cicchini, C., Battistelli, C., Ogrodowicz, R.W., Christodoulou, E., Martin,
S.R., Castello, A., Tripodi, M., Taylor, I.A., and Ramos, A. (2018). A cryptic RNA-binding domain
mediates Syncrip recognition and exosomal partitioning of miRNA targets. Nat Commun **9,** 831.
Hotto, A.M., Huston, Z.E., and Stern, D.B. (2010). Overexpression of a natural chloroplast-encoded antisense
RNA in tobacco destabilizes 5S rRNA and retards plant growth. BMC Plant Biol **10,** 213.
Hotto, A.M., Schmitz, R.J., Fei, Z., Ecker, J.R., and Stern, D.B. (2011). Unexpected Diversity of Chloroplast
Noncoding RNAs as Revealed by Deep Sequencing of the Arabidopsis Transcriptome. G3 (Bethesda) **1,**
559-570.
Ivanov, A.G., Rosso, D., Savitch, L.V., Stachula, P., Rosembert, M., Oquist, G., Hurry, V., and Huner, N.P.
(2012). Implications of alternative electron sinks in increased resistance of PSII and PSI photochemistry to
high light stress in cold-acclimated Arabidopsis thaliana. Photosynth Res **113,** 191-206.
Keene, J.D. (2007). RNA regulons: coordination of post-transcriptional events. Nat Rev Genet **8,** 533-543.
Klaff, P., and Gruissem, W. (1991). Changes in chloroplast mRNA stability during leaf development. Plant Cell **3,**
517-529.
Klein, R.R. (1991). Regulation of light-induced chloroplast transcription and translation in eight-day- old dark-
grown barley seedlings. Plant Physiol. **97,** 335-342.
Kupsch, C., Ruwe, H., Gusewski, S., Tillich, M., Small, I., and Schmitz-Linneweber, C. (2012). Arabidopsis
Chloroplast RNA Binding Proteins CP31A and CP29A Associate with Large Transcript Pools and Confer
Cold Stress Tolerance by Influencing Multiple Chloroplast RNA Processing Steps. Plant Cell **10,** 4266-
4280.
Li, X.G., Duan, W., Meng, Q.W., Zou, Q., and Zhao, S.J. (2004). The function of chloroplastic NAD(P)H
dehydrogenase in tobacco during chilling stress under low irradiance. Plant Cell Physiol **45,** 103-108.
Li, Y.Q., and Sugiura, M. (1990). Three distinct ribonucleoproteins from tobacco chloroplasts: each contains a
unique amino terminal acidic domain and two ribonucleoprotein consensus motifs. Embo J **9,** 3059-3066.
Li, Y.Q., and Sugiura, M. (1991). Nucleic acid-binding specificities of tobacco chloroplast ribonucleoproteins.
Nucleic Acids Res **19,** 2893-2896.
Lin, B.C., Defenbaugh, D.A., and Casey, J.L. (2010). Multimerization of hepatitis delta antigen is a critical
determinant of RNA binding specificity. J Virol **84,** 1406-1413.
Lisitsky, I., and Schuster, G. (1995). Phosphorylation of a chloroplast RNA-binding protein changes its affinity to
RNA. Nucl. Acids Res. **23,** 2506-2511.
Lisitsky, I., Liveanu, V., and Schuster, G. (1995). RNA-Binding Characteristics of a Ribonucleoprotein from
Spinach Chloroplast. Plant Physiol. **107,** 933-941.
Loza-Tavera, H., Vargas-Suarez, M., Diaz-Mireles, E., Torres-Marquez, M.E., Gonzalez de la Vara, L.E.,
Moreno-Sanchez, R., and Gruissem, W. (2006). Phosphorylation of the spinach chloroplast 24 kDa RNA-
binding protein (24RNP) increases its binding to petD and psbA 3' untranslated regions. Biochimie **88,**
1217-1228.
Lunde, B.M., Moore, C., and Varani, G. (2007). RNA-binding proteins: modular design for efficient function. Nat
Rev Mol Cell Biol **8,** 479-490.
Lykke-Andersen, J., and Wagner, E. (2005). Recruitment and activation of mRNA decay enzymes by two ARE-
mediated decay activation domains in the proteins TTP and BRF-1. Genes Dev **19,** 351-361.
Malay, A.D., Watanabe, M., Heddle, J.G., and Tame, J.R. (2011). Crystal structure of unliganded TRAP:
implications for dynamic allostery. Biochem J **434,** 427-434.

Manavski, N., Schmid, L.M., and Meurer, J. (2018). RNA-stabilization factors in chloroplasts of vascular plants.
Essays Biochem **62,** 51-64.
Manavski, N., Torabi, S., Lezhneva, L., Arif, M.A., Frank, W., and Meurer, J. (2015). HIGH CHLOROPHYLL
FLUORESCENCE145 Binds to and Stabilizes the psaA 5' UTR via a Newly Defined Repeat Motif in
Embryophyta. Plant Cell **27,** 2600-2615.
Mentzen, W.I., and Wurtele, E.S. (2008). Regulon organization of Arabidopsis. BMC Plant Biol **8,** 99.
Muino, J.M., Kaufmann, K., van Ham, R.C., Angenent, G.C., and Krajewski, P. (2011). ChIP-seq Analysis in R
(CSAR): An R package for the statistical detection of protein-bound genomic regions. Plant Methods **7,** 11.
Munteanu, A., Mukherjee, N., and Ohler, U. (2018). SSMART: sequence-structure motif identification for RNA-
binding proteins. Bioinformatics **34,** 3990-3998.
Nakamura, T., Ohta, M., Sugiura, M., and Sugita, M. (1999). Chloroplast ribonucleoproteins are associated with
both mRNAs and intron-containing precursor tRNAs. FEBS Lett **460,** 437-441.
Nakamura, T., Ohta, M., Sugiura, M., and Sugita, M. (2001). Chloroplast ribonucleoproteins function as a
stabilizing factor of ribosome-free mRNAs in the stroma. J Biol Chem **276,** 147-152.
**Nakamura, T., Furuhashi, Y., Hasegawa, K., Hashimoto, H., Watanabe, K., Obokata, J., Sugita, M., and
Sugiura, M.** (2003). Array-based analysis on tobacco plastid transcripts: preparation of a genomic
microarray containing all genes and all intergenic regions. Plant Cell Physiol **44,** 861-867.
Nickelsen, J., and Link, G. (1993). The 54 kDa RNA-binding protein from mustard chloroplasts mediates
endonucleolytic transcript 3' end formation in vitro. Plant J **3,** 537-544.
**Paieri, F., Tadini, L., Manavski, N., Kleine, T., Ferrari, R., Morandini, P., Pesaresi, P., Meurer, J., and
Leister, D.** (2018). The DEAD-box RNA Helicase RH50 Is a 23S-4.5S rRNA Maturation Factor that
Functionally Overlaps with the Plastid Signaling Factor GUN1. Plant Physiol **176,** 634-648.
Pfannschmidt, T. (2003). Chloroplast redox signals: how photosynthesis controls its own genes. Trends Plant Sci **8,**
33-41.
Pulido, P., Zagari, N., Manavski, N., Gawronski, P., Matthes, A., Scharff, L.B., Meurer, J., and Leister, D.
(2018). CHLOROPLAST RIBOSOME ASSOCIATED Supports Translation under Stress and Interacts with
the Ribosomal 30S Subunit. Plant Physiol **177,** 1539-1554.
Quaresma, A.J., Oyama, S., Jr., Barbosa, J.A., and Kobarg, J. (2006). The acidic domain of hnRNPQ (NSAP1)
has structural similarity to Barstar and binds to Apobec1. Biochemical and biophysical research
communications **350,** 288-297.
**Reiland, S., Messerli, G., Baerenfaller, K., Gerrits, B., Endler, A., Grossmann, J., Gruissem, W., and
Baginsky, S.** (2009). Large-scale Arabidopsis phosphoproteome profiling reveals novel chloroplast kinase
substrates and phosphorylation networks. Plant Physiol **150,** 889-903.
Rogalski, M., Ruf, S., and Bock, R. (2006). Tobacco plastid ribosomal protein S18 is essential for cell survival.
Nucleic Acids Res **34,** 4537-4545.
Rogalski, M., Schottler, M.A., Thiele, W., Schulze, W.X., and Bock, R. (2008). Rpl33, a nonessential plastid-
encoded ribosomal protein in tobacco, is required under cold stress conditions. Plant Cell **20,** 2221-2237.
Rousseau, S., Morrice, N., Peggie, M., Campbell, D.G., Gaestel, M., and Cohen, P. (2002). Inhibition of
SAPK2a/p38 prevents hnRNP A0 phosphorylation by MAPKAP-K2 and its interaction with cytokine
mRNAs. EMBO J **21,** 6505-6514.
Ruwe, H., Kupsch, C., Teubner, M., and Schmitz-Linneweber, C. (2011). The RNA-recognition motif in
chloroplasts. J Plant Physiol **168,** 1361-1371.
**Schonberg, A., Bergner, E., Helm, S., Agne, B., Dunschede, B., Schunemann, D., Schutkowski, M., and
Baginsky, S.** (2014). The peptide microarray "ChloroPhos1.0" identifies new phosphorylation targets of
plastid casein kinase II (pCKII) in Arabidopsis thaliana. PLoS One **9,** e108344.
Schuster, G., and Gruissem, W. (1991). Chloroplast mRNA 3' end processing requires a nuclear-encoded RNA-
binding protein. EMBO J. **10,** 1493-1502.
Schuster, G., Lisitsky, I., and Klaff, P. (1999). Polyadenylation and degradation of mRNA in the chloroplast. Plant
Physiol **120,** 937-944.
Selinger, D.W., Saxena, R.M., Cheung, K.J., Church, G.M., and Rosenow, C. (2003). Global RNA half-life
analysis in Escherichia coli reveals positional patterns of transcript degradation. Genome Res **13,** 216-223.
Sharwood, R.E., Hotto, A.M., Bollenbach, T.J., and Stern, D.B. (2011). Overaccumulation of the chloroplast
antisense RNA AS5 is correlated with decreased abundance of 5S rRNA in vivo and inefficient 5S rRNA
maturation in vitro. RNA **17,** 230-243.
Stern, D.B., Jones, H., and Gruissem, W. (1989). Function of plastid mRNA 3'inverted repeats. RNA stabilization
and gene specific protein binding. J. Biol. Chem. **264,** 18742-18750.

Stoppel, R., Lezhneva, L., Schwenkert, S., Torabi, S., Felder, S., Meierhoff, K., Westhoff, P., and Meurer, J.
(2011). Recruitment of a ribosomal release factor for light- and stress-dependent regulation of petB
transcript stability in Arabidopsis chloroplasts. Plant Cell **23,** 2680-2695.
Teubner, M., Fuss, J., Kuhn, K., Krause, K., and Schmitz-Linneweber, C. (2017). The RNA recognition motif
protein CP33A is a global ligand of chloroplast mRNAs and is essential for plastid biogenesis and plant
development. Plant J **89,** 472-485.
Tillich, M., Hardel, S.L., Kupsch, C., Armbruster, U., Delannoy, E., Gualberto, J.M., Lehwark, P., Leister, D.,
Small, I.D., and Schmitz-Linneweber, C. (2009). Chloroplast ribonucleoprotein CP31A is required for
editing and stability of specific chloroplast mRNAs. Proc Natl Acad Sci U S A **106,** 6002-6007.
Townley-Tilson, W.H., Pendergrass, S.A., Marzluff, W.F., and Whitfield, M.L. (2006). Genome-wide analysis
of mRNAs bound to the histone stem-loop binding protein. RNA **12,** 1853-1867.
Tsunoyama, Y., Ishizaki, Y., Morikawa, K., Kobori, M., Nakahira, Y., Takeba, G., Toyoshima, Y., and Shiina,
T. (2004). Blue light-induced transcription of plastid-encoded psbD gene is mediated by a nuclear-encoded
transcription initiation factor, AtSig5. Proc Natl Acad Sci U S A **101,** 3304-3309.
Udy, D.B., Belcher, S., Williams-Carrier, R., Gualberto, J.M., and Barkan, A. (2012). Effects of reduced
chloroplast gene copy number on chloroplast gene expression in maize. Plant Physiol **160,** 1420-1431.
Wang, B.C., Wang, H.X., Feng, J.X., Meng, D.Z., Qu, L.J., and Zhu, Y.X. (2006). Post-translational
modifications, but not transcriptional regulation, of major chloroplast RNA-binding proteins are related to
Arabidopsis seedling development. Proteomics **6,** 2555-2563.
Wang, S., Bai, G., Wang, S., Yang, L., Yang, F., Wang, Y., Zhu, J.K., and Hua, J. (2016). Chloroplast RNA-
Binding Protein RBD1 Promotes Chilling Tolerance through 23S rRNA Processing in Arabidopsis. PLoS
Genet **12,** e1006027.
Yamori, W., Sakata, N., Suzuki, Y., Shikanai, T., and Makino, A. (2011). Cyclic electron flow around
photosystem I via chloroplast NAD(P)H dehydrogenase (NDH) complex performs a significant
physiological role during photosynthesis and plant growth at low temperature in rice. Plant J **68,** 966-976.
Ye, L., and Sugiura, M. (1992). Domains required for nucleic acid binding activities in chloroplast
ribonucleoproteins. Nucleic Acids Res **20,** 6275-6279.
Zhang, J., Yuan, H., Yang, Y., Fish, T., Lyi, S.M., Thannhauser, T.W., Zhang, L., and Li, L. (2016). Plastid
ribosomal protein S5 is involved in photosynthesis, plant development, and cold stress tolerance in
Arabidopsis. J Exp Bot **67,** 2731-2744.
Zoschke, R., and Bock, R. (2018). Chloroplast Translation: Structural and Functional Organization, Operational
Control, and Regulation. Plant Cell **30,** 745-770.