1
2   **Title:**
3   Single-dimensional human revealed stochastic preferences for multi-dimensional economic
4   choice options
5
6   **Authors:**
7   Konstantin Volkmann*, Alexandre Pastor-Bernier*, Arkadiusz Stasiak, Fabian Grabenhorst
8   and Wolfram Schultz
9
10  * These authors contributed equally to the study.
11
12  Department of Physiology
13  Development and Neuroscience
14  University of Cambridge
15  Cambridge CB2 3DY
16  United Kingdom
17
**Corresponding author**:
Wolfram Schultz
Department of Physiology, Development and Neuroscience
University of Cambridge
Cambridge CB2 3DY
United Kingdom
Email: ws234direct@gmail.com
25
**Email addresses of all authors:**
Konstantin Volkmann: konstantin.volkmann@googlemail.com
Alexandre Pastor-Bernier: pastor.bernier@gmail.com
Arkadiusz Stasiak: as863@cam.ac.uk
Fabian Grabenhorst: fabian.grabenhorst@googlemail.com
Wolfram Schultz: ws234direct@gmail.com
32
33

**Abstract**

Realistic, everyday rewards contain multiple components. An apple has taste and size.
However, we choose in single dimensions, simply preferring some apples to others. How can
such single-dimensional relationships refer to multi-dimensional choice options? Here, we
investigated stochastic choices of two-component milkshakes. The revealed preferences were
intuitively graphed as indifference curves that indicated how much of one component was
traded-in for obtaining one unit of the other component without a change in preference, thus
defining the orderly integration of multiple components into single-dimensional estimates.
Options on higher indifference curves were preferred to those on lower curves. The
systematic, non-overlapping curves satisfied leave-one-out tests, followed decoder
predictions and correlated with Becker-DeGroot-Marschak auction-like bids. These single-
dimensional estimates of multi-component options complied with rigorous concepts of
Revealed Preference Theory and encourage formal investigations of normal, irrational and
pathological decisions and their neural signals.

**Keywords**: Bundles, decision-making, reward, stochastic choice, psychophysics

## Introduction (960 words with refs)

We like apples, in particular if they are sweet. We can state our preference in words, but this may not be accurate because of poor introspection, faulty memory or erroneous report. It would be more accurate to observe our choice between different apples by which we reveal our preference at that moment. But how do we make that choice? When choosing between apples, we may prefer a sweeter one even if it is a bit smaller; we would trade-in some size for more sweetness. As the trade-off illustrates, our preference among apples does not come from any component alone but from their combination. Every reward or economic good constitutes a bundle with multiple components, attributes or dimensions, and thus is formally a vector. The bundle components may be integral parts of a good, like the sweetness and size of the apple, or consist of distinct entities, like the steak and vegetable of a meal. Importantly, each component contributes to the choice of the bundle. Without considering the multi-component nature of choice options, we can only study gross choices or exchanges, like between an apple and a pear (not really a choice for an apple lover), or between a movie ticket and a meal (not good when hungry). Thus, to understand realistic, fine-grained choices, we should consider that choice options have multiple components.

In contrast to the multi-dimensionality of realistic, vectorial choice options, revealed preferences and subjective reward value ('utility') are single-dimensional. When faced with two options, a rational decision maker can only prefer one option or its alternative or be indifferent to the options (completeness axiom; see Mas-Colell et al. 1995). With repeated, stochastic choices, preferences are revealed by choice probability (McFadden 2004); the probability of choosing one option over its alternative varies in a graded, scalar manner. Correspondingly, the utility of one choice option can only be higher, lower or equal to that of its alternative. Further, neural signals reflecting choice options vary only along a single dimension at any given moment and thus are also scalar. Hence the question: how can single-dimensional revealed preferences, utility and neural signals emerge from vectorial, multi-component choice options? In particular, can we test the issue empirically using well worked-out theoretical concepts that should reduce possible confounds?

The issue of vectorial-to-scalar transformation can be formally approached by considering that the same scalar measure may arise from oppositely varying vector components. I may trade-in a larger, less sweet apple for a smaller but sweeter one without loss or gain in utility (I do not want to lose anything, nor is it easy to gain anything). Thus, the increase in sweetness is compensated by the decrease in size. This trade-off is captured by Revealed Preference Theory, whose two-dimensional indifference curves (IC) provide a convenient graphic formalism (Fisher 1892; Samuelson 1937; 1938). Equally revealed preferred but differently composed multi-component bundles are plotted on one and the same IC. The continuous nature of these ICs is an important indicator for the systematic trade-off that indicates the well-ordered integration of the relative values of multiple bundle components into single-dimensional preference and utility. For these reasons, ICs are a mainstay of economic theory and textbooks (Laidler & Estrin 1989; Kreps 1990; Varian 1992; Mas-Colell et al. 1995; Perloff 2009).

Previous research investigated multi-component choice options without referring to the IC concept. The tests include the quality and price of television sets (Simonson 1989), comfort and fuel consumption of cars (Simonson 1989), payoff amount and probability (Tversky 1969; Soltani et al. 2012; Levy et al. 2012), shape and color of cards (Fellows 2016), artificial objects and appendages (Pelletier & Fellows 2019), various food components (Suzuki et al. 2017; DiFeliceantonio et al. 2018; Harris et al. 2018), hats and shoes (Thurstone 1931), pastries and payoffs (MacCrimmon & Toda 1969), and monetary token from two accounts (Choi et al. 2007; Kurtz-David at al. 2019). More formal studies used ICs to represent hypothetical outcomes, such as hats and shoes (Thurstone 1931), pencils or pastries and payoff (MacCrimmon & Toda 1969), and monetary token (Choi et al. 2007; Kurtz-David et al. 2019). The formal IC scheme is also useful for explaining inconsistencies

of economic choices (Rieskamp et al. 2006; Chung et al. 2017; Gluth et al. 2017). However,
what is missing from past work are empirical designs that apply the IC formalisms of
Revealed Preference Theory to controlled laboratory situations for a wide range of studies.
In building on previous research, the purpose of the present study was to empirically
estimate formal ICs of revealed preference relationships that take all components properly
into account. The design used the rigorous concepts of Revealed Preference Theory and
should be suitable for testing further aspects of revealed preferences in normal humans, for
detecting fine-grained 'irrational' anomalies in normal and brain-damaged humans who fail
to properly consider all components of choice options, for studying underlying normal and
pathological brain processes, and for comparing the data with behavioural choices and high-
precision single-neuron recordings in animals. These goals imposed substantial constraints,
including specific event timing and the use of types and amounts of payouts that are
comparable to the immediate and tangible liquid and food rewards of animal studies (Kagel et
al. 1975; Pastor-Bernier et al. 2017). The repeated trials necessary for statistical data analysis
of human imaging (and animal neurophysiology) required us to study stochastic preferences
(McFadden 2004; Stott 2006). Sitting in front of a computer monitor, our participants viewed
two discrete choice options (rather than verbal information) that contained the same two fatty
and sugary milkshakes in independently varying quantities. We varied bundle composition
and psychophysically estimated indifference points (IP) at which an original and a changed
bundle were chosen with equal probability.
126
127  **Results**
128
**Basic properties of indifference curves.** Participants repeatedly chose between two
composite visual stimuli that represented two-component bundles (see Methods notion 1; Fig.
1a). Each bundle stimulus consisted of two colored vertical rectangles; blue indicated
component A, a low-sugar high-fat milkshake; red indicated component B, a high-sugar low-
fat milkshake; a vertically positioned bar inside each rectangle indicated the amount of each
milkshake (higher was more). We assessed choices between a preset Reference bundle and a
Variable bundle whose component A was set to a specific test amount and whose component
B varied pseudorandomly across a wide range (Fig. 1b). The choice probabilities in all 24
participants followed the change of component B in an orderly fashion, thus revealing
monotonic preference relationships (notions 2 - 4; Fig. 1c). We derived the indifference point
(IP; $P = 0.5$ for each bundle) from six repetitions using the probit choice function (Eqs. 1, 2);
each IP required 42 choices. Thus, a two-dimensional IP indicated the amounts of the two
components of the Variable Bundle that was as much revealed preferred, and had the same
utility, as the constant Reference Bundle. To obtain a series of IPs, we repeated this
procedure by setting increasing amounts of component A in the Variable Bundle while
keeping the Reference Bundle constant. All IPs of such a series were as much revealed
preferred as the constant Reference Bundle and, by transitivity, equally revealed preferred to
each other.
We obtained single ICs from such a series of five equally revealed preferred IPs by
hyperbolic fitting (Eqs. 3, 3a) (notions 5 and 6; Fig. 1d, e). Such an IC defined the trade-off
between the two bundle components at equal preference: it indicated how much of
component B a participant gave up for obtaining one unit of component A. The continuous
ICs were asymmetric between the x-axis and y-axis, indicating different subjective weighting
of the two milkshakes; the convex IC curvature suggests that lower amounts of both
milkshakes together were as much preferred as higher amounts of singular milkshakes
(possibly reflecting gradually flattening, concave utility functions and/or complementarity
between high-sugar and high-fat components).
We estimated three sets of 5 IPs with their respective, hyperbolically fitted ICs by
presetting the Reference Bundle to three different amounts of component B (2.0 ml, 5.0 ml or
8.0 ml; component A being always 0.0 ml). Fig. 2a (inset) shows the psychophysical

assessment of an example IP. Although the ICs varied in slope and curvature between
participants (for two example participants, see Fig. 2a, b; for all participants, see Fig. 3), the
ICs of bundles with larger reward amounts were located farther away from the origin, thus
reflecting higher revealed preference (notion 7). The three ICs were well ordered and failed
to overlap. Only the 95% confidence intervals overlapped partly in four of the 24 participants
(17%). Closer inspection of Figs. 2a, b and 3 shows that some bundles on higher ICs had a
smaller amount of component B than bundles on lower ICs, indicating revealed preference
despite partial physical non-dominance (notion 8; involving overcompensation by higher
amount of component A).
Taken together, the continuous nature of the ICs with their multiple IPs indicated the
orderly and systematic emergence of single-dimensional utility and stochastic preference
from differently composed vectorial bundles, in compliance with the basic concepts.
171
**IC coefficients.** The shape of ICs reflects the trade-off between the components and can
be quantified by slope and curvature coefficients of hyperbolic fits (Eq. 3a). The global IC
slope, between y-axis and x-axis intercepts, was measured as ratio of the two regression
coefficients $\beta_2 / \beta_1$ in Eq. 3; it indicated how much the participant was globally willing to
give up in order to obtain one unit of the other component; the measure reflected the relative
utility (currency) of the two bundle components. The IC slopes steeper than -45° indicate that
a participant gave up a higher amount of component B (high-sugar, y-axis) for a smaller
amount of component A (high-fat, x-axis), thus indicating higher subjective value of fatty
than sugary milkshake. The IC slope was -71° ± 6.5° (mean ± standard error of the mean,
SEM; range -45° to -76°; $N = 24$ participants; Fig. 2c). The higher valuation of high-fat
component A over high-sugar component B amounted to a factor of 3:1 in 18 of the 24
participants (75%). The predominantly asymmetric trade-off between the two milkshake
components documents that each component contributed to bundle preference in its own
distinct way.
The IC curvature showed substantial convexity in 18 of the 24 participants (75%), as
indicated by $\beta_3$ coefficients from the hyperbolic fit (Eq. 3) that were significantly larger than
0.0 (8.89; mean ± 5.9 SEM; $p<0.05$, t-test; Fig. 2d). For graphically assessing IC curvature,
we measured the distance between the IC center and a straight line connecting equally
revealed preferred bundles at the x-and y-intercepts, in units of ml on the y-axis (Fig. 2e).
This distance ranged from 0.09 ml (quasi-linear IC) to 3.76 ml (most convex IC) (mean of
1.28 ml ± 0.19 SEM; Fig. 2f). The distribution of the IC distance was overall similar to that
of the $\beta_3$ curvature coefficient from Eq. 3. The two highest histogram bars in Fig. 2f show
data from six participants with rather similar, considerably convex IC, and from six other
participants with rather similar but quasi-linear IC. Thus, the coefficients confirmed
numerically the well-ordered nature of the representation of revealed preferences by the ICs.
197
**Control for other choice variables.** To test whether the choices reflected the
components of the bundles rather than other, unrelated factors, we performed a logistic
regression analysis separately for each of the 24 participants, using the following regressors:
amount of each bundle component, reaction time, Reference Bundle position on participant's
monitor, and previous trial choice (Eq. 5). The standardized beta (β) coefficients and p-values
were assessed for each participant and then averaged across all 24 participants; they
demonstrated that the choice of the Variable Bundle was negatively correlated with the
amount of component B in the Reference Bundle (RefB: $\beta = -0.43 \pm 0.16$, mean ± SEM; $P =$
$0.020 \pm 0.005$) (component A was constant 0.0 ml, see Methods) and positively correlated
with both components A and B in the Variable Bundle (VarA: $\beta = 0.67 \pm 0.16$; $P = 0.009 \pm$
$0.004$; VarB: $\beta = 0.94 \pm 0.33$; $P = 0.012 \pm 0.009$) (Fig. 2g). The beta (β) coefficients for these
three variables differed significantly from 0 ($P = 0.012$, $P = 0.00088$ and $P = 0.00028$,
respectively; one-sample t-test), confirming the validity of the β's. Thus, more frequent
choice of the Variable Bundle correlated with lower amounts in the Reference Bundle and

with higher amounts of either component in the Variable Bundle. The result suggests that
both bundle components, rather than a single component alone, were important for the
observed preference relationships. The remaining variables, including reaction time, position
of Reference Bundle on the monitor and previous trial choice, failed to significantly account
for current choice of the Variable Bundle ($P = 0.754 – 0.988 \pm 0.003 – 0.290$). Thus, the
revealed preference relationships concerned the bundles with their two components rather
than other task factors.
To assess potential consumption effects, we searched for signs of satiety. We followed
choice probability across the total test duration in each of the 24 participants. We selected
two bundles that were situated above and below the IP, respectively. These two bundles
contained high-sugar low-fat (above IP) and low-sugar high-fat (below IP) milkshakes. We
plotted choice probability over six repeated test steps (Extended Data Fig. 1). Choice
probabilities fluctuated only insignificantly, without conspicuous upward or downward trend
(above IP: F (5, 41) = 0.28, $P > 0.05$; below IP: F (5, 41) = 1.53, $P > 0.05$; 1-way repeated
measures Anova with post-hoc Tukey Test). Even at the final, sixth step, choice probability
differed only insignificantly from any other step. Thus, the revealed references did not seem
to be importantly confounded by satiety for neither sugar nor fat within the amounts and
concentrations used in our experiment.
230
**Internal validation of indifference curves.** We performed a leave-one-out analysis to
assess the contribution of individual bundles to the ICs obtained from hyperbolic fits to the
empirically estimated IPs. We removed one IP at a time (except the initial Reference Bundle
at x = 0.0) from the set of five IPs to which a given IC had been fitted, and then refitted the
IC with the remaining four IPs using the same hyperbolic model (see Methods, Eqs. 3, 3a).
This was done separately for each of the three ICs, separately in all 24 participants (total of
288 IPs removed from 72 ICs). We found consistency in the refitted ICs in four measures
(Fig. 4a). First, none of the 72 refitted ICs overlapped with the refitted ICs at different levels
in the same participant, thus demonstrating maintained IC separation despite one left-out IP.
Second, none of the 72 refitted ICs overlapped with the 95% confidence intervals of original
ICs at different levels, confirming IC separation despite one left-out IP. Third, most refitted
ICs (66 of 72 ICs, 92%) fell inside the 95% confidence intervals of the original ICs, and the
remainder curves (6 of 72 ICs, 8%) showed only some portions outside the 95% confidence
intervals of the original ICs, thus refuting possible overweighted influence of individual IPs
on ICs. Fourth, numeric comparisons showed only insignificant deviations between refitted
ICs and the IPs that had been left out when refitting the curves (vertical distance of $0.05 \pm$
$0.13$ ml in all 24 participants; mean $\pm$ standard error of the mean, SEM; $N = 336$; $P = 0.98$
against normal distribution; t-test) (Fig. 4b, c), confirming absence of overweighted IP
influence on ICs. These four results suggest that the hyperbolically fitted ICs captured the IPs
consistently and provided valid representations of the revealed preferences.
We used a linear support vector machine (SVM) decoder as different statistical
procedure to confirm the contribution of each IP to the two-dimensional representation of
revealed preferences by ICs. In each participant, we set a given test bundle to one of the
psychophysically determined IPs (Fig. 5a) and assessed the accuracy with which the decoder
assigned that bundle to its original preference level (each preference level was defined by a
series of empirically estimated IPs but was not a fitted IC). SVM decoding accuracy ranged
largely from 70% to 100% ($P = 2.055$ x $10^{-101}$), although a few lower values were observed
(Table 1 left).
We supplemented the SVM analysis by visualization of decoding using linear
discriminant analysis (LDA). The considerable amount of data necessary for reasonable LDA
required us to pool data from several participants, which violates a basic tenet of economic
theory that prohibits pooling of subjective preferences across individual participants. To
somewhat contain expected inaccuracies, we normalized IPs across participants (z-score
normalization for reward B along the y-axis; reward A had been set experimentally to

identical values on the x-axis) and restricted the analysis to specific subsets of participants.
The LDA confirmed the SVM results in all subsets (Fig. 5); the first linear discriminant
assigned bundles to the three revealed preference levels, as shown by spatial separation of the
three colored groups, with a numeric accuracy of 80-100% ($P = 1.148$ x $10^{-97}$). By contrast,
the second discriminant failed to accurately assign bundles to different positions on same
preference levels, as shown by the mix of the five shapes representing bundle position. These
characteristics were seen in six participants whose fitted ICs showed the highest similarity in
convexity (Fig. 5b, c), in six participants with linear ICs (Fig. 5d, e) and, for comparison, in
all 24 participants (Fig. 5f, g) (for distinction of participants based on IC curvature, see two
highest bars in Fig. 2f). Thus, LDA decoding followed the fundamentals of ICs: preference
between bundles across ICs but indifference along ICs.
276           Taken together, the two decoders confirmed three distinguishable levels of IPs, and
277     LDA confirmed indifference between IPs on same levels. As these IPs constituted the basis
278     for the hyperbolic fitting of the three ICs, the decoder results validated also the fitting
279     procedure and confirmed that the estimated ICs represented well the revealed preferences.
280
**Mechanism-independent validation of indifference curves.** The ordered
representation of revealed preferences by our ICs can be further validated by comparison of
the key features of ICs with subjective values inferred from a different estimation
mechanism. To this end, we used a monetary Becker-DeGroot-Marschak (BDM) bidding task
that is equivalent to a second-price auction and estimates participants' truthful, 'incentive
compatible' bids on a trial-by-trial basis. The property of truthful revelation makes BDM an
indispensable tool of experimental economics and explains its increasing popularity in
neuroscientific studies of human decision making (Plassmann, O'Doherty & Rangel 2007;
Medic et al. 2014).
290           For these comparisons, the same 24 participants made monetary BDM bids (UK pence)
for 15 bundles that had been set, individually in each participant, to five psychophysically
estimated IPs on each of the three preference levels (in ml of each of the two component
milkshakes). Each preference level was defined by IP bundles that were equally revealed
preferred to the initial Reference Bundle with component A set to 0.0 ml (Fig. 6a; the same
15 bundles were used for hyperbolic IC fitting). Although BDM estimates typically derive
from single-shot choice, we obtained averaged BDM estimates for each bundle from 12
repetitions, thus aiming to approach the nature of data collection during stochastic choice
with psychophysical estimation of each IP (42 trials).
299           The BDM bids followed the ordered preference levels, as shown by significant positive
correlation between the bids for same-level bundles (means from all five bundles) and the
three levels (Spearman rank correlation: rho $= 0.60 \pm 0.05$; mean $\pm$ SEM; $N = 24$ participants;
$P < 0.01$; confirmed by one-way Anova with post-hoc Tukey-Kramer test between all
bundles across the three levels: $P < 0.01$) (Fig. 6b; red, green, blue). A random-effects
analysis separately for each of the 24 participants (Eq. 8) demonstrated a relationship of
BDM bids to preference level (PrefLev: $\beta = 0.47 \pm 0.09$, mean across all 24 participants $\pm$
SEM; $P = 0.016 \pm 0.015$; $\beta$-coefficient difference from 0: $P = 0.000026$, one-sample t-test)
and bundle amount (AmBundle: $\beta = 0.15 \pm 0.13$; $P = 0.020 \pm 0.017$; $P = 0.0278$), rather than
trial number (TrialN: $\beta = -0.10 \pm 0.25$; $P = 0.726 \pm 0.354$), previous trial bid (PrevBid: $\beta =$
$0.12 \pm 0.11$; $P = 0.676 \pm 0.427$) or consumption history (Consum: $\beta = 0.12 \pm 0.11$; $P = 0.224$
$\pm 0.185$) (all $\beta$'s were standardized) (Fig. 6c). A specific analysis (Eq. 9) demonstrated that
the BDM bids reflected both bundle components (component A: $\beta = 0.6534 \pm 0.0866$, mean
$\pm$ SEM; $P = 0.0324 \pm 0.0150$; $\beta$ difference from 0: $P = 1.1613\,0884$ x $10^{-7}$, one-sample t-test;
component B: $\beta = 0.6425 \pm 0.0585$, $P = 0.0289 \pm 0.0202$; $\beta$ difference from 0: $P = 1.2770$ x
$10^{-10}$). Thus, the BDM bids followed well the revealed preference levels and took both
bundle components into account.
316           The BDM bids were similar for equally revealed preferred bundles. To comply with the
notion of trade-off between different bundle components, equally revealed preferred bundles

should elicit similar BDM bids despite varying bundle composition. We conducted a one-way
Anova on all BDM bids for bundles positioned on the same preference level, separately for
each level and each participant. The bids for bundles located on the same revealed preference
level failed to differ statistically (low level: $P = 0.54 \pm 0.06$; medium level: $P = 0.69 \pm 0.06$;
high level: $P = 0.78 \pm 0.05$; mean $\pm$ SEM, $N = 24$ participants). Only three participants
showed significantly different bids for bundles on the medium preference level, and one
participant showed different bids for bundles on the high preference level ($P < 0.05$). Thus,
most participants showed similar BDM bids for equally revealed preferred bundles on same
revealed preference levels (while being sensitive to bundle amounts, as shown by increasing
bids across levels). The BDM bids confirmed the equal-preference trade-off between bundle
components independent of the binary choice mechanism used for assessing revealed
preferences.
The BDM bids were internally consistent. We used decoders to test the distinction of
bundle position across but not along preference levels. Using BDM bids, a binary SVM
decoder showed good accuracy of assigning a test bundle to its original preference level in
individual participants (mostly 50-70%; $P = 3.789 \times 10^{-9}$) but failed to distinguish bundles
along same levels (43-51%; $P = 0.1433$) (Table 1 right). We used an LDA, which is able to
decode several levels together, as supplementary test and found good visual assignment of the
test bundles (Fig. 6a) to the three revealed preference levels in our population of 24
participants (first discriminant; numeric accuracy of 88-100%; $P = 9.46 \times 10^{-12}$; Fig. 6d; three
colored symbol groups) but not to different positions on same preference levels (second
discriminant; numeric accuracy of 43-51%; mix of shapes) (note the reservations above when
combining data from multiple participants). Thus, the BDM bids followed well the two-
dimensional scheme of revealed preference represented by the ICs.
The BDM bids matched well the revealed preference ICs in direct comparison. A
stronger mechanism-independent validation of the IC scheme may be achieved by direct
graphic comparison between BDM bids and hyperbolically fitted ICs. To this end, we
estimated isolines that were fit to BDM bids using Eqs. 9, 9a and compared them with ICs
that had been hyperbolically fitted to IPs (Eqs. 3, 3a). The BDM isolines derived from
averaged 12 BDM bids, and the revealed preference ICs derived from fits to IPs estimated
from 42 stochastic choices. We found that the BDM isolines increased for IPs on increasing
preference levels (farther away from the origin), but were similar for IPs on the same
revealed preference level (same distance from origin) (Fig. 7a). The BDM isolines matched
the revealed preference ICs within their 95% confidence intervals (CIs) in every one of the 24
participants (Fig. 7b). Statistical comparisons showed significantly higher CIs of BDM
isolines compared to revealed preference ICs (Fig. 7c; $P < 0.0884 \times 10^{-8}$, Wilcoxon paired
test; $N = 24$ participants). Despite their larger variability, both BDM isoline slope and
curvature coefficients, derived from the respective $\beta_2 / \beta_1$-ratio and $\beta_3$ regression coefficient
in Eq. 9, failed to differ significantly from the respective slope and curvature coefficients of
revealed preference ICs (Eq. 3; Fig. 7d, e; both $P > 0.05$, Wilcoxon test on BDM vs. IC
coefficients paired from each participant; $N = 24$). Thus, despite larger variability, the BDM
bids matched well the revealed preference ICs when assessed in a comparable way.
Taken together, the different estimation mechanism of BDM bidding provided good
validation of the estimated revealed preferences and their graphic representation by ICs.
362
363    ## Discussion
364
This study shows that ICs representing revealed preferences among multi-dimensional choice
options can be reliably estimated in a controlled and unequivocal manner using immediate
consumption of small payouts in repeated trials. The preferences followed critical
assumptions of formal theory that define the emergence of scalar utility and preference
measures (choice probability) from vectorial choice options, in particular the continuous,
graded trade-off between bundle components that transcends physical bundle composition.

These preferences were graphically well represented by the asymmetric and nonlinear ICs.
Lack of overlap between ICs, validation by leave-one-out statistics and accuracy of SVM and
LDA decoding demonstrate proportionate contribution of individual bundles to robust
preference representations by ICs. The BDM bids corresponding to the ICs provided a
mechanism-independent validation of the revealed preferences. These empirical results
demonstrate revealed stochastic preferences in humans in controlled, non-verbal, laboratory
setting. Further, the systematic ICs correspond well to, and thus implement, the abstract
schematic graphs in economic textbooks and decision research that are used for explaining
human choices (Laidler & Estrin 1989; Kreps 1990; Varian 1992; Mas-Colell et al. 1995;
Perloff 2009). In following these concepts, the currently estimated, well-structured ICs
represent orderly revealed preferences that would allow the distinction from irrational
preferences not fully capturing the multi-component nature of choice options, including
lexicographic preferences restricted to single components. These properties make the
currently developed tests suitable for concept-driven studies of rational and irrational
behaviour in normal and brain-damaged humans.

The representation of vectorial, multi-component choice options by single-dimensional
neural signals is an open question in neuroeconomics. Our behavioural tests provide a formal,
concept-driven foundation for investigating such signals in humans. These tests require
repeated trials rather than one-shot tests typical for experimental economics; the multi-trial
nature is well captured by stochastic choice theories that facilitate data interpretation due to
decades of decision research. The immediately consumed, small payouts facilitate
comparisons with animal studies and control for temporal discounting. The tangible payouts
after small sets of trials do not rely on language and should reduce mental ruminations about
future rewards and assure reliable cooperation by the participants, thus reducing interfering
and confounding brain activity. We were particularly interested in estimating whole maps of
well-ordered, fitted ICs derived from multiple IPs that conform to predictive mathematical
functions, rather than testing preferences for a few bundles with limited general validity. As a
result, neuroeconomic work using the demonstrated empirical tests could investigate decision
mechanisms not resolvable by purely behavioural examination. For example, it is unknown
whether the bundle components are combined into a common neural utility estimate or are
coded separately and integrated at a later stage. The ensuing neural decision process might
occur between whole bundles with all components integrated, or involve hierarchical or
parallel component-by-component comparisons. The systematic ICs may also allow to
investigate neural underpinnings of specific theories, such as the switching of attentional
processes between components conceptualized by multialternative decision field theory (Roe
et al. 2001). Finally, our detailed ICs would be helpful for investigating neural mechanisms
underlying inconsistent decision making when choice options are added (independence of
irrelevant alternatives, IIA), beyond the decoy and attraction effects already being addressed
(Chung et al. 2017; Gluth et al. 2017).

Our quantitative, psychophysically controlled preference assessment required repeated
choices (Green & Swets 1966). Such a multi-trial approach is dictated by statistical
requirements of neural studies, corresponds to the standard elicitation of choice functions in
reinforcement learning (Sutton & Barto 1998) and allows comparison with animal studies
requiring similar schedules (Kagel et al. 1975; Pastor-Bernier et al. 2017). In humans and
monkeys, these methods deliver systematically varying choice probabilities rather than
single, all-or-none selection. The visible trial-by-trial variations are assumed to reflect
underlying random processes that make the choice process stochastic, as captured by choice
probability and probabilistic choice functions in random-choice theories (Luce 1959;
McFadden & Richter 1990; McFadden 2004; Stott 2006). We appreciate that our multi-trial
schedule is at odds with the frequently employed, standard, single-shot, deterministic
assessment of ICs in experimental economics (Thurstone 1931; MacCrimmon & Toda 1969),
but we believe that the obtained consistent and robust ICs validate the approach.

Economic choice often involves substantial but imaginary sums of money and consumer items, or random, singular payouts (Simonson 1989; Simonson, Tversky 1993; Rieskamp et al. 2006). By contrast, our payout schedule was tailored to requirements of neural studies and allowed the participants to actually consume the chosen rewards over tens and hundreds of trials (while controlling for satiety effects), as is typically required for animal experiments. These behavioural choices resembled small daily activities, such as consuming snacks and drinks, and were met by the good motivation of our participants. In this way, we obtained well-ordered ICs that provided valid representations of revealed preferences, without requiring imagined items or unreasonably large sums of money.

Empirical investigations of economic choice of multi-component options have revealed several inconsistencies, including reference biases, differences between willingness-to-pay and willingness-to-sell, and violations of independence of irrelevant alternatives such as the similarity, compromise, asymmetric dominance and attraction effects (Tversky 1972; Knetsch & Sinden 1984; Simonson 1989; Tversky & Simonson 1993; Bateman et al. 1997; Rieskamp et al. 2006). These phenomena may well be due to the intrinsically adaptive nature of biological processes (Soltani et al. 2012; Li et al. 2018), rather than invalidating the revealed preference concept itself. We aimed to avoid interference from adaptive processes by designing stable and highly reproducible test conditions in a well-controlled laboratory environment, non-reliance on verbal report, single, uninterrupted test sessions, singular changes of bundle components, constant direction of testing (from top left to bottom right on the revealed preference map), and preventing satiety by limiting total milkshake intake to 200 ml. As a result, our IPs remained stable over successive testing steps. We used the exact same conditions for eliciting BDM bids, which may have facilitated their correspondence to revealed preference ICs. With these testing conditions, we avoided some known compromising factors, which might help to identify more basic factors underlying decision irregularities.

The coefficients of hyperbolic fits to the ICs characterize numerically the representation of revealed preferences. The slope coefficient indicates the relative weighting of the two milkshakes. For example, the amount of equally revealed preferred single milkshakes (graphed at the respective x-axis and y-axis intercepts of the two-dimensional map) was lower for high-fat (component A) than high-sugar (component B) milkshakes in our participants, which was represented by a IC slope steeper than -45 degrees; thus, participants would have revealed preferred high-fat over high-sugar milkshakes if they came in same amounts. Such asymmetric IC slopes are also seen with various bundles in primates (Pastor-Bernier et al. 2017). A key reason for the asymmetry may be the use of a physical amount scale; in addition, the revealed preference among the milkshake bundles depended also on their fat and sugar content, for which there is no simple common physical scale. Thus, the scaling of milkshakes in units of physical amount would explain the IC asymmetry. Further, the slope for identical bundle compositions varied between our participants, which demonstrates an additional subjective component in revealed preferences. Despite these scaling and subjectivity issues, our estimated ICs had well-ordered slopes and failed to overlap.

Further, the convex IC curvature in most of our participants indicated that unproportionately smaller amounts of combined sugar-fat milkshakes (at IC center) were equally revealed preferred as larger amounts of milkshakes containing primarily sugar or fat (positioned closer to IC boundaries near x- and y-intercepts). The BDM isolines confirmed the convex curvature; unproportionately smaller amounts of combined sugar-fat milkshakes (at isoline center) elicited the same bids as larger amounts of milkshakes containing primarily sugar or fat (positioned closer to isoline x- and y- intercepts). A previous study found comparable data for BDM bids; food snacks containing sugar-fat combinations elicited higher bids than snacks with similar calories but derived only from fat or sugar (DiFeliceantonio et al. 2018). However, not all ICs must be convex; bundles with one unattractive component showed concave ICs in primates (Pastor-Bernier et al. 2017). In

addition, IC convexity may be ascribed to concave utility of each bundle component (Perloff
2009); more of the same component (closer to IC x- or y-intercepts) has lower marginal gain,
and therefore the decision-maker is willing to trade in more of a component of which she has
much for one unit of the other component of which she has little. This may be the reason why
most participants showed convex ICs. Thus, the systematic estimates of the defining slope
and curvature coefficients suggest that the ICs reliably represented well-ordered preferences
for the milkshake bundles.
The estimating mechanism for BDM bids differed substantially from that underlying
revealed preference ICs. Both the auction-like bidding and the mechanics of cursor
movement differed from the choice between two simultaneously presented options. BDM
bidding is incentive compatible, such that erroneous or deceiving bids lead to suboptimal
outcome, as conceptualized by the expected cost of misbehaviour (Lusk et al. 2007); bidders
should state exactly their subjective value to avoid paying an exaggerated price or foregoing a
desired object. With these properties, BDM bidding constituted a well-controlled,
authoritative test for eliciting true subjective values, thus providing a useful validation
mechanism for preferences that were revealed by binary choice. Indeed, the obtained SVM-
and LDA-consistent BDM bids followed the preference scheme of ICs, namely higher bids
for revealed preferred bundles and similar bids with choice indifference despite varying
bundle composition. Most strikingly, hyperbolically fitted BDM isolines closely resembled
the hyperbolically fitted revealed preference ICs, both in terms of graphics and numeric
coefficients. The only notable difference was higher BDM bid variability. Our data align well
with, and extend, the previously noted correspondence between binary choices and BDM bids
for single-component options and bundle choices on paper or via verbal communication
(Roosen et al. 2017). Thus, the BDM estimating mechanism resulted in bids that
corresponded well to the graphic scheme of ICs conceptualized by Revealed Preference
Theory and thus validated our empirical IC assessments.
502
503

## Methods

**Participants.** A total of 24 participants (11 males, 13 females; mean age 25.4 years, range 19-36 years) completed the binary choice task for measuring revealed preferences and performed the BDM control task. None of the participants had diabetes or lactose intolerance, nor did they require specific diets, to avoid medical and cultural interference. All participants had a known appetite for milkshakes and provided written informed consent based on a detailed information sheet. The Local Research Ethics Committee of the Cambridgeshire Health Authority approved the study.

**Implementation of concepts.** Revealed Preference Theory conceptualizes observable preference relationships between multi-component choice options. Decision makers reveal their preference at the moment of choice. We respected general notions of discrete choice models: the employed choice sets were exhaustive and had finite numbers of options (two), and their options were mutually exclusive. The statistical analysis of neural responses in humans and animals requires the use of multiple trials. Therefore, we used a version of Revealed Preference Theory in which preferences are stochastic (McFadden 2004). Thus, we assessed revealed preference from the probability of multiple choices, rather than by traditional single-shot economic tests. The probability of choosing one option over its alternative varies in a graded manner and thus is scalar. As in our parallel behavioural and neurophysiological work on monkeys (Pastor-Bernier et al. 2017; 2019), we implemented these concepts using the following notions (Fig. 1):

(1) Participants made choices between two bundles that contained the same two distinct milkshakes in independently varying scalar amounts (component A, component B, each measured in ml; Fig. 1a). Thus, each choice option consists of a two-dimensional bundle vector. We kept effort cost for obtaining any bundle and budget constant for all choice options for better task control and data interpretation (single keyboard button press).

(2) Revealed stochastic preference is inferred from the measured probability of repeated choice. The probability of choosing a given bundle depends on all components of all bundles present on that trial (as opposed to 'lexicographic' preferences that follow only one component) (Fig. 1b, c).

(3) Equal choice probability for two bundles indicates equal revealed stochastic preference ($P = 0.5$ each bundle). The point at which both bundles are chosen with equal probability (point of subjective equivalence or choice indifference point, IP) is estimated from an S-shaped psychophysical function fitted to the probabilities of repeated choices while varying one component of one bundle and keeping all other components constant (Fig. 1c).

(4) Every bundle has a subjective value ('utility') for the decision maker that depends only on the reward amounts of both milkshakes. A bundle is chosen with a higher probability than any other bundle in the same option set if and only if its utility is higher than in any other bundle in that choice set. In other words, the preference relationship between two bundles is monotone if the higher utility of one bundle implies that it is preferred to its alternative (Mas-Colell et al. 1995). A bundle is chosen with equal probability of $P = 0.5$ against another bundle if the two bundles have the same utility. Notions (2) - (4) are compatible with basic assumptions of stochastic choice theories (Luce 1959; McFadden & Richter 1990; McFadden 2004; Stott 2006) (although we analyzed the choices with the probit model rather than McFadden's logit model because of its less restrictive assumptions; see below).

(5) Each bundle is graphically displayed at the intersection of the x-coordinate (component A) and y-coordinate (component B) of a two-dimensional graph (Fig. 1d). Equally revealed preferred bundles are graphically represented as two-dimensional IPs and align on an indifference curve (IC) (Fig. 1e). Multiple IPs aligned on a single, continuous IC reflect the orderly and systematic emergence of scalar utility and choice probability from the vectorial bundle components.

(6) Bundles can be equally revealed preferred despite variation in bundle composition,
such that some amount of one component is given up in order to gain one unit of the other
component without change in preference (marginal rate of substitution, MRS). This trade-off
is graphically characterized by two parameters: (i) the IC slope, which reflects the relative
utility (currency) of the two bundle components and can be variable and asymmetric between
x-axis and y-axis; (ii) the curvature, which captures any slope change between IC center and
IC periphery (Fig. 1d).
(7) Bundles with larger amounts of both components are revealed preferred to bundles
with smaller amounts (as long as the 'value function' for each component does not decrease
with amount, indicating that 'more is better') (Fig. 1e). Any bundle that lies above an IC
(farther away from the origin) is revealed preferred to any bundle that lies on that IC, any
bundle that lies on an IC is revealed preferred to any bundle that lies below that IC (closer to
the origin), and thus any bundle on a higher IC is revealed preferred to any bundle on a lower
IC.
(8) The preference relationship between two bundles may hold even when one
component of the revealed preferred bundle comes with a smaller amount than the alternative
bundle (physical non-dominance, requiring overcompensation by the other bundle
component) (Fig. 1e, arrows).
575
**Experimental design.** We measured repeated choices between two options, each of which
contained two milkshakes; the liquids constituted rewards, as evidenced by the participants'
voluntary consumption. A more complete and traditional description of Revealed Preference
Theory includes budget constraints (Mas-Colell et al. 1995); however, in this initial empirical
study, we aimed to obtain as straightforward and easily interpretable empirical data as
possible and therefore did not test aspects of budget constraint. In particular, the acquisition
of any chosen bundle always required exactly one single finger movement (button press on a
computer keyboard).
We used quantitative stimuli to represent the two milkshakes and their amounts in each
of the two bundles (Fig. 1a). Each bundle stimulus consisted of two vertically aligned
rectangles. The colour of each rectangle indicated the bundle component. After extensive
piloting with various liquids and liquidized foods, we found milkshakes with a controlled mix
of sugar and fat to give the most reliable behavioural performance. As the milkshakes were
delivered separately, with a 500 ms interval (see below), drinks containing either only sugar
or only fat were deemed to be too unnatural. Thus, in our bundles, component A (top, blue)
consisted of a low-sugar high-fat milkshake (25% whole milk, 75% double cream, no sugar),
and component B (bottom, red) consisted of a high-sugar low-fat milkshake (10% sugar in
skimmed milk). The vertical position of a bar in each rectangle indicated the physical amount
of each component (higher was more). The milkshakes were delivered using a custom-made
syringe pump system with silicone tubing approved for delivery of food stuffs (VWR
International Ltd). The pump was controlled using a National Instruments card (NI-USB-
6009) via the Matlab Data Acquisition Toolbox. The Psychtoolbox in Matlab (Version
R2015b) running on a Dell Windows computer served for stimulus display and recording of
behavioural choices.
In the choice task, each participant revealed her preference in choices between the two
composite stimuli representing the two bundles (Fig. 1a). The two bundle stimuli appeared
simultaneously at pseudorandomly alternating fixed left and right positions on a computer
monitor in front of the participant; each bundle stimulus contained the same two milkshakes
with independently set amounts. Both bundle stimuli appeared after a short, pseudorandomly
varying interval (mean 0.5 s) following an initial central fixation cross. The participant made
a choice between the two bundle stimuli by pressing a single button once (left or right
computer keyboard arrow for corresponding choice of left or right bundle). Reaction time
was defined as the interval between the appearance of the two bundle stimuli and the
participant's key press. The two milkshakes from the chosen bundle were delivered together

with a probability of $P = 0.2$; i. e. every fifth chosen bundle was paid out on average using a
Poisson distribution, and there was no payout of any milkshake on the remaining trials.
Component B was delivered after a constant interval of 500 ms after component A. This
constant delay, rather than simultaneous delivery or pseudo-randomly alternating sequential
liquid delivery, prevented uncontrolled liquid interactions, maintained discriminability of the
liquids and amounted to constant temporal discounting. Thus, the utility for component B
derived not only from this milkshake alone but also from temporal discounting due to longer
delay. Although participants were instructed to not eat or drink up to four hours prior to the
testing, satiety was a concern due to the high sugar and fat content of the milkshakes. We
addressed the issue by the $P = 0.2$ payout schedule, by limiting each payout to maximally
10.0 ml, and by delivering not more than 200 ml of total liquid to each participant on each
session.
We used a standard psychophysical staircase procedure (Green & Swets 1966) to
estimate the IP at which each of the two bundles were chosen with equal probability ($P = 0.5$
each option), revealing equal preference for each option. The procedure required repeated
testing, which was also required for the subsequent neuroimaging experiment with the same
participants (to be reported elsewhere).
We started the psychophysical procedure by setting, in the Reference Bundle,
component A to 0.0 ml and component B to 2.0 ml, 5.0 ml or 8.0 ml (Fig. 1b). We then
varied the alternative Variable Bundle in a systematic fashion; first, we set its component A
one unit higher (mostly 0.5 ml, 1.0 ml or 2.0 ml higher), thus specifying the amount of
component A gained by the participant from the choice; then we selected randomly (without
replacement) one amount from a mean of 7 fixed amounts of component B (multiples of 0.5
ml), spanning the whole, constant range of tested amounts; we repeated the selection until all
7 amounts had been tested once (Fig. 1c). Then we estimated each IP from 6 repetitions using
sigmoid fitting (see Eqs. 1, 2 below), requiring 42 choices per IP. At the IP, the amount of
component B was usually lower in the Variable Bundle compared to the Reference Bundle.
In this way, we assessed at each IP how much of component B the participant was willing to
give up in order to gain one unit of component A, always relative to the constant Reference
Bundle.
We obtained further IPs in choices between the constant Reference Bundle and the
Variable Bundle whose amounts of component A increased step-wise, thus advancing from
top left to bottom right on the two-dimensional x-y indifference map. We are aware that the
unidirectional progression of testing may lead to somewhat different IP estimates than testing
in the opposite direction or in random sequence (Knetsch 1989). However, in this initial
study, we were primarily interested in the systematic assessment of consistent IPs rather than
exploring potential pitfalls.
To obtain three levels of revealed preference, we used three starting amounts of
component B in the Reference Bundle (2.0 ml, 5.0 ml, or 8.0 ml). In total, we estimated 4 IPs
(from 5 test bundles) at each of 3 levels of revealed preference, resulting in a total of 12 IPs,
derived from 504 choices among 84 different option sets in each participant (7
psychophysically tested amounts for each of 12 IPs; 6 repetitions).
652
**Statistical analysis.** In order to estimate the choice IPs numerically, we obtained a
sigmoid fit to the empirically assessed choice frequencies via a general linear regression. To
do so, we used the Matlab function *glmfit* (Matlab R2015b; Mathworks) on a binomial
distribution with a probit link function, which is the inverse of the normal cumulative
distribution function (G). Specifically, the generalized linear regression $y = \beta_0 + \beta_1 B_{var} + \varepsilon$
can be rewritten after applying the link function as:
659
660  $G(y) = \beta_0 + \beta_1 B_{var} + \varepsilon$                                         Eq. 1
661

with y as number of times the subject chose the Variable Bundle in the current block from a
series of six repetitions, $\beta_0$ as offset constant, $\beta_1$ as regression slope coefficient, $B_{var}$ as
reward amount (ml) of component B in the Variable Bundle, and $\epsilon$ as residual error. We
choose the probit model because it assumes that random errors have a multivariate normal
distribution, which makes it attractive as the normal distribution provides a good
approximation to many other distributions. The model does not rely on the assumption of
error independence and is used frequently by econometricians (Razzaghi 2013). By
comparison, the logit model is computationally simpler but has more restrictive assumptions
of error independence. As preliminary data had revealed a similar fit for the logit as for the
probit model we used the probit model due to its less restrictive assumptions. Thus, we
estimated the IPs from the sigmoid fit provided by the probit model, using the following
equation:

Indifference Point = - ($\beta_0$ / $\beta_1$)                                        Eq. 2

with $\beta_0$ and $\beta_1$ as coefficients of the general linear regression (Eq. 1). By using a probit
instead of a logit function, we obtained the coefficients from the output of the probit analysis
(Amemiya 1981).
We obtained single ICs, separately for each individual participant, from a set of
individual IPs by weighted least-mean-square, non-linear regression (as opposed to the probit
regression for estimating each IP). We applied a weight in order to account for within-
participant choice variability; the weight was the inverse of the standard deviation of the
titrated amount of the B-component at the corresponding IP (the IP having been estimated by
the probit regression analysis). We estimated the best-fitting $\beta$ coefficients from least-mean-
square fitting to obtain the equal-revealed-preference IC (which is a utility level) and wrote
the basic hyperbolic equation in our notation as:

IC  = $\beta_0$ + $\beta_1$B + $\beta_2$A + $\beta_3$BA + $\epsilon$                              Eq. 3

with A and B as amounts of components A and B (ml) referring to the x-axis and y-axis,
respectively, $\beta_1$ and $\beta_2$ as slope coefficients of the least-mean-square, non-linear regression,
and $\beta_3$ as curvature coefficients. As IC is a constant, we merged the other constants offset
($\beta_0$) and error ($\epsilon$) into a common final constant k. To draw the ICs, we computed the amount
of component B as a function of component A from the derived equation:

B = k – ($\beta_2$/$\beta_1$)A + $\beta_3$A                                        Eq. 3a

To graphically display a fitted IC (Fig. 2a, b), we plotted the preset amount of component A
on the x-axis, and the computed fitted amount of component B (Eq. 3a) on the y-axis. The
error on the hyperbolic curve was measured as 95% confidence interval. The higher the error
around an IP the less weight was given to this point when the IC was calculated. This model
resulted in good fits in earlier work (Pastor-Bernier et al. 2017). In this way, the IPs of 5
equally revealed preferred but differently composed bundles aligned as a single fitted IC. The
three ICs representing increasing revealed preference levels (low, medium, high) were
located increasingly farther away from the origin (Figs. 1e; 2a, b). The indifference map of 3
x 5 IPs was unique for each participant (Fig. 3).
The IC shape is derived from the hyperbolic fit and is quantified by two coefficients:
slope and curvature. The IC slope coefficient, derived from the ratio of regression slope
coefficients ($\beta_2$ / $\beta_1$), reflects the currency relationship between the components and
describes the participant's preference for component A relative to component B. For
example, an IC slope of - $60^0$ indicates that component A is valued twice as much as the
same ml amount of component B. The curvature coefficient ($\beta_3$) quantifies the constancy in
the trade-off between bundle components. A linear curve (curvature coefficient = 0) indicates

a constant rate of exchange for the bundle components, indicating that the components are
perfect substitutes. A more convex IC (curvature coefficient > 0) indicates a varying rate of
exchange, suggesting that the participant is giving up lesser amounts of component B to
obtain one unit of component A when advancing on the IC from top left to bottom right. For
a more intuitive measure, we quantified the curvature by measuring the largest perpendicular
distance between the IC and the line between the x-axis and y-axis intercepts (Fig. 2e):
721
722 $d = \max (B_{IC} - B_{linearIC})$ Eq. 4
723
with d as maximal perpendicular distance (ml) (whereas $\beta_3$ is a best-fitted, estimated
parameter, and thus less conservative), $B_{IC}$ as amount of component B on the IC (ml), and
$B_{linearIC}$ as amount of component B at the line connecting the x- and y-intercepts (constant
amount of component A, x-axis; ml). This simple curvature measure indicates the effect of
curvature on the trade-off between the two components, in ml of component B.
We used logistic regression on trial-by-trial choices to confirm that the measured
choices were explained by bundle components rather than other factors. In a random-effects
analysis, we fitted a logistic regression to the data from each individual participant and then
averaged the obtained β coefficients and p-values across all participants. We used the
following regression:
734
735 $y = \beta_0 + \beta_1 RefB + \beta_2 VarA + \beta_3 VarB + \beta_4 RT + \beta_5 VarPos + \beta_6 PChoice + \varepsilon$ Eq. 5
736
with y as either 0 or 1 (not-choice or choice of Variable Bundle), A and B as amounts of
bundle components A and B (ml), RefB as amount of component B in the Reference Bundle
(ml), VarA and VarB as amount of components A and B in the Variable Bundle (ml), RT as
reaction time (ms), VarPos as left or right position of the Variable Bundle on the computer
monitor relative to the Reference Bundle (0 or 1), and PChoice as choice in the previous trial
(0 or 1) (Fig. 2g). Each β coefficient was standardized by multiplication with standard
deviation of the respective independent variable and division by standard deviation of y
(dependent variable). A subsequent one-sample t-test against 0 served to assess the
significance of the beta (β) coefficients in the population of the 24 participants.
Satiety may have occurred and could have affected the preferences for the two bundle
components in an uncontrolled manner, even though the bundle rewards were only paid out
on every fifth trial on average and were limited to a total of 200 ml. A prime suspected effect
might have been a differential devaluation between the two bundle components that would
result in changed currency relationship between the two components. Such change between
the two components should be manifested as gradual change in instantaneous choice
probability near the IPs over repeated test steps of 42 trials each (Extended Data Fig. 1). We
calculated the instantaneous choice probability as:
754
755 $y = \Sigma$ (n=1 to 6) (CV / TS) Eq. 6
756
with y as instantaneous probability ($P = 0.0$ to $1.0$), CV as not-choice or choice of Variable
Bundle (0 or 1), and TS as test step (1-6).
We used a leave-one-out analysis to assess the meaningful representation of revealed
preferences by the fitted ICs and to test the accuracy of the hyperbolic IC fit to the IPs. In this
analysis, we removed one IP per IC (but not the initial Reference Bundle set at x = 0), fitted
the curve again with the hyperbolic model and assessed the deviation between the original IC
and the IC with the left-out IP. We defined the deviation as the difference of component B
between the original, left-out IP and the refitted IC (Fig. 4b):
765
766 $d = B_{IP} - B_{refit}$ Eq. 7
767

with d as difference (in ml; y-axis), $B_{IP}$ as amount of component B of the left-out IP (ml), and
$B_{refit}$ as amount of component B on the refitted IC (ml). Thus, a difference of 0 ml suggested
that removal of one IP did not affect the shape of the IC at all, whereas any difference
unequal to 0 ml quantified the violation of this assumption.

**Decoding of preference levels.** To confirm the contribution of each IP to the two-
dimensional representation of revealed preferences, we determined the accuracy (in percent
correct) with which a randomly selected bundle, defined by the amounts of the two
components A and B (in ml), could be assigned to its original revealed preference level as
opposed to any one other level (binary distinction). By definition, each bundle that was
psychophysically estimated to be as much revealed preferred as the Reference bundle
constituted an IP; all bundles to which participants were choice indifferent against the same
Reference Bundle constituted a series of IPs. In our experiment, three different Reference
Bundles defined three preference levels (low, medium, high: component B: 2.0 ml, 5.0 ml or
8.0 ml, respectively; component A was always 0.0 ml; Fig. 5a). The decoder used as inputs
only bundles at the psychophysically estimated IPs (to which an IC was fitted using Eqs. 3,
3a), rather than bundles positioned on the fitted ICs.
Our main test employed a binary support vector machine (SVM) decoder separately on
each individual participant. We used similar methods as previously described for predicting
choice from neuronal activity (Tsutsui et al. 2016). The SVM algorithm considered 5 IP
bundles from each of 2 revealed preference levels (total of 10 IPs that had been assessed 12
times at each position in each participant) (Fig. 5a). Each of the 2 preference levels was
associated with a matrix of 2 columns (containing the x- and y-coordinates of bundle
components A and B, respectively) and 5 rows (containing the 5 bundles). The 5 bundles
were randomly selected (with replacement) from 60 bundles on each level (due to the random
procedure with replacement, some bundles may have entered the algorithm multiple times,
and not all five bundles may have been used for a given analysis). We left out 1 randomly
selected bundle from the 10 bundles, trained the SVM algorithm with the remaining 9
bundles, and assessed whether the SVM decoder assigned the left-out bundle to its original
revealed preference level or to another level. Thus we used 90% of the data for training the
decoder and 10% for testing its classification performance. We repeated this procedure 10
times with the same selected 2 x 5 bundles but with a new randomly selected left-out bundle
and calculated decoder accuracy as percent correct classification in these 10 trials. We
repeated the random selection of the 2 x 5 bundles and the 10-trial accuracy assessment 150
times. For final decoding accuracy, we averaged the percentages from these 150 iterations
(Table 1). We applied this procedure separately to all three possible combinations of two
revealed preference levels (i. e. low and medium, medium and high, low and high). The SVM
was implemented with custom written software in Matlab R2015b (Mathworks) using the
functions *svmtrain* and *svmclassify* with linear kernel (our previous work had shown that use
of nonlinear kernels did not improve decoder performance; Tsutsui et al. 2016).
We supplemented the SVM procedure with binary linear discriminant analysis (LDA)
that provided visualization of the different levels of revealed preference (Fig. 5). We used the
same IPs and the same data matrices as for the SVM analysis (and the same IPs as used for
the hyperbolic fitting of the three indifference curves, ICs). We obtained two variances; the
discriminant 1 eigenvector captured the best separation between the three revealed preference
levels as 'across-level variance' (colors in Fig. 5); the discriminant 2 eigenvector captured the
best within-level separation between five bundles on each of the three preference levels as
'within-level variance' (symbols in Fig. 5). The results indicate visually the discrimination
accuracy on the two axes of the 2-dimensional plots. We also assessed the numeric accuracy
of decoding as percent of correctly assigning a randomly selected bundle to its original
revealed preference level. The decoder used the Matlab functions *fitcdiscr* and *predic* on z-
normalised data from individual participants. For the LDA, our limited data required pooling
from multiple participants. As revealed preferences are private and subjective, and therefore

difficult to compare between individual participants, the LDA results should be considered as
merely supportive and not as stand-alone data.
823
**Mechanism-independent validation of revealed preferences.** To relate the
estimated revealed preferences to inferred utility, we implemented a Becker-DeGroot-
Marschak (BDM) mechanism akin to a second price auction (Becker et al. 1964). The BDM
value bids were then compared with the three levels of revealed preference within each
participant.
In the BDM, the participant received a fresh monetary endowment (20 UK pence) on
each trial. The participant bid for a bundle against a pseudorandomly set computer bid
(retrieved from a normal distribution with replacement). If the participant's bid was higher
than or equal to the computer bid then she received both component rewards of the bundle
and paid an amount equal to the computer bid. If her bid was lower than the computer bid,
she lost the auction, paid nothing and did not receive any bundle reward. The participant was
informed about a win or a loss immediately after placing the bid; when winning the bid, the
participant received the bundle rewards in the same sequence and frequency (every fifth trial
on average) as in the choice task assessing revealed preferences.
We showed each participant single bundles that were randomly selected (without
replacement) from a set of 15 bundles (5 equally revealed preferred bundles at each of 3
levels; the 15 bundles had been used to fit the 3 ICs shown in Fig. 2a, b and Fig. 3). A given
bundle was set to the participant's psychophysically estimated IP (Fig. 6a). We presented
each of the 15 bundles 12 times, resulting in 180 trials in total, and considered the mean of
these bids as the BDM-estimated utility. The participant indicated her bid by moving a cursor
horizontally on the computer monitor with left and right keyboard arrows (Fig. 6a inset). The
BDM bid was registered from the cursor position at 5.0 s after onset of presentation of the
horizontal bidding scale.
We first assessed the basic question whether the monetary bids increased for higher
valued bundles but were similar for equally valued bundles (which constituted IPs), using
Spearman rank correlation analysis confirmed by one-way Anova with post-hoc Tukey-
Kramer test. Then we performed a random-effects analysis with a general linear regression
with a normal (Gaussian) link function on separate data from each participant and averaged
the obtained β coefficients and their p-values across participants. We used the following
regression:
854
855    $y = \beta_0 + \beta_1 \, \text{PrefLev} + \beta_2 \, \text{AmBundle} + \beta_3 \, \text{TrialN} + \beta_4 \, \text{PrevBid} + \beta_5 \, \text{Consum} + \varepsilon$   Eq. 8
856
with y as monetary bid, PrefLev as revealed preference level (low, medium, high),
AmBundle as summed ml amount of components A and B in the currency of component A
$(A + (k - (\beta_2/\beta_1)A + \beta_3 A)$ as in Eq. 3a), TrialN as trial number, PrevBid as BDM bid in
previous trial (UK pence), and Consum as accumulated drinks consumption for component A
and component B up to this point in the experiment (ml) (Fig. 6c). Each β coefficient was
standardized by multiplication with standard deviation of the respective independent variable
and division by standard deviation of y (dependent variable). A subsequent one-sample t-test
against 0 assessed the significance of the beta (β) coefficients in all 24 participants.
To assess the internal consistency of BDM bids, we used binary SVM analysis on bids
from individual participants in analogy to SVM decoding of preference levels. We tested the
same IPs as used for hyperbolic IC fitting (Fig. 6a). Each of the 2 preference levels was
associated with a matrix of 1 column (containing the bids to each bundle) and 5 rows
(containing the 5 bundles). The remainder of the bundle selection, leave-out and repetition
procedure was identical to that used for the SVM decoding of preference levels (see above).
Thus, the SVM decoder for BDM bids assessed the accuracy with which the left-out bundle
belonged to its original revealed preference level. We supplemented the SVM analysis of the
BDM bids with analogous LDA for supportive visualization.

874    Finally, we compared hyperbolically fitted BDM isolines directly with hyperbolically
875 fitted revealed preference ICs (rather than with revealed preference levels just described),
876 separately for each individual participant. This procedure required to present BDM bids on
877 the same scale as revealed preference ICs. To this end, we fitted isolines of same BDM-bids
878 in analogy to fitting same-preference ICs. We fitted a hyperbolic function to the measured
879 mean BDM bids in analogy to Eq. 3:

881 $BDMBid = \beta_0 + \beta_1 B + \beta_2 A + \beta_3 BA + \varepsilon$         Eq. 9

883 with $\beta_1$ and $\beta_2$ as regression slopes, and $\beta_3$ as curvature coefficients, and A and B as amounts
884 of components A and B (ml), respectively. Coefficients $\beta_1$ and $\beta_2$ were standardized by
885 multiplication with standard deviation of components B and A, respectively (independent
886 variables), and division by standard deviation of BDMBid (dependent variable). We obtained
887 separate $\beta$ coefficients from all participants and averaged them and their p-values across
888 participants. A subsequent one-sample t-test used the individual beta ($\beta$) coefficients from all
889 24 participants to test overall significance against 0.
890    To compare BDM bids with ICs, we graphically displayed BDM isolines along which
891 all mean BDM bids were equal. As a BDM isoline is a constant, we merged the constants
892 offset ($\beta_0$) and error ($\varepsilon$) into a common final constant k. To draw the BDM isolines, we
893 computed the amount of component B as a function of component A from the derived
894 equation:

896 $B = k - (\beta_2/\beta_1)A + \beta_3 A$            Eq. 9a

898 To display a three-dimensional map, we graphed colored BDM isoline zones on the z-axis as
899 a function of the amounts of components A (x-axis) and B (y-axis) (Fig. 7a). For a two-
900 dimensional map of BDM isolines, we plotted the preset amount of component A on the x-
901 axis and the amount of component B computed from the isolines (Eq. 9a) on the y-axis (Fig.
902 7b). For comparison, we plotted the revealed-preference ICs on the same two-dimensional
903 map using the same scale. We also compared numerically, separately for each participant,
904 confidence intervals and slope and curvature coefficients between hyperbolically fitted
905 BDMBids (Eq. 9a) and hyperbolically fitted revealed preference ICs (Eq. 3a), using the
906 paired Wilcoxon test (Fig. 7c-e).

908 **Data availability:** Data are available from the authors upon reasonable request. A
909 Reporting Summary for this article is available as a Supplementary Information file.

913
914

## Fig. 1 | Binary choice task and experimental design.

**a**, Bundle stimuli for binary choice. Each bundle contained two components with amounts
between 0.0 and 10.0 ml. Component (Comp) A: low-sugar high-fat milkshake, Comp B:
high-sugar low-fat milkshake. Participants chose between the Reference Bundle and the
Variable Bundle, whose locations on a computer monitor alternated pseudorandomly between
fixed left and right positions.
**b**, Psychophysical test design. In the Reference Bundle, components A and B were set to 0.0
ml and 8.0 ml, respectively; in the Variable bundle, component A was set to a specific test
amount while the amount of component B was psychophysically varied (dashed arrows).
**c**, Psychophysical assessment of two example choice indifference points (IP; choice
probability P = 0.5 each option; yellow and green) in a typical participant. During repeated
trials, the participant chose between the preset Reference Bundle and the Variable Bundle.
Component A of the Variable Bundle was set to specific test amounts (here 0.5 ml and 2.0
ml, respectively). Component B of the Variable Bundle varied between seven randomly
selected amounts of component B. IPs were estimated from choice probabilities, p (Bundle 1)
and p (Bundle 2), respectively, using the probit model (Eqs. 1, 2).
**d**, Schematic indifference curve (IC), fitted by a hyperbola (Eqs. 3, 3a) to all equally revealed
preferred but differently composed bundles. Yellow and green dots show IPs relative to the
bundle marked by blue dot, as estimated in B, C.
**e**, Schematic map of three ICs. Increasing distance from origin represents higher utility; all
bundles on higher ICs are revealed preferred to all bundles on lower ICs. Heavy dots denote a
preference relationship between two bundles with oppositely varying physical amounts of
component A (top-IC bundle is revealed preferred to mid-IC bundle despite lower physical
amount of component A; partial physical non-dominance).
938
939
940

942
943

**Fig. 2 | Empirical indifference curves (IC) representing revealed preferences.**
**a**, Typical convex ICs from an example participant, as seen in 18 of the 24 participants.
Component A was a low-sugar high-fat milkshake; component B was a high-sugar low-fat
milkshake. Solid lines show hyperbolically fitted ICs, dotted lines show 95% confidence
intervals of fits. Dots show bundles that are equally preferred on the same IC (IPs). Inset:
psychophysical assessment of indifference point (IP) marked on highest IC (test points in
blue, IP estimated by probit regression in red).
**b**, Typical linear ICs from another example participant, as seen in 6 of the 24 participants.
**c, d**, Distributions of slope and curvature, respectively, of hyperbolically fitted ICs from all
24 participants (coefficients $\beta_2 / \beta_1$ and $\beta_3$ in Eq. 3, respectively). N = number of
participants.
**E**, Scheme of intuitive numeric assessment of IC curvature: maximal vertical distance (ml of
component B on y-axis) between fitted IC (curve) and a straight line connecting the x-axis
and y-axis intercepts. A distance of > 0.0 ml indicates convexity, whereas a 0.0 ml distance
indicates perfect linearity.
**f**. Distribution of convex curvature, as measured using the scheme shown in E. The two
peaks indicate six participants each with similarly near-linear ICs and similarly convex ICs,
respectively.
**g**, Specificity of bundle choice, as opposed to unrelated parameters. Bar graph showing
standardized beta (β) regression coefficients for choice of Variable Bundle over Reference
Bundle (Eq. 5), derived from each individual participant and averaged across all 24
participants. RefB, component B in Reference Bundle; VarA and VarB, components A and B
in Variable Bundle; RT, reaction time; VarPos, left-right position of Variable Bundle
stimulus; PChoice, choice in previous trial. Error bars show standard error of the mean
(SEM). $*P \leq 0.02$.

969
970

972
973
974 **Fig. 3 | Empirical indifference curves (IC) from all 24 participants.** Note the
975 fanning out of the confidence intervals towards the bottom right in each graph (thin lines),
976 which likely reflects the progression of choice testing: the Reference Bundle was kept
977 constant at the y-axis intercept (x = 0), whereas testing with the Variable Bundle progressed
978 from top left to bottom right. Same conventions as for Fig. 2.
979
980

982
983
**Fig. 4 | Leave-one-out validation of estimated indifference curves (ICs).**
**a**, Graphic assessment with an example participant: hyperbolically refitted ICs with one left-
out indifference point (IP) (solid lines), plotted together with 95% confidence intervals of the
original hyperbolically fitted ICs (dotted lines). The refitting resulted in 4 new ICs (partly
overlapping) at each of three levels. None of the refitted IPs fell outside the original
confidence intervals.
**b**, Scheme of numeric assessment: distance in ml (in ml on y-axis, red) between the left-out
IP on the original IC (heavy black dot on black curve) and the refitted IC (green).
**c**, Histogram of distance between refitted ICs and left-out IPs across all subjects. Skewness
was -0.12, suggesting rather symmetric distribution around the mean (modeled in red).
994
995

997
998

**Fig. 5 | Visualization of bundle decoding using Linear Discriminant Analysis**
**(LDA).**
**a**, Schematics of bundle decoding at psychophysically estimated points of equal revealed
preference (indifference points, IPs, plotted along the dotted lines). Following the notions of
revealed preference, LDA should show accurate decoding across the three preference levels
(green, blue, red) but not along each level.
**b**, LDA bundle distinction between three levels of revealed preference (81-100% binary
numeric decoding accuracy between two levels; first discriminant) but not along same
preference levels (second discriminant) in N = 80 bundles from six participants (6P) with
similar convex ICs (same convention applies to all panels). Bundles on the three preference
levels are colored blue, green and red according to distance from origin, red being highest.
The five bundles on each preference level are marked from top left to bottom right with 'o',
'*', '+', 'x' and '  ' symbols. Due to the arbitrariness of the scale, numbers are not indicated.
**c**, As **b** but for partly physically non-dominating bundles (one lower component in preferred
bundle than in alternative bundle (97-100% accuracy; first discriminant).
**d**, As **b** but for six participants with linear ICs (90-100% accuracy).
**e**, As **d** but for partly physically non-dominating bundles (96-100% accuracy).
**f**, As **b** but for all 24 participants (18 with convex IC, six with linear ICs) (83-100%
accuracy).
**g**, As **f** but for partly physically non-dominating bundles (97-100% accuracy).
1019
1020

1021

1022
1023
**Fig. 6 | Characteristics of BDM bids for bundles at different revealed**
**preference levels.**
**a**, Schematics of positions of bundles used for eliciting BDM bids at psychophysically
estimated points of equal revealed preference (indifference points, IPs, plotted along the
dotted lines). Following the notions of trade-off and increasing revealed preference, BDM
bids should be similar for equally valued bundles (along the dotted lines) but higher for
revealed preferred bundles (farther away from origin). We tested 5 bundles per level, 3 levels,
12 repetitions, total of 180 bids. Inset: BDM task. Each participant bid for the visually
presented two-component (A, B) bundle by moving the black dot cursor using the leftward
and rightward horizontal arrows on a computer keyboard. Numbers indicate example bids (in
UK pence).
**b**, Mean BDM bids from a typical participant. The bids were rank ordered (Spearman Rho =
0.83, P < 0.001) across bundles positioned on increasing indifference curves (blue, green,
red) and failed to overlap between lowest, medium and highest levels. There was a significant
difference for BDM bids between levels but not within levels (*P < 0.001; two-way Anova
followed by Tukey-Kramer test). Data are shown as mean ± SEM (standard error of the
mean), N = 12 bids per bar.
**c**, Specificity of monetary BDM bids, as opposed to unrelated parameters. Bar graph showing
the standardized beta (β) regression coefficients for BDM bids (Eq. 8), derived from each
individual participant and averaged across all 24 participants. Abbreviations: PrevLev,
revealed preference level (low, medium, high); AmBundle, summed currency-adjusted
amount of both bundle components; TrialN, trial number; PrevBid, BDM bid in previous
trial; Consum, accumulated drinks consumption. Error bars show SEMs. * P ≤ 0.020.
**d,**) Visual decoding by Linear Discriminant Analysis (LDA) of BDM bids for bundles on
different preference levels (first discriminant; 88-100% numeric decoding accuracy; $P = 9.46$
$\times 10^{-12}$) and along same preference levels (second discriminant; 43-51% accuracy) (N = 243
bundles; all 24 participants). The LDA used scalar BDM bids for bundles positioned at IPs
shown in Fig. 6a. Same conventions as for Fig. 5a.

1052
1053

1054
1055

1056
1057

Fig. 7 | Comparison between BDM bids and indifference curves (IC).

**a**, Three-dimensional representation of hyperbolically fitted BDM isolines from a typical participant (mean ± SEM). The BDM bids were made for bundles that had been placed on psychophysically estimated revealed preference indifference points (black dots). The BDM isolines were similar along same levels and increased across revealed preference levels.

**b**, Match between hyperbolically fitted isolines of mean BDM bids (black; Eqs. 9, 9a) and hyperbolically fitted revealed preference ICs (blue, green, red; ± 95% confidence intervals, CI, shaded) from a typical participant. Thus, the BDM isolines fell within the respective CIs of the revealed preference ICs.

**c**, Confidence intervals of fitted BDM isolines (red) and revealed preference ICs (blue) from all 24 participants (averaged along each isoline/IC). The larger 95%CIs of BDM isolines (BDMiso) suggest more variability compared to revealed preference ICs.

**d, e**, Comparison of slope and curvature coefficient estimates, respectively, between BDM isolines (red) and revealed preference ICs (blue; same data as shown in Fig. 2c, d) from all 24 participants.

1075
1076    **Table 1.** Decoding accuracy, as assessed with support vector machine.
1077

| Participant<br>C – convex IC<br>L – linear IC | Bundles | | | BDM bids | | |
|---|---|---|---|---|---|---|
| | Lev1<br>vs.<br>Lev2 | Lev1<br>vs.<br>Lev3 | Lev2<br>vs.<br>Lev3 | Lev1<br>vs.<br>Lev2 | Lev1<br>vs.<br>Lev3 | Lev2<br>vs.<br>Lev3 |
| C1 | 83.9 | 100.0 | 68.8 | 70.6 | 81.0 | 54.4 |
| C2 | 86.8 | 99.8 | 68.7 | 44.0 | 51.0 | 47.4 |
| C3 | 72.7 | 96.0 | 58.7 | 56.6 | 71.3 | 59.3 |
| C4 | 88.0 | 99.5 | 67.6 | 67.7 | 76.3 | 62.5 |
| C5 | 99.1 | 100.0 | 81.4 | 64.8 | 69.9 | 49.8 |
| C6 | 93.7 | 99.5 | 61.9 | 53.5 | 65.9 | 54.0 |
| C7 | 66.2 | 75.5 | 49.5 | 70.0 | 72.2 | 43.6 |
| C8 | 72.9 | 81.8 | 52.6 | 49.8 | 67.2 | 62.6 |
| C9 | 80.0 | 100.0 | 80.0 | 68.4 | 83.8 | 62.1 |
| C10 | 99.6 | 100.0 | 69.8 | 47.0 | 46.6 | 43.7 |
| C11 | 90.2 | 99.5 | 55.4 | 68.9 | 75.7 | 53.1 |
| C12 | 96.8 | 100.0 | 71.2 | 47.3 | 57.9 | 47.1 |
| C12 | 86.1 | 87.7 | 56.7 | 58.7 | 60.9 | 48.2 |
| C14 | 80.0 | 100.0 | 80.0 | 63.2 | 74.3 | 56.8 |
| C15 | 99.8 | 100.0 | 85.5 | 67.8 | 78.6 | 54.8 |
| C16 | 100.0 | 100.0 | 90.0 | 57.8 | 62.5 | 50.4 |
| C17 | 100.0 | 100.0 | 100.0 | 44.8 | 70.9 | 68.4 |
| C18 | 99.0 | 99.9 | 55.3 | 68.1 | 75.3 | 51.7 |
| L19 | 100.0 | 100.0 | 88.6 | 43.5 | 65.8 | 59.6 |
| L20 | 100.0 | 100.0 | 95.9 | 69.2 | 84.9 | 70.1 |
| L21 | 100.0 | 100.0 | 100.0 | 62.2 | 66.0 | 44.4 |
| L22 | 100.0 | 100.0 | 83.6 | 54.8 | 52.5 | 45.9 |
| L23 | 100.0 | 100.0 | 100.0 | 63.2 | 58.4 | 47.8 |
| L24 | 100.0 | 100.0 | 80.0 | 61.4 | 68.5 | 57.7 |

1079
Accuracy is shown in % of numbers of bundles and BDM bids correctly assigned to one of
the two tested revealed preference levels (averages from 150 iterations of classification of 10
pseudorandomly selected bundles / BDM bids from two tested revealed preference levels).
Bundles were tested at psychophysically assessed choice indifference points (IP) on one of
three revealed preference levels (corresponding to the three indifference curves, IC, fitted to
the IPs). The tested bundles had been pseudorandomly selected from 5 unique estimated IPs
on each of two revealed preference levels under study. BDM: Becker-DeGroot-Marschak
auction-like bidding mechanism. Lev1, Lev2, Lev3: revealed preference levels, numbered
according to distance from origin. The 24 participants were labelled as C or L according to
their IPs being fitted best to convex or linear ICs.
1090
1091

## References

Alphonce, R. & Alfnes, F. Eliciting consumer WTP for food characteristics in a developing context: Application of four valuation methods in an African market. *J. Agricult. Econ.* **68**, 123-142 (2017).

Amemiya, T. Qualitative response models: A survey. *J. Econ. Lit.* **19**, 1483-1536 (1981).

Bateman, I., Munro, A., Rhodes, B., Starmer, C. & Sugden, R. A test of the theory of reference-dependent preferences. *Quart. J. Econ.* **112**, 479-505 (1997).

Becker, G. M., DeGroot, M. H. & Marschak, J. Measuring utility by a single–response sequential method. *Behav. Sci.* **9**, 226-232 (1964).

Choi, S., Fisman, R., Gale, D. & Kariv, S. Consistency and heterogeneity of individual behaviour under uncertainty. *Am. Econ. Rev.* **97**, 1921–1938 (2007).

Chung, H.-K., Sjöström, T., Lee, H.-J., Lu, Y.-T., Tsuo, F.-Y., Chen, T.-S., Chang, C.-F., Juan, C.-H., Kuo, W.-J. & Huang, C.-Y. Why Do Irrelevant Alternatives Matter? An fMRI-TMS study of context-dependent preferences. *J. Neurosci.* **37**, 11647–11661 (2017).

DiFeliceantonio, A. G., Coppin, G., Rigoux, L., Thanarajah, S. E., Dagher, A., Tittgemeyer, M. & Small, D. M. Supra-additive effects of combining fat and carbohydrate on food reward. *Cell. Metab.* **28**, 33-44 (2018).

Fellows, L. K. Deciding how to decide: ventromedial frontal lobe damage affects information acquisition in multi-attribute decision making. *Brain* **129**, 944-952 (2006).

Fisher, I. Mathematical Investigations in the theory of value and prices. *Trans. Connecticut Acad.* **9**, 1–124 (1892).

Gluth, S., Hotaling, J. M. & Rieskamp, J. The attraction effect modulates reward prediction errors and intertemporal choices. *J. Neurosci.* **37**, 371-382 (2017).

Green, D. M. & Swets, J. *Signal detection theory and psychophysics*: New York: Wiley, (1966).

Harris, A., Clithero, J. A., Cendri, A. & Hutcherson, C. A. Accounting for Taste: A multi-component neurocomputational model explains the neural dynamics of choices for self and others. *J. Neurosci.* **38**, 7952–7968 (2018).

Kagel, J. H., Battalio, R. C., Rachlin, H., Basmann, R. L., Green, L. & Klemm, W. R. Experimental studies of consumer demand behaviour using laboratory animals. *Econ. Inquiry* **13**, 22-38 (1975).

Knetsch, J. L. The endowment effect and evidence of nonreversible indifference curves. *Am. Econ. Rev.* **79**, 1277-1288 (1989).

Knetsch, J. L. & Sinden, J. A. Willingness to pay and compensation demanded: Experimental evidence of an unexpected disparity in measures of value. *Quart. J. Econ.* **99**, 507-521 (1984).

Kreps, D. M. *A Course in Microeconomic Theory*. Harlow: Pearson Education Ltd (1990).

Kurtz-David, V., Persitz, D., Webb, R. & Levy, D. J. The neural computation of inconsistent choice behaviour. *Nat. Comm.* **10**, 1583 (2019).

Laidler, D. E. W. & Estrin, S. *Introduction to Microeconomics*. New York: Philip Allan (1989).

Levy, I., Belmaker, L., Manson, K., Tymula, A. & Glimcher, P. W. Measuring the subjective value of risky and ambiguous options using experimental economics and functional MRI methods. *J. Vis. Exp.* **67**, e3724 (2012).

Li, V., Michael, E., Balaguera, J., Castañón, S. H. & Summerfield, C. Gain control explains the effect of distraction in human perceptual, cognitive, and economic decision making. *Proc. Natl. Acad. Sci.* **115**, E8825-E8834 (2018).

Luce, R. D. *Individual Choice Behaviour: A Theoretical Analysis*. New York: Wiley (1959).

Lusk, J. L., Alexander, C. & Rousu, M. C. Designing experimental auctions for marketing research: The effect of values, distributions, and mechanisms on incentives for truthful bidding. *Rev. Market Sci.* **5**, 1-32 (2007).

MacCrimmon, K. R. & Toda, M. The experimental determination of indifference curves. *Rev. Econ. Stud.* **36**, 433-451 (1969).

Mas-Colell, A., Whinston, M. D. & Green, J. R. *Microeconomic theory.* New York: Oxford Univ Press (1995).

McFadden, D. & Richter, M. K. *Stochastic rationality and revealed stochastic preference. Preferences, Uncertainty, and Optimality, Essays in Honor of Leo Hurwicz,* Westview Press: Boulder, CO, 161-186 (1990).

McFadden, D. L. Revealed stochastic preference: A synthesis. *Econ. Theory* **26**, 245–264 (2004).

Medic, N., Ziauddeen, H., Vestergaard, M. D., Henning, E., Schultz, W., Farooqi, I. S. & Fletcher, P. C. Dopamine modulates the neural representation of subjective value of food in hungry subjects. *J. Neurosci.* **34**, 16856-16864 (2014).

Pastor-Bernier, A., Plott, C. R. & Schultz W. Monkeys choose as if maximizing utility compatible with basic principles of revealed preference theory. *Proc. Natl. Acad. Sci.* **114**, E1766-E1775 (2017).

Pastor-Bernier, A., Stasiak, A. & Schultz, W. Orbitofrontal signals for two-component choice options follow indifference curves of Revealed Preference Theory. *Nat. Comm.* **10**, 4885 (2019).

Perloff, J. M. *Microeconomics.* Boston: Addison-Wesley (2009).

Pelletier, G., & Fellows, L. K. A critical role for human ventromedial frontal lobe in value comparison of complex objects based on attribute configuration. *J. Neurosci.* **39**, 4124-4132 (2019).

Plassmann, H., O'Doherty, J.,& Rangel, A. Orbitofrontal cortex encodes willingness to pay in everyday economic transactions. *J. Neurosci.* **27**, 9984-9988 (2007).

Razzaghi, M. The Probit Link Function in Generalized Linear Models for Data Mining Applications. *J. Mod. Appl. Stat. Meth.* **12**, Issue 1, Article 19 (2013).

Rieskamp, J., Busemeyer, J. R. & Mellers, B. A. Extending the bounds of rationality: Evidence and theories of preferential choice. *J. Econ. Lit.* **44**, 631-661 (2006).

Roe, R. M., Busemeyer, J. R. & Townsend, J. T. Multialternative decision field theory: a dynamic connectionist model of decision making. *Psychol. Rev.* **108**, 370-392 (2001).

Roosen, J., Marette, S. & Blanchemanche, S. Value elicitation using BDM and a discrete choice mechanism. *Econ. Bull.* **30**, 1554-1563 (2010).

Samuelson, P. A. A note on measurement of utility. *Rev. Econ. Stud.* **4**, 155-161 (1937).

Samuelson, P. A. A note on the pure theory of consumer's behaviour. *Economica* **5**, 61-71 (1938).

Simonson, I. Choice based on reasons: The case of attraction and compromise effects. *J Consum. Res.* **16**, 158-174 (1989).

Soltani, A., De Martino, B. & Camerer, C. A range-normalization model of context-dependent choice: A new model and evidence. *PLOS Comput. Biol.* **8**, e1002607 (2012).

Stott, H. P. Cumulative prospect theory's functional menagerie. *J. Risk Uncert.* **32**, 101-130 (2006).

Sutton, R. S. & Barto, A. G. *Reinforcement Learning: An Introduction.* Cambridge MA: MIT Press (1998).

Suzuki, S., Cross, L. & O'Doherty, J. P. Elucidating the underlying components of food valuation in the human orbitofrontal cortex. *Nat. Neurosci.* **20**, 1780–1786 (2017).

Thurstone, L.L. The indifference function. *J. Soc. Psychol.* **2**, 139-167 (1931).

Tsutsui, K.-I., Grabenhorst, F., Kobayashi, S. & Schultz, W. A dynamic code for economic object valuation in prefrontal cortex neurons. *Nat. Comm.* **7**, 12554 (2016).

Tversky, A. Intransitivity of preferences. *Psychol. Rev.* **76**, 31-48 (1969).

Tversky, A. Elimination by aspects: A theory of choice. *Psychol. Rev.* **79**, 281-99 (1972).

Tversky, A. & Simonson, I. Context-dependent preferences. *Management Sci.* **39**, 1179-1189 (1993).

Varian, H. R. *Microeconomic Analysis.* New York: Norton (1992).

1198
1199 **Acknowledgements**. We thank Charles R. Plott for discussions and conceptual support,
1200 Steve Edgley for help and logistic support, Simone Ferrari-Toniolo for comments on
1201 experimental economics and Francine Ganter-Restrepo for engineering expertise. This work
1202 was supported by the Wellcome Trust (WT 095495, WT 204811).
1203

1204 **Author contributions**
1205 K. V., A. P.-B., F. G. and W.S. designed the research, K. V. performed the experiments, K.
1206 V., A. P.-B. and A.S. analyzed the data, K. V. and W.S. wrote the manuscript.
1207

1208 **Competing Interests**: The authors declare no competing interests.

1209
1210

1211

1212
1213
**Extended Data Fig. 1 | Satiety control.**
**A,** Choice function obtained by sigmoid fit with the probit model. From seven randomly
selected fixed amounts of component B in the Variable Bundle, the two colored dots show
two amounts associated with choice probabilities closest to the indifference point (IP).
**b,** Lack of satiety across typical test duration. Choice probabilities varied insignificantly
above (blue) and below (orange) the IP across six repeated choice steps, using 42 trials at
each step. Data were averaged from all 24 participants. The total duration of the six steps was
$20 \pm 1.38$ minutes (mean $\pm$ SEM).
1222