1 **Full title:** Genetic alteration of ferredoxin NADP[+]-reductase or cysteine desulfurase in
2 piperaquine-resistant *Plasmodium berghei* restores susceptibility to lumefantrine and
3 abolishes the impact of probenecid, verapamil, and cyproheptadine
4
5 **Short title:** Mechanisms of lumefantrine resistance in piperaquine-resistant *Plasmodium*
6 *berghei* parasites
7

8 **Authors**: Fagdéba David Bara[1,2,3], Loise Ndung'u[2], Noah Machuki Onchieku[1], Beatrice Irungu[2],
9 Simplice Damintoti Karou[3], Francis Kimani[4], Damaris Matoke-Muhia[4], Peter Mwitari[2], Gabriel
10 Magoma[1,5], Alexis Nzila[6], Daniel Kiboi[5]#
11

Affiliations:
[1]Department of Molecular Biology and Biotechnology, Pan African University Institute for
Basic Sciences, Technology and Innovation (PAUSTI), Nairobi, Kenya.
[2]Centre for Traditional Medicine and Drug Research, Kenya Medical Research Institute,
Nairobi, Kenya.
[3]School of Food and Biology Technology, Universite du Lome, Lome, Togo.
[4]Centre for Biotechnology Research and Development, Kenya Medical Research Institute,
Nairobi, Kenya.
[5]Department of Biochemistry, Jomo Kenyatta University of Agriculture and Technology
(JKUAT), Nairobi, Kenya.
[6]Department of Life Sciences, King Fahd University of Petroleum and Minerals, Dharam,
Saudi Arabia.
24
25 #**Address correspondence to Daniel Kiboi:** dmuthui@jkuat.ac.ke (DK);
26 muthuikiboi@gmail.com

Abstract

Chemotherapy remains central in the control of malaria; however, resistance has consistently thwarted these efforts. Currently, lumefantrine (LM), and piperaquine (PQ) drugs, are essential components in the mainstay artemisinin-based therapies used for the treatment of malaria globally. Using LM and PQ-resistant *Plasmodium berghei,* we measured the effect of known chemosensitizers: probenecid, verapamil, or cyproheptadine on the activity of LM or PQ. Using PlasmoGEM vectors, we then evaluated the impact of deleting cysteine desulfurase (SufS) or over-expressing Ferredoxin NADP+ reductase (FNR), genes that mediate drug action. Our data showed that, only cyproheptadine at 5mgkg^{-1} restored LM activity by above 65% against the LM-resistant parasites (LM^R) but failed to reinstate PQ activity against the PQ-resistant parasites (PQ^R). Whereas the PQ^R had lost significant susceptibility to LM, the three chemosensitizers; cyproheptadine, probenecid, and verapamil, restored LM potency against the PQ^R by above 70%, 60%, and 55% respectively. We thus focused on LM resistance in PQ^R. Deletion of the SufS or overexpression of the FNR genes in the PQ^R abolished the impact of the chemosensitizers on the LM activity, and restored the susceptibility of the PQ^R parasites to LM. Taken together, we demonstrate the association between SufS or FNR genes with the action of LM and chemosensitizers in PQ^R parasites. There is, however, need to interrogate the impact of the chemosensitizers and the role of SufS or FNR genes on LM action in the human malaria parasite, *Plasmodium falciparum*.

Keywords: Malaria; Lumefantrine, Resistance; Reversal; *Plasmodium berghei*; Ferredoxin NADP+ reductase, cysteine desulfurase

INTRODUCTION

Malaria affects more than one billion people worldwide [1]. Presently, five malaria parasite species infect humans. *Plasmodium falciparum* remains a significant contributor to the global disease burden, with an estimated 228 million cases and 405 000 deaths annually [1]. An enormous proportion of this burden affects children under five years of age and pregnant women. In Africa, over 70% of the population is still at risk of infection by the malaria parasites [1,2]. To date, the use of drugs is central to the control and management of malaria. However, this approach is hampered by the ability of the parasite to rapidly develop resistance against antimalarial drugs. Currently, the artemisinin-based combination therapies (ACTs) are the mainstay drugs for the treatment and management of uncomplicated *P. falciparum* malaria. The ACTs comprise a short-acting artemisinin derivative and a long-acting partner drug [2]. The long-acting partner drugs reduce the remaining parasite biomass after artemisinin clearance and simultaneously protect against reinfection, especially in high transmission settings [3]. Thus, the long-acting drug components within the ACTs are of primary importance in the control of subsequent malaria infection in sub-Saharan Africa.

In many African countries, the long-acting antimalarial drugs: lumefantrine (LM) and piperaquine (PQ) are essential components in the mainstay ACTs; Coartem™ and Artekin™, respectively [4]. Unfortunately, the high transmission of malaria in endemic regions coupled with the long half-life portend intense selection pressure [5], a recipe for the rapid emergence of LM and PQ resistant parasites. It is therefore imperative to understand how parasites may evade LM and PQ action. Polymorphisms in two genes, primarily mediate resistance to the 4-aminoquinolines and chemically related drugs such as PQ. The *chloroquine resistance transporter (crt)*, the fundamental determinant of CQ resistance in *P. falciparum,* which can carry Lys76Thr mutation [6] and *multidrug-resistant 1 (mdr1)* gene that encodes a P-glycoprotein homolog 1 (*Pgh-1*) which modulate resistance to CQ and other quinolines drugs such as AQ and PQ [7,8]. In recent studies, Cys101Phe mutation in *chloroquine resistance transporter* (*crt*) conferred resistance to both PQ and CQ in *P. falciparum in vitro* [9]. Also, using field isolates, a nonsynonymous mutation, Glu415Gly in the exonuclease (PF3D7_1362500), and amplification of Plasmepsin II and III, proteases involved in the heme degradation within the digestive vacuole are linked with PQ resistance [10]. LM, on the other hand, is an aryl alcohol antimalarial drug, chemically related to mefloquine and quinine. Although, like the 4-aminoquinoline, LM is predicted to inhibit heme detoxification, several studies have associated reciprocal resistance between CQ and LM [11], suggesting potentially different mechanisms of resistance and action. To date, many questions regarding how LM exerts its

antimalarial action and how the parasites might evolve to evade such action remain
unanswered.

One archetypical method of studying the mechanisms by which parasites evade drug
action and simultaneously generate new drugs is the use of a chemosensitizer (a drug with
the ability to enhance/restore activity) in association with antimalarial drugs for which the
parasite is resistant [12]. This approach has been utilized previously to generate prophylactic
drugs against malaria parasites [13,14]. Although many chemosensitizers have no intrinsic
antimalarial potency, an antihistamine cyproheptadine can reverse resistance and kill asexual
blood-stage malaria parasite[15]. Chemosensitizers reverse resistance mechanisms by either
enhancing drug uptake or inhibiting the efflux of drugs from the target site [16–18]. Three
chemosensitizers have been extensively used to study the resistance mechanisms and design
next-generation drugs in both malaria and cancer [19–22]. In malaria, verapamil, a calcium (Ca^{+2})
channel blocker, partially reverses CQ resistance in *P. falciparum in vitro* and in clinical
isolates via increasing net CQ within the infected erythrocyte parasites [23]. Probenecid, anion
transporter inhibitor, reverses methotrexate resistance in cancer [24], CQ, and antifolates
resistance in *P. falciparum* [14]. Cyproheptadine and a related first-generation antihistamine;
chlorpheniramine restore CQ sensitivity in CQ resistant *P. yoelii* and *P. falciparum in vitro* [15,25].

Using a model malaria parasite, *Plasmodium berghei* ANKA, we have previously
selected PQ- and LM-resistant parasites[26]. Further analysis of the PQ^R parasites showed that
the parasite had lost significant susceptibility to LM, amodiaquine (AQ), primaquine (PMQ),
and dihydroartemisinin (DHA)[26]. On the other hand, although LM^R parasites retained
sensitivity to PQ, the parasites lost significant susceptibility to DHA, PMQ, and CQ [26,27]. Also,
the resistant parasites acquired significant growth fitness cost probably associated with the
changes within the genomes that affect the essential genes necessary for the growth of the
asexual blood-stage parasites [28]. In the current study, we hypothesized that: i) a combination
of probenecid, verapamil, or cyproheptadine with PQ might restore the efficacy of PQ against
the PQ^R parasites; ii) a combination of probenecid, verapamil, or cyproheptadine with LM
might restore the efficacy of LM against both PQ^R and LM^R parasites; iii) PQ^R and LM^R
parasites are accompanied by differential mRNA expression of a panel of plausible genes that
mediate drug action and transport quinoline and aryl alcohol drugs. These are; *crt*
(PBANKA_1219500)*, mdr1* (PBANKA_1237800)*, multidrug resistance-associated protein 2*
(mrp2) (PBANKA_1443800)*, pantothenate kinase (pank)* (PBANKA_1022600)*, ferredoxin*
NADP$^+$-reductase (fnr) (PBANKA_1122100)*, cysteine desulfurase (SufS)*
(PBANKA_0614300)*, Plasmepsin IV (pmiv)* (PBANKA_1034400)*, Plasmepsin IX (pmix)*
(PBANKA_1014500)*, Plasmepsin X (pmx) or phosphatidylinositol 3-kinase (pi3k)*

(PBANKA_1114900); iv) deletion or overexpression of selected plausible genes would restore
parasite susceptibility to PQ or LM. Here, we provide insights into the mechanisms of LM
resistance and reversal in *Plasmodium berghei,* which are critical for future rational drug
design.

RESULTS

Cyproheptadine selectively restores LM activity against lumefantrine and piperaquine
resistant parasites

To select the appropriate concentrations for the drug combination studies, we first
separately conducted cyproheptadine, LM or PQ activity assays against the resistant (PQ[R]
and LM[R]) and wildtype (WT) parasites. Since the LD_{50} of cyproheptadine is 123mg/kg in
mice[29], we used 30mg/kg as the highest concentration. Surprisingly, 30mg/kg of
cyproheptadine killed 90% of the LM[R] (Fig 1A) and 40% of the PQ[R] parasites (Fig 1B).
However, at 5mg/kg, cyproheptadine had no significant activity against either the WT (4.4%),
LM[R] (0.6%) or PQ[R] parasites (0.7%) (Fig 1A and Fig 1B). Based on these cyproheptadine data,
a concentration of 5mg/kg or 2.5mg/kg was selected for subsequent drug combination
assays. To select the appropriate concentrations of LM or PQ for use with cyproheptadine,
we measured the activity profiles of LM against LM[R] or WT and PQ against PQ[R] parasites. As
expected, 50mg/kg of LM yielded a marginal effect (1.66% parasite killing) on the LM[R]
parasites. Against the WT, 0.375mg/kg of LM showed insignificant parasite killing (17%)
compared to 5mg/kg, which yielded 100% parasite killing (Fig 1A, Table 1). On the other hand,
75mg/kg of PQ killed a meager 10.1% of the PQ[R] parasites as expected (Fig 1B). Based on
these LM and PQ activity results, we selected 100mg/kg, 50mg/kg or 25mg/kg of LM for LM[R]
parasites assays, 0.375mg/kg of LM for WT parasites assays, and 75mg/kg of PQ for PQ[R]
parasites assays, for subsequent drug combination experiments.

We first measured the impact of cyproheptadine on LM activity against the LM[R]
parasites. Here, a combination of cyproheptadine [5mg/kg] and LM [100mg/kg] significantly
restored the LM potency from 23% to 67% (p<0.001) with a surprisingly high standard
deviation (Fig 2A). A lower cyproheptadine concentration [2.5mg/kg] and LM [100mg/kg]
yielded 91% parasite killing with low standard deviation (Fig 2A). Next, we hypothesized that
the restoration of LM potency by cyproheptadine was dependent on the concentration of LM
or cyproheptadine. To dissect this hypothesis, we evaluated cyproheptadine at 2.5 and
5mg/kg with 25 or 50mg/kg of LM. Cyproheptadine [2.5mg/kg] and LM [25mg/kg] yielded
67%, cyproheptadine [2.5mg/kg] and LM [50mg/kg] yielded 60.6%, cyproheptadine [5mg/kg]
and LM [25mg/kg] yielded 61% while cyproheptadine [5mg/kg] and LM [50mg/kg] yielded
68% of parasite clearance. Statistical analysis revealed that the % parasite killing for the

different combinations was not significantly different (p<0.08) (Fig 2B), signifying that the
restoration of LM activity by cyproheptadine was dose-independent. These data also
reaffirmed 2.5mg/kg of cyproheptadine was sufficient low concentration to restore at least
two-thirds of LM efficacy against the LM[R] parasites without obvious reciprocal drug
interaction. We next reasoned that the reversal of LM resistance was selective only in the
resistant parasites. To evaluate the selectivity of cyproheptadine on the resistant parasites,
2.5 and 5mg/kg of cyproheptadine was combined with 0.375mg/kg of LM against the wildtype
parasites. Cyproheptadine [2.5mg/kg] and LM [0.375mg/kg] killed a paltry 6% of the
parasites while cyproheptadine [5mg/kg] and LM [0.375mg/kg] yielded 20% compared to 17%
of LM [0.375mg/kg] alone (Table 2). Neither of the cyproheptadine concentration increased
the susceptibility of the wildtype parasites to LM (p<0.09), meaning that the potentiation of
LM activity was selective only in the resistant parasites.

Our next combination assays measured the impact of cyproheptadine on PQ or LM
activity against the PQ[R] parasites. On average, 75mg/kg of PQ yielded 8% parasite killing, a
combination of cyproheptadine [5mg/kg] and PQ [75mg/kg] yielded 14% while
cyproheptadine [2.5mg/kg] and PQ [75mg/kg] showed 8% parasite killing. We thus concluded
that cyproheptadine failed to restore the activity of PQ against PQ[R] parasites (Table 3).

Our previous studies have shown that a significant loss of LM activity accompanies
the selection of PQ resistance[26]. We, therefore, investigated whether cyproheptadine might
restore LM activity against the PQ[R] parasites. As expected, 50mg/kg of LM alone had no
measurable antimalarial activity (0%) against the PQ[R] parasites (Fig 3A). Surprisingly, LM
activity was re-established by 5mg/kg of cyproheptadine to above 70% (p<0.001) (Fig 3A).
The overall results reveal that cyproheptadine selectively reverses LM resistance in LM[R]
parasites but does not restore PQ activity against PQ[R] parasites. However, it selectively
restores LM potency against PQ[R] parasites.

Probenecid and verapamil reinstate LM potency against the PQ[R] parasites but not
against the LM[R] parasites

We screened for the effect of probenecid or verapamil on the growth of the asexual
blood-stage of LM[R] and PQ[R] parasites (Table 2). First, we assessed the activity of PQ against
the WT parasites. PQ at 6mg/kg and 1.5mg/kg killed 98.69% and 57.96%, respectively,
confirming the potency of the drug (Table 3). Verapamil [50mg/kg] yielded 0% and 0.81%
against the LM[R] and PQ[R] parasites, respectively. Probenecid [400mg/kg] yielded 8.44 % and
8.31% against LM[R] and PQ[R] parasites. Probenecid [400mg/kg] and verapamil [50mg/kg] were
therefore selected for combination assays with either LM or PQ. We first tested the impact
of probenecid or verapamil on LM potency against the LM[R] parasites (Fig 3B). While

100mg/kg of LM yielded 23% parasite killing, a combination of probenecid [400mg/kg] and
LM [100mg/kg] yielded 28.2%. Verapamil [50mg/kg] and LM [100mg/kg] yielded 26.5%.
These results suggest that probenecid or verapamil failed to restore LM potency against LM[R]
parasites. Next, we measured the impact of probenecid or verapamil on PQ activity against
the PQ[R] parasites (Table 3). While PQ [75mg/kg] showed a meager 8.39% parasite killing, a
combination of PQ [75mg/kg] with either probenecid [400mg/kg] or verapamil [50mg/kg]
yielded 0% parasite killing (Table 3). These results show that both probenecid and verapamil
did not potentiate PQ activity against the PQ[R] parasites.

Since the PQ[R] parasites are also highly resistant to LM[26], we evaluated whether
probenecid or verapamil could restore LM potency against PQ[R] parasites. As expected, LM
[50mg/kg] had 0% parasite killing. To our surprise, probenecid at 400mg/kg reinstated the
parasite killing potency of LM [50mg/kg] from 0% to above 60% (p<0.0001) (Fig 3B). In a
familiar trend with other reversing agents, 50mg/kg of verapamil significantly re-established
LM [50mg/kg] potency against the PQ[R] parasites from 0% to above 55% (p<0.0001) (Fig 3B).
This selective restoration of LM by probenecid, cyproheptadine, and verapamil against the
PQ[R] parasites formed the basis for our subsequent molecular studies.

205 Essential drugs transporters and enzymes are differentially expressed between drug-
206   resistant and wildtype parasites

To dissect the mechanisms of PQ and LM resistance and reversal, we hypothesized
that major transporters and enzymes involved in the transport and facilitation of drug action
could mediate LM and PQ resistance. These genes include; the *crt, mdr1, mrp2, pank, fnr,*
pmiv, pmix, pmx, SufS, and pi3k. In PQ[R] parasites, the mRNA profile showed that the
expression of two genes, *SufS,* and *pmix* was upregulated by 1.47-fold (p<0.018) and 2.08-
fold (p<0.007), respectively (Fig 4). The mRNA expression was suppressed in nine genes in
the LM[R] parasites, save for the pmiv gene, which showed marginal elevation of the mRNA
amounts (1.04fold (p<0.56) (Fig S1 https://osf.io/rfvgp). After profiling and comparing the
gene expression trends between the LM[R] and PQ[R] parasites, we found that SufS, pmiv, and
pmix had differential expression trends between the two drug-resistant parasites. Overall, we
recorded a lower expression of mRNA in *crt, mdr1, mrp2, fnr, pmx,* and *pi3k* genes in both
LM[R] and PQ[R] parasites. Two genes, pmix, and sufs were uniquely elevated and warranted
further investigations. To date, pmix has not been associated with the susceptibility of current
antimalarial drugs; however, genes in Fe-S biogenesis and FNR redox systems have been
associated with ACTs susceptibility such as artemether and lumefantrine [30,31]. Additionally,
we recently mapped a nonsynonymous mutation in the FNR protein that we associate with
LM and artemisinin (ART) susceptibility in *P. berghei* (Kiboi et al. unpublished data). The FAD

containing FNR is a redox couple protein of the ferredoxin (Fd) in the FNR redox system[32]. To
function, the FNR redox system relies on the supply of Fe-S clusters supplied by SufS and
associated enzyme SufE within the apicoplast [33]. In this study, we selected FNR and SufS
protein for further evaluation. Since FNR was downregulated while SufS was overexpressed
in the PQ^R parasites, we sought to reverse these mRNA expression trends and then measure
the impact on drug susceptibility.

The successful generation of SufS knockdown or FNR over-expression transgenic parasites

We sought to investigate the impact of altering mRNA expression of FNR or SufS in
the PQ^R parasites on LM susceptibility. We successfully generated by transfection two
transgenic parasites: one with a deleted SufS gene (PQ^R_SufS_KD) and the other with an
overexpressed FNR gene (PQ^R_FNR_OE) (Fig 5). Recovered parasites after transfection were
cloned by limiting dilution to generate genetically homogenous parasites and genotyped using
three sets of primers: QCR2/GW2, GT/GW1, and QCR1/QCR2. As expected, using the vector-
specific QCR2 and the standard GW2 primers, we obtained a fragment of 0.9kb, and 1.5kb,
respectively, in PQ^R_SufS_KD (Fig 5A) and PQ^R_FNR_OE parasites (Fig 5B). PCR amplification
analysis using vector-specific quality control primers, the QCR1, and QCR2 amplified in both
PQ^R parasites and PQ^R_SufS_KD parasites (Fig 5A) as well as in both PQ^R and PQ^R_FNR_OE
parasites (Fig 5B) confirming that the SufS knock out vector, as highlighted from PlasmoGEM
[34], does not cover the entire gene thus a possibility of generating a knockdown transgenic
parasites. To confirm integration of the vector into the correct chromosome and position, PCR
analysis using standard primers GW1 and the vector-specific primers GT yielded, as
anticipated, PCR products of 1.9kb and 3.9kb for PQ^R_SufS_KD (Fig 5A) and PQ^R_FNR_OE
parasites, respectively (Fig 5B). To confirm the overexpression of the FNR, we quantified the
mRNA expression in PQ^R_FNR_OE line relative to the PQ^R parasites. The results indicated a
5-fold (p<0.0046) increase in the mRNA amount of the FNR gene in the PQ^R_FNR_OE
parasites (Fig 5C). To assess the partial disruption of the SufS gene in PQ^R_SufS_KD
parasites, the mRNA expression was quantified and robustly demonstrated a reduction in the
expression of the SufS mRNA (Fig 5C). Overall, these results affirmed the successful
generation of the transgenic parasites. The transgenic parasites were then utilized to
measure the impact of deleting SufS or overexpressing FNR genes on the drug responses
relative to the PQ^R and wildtype parasites.

Alteration of SufS and FNR genes significantly attenuates the growth rate of the asexual blood-stage parasites.

Here we hypothesized that alteration of a gene in parasites might concomitantly alter the growth rates of the asexual blood-stage parasites. To prosecute this hypothesis, we assessed the growth rates of the transgenic parasites (Fig 6). The wildtype parasites grew as expected with parasitemia reaching an average of 15% on Day four post parasite inoculation (D4 PI). Similarly, the PQ^R parasites had an average parasitemia of 13% on D4 PI. However, the growth rate of the transgenic parasites was significantly low. For instance, PQ^R_FNR_OE or PQ^R_SufS_KD yielded an average parasitemia of 8% or 10% respectively, suggesting a fitness cost on the growth of the asexual blood-stage parasites. After quantifying the growth fitness cost, the study revealed that alteration of FNR or SufS significantly reduces normal parasite growth rate phenotype by 47% and 33%, respectively.

Overexpression of FNR or knockdown of SufS enhances PQ^R susceptibility to LM alone or in combination with the RA.

To query the impact of partially deleting SufS and overexpressing FNR genes in the PQ^R parasites on LM and RA responses, we assessed the activity of LM alone or in combination with the RA against the transgenic and PQ^R parasites (Fig 7A). As expected, LM [50mg/kg] showed negligible parasite killing at 4.8% against the PQ^R parasites. Surprisingly, similar LM concentration regained potency against PQ^R_FNR_OE (64.5%) and PQ^R_SufS_KD (67.1%) parasites. Based on these results and if RA enhanced LM susceptibility via SufS and/or FNR in PQ^R parasites, we reasoned that the deletion of SufS and overexpressing FNR might have two biological impacts; RA might enhance LM potency against PQ^R parasites or the impact of RA on LM potency might be abolished. Against the PQ^R_FNR_OE parasites, combination of LM [50mg/kg] and cyproheptadine [5mg/kg] or probenecid [200mg/kg] or verapamil [50mg/kg] yielded 62.5%, 71.8% and 53.1% respectively. In a similar trend, against the PQ^R_SufS_KD parasites, combination of LM [50mg/kg] and cyproheptadine [5mg/kg] or probenecid [200mg/kg] or verapamil [50mg/kg] yielded 84.2%, 73.1% and 78.5% respectively. The percentage parasite killing of the RA-LM combination was not significantly different from the potency of LM alone on either PQ^R_FNR_OE or PQ^R_SufS_KD parasites. These results indicate that the partial deletion of SufS or overexpression of FNR in PQ^R parasites restored PQ^R parasites susceptibility to LM and abolished the action of RA on LM activity.

Lumefantrine and cyproheptadine exhibited higher binding affinities for both FNR and SufS proteins than verapamil or probenecid.

To confirm the *in vivo* drug susceptibility results and investigate the potential interaction of LM or RA with either SufS or FNR, we used *in-silico* bioinformatics tools and

approaches. Here, for all the structures modeled, the Z-scores from the PROSA-web server that indicates the degree of correctness showed scores of -8.49 and -8.67 for FNR and SufS, respectively (Table 4). The results of the Z-scores obtained imply that the modeled protein structures were within the range of scores typical of the experimentally defined proteins. As expected, LM yielded high but equal binding affinities of -7.9kcal/mol on both SufS and FNR, indicating a better binding between LM and the proteins. Surprisingly, cyproheptadine showed the highest binding affinity to SufS (-9.5kcal/mol). The lowest binding affinity of -6.3 kcal/mol was between FNR and verapamil (Fig 8A and Fig 8B).

DISCUSSION

Here, we unveil new evidence that chemosensitizers: cyproheptadine, verapamil, or probenecid selectively restore LM activity against PQ[R] parasites. We also show this selective reversal of LM resistance is abolished by the deletion of SufS and overexpression of the FNR gene. We have used parasites previously selected by submitting *P. berghei* parasites to PQ or LM for at least forty mechanical passages[35] to systematically dissect reversal and the mechanisms of LM and PQ resistance in *P. berghei* ANKA. During the selection process, the PQ[R] parasites were submitted to PQ pressure alone, but resultant phenotypes had significantly reduced susceptibility to CQ, AQ, LM, and DHA [26], displayed stability in the absence of PQ pressure and after freeze-thawing process [28,36]. These prior studies affirmed the multidrug resistance nature of the PQ[R] parasites. In the current study, all the chemosensitisers restored LM potency against the PQ[R] parasites, but only cyproheptadine restored LM activity against the LM[R] parasites. The current study, therefore, studied potential molecular mechanisms of LM resistance and subsequent potentiation of LM efficacy by the three chemosensitisers in the PQ[R] parasites.

Our results show that cyproheptadine has at least two functions; first, and as expected, is the parasite killing ability against both the LM[R] and the wildtype parasites, possibly by inhibition of heme polymerization as previously elaborated[15]. Second, cyproheptadine enhances parasite killing by selectively restoring LM activity against both the LM[R] and PQ[R] parasites. PQ and cyproheptadine exert parasite killing by inhibiting detoxification of heme in the digestive vacuole [15]. Therefore, the low susceptibility of the PQ[R] parasites to cyproheptadine compared to the LM[R] and the wildtype parasites suggest that PQ and cyproheptadine also share some common resistance mechanisms. The failure of cyproheptadine to restore PQ activity against PQ[R] parasites suggests that the PQ[R] parasites possess at least two types of resistance mechanisms. The first mechanism codes for PQ resistance and is cyproheptadine insensitive and second, which mediates LM resistance and is cyproheptadine susceptible. It is unclear whether the genes associated with

cyproheptadine modifiable mechanisms in both LM^R and PQ^R are similar or different. A lower

dose of cyproheptadine [5mg/kg] had only one biological function of restoring LM activity

against both LM^R and PQ^R parasites. We also note that resistance mechanisms mediating the

loss of LM activity against the PQ^R parasites are verapamil susceptible, but the loss of LM

activity against LM^R parasites are independent of verapamil action suggesting a complex

emergence of LM resistance. Resistance to quinoline antimalarials such as PQ is associated

with acidification of the digestive vacuole (DV) [12]. Verapamil mediates reversal by

replacement of positive charge and normalization of pH, blocking efflux of quinoline drugs

from digestive vacuole (DV) via the mutated PfCRT protein [37]. The mechanism of action of PQ

and LM is predicted to be similar to that of standard quinoline; chloroquine [38]. Thus, these

drugs may share some resistance mechanisms. However, *PbCRT* in PQ^R or LM^R parasites

were found not to carry mutations [36], suggesting that verapamil may use alternative proteins

to enhance drug activity. Our data provide clues that verapamil may interact with FNR and

SufS protein to restore and enhance LM activity in PQ^R parasites.

Like with verapamil and cyproheptadine, we provide compelling evidence that

mechanisms of LM resistance in PQ^R parasites are associated with proteins that interact with

organic anion inhibitors, probenecid. Probenecid is a substrate for multidrug resistance-

associated protein; for instance, in *P. falciparum*, probenecid selectively sensitizes

chloroquine-resistant isolates by increasing the level of chloroquine accumulation *in vitro* [14].

Also, probenecid unselectively potentiates the activities of antifolates agents against both

antifolates resistant and sensitive *P. falciparum*. This reversal of antifolates resistance is

associated with inhibition of multi-resistance-associated proteins [39] and inhibition of

endogenous folate transport [14]. Multi-resistance associated proteins are also implicated in

resistance to quinoline antimalarial drugs in both *P. falciparum* and *P. berghei* [40]. We have

previously confirmed PQ^R parasites as multidrug-resistant parasites; thus, probenecid

mediated inhibition of multi-resistance proteins in re-establishing LM activity against PQ^R

parasites is a possible alternative mechanism.

The next generation of antimalarial drugs has envisioned the incorporation of three

drugs comprising of two long-acting drugs and a short-acting artemisinin drug [41]. Currently,

cyproheptadine, an H-1 antagonist, at 4mg/kg, is recommended for the treatment of allergic

reactions and improvement of appetite (www.medscape.com). We have shown that 2.5mg/kg

of cyproheptadine is a potent modulator of lumefantrine resistance in both PQ^R and LM^R

parasites, at least in *P. berghei*. In humans, the recommended dosage for verapamil and

probenecid is 80mg/kg and 500mg/kg, respectively (www.medscape.com). Studies have

supported a combination of probenecid and SP for intermittent prevention of malaria in

pregnancy (IPTp) [13]. Here, we have used a lower dose to restore the activity of LM against

PQ^R parasites. If the mechanisms of LM resistance in *P. berghei* and *P. falciparum* are similar,

then our data demonstrate that incorporation of cyproheptadine into antimalarial combination

therapy that comprises LM might achieve two aims: First is to boost LM activity and possibly

delay the emergence of LM resistance. Second, an optimized cyproheptadine concentration

might offer an added advantage in the combination by exerting extra antimalarial. The data

also confirms the existence of more than one control channels for PQ resistance in PQ^R

parasites. The first is associated with LM resistance and is verapamil, probenecid, or

cyproheptadine inhibitable channel only in the presence of LM. The other control is linked

with PQ resistance, which is verapamil, probenecid or cyproheptadine insensitive

mechanisms. Currently, LM and PQ are used in ACTs therapies that are mainstay drugs for

the treatment of malaria in many African countries [42]. Again, if mechanisms of resistance to

LM in *P. berghei* are the same as those in *P. falciparum*, then the emergence of LM resistance

may proceed through two possible mechanisms; first is through the indirect PQ selection

pressure via the PQ/DHA combination. The second mechanisms are by the direct LM

pressure via the LM/ATM combination. The selective potentiation of LM potency by

probenecid, verapamil, and cyproheptadine against the PQ^R parasites is appealing since

parasites are multidrug resistant parasites and might expose shared resistance

mechanisms[26]. Our data suggest that the incorporation of any of the reversing agent into an

antimalarial combination therapy that comprises LM as one of the components might

concomitantly antagonize the emergence of LM resistance derived from PQ selection

pressure.

The impact of chemosensitizers on LM activity against both PQ^R and LM^R parasites

suggests that LM resistance is under the control of more than one molecular signature and

pathway. Our data show that changes in mRNA expression of SufS, a unique protein in the

Fe-S biogenesis [30], and FNR in the FNR redox system; [43] are associated with PQ and LM

resistance. The fact that an increase of the mRNA amount of FNR correlated with increased

susceptibility of PQ^R parasites to LM and concomitantly abolished the impact of RA on LM

activity confirm the interaction between RA and LM with FNR. Indeed, our *in-silico* binding

results show significant binding power between the RA, LM, and FNR. Several studies have

established that the Fd/FNR redox system, which supplies reducing power for essential

biosynthetic pathways in the malaria parasite apicoplast [44], is a potential target for the

development of specific new antimalarial agents [43]. Validation of the FNR redox system as a

potential target and resistance marker for the artemisinins was revealed by the

nonsynonymous background mutation (Asp193Tyr) in the *Fd* protein in parasite exposed to

the ACTs[31]. We provide a clear link between LM activity, RA action, and overexpression of the FNR gene. Redox enzymes such as FNR convert inactive compounds into active molecules [45],[46]. Two possible assumptions exist; First, FNR may be a target for LM. Reduced expression of FNR in the PQ^R parasites may potentially result in reduced binding targets for LM. The high FNR expression in the transgenic parasites PQ^R_FNR_OE seems to restore the concentration of the FNR, thus more binding targets for LM and a consequent high susceptibility of PQ^R_FNR_OE to LM. Second, the FNR is a potential activator for LM, therefore reduced expression of FNR in the PQ^R is accompanied by reduced active metabolites from LM, thus less activity. The increase in the FNR, as confirmed in the PQ^R_FNR_OE transgenic parasites, seems to reinstate the activator; therefore, more of the LM metabolite is generated restoring LM activity in the PQ^R_FNR_OE transgenic parasites as compared to PQ^R parasites. The molecular mechanism of drug-target interaction between the RA and LM with FNR requires further in-depth investigation. The reversing agents restore the activity of antimalarial drugs by enhancing the accumulation of drugs within the target site [18]. It is likely that in the event the FNR amount is low, the threshold concentration for LM action is not achieved, the threshold may be enhanced by a RA possibly by increasing the uptake of the few active LM metabolites.

The FNR redox system relies on the Fe-S clusters for its action [47]. Both the FNR redox system and Fe-S cluster biogenesis are essential pathways for parasite survival within the red blood cells [33],[43], the site of action of LM and PQ. Plasmodium species possess two Fe-S cluster biogenesis pathways: the mitochondria localized ISC pathway and the SUF pathway within the apicoplast. Here, we focused on apicoplast exclusive SufS enzyme [48]. SufS converts L-cysteine to L-alanine and sulfane sulfur via the formation of an enzyme-bound persulfide intermediate within the apicoplast compartment[49]. Here, we first confirm a significant attenuation of the growth rates on the deletion of the SufS gene, thus essential for the growth of asexual blood-stage parasites. Second, the results presented here show that an increase in the RNA amount of SufS correlates with decreased LM susceptibility in PQ^R parasites.

We have validated these findings through the deletion of the SufS gene in PQ^R parasites yielding parasite lines: PQ^R_SufS_KD that showed significant susceptibility to LM and simultaneously obliterated the impact of cyproheptadine, verapamil, and probenecid. Taken together, our results suggest that LM interacts with SufS and exert its antimalarial action through the protein. However, this mechanism seems to be solely in the PQ^R parasites. We reveal clues that SufS may be a target for LM. Thus, the low expression of SufS in the LM^R parasites may be adaptive mechanisms for the parasites to reduce potential binding targets for the LM. Indeed, the adaptive parasite mechanism is supported by the low growth

profiles of the parasite observed in this study after partial deletion of the SufS in the PQ[R]

parasites. Abolishing of the impact of chemosensitiser on LM activity, expose additional route

for reversal of resistance using chemosensitisers. Our results on mRNA expression of SufS

seemed antagonistic between PQ[R] and LM[R] parasites. These findings reveal that LM

resistance may associate with SufS if the selection pressure applied is PQ; however, LM

pressure does not select high expression of SufS, and thus resistance to LM through LM

pressure is independent of SufS protein.

In conclusion, this study provides critical evidence regarding LM resistance in a PQ

resistant *P. berghei* ANKA; As highlighted earlier, PQ pressure yields multidrug resistant PQ[R]

parasites that are accompanied by the loss of LM efficacy [26,28]. First, we have restored LM

efficacy against the PQ[R] parasites by combining with either cyproheptadine, verapamil, or

probenecid. SufS deletion and FNR overexpression in the PQ[R] parasites abolishes the impact

of the chemosensitizers. We provide essential clues and propose the association of SufS and

FNR with the action of LM and chemosensitizers. Taken together, if mechanisms of LM

resistance between *P. berghei* and *P. falciparum* are similar, then the mRNA expression of

SufS and FNR genes may be predictors of the emergence of LM resistance in the *P.*

falciparum field isolates. Also, our data suggest that the incorporation of any of the

chemosensitizers into an antimalarial combination that comprises LM might augment LM

activity and concurrently antagonize the emergence of LM resistance derived from PQ

pressure. The in-depth mechanisms of how SufS and FNR mediate LM resistance and the

impact of their differential expression in drug susceptibility should be investigated in the

human malaria parasite *P. falciparum*.

MATERIALS AND METHODS

Parasites, Host, and Compounds

We used two parasite reference lines *of P. berghei* ANKA, the MRA-865 and MRA-868

reference lines obtained from MR4, ATCC® Manassas, Virginia, which were used as the drug-

susceptible wildtype (WT) parasites. Also, two multidrug-resistant parasite lines; the PQ-

resistant (PQ[R]) and LM-resistant (LM[R]) *P. berghei* ANKA parasites previously selected from

MRA-865 and MRA-868 lines, respectively[26] were studied. The two parasite lines were

confirmed as stable resistant parasites[26,28]. Male Swiss albino mice weighing 20 ± 2g outbred

at KEMRI, Animal house Nairobi were utilized, housed, and fed as previously described[50]. The

knockout (KO) vector (PbGEM-018972), as well as the overexpression (OE) vector (PbGEM-

456502), were kindly provided by the *Plasmo*GEM project under an agreement of a material

transfer (PG-MTA-0093). Graphical overview of KO and OE vector design are outlined in

PlasmoGEM (https://plasmogem.sanger.ac.uk/), and experimental strategies have

previously been detailed[34]. Verapamil, probenecid, and cyproheptadine were purchased from

Carramore international limited (UK). LM and PQ were donated from Universal Corporation,

Kikuyu, Kenya. All drugs, PQ, LM, verapamil, probenecid, and cyproheptadine, were freshly

prepared and administered as previously described[36].

Ethical consideration

This study was conducted at KEMRI and JKUAT. All mouse experiments were carried

out as per relevant national and international standards (The ARRIVE Guidelines) and as

approved by KEMRI-Animal Use and Care Committee. This study was permitted and ethically

approved by KEMRI under the SERU No 3914.

Drug sensitivity assays

The activity profiles of probenecid, verapamil, or cyproheptadine alone and in

combination with PQ, or LM against the wildtype, resistant, or the transgenic parasites were

assessed in the standard 4-day suppressive test (4-DT) in which the parasite is exposed to

four daily drug doses[51]. Infection, randomization of experimental mice, drug administration,

estimation of parasite densities, and calculation of percentage drug killing was performed

and determined as previously described[26,50,52].

Extraction of RNA, synthesis of cDNA, and qPCR assays

The quantification of mRNA transcripts of *Pbcrt, Pbmdr1, Pbmrp2, Pbpank, Pbfnr, PbSufS,*

Pbpmiv, Pbpmix, Pbpmx, and *Pbpi3k* was carried out using primer listed on Table S1

https://osf.io/j5w3e. Extraction of total RNA from the wildtype, LM^R, PQ^{R,} and transgenic

parasites and synthesis of respective first-strand cDNA was performed as previously

described[50]. Using the *Pbβ-actin I*, as the housekeeping gene, the relative expression profile

of the ten genes was performed in a final volume of 20 μ l using Maxima SYBR Green/ROX

qPCR Master Mix (Thermo Scientific™). Briefly, 12µl of Maxima SYBR mix, 3µl water were

mixed with 2.0µl (0.25µM) of forward and reverse primers each, and 1µl cDNA. The reaction

mix was run for pre-treatment at 50° C, for 2 min; initial denaturation at 95° C for 10 min;

denaturation at 95° C for 15 secs; and annealing at 60° C for 60 secs for 45 cycles.

Generation of the transgenic parasites

Isolation, digestion, and purification of the vector DNA

Using the QIAfilter Plasmid Midi Kit (Qiagen™), Plasmid DNA for each of the vectors

was isolated from cultures of *E. coli* (vector hosts) in terrific broth (TB) supplemented with

kanamycin after overnight culture. Following a standard protocol, the extracted plasmid DNA

was restricted using Not I enzyme to release the vector backbone, followed by concentration

using conventional ethanol precipitation protocol[53]. The vector was dried for 5 min at 65˚C

and then dissolved in $10\,\mu$l of PCR water. Besides Not I restriction digest, a diagnostic PCR

was performed using standard primer pair (GW2 and a vector-specific QCR2)[34] to verify the

isolated vector of interest.

Purification of the *P. berghei* schizonts for transfection

Using standard *in vitro* culture protocols as described by [54], we collected *P. berghei*

schizonts for transfection. Briefly, at least three mice were used for schizont culture for each

of the vectors. Propagation of *P. berghei* parasites was done by intraperitoneal (IP) injection

into the mice, and parasites were harvested using cardiac puncture for schizont culture at 3%

parasitemia. For schizonts propagation and subsequent purification, rings stage parasites

(the asexual blood-stages) were incubated at 37˚C in vitro as reported previously[52,55].

Harvested and purified schizonts were re-suspended in schizont culture media, ready for

transfection.

Transfection and selection of transgenic parasite lines

A SufS specific knockout vector (PbGEM-018972) and an FNR overexpression vector

(PbGEM-456502) were separately transfected in the PQ[r] parasites. Each of the transfection

mixes contained, $20\,\mu$l of schizonts, $10\,\mu$l ($5\,\mu$g) of plasmid DNA, and 100ml of supplemented

nucleofector solution (Amaxa™). Exactly $100\,\mu$l of the transfection mix in Amaxa cuvette was

electroporated in a Nucleofector 2B Device (Lonza™) using the program U33. A naïve mouse

was intravenously injected with the electroplated parasites. Three transfections were

performed for each of the vectors to increase the chance of recovering transgenic parasites.

Pyrimethamine drug at 7 mg/mL, for the selection of transgenic parasites, was provided in

drinking water, 24hours post-parasite inoculation for a total of 12 days. Freshly prepared

pyrimethamine drug was provided every 72 hours. After the recovery of transgenic parasites,

genetically homogenous parasite lines were generated by dilution cloning protocol [36,54].

Genotyping of transgenic parasites

Genomic DNA (gDNA) was extracted from the transgenic parasite lines using the

QIAamp DNA Blood Mini Kit (Qiagen™) following manufacturer instructions. Three pairs of

primers were used for diagnosis and genotyping, as detailed in Table S2 https://osf.io/zkxyj:

i) QCR2/GW2 pair ii) QCR1/QCR2 and iii) GT/GW1 or GW2. Each reaction mixes contained

1µl of gDNA, 12.5µl of 2xGoTag Green master mix (Promega™), 6.5µl of water and 2.5µl of

each of the primers. The PCR reactions were run using a 2xGoTag Green master mix for thirty

cycles with an annealing temperature of 50° C and 62° C as the extension temperature. The

PCR products were resolved on a 1% agarose gel.

Molecular docking

Homology modeling

The 3D structures of the SufS; PBANKA_0614300, and FNR; PBANKA_1122100, were predicted by selecting as templates, the best structures with the highest similarity to their sequences available in NCBI protein database http://ncbi.nlm.nih.gov and *Plasmodium* database http://plasmodb.org/. SWISS-MODEL [56] available at https://swissmodel.expasy.org/ was used to predict the homology models. The SWISS-MODEL was a preferred modeling server because it annotates essential cofactors and ligands as well as quaternary structures, allowing for modeling of complete structures with less complicated software packages [56]. The structures were then downloaded and saved in PDB format.

Structure validation

The models were analyzed using PROCHECK [57] and PRoSA-web [58] to determine the quality of the modeled structures. PROCHECK was used to evaluate the general stereochemistry of the protein, while PRoSA-web was employed to assess for potential errors in the 3D structures. Verify_3D was used to determine the compatibility of an atomic model (3D) with its amino acid sequence (1D)[59].

Ligand selection

To perform docking experiments, the most potent RA, as well as LM, were selected as test ligands. The molecular structures of the four chosen ligands were downloaded from the ChemSpider (http://www.chemspider.com/), an online database to access unique chemical compounds [60]. For compatibility with the docking software (https://cactus.nci.nih.gov/translate/), we converted the chemical structures from the Mol2 format into PDB format using the CADD Group's Chemoinformatics Tools and User Services

Binding site analysis and ligand docking

PyMOL version 1.6.x (Delano, 2002) was used to visualize the modeled structures and binding sites. The grid box parameters of the ligands and the receptors optimization were executed using scripts within AutoDock Tools (ADT). The files were then saved in PDBQT format, and their corresponding coordinates rewrote into a configuration file used for docking. The configuration file specified the pdbqt files for both the ligand and the proteins as well as the docking parameters (dimensions and spacing angstrom). Optimization of the inputs involved the removal of water molecules from the receptor and the addition of missing

hydrogen ions. Docking of a single receptor and a single ligand was carried out using the
Autodock Vina [61,62]. Autodock Vina employs the Lamarckian genetic algorithm (LGA). FNR and
SufS, whose 3D structures were available in the protein data bank (PDB;
http://www.rcsb.org/pdb/explore.do?structureId=2hvp) were also included. The binding
energies and the positional root-mean-square deviation (RMSD) of the proteins and the
ligands were presented on an output file. A ligand orientation with low binding energy signifies
a better affinity towards a receptor.

Data presentation and statistical analysis

The percentage parasitemia and the percentage of drug killing were analyzed using the
Nonparametric Mann-Whitney U Test in the R statistical software with a p-value set at 0.05.
The means for the gene expression levels from three independent experiments and triplicate
assays were compared using the Mann-Whitney U test with a p-value set at 0.05. The relative
expression level data were normalized using β-actin I as the housekeeping gene based on
the formula $2^{\Delta\Delta CT}$[63].

Data Availability

All lumefantrine reversal datasheet files are available from the https://osf.io/brz63/
All piperaquine reversal datasheet files are available from the https://osf.io/5trab/
All PQR_FNR_OE parasites datasheet files are available from the https://osf.io/pdvj6/
All PQr_SufS_KD parasites datasheet files are available from the https://osf.io/8upkc/

Acknowledgments

We thank the Wellcome Trust Sanger Institute *Plasmo*GEM project team for providing the
highly efficient gene modification resources used in this study.

References

1. Global Malaria Programme: WHO Global. *World Malaria Report 2019.*; 2019. https://www.who.int/news-room/fact-sheets/detail/malaria.

2. WHO. *Guidelines for the Treatment of Malaria, 3rd Edition. World Health Organization.*; 2016.

3. Tse EG, Korsik M, Todd MH. The past, present and future of anti-malarial medicines. *Malar J.* 2019;18(1):93. doi:10.1186/s12936-019-2724-z

4. Nosten F, White NJ. Artemisinin-based combination treatment of falciparum malaria. *Am J Trop Med Hyg.* 2007. doi:10.4269/ajtmh.2007.77.181

5. White NJ. Malaria parasite clearance. *Malar J.* 2017;16(1):1-14. doi:10.1186/s12936-017-1731-1

6. Fidock DA, Nomura T, Talley AK, et al. Mutations in the P. falciparum digestive vacuole transmembrane protein PfCRT and evidence for their role in chloroquine resistance. *Mol Cell.* 2000;6(4):861-871. doi:10.1016/S1097-2765(05)00077-8

7. Sidhu ABS, Uhlemann A, Valderramos SG, Valderramos J, Krishna S, Fidock DA. Decreasing pfmdr1 Copy Number in Plasmodium falciparum Malaria Heightens Susceptibility to Mefloquine, Lumefantrine, Halofantrine, Quinine, and Artemisinin. *J Infect Dis.* 2006;194(4):528-535. doi:10.1086/507115

8. Sidhu ABS, Valderramos SG, Fidock DA. pfmdr1 mutations contribute to quinine resistance and enhance mefloquine and artemisinin sensitivity in Plasmodium falciparum. *Mol Microbiol.* 2005;57(4):913-926. doi:10.1111/j.1365-2958.2005.04729.x

9. Dhingra SK, Redhi D, Combrinck JM, et al. A variant pfcrt isoform can contribute to plasmodium falciparum resistance to the first-line partner drug piperaquine. *MBio.* 2017;8(3). doi:10.1128/mBio.00303-17

10. Amato R, Lim P, Miotto O, et al. Genetic markers associated with dihydroartemisinin–piperaquine failure in Plasmodium falciparum malaria in Cambodia: a genotype–phenotype association study. *Lancet Infect Dis.* 2017;17(2):164-173. doi:10.1016/S1473-3099(16)30409-1

11. Nzila A, Okombo J, Ohuma E, Al-Thukair A. Update on the in vivo tolerance and in vitro reduced susceptibility to the antimalarial lumefantrine. *J Antimicrob Chemother.* 2012;67(10):2309-2315. doi:10.1093/jac/dks252

12. Henry M, Alibert S, Rogier C, Barbe J, Pradines B. Inhibition of Efflux of Quinolines as New Therapeutic Strategy in Malaria. *Curr Top Med Chem.* 2008. doi:10.2174/156802608783955593

13.   Gutman J, Kachur S, Slutsker L, Nzila A, Mutabingwa T. Combination of probenecid-
sulphadoxine-pyrimethamine for intermittent preventive treatment in pregnancy.
Malar J. 2012;11. doi:10.1186/1475-2875-11-39

14.   Nzila A, Mberu E, Bray P, et al. Chemosensitization of Plasmodium falciparum by
probenecid in vitro. *Antimicrob Agents Chemother*. 2003;47(7):2108-2112.
doi:10.1128/AAC.47.7.2108-2112.2003

15.   Agrawal R, Tripathi R, Tekwani BL, Jain SK, Dutta GP, Shukla OP. Haem polymerase
as a novel target of antimalarial action of cyproheptadine. *Biochem Pharmacol*.
2002;64(9):1399-1406. doi:10.1016/S0006-2952(02)01299-6

16.   Bates SE, Wilson WH, Fojo AT, et al. Clinical reversal of multidrug resistance. *Stem
Cells*. 1996;14(1):56-63. doi:10.1002/stem.140056

17.   Lehane AM, Hayward R, Saliba KJ, Kirk K. A verapamil-sensitive chloroquine-
associated H+ leak from the digestive vacuole in chloroquine-resistant malaria
parasites. *J Cell Sci*. 2008;121(10):1624-1632. doi:10.1242/jcs.016758

18.   Warhurst DC, Craig JC, Adagu IS. Lysosomes and drug resistance in malaria. *Lancet
(London, England)*. 2002;360(9345):1527-1529. doi:10.1016/S0140-6736(02)11577-7

19.   Zheng W, Li M, Lin Y, Zhan X. Encapsulation of verapamil and doxorubicin by MPEG-
PLA to reverse drug resistance in ovarian cancer. *Biomed Pharmacother*. 2018.
doi:10.1016/j.biopha.2018.09.039

20.   Berger D, Citarella R, Dutia M, et al. Novel multidrug resistance reversal agents. *J
Med Chem*. 1999. doi:10.1021/jm9804477

21.   De Luca A, Parker LJ, Ang WH, et al. A structure-based mechanism of cisplatin
resistance mediated by glutathione transferase P1-1. *Proc Natl Acad Sci U S A*. 2019.
doi:10.1073/pnas.1903297116

22.   H. Peyton D. Reversed Chloroquine Molecules as a Strategy to Overcome Resistance
in Malaria. *Curr Top Med Chem*. 2012. doi:10.2174/156802612799362968

23.   Martiney JA, Cerami A, Slater AFG. Verapamil reversal of chloroquine resistance in
the malaria parasite Plasmodium falciparum is specific for resistant parasites and
independent of the weak base effect. *J Biol Chem*. 1995.
doi:10.1074/jbc.270.38.22393

24.   Sirotnak FM, Wendel HG, Bornmann WGB, et al. Co-administration of probenecid, an
inhibitor of a cMOAT/MRP-like plasma membrane ATPase, greatly enhanced the
efficacy of a new 10-deazaaminopterin against human solid tumors in vivo. *Clin
Cancer Res*. 2000.

25.   van Schalkwyk DA, Egan TJ. Quinoline-resistance reversing agents for the malaria

657  parasite Plasmodium falciparum. *Drug Resist Updat*. 2006;9(4-5):211-226.

658  doi:10.1016/j.drup.2006.09.002

659 26. Kiboi DM, Irungu BN, Langat B, et al. Plasmodium berghei ANKA: Selection of

660  resistance to piperaquine and lumefantrine in a mouse model. *Exp Parasitol*.

661  2009;122(3):196-202. doi:10.1016/j.exppara.2009.03.010

662 27. Langat B, Kiboi D, Irungu B, Kimoloi S. Lumefantrine-resistant and Piperaquine-

663  resistant Plasmodium berghei show cross-resistance to Primaquine but not to

664  Atovaquone. *African J Pharmacol Ther*. 2012;1(2):35-40.

665 28. Gimode WR, Kiboi DM, Kimani FT, Wamakima HN, Burugu MW, Muregi FW. Fitness

666  cost of resistance for lumefantrine and piperaquine-resistant Plasmodium berghei in

667  a mouse model. *Malar J*. 2015;14(1). doi:10.1186/s12936-015-0550-5

668 29. Fda. CYPROHEPTADINE HYDROCHLORIDE LEAFLET CYPROHEPTADINE

669  HYDROCHLORIDE TABLETS USP Rx only. 2007.

670  https://www.accessdata.fda.gov/drugsatfda_docs/label/2009/087056s045lbl.pdf.

671 30. Amambua-ngwa A, Jeffries D, Amato R, et al. Consistent signatures of selection from

672  genomic analysis of pairs of temporal and spatial Plasmodium falciparum populations

673  from The Gambia. 2018;(June):1-10. doi:10.1038/s41598-018-28017-5

674 31. Miotto O, Amato R, Ashley EA, et al. Genetic architecture of artemisinin-resistant

675  Plasmodium falciparum. *Nat Genet*. 2015;47(3):226-234. doi:10.1038/ng.3189

676 32. Pandini V, Caprini G, Thomsen N, Aliverti A, Seeber F, Zanetti G. Ferredoxin-NADP+

677  reductase and ferredoxin of the protozoan parasite Toxoplasma gondii interact

678  productively in vitro and in vivo. *J Biol Chem*. 2002;277(50):48463-48471.

679  doi:10.1074/jbc.M209388200

680 33. Pala ZR, Saxena V, Saggu GS, Garg S. Recent Advances in the [Fe–S] Cluster

681  Biogenesis (SUF) Pathway Functional in the Apicoplast of Plasmodium. *Trends*

682  *Parasitol*. 2018;34(9):800-809. doi:10.1016/j.pt.2018.05.010

683 34. Schwach F, Bushell E, Gomes AR, et al. PlasmoGEM, a database supporting a

684  community resource for large-scale experimental genetics in malaria parasites.

685  *Nucleic Acids Res*. 2015. doi:10.1093/nar/gku1143

686 35. Irungu B, Kiboi D, Langat B, Rukunga G, Wittlin S, Nzila A. Methotrexate and

687  aminopterin lack in vivo antimalarial activity against murine malaria species. *Exp*

688  *Parasitol*. 2009;123(2):118-121. doi:10.1016/j.exppara.2009.06.007

689 36. Kiboi D, Irungu B, Orwa J, et al. Piperaquine and lumefantrine resistance in

690  Plasmodium berghei ANKA associated with increased expression of

691  Ca2+/H+antiporter and glutathione associated enzymes. *Exp Parasitol*. 2014;147:23-

32. doi:10.1016/j.exppara.2014.10.008

37.   Warhurst DC, Steele JCP, Adagu IS, Craig JC, Cullander C. Hydroxychloroquine is

much less active than chloroquine against chloroquine-resistant Plasmodium

falciparum, in agreement with its physicochemical properties. *J Antimicrob*

Chemother. 2003. doi:10.1093/jac/dkg319

38.   Robert A, Benoit-Vical F, Dechy-Cabaret O, Meunier B. From classical antimalarial

drugs to new compounds based on the mechanism of action of artemisinin. *Pure Appl*

Chem. 2001;73(7):1173-1188. doi:10.1351/pac200173071173

39.   Hooijberg JH, Broxterman HJ, Kool M, et al. Antifolate resistance mediated by the

multidrug resistance proteins MRP1 and MRP2. *Cancer Res*. 1999.

40.   Valderramos SG, Fidock DA. Transporters involved in resistance to antimalarial drugs.

Trends Pharmacol Sci. 2006;27(11):594-601. doi:10.1016/j.tips.2006.09.005

41.   J KM, Krishna S. Triple artemisinin - containing combination anti - malarial

treatments should be implemented now to delay the emergence of resistance : the

case against. *Malar J*. 2019:1-3. doi:10.1186/s12936-019-2976-7

42.   Eastman RT, Fidock DA. Artemisinin-based combination therapies: A vital tool in

efforts to eliminate malaria. *Nat Rev Microbiol*. 2009;7(12):864-874.

doi:10.1038/nrmicro2239

43.   Suwito H, Jumina, Mustofa, et al. Design and synthesis of chalcone derivatives as

inhibitors of the ferredoxin - Ferredoxin-NADP+ reductase interaction of Plasmodium

falciparum: Pursuing new antimalarial agents. *Molecules*. 2014;19(12):21473-21488.

doi:10.3390/molecules191221473

44.   Balconi E, Pennati A, Crobu D, et al. Balconi et al 2009 The ferredoxin-NADP+

reductase∕ferredoxin electron.pdf. *FEBS J Authors J Compil*. 2009;276:3825-3836.

doi:10.1111/j.1742-4658.2009.07100.x

45.   Graf N, Lippard SJ. Redox activation of metal-based prodrugs as a strategy for drug

delivery. *Adv Drug Deliv Rev*. 2012;64(11):993-1004. doi:10.1016/j.addr.2012.01.007

46.   Lesanavičius M, Aliverti A, Šarlauskas J, Čėnas N. Reactions of plasmodium

falciparum ferredoxin:Nadp+ oxidoreductase with redox cycling xenobiotics: A

mechanistic study. *Int J Mol Sci*. 2020;21(9):9-11. doi:10.3390/ijms21093234

47.   Charan M, Singh N, Kumar B, Srivastava K, Siddiqi MI, Habib S. Sulfur mobilization for

fe-s cluster assembly by the essential suf pathway in the plasmodium falciparum

apicoplast and its inhibition. *Antimicrob Agents Chemother*. 2014.

doi:10.1128/AAC.02711-13

48.   Pala ZR, Saxena V, Saggu GS, et al. Functional analysis of iron-sulfur cluster

biogenesis (SUF pathway) from Plasmodium vivax clinical isolates. *Exp Parasitol*.
2019. doi:10.1016/j.exppara.2019.01.015

49.   Gisselberg JE, Dellibovi-Ragheb TA, Matthews KA, Bosch G, Prigge ST. The Suf Iron-
Sulfur Cluster Synthesis Pathway Is Required for Apicoplast Maintenance in Malaria
Parasites. *PLoS Pathog*. 2013. doi:10.1371/journal.ppat.1003655

50.   Ndung'u L, Langat B, Magiri E, et al. Amodiaquine resistance in plasmodium berghei
is associated with PbCRT His95Pro mutation, loss of chloroquine, artemisinin and
primaquine sensitivity, and high transcript levels of key transporters. *Wellcome Open*
Res. 2018;2(0). doi:10.12688/wellcomeopenres.11768.2

51.   Fidock DA, Rosenthal PJ, Croft SL, Brun R, Nwaka S. Antimalarial drug discovery:
Efficacy models for compound screening. *Nat Rev Drug Discov*. 2004;3(6):509-520.
doi:10.1038/nrd1416

52.   Onchieku NM, Mogire R, Ndung'u L, et al. Deciphering the targets of retroviral
protease inhibitors in Plasmodium berghei. *PLoS One*. 2018;13(8):1-16.
doi:10.1371/journal.pone.0201556

53.   Pfander C, Anar B, Schwach F, et al. A scalable pipeline for highly effective genetic
modification of a malaria parasite. *Nat Methods*. 2011;8(12):1078-1084.
doi:10.1038/nmeth.1742

54.   Janse CJ, Ramesar J, Waters AP. High-efficiency transfection and drug selection of
genetically transformed blood stages of the rodent malaria parasite Plasmodium
berghei. *Nat Protoc*. 2006;1(1):346-356. doi:10.1038/nprot.2006.53

55.   Gomes AR, Bushell E, Schwach F, et al. A genome-scale vector resource enables
high-throughput reverse genetic screening in a malaria parasite. *Cell Host Microbe*.
2015. doi:10.1016/j.chom.2015.01.014

56.   Biasini M, Bienert S, Waterhouse A, et al. SWISS-MODEL: Modelling protein tertiary
and quaternary structure using evolutionary information. *Nucleic Acids Res*.
2014;42(W1):252-258. doi:10.1093/nar/gku340

57.   Laskowski RA, MacArthur MW, Moss DS, Thornton JM. PROCHECK: a program to
check the stereochemical quality of protein structures. *J Appl Crystallogr*. 1993.
doi:10.1107/s0021889892009944

58.   Wiederstein M, Sippl MJ. ProSA-web : interactive web service for the recognition of
errors in three-dimensional structures of proteins. *Nucleic Acids Res*. 2007;35(2):407-
410. doi:10.1093/nar/gkm290

59.   Bowie JU, Lüthy R, Eisenberg D. A Method to Identify Protein Sequences That into
Structure a Known Three-Dimensional Structure. *Science (80-)*. 1991;253(5016):164-

170.

60.   Williams AJ, Pence HE. ChemSpider : An Online Chemical Information Resource. *J*

Chem Educ. 2010;87(11). doi:10.1021/ed100697w

61.   Forli S, Huey R, Pique ME, Sanner M, Goodsell DS, Arthur J. 00006565-201002000-

00017. 2016;11(5):905-919. doi:10.1038/nprot.2016.051.Computational

62.   Trott O, Olson A. Autodock vina: improving the speed and accuracy of docking. *J*

Comput Chem. 2010;31(2):455-461. doi:10.1002/jcc.21334.AutoDock

63.   Livak KJ, Schmittgen TD. Analysis of Relative Gene Expression Data Using Real- Time

Quantitative PCR and the 2 $\mathbf{x}$ $\boxtimes\boxtimes$ C T Method. 2001;408:402-408.

doi:10.1006/meth.2001.1262

Fig 1: Activity profile of cyproheptadine (CYP), Lumefantrine (LM) and piperaquine (PQ) as assessed in the 4-Day Suppressive Test. **(A)** CYP and LM against the lumefantrine resistant (LM[R]) and wildtype parasites (WT). **(B)** CYP and PQ against the piperaquine resistant (PQ[R]) and wildtype (WT) parasites. The results are presented as percentage (%) drug killing of parasites in at least five mice per dose.

Fig 2: Effect of cyproheptadine (Cyp) on lumefantrine (LM) against the lumefantrine resistant (LM[R]) parasites as assessed in the 4-Day Suppressive Test. (A) Impact of varying concentration of cyproheptadine on LM activity against LM[R] parasites **(B)** Cyproheptadine restores LM potency against LM[R] parasites independent of synergy and additive effects. The results are presented as percentage (%) drug killing of parasites in at least five mice per dose.

Fig 3: Impact of probenecid (Prob), Verapamil (Ver), or Cyproheptadine (Cyp) on lumefantrine (LM) activity as assessed in the 4-Day Suppressive Test. (A) Probenecid, verapamil and cyproheptadine restores LM potency against the piperaquine resistant (PQ[R]) parasites **(B)** Probenecid or verapamil fails to restore LM potency against lumefantrine resistant (LM[R]) parasites. The results are presented as percentage (%) drug killing of parasites in at least five mice per dose.

Fig 4: Gene expression profiles of transporters and enzymes: Transporters: chloroquine resistance transporter (CRT), multidrug resistance protein (MDR1), multidrug resistance-associated protein 2 (MRP2). Enzymes**:** Plasmepsin IV (PMIV), Plasmepsin IX (PMIX), Plasmepsin X (PMX) or phosphatidylinositol 3-kinase (PI3K), pantothenate kinase (PANK), ferredoxin NADP-reductase (FNR), cysteine desulfurase (SUFS), as measured from cDNA amount derived from $5\,\mu\text{g/}\mu\text{l}$ of total RNA isolated from piperaquine resistant clones (PQR) relative to the wild-type piperaquine sensitive (PQS). The significance of differential gene expression from a mean of three independent experiments was computed using the Mann–Whitney U test with p-value set at 0.05.

Fig 5: PCR genotyping of the SUFS-knockdown (KD) and FNR Over-expression (OE) parasites; **(A)** SUFS_KD, **(B)** FNR_OE and the wildtype (WT) control lines. The PCR amplification used three sets of primer pairs; QCR2/GW2 to confirm the presence of the vector, GT/GW1, to verify the integration of the vector into the parasite genome and QCR1/QCR2 to confirm the modification of the specific genes. The QCR1, QCR2, and GT are vector-specific primers, while GW2 and GW1 primers are standard primers. **(C)** Expression profiles in ferredoxin NADP-reductase (FNR), cysteine desulfurase (SUFS) as measured from cDNA amount derived from $5\,\mu g$/ul of total RNA isolated from transgenic parasites (PQ^R_FNR_OE and PQ^R_SUFS_KD) relative to the drug-resistant parent piperaquine resistant parasites (PQ^R). The differential expression from a mean of three independent experiments was significantly different for FNR ($p<0.0046$), and SUFS ($p<0.03$) using Mann–Whitney U test with p-value set at 0.05.

Fig 6: The growth rates of transgenic parasites: Average growth rate profiles of PQ^R_SufS_KD or PQ^R_FNR_OE relative to the wild type drug-sensitive (WT), and the piperaquine resistant (PQ^R) parasites as determined on day 1, 2, 3, day 4, and day 7 post parasite inoculation.

Fig 7: Impact of overexpressing FNR gene and deletion of SufS gene on drug response profiles as assessed in the 4-Day Suppressive Test. (A) The activity profile of LM alone against piperaquine-resistant parasites (PQ[R]) and transgenic parasites with an overexpressed FNR gene (PQ[R]_FNR_OE) and parasites with SufS knock down (PQ[R]_SufS_KD). **(B)** Impact of overexpressing FNR gene in the PQ[R] on the activity of LM in combination with probenecid (Prob), verapamil (Ver), or cyproheptadine (Cyp). **(C)** Impact of knocking down the SufS gene in the PQ[R] on the activity of LM in combination with Prob, Ver, or Cyp. The results are presented as percentage (%) drug killing of parasites from at least five mice per dose.

Fig 8: **Drug binding profiles on target proteins; FNR and SUFS: (A)** Ligand best docking modes with respective binding energies towards FNR and SUFS: LM exhibited a high binding affinity to the FNR and SUFS proteins. The binding geometries for docking of a single ligand with a single receptor as modelled using the Autodock Vina. The binding energies for LM to the FNR and SUFS were the lowest, suggesting a better affinity to the protein. Cyproheptadine yielded higher binding affinity to both FNR and SUFS as compared to the other RA, verapamil, and probenecid **(B)** Interaction of LM with the FNR and SUFS. **TOP:** The crystallographic structure of (1) FNR and (2) SUFS in complex with LM as visualised using PyMOL. **Bottom:** 2D: (1) LM-FNR and (2) LM-SUFS interaction with the oxygen atoms shown in red, nitrogen atoms shown in blue, while hydrogen bonds are shown as olive-green dotted parasites. The interaction plots were generated using LigPlot+.

Table 1: Effects of cyproheptadine (Cyp) or lumefantrine (LM) alone or in combination against the Wildtype (WT) *Plasmodium berghei* ANKA as assessed in the 4-Day-Test. Results presented as percentage (%) parasite killing ± SD (standard deviation) of five mice per dose

Drug	Dose (mg/kg)	% chemo-suppression ± SE
Cyp	5	4.44 ± 1.65
Cyp	2.5	5.32 ± 4.12
LM	0.375	17.12 ± 4.12
LM	5	100.0
LM + Cyp	0.375 + 5	20.77 ± 5.25
LM + Cyp	0.375 + 2.5	5.93 ± 2.57

Table 2: Effect of verapamil (Ver) and probenecid (Prob) alone against piperaquine resistant (PQ[R]) and lumefantrine resistant (LM[R]) *Plasmodium berghei* ANKA parasites as assessed in the 4-Day Test. Results presented as percentage (%) parasite killing $\pm$ SD (Standard Deviation) of five mice per dose.

Reversing agent	Dose (mg/kg)	% chemo-suppression $\pm$ SE (LM[R])	% chemo-suppression $\pm$ SE (PQ[R])
Probenecid (Prob)	400	8.44 $\pm$ 3.07	8.03 $\pm$ 1.86
	200	0.00	4.06 $\pm$ 2.75
Verapamil (Ver)	50	0.00	0.81 $\pm$ 1.31
	25	0.00	0.78 $\pm$ 1.56

Table 3: Activity profile of piperaquine (PQ) alone against the wild type (WT) and PQ-resistant (PQ[R]) *Plasmodium berghei* ANKA or PQ in combination with probenecid (Prob), verapamil (Ver), or cyproheptadine (Cyp) against the PQ[R] parasites assessed in the 4-day Test. Results presented as percentage (%) parasite killing ± SD (standard deviation) of five mice per dose

Drugs	Dose (mg/kg)	% parasite killing± SE (PQ[R])	% parasite killing ± SE (WT)
PQ	6	0.00	98.69± 1.06
PQ	1.5	nd	57.97±9.61
PQ	75	8.39±3.85	nd
Prob + PQ	400 + 75	0.00	nd
	200 + 75	0.00	nd
Ver + PQ	50 + 75	0.00	nd
	25 + 75	0.00	nd
Cyp + PQ	5.0 + 75	14.18±1.77	nd
	2.5 + 75	8.71±3.85	nd

Table 4: Validation of the modelled 3D structures of the *Plasmodium berghei* ferredoxin NADP+ reductase (*PbFNR*) and *Plasmodium berghei* cysteine desulfurase (*PbSUFS*) proteins as determined using PROSA-web server.

Protein	Gene ID	Z-Score	score Averaged 3D-1D score > = 0.2 (%)	Residues in most favored regions (%)	Residues in additional allowed regions (%)	Residues in disallowed regions (%)
PbFNR	PBANKA_112210	-8.49	86.21	85.9	13.2	0.4
PbSUFS	PBANKA_061430	-8.67	75.11	82.2	14.1	1.6

Fig S1: Gene expression profiles of transporters and enzymes in lumefantrine resistant Plasmodium berghei ANKA**:** Transporters: chloroquine resistance transporter (CRT), multidrug resistance protein (MDR1), multidrug resistance-associated protein 1 (MRP2). Enzymes: Plasmepsin IV (PMIV), Plasmepsin IX (PMIX), Plasmepsin X (PMX) or phosphatidylinositol 3-kinase (PI3K), pantothenate kinase (PANK), ferredoxin NADP-reductase (FNR), cysteine desulfurase (SUFS), as measured from cDNA amount derived from $5\,\mu g$/µl of total RNA isolated from lumefantrine resistant clones (LMR) relative to the wild type lumefantrine sensitive (LMS). The differential expression from a mean of three independent experiments were significantly different for CRT ($p<0.0013$), MDR1 ($p<0.039$), MRP2 ($p<0.001$), FNR ($p<0.0026$), PM9 ($p<0.023$), PM10 ($p<0.008$), PANK ($p<0.046$), and PI3K ($p<0.013$) but not significantly different for SUFS ($p<0.095$), PM4 ($p<0.56$) using Mann–Whitney U test with p-value set at 0.05.

Table S1: Oligonucleotide primers sequences used to measure the transcriptional level profiles of chloroquine resistance transporter (*crt*) multidrug-resistant 1 (*mdr1*), multidrug resistance-associated protein 2 (*mrp2*), pantothenate kinase (*pank*), ferredoxin NADP$^+$-reductase (*fnr*), cysteine desulfurase (*sufs*), Plasmepsin IV (*pmiv*), Plasmepsin IX (*pmix*), Plasmepsin X (*pmx*) or phosphatidylinositol 3-kinase (*pi3k*) with β-actin I as the housekeeping gene using Maxima SYBR Green chemistry in qPCR.

Name	Primer sequence (5' – 3')	Position	Tm
Pbcrt – Forward	TTA GTA TGA TTG GTC TCA CAA G	431 - 452	56.4° C
Pbcrt – Reverse	TGA TTG AGT TTT CAC CTT GTG TT	637 - 615	57.6° C
Pbmdr1 – Forward	GCA AGT TAT TGT GGA GAA AGT GT	721 - 743	59.2° C
Pbmdr1 – Reverse	CCA CTA CCG TAA TTC AAA GTT T	926 – 905	56.4° C
Pbmrp2 – Forward	ACT GAT AAT TGA TAT GGA GGT AAC	603 - 626	58.3° C
Pbmrp2 – Reverse	TTG GAT GTG CCA TGT GTA TTA TT	797 - 775	57.6° C
Pbpank – Forward	GAA GTT TGT AGA AAA GAT GAA ATG	872 - 895	56.6° C
Pbpank – Reverse	AAT CAT CAA AAG AAA TTG TCC CAA	1057 - 1034	56.6° C
Pbfnr – Forward	ATT AGT TAG GGA AAA TGC AAA GC	249 - 271	57.6° C
Pbfnr – Reverse	TAA CAA TTA CCT GAC GAT ATA GAA	479 – 456	56.6° C
Pbpi3k – Forward	AAC CGT TTT CAC CTC CTA TGA A	4985 - 5006	58.4° C
Pbpi3k – Reverse	ATT CAG GTG CCG TTT TCA TAT C	5182 - 5161	58.4° C
Pbpmiv– Forward	TTC ATT TAT CCT TGA TAC GGG T	459 – 480	56.4° C
Pbpmiv – Reverse	TTG TAT GGT ATT GTG TGA TGT C	674 – 653	56.4° C
Pbpmix – Forward	AAC AGC TGC AAG ATA GTC AAT AT	602 – 624	57.6° C
Pbpmix – Reverse	AAT ATC TAA ATT AGT TCG TGG GGT	807 – 784	58.3° C
Pbpmx – Forward	AAT GCT TCA GGA AAT GAC ATA AAC	319 - 342	58.3° C
Pbpmx – Reverse	AAC TAC AGA TTT ATC AAC AAC TTG	564 – 541	56.6° C
Pbsufs – Forward	TTG TGA TAG TTC AGA TGA GAT TAT	396 – 419	56.6° C
Pbsufs – Reverse	ATT TCG TAT ATT TTC TTG CCA TGG	564 – 541	58.3° C
*Pbβ-actin I_* Forward	TGT TCC AGC AAT GTA TGT AGC A	387-408	58.4° C
*Pb β-actin I_* Reverse	AGA TGT CGA AAA TCC ATA TCC T	612-591	56.4° C

Table S2: Oligonucleotide primer sequences used for genotyping of the transgenic parasite lines. (A) PQ^R_SUFS_KD transgenic parasites. (B) PQ^R_FNR_OE transgenic parasites.

2A

Purpose	Primer pair	Primer Sequence
Vector integration	GT	TGT GCT CCC CTT TTC GCA C
	GW1	CAT ACT AGC CAT TTT ATG TG
Vector resistance	QCR2	TGC TGT GCA ACA TGT GAG TC
marker cassette	GW2	CTT TGG TGA CAG ATA CTA C
WT locus	QCR1	AGC CAG ACA AAA TGC GCA CA
	QCR2	TGC TGT GCA ACA TGT GAG TC

2B

Purpose	Primer pair	Primer Sequence
Vector integration	GT	AGC GGG TTG TAA AAA TGG GAA
	GW1	CAT ACT AGC CAT TTT ATG TG
Vector resistance	QCR2	TTC CAC ATG ATT GCC CTT CT
marker cassette	GW2	CTT TGG TGA CAG ATA CTA C
WT locus	QCR1	ACA TTT TAT TCT GCA GAG TCG CT
	QCR2	TTC CAC ATG ATT GCC CTT CT