Manuscript Title

Neuroprogesterone synthesis in hypothalamic astrocytes is associated with a pubertal

increase in membrane ERα levels

Abbreviated Title

Estrogen signaling in astrocytes across puberty

Mohr, M.A.[1], Falcy, B.A.[1], Laham, B.J.[2], Wong, AM[1], Micevych, P.E[1]

1: UCLA DGSOM Dept of Neurobiology, 650 Charles E Young Dr. S, Los Angeles, CA 90095,

Laboratory of Neuroendocrinology of the Brain Research Institute, United States of America

2:Current address: Princeton Neuroscience Institute and Department of Psychology,

Princeton University, Princeton, New Jersey 085444.

Author Contributions: Each author must be identified with at least one of the following:

MM and PM Designed research; MM, BF, BL, and AM performed research and analyzed

data; MM, BF, and PM wrote the paper.

Correspondence should be addressed to Margaret Mohr, mmohr@mednet.ucla.edu

Acknowledgements: The authors wish to thank Carolina Thörn Perez for technical assistance

during calcium imaging. This research was supported by NIH National Center for Advancing

Translational Science (NCATS) UCLA CTSI grant number UL1TR001881 (MAM), and NIH

Eunice Kennedy Shriver National Institute Of Child Health & Human Development grant number

HD042635 (PEM).

Conflict of Interest: The authors declare no competing financial interests.

Funding sources: National Center for Advancing Translational Sciences UCLA CTSI Grant

UL1TR001881 (MAM), Eunice Kennedy Shriver National Institute Of Child Health & Human

Development grant number HD042635 (PEM).

ABSTRACT

The development of estrogen positive feedback is a hallmark of female puberty. Both estrogen and progesterone signaling are required for the functioning of this neuroendocrine feedback loop but the physiological changes that underlie the emergence of estrogen positive feedback remain unknown. Only after puberty does estradiol (E2) facilitate progesterone synthesis in female hypothalamic astrocytes (neuroP) (Mohr et al. 2018), an event critical for estrogen positive feedback and the LH surge. We hypothesize that prior to puberty, these astrocytes have low levels of membrane estrogen receptor alpha (ERα), making them unable to respond to E2 with increased neuroP synthesis prior to puberty. To test this hypothesis, pure populations of primary astrocyte cultures were derived from female mice at three different stages of development: pre-puberty (postnatal week 3), pubertal onset (week 5), and post-puberty (week 8). Hypothalamic astrocyte responses were measured after treatment with E2. Hypothalamic astrocytes increased progesterone synthesis across pubertal development. Prior to puberty, mERα expression was low in hypothalamic astrocytes, but expression increased across puberty. The increase in mERα expression in hypothalamic astrocytes also corresponded with an increase in caveolin-1 protein, PKA phosphorylation, and a more rapid $[Ca^{2+}]_i$ flux in response to E2. Together, these results indicate that increased mERα in hypothalamic astrocytes contributed to the post-pubertal response to E2 that results in neuroP synthesis, critical for ovulation.

SIGNIFICANCE STATEMENT

Hypothalamic astrocytes, when exposed to estradiol, make progesterone that is necessary for estrogen positive feedback during adulthood in the female rodent. However, little is known about what cellular changes occur during puberty that allow for the estradiol facilitation of progesterone synthesis. In this study, we compared cell excitability, progesterone synthesis, and levels of membrane estrogen receptor in hypothalamic astrocytes derived from pre-, mid-, and post-pubertal female mice to characterize their maturation that allows them to increase progesterone synthesis facilitating estrogen positive feedback. This study suggests that hypothalamic astrocytes acquire the necessary cellular machinery during puberty to enable E2-facilitated progesterone synthesis.

INTRODUCTION

Puberty is a complex developmental stage that elicits a barrage of neuroendocrine changes

leading to reproductive maturation. In females, the development of reproductive circuits during

puberty culminates in the brain responding positively to ovarian estradiol (E2), triggering the

luteinizing hormone (LH) surge that underlies ovulation. There is mounting evidence that

alongside E2, brain-derived progesterone (neuroP) is needed for full activation of estrogen

positive feedback: 1) astrocyte-derived neuroP is crucial for fully stimulating kisspeptin release

in an in vitro model of adult anterior hypothalamic kisspeptin neurons (Mittelman-Smith et al.

2015, Mittelman-Smith et al. 2018), 2) blocking neuroP synthesis arrests estrous cyclicity and

decreases the LH surge (Micevych et al. 2003, Micevych et al. 2008), and 3) deleting the

nuclear progesterone receptor (PGR) from kisspeptin neurons diminishes the LH surge and

decreases fertility in mice (Stephens et al. 2015). What leads to the development of neuroP

synthesis during puberty remains unknown.

Steroid hormones, including E2 and P4, act directly in the nucleus as well as at the

membrane, initiating non-classical signaling that elicits intracellular changes that can affect

transcription (Mittelman-Smith et al. 2017). E2 membrane-initiated signaling elicits elevated

intracellular calcium levels and rapid P4 synthesis in hypothalamic astrocytes (Micevych et al.

2007), which are the source of neuroP. E2-facilitated neuroP synthesis in astrocytes is initiated

via membrane-associated ER-alpha (mERα) transactivation of metabotropic glutamate receptor

1a (mGluR1a; (Micevych et al. 2017)). The shuttling of mERα to the plasma membrane and

subsequent coupling to an mGluR subtype is regulated by a class of lipid raft proteins, called

caveolins (Luoma et al. 2008). In a process necessary for neuroP synthesis, caveolin-1 shuttles

full-length mERα to the plasma membrane in association with mGluR1a. E2 binding initiates

excitatory downstream signaling cascades, including IP_3 receptor-dependent intracellular Ca^{2+}

flux (Chaban et al. 2004) and the phosphorylation of PKA (Chen et al. 2014). Activated

steroidogenic acute regulatory protein (StAR) and translocator protein (TSPO) mediate the

transport of cholesterol into the mitochondria where P450scc converts it into pregnenolone.

Finally, 3β-hydroxysteroid dehydrogenase (3β-HSD) converts pregnenolone into P4. Rapid E2

signaling in female hypothalamic astrocytes, eliciting intracellular Ca^{2+} flux and subsequent P4

synthesis, occurs within minutes of E2 exposure (Kuo et al. 2010).

Like the LH surge, E2-facilitated neuroP synthesis is observed solely in post-pubertal

females. Indeed, astrocytes derived from the adult female hypothalamus respond to E2

stimulation with an increase in neuroP synthesis. Recently, Mohr et al. 2018, formally tested

whether systemic administration of E2 to pre- and post-pubertal ovariectomized rats induced

neuroP in the hypothalamus. Only post-pubertal female rats had elevated hypothalamic neuroP

levels (Mohr et al. 2018). Moreover, neuroP levels were elevated on proestrus in gonadally

intact female rats. Thus, in vivo and in vitro studies suggest that during puberty, hypothalamic

astrocytes develop the cellular machinery to respond to E2 with neuroP synthesis. To explore

this hypothesis, the present study examined neuroP synthesis, and the sequelae of E2 action in

astrocytes that are associated with neuroP synthesis: E2-induced $[Ca^{2+}]_i$, levels of membrane

ERα and expression of proteins associated with steroidogenesis at three specific time points in

pubertal development (postnatal week 3 - pre-puberty; week 5 - pubertal onset and week 8 -

post-puberty).

MATERIALS AND METHODS

All experimental procedures were performed in accordance with the principles and procedures

of the National Institutes of Health Guide for the Care and Use of Laboratory Animals and

approved by University of California, Los Angeles.

Primary astrocyte culture.

Primary hypothalamic astrocyte cultures were prepared from a total of 93 C57Bl/6 female mice

(Jackson) consisting of 3 ages: postnatal day (PND) 23 (n = 34), 35 (n = 30), and 60 (n = 29).

Cultures were generated from cohorts containing 8-12 animals (n = 4/age) at a time. In some

cases where data are represented as fold change from PND 23 control, the cohort consisted of

a PND 23 group with either a PND 35 or PND 60 group. Hypothalamic blocks were dissected

with the following boundaries: rostral extent of the optic chiasm, rostral extent of the mammillary

bodies, lateral edges of the tuber cinereum, and the top of the third ventricle. The tissue was

minced, dissociated by enzymatic treatment with 2.5% trypsin solution (Invitrogen) with DNase I

(Sigma #D4263) and incubated in a 37°C water bath where it was agitated every 10 min for 30

min. The trypsin was removed and 5 ml DMEM/F12 (Mediatech), supplemented with 10% fetal

bovine serum (FBS) (Invitrogen), 1% penicillin (10000 IU/mL)/streptomycin (10000 µg/mL)

solution (Mediatech), and 2 mM L-glutamine (Hyclone) was added. The tissue was triturated 6

times with a 5 ml pipette. Cells were plated in T-75 flasks with 10 ml DMEM/F12 (Mediatech),

supplemented with 20% fetal bovine serum (FBS) (Invitrogen), 1% penicillin (10000

IU/mL)/streptomycin (10000 µg/mL) solution (Mediatech), and 2 mM L-glutamine (Hyclone) and

maintained at 37°C with 5% CO2 for 3 days before the medium was replaced. After 1 week,

FBS concentration was decreased to 15%. After 2 weeks, FBS concentration was decreased to

10%. Astrocyte cultures were grown to confluence and shaken on an orbital shaker at 200 rpm

for 4 hours to eliminate oligodendrocytes and microglia, yielding cultures with >95% glial

fibrillary acid protein-immunoreactive astrocytes. Before experimental manipulations, astrocytes

were steroid- starved for 2 hours with 5% charcoal-stripped dextran-treated FBS in phenol-free

DMEM/F12 medium. Cultures were then treated with vehicle dimethylsulfoxide (DMSO; Sigma-

Aldrich, cat# D2650) or 17β-estradiol (1 or 10 nM; Sigma-Aldrich, cat #: E-8875) in steroid-free

media (see above) for 1 hour at 37°C. After E2 treatment, media was collected for P4 analysis

and cells were collected for immunoblotting.

Progesterone Extraction and ELISA.

NeuroP was extracted from conditioned astrocyte media using diethyl ether and frozen at -20°C.

Frozen steroids were reconstituted in assay buffer (Cat# 80-0010). Enzo Scientific Progesterone

ELISA kit (cat #ADI-900-011) was used to measure [P4], according to manufacturer's

instructions. Positive (blank media spiked with 50 ng/ml P4) and negative controls (blank media

alone) were run with the samples. Standard curve ($R2=0.996$). The detectable range of P4 was

8.57 - 500 pg/ml. Inter-assay coefficients of variation were between 2.7% and 8.3%, and intra-

assay coefficients of variation were between 4.9% and 7.6%.

Calcium Imaging.

Astrocytes (375,000-570,000) were plated onto poly-d-lysine (0.1 mg/ml; Sigma-Aldrich)-coated

glass coverslips and 1-2 days later pretreated for 2 h with 5% charcoal-stripped FBS DMEM/F-

12 medium. Before imaging, cells were loaded with the calcium indicator Fluo 4-AM (3 µM;

Thermo-Fisher; F14201) dissolved in HBSS with 0.3% DMSO and 0.06% Pluronic F-127

(Thermo-Fisher; P3000MP) for 45 min at 37°C. HBSS was used to wash the cells and then

glass coverslips were mounted onto a 50-mm chamber insert (Warner Instruments) fixed into 60

x 15-mm cell culture dish (Dow Corning) and placed into a QE-2 quick exchange platform

(Warner Instruments) for imaging on a Zeiss Examiner D1 microscope, using an Achroplan

20/0,50 water immersion objective (Zeiss). A perfusion pump (1 ml/min, 37°C) was used to

deliver HBSS before and after application of E2 (100 pM diluted from DMSO-stocks in HBSS,

17β-estradiol, Sigma-Aldrich, cat # E-8875) or ATP (30 uM diluted in HBSS; Sigma Aldrich, cat#

A2383). Fluo-4AM fluorescence was excited with the 488-nm argon laser, and emission signals

above 505 nm were collected. Videos consisted of 1200 frames (512 x 512) taken, on average,

every 161.08 ms (6.2 Hz, +/- 0.4 ms). Calcium flux was measured as Δ F/F, or the change in

fluorescence intensity after E2 or ATP treatment over the fluorescent intensity during HBSS

treatment before stimulation. Peak values were defined as the highest Δ F/F value obtained

after the perfusion with E2 or ATP. Responding cells were those whose peak values were at

least 20% above the baseline value prior to E2 or ATP delivery.

Cell Surface Biotinylation.

Astrocytes were pretreated for 2 h with 5% charcoal-stripped FBS DMEM/F-12 medium and

incubated with DMSO vehicle or in the presence 1 nM 17-estradiol (Sigma-Aldrich) in 5%

charcoal-stripped FBS DMEM/F-12 medium for 1 h at 37°C. Cells were then washed three times

with ice-cold HBSS with Ca^{2+} and Mg^{2+} buffer and incubated with freshly prepared 0.5 mg/ml

EZ-Link Sulfo-NHS-LC-Biotin (Pierce) in HBSS at 4°C for 30 min with gentle agitation. Excess

biotin reagent was quenched by rinsing the cells three times with ice-cold glycine buffer (50 mM

glycine in HBSS). Cells were scraped into 10 ml of HBSS solution, transferred into a 50 ml

conical tube, and centrifuged at 850 x g for 5 min. The pellet was resuspended in 200 µl of RIPA

Lysis Buffer (Santa Cruz Biotechnology) containing the following proteases inhibitors: 1 mM

phenylmethylsulfonyl fluoride, 1 mM EDTA, 1 g/ml pepstatin, 1 g/ml leupeptin, 1 g/ml aprotinin,

and 1 mM sodium orthovanadate (all inhibitors were from Santa Cruz Biotechnology). The cells

were homogenized by passing through a 25 gauge needle 10 times every 10 min for a total of

30 min. The cell extract was centrifuged at 14,000 rpm for 5 min at 4°C, and the protein

concentration of the supernatant was determined using the bicinchoninic acid (BCA) Assay

(Pierce). Two hundred microliters of each sample with 1500 g/ml protein concentration was

incubated with 200 µl of Immobilized NeutrAvidin Gel (Pierce) overnight at 4°C and spun for 1

min at 1000 x g. The beads were washed four times with 500 µl of radioimmunoprecipitation

assay (RIPA) buffer (Santa Cruz Biotechnology) containing the same proteases inhibitors

mentioned previously. The bound proteins were eluted with Laemmli Buffer containing BME

(950 µl Laemmli, 50 µl of BME) for 2 h at 37°C before centrifugation at 14000 rpm for 5 min.

Immunoblotting.

For Western blotting, astrocyte cultures were washed three times with 10 mL PBS buffer and collected by mechanical scraping (Corning). The samples were centrifuged (850 x g for 5 minutes), aspirated of PBS, and lysed in RIPA lysis buffer (Santa Cruz) with protease inhibitor and phosphatase inhibitor (Invitrogen). Pellets were sonicated at 3 x 10 pulses and kept on ice for 30 min before 4°C centrifugation at 14000 x g for 5 min. Protein concentrations were determined using the BCA Assay (Pierce). Protein (10 μg of whole-cell lysates; 40 μl of surface biotinylated lysates) of each sample was loaded on an 8-16% (for TSPO, 3b-HSD, and StAR) or 4-20% (for mERα, caveolin-1, and PKA) SDS precast gels (Bio-Rad) and separated by electrophoresis. Proteins were transferred onto PVDF (GE) membranes using electrophoresis at 11 V for 18 hours (Invitrogen). Because GAPDH protein changed across puberty (data not shown) a Ponceau S (0.5% Ponceau S with 1% acetic acid) stain was done for westerns using whole-cell lysates and the 45 kDa band was imaged and used as a loading control (Olesen and Auger 2005). Membranes were probed with primary rabbit or mouse monoclonal or polyclonal antibodies diluted in 5% bovine serum albumin or nonfat dry milk in Tris-buffered saline/Tween 20 (0.1% [vol/vol]) according to the suppliers' recommendations (see Table 1). Primary incubations lasted overnight, except for ERα (3 nights) and caveolin-1 (2 nights). Blots were then incubated with goat anti-rabbit or mouse IgG horseradish peroxidase (1:10,000; Vector Laboratories) for 2 hours at room temperature. Bands were visualized using an enhanced chemiluminescence (ECL) kit (Thermo Fisher Scientific). Ponceau S (whole-cell) or Flotillin (biotinylation) was used as a loading control, and immunoreactive bands were normalized to obtain the percentage of protein to Ponceau S or Flotillin ratio. Bands were immediately visualized using FluorChem E and analyzed by Alphaview software (Cell Bioscience). To normalize the data as fold change from PND23 vehicle-treated for each target, the same PND23 vehicle-treated lysates were included across blots. Samples from similar treatment groups were pooled to achieve a minimum of 10 μg total protein per western blot lane.

Reverse Transcription Polymerase Chain Reaction (PCR).

RNA was extracted from astrocytes using TRIzol (Invitrogen) reagent, according to manufacturer's protocol. Following chloroform extraction of RNA and precipitation with 100% isopropanol, RNA pellet was washed with 75% ethanol/DEPC-treated water. RNA pellets dried for 10 min at room temperature and were re-suspended in DEPC-treated water. RNA

concentration and quality was assessed using a spectrophotometer (NanoDrop 1000). RNA was
then converted to cDNA using Superscript III First Strand Synthesis Kit (Invitrogen, cat #18080-
051). cDNA was made using Oligo dTs for Tspo, Esr1, and HSD3B1, and random hexamers for
StAR using 1-2 µg RNA. The RT reaction was performed using a thermal cycler (Agilent
Technologies, Stratagene Mx3000P) with the following parameters: for Oligo dTs, 50 min at
50°C and 5 min at 85°C; for random hexamers, 10 min at 25°C, 1 hr at 50°C, 10 min at 85°C.
cDNA was used immediately for qPCR or stored at -20°C for ≤ 1 month.

See Table 2 for primer information. PCR reactions were run in duplicate (if using SYBR
GreenER primers) or triplicate (if using TaqMan primers), and at least 3 duplicates/triplicates per
condition were included in each PCR assay. No template controls (NTC; cDNA reverse
transcribed without RNA) were run alongside cDNA for each primer pair in every PCR
experiment. No amplification was observed in NTC wells. PCR reactions were prepared using
2 µL of cDNA in a 20 µL reaction with SYBR GreenER qPCR SuperMix Universal (Invitrogen;
Carlsbad, CA) or Taqman Gene Expression Master Mix (Thermo Fisher, cat # 4369016).
Conditions for amplification using SYBR GreenER primers were as follows: 2 min at 50°C (UDG
incubation), 10 min at 95° C for UDG inactivation and DNA polymerase activation, then 50
cycles each (SYBR GreenER primers) or 40 cycles each (TaqMan primers) consisting of 15 sec
at 95° C and 60 sec at the annealing temperature (60°C).
241

Statistics.
Data are presented as means ±SEM. Statistical comparisons were made using GraphPad Prism
7.01 Software (La Jolla, California). One-way ANOVAs were performed to determine if mRNA
changes as a function of age. Two-way ANOVAs with Tukey's multiple comparisons post hoc
tests were performed to determine if protein levels or P4 synthesis change as a function of age
and/or estrogen treatment. Statistical significance was considered if $p \leq 0.05$.
248

RESULTS

P4 levels increased across puberty in female mouse hypothalamic astrocytes
After treating female mouse hypothalamic astrocytes with 10 nM E2 for 3 hrs, media was
collected, steroids were extracted, and neuroP was measured to determine if P4 synthesis
increases across pubertal development. Two-way ANOVA with age and hormone treatment as

independent variables and neuroP (pg/ml) as the dependent variable was performed and

revealed a significant main effect of age ($F_{(2, 16)} = 3.9$, $p = 0.0418$). Tukey post-hoc tests

comparing [neuroP] at different ages indicated a nearly significant increase in P4 from PND 23

to PND 60 ($p = 0.053$). n= media from 4 T75 flasks/grp for PND23 and PND35, n = media from

3 T75 flasks/grp for PND 60.

E2-induced $[Ca^{2+}]_i$ in hypothalamic astrocytes across puberty

To determine if the response to E2 changes during pubertal development, prepubertal and adult

female mouse hypothalamic astrocytes loaded with a calcium indicator (Fluo-4 AM) were

treated with E2 and the change in fluorescence from baseline was monitored using epi-

fluorescence. Pre-pubertal hypothalamic astrocytes exhibit both an attenuated and sustained

release of intracellular calcium in response to E2 treatment, compared with the rapid and

transient response of these cells after puberty (Figure 2B-D). Unpaired t-tests comparing peak

delta F/F indicated that adult responses to 100 pM E2 were significantly higher than the

prepubertal responses to 100 pM E2 ($t_{(20)} = 7.517$, $p < 0.0001$; Figure 2C). Using ATP as a

control for inducing intracellular calcium levels, both pre- and post-pubertal astrocytes were

capable of having rapid and equally large responses, however pre-pubertal astrocytes were

slower to recover, as indicated by their sustained response (Figure 2E).

Steroidogenic proteins

We did not observe any changes in steroidogenic protein expression across pubertal

development. Neither levels of mRNA coding for StAR, TSPO, and 3β-HSD, nor the resulting

proteins differed at the time points examined. Separate two-way ANOVAs with age and 10 nM

E2 treatment as independent variables and fold change in protein (normalized to Ponceau S)

from PND 23 vehicle-treated as the dependent variable was performed for StAR, TSPO, and

3β-HSD. StAR protein levels did not change across pubertal development in female mouse

hypothalamic astrocytes. While age significantly affected TSPO protein levels ($F_{(2,34)} = 4.127$,

$p = 0.0249$), Tukey post-hoc tests indicated a significant decrease from pre-puberty to puberty

($p = 0.0487$), whereas prepubertal and adult female astrocytes did not differ in TSPO protein

levels. E2 treatment significantly altered 3β-HSD protein levels ($F_{(1,33)} = 4.61$, $p = 0.0392$),

without any effects of age. Using Bonferroni post-hoc tests to compare the effect of E2 on 3β-

HSD expression at each age separately revealed no significant differences. One-way ANOVAs

were performed with age as the independent variable and fold change in mRNA from PND23
astrocytes as the dependent variable revealed no significant differences in levels of StAR,
TSPO, or 3β-HSD mRNA across pubertal development in female mouse hypothalamic
astrocytes in vitro.
290
ERα mRNA and membrane ERα
Esr1 mRNA levels in female mouse hypothalamic astrocytes harvested from prepubertal,
pubertal, and adult female mice were compared using PCR. No changes in Esr1 mRNA were
detected across the time points analyzed (one-way ANOVA; p = 0.105; Figure 4A).
Surface biotinylation and western blotting revealed that the protein levels of the full-length
(66kDa) ERα changed across pubertal development but not in response to 1 nM E2 treatment
(Figure 4B). Two-way ANOVA with age and hormone treatment as independent variables and
fold change from P23 vehicle-treated mERα protein level as the dependent variable revealed a
significant main effect of age on mERα levels (F(2, 55) = 4.403, p = 0.0168). Tukey post-hoc
tests comparing mERα at different ages indicated that mERα protein concentration in female
mouse hypothalamic astrocytes increased 3.1-fold from pre- to mid-puberty (1.56 ± 0.17 to 4.87
± 1.51, p = 0.0316) and increased 3.2-fold from pre-puberty to adulthood (1.56 ± 0.17 to 5.02 ±
0.38, p = 0.0264). Mid-pubertal and adult astrocytes did not differ in mERα concentration (p =
0.997).
305
Caveolin-1 protein
Caveolin proteins are required for ER trafficking to the membrane (Meitzen and Mermelstein,
2011). Because ERα increased across development, we also examined the expression of
caveolin proteins. Whole-cell lysates of E2 or vehicle treated hypothalamic astrocytes at
different ages were analyzed by western blotting (Figure 5). Two-way ANOVA with age and
hormone treatment as independent variables and the fold change in caveolin-1 protein
(normalized to Ponceau S) from vehicle-treated PND 23 astrocytes as the dependent variable
revealed a significant main effect of age (F(2, 22) = 7.814, p = 0.0027). Tukey post-hoc tests
indicate a significant 3-fold increase in caveolin-1 protein from prepuberty/puberty to adulthood
(PND23 < PND60, p = 0.0064; PND35 < PND60, p = 0.0051).
316

Phosphorylated PKA increased across puberty

We did not observe any changes in steroidogenic protein levels from prepuberty into adulthood,

but there was a developmental increase in neuroP levels, suggesting that the activation of

steroidogenic proteins might be increasing. Membrane ERα signaling activates PKA to

phosphorylate StAR and TSPO (Chen et al. 2014) to then increase the transport of cholesterol

into the mitochondrial matrix (Papadopoulos et al. 2006), the rate-limiting step of

steroidogenesis. Levels of phospho-PKA from whole-astrocyte lysates were normalized to total

PKA at each developmental time point (Figure 6). Two-way ANOVA, with age and hormone

treatment as independent variables and the fold change in phospho-PKA/total PKA from PND

23 vehicle-treated as the dependent variable, revealed a significant main effect of age ($F(2, 43)$

$= 5.29$, $p = 0.0088$). Tukey post-hoc tests indicate a significant ~2-fold increase in phospho-

PKA/total PKA from prepuberty/puberty to adulthood (PND23 < PND60, $p = 0.0054$; PND35 <

PND60, $p = 0.0167$).

DISCUSSION

The major findings of the present study are that as hypothalamic astrocytes respond to E2

with augmented neuroP synthesis across pubertal development, as the levels of mERα

increased. The coordinate increase in mERα levels was associated with an increase in PKA

activation and E2-induced neuroP. The current results with hypothalamic astrocytes parallel

earlier results in the female rat that demonstrate neuroP synthesis increases across puberty

(Mohr et al. 2018). Our current understanding of the female neuroendocrine circuit implicates

neuroP in augmenting E2-induced kisspeptin expression and release. This kisspeptin in turn

stimulates GnRH neurons and the released GnRH induces a LH surge. Indeed, knocking out

progesterone receptor from kisspeptin neurons attenuates the LH surge (Stephens et al. 2015).

Our finding suggests that post-pubertal female hypothalamic astrocytes are able to increase P4

compared with pre-pubertal female hypothalamic astrocytes. Our study aimed to further

elucidate this mechanism by comparing the cellular machinery involved in neuroP synthesis in

hypothalamic astrocytes harvested from female mice at different stages of pubertal

development.

Both pre- and post-pubertal female mouse hypothalamic astrocytes responded to E2

administration with increased $[Ca^{2+}]_i$, however their responses are different. Prepubertal

hypothalamic astrocytes exhibited a smaller but longer-lasting E2-induced intracellular calcium

release. Adult hypothalamic astrocytes, on the other hand, demonstrated a rapid, higher and

more transient $[Ca^{2+}]_i$ response to E2. Additionally, the adult astrocytes responded more

robustly to E2 than the pre-pubertal astrocytes. Whether these differences have functional

significance remains unknown. When Kuo et al., (2010) compared $[Ca^{2+}]_i$ responses to E2 in

adult male astrocytes with responses in adult female astrocytes a similar effect was observed –

males had an attenuated $[Ca^{2+}]_i$ response. Moreover, male astrocytes, which synthesize a low

level of neuroP, could not be stimulated by E2 to increase their neuroP synthesis. The response

to E2 is similar in prepubertal female astrocytes and adult male astrocytes, further suggesting

that adult female hypothalamic astrocytes undergo pubertal changes to allow them to

synthesize neuroP. The present experiments revealed that only in post-pubertal astrocytes was

PKA activated, which is needed for the E2 augmentation of neuroP synthesis (Chen et al.

2014). These findings suggest that intracellular signaling mechanisms in pre-pubertal astrocytes

are different from those in adult females. One possibility is that mER, through which E2 signals,

changes during development.

Estradiol signaling at the membrane is dependent on the mGluR transactivated by mER

(reviewed in (Meitzen et al. 2017)). For E2- facilitated neuroP synthesis to occur, the

mERα/mGluR1a complex is trafficked to the cell membrane by caveolin-1 (Christensen and

Micevych 2012) and signals through an excitatory Gq to activate PLC and the release of Ca^{2+}

from intracellular stores (Chaban et al. 2004). Conversely, inhibitory E2 signaling is mediated

by mGluR2/3 activation of a Gi (Meitzen and Mermelstein 2011). Indeed, it has been shown that

inhibitory E2 signaling in the hypothalamus was mediated by mGluR2/3 that was transactivated

by a 52 kDa splice variant of ERα missing exon 4, the ERαΔ4 (Wong A.M., Scott A.K., Johnson

C.S., Mohr M.A., Mittelman-Smith M.A., Micevych P.E, unpublished observations) . In the

present study, estradiol was able to elicit an attenuated $[Ca^{2+}]_i$ in prepubertal astrocytes,

perhaps due to the low-levels of full-length mERα. In these astrocytes, E2 could be signaling

through an entirely different membrane-associated receptor, GPR30, whose signaling can

mediate rapid E2 effects (Kuo et al. 2010). As mERα and caveolin-1 increase with

development, the E2 increase in hypothalamic neuroP synthesis also increases.

We quantified mRNA and proteins levels of steroidogenic proteins to assess whether our

observed developmental changes in neuroP synthesis were due to transcriptional or

translational modulation. We observed that only protein levels of TSPO decreased significantly

from pre-puberty to puberty, with no change seen in response to E2 treatment. TSPO's role in

steroidogenesis remains controversial, but TSPO knockouts have implicated this protein in

hormone-induced steroidogenesis (Fan et al. 2015). However, the relatively weak induction of

TSPO, and lack of effect on StAR or 3β-HSD suggest that the post-pubertal responsiveness to

E2 is not likely attributable to up-regulation of steroidogenically relevant proteins. Given that

PKA phosphorylation significantly increased post-pubertally, and that PKA phosphorylation is

necessary for the phosphorylation of TSPO and StAR, we predict that TSPO and StAR

phosphorylation activation should increase post-pubertally, which would increase neuroP

synthesis.

In the present study, the developmental effects on hypothalamic astrocytes were

stronger than the effects of E2 stimulation, suggesting that these cells undergo dramatic change

during puberty. One such change was that the sensitivity to E2 that resulted in neuroP synthesis

increased across pubertal development in hypothalamic astrocytes. While prepubertal

astrocytes respond rapidly to E2, albeit weakly, only adult astrocytes responded robustly and

rapidly to E2 that resulted in PKA phosphorylation and neuroP synthesis. When neuroP is

measured in hypothalamic tissue, either in gonadectomized and EB-replaced or gonad-intact

rats, E2-facilitated P4 synthesis occurred only post-pubertally (Mohr et al. 2018), and could be

explained by the presence of ERα on the membrane only after puberty in hypothalamic

astrocytes.

Conclusion

In the present study hypothalamic astrocytes harvested from female mice at different ages were

used to compare estradiol responsiveness across pubertal development. The response to E2 in

prepubertal hypothalamic astrocytes resulted in $[Ca^{2+}]_i$ increase but due to low amounts of full-

length mERα, PKA was not phosphorylated and P4 synthesis remained low. The response to

E2 in post-pubertal hypothalamic astrocytes was rapid, resulting in $[Ca^{2+}]_i$ flux mediated by full-

length mERα, which led to PKA phosphorylation and neuroP synthesis. The expression of full-

length mERα after puberty in hypothalamic astrocytes likely underlies the female's distinctly

post-pubertal response to E2 to produce neuroP, which is critical for ovulation.

References

Chaban, V. V., A. J. Lakhter and P. Micevych (2004). "A membrane estrogen receptor mediates
intracellular calcium release in astrocytes." Endocrinology **145**(8): 3788-3795.

Chen, C., J. Kuo, A. Wong and P. Micevych (2014). "Estradiol Modulates Translocator Protein
(TSPO) and Steroid Acute Regulatory Protein (StAR) via Protein Kinase A (PKA)
signaling in Hypothalamic Astrocytes." Endocrinology **155**(8): 2976-2985.

Christensen, A. and P. Micevych (2012). "CAV1 siRNA reduces membrane estrogen receptor-
alpha levels and attenuates sexual receptivity." Endocrinology **153**(8): 3872-3877.

Fan, J., E. Campioli, A. Midzak, M. Culty and V. Papadopoulos (2015). "Conditional
steroidogenic cell-targeted deletion of TSPO unveils a crucial role in viability and
hormone-dependent steroid formation." Proceedings of the National Academy of
Sciences **112**(23): 7261.

Kuo, J., N. Hamid, G. Bondar, E. R. Prossnitz and P. Micevych (2010). "Membrane estrogen
receptors stimulate intracellular calcium release and progesterone synthesis in
hypothalamic astrocytes." Journal of Neuroscience **30**(39): 12950-12957.

Luoma, J. I., M. I. Boulware and P. G. Mermelstein (2008). "Caveolin proteins and estrogen
signaling in the brain." Molecular and cellular endocrinology **290**(1-2): 8-13.

Meitzen, J., K. A. Britson, K. Tuomela and P. G. Mermelstein (2017). "The expression of select
genes necessary for membrane-associated estrogen receptor signaling differ by sex in
adult rat hippocampus." Steroids.

Meitzen, J. and P. G. Mermelstein (2011). "Estrogen receptors stimulate brain region specific
metabotropic glutamate receptors to rapidly initiate signal transduction pathways." J
Chem Neuroanat **42**(4): 236-241.

Micevych, P., K. Sinchak, R. H. Mills, L. Tao, P. LaPolt and J. K. Lu (2003). "The luteinizing
hormone surge is preceded by an estrogen-induced increase of hypothalamic
progesterone in ovariectomized and adrenalectomized rats." Neuroendocrinology **78**(1):
29-35.

Micevych, P., K. K. Soma and K. Sinchak (2008). "Neuroprogesterone: Key to estrogen positive
feedback?" Brain Res Rev **57**(2): 470-480.

Micevych, P. E., V. Chaban, J. Ogi, P. Dewing, J. K. Lu and K. Sinchak (2007). "Estradiol
stimulates progesterone synthesis in hypothalamic astrocyte cultures." Endocrinology
148(2): 782-789.

Micevych, P. E., P. G. Mermelstein and K. Sinchak (2017). "Estradiol Membrane-Initiated
Signaling in the Brain Mediates Reproduction." Trends Neurosci **40**(11): 654-666.

Mittelman-Smith, M. A., L. M. Rudolph, M. A. Mohr and P. E. Micevych (2017). "Rodent Models
of Non-classical Progesterone Action Regulating Ovulation." <u>Front Endocrinol</u>
<u>(Lausanne)</u> **8**: 165.

Mittelman-Smith, M. A., A. M. Wong, A. S. Kathiresan and P. E. Micevych (2015). "Classical and
membrane-initiated estrogen signaling in an in vitro model of anterior hypothalamic
kisspeptin neurons." <u>Endocrinology</u> **156**(6): 2162-2173.

Mittelman-Smith, M. A., A. M. Wong and P. E. Micevych (2018). "Estrogen and Progesterone
Integration in an in vitro Model of RP3V Kisspeptin Neurons." <u>Neuroendocrinology</u>
106(2): 101-115.

Mohr, M. A., A. M. Wong, R. J. Tomm, K. K. Soma and P. E. Micevych (2018). "Pubertal
development of estradiol-induced hypothalamic progesterone synthesis." <u>Horm Behav</u>.

Olesen, K. M. and A. P. Auger (2005). "Sex differences in Fos protein expression in the
neonatal rat brain." <u>J Neuroendocrinol</u> **17**(4): 255-261.

Papadopoulos, V., M. Baraldi, T. R. Guilarte, T. B. Knudsen, J. J. Lacapere, P. Lindemann, M.
D. Norenberg, D. Nutt, A. Weizman, M. R. Zhang and M. Gavish (2006). "Translocator
protein (18kDa): new nomenclature for the peripheral-type benzodiazepine receptor
based on its structure and molecular function." <u>Trends in pharmacological sciences</u>
27(8): 402-409.

Stephens, S. B., K. P. Tolson, M. L. Rouse, Jr., M. C. Poling, M. K. Hashimoto-Partyka, P. L.
Mellon and A. S. Kauffman (2015). "Absent Progesterone Signaling in Kisspeptin
Neurons Disrupts the LH Surge and Impairs Fertility in Female Mice." <u>Endocrinology</u>
156(9): 3091-3097.

 Figure Legends

Figure 1. E2-facilitated P4 synthesis increases in female mouse hypothalamic astrocytes
across puberty. Hypothalamic astrocytes harvested from 60 day-old female mice synthesized
significantly more progesterone compared with astrocytes harvested from 23 day-old female
mice after 10 nM E2 treatment for 3 hrs. Data indicate the amount of progesterone (pg/ml)
present in the media of astrocytes harvested from female mice at different stages of pubertal
development ± SEM. n = 3-4/group. * $p < 0.05$.

Figure 2. Prepubertal and adult astrocytes respond differently to E2. A) Representative
image of a hypothalamic astrocyte loaded with calcium indicator Fluo-4 AM. Scale bar = 10 µm.
B) Representative traces of $[Ca^{2+}]_i$ in hypothalamic astrocytes harvested from postnatal day 23
(prepubertal) or 60 (adult) female mice loaded with Fluo-4 AM and treated with 100 pM E2. C)
An unpaired t-test comparing the average peak in fluorescence indicated that adult astrocytes
treated with 100 pM E2 elicited the most $[Ca^{2+}]_i$ compared with the prepubertal astrocytes. D)
Percentage of astrocytes not responding to E2 (p delta F/F < 1.2), responding with a delay (p
delta F/F > 1.2 occurring after 50 sec), and responding rapidly (p delta F/F > 1.2 occurring within
50 sec of E2 application). E) Prepubertal and adult astrocytes exhibited a robust $[Ca^{2+}]_i$ flux
when treated with 30 µM ATP. An unpaired t-test indicated no difference in peak delta F/F
between prepubertal and adult astrocytes ($p = 0.087$). Dotted lines indicate baseline
fluorescence. Sample size (n) for each condition is indicated inside bars. **** $p < 0.0001$.

Figure 3. Steroidogenic proteins were unremarkable in hypothalamic astrocytes
harvested from female mice at different stages of pubertal development. Protein and
mRNA levels of StAR, TSPO, and 3β-HSD remained relatively stable in hypothalamic astrocytes
across pubertal development. A) StAR protein, regardless of age or E2 treatment, did not
change across puberty in hypothalamic whole-astrocyte lysates. B) TSPO protein decreased
from pre-puberty to puberty, with adult levels intermediate in female mouse hypothalamic
astrocytes. C) Regardless of age, E2 treatment increased levels of 3β-HSD protein in female
mouse hypothalamic astrocytes. D-F) StAR, TSPO, and 3β-HSD mRNA levels did not change in
hypothalamic astrocytes harvested from prepubertal, pubertal, or adult female mice. Data are
represented as fold change in protein (normalized to Ponceau S) or mRNA (2^-ddCt) from
postnatal day 23 vehicle-treated hypothalamic astrocytes ± SEM. Sample size (n) for each
condition is indicated within bars.

Figure 4. Esr1 mRNA levels were stable while mERα increased across pubertal
development in female mouse hypothalamic astrocytes.

A) ERα (Esr1) mRNA did not differ in hypothalamic astrocytes taken from female mice pre,
during, or post-puberty. n = RNA from individual T75 flasks, indicated within bars. B)
Independent of E2-treatment, pubertal development significantly altered membrane ERα protein
levels in surface biotinylated female mouse hypothalamic astrocytes. Full-length mERα levels
increased after puberty in surface biotinylated lysates taken from female mouse hypothalamic
astrocytes (ME of age, $p = 0.0168$). Tukey post-hoc tests indicated a significant increase in
mERα from pre-puberty to mid-puberty ($p = 0.0316$) and from pre-puberty to adulthood ($p =$
0.0264). Data are plotted as fold change from P23 vehicle mERα normalized to flotillin ± SEM.
* $p < 0.05$.

Figure 5. Caveolin-1 protein increased in adulthood in female mouse hypothalamic
astrocytes. Independent of E2-treatment, caveolin-1 protein levels significantly increased after
puberty in whole-cell lysates taken from female mouse hypothalamic astrocytes (ME of age, p =
0.003). Data are plotted as fold change in caveolin-1 protein from P23 vehicle-treated ± SEM.
**p < 0.01. Sample size (n) for each condition is indicated within bars.
529
Figure 6. PKA phosphorylation increased across puberty in female mouse hypothalamic
astrocytes. Independent of E2-treatment, levels of phosphorylated PKA protein significantly
increased after puberty in whole-cell lysates taken from female mouse hypothalamic astrocytes
(ME of age, p = 0.0088). Data are plotted as fold change in phosphorylated PKA/total PKA
protein from P23 vehicle-treated ± SEM. **p < 0.01. *p < 0.05. Sample size (n) for each
condition is indicated within bars.
536

537   Figure 1

575   Figure 2

607　Figure 3

637　Figure 4
638

661   Figure 5

707   Figure 6

754 TABLES

755 **Table 1. Antibody information**

Antibody	Manufacturer	Catalog Number	RRID	Dilution
Rabbit monoclonal anti-3βHSD	Abcam	ab150384	n/a	1:500 in 5% milk
Mouse monoclonal anti-Caveolin 1	BD Laboratories	610407	AB_397789	1:1,000 in 5% milk
Rabbit polyclonal anti-ERα (C1355)	Millipore	06-935	AB_310305	1:1,000 in 5% BSA
Rabbit polyclonal anti-Flotillin 1	Abcam	ab41927	AB_941621	1:1,000 in 5% BSA
Goat anti-Mouse HRP	Invitrogen	A16072	AB_2534745	1:10,000 in 5% milk
Goat anti-Rabbit HRP	Invitrogen	A16104	AB_2534776	1:10,000 in 5% milk
Rabbit monoclonal anti-P-PKA C	Cell Signaling	5661	AB_10707163	1:1,000 in 5% BSA
Rabbit polyclonal anti-PKA C-α	Cell Signaling	4782	AB_2170170	1:1,000 in 5% BSA
Rabbit polyclonal anti-StAR	Abcam	ab96637	AB_10678397	1:1,000 in 5% milk
Rabbit polyclonal anti-TSPO	Cell Signaling	9530	n/a	1:1,000 in 5% BSA

766   **Table 2. PCR Primer Information**

Primer	Gene Accession #	FW Primer	RV Primer	Amplicon Size (BP)	Annealing Temperature	Ref/Assay #
Hsd3b1 (SYBR Green)	NM_008293	TGG ACA AGG TAT TCC GAC CAG A	GGC ACA CTT GCT TGA ACA CAG	250	60°C	Sarraj et al., 2010
Esr1 (SYBR Green)	NM_007956	TGG GCT TAT TGA CCA ACC TAG CA	AGA ATC TCC AGC CAG GCA CAC	130	60°C	n/a Li et al., 2013
GAPDH (SYBR Green)	NM_017008	GCA CAG TCA AGG CCG AGA AT	GCC TTC TCC ATG GTG GTG AA	150	60°C	n/a Chen et al., 2006
StAR (Taqman)	NM_011485	Exon 6	Exon 7	73	60°C	Mm00441558_m1
Tspo (Taqman)	NM_009775	Exon 3	Exon 4	116	60°C	Mm00437828_m1
GAPDH (Taqman)	NM_008084	Exon 2	Exon 3	107	60°C	4352339E