How Array Design Affects SNP Ascertainment Bias

Johannes Geibel[*†], Christian Reimer[*†], Steffen Weigend[†‡], Annett Weigend[‡], Torsten Pook[*†]

& Henner Simianer[*†]

[*]University of Goettingen, Department of Animal Sciences, Animal Breeding and Genetics

Group, Albrecht-Thaer-Weg 3, 37075 Göttingen, Germany

[†]University of Goettingen, Center for Integrated Breeding Research, Albrecht-Thaer-Weg 3,

37075 Göttingen, Germany

[‡]Friedrich-Loeffler-Institut, Institute of Farm Animal Genetics, Höltystrasse 10, 31535 Neu-

stadt-Mariensee, Germany

Data Availability: Additional to the sequences already published by Malomane *et al.* (2018)
or Qanbari *et al.* (2019), sequencing reads of *Gallus varius* (GV), Rhode Island Red (RI) and
White Rock (WR) pools were uploaded to European Nucleotide Sequence Database (ENA;
https://www.ebi.ac.uk/ena/data/view/PRJEB31780) and will made publically available upon
acceptance of the paper. The study accession numbers of former and newly submitted reads
can be found in supplementary file 1. Note that Malomane *et al.* (2018) already published a
subset of the individually sequenced white layers, published in total by Qanbari *et al.* (2019).
Supplementary material can be found on figshare.

Array Design and SNP Ascertainment Bias

KEYWORDS

ascertainment bias; array design; whole genome re-sequencing; chicken

CORRESPONDING AUTHOR

Johannes Geibel
Albrecht-Thaer-Weg 3
37075 Göttingen, Germany
+49 551 3925609
johannes.geibel@uni-goettingen.de

31
32
33

ABSTRACT

Single nucleotide polymorphisms (SNPs), genotyped with SNP arrays, have become the

most widely used marker types in population genetic analyses over the last 10 years. How-

ever, compared to whole genome re-sequencing data, arrays are known to lack a substantial

proportion of globally rare variants and tend to be biased towards variants present in popu-

lations involved in the development process of the respective array. This affects population

genetic estimators and is known as SNP ascertainment bias. We investigated factors contrib-

uting to ascertainment bias in array development by redesigning the Axiom™ Genome-Wide

Chicken Array *in silico* and evaluating changes in allele frequency spectra and heterozygosity

estimates in a stepwise manner. A sequential reduction of rare alleles during the develop-

ment process was shown with main influencing factors being the identification of SNPs in a

limited set of discovery populations and a within-line selection of SNPs when aiming for

equidistant spacing. These effects were shown to be less severe with a larger discovery pan-

el. Additionally, a generally massive overestimation of expected heterozygosity for the ascer-

tained SNP sets was shown. This overestimation was 28 % higher for populations involved in

the discovery process than not involved populations for the original array. The same was

observed after the SNP discovery step in the redesign. However, an unequal contribution of

populations in the equal spacing process can mask this effect but also adds uncertainty. Fi-

nally, we make suggestions for the design of specialized arrays for large scale projects where

whole genome re-sequencing techniques are still too expensive.

                                          INTRODUCTION

Starting in the first decade of this century, the possibility of cost effectively genotyping

high numbers of Single Nucleotide Polymorphisms (SNP) for many individuals in parallel via

SNP arrays led to an increased usage of them for population genetic analyses in humans

(Novembre *et al.* 2008; Patterson *et al.* 2012), model species (Laurie *et al.* 2007; Platt *et al.*

2010), plants (Travis *et al.* 2015; Mayer *et al.* 2017) and livestock (Muir *et al.* 2008; Gibbs *et*

al. 2009; Kijas *et al.* 2009; Gautier *et al.* 2010; Qanbari *et al.* 2010; McTavish *et al.* 2013;

Malomane *et al.* 2019).

Various SNP arrays exist for humans (Perkel 2008), plants (Unterseer *et al.* 2014; Singh *et*

al. 2015) and all major livestock species (Matukumalli *et al.* 2009; Ramos *et al.* 2009;

Groenen *et al.* 2011; Boichard *et al.* 2012; Kranis *et al.* 2013; Tosser-Klopp *et al.* 2014; San-

denbergh *et al.* 2016). SNP numbers within these arrays range from 10 k SNPs (Boichard *et*

al. 2012) over approximately 50 k (Groenen *et al.* 2011; Matukumalli *et al.* 2009; Tosser-

Klopp *et al.* 2014; Singh *et al.* 2015) up to 600 k (Kranis *et al.* 2013; Matukumalli *et al.* 2009;

Unterseer *et al.* 2014). The design process of every array has an initial step of SNP discovery

in common, where SNPs are identified from existing databases and/or from a small set of

sequenced individuals. SNPs are then selected based on different quality criteria like minor

allele frequency (MAF) thresholds and platform specific design scores (Fan *et al.* 2010). Addi-

tional criteria like equidistant spacing over the genome (Kranis *et al.* 2013), overrepresenta-

tion of some areas (Boichard *et al.* 2012) or increased overrepresentation of high MAF SNPs

(Matukumalli *et al.* 2009) are applied dependent on the design intentions. In the end, draft

arrays are validated either on the discovery set itself (Ramos *et al.* 2009) and/or on a broad

set of individuals from different populations (Kranis *et al.* 2013; Fan *et al.* 2010).

In contrast to whole genome re-sequencing (WGS) data, SNP arrays often show a clear

underrepresentation of SNP with extreme allele frequencies (Nielsen 2004). As population

genetic statistics are mostly based on estimates of allele frequencies, this context leads to

biased population genetic estimators (Nielsen 2004) and is known as SNP ascertainment bi-

as.

The absence of rare alleles is mainly driven by two factors in the array design process

where SNPs are selected (ascertained) based on different requirements and decisions (Eller

2001). The first factor is a relatively small panel of individuals used for discovery of SNPs,

leading to a large proportion of globally rare variants not being selected, since they appear

monomorphic in the discovery panel (Nielsen and Signorovitch 2003; Clark *et al.* 2005). The

second factor is the across population use of arrays. Arrays are developed based on the vari-

ation within the discovery panel, thus missing variation present in distantly related individu-

als or populations (Nielsen 2004; Eller 2001). This second source of bias was shown to be of

relatively high importance for livestock studies, where arrays are usually developed for large

commercial breeds and later used to genotype diverse sets of local breeds all over the world

(McTavish and Hillis 2015; Malomane *et al.* 2018).

Besides different strategies to minimize the impact of ascertainment bias (Malomane *et*

al. 2018; Lachance and Tishkoff 2013), there are some attempts to correct the allele fre-

quency spectrum via Bayesian methods (Nielsen and Signorovitch 2003; Nielsen *et al.* 2004;

Nielsen 2004). However, those corrections highly rely on detailed statistical assumptions of

the ascertainment process (Guillot and Foll 2009; Albrechtsen *et al.* 2010) and are currently

only tested for corrections of the first source of ascertainment bias, the small discovery pan-

el (Nielsen 2004; Nielsen *et al.* 2004; Nielsen and Signorovitch 2003). In practice, detailed

information on the design process is limited (Albrechtsen *et al.* 2010) and the complexity of

the processes makes statistical models for the corrections inaccurate.

Given the complexity of modern array design processes, this study aims at highlighting

the mechanisms which promote the bias by illustrating the effects of the different steps of

the array design process on ascertainment bias, using real data in a typical setting from live-

stock sciences. For this purpose, the design process of the Axiom™ Genome-Wide Chicken

Array (www.affymetrix.com) was remodeled in a set of diverse chicken WGS data. Allele fre-

quency spectra as well as expected heterozygosity (H_{exp}) were compared to the WGS data

and the SNPs of the Axiom™ Genome-Wide Chicken Array. Finally, some recommendations

are made on how to design an array for monitoring genetic diversity.

110   **MATERIAL AND METHODS**

Populations and sequencing: The analysis is based on WGS data of a diverse set of 46 com-

mercial, non-commercial and wild chicken populations, sampled within the framework of the

projects AVIANDIV (www.aviandiv.fli.de) and SYNBREED (www.synbreed.tum.de). Commer-

cial brown (BL) and white layer (WL) populations consist of 25 individually re-sequenced an-

imals each, while the two commercial broiler lines (BR1 and BR2) include 20 individually se-

quenced animals each. For 41 populations, pooled DNA from 9 - 11 animals per population

was sequenced, while *Gallus varius* (GV) samples of only two animals were sequenced as a

pool. More detailed information about the samples can be found in supplementary file 1 and

two previously published papers, Malomane *et al.* (2018) and Qanbari *et al.* (2019). Cover-

age ranged between 7X and 10X for the individual sequences, while DNA pools were se-

quenced with 15X to 70X coverage. Paired-end sequencing was conducted on Illumina HiSeq

machines at the Helmholtz Zentrum, German Research Center for Environmental Health in

Munich, Germany.

Raw data preparation and SNP calling: Sequences were aligned to the reference genome

Gallus gallus 5.0 (UCSC 2016; Warren *et al.* 2017) and the SNP calling was conducted accord-

ing to GATK Best Practices guidelines (DePristo *et al.* 2011; van der Auwera *et al.* 2013). BWA

0.7.12 (Li 2013) was used for the alignment step, duplicates were marked using Picard

Tools 2.0.1 (Broad Institute 2015) MarkDuplicatesWithMateCigar and base qualities were

recalibrated with GATK 3.7 (McKenna *et al.* 2010) BaseQualityRecalibrator. The set of known

SNPs, necessary for base quality score recalibration, was downloaded from ENSEMBL release

87 (ENSEMBL 2016). SNPs were called for all samples separately using the GATK 3.7 Haplo-

typeCaller and later on simultaneously genotyped across samples with GATK 3.7 Geno-

typeGVCFs. Due to computational limitations, a ploidy of two instead of the higher true

ploidy of the pooled sequences had to be assumed, which led to calling slightly less rare al-

leles. However, effects of this limitation are negligible (supplementary file 2; Figure S 1).

SNP filtering was conducted using GATK 3.7 VariantRecalibrator, which filtered the called

SNPs by a machine learning approach (use of a Gaussian mixture model), which uses both, a

set of previous known (low confidence needed) and a set of highly reliable (assumed to be

true) variants as training resources (Broad Institute 2018). The source for known SNPs (pri-

or 2) provided to VariantRecalibrator was again ENSEMBL (release 87) and the SNPs of the

Axiom™ Genome-Wide Chicken Array were defined as true training set (prior 15). The algo-

rithm was trained on the quality parameters DP, QD, FS, SOR, MQ and MQRankSum. Filters

were set to recover 99 % of the training SNPs in the filtered set, which resulted in an Transi-

tion/Transversion ratio of 2.52 for known SNPs, and an Transition/Transversion ratio of 2.26

for novel SNPs. Only biallelic autosomal SNPs were used in all further analyses.

Identification of the ancestral allele: Ancestral alleles were defined using allele frequency

information from the three wild populations *Gallus gallus gallus* (GG), *Gallus gallus spadi-*

ceus (GS) and *Gallus varius* (GV) by an approach comparable to Rocha *et al.* (2014). It was

assumed that the *Gallus gallus* and *Galllus varius* species emerged from a common ancestor

and *Gallus gallus* later split into *Gallus gallus gallus* and *Gallus gallus spadiceus* subspecies.

Additionally, assuming neutral molecular evolution (Kimura 1991), the ancestral allele was

most likely the major allele within those three populations, when weighting the allele fre-

quency of *Gallus varius* twice. This procedure assigned the ancestral status to the reference

allele for 86 % of the SNPs and to the alternative allele for 14 % of the SNPs, causing close to

no change in the allele frequency spectrum (supplementary file 2; Figure S 2).

Reference Sets: Three different reference sets were defined as follows: the **unfiltered WGS**

SNPs (28.5 M SNPs), SNPs filtered using GATK 3.7 (McKenna *et al.* 2010) VariantRecalibrator

(20.9 M SNPs; **filtered WGS**) and **array SNPs** (540 k SNPs), which are the intersection of the

unfiltered SNPs and the SNPs of the Axiom™ Genome-Wide Chicken Array. The separate use

of unfiltered and filtered WGS SNPs was done to assess the effect of filtering (especially the

use of an ascertained SNP set as the true set) on ascertainment bias.

Redesigning the SNP Array: The populations were divided into four groups for the array de-

sign. Eight populations were chosen for SNP discovery (Figure 1): two non-commercial popu-

lations (LE, NH), one commercial white (WL) and three commercial brown layer lines (BL, RI,

WR), and two commercial broiler lines (BR1 and BR2). Further, the discovery set had to be

split into broilers (BR1, BR2, NH) and layers (WL, LE, BL, RI, WR) for the equal spacing step.

Even though the New Hampshire (NH) population is a dual purpose breed, it was chosen as

broiler as it is assumed that the breed has contributed to founders of contemporary com-

mercial broiler lines and shows the smallest genetic distances (Nei 1972) to the two broiler

populations in this study. From the remaining populations, 19 were randomly chosen for SNP

validation, 18 populations were used for an application of the array, and *Gallus varius* as a

different species was defined as outgroup.

Figure 1: Neighbor Joining phylogenetic tree based on pairwise Nei's standard genetic dis-
tances (Nei 1972). The tree was calculated from the filtered WGS SNPs and rooted by using
Gallus varius (GV) as   outgroup. Layer and broiler populations which form in total the dis-
covery set for the array design according to Kranis *et al.* (2013) are highlighted. The plot was
produced using the R package ape (Paradis *et al.* 2004)and the abbreviations for the popula-
tions can be found in supplementary file 1.

Based on the unfiltered SNP set, the sampling of the SNPs for an approximately 600 k

sized array was remodeled *in silico* in five consecutive steps according to Kranis *et al.* (2013),

starting from the unfiltered SNP set:

1) **SNP discovery → 10.8 M SNPs**

Discovery of SNPs fulfilling basic criteria (quality ≥ 60; MAF ≥ 0.05; cover-

age ≤ mean + three standard deviations) within the discovery populations.

2) **Cluster removal → 8.8 M SNPs**

SNP clusters with less than 4 bp invariant sites at one side of a SNP (no matter of

whether 3' or 5' direction) and less than 10 bp fixation at the other side of the SNP

were removed.

3) **Equal spacing → 2.3 M SNPs**

Reduction of SNPs to achieve approximately equidistant spacing between SNPs

within discovery populations based on genetic distances. The used algorithm was

modeled according to Kranis *et al.* (2013) and initially sets up a backbone of SNPs

segregating either in all layer populations or in all broiler populations and then fills

up the gaps while aiming for a predefined local target density of 667 equidistant

segregating SNPs/cM (linkage map taken from Groenen *et al.* (2011)) within each

population.

4) **SNP validation → 1.9 M SNPs**

Removing SNPs (~ 20 %) which were not variable in at least 8 of the 19 validation

populations.

5) **Downsampling → 600 k SNPs**

Downsampling of SNPs comparable to step 3, but without any initial backbone.

205   Keeping all exonic SNPs (annotation using Ensembl VEP 89.7; McLaren *et al.* 2016)

206    and setting the target density in broiler lines as three times the target density of

207    the layer lines.

208 **Variation of the design process:** The whole design process was repeated 50 times with pop-

209 ulations being randomly assigned to be discovery, validation or application populations,

210 while the *Gallus varius* population was always kept as the outgroup.

211  To assess the impact of the number of discovery populations on the design process, the

212 number of discovery populations was varied in additional runs from 4 to 40 randomly chosen

213 populations (while assigning the remaining populations, except *Gallus varius,* to validation

214 and application groups of equal size) with 20 random replicates for each number of discov-

215 ery populations. In a last scenario, equal spacing was varied with respect to the target densi-

216 ty (33 – 3333 SNPs/cM) with 20 independent population groupings for each target density,

217 with or without the initial backbone. As the number of SNPs from the backbone was con-

218 stant, the increase of target density led to a higher number of SNPs chosen by the algorithm

219 due to the equal spacing itself and hence the relative influence of the fixed number of com-

220 mon backbone SNPs decreased.

221 **Analyses of the results:** Per-locus-allele frequencies for individually sequenced populations

222 were estimated from genotypes, whereas the estimation for the sequenced DNA-pools was

223 based on the allelic depth. Influences on the allele frequency spectra were examined by

224 comparing density estimates of derived allele frequency spectra (unfolded frequency spec-

225 trum). Further H_{exp}, the expected heterozygosity assuming Hardy Weinberg frequencies of

226 the genotypes, for the different populations were used as summary statistics of the within

population allele frequency spectra and calculated as in equation (1), where $p_{ref;l}$ denotes

the frequency of the reference allele at locus l and L the total number of loci.

$$H_{\exp} = \frac{\sum_l 2p_{ref;l}(1-p_{ref;l})}{L} \tag{1}$$

Deviations in the estimation of H_{exp} from the various SNP sets were quantified as differ-

ences between the H_{exp} calculated from the respective SNP set and the H_{exp} calculated from

the filtered WGS SNPs relative to the H_{exp} from the filtered WGS SNPs, further called overes-

timation of H_{exp} (OHE; equation (2)). An OHE of zero means that the estimates are equal,

while an OHE of one describes doubling of the unbiased estimate.

$$OHE = \frac{H_{\exp;\,\text{SNP set}} - H_{\exp;\,\text{filtered WGS SNPs}}}{H_{\exp;\,\text{filtered WGS SNPs}}} \tag{2}$$

The effects of the population group assignments on the OHE of the random population

assignments were evaluated by pairwise comparisons of least square means (LSMEANS; cal-

culated with the R package emmeans (Lenth 2019; R Core Team 2017) by using Tukey cor-

rection for multiple pairwise contrasts) of the population groups. An underlying mixed linear

model for the estimation of LSMEANS was fitted using the R package lme4 (Bates *et al.* 2015)

as shown in equation (3), where the OHE depended on an overall mean μ, the fixed effect

of the population group $popG_i$ (i can be discovery-, validation-, application- or outgroup), a

random effect for the j[th] repetition of random population grouping ($rep_j \sim N(0, I\sigma_{rep}^2)$) and

a random error $e_{ijk} \sim N(0, I\sigma_e^2)$. The procedure is comparable to simple pairwise compari-

sons of group means, the correction by the repetition only reduces the error variance and

thus decreases the confidence intervals.

247
$$OHE_{ijk} = \mu + popG_i + rep_j + e_{ijk} \qquad\qquad (3)$$

Data Availability: Additional to the sequences already published by Malomane *et al.*

(2018) or Qanbari *et al.* (2019), sequencing reads of *Gallus varius* (GV), Rhode Island Red (RI)

and White Rock (WR) pools were uploaded to European Nucleotide Sequence Database

(ENA; https://www.ebi.ac.uk/ena/data/view/PRJEB31780). The study accession numbers of

former and newly submitted reads can be found in supplementary file 1. Note that Maloma-

ne *et al.* (2018) already published a subset of the individually sequenced white layers, pub-

lished in total by Qanbari *et al.* (2019). Supplementary material can be found on figshare.

RESULTS

Numbers of SNPs: The SNP calling identified 28.5 M biallelic autosomal SNPs from which 20.9 M SNPs passed GATK's filtering procedure. 540 k SNPs from the unfiltered WGS SNP set are also mapped on the original Axiom™ Genome-Wide 580 k Chicken Array. The remodeling of the array according to Kranis *et al.* (2013) returned 10.8 M SNPs from the discovery step, which were reduced to approximately 600 k in steps as described. Numbers of identified SNPs for the additional runs differed depending on the populations and settings used and are listed in Table S 1. Additional information about the identified number of SNPs depending on the number of discovery populations and target density as well as information about the share of SNPs of different random runs can be found from Figure S 3 – Figure S 5.

Underrepresentation of rare SNPs: A clear underrepresentation of rare SNPs in all ascertained SNP sets compared to WGS is evident from the allele frequency spectra (Figure 2). Major changes in the allele frequency spectra during the array development process were especially observed after the SNP discovery step and the equal spacing step. The SNP discovery led to an underrepresentation of rare SNPs compared to sequence data, which was intensified by the equal spacing step (Figure 2) and resulted finally in a spectrum close to the spectrum of the original array. Randomly choosing populations as discovery populations confirmed the shape of the first remodeling, where the population groups were chosen according to Kranis *et al.* (2013). As major changes in the spectra mainly occurred after the SNP discovery and equal spacing, further results will concentrate on those steps.

Figure 2: Derived allele frequency spectra for the different SNP sets. For the remodeled sets, areas show the modelling according to Kranis *et al.* (2013) while grey lines represent the 50 random population groupings.

The allele frequency spectra (Figure 3) within discovery populations, compared to the spectra over all populations, clearly showed the cutoff from the MAF 0.05 filter. After the equal spacing, they additionally revealed a higher share of common SNP than the overall spectra. In contrast, the spectra within validation- and application populations showed less pronounced peaks after the discovery step and the outgroup (*Gallus varius*) revealed fixation of most SNPs variable in the discovery populations.

Figure 3: Derived allele frequency spectra within the population groups.

Influence of number of discovery populations and target density on allele frequency spec-

tra: Not surprisingly, an increased number of discovery populations resulted in a higher

number of rare alleles after the discovery step, and thus an allele frequency spectrum with a

more pronounced peak of rare alleles (Figure 4). Apparently, the shift of the allele frequency

spectrum after the equal spacing step was dependent on the number of discovery popula-

tions, as an increase in the number of discovery populations shifted the allele frequency

spectra towards a higher proportion of alleles with a low derived allele frequency. With an

increasing number of discovery populations, the shape of the allele frequency spectra got

closer to the spectrum of the original array.

Figure 4: Derived allele frequency spectra after SNP discovery (blue) and equal spacing (red)
for varying numbers of populations in the discovery set. Different numbers of populations in
the discovery set (4 to 40) are indicated by an intensifying color gradient and only one ran-
domly picked run per population number is shown. Arrows in the respective color are indi-
cating the shift of the spectra with increasing numbers of discovery populations.

A very low target density, indicating that SNPs were mostly called due to being common

backbone SNPs, resulted in an allele frequency spectrum with the majority of alleles having a

MAF of around 0.5 (Figure 5). Increasing the target density for the equal spacing and thus

reducing the influence of the initial backbone of common SNPs shifted the peak of the allele

frequency spectrum left towards a higher proportion of alleles with small derived allele fre-

quencies. Using only the backbone SNPs common over all discovery populations and thus

calling SNPs mostly by the equal spacing procedure resulted, independently from the target

density, in a spectrum similar to the one obtained with a high target density with backbone

(Figure 5).

Figure 5: Derived allele frequency spectra for different target densities [SNPs/cM] after the equal spacing step. Blue spectra show the runs with the initial backbone included, red ones without backbone. Different target densities are indicated by a log scaled increase in color intensity. Only one representative run per target density is shown.

Overestimation of H_{exp}: Figure 6 shows the H_{exp} of different SNP sets by population. The H_{exp}

obtained from the filtered WGS SNPs were slightly higher than from the unfiltered WGS

SNPs. H_{exp} obtained from the ascertained SNP sets showed an even more pronounced over-

estimation together with an increase during the design steps. In general, the correlations

between the H_{exp} obtained in the different SNP sets were relatively high (≥ 0.94; Table S 2).

Especially the H_{exp} of the two WGS SNP sets showed a nearly perfect correlation of > 0.99,

which led to an almost constant OHE of -0.23 (Table 1) for the unfiltered WGS SNPs. As al-

ready recognizable from the H_{exp} themselves, the OHE was positive for all ascertained SNP

sets (0.66 – 1.29), which at the same time showed a slightly reduced correlation to the fil-

tered WGS SNP set (0.95 – 0.97). Comparable to the allele frequency spectra, the most pro-

nounced increase of the OHE was caused by the SNP discovery and followed by the equal

spacing step (OHE increased by 0.66), while the OHE from the original array SNPs (1.41; Fig-

ure 6; Table 1) lay in the range covered by the remodeling steps.

339
340   Figure 6: Expected Heterozygosity (H_{exp})by population and SNP set. Populations are ordered
341   by the H_{exp} of the unfiltered WGS SNP set. Only the reference sets and relevant steps of the
342   array design are shown. Discovery populations are shaded with a darker background.

Table 1: OHE of the SNP sets from the first run

Populations	Unfiltered WGS	Array SNPs	1 – SNP discovery	2 - cluster removal	3 – equal spacing	4 – validation	5 – down-sampling
All	$-0.23_{\pm 0.01}$	$0.84_{\pm 0.30}$	$0.66_{\pm 0.26}$	$0.66_{\pm 0.26}$	$1.09_{\pm 0.32}$	$1.29_{\pm 0.35}$	$1.27_{\pm 0.34}$
Discovery	$-0.23_{\pm 0.00}$	$1.08_{\pm 0.12}$	$0.91_{\pm 0.05}$	$0.90_{\pm 0.05}$	$1.15_{\pm 0.22}$	$1.26_{\pm 0.21}$	$1.28_{\pm 0.13}$
Validation	$-0.23_{\pm 0.00}$	$0.84_{\pm 0.11}$	$0.65_{\pm 0.07}$	$0.65_{\pm 0.07}$	$1.13_{\pm 0.11}$	$1.38_{\pm 0.13}$	$1.35_{\pm 0.14}$
Application	$-0.23_{\pm 0.00}$	$0.83_{\pm 0.11}$	$0.65_{\pm 0.07}$	$0.65_{\pm 0.07}$	$1.13_{\pm 0.10}$	$1.32_{\pm 0.12}$	$1.30_{\pm 0.13}$
Outgroup	-0.17	-0.88	-0.85	-0.86	-0.85	-0.85	-0.84

Mean OHE ± standard deviation
An OHE of zero means no bias and an OHE of 1 means doubling the H_{exp}

Averaging the OHE within the population groups revealed a 40 % higher OHE of the discovery populations compared to validation and application populations after the discovery step. The equal spacing step reduced this difference to an only 1 % larger OHE for discovery populations, while it came with a substantial increase of the variance of OHE, which was larger for the discovery populations than validation and application populations. The validation step then increased the OHE of the validation populations more than the OHE of discovery and application populations. This stronger OHE of discovery populations was also apparent within the array SNPs (28 % higher). In contrast to the other populations, the outgroup showed an underestimation of the H_{exp}, resulting in an OHE of < -0.84 for all ascertained SNP sets (Figure 6; Table 1).

A closer look on the contribution of the sub-steps during the equal spacing step revealed that 65 % of the SNPs which were preserved during equal spacing were variable in all of the three closely related broiler populations (BR1, BR2, NH; maximum pairwise Nei's distances (Nei 1972) of 0.07 in the unfiltered SNP set), while only 1.2 % of the SNPs were retained due to being variable in all of the five less closely related layer populations (WL, LE, WR, BL, RI; maximum pairwise Nei's distances of 0.15 in the unfiltered SNP set). The first population

362 used to fill in the gaps in the backbone (WL) contributed 17 % of the SNPs, while the other

363 populations contributed < 5 %.

365 Table 2: OHE of the SNP sets out of the 50 random population groupings

Populations	1 – SNP discovery	2 – cluster removal	3 – equal spacing	4 – validation	5 – down-sampling
Discovery	$0.76_{\pm0.004}{}^{a}$	$0.75_{\pm0.004}{}^{a}$	$1.13_{\pm0.006}{}^{a}$	$1.28_{\pm0.006}{}^{b}$	$1.33_{\pm0.007}{}^{a}$
Validation	$0.61_{\pm0.003}{}^{b}$	$0.60_{\pm0.003}{}^{b}$	$1.11_{\pm0.004}{}^{b}$	$1.29_{\pm0.004}{}^{b}$	$1.29_{\pm0.005}{}^{b}$
Application	$0.61_{\pm0.003}{}^{b}$	$0.61_{\pm0.003}{}^{b}$	$1.12_{\pm0.004}{}^{ab}$	$1.35_{\pm0.004}{}^{a}$	$1.34_{\pm0.004}{}^{a}$
Outgroup	$-0.85_{\pm0.008}{}^{c}$	$-0.86_{\pm0.008}{}^{c}$	$-0.85_{\pm0.015}{}^{c}$	$-0.84_{\pm0.017}{}^{c}$	$-0.84_{\pm0.019}{}^{c}$

LSMEANS for OHE ± standard error
An OHE of zero means no bias and an OHE of 1 means doubling the H_{exp}
Different lowercase letters within columns indicate significant differences to the 5 % level

These findings were supported by the 50 random groupings (Figure 7). The LSMEANS (Ta-

ble 2) of the population groups revealed 24 % larger OHE for discovery populations than for

validation and application populations after discovery and cluster removal step, which was

decreased to a numerically insignificant difference after the equal spacing step. Interestingly,

and in contrast to the findings from the first remodeling, SNP validation led to a significantly

higher OHE (5 % larger) for application populations than discovery and validation popula-

tions.

Figure 7: OHE for discovery validation and application populations after the five steps of ar-
ray design. Discovery populations are chosen to represent populations which are compara-
ble to the original array (blue) or 50 times randomly sampled (grey.

Influence of number of discovery populations and target density on H_exp: Figure 8 A shows

that increasing the number of discovery populations reduces the mean OHE of discovery

populations after SNP discovery while not affecting the OHE of validation and application

populations. After equal spacing (Figure 8 B), no dependency of OHE on the number of dis-

covery populations remained. As, due to the limited number of populations in the complete

set, the number of validation populations decreased with more populations in the discovery

set, the individual validation populations gained more impact on the ascertainment. This led

to a higher OHE of validation populations with a high number of discovery populations (Fig-

ure S 6 D), comparable to the higher OHE of discovery populations for a small number of

discovery populations.

Figure 8: Relation of the OHE as a function of the number of discovery populations.
A - discovery, **B**– equal spacing. The boxplots are only shown for a subset of the number of
discovery populations, while the smoothing lines, which show the trend, are calculated from
all observations. Plots for all five steps can be found in Figure S 6.

In the equal spacing step, using only backbone SNPs resulted in a higher OHE for discov-

ery than for non- discovery populations. Increasing the target density and thus increasing the

proportion of SNPs due to the equal spacing part of the algorithm reduced the difference in

OHE between the population groups (Figure 9 A). If the SNPs from the initial backbone were

not used, no difference of OHE between discovery and non- discovery populations was pre-

sent, regardless of the target density (Figure 9 B).

Figure 9: OHE after equal spacing (step 3) by target density in SNPs/cM and population
group. The smoothing lines show the trend and the dashed lines the target density of 667
SNPs/cM, used for the remodeling according to Kranis *et al.* (2013). The algorithm was run
including the initial backbone SNPs (**A**) or not including them (**B**). Gallus varius is not includ-
ed, as it is constantly underestimated.

DISCUSSION

In this study we used a uniquely diverse collection of sequenced wild, commercial and non-commercial chicken populations, mainly based on samples of the Synbreed Chicken Diversity Panel (Malomane *et al.* 2019). Parts of our set were also involved in the development process of the Axiom™ Genome-Wide 580 k Chicken Array (Kranis *et al.* 2013). This offered an excellent possibility for assessing the impact of ascertainment bias on real data in a complex scenario.

Potential impacts of the SNP calling pipeline: As the state of the art pipeline of GATK relies on a supervised machine learning approach for filtering the SNP calls, which needs a highly reliable set of known SNPs, we started with examining potential impacts of the filtering procedure on ascertainment bias. The number of rare variants was slightly reduced by the filtering procedure and thus increased estimates of H_{exp} were obtained in the filtered WGS set. As rare variants have a higher risk to be discarded as sequencing errors (Heslot *et al.* 2013), this reduction is expected when applying quality filters. However, a clear assessment of correctly and falsely filtered variants is not possible here and one has to balance this tradeoff based on the study purpose.

Another source of ascertainment bias could be the use of array SNPs as training set for GATK RecalibrateVariants, which potentially leads to discarding rare variants more likely if they are not present in the discovery populations of the used array. As the correlation between the H_{exp} of the unfiltered and filtered WGS SNPs was nearly one, this source seems to be negligible and the use of array variants as a highly reliable training set seems to be unproblematic.

Additionally, due to computational limitations, we had to assume a ploidy of two for

pools during the SNP calling process, which resulted in a minimal reduction of rare alleles.

However, this effect was shown to have a very minor impact on the findings of this study

(supplementary file 2).

General impact over all groups: The general reduction of rare alleles in array data compared

to WGS data and the resulting overestimation of H_{exp} supports findings of previous studies

(Nielsen 2004; Clark *et al.* 2005; Albrechtsen *et al.* 2010; Malomane *et al.* 2018). This reduc-

tion of rare alleles was mainly seen at steps where selection was explicitly biased towards

high MAF alleles (MAF filter for quality control in discovery step and use of common alleles

for the backbone in the equal spacing step) and/or was applied to a small number of popula-

tions (small discovery set vs. small validation set). Thereby, the strongest shifts of the allele

frequency spectra and increases of H_{exp} are observed after SNP discovery and equal spacing.

Both, cluster removal and second downsampling had almost no effect on allele frequency

spectra and H_{exp}, while the validation step slightly decreased the share of rare SNPs.

The discovery step had the strongest impact on discovery populations, when a small set

of discovery populations was used (Figure 8 A). Similarly, the influence of the validation step

on validation populations was strongest in case of a small number of validation populations

(Figure S 6D). A balancing of these two groups of samples is therefore necessary, if the num-

ber of available DNA samples for array development is limited. We rather suggest the use of

an extended discovery set for validation as best practice instead of using completely differ-

ent populations.

If the equal spacing step contains a preselection of SNPs based on their variability within

population groups, the bias is stronger towards high MAF SNPs and thus yields a higher OHE.

This effect was reduced by increasing the target density and thus selecting relatively more

SNPs due to the equal spacing instead of common occurrence.

Differences between groups: If allele frequency spectra are changed in the same way for all

populations and are therefore biasing heterozygosity estimates to the same amount, find-

ings for between population comparisons will be little affected. Ascertainment bias then is

only of importance if one compares populations based on different arrays, and corrections

of the allele frequency spectrum as reviewed by Nielsen (2004) should be possible. As corre-

lations between H_{exp} of ascertained SNP sets and unfiltered/ filtered WGS SNP sets were

consistently high (> 0.94), arrays designed in the way as performed in this paper should

mostly be suitable for robust and cost efficient analyses. Unbiasedness of estimates could be

increased even more by considering filter strategies according to Malomane *et al.* (2018).

However, we could show that a structural bias due to the second source of ascertainment

bias, the across population use of arrays, exists. This structural bias was already shown to

have severe impact on findings from SNP arrays. For example Bradbury *et al.* (2011) found a

demographic decline up to an approximately 30 % lower H_{exp} for Atlantic cod based on the

distance to the sampling location of the discovery panel and McTavish and Hillis (2015)

showed strong deviations between simulated and observed polymorphisms for different

combinations of migration- and ascertainment scenarios on simulated cattle populations. In

our study, populations which are closely related to the discovery populations of the original

array on average showed a 28 % higher OHE than validation and application populations for

the original array, which supports the mentioned findings from literature.

This structural bias was mainly introduced by the initial discovery step. It was also ob-

served for the random population groupings, but to a slightly smaller extent. The difference

in overestimation decreased with an increase in the number of discovery populations (Figure

8) and was smallest if the discovery populations showed minimum distance to the applica-

tion and validation populations (results not shown). Comparable observations were already

made by Frascaroli *et al.* (2013) who found very small ascertainment bias for European elite

maize lines when using a SNP panel discovered in a very diverse set of maize and inbred

lines, but strong ascertainment bias when using SNP discovered in American elite lines.

Therefore, we suggest to ideally choose an array where the discovery panel does span the

scope of populations it will be applied to, and by this covers the existing variation in a most

representative way, or to design such an array for oneself if it does not exist.

The equal spacing step removed the structural bias in most of our remodeling scenarios,

but obviously not in case of the original array. In the remodeling, we saw this structural bias

only with a low target density and thus calling SNPs in the equal spacing step mainly due to

being common over many populations (Figure 9 A). However, the equal spacing step also

increased the variance of OHE in the discovery panel, meaning that the OHE was increased

more for some of the discovery populations than for others, causing more uncertainty for

resulting effects. This effect is driven by the unequal contribution of populations during the

equal spacing step. It increases the bias for some of the discovery populations, while it de-

creases it for others, and hence it does not remove the structural bias. This means that the

knowledge of which discovery populations were used alone is not sufficient to draw conclu-

sions regarding a possible ascertainment bias, since their relative contribution varies through

the described pipeline.

Outgroup: *Gallus varius* as an outgroup showed a different behavior than all other popula-

tions. It already exhibited the lowest H_{exp} in the unfiltered WGS SNP set, which was most

likely driven by the small number of only two samples in the pool, and showed less upward

bias of H_{exp} in the filtered WGS SNP set than all other populations. The *Gallus varius* se-

quence reads on average showed weak Phred-scaled mapping quality scores of 19 (1.3 %

probability of misalignment), while the mean quality scores of the other populations ranged

from 25 (0.3 %) to 28 (0.1 %) – meaning that variation, only present in *Gallus varius*, will be

more likely missed due to misplacement of the reads or discarded as possible sequencing

errors. Additionally, every ascertained SNP set showed an OHE for *Gallus varius* of < -0.84, as

variation being present only in *Gallus varius* was not found in *Gallus gallus* discovery panels

and contrary variants from *Gallus gallus* were not variable in *Gallus varius* (Figure 3). This

demonstrates that arrays should not be used if different species (even closely related ones)

are included in the research project. Even sequence based estimates can be slightly biased, if

the reference genome does not fit properly.

Further recommendations for future studies: We showed that existing arrays come with a

large, and due to a diverse set of promoting factors barely predictable, potential for ascer-

tainment bias. Strongly decreasing costs for WGS and increasing availability of powerful

computing resources therefore promote an intensified use of WGS for population genetic

analyses, especially when diverse populations are included in the studies. However, costs

and computational effort will still be substantial for large scale projects. Possible cost effec-

tive alternatives could be reduced library sequencing approaches like Genotyping-by-

Sequencing (Elshire *et al.* 2011; Heslot *et al.* 2013), even though such methods introduce

other problems related to the use of restriction enzymes which are reviewed in Andrews *et*

al. (2016).

For the purpose of monitoring genetic diversity in a large set of small non-commercial

populations, the development of a specialized new array for high throughput cost effective

genotyping could be still a good option. For the design of such an array, it is crucial to extend

the discovery set in a way which represents the total variability over all populations as bal-

anced as possible. The use of publicly available sequences can be helpful to reach this goal.

The ascertainment of the SNPs should then be done preferably over all available populations

instead of biasing the process towards common alleles by performing within population as-

certainment.

However, when considering the findings of this study, choosing an array for which the

SNP ascertainment process is well documented and a discovery panel that matches the pop-

ulations to be studied, should give sufficient results when robust estimators are used.

ABBREVIATIONS

535   H_{exp} – expected heterozygosity

536   MAF – minor allele frequency

537   OHE – overestimation of expected heterozygosity

538   SNP – single nucleotide polymorphism

539   WGS – whole genome re-sequencing

DECLARATIONS

541 Ethics approval and consent to participate

Sampling was approved by the animal welfare committees of the involved institutions

and was done in strict accordance to the German Animal Welfare regulations and with writ-

ten consent of the animal owners. In case of the samples which were taken during the SYN-

BREED project, according to a change in German regulations, notice was additionally given to

the Lower Saxony State Office for Consumer Protection and Food Safety (33.9-42502-05-

10A064).

548 Funding

549   The sampling and sequencing of the chicken populations was funded by the EC project

550 AVIANDIV (EC Contract No. BIO4-CT98_0342; 1998 – 2000; www.aviandiv.fli.de) and the

551 German Federal Ministry of Education and Research (BMBF) via the SYNBREED project (FKZ

552 0315528E; 2009 – 2014; www.synbreed.tum.de).

Authors' contributions

JG designed the study, analyzed and interpreted the data, wrote the final draft and pre-

pared the manuscript. HS supervised the study, substantially contributed to design and con-

ception of the study as well as revision and editing of the manuscript. CR substantially con-

tributed to design and conception of the study, participated in interpreting the data and was

involved in revision and editing of the manuscript. SW contributed to design and conception

of the study, participated in the provision of the data as well as in revision and editing of the

manuscript. AW participated in provision and preparation of the data. TP wrote a part of the

equal spacing algorithm and participated in revision and editing of the manuscript. All au-

thors have read and approved the final manuscript.

Acknowledgments

564   We are grateful to all the participating breeders for their assistance in sampling.

565   **REFERENCES**

Albrechtsen, A., F. C. Nielsen, and R. Nielsen, 2010 Ascertainment biases in SNP chips affect
measures of population divergence. Mol Biol Evol 27 (11): 2534–2547.
Andrews, K. R., J. M. Good, M. R. Miller, G. Luikart, and P. A. Hohenlohe, 2016 Harnessing
the power of RADseq for ecological and evolutionary genomics. Nat Rev Genet 17 (2): 81.
Bates, D., M. Mächler, B. Bolker, and S. Walker, 2015 Fitting Linear Mixed-Effects Models
Using   lme4.   Journal   of   Statistical   Software   67   (1):   1–48.
https://doi.org/10.18637/jss.v067.i01
Boichard, D. A., H. Chung, R. Dassonneville, X. David, A. Eggen *et al.*, 2012 Design of a bo-
vine low-density SNP array optimized for imputation. PLoS One 7 (3): e34130.
Bradbury, I. R., S. Hubert, B. Higgins, S. Bowman, I. G. Paterson *et al.*, 2011 Evaluating SNP
ascertainment bias and its impact on population assignment in Atlantic cod, Gadus
morhua. Mol Ecol Res 11 (s1): 218–25.
Broad Institute, 2015 Picard Tools 2.0.1. https://broadinstitute.github.io/picard/.
Broad   Institute,   2018 GATK   User   Guide.   Accessed   March   20,   2018.
https://software.broadinstitute.org/gatk/documentation/.

Clark, A. G., M. J. Hubisz, C. D. Bustamante, S. H. Williamson, and R. Nielsen, 2005 Ascertainment bias in studies of human genome-wide polymorphism. Genome Res 15 (11): 1496–1502.

DePristo, M. A., E. Banks, R. Poplin, K. V. Garimella, J. R. Maguire *et al.*, 2011 A framework for variation discovery and genotyping using next-generation DNA sequencing data. Nat Genet 43: 491. https://doi.org/10.1038/ng.806

Eller, E., 2001 Effects of Ascertainment Bias on Recovering Human Demographic History. Human Biology 73 (3): 411–27. https://doi.org/10.1353/hub.2001.0034

Elshire, R. J., J. C. Glaubitz, Q. Sun, J. A. Poland, K. Kawamoto *et al.*, 2011 A robust, simple genotyping-by-sequencing (GBS) approach for high diversity species. PLoS One 6 (5): e19379.

ENSEMBL, 2016 Chicken Germline SNP and INDELS. Accessed January 06, 2017. http://e87.ensembl.org/Gallus_gallus.

Fan, B., Z.-Q. Du, D. M. Gorbach, and M. F. Rothschild, 2010 Development and application of high-density SNP arrays in genomic studies of domestic animals. Asian-Australas J Anim Sci 23 (7): 833–47.

Frascaroli, E., T. A. Schrag, and A. E. Melchinger, 2013 Genetic diversity analysis of elite European maize (Zea mays L.) inbred lines using AFLP, SSR, and SNP markers reveals ascertainment bias for a subset of SNPs. Theoretical and Applied Genetics 126 (1): 133–41. https://doi.org/10.1007/s00122-012-1968-6

Gautier, M., D. Laloe, and K. Moazami-Goudarzi, 2010 Insights into the genetic history of French cattle from dense SNP data on 47 worldwide breeds. PLoS One 5 (9). https://doi.org/10.1371/journal.pone.0013038

Gibbs, R. A., J. F. Taylor, C. P. van Tassell, W. Barendse, K. A. Eversole *et al.*, 2009 Genome-wide survey of SNP variation uncovers the genetic structure of cattle breeds. Science 324 (5926): 528–32. https://doi.org/10.1126/science.1167936

Groenen, M. A. M., H.-J. Megens, Y. Zare, W. C. Warren, L. W. Hillier *et al.*, 2011 The development and characterization of a 60K SNP chip for chicken. BMC Genomics 12 (1): 274.

Guillot, G., and M. Foll, 2009 Correcting for ascertainment bias in the inference of population structure. Bioinformatics 25 (4): 552–54. https://doi.org/10.1093/bioinformatics/btn665

Heslot, N., J. Rutkoski, J. A. Poland, J.-L. Jannink, and M. E. Sorrells, 2013 Impact of marker ascertainment bias on genomic selection accuracy and estimates of genetic diversity. PLoS One 8 (9): e74612.

Kijas, J. W., D. Townley, B. P. Dalrymple, M. P. Heaton, J. F. Maddox *et al.*, 2009 A genome wide survey of SNP variation reveals the genetic structure of sheep breeds. PLoS One 4 (3): e4668.

Kimura, M., 1991 The neutral theory of molecular evolution: A review of recent evidence. Idengaku zasshi 66 (4): 367–86.

Kranis, A., A. A. Gheyas, C. Boschiero, F. Turner, Le Yu *et al.*, 2013 Development of a high density 600K SNP genotyping array for chicken. BMC Genomics 14 (1): 59. https://doi.org/10.1186/1471-2164-14-59

Lachance, J., and S. A. Tishkoff, 2013 SNP ascertainment bias in population genetic analyses: why it is important, and how to correct it. Bioessays 35 (9): 780–86.

Laurie, C. C., D. A. Nickerson, A. D. Anderson, B. S. Weir, R. J. Livingston *et al.*, 2007 Linkage
Disequilibrium in Wild Mice. PLoS Genet 3 (8): e144.
https://doi.org/10.1371/journal.pgen.0030144

Lenth, R., 2019 *emmeans: Estimated Marginal Means, aka Least-Squares Means*.

Li, H., 2013 Aligning sequence reads, clone sequences and assembly contigs with BWA-
MEM. Accessed March 21, 2019. https://arxiv.org/pdf/1303.3997.

Malomane, D. K., C. Reimer, S. Weigend, A. Weigend, A. R. Sharifi *et al.*, 2018 Efficiency of
different strategies to mitigate ascertainment bias when using SNP panels in diversity
studies. BMC Genomics 19 (1): 22. https://doi.org/10.1186/s12864-017-4416-9

Malomane, D. K., H. Simianer, A. Weigend, C. Reimer, A. O. Schmitt *et al.*, 2019 The SYN-
BREED chicken diversity panel: A global resource to assess chicken diversity at high ge-
nomic resolution. BMC genomics 20 (1): 345. https://doi.org/10.1186/s12864-019-5727-9

Matukumalli, L. K., C. T. Lawley, R. D. Schnabel, J. F. Taylor, M. F. Allan *et al.*,
2009 Development and characterization of a high density SNP genotyping assay for cat-
tle. PLoS One 4 (4): e5350.

Mayer, M., S. Unterseer, E. Bauer, N. de Leon, B. Ordas *et al.*, 2017 Is there an optimum
level of diversity in utilization of genetic resources? Theor Appl Genet 130 (11): 2283–95.
https://doi.org/10.1007/s00122-017-2959-4

McKenna, A., M. Hanna, E. Banks, A. Sivachenko, K. Cibulskis *et al.*, 2010 The Genome Anal-
ysis Toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data.
Genome Res 20 (9): 1297–1303. https://doi.org/10.1101/gr.107524.110

McLaren, W., L. Gil, S. E. Hunt, H. S. Riat, G. R. S. Ritchie *et al.*, 2016 The Ensembl Variant
Effect Predictor. Genome Biol 17 (1): 122. https://doi.org/10.1186/s13059-016-0974-4

McTavish, E. J., J. E. Decker, R. D. Schnabel, J. F. Taylor, and D. M. Hillis, 2013 New World
cattle show ancestry from multiple independent domestication events. Proc Natl Acad Sci
110 (15): E1398-E1406.

McTavish, E. J., and D. M. Hillis, 2015 How do SNP ascertainment schemes and population
demographics affect inferences about population history? BMC Genomics 16 (1): 1.

Muir, W. M., G. K.-S. Wong, Y. Zhang, J. Wang, M. A. M. Groenen *et al.*, 2008 Genome-wide
assessment of worldwide chicken SNP genetic diversity indicates significant absence of ra-
re alleles in commercial breeds. Proc Natl Acad Sci 105 (45): 17312–17.
https://doi.org/10.1073/pnas.0806569105

Nei, M., 1972 Genetic Distance between Populations. The American Naturalist 106 (949):
283–92.

Nielsen, R., 2004 Population genetic analysis of ascertained SNP data. Hum Genomics 1 (3):
1.

Nielsen, R., M. J. Hubisz, and A. G. Clark, 2004 Reconstituting the frequency spectrum of
ascertained single-nucleotide polymorphism data. Genetics 168 (4): 2373–82.

Nielsen, R., and J. Signorovitch, 2003 Correcting for ascertainment biases when analyzing
SNP data: applications to the estimation of linkage disequilibrium. Theor Popul Biol 63 (3):
245–55.

Novembre, J., T. Johnson, K. Bryc, Z. Kutalik, A. R. Boyko *et al.*, 2008 Genes mirror geogra-
phy within Europe. Nature 456: 98. https://doi.org/10.1038/nature07331

Paradis, E., J. Claude, and K. Strimmer, 2004 APE: Analyses of phylogenetics and evolution
in R language. Bioinformatics 20: 289–90.

Patterson, N., P. Moorjani, Y. Luo, S. Mallick, N. Rohland *et al.*, 2012 Ancient admixture in
human history. Genetics 192 (3): 1065–93. https://doi.org/10.1534/genetics.112.145037

Perkel, J., 2008 SNP genotyping: Six technologies that keyed a revolution. Nature Methods
5: 447. https://doi.org/10.1038/nmeth0508-447

Platt, A., M. Horton, Y. S. Huang, Y. Li, A. E. Anastasio *et al.*, 2010 The Scale of Population
Structure in Arabidopsis thaliana. PLoS Genet 6 (2): e1000843.
https://doi.org/10.1371/journal.pgen.1000843

Qanbari, S., E. C.G. Pimentel, J. Tetens, G. Thaller, P. Lichtner *et al.*, 2010 A genome-wide
scan for signatures of recent selection in Holstein cattle. Animal genetics 41 (4): 377–89.

Qanbari, S., C.-J. Rubin, K. Maqbool, S. Weigend, A. Weigend *et al.*, 2019 Massive whole
genome sequencing reveals strong purifying selection in commercial chicken: - accepted
but not yet published. PLoS Genet.

Qanbari, S., M. Seidel, T.-M. Strom, K. F. X. Mayer, R. Preisinger *et al.*, 2015 Parallel Selec-
tion Revealed by Population Sequencing in Chicken. Genome Biol Evol 7 (12): 3299–3306.
https://doi.org/10.1093/gbe/evv222

R Core Team, 2017 *R: A Language and Environment for Statistical Computing.* Vienna, Aus-
tria.

Ramos, A. M., R. P. M. A. Crooijmans, N. A. Affara, A. J. Amaral, A. L. Archibald *et al.*,
2009 Design of a high density SNP genotyping assay in the pig using SNPs identified and
characterized by next generation sequencing technology. PLoS One 4 (8): e6524.

Rocha, D., C. Billerey, F. Samson, D. Boichard, and M. Boussaha, 2014 Identification of the
putative ancestral allele of bovine single-nucleotide polymorphisms. J Anim Breed Genet
131 (6): 483–86. https://doi.org/10.1111/jbg.12095

Sandenbergh, L., S. W.P. Cloete, R. Roodt-Wilding, M. A. Snyman, and A. E. Bester-van der
Merwe, 2016 Evaluation of the OvineSNP50 chip for use in four South African sheep
breeds. S Afr J Anim Sci 46 (1): 89–93.

Singh, N., P. K. Jayaswal, K. Panda, P. Mandal, V. Kumar *et al.*, 2015 Single-copy gene based
50 K SNP chip for genetic studies and molecular breeding in rice. Sci Rep 5: 11600.

Tosser-Klopp, G., P. Bardou, O. Bouchez, C. Cabau, R. P. M. A. Crooijmans *et al.*, 2014 Design
and characterization of a 52K SNP chip for goats. PLoS One 9 (1): e86227.

Travis, A. J., G. J. Norton, S. Datta, R. Sarma, T. Dasgupta *et al.*, 2015 Assessing the genetic
diversity of rice originating from Bangladesh, Assam and West Bengal. Rice 8 (1): 35.
https://doi.org/10.1186/s12284-015-0068-z

UCSC, 2016 Reference Genome Gallus gallus 5.0. Accessed October 25, 2016.
http://hgdownload.soe.ucsc.edu/goldenPath/galGal5/bigZips/galGal5.fa.gz.

Unterseer, S., E. Bauer, G. Haberer, M. Seidel, C. Knaak *et al.*, 2014 A powerful tool for ge-
nome analysis in maize: development and evaluation of the high density 600 k SNP geno-
typing array. BMC Genomics 15 (1): 823.

van der Auwera, G. A., M. O. Carneiro, C. Hartl, R. Poplin, G. Del Angel *et al.*, 2013 From
FastQ data to high confidence variant calls: The Genome Analysis Toolkit best practices
pipeline. Current protocols in bioinformatics 43 (1): 11.10.1-11.10.33.
https://doi.org/10.1002/0471250953.bi1110s43

Warren, W. C., L. W. Hillier, C. Tomlinson, P. Minx, M. Kremitzki *et al.*, 2017 A New Chicken
Genome Assembly Provides Insight into Avian Genome Structure. G3 7 (1): 109–17.
https://doi.org/10.1534/g3.116.035923

715
716

AUTHORS SUMMARY

Compared to whole genome sequencing data, SNP derived by genotyping arrays are bi-

ased towards common variants. This effect is widely known as SNP ascertainment bias. Es-

pecially the different strength of the bias for different populations, depending on whether

they were involved in the development process or not, can strongly affect population genet-

ic estimators. In this work Geibel et al. investigate the introduction of the bias in its magni-

tude during array development by redesigning a commercial chicken genotyping array under

usage of a diverse set of sequenced chicken populations and give recommendations for de-

veloping bias minimized arrays.

SUPPLEMENTARY FILE 2

Correct ploidy calling vs. ploidy two calling: Due to the large diversity of the chicken popula-

tions and a series of bugs in GATK (up to version 4.2), it was only possible to call some inter-

vals, spanning 28.2 % of the autosomal region, with the correct ploidy. Compared to the call-

ing with the correct ploidy, assuming ploidy two for the pools came along with a slight re-

duction of SNPs with very rare alternative allele frequencies of ≤ 0.02 in the pools (Figure S 1

C). As those SNPs had to be invariable in the individually sequenced populations, a reduction

of invariable alleles is present in the individually sequenced populations (Figure S 1 B). When

comparing expected heterozygosities (H_{exp}) from the two callsets, the ploidy two calling

comes along with a slight increase of H_{exp}, which is minimally larger for individually se-

quenced populations (Figure S 1 A), as the lost SNPs are invariable rather than rare in those

populations. However, as the correlation between the two callsets is > 0.99 and the mean

overestimation of H_{exp} (OHE) for the pools is 0.02 and 0.05 for the individual sequenced

populations, the effect is negligible compared to the effects of ascertainment bias.

Figure S 1: Pooled vs. ploidy two calling. Expected Heterozygosity (H_{exp}) from the calling with
assuming the correct ploidy vs. assuming ploidy two for all samples (A) and alternative allele

frequency distributions of called alleles for individually sequenced (B) and pooled sequenced

(C) populations.

Alternative allele frequency spectrum vs. derived allele frequency spectrum: As the chicken

reference genome is based on an inbred Red Jungle Fowl (*Gallus gallus gallus*; Warren *et al.*

2017), many publications handle the reference allele as ancestral allele (e.g. Qanbari *et al.*

2015). However, this assumption does not account for mutation, selection and inbreeding

happening in the Red Jungle Fowl after separation from the domesticated strains as well as

variation already present before separation. Therefore, ancestral alleles were defined by an

approach comparable to Rocha *et al.* (2014) using allele frequency information from the

three wild populations *Gallus gallus gallus* (GG), *Gallus gallus spadiceus* (GS) and *Gallus vari-*

us (GV). It was assumed that the *Gallus gallus* and *Galllus varius* species emerged from a

common ancestor and *Gallus gallus* later split up into *Gallus gallus gallus* and *Gallus gallus*

spadiceus subspecies. Additionally, assuming neutral molecular evolution (Kimura 1991), the

ancestral allele was most likely the major allele within those three populations, when

weighting the allele frequency of *Gallus varius* twice. This procedure assigned the ancestral

status to the reference allele for 86 % of the SNPs and to the alternative allele for 14 % of

the SNPs. SNPs where the ancestral state was assigned to the alternative allele were approx-

imately equal distributed over the range of allele frequencies (Figure S 2 A). Only alleles be-

ing close to fixation for the alternative allele were overrepresented, possibly being artefacts

of selection on de novo mutations in ancestors of the reference individual or sequencing

errors in the reference genome. Therefore, a comparison between alternative (Figure S 2 B)

and derived (Figure S 2 C) allele frequency spectra in the complete callset does only show

the reduction of a tiny peak being present at the end of the alternative allele frequency

spectrum.

774 Figure S 2: Alternative allele frequency spectrum for SNPs where the reference allele is not
775 the ancestral allele (A) and alternative (B) vs. derived (C) allele frequency spectrum of all
776 SNPs.

778 Table S 1: Number of SNPs from the remodeling processes

	1 – SNP discovery	2 - cluster removal	3 – equal spacing	4 – validation	5 – downsampling
Original Array (Kranis *et al.* 2013)	23 998	10 029	1 829	1 187	581
First Run	10 834	8 788	2 331	1 910	600
Repeated Remodeling	11 603 ± 226	9 328 ± 157	2 032 ± 147	1 698 ± 147	662 ± 33

Number of SNPs in 1,000
In case of repeated remodeling mean of the 50 runs ± standard deviation

780 Table S 2: Pearson correlations between the H_{exp} of the different SNP sets

	Filtered WGS	Array SNPs	1 – SNP discovery	2 - cluster removal	3 – equal spacing	4 – validation	5 – downsampling
Unfiltered WGS	1.00	0.94	0.95	0.95	0.96	0.96	0.96
Filtered WGS		0.95	0.96	0.95	0.97	0.96	0.96
Array SNPs			1.00	1.00	0.98	0.97	0.98
1 – SNP discovery				1.00	0.98	0.97	0.97
2 - cluster					0.98	0.96	0.97

removal		
3 – equal spacing	1.00	0.99
4 – vali-dation		1.00

784   Figure S 3: SNPs retained by number of populations in the discovery set.

788   Figure S 4: Number of SNPs retained after the equal spacing step by target density.

791   Figure S 5: Relative amount of SNPs shared by a specific number of runs from 50 random
792   population assignments.

Figure S 6: Relation of the OHE as a function of the number of discovery populations. **A** - discovery, **B** – cluster removal, **C** – equal spacing, **D** – validation, **E** – downsampling. The Boxplots are only shown for a subset of the number of discovery populations, while the smoothing lines, which show the trend, are calculated from all observations.