How Array Design affects SNP Ascertainment Bias

Johannes Geibel[1,2*], Christian Reimer[1,2], Steffen Weigend[2,3], Annett Weigend[3], Torsten

Pook[1,2], Henner Simianer[1,2]

[1] Department of Animal Sciences, Animal Breeding and Genetics Group, University of
Goettingen, Göttingen, Germany

[2] Center for Integrated Breeding Research, University of Goettingen, Göttingen, Germany

[3] Institute of Farm Animal Genetics, Friedrich-Loeffler-Institut, Neustadt-Mariensee, Germa-
ny

* Corresponding author

E-Mail: johannes.geibel@uni-goettingen.de (JG)

Abstract

Single nucleotide polymorphisms (SNPs), genotyped with SNP arrays, have become a widely used marker type in population genetic analyses over the last 10 years. However, compared to whole genome re-sequencing data, arrays are known to lack a substantial proportion of globally rare variants and tend to be biased towards variants present in populations involved in the development process of the respective array. This affects population genetic estimators and is known as SNP ascertainment bias. We investigated factors contributing to ascertainment bias in array development by redesigning the Axiom™ Genome-Wide Chicken Array *in silico* and evaluating changes in allele frequency spectra and heterozygosity estimates in a stepwise manner. A sequential reduction of rare alleles during the development process was shown with main influencing factors being the identification of SNPs in a limited set of populations and a within-population selection of common SNPs when aiming for equidistant spacing. These effects were shown to be less severe with a larger discovery panel. Additionally, a generally massive overestimation of expected heterozygosity for the ascertained SNP sets was shown. This overestimation was 24 % higher for populations involved in the discovery process than not involved populations in case of the original array. The same was observed after the SNP discovery step in the redesign. However, an unequal contribution of populations during the SNP selection can mask this effect but also adds uncertainty. Finally, we make suggestions for the design of specialized arrays for large scale projects where whole genome re-sequencing techniques are still too expensive.

Introduction

Starting in the first decade of this century, the possibility of cost effectively genotyping high numbers of Single Nucleotide Polymorphisms (SNP) for many individuals in parallel via SNP arrays led to an increased usage of them for population genetic analyses in humans [1,2], model species [3,4], plants [5,6] and livestock [7–13].

Various SNP arrays exist for humans [14], plants [15,16] and all major livestock species [17–23]. SNP numbers within these arrays range from 10 k SNPs [20] over a range from 40 to 60 k [16,17,19,22] up to 600 k [15,21]. The design process of every array has an initial step of SNP discovery in common, where SNPs are identified from existing databases and/or from a small set of sequenced individuals. SNPs are then selected based on different quality criteria like minor allele frequency (MAF) thresholds and platform specific design scores [24]. Additional criteria like equidistant spacing over the genome [21], overrepresentation of some areas [20] or increased overrepresentation of high MAF SNPs [17] are applied dependent on the design intentions. In the end, draft arrays are validated either on the set of populations used for the SNP discovery itself [18] and/or on a broad set of individuals from different populations [21,24].

In contrast to whole genome re-sequencing (WGS) data, SNP arrays often show a clear underrepresentation of SNP with extreme allele frequencies [25]. As population genetic statistics are mostly based on estimates of allele frequencies, this context leads to biased population genetic estimators [25,26] and is known as SNP ascertainment bias.

The absence of rare alleles is mainly driven by two factors in the array design process where SNPs are selected (ascertained) based on different requirements and decisions [27].

The first factor is a relatively small panel of individuals being used for discovery of SNPs,

leading to a large proportion of globally rare variants not being selected, since they appear

monomorphic in the discovery panel [26,28]. The second factor is the across population use

of arrays. Arrays are developed based on the variation within the discovery panel, thus miss-

ing variation present in distantly related individuals or populations [25,27]. This second

source of bias was shown to be of relatively high importance for livestock studies, where

arrays are usually developed for large commercial breeds and later used to genotype diverse

sets of local breeds all over the world [29,30].

Besides different strategies to minimize the impact of ascertainment bias [30,31], there

are some attempts to correct the allele frequency spectrum via Bayesian methods

[25,28,32]. However, those corrections highly rely on detailed statistical assumptions of the

ascertainment process [33,34] and are currently only tested for corrections of the first

source of ascertainment bias, the small discovery panel [25,28,32]. In practice, detailed in-

formation on the design process is limited [34] and the complexity of the processes makes

statistical models for the corrections inaccurate.

Agricultural species such as chickens often show a complex domestication history, and

therefore allow for few prior assumptions on ascertainment bias. Domestic chickens are as-

sumed to originate from red jungle fowl (*Gallus gallus*) ancestors in Southeast Asia [35,36],

represented by the five subspecies *G. g. gallus*, *G. g. spadiceus*, *G. g. murghi*, *G. g. bankiva*

and *G. g. jabouillei* [37]. Additionally, some hybridization events with other *Gallus* species

(e.g. grey jungle fowl; *Gallus sonneratii*) have been suggested [36,38]. The diversity of to-

day's local breeds of chickens in Europe originates from chickens that reached the continent

about 3000 years ago via a northern and a southern route, followed by selection and cross-

ing with Asian chicken breeds introduced in the 19th century [37]. While commercial white

layers were derived solely by intensive directional selection of a single breed, the White Leg-

horn, commercial brown layers are derived from a broader genetic basis (e.g. Rhode Island

Red, New Hampshire, Barred Plymouth Rock). Commercial broilers are derived by cross-

breeding of paternal lines (e.g. White Cornish) with maternal lines which descend from a

comparable basis as brown layers (e.g. White Plymouth Rock) [39]. For more detailed infor-

mation on chicken ancestry we refer to Qanbari *et al.* [36] and for a comprehensive overview

on diversity and population structure of domesticated chickens to Malomane *et al.* [13].

Given the complexity of modern array design processes and the chicken population struc-

ture, this study aims at highlighting the mechanisms which promote the bias by illustrating

the effects of the different steps of the array design process on ascertainment bias, using

real data in a typical setting from livestock sciences. For this purpose, the design process of

the Axiom™ Genome-Wide Chicken Array [21] was simulated in a set of diverse chicken WGS

data. Allele frequency spectra as well as expected heterozygosity (H_{exp}) were compared to

the WGS data and the SNPs of the Axiom™ Genome-Wide Chicken Array. Finally, some rec-

ommendations are made to design an array for monitoring genetic diversity.

Material and methods

Populations and sequencing

The analysis is based on WGS data of a diverse set of 46 commercial, non-commercial and

wild chicken populations, sampled within the framework of the projects AVIANDIV

(www.aviandiv.fli.de) and SYNBREED (www.synbreed.tum.de). Commercial brown (BL) and

white layer (WL) populations consist of 25 individually re-sequenced animals each, while the

two commercial broiler lines (BR1 and BR2) include 20 individually sequenced animals each.

For 41 populations, pooled DNA from 9 - 11 animals per population was sequenced, while

Gallus varius (green jungle fowl; GV) samples of only two animals were sequenced as a pool.

More detailed information about the samples can be found in **S1 File** and two previously

published papers, from Malomane *et al.* [30] and Qanbari *et al.* [40]. Coverage was between

7X and 10X for the individual sequences, while DNA pools were sequenced with 15X to 70X

coverage. Sequencing was conducted on Illumina HiSeq machines at the Helmholtz Zentrum,

German Research Center for Environmental Health in Munich, Germany.

Raw data preparation and SNP calling

Sequences were aligned to the reference genome Gallus gallus 5.0 [41,42] and the SNP

calling was conducted according to GATK Best Practices guidelines [43,44]. BWA-MEM 0.7.12

[45] was used for the alignment step, duplicates were marked using Picard Tools 2.0.1 [46]

MarkDuplicatesWithMateCigar and base qualities were recalibrated with GATK 3.7 [47]

BaseQualityRecalibrator. The set of known SNPs, necessary for base quality score recalibra-

tion, was downloaded from ENSEMBL release 87 [48]. SNPs were called for all samples sepa-

rately using the GATK 3.7 HaplotypeCaller and later on simultaneously genotyped across

samples with GATK 3.7 GenotypeGVCFs. Due to computational limitations, a ploidy of two

instead of the higher true ploidy of the pooled sequences had to be assumed, which led to

calling slightly less rare alleles. However, effects of this limitation are negligible (**S2 File; S1**

Fig).

SNP filtering was conducted using GATK 3.7 VariantRecalibrator, which filtered the called

SNPs by a machine learning approach (use of a Gaussian mixture model), which uses both, a

set of previous known (low confidence needed) and a set of highly reliable (assumed to be

true) variants as training resources [49]. The source for known SNPs (prior 2) provided to

VariantRecalibrator was again ENSEMBL (release 87) and the SNPs of the Axiom™ Genome-

Wide Chicken Array were defined as true training set (prior 15). The algorithm was trained

on the quality parameters DP, QD, FS, SOR, MQ and MQRankSum. Filters were set to recover

99 % of the training SNPs in the filtered set, which resulted in a Transition/Transversion ratio

of 2.52 for known SNPs, and a Transition/Transversion ratio of 2.26 for novel SNPs. Only

biallelic autosomal SNPs were used in all further analyses.

138   Identification of the ancestral allele

Ancestral alleles were defined using allele frequency information from the three wild

populations *Gallus gallus gallus* (GG), *Gallus gallus spadiceus* (GS) and *Gallus varius* (GV) by

an approach comparable to 50 [50]. It was assumed that the *Gallus gallus* and *Galllus varius*

species emerged from a common ancestor and *Gallus gallus* later split into *Gallus gallus gal-*

lus and *Gallus gallus spadiceus* subspecies. Additionally, assuming neutral molecular evolu-

tion [51], the ancestral allele was most likely the major allele within those three populations,

when weighting the allele frequency of *Gallus varius* twice. This procedure assigned the an-

cestral status to the reference allele for 86 % of the SNPs and to the alternative allele for 14

% of the SNPs. The change in the allele frequency spectrum was only relevant for the interval

from 0.95 – 1.00, which was reduced by 111'851 SNPs (0.39 % of all SNPs) when switching

from alternative to derived allele frequency (**S2 File; S2 Fig**).

Reference Sets

Three different reference sets were defined as follows: the **unfiltered WGS** SNPs (28.5 M

SNPs), SNPs filtered using GATK 3.7 [47] VariantRecalibrator (20.9 M SNPs; **filtered WGS**) and

array SNPs (540 k SNPs), which are the intersection of the unfiltered SNPs and the SNPs of

the Axiom™ Genome-Wide Chicken Array. The separate use of unfiltered and filtered WGS

SNPs was done to assess the effect of filtering (especially the use of an ascertained SNP set

as the true set) on ascertainment bias.

Redesigning the SNP Array

The populations were divided into four groups for the array design:

1) Discovery populations (8)

2) Validation populations (19)

3) Application populations (18)

4) Outgroup (1)

For SNP discovery firstly the 4 commercial lines (commercial white layers, WL; commer-

cial brown layers, BL and the two commercial broiler lines, BR1 and BR2) were used. The set

was then extended by additionally selecting those populations that were closest related to

each of the commercial populations based on pairwise Nei's standard genetic distance [52]

As the two broiler populations were closest related (**Fig 1**), the next two closest populations

were chosen. This resulted in the inclusion of White Leghorn (LE), Rhode Island Red (RI),

Marans (MR) and Rumpless Araucana (AR). Note that these populations are closely related

to the populations used as discovery populations for the development of the Axiom™ Ge-

nome-Wide Chicken Array [21] with exception to some inbred lines from the Roslin Institute

in Edinburgh we do not know the genetic origin of. The discovery set used for the original

array [21] additionally consisted of more animals than ours. Further, the discovery set had to

be split into broilers (BR1, BR2, MR, AR) and layers (WL, LE, BL, RI) for the equal spacing step.

From the remaining populations, 19 were randomly chosen for SNP validation (validation

populations), 18 populations (which were not included in the array development) were used

as showcase for an application of the array (application populations), and *Gallus varius* as a

different species was defined as outgroup. The interested reader can find all underlying

pairwise Nei's standard genetic distances [52] in **S3 File** and additionally pairwise F_{ST} values

[53] in **S4 File**.

Fig 1. UPGMA tree based on pairwise Nei's standard genetic distances. The tree was calculated from the fil-
tered WGS SNPs. Populations defined as layers or broilers, which form in total the discovery set for the array
design close to the original array, are highlighted. The plot was produced using the R package ape [54]. Note
that the plot is only supposed to reveal close clustering chicken populations and cannot be interpreted in depth
as chickens show a rich history of hybridization events. The interested reader can find all underlying pairwise
Nei's standard genetic distances [52] in **S3 File** and additionally pairwise FST values [53] in **S4 File**.

Based on the unfiltered SNP set, the sampling of the SNPs for an approximately 600 k

sized array was remodeled *in silico* in five consecutive steps according to the design process

of the original array which was described by Kranis *et al.* [21], starting from the unfiltered

SNP set:

1) **SNP discovery → 10.9 M SNPs**

194   Discovery of SNPs fulfilling basic criteria (quality ≥ 60; MAF ≥ 0.05; cover-

195   age ≤ mean + three standard deviations) within the discovery populations.

196   2) **Cluster removal → 8.8 M SNPs**

197   SNP clusters with less than 4 bp invariant sites at one side of a SNP (no matter of

whether 3' or 5' direction) and less than 10 bp invariant sites at the other side of

the SNP within the discovery populations were removed. This step is justified ra-

ther technically to enable probe binding, but could also lead to an overrepresenta-

tion of conserved regions compared to highly variable regions of the genome.

3) **Equal spacing → 2.1 M SNPs**

Reduction of SNPs to achieve approximately equidistant spacing between variable

SNPs within discovery populations based on genetic distances. The used algorithm

was modeled according to Kranis *et al.* [21] and followed a two-step procedure.

The first step was setting up an initial backbone of common SNPs (three sub

steps). It started with selecting SNPs which segregated in all discovery populations

(MAF within each population > 0) while requiring a minimal distance of 2 kb, re-

sulting in about 8 k SNPs. This was complemented by a backbone of SNPs which

segregated in all layer populations and another one of SNPs which segregated in

all broiler populations. Note that Kranis *et al.* [21] additionally constructed a back-

bone from a group of inbred lines for which no comparable samples were availa-

ble for this study. In the second step, the algorithm iterated over all single popula-

tions and filled in potential gaps between backbone SNPs which are variable with-

in the according population. This was done by choosing the SNPs closest to equi-

distant positions within the gap while aiming for a predefined local target density

of 667 segregating SNPs/cM (linkage map taken from [55]). See **S3 Fig** for the de-

tailed contribution of additional SNPs from each sub step of the algorithm.

4) **SNP validation → 1.7 M SNPs**

Removing SNPs (~ 20 %) which were not variable in at least 8 of the 19 validation

populations. This step would in reality be done by genotyping with preliminary

test arrays and therefore allows the use of a broader set of populations than the

discovery step.

5) **Downsampling → 580 k SNPs**

Downsampling of SNPs comparable to step 3, but without adding the broiler/ layer

specific backbones and instead keeping all exonic SNPs (annotation using Ensembl

VEP 89.7; 56). Additionally, the target density in broiler lines was set as three

times the target density of the layer lines. The increased target density in broilers

is intended to account for lower levels of linkage disequilibrium in these lines.

Variation of the design process

The whole design process was repeated 50 times with populations being randomly as-

signed to be discovery, validation or application populations, while the *Gallus varius* popula-

tion was always kept as the outgroup. In this process, the number of populations per group

was the same as in the previous scenario.

To assess the impact of the number of discovery populations on the design process, the

number of discovery populations was varied in additional runs from 4 to 40 randomly chosen

populations (while assigning the remaining populations, except *Gallus varius,* to validation

and application groups of equal size) with 20 random replicates for each number of discov-

ery populations. In a last scenario, equal spacing was varied with respect to the target densi-

ty (33 − 3333 SNPs/cM) with 20 independent population groupings for each target density,

with or without the initial backbone. As the number of SNPs from the backbone was con-

stant, the increase of the target density led to a higher number of SNPs chosen by the algo-

rithm due to the equal spacing itself and hence the relative influence of the fixed number of

common backbone SNPs decreased.

Analyses of the results

Per-locus-allele frequencies for individually sequenced populations were estimated from

genotypes, whereas the estimation for the sequenced DNA-pools was based on the allelic

depth. Influences on the allele frequency spectra were examined by comparing density esti-

mates of derived allele frequency spectra (unfolded frequency spectrum). Further H_{exp}, the

expected heterozygosity assuming Hardy Weinberg frequencies of the genotypes, for the

different populations were used as summary statistics of the within population allele fre-

quency spectra and calculated as in equation (1), where $p_{ref;l}$ denotes the frequency of the

reference allele at locus l and L the total number of loci.

254
$$H_{exp} = \frac{\sum_{l} 2 p_{ref;l}(1 - p_{ref;l})}{L} \qquad (1)$$

255   Deviations in the estimation of H_{exp} from the various SNP sets were quantified as differ-

256 ences between the H_{exp} calculated from the respective SNP set and the H_{exp} calculated from

257 the filtered WGS SNPs relative to the H_{exp} from the filtered WGS SNPs, further called overes-

258 timation of H_{exp} (OHE; equation (2)). An OHE of zero means that the estimates are equal,

259 while an OHE of one describes doubling of the unbiased estimate.

260
$$OHE = \frac{H_{exp;\ SNP\ set} - H_{exp;\ filtered\ WGS\ SNPs}}{H_{exp;\ filtered\ WGS\ SNPs}} \qquad (2)$$

The effects of the population group assignments on the OHE of the random population

assignments were evaluated by pairwise comparisons of least square means (LSMEANS; cal-

culated with the R package emmeans [57,58] by using Tukey correction for multiple pairwise

contrasts) of the population groups. An underlying mixed linear model for the estimation of

LSMEANS was fitted using the R package lme4 [59] as shown in equation (3), where the OHE

depended on an overall mean μ, the fixed effect of the population group $popG_i$ (i can be

discovery-, validation-, application- or outgroup), a random effect for the j[th] repetition of

random population grouping ($rep_j \sim N(0, I\sigma^2_{rep})$) and a random error $e_{ijk} \sim N(0, I\sigma^2_e)$. The

procedure is comparable to simple pairwise comparisons of group means, the correction by

the repetition only reduces the error variance and thus decreases the confidence intervals.

$$OHE_{ijk} = \mu + popG_i + rep_j + e_{ijk} \qquad (3)$$

Results

Numbers of SNPs

The SNP calling identified 28.5 M biallelic autosomal SNPs from which 20.9 M SNPs

passed GATK's filtering procedure. 540 k SNPs from the unfiltered WGS SNP set are also

mapped on the original Axiom™ Genome-Wide 580 k Chicken Array. The remodeling of the

array according to the design process of the original array returned 10.9 M SNPs from the

discovery step, which were reduced to approximately 580 k in steps as described. Numbers

of identified SNPs for the additional runs differed depending on the populations and settings

used and are listed in **S1 Table**. It has to be noted that the different sub steps of the equal

spacing algorithm contributed with different amounts of SNPs (**S3 Fig**). Especially the much

higher contribution of SNPs which were segregating in all broiler populations compared to

SNPs segregating in all layer populations in the first remodeling, which is due to closer rela-

tionships between the broiler populations and their generally higher heterozygosity, was

remarkable. Additional information about the identified number of SNPs depending on the

number of discovery populations and target density as well as information about the share

of SNPs of different random runs can be found in **S4 – S6 Figs**.

Underrepresentation of rare SNPs

A clear underrepresentation of rare SNPs in all ascertained SNP sets compared to WGS is

evident from the allele frequency spectra (**Fig 2**). Major changes in the allele frequency spec-

tra during the array development process were observed after the SNP discovery step and

the equal spacing step. The SNP discovery led to an underrepresentation of rare SNPs com-

pared to sequence data, which was intensified by the equal spacing step (**Fig 2**) and resulted

finally in a spectrum close to the spectrum of the original array. Randomly choosing popula-

tions as discovery populations confirmed the shape of the first remodeling, where the popu-

lation groups were chosen according to the original array [21]. As major changes in the spec-

tra mainly occurred after the SNP discovery and equal spacing, further results will concen-

trate on those steps.

Fig 2. Derived allele frequency spectra for the different SNP sets. For the remodeled sets, areas show the
modelling according to the original array [21] while grey lines represent the 50 random population groupings.

The allele frequency spectra (**Fig 3**) within discovery populations, compared to the spectra

over all populations, clearly showed the cutoff from the MAF 0.05 filter. Furthermore, the

allele frequency spectra of the discovery populations revealed a higher share of common

SNPs than the overall spectra after equal spacing. In contrast, the spectra within validation-

and application populations showed less pronounced peaks after the discovery step and the

outgroup (*Gallus varius*) revealed fixation of most SNPs variable in the discovery popula-

tions.

Fig 3. Derived allele frequency spectra within the population groups.

Influence of number of discovery populations and target density on allele frequency spectra

Not surprisingly, an increased number of discovery populations resulted in a higher num-

ber of rare alleles after the discovery step, and thus an allele frequency spectrum with a

more pronounced peak of rare alleles (**Fig 4**). Apparently, the shift of the allele frequency

spectrum after the equal spacing step was dependent on the number of discovery popula-

tions, as an increase in the number of discovery populations shifted the allele frequency

spectra towards a higher proportion of alleles with a low derived allele frequency. With an

increasing number of discovery populations, the shape of the allele frequency spectra got

closer to the spectrum of the original array.

Fig 4. Derived allele frequency spectra after SNP discovery (blue) and equal spacing (red) for varying numbers
of populations in the discovery set. Different numbers of populations in the discovery set (4 to 40) are indicat-
ed by an intensifying color gradient and only one randomly picked run per population number is shown. As the
differences in the color gradients are hard to distinguish, arrows in the respective color are indicating the shift
of the spectra with increasing numbers of discovery populations.

A very low target density, indicating that SNPs were mostly called due to being common

backbone SNPs, resulted in an allele frequency spectrum with the majority of alleles having a

MAF of around 0.5 (**Fig 5**). Increasing the target density for the equal spacing and thus reduc-

ing the influence of the initial backbone of common SNPs shifted the peak of the allele fre-

quency spectrum left towards a higher proportion of alleles with small derived allele fre-

quencies. Using only the backbone SNPs common over all discovery populations and thus

calling SNPs mostly by the equal spacing procedure resulted, independently from the target

density, in a spectrum similar to the one obtained with a high target density with backbone

(**Fig 5**).

Fig 5. Derived allele frequency spectra for different target densities [SNPs/cM] after the equal spacing step.
Blue spectra show the runs with the initial backbone included, red ones without backbone. Different target
densities are indicated by a log scaled increase in color intensity. Only one representative run per target density
is shown.

Overestimation of H_{exp}

Fig 6 shows the H_{exp} of different SNP sets by population. The H_{exp} obtained from the fil-

tered WGS SNPs were slightly higher than from the unfiltered WGS SNPs. H_{exp} obtained from

the ascertained SNP sets showed an even more pronounced overestimation together with

an increase during the design steps. In general, the correlations between the H_{exp} obtained

in the different SNP sets were relatively high ($\geq$ 0.95; **S2 Table**). Especially the H_{exp} of the two

WGS SNP sets showed a nearly perfect correlation of > 0.99, which led to an almost constant

OHE of -0.23 (**Table 1**) for the unfiltered WGS SNPs. As already recognizable from the H_{exp}

themselves, the OHE was positive for all ascertained SNP sets (0.66 − 1.29), which at the

same time showed a slightly reduced correlation to the filtered WGS SNP set (0.95 − 0.97).

Comparable to the allele frequency spectra, the most pronounced increase of the OHE was

caused by the SNP discovery and followed by the equal spacing step (OHE increased by

0.66), while the OHE from the original array SNPs (1.41; **Fig 6**; **Table 1**) laid in the range cov-

ered by the remodeling steps.

Fig 6. Expected Heterozygosity (H_{exp}) by population and SNP set. Populations are ordered by the H_{exp} of the
unfiltered WGS SNP set. Only the reference sets and relevant steps of the array design are shown. Discovery
populations are shaded with a darker background.

Table 1. OHE of the SNP sets from the first run

Populations	Unfiltered WGS	Array SNPs	1 – SNP discovery	2 - cluster removal	3 – equal spacing	4 – validation	5 – downsampling
All	$-0.23_{\pm 0.01}$	$0.84_{\pm 0.30}$	$0.66_{\pm 0.26}$	$0.66_{\pm 0.26}$	$1.09_{\pm 0.32}$	$1.29_{\pm 0.35}$	$1.27_{\pm 0.34}$
Discovery	$-0.23_{\pm 0.00}$	$1.05_{\pm 0.10}$	$0.86_{\pm 0.10}$	$0.86_{\pm 0.10}$	$1.15_{\pm 0.15}$	$1.28_{\pm 0.17}$	$1.32_{\pm 0.13}$
Validation	$-0.23_{\pm 0.00}$	$0.87_{\pm 0.13}$	$0.68_{\pm 0.10}$	$0.67_{\pm 0.10}$	$1.15_{\pm 0.13}$	$1.36_{\pm 0.14}$	$1.33_{\pm 0.12}$
Application	$-0.23_{\pm 0.00}$	$0.83_{\pm 0.12}$	$0.64_{\pm 0.07}$	$0.63_{\pm 0.07}$	$1.10_{\pm 0.11}$	$1.33_{\pm 0.14}$	$1.30_{\pm 0.15}$
Outgroup	-0.17	-0.88	-0.85	-0.86	-0.85	-0.85	-0.84

365 Mean OHE ± standard deviation.
366 An OHE of zero means no bias and an OHE of 1 means doubling the H_{exp}.
367

Averaging the OHE within the population groups revealed a 30 % higher OHE of the dis-

covery populations compared to validation and application populations after the discovery

step. The equal spacing step reduced this difference to an only 1 % larger OHE for discovery

populations, while it came with a substantial increase of the variance of OHE, which was

larger for the discovery populations than validation and application populations. The valida-

tion step then increased the OHE of the validation populations more than the OHE of discov-

ery and application populations. This stronger OHE of discovery populations was also appar-

ent within the array SNPs (24 % higher). In contrast to the other populations, the outgroup

showed an underestimation of the H_{exp}, resulting in an OHE of < -0.84 for all ascertained SNP

sets (**Fig 6**; **Table 1**).

A closer look on the contribution of the sub-steps during the equal spacing step revealed

that 62 % of the SNPs which were preserved during equal spacing were variable in all of the

four closely related broiler populations (BR1, BR2, MR, AR; maximum pairwise Nei's distance

of 0.06 and FST of 0.17 in the filtered SNP set), while only 3 % of the SNPs were retained due

to being variable in all of the four less closely related layer populations (WL, LE, BL, RI; max-

imum pairwise Nei's distance of 0.15 and FST of 0.48 in the filtered SNP set). The first popu-

lation used to fill in the gaps in the backbone (WL) contributed 17 % of the SNPs, while the

other populations contributed < 8 %.

Table 2. OHE of the SNP sets out of the 50 random population groupings

Populations	1 – SNP discovery	2 – cluster removal	3 – equal spacing	4 – validation	5 – down-sampling
Discovery	$0.76_{\pm0.004}^{a}$	$0.75_{\pm0.004}^{a}$	$1.13_{\pm0.006}^{a}$	$1.28_{\pm0.006}^{b}$	$1.33_{\pm0.007}^{a}$
Validation	$0.61_{\pm0.003}^{b}$	$0.60_{\pm0.003}^{b}$	$1.11_{\pm0.004}^{b}$	$1.29_{\pm0.004}^{b}$	$1.29_{\pm0.005}^{b}$
Application	$0.61_{\pm0.003}^{b}$	$0.61_{\pm0.003}^{b}$	$1.12_{\pm0.004}^{ab}$	$1.35_{\pm0.004}^{a}$	$1.34_{\pm0.004}^{a}$
Outgroup	$-0.85_{\pm0.008}^{c}$	$-0.86_{\pm0.008}^{c}$	$-0.85_{\pm0.015}^{c}$	$-0.84_{\pm0.017}^{c}$	$-0.84_{\pm0.019}^{c}$

388 LSMEANS for OHE ± standard error.
389 An OHE of zero means no bias and an OHE of 1 means doubling the H_{exp}.
390 Different lowercase letters within columns indicate significant differences to the 5 % level.
391

These findings were supported by the 50 random groupings (**Fig 7**). The LSMEANS (**Table**

2) of the population groups revealed 24 % larger OHE for discovery populations than for val-

idation and application populations after discovery and cluster removal step, which was de-

creased to a numerically insignificant difference after the equal spacing step. Interestingly,

and in contrast to the findings from the first remodeling, SNP validation led to a significantly

higher OHE (5 % larger) for application populations than discovery and validation popula-

tions.

Fig 7. OHE for discovery validation and application populations after the five steps of array design. Discovery
populations are chosen to represent populations which are comparable to the original array (blue) or 50 times
random sampled (grey).

404 Influence of number of discovery populations and target density on

H_{exp}

Fig 8 A shows that increasing the number of discovery populations reduces the mean OHE

of discovery populations after SNP discovery while not affecting the OHE of validation and

application populations. After equal spacing (**Fig 8 B**), no dependency of OHE on the number

of discovery populations remained. As, due to the limited number of populations in the

complete set, the number of validation populations decreased with more populations in the

discovery set, the individual validation populations gained more impact on the ascertain-

ment. This led to a higher OHE of validation populations with a high number of discovery

populations (**S7 D Fig**), comparable to the higher OHE of discovery populations for a small

number of discovery populations.

Fig 8. Relation of the OHE as a function of the number of discovery populations. A - discovery, **B** - equal spac-
ing. The Boxplots are only shown for a subset of the number of discovery populations, while the smoothing
lines, which show the trend, are calculated from all observations. Plots for all five steps can be found in **S7 Fig**.

In the equal spacing step, using only backbone SNPs resulted in a higher OHE for discov-

ery than for non- discovery populations. Increasing the target density and thus increasing the

proportion of SNPs due to the equal spacing part of the algorithm reduced the difference in

OHE between the population groups (**Fig 9 A**). If the SNPs from the initial backbone were not

used, no difference of OHE between discovery and non- discovery populations was present,

regardless of the target density (**Fig 9 B**).

Fig 9. OHE after equal spacing (step 3) by target density in SNPs/cM and population group. The smoothing
lines show the trend and the dashed lines the target density of 667 SNPs/cM, used for the remodeling accord-
ing to the original array [21]. The algorithm was run including the initial backbone SNPs (**A**) or not including
them (**B**). *Gallus varius* is not included, as it is constantly underestimated.

Discussion

In this study we used a uniquely diverse collection of sequenced wild, commercial and

non-commercial chicken populations, mainly based on samples of the Synbreed Chicken Di-

versity Panel [13]. Parts of our set were also involved in the development process of the Axi-

om™ Genome-Wide 580 k Chicken Array [21]. This offered an excellent possibility for as-

sessing the impact of ascertainment bias on real data in a complex scenario. In general, re-

sults derived in this study should therefore be comparable to other species. However, do-

mestic chickens show a rich history of hybridization and crossbreeding events [13]. The ef-

fects of using a discovery set closely related to the commercial populations and distributing

the discovery set randomly across the spectrum of populations were therefore comparably

small in this study. Special patterns of population structure e.g. the stronger differentiation

in cattle due to the two subspecies *Bos taurus* and *Bos indicus* [60] accompanied by limiting

the discovery set to one of the two clades, should increase the impact of population struc-

ture dependent ascertainment bias.

Potential impacts of the SNP calling pipeline

As the state of the art pipeline of GATK relies on a supervised machine learning approach

for filtering the SNP calls, which needs a highly reliable set of known SNPs, we started with

examining potential impacts of the filtering procedure on ascertainment bias. The number of

rare variants was slightly reduced by the filtering procedure and thus increased estimates of

H_{exp} were obtained in the filtered WGS set. As rare variants have a higher risk to be discard-

ed as sequencing errors [61], this reduction is expected when applying quality filters. How-

ever, a clear assessment of correctly and falsely filtered variants is not possible here and one

has to balance this tradeoff based on the study purpose.

Another source of ascertainment bias could be the use of array SNPs as training set for

GATK RecalibrateVariants, which potentially leads to discarding rare variants more likely if

they are not present in the discovery populations of the used array. As the correlation be-

tween the H_{exp} of the unfiltered and filtered WGS SNPs was nearly one, this source seems to

be negligible and the use of array variants as a highly reliable training set seems to be un-

problematic.

Due to computational limitations, we had to assume a ploidy of two for pools during the

SNP calling process, which resulted in a minimal reduction of rare alleles. However, this ef-

fect was shown to have a very minor impact on the findings of this study (**S2 File**).

General impact over all groups

The general reduction of rare alleles in array data compared to WGS data and the result-

ing overestimation of H_{exp} supports findings of previous studies [25,26,30,34]. This reduction

of rare alleles was mainly seen at steps where selection was explicitly biased towards high

MAF alleles (MAF filter for quality control in discovery step and use of common alleles for

the backbone in the equal spacing step) and/ or was applied to a small number of popula-

tions (small discovery set vs. small validation set). Thereby, the strongest shifts of the allele

frequency spectra and increases of H_{exp} are observed after SNP discovery and equal spacing.

Both, cluster removal and second downsampling had almost no effect on the allele frequen-

cy spectra and H_{exp}, while the validation step slightly decreased the share of rare SNPs.

The discovery step had the strongest impact on discovery populations, when a small set

of discovery populations was used (**Fig 8 A**). Similarly, the influence of the validation step on

validation populations was strongest in case of a small number of validation populations

(**S7D Fig**). A balancing of these two groups of samples is therefore necessary, if the number

of available DNA samples for array development is limited. We rather suggest the use of an

extended discovery set for validation as best practice instead of using completely different

populations.

If the equal spacing step contains a preselection of SNPs based on their variability within

population groups, the bias is stronger towards high MAF SNPs and thus yields a higher OHE.

This effect was reduced by increasing the target density and thus selecting relatively more

SNPs due to the equal spacing instead of common occurrence.

Differences between groups

If allele frequency spectra are changed in the same way for all populations and are there-

fore biasing heterozygosity estimates to the same extent, findings for between population

comparisons will be little affected. Ascertainment bias then is only of importance if one

compares populations based on different arrays, and corrections of the allele frequency

spectrum as reviewed by Nielsen [25] should be possible. As correlations between H_{exp} of

ascertained SNP sets and unfiltered/ filtered WGS SNP sets were consistently high (> 0.94),

arrays designed in the way as performed in this paper should mostly be suitable for robust

and cost efficient analyses. Biasedness of estimates could be reduced even more by consid-

ering filter strategies according to Malomane *et al.* [30].

However, we could show that the bias acts with different extent on different population

groups (population structure dependent bias) and therefore changes ranking of populations

and can affect conclusions. This population structure dependent bias was already shown to

have severe impact on findings from SNP arrays. For example, Bradbury *et al.* [62] found a

demographic decline up to an approximately 30 % lower H_{exp} for Atlantic cod based on the

distance to the sampling location of the discovery panel and McTravish and Hillis [29]

showed strong deviations between simulated and observed polymorphisms for different

combinations of migration- and ascertainment scenarios on simulated cattle populations. In

concordance with this, populations which are closely related to the discovery populations of

the original array in our study on average showed a 24 % higher OHE than validation and

application populations for the original array.

This population structure dependent bias was mainly introduced by the initial discovery

step. It was also observed for the random population groupings, but to a slightly different

extent. The difference in overestimation decreased with an increase in the number of dis-

covery populations (**Fig 8**) and was smallest if the discovery populations showed minimum

distance to the application and validation populations (results not shown). Comparable ob-

servations were already made by Frascaroli *et al.* [63] which found very small ascertainment

bias for European elite maize lines when using a SNP panel discovered in a combination of a

maize diversity set and inbred lines, but strong ascertainment bias when using SNPs which

were discovered in American elite lines. Therefore, we suggest to ideally choose an array

where the discovery panel does span the scope of populations it will be applied to, and by

this covers the existing variation in a most representative way, or to design such an array for

oneself if it does not exist.

The equal spacing step lowered the difference in mean OHE between population groups

in most of our remodeling scenarios, but obviously not in case of the original array. In the

remodeling, we saw this difference only with a low target density and thus calling SNPs in

the equal spacing step mainly due to being common over many populations (**Fig 9 A**). How-

ever, the equal spacing step also increased the variance of OHE in the discovery panel,

meaning that the OHE was increased more for some of the discovery populations than for

others, thus causing more uncertainty for resulting effects. This effect is driven by the une-

qual contribution of variable SNPs to the chosen SNP set by the different populations during

the equal spacing step (**S3 Fig**). The equal spacing step increases the OHE for some of the

discovery populations, while it decreases it for others, and hence it does not remove the

population structure dependent bias. This means that the knowledge of which discovery

populations were used is not sufficient to draw conclusions regarding a possible ascertain-

ment bias, since their relative contribution varies through the described pipeline.

Outgroup

Gallus varius as an outgroup showed a different behavior than all other populations. It al-

ready exhibited the lowest H_{exp} in the unfiltered WGS SNP set, which was most likely driven

by the small number of only two samples in the pool, and showed less upward bias of H_{exp} in

the filtered WGS SNP set than all other populations. The *Gallus varius* sequence reads on

average showed weak Phred-scaled mapping quality scores of 19 (1.3 % probability of misa-

lignment), while the mean quality scores of the other populations ranged from 25 (0.3 %) to

28 (0.1 %) – meaning that variation, only present in *Gallus varius*, will be more likely missed

due to misplacement of the reads or discarded as possible sequencing errors. Additionally,

every ascertained SNP set showed an OHE for *Gallus varius* of < -0.84, as variation being pre-

sent only in *Gallus varius* was not found in *Gallus gallus* discovery panels and contrary vari-

ants from *Gallus gallus* were not variable in *Gallus varius* (**Fig 3**). This demonstrates that ar-

rays should not be used if different species (even closely related ones) are included in the

research project. Even sequence based estimates can be slightly biased, if the reference ge-

nome does not fit properly.

Potential impact on other breeding applications

In general, we cannot infer the impact on breeding applications which require phenotypic

data (e.g. genomic selection [64] or genome wide association studies [65]) and/or individual-

ly sequenced or genotyped individuals (e.g. linkage disequilibrium decay [66] or runs of

homozygosity analyses [67]) from this study. However, literature highlights the increased

power of high MAF SNPs to capture/ detect effects which are caused by common variants

due to stronger linkage disequilibrium and higher levels of variance explained. Therefore,

increasing MAF in a first instance increases prediction accuracy when the number of SNPs is

limited [68] and therefore some SNP ascertainment schemes intentionally bias the used

SNPs towards high MAF within the desired populations [17]. The switch to WGS data, and

therefore the additional inclusion of rare alleles, is then expected to increase the possibility

of capturing the effects of rare alleles [68–70]. However, the increase in efficiency by higher

numbers of SNPs levels off when going towards WGS data [71]. Nevertheless, we would ex-

pect negative impacts of ascertainment bias due to the across population use of the arrays.

When biasing the genotyped variation towards the discovery population, the variability in

populations, which are less related to the discovery populations, is less increased or even

reduced, and arrays therefore become less valuable in non-target populations. Slight effects

of this were demonstrated by simulation [68] and we can clearly support these findings by

the levels of differences in the genotyped heterozygosity which we observed in this study.

For the effect of ascertainment bias on a broader set of applications, we further refer the

interested reader to studies which specifically address those issues (e.g. 30,31,66).

Further recommendations for future studies

We showed that existing arrays come with a large, and due to a diverse set of promoting

factors barely predictable, potential for ascertainment bias. Strongly decreasing costs for

WGS and increasing availability of powerful computing resources therefore promote an in-

tensified use of WGS for population genetic analyses, especially when diverse populations

are included in the studies. However, costs and computational effort will still be substantial

for large scale projects. Possible cost effective alternatives could be reduced library sequenc-

ing approaches like Genotyping-by-Sequencing [61,72], even though such methods intro-

duce other problems related to the use of restriction enzymes which are reviewed by An-

drews *et al.* [73].

For the purpose of monitoring genetic diversity in a large set of small non-commercial

populations, the development of a specialized new array for cost effective high throughput

genotyping could be still a good option. For the design of such an array, it is crucial to extend

the discovery set in a way which represents the total variability over all populations as bal-

anced as possible. The use of publicly available sequences can be helpful to reach this goal.

The ascertainment of the SNPs should then be done preferably over a large set of highly di-

verse populations covering a wide spectrum of the diversity within a species available popu-

lations instead of biasing the process towards common alleles by performing within popula-

tion ascertainment.

However, when considering the findings of this study, choosing an array for which the

SNP ascertainment process is well documented and a discovery panel that matches the pop-

ulations to be studied, should enable researchers to derive appropriate conclusions when

robust estimators are used.

590 Acknowledgments

591 We are grateful to all the participating breeders for their assistance in sampling.

592 References

1. Novembre J, Johnson T, Bryc K, Kutalik Z, Boyko AR, Auton A, et al. Genes mirror geog-
raphy within Europe. Nature. 2008; 456:98. doi: 10.1038/nature07331.
2. Patterson N, Moorjani P, Luo Y, Mallick S, Rohland N, Zhan Y, et al. Ancient admixture in
human history. Genetics. 2012; 192:1065–93. doi: 10.1534/genetics.112.145037 PMID:
22960212.
3. Laurie CC, Nickerson DA, Anderson AD, Weir BS, Livingston RJ, Dean MD, et al. Linkage
Disequilibrium   in   Wild   Mice.   PLoS   Genet.   2007;   3:e144.
doi: 10.1371/journal.pgen.0030144.
4. Platt A, Horton M, Huang YS, Li Y, Anastasio AE, Mulyati NW, et al. The Scale of Popula-
tion   Structure   in   Arabidopsis   thaliana.   PLoS   Genet.   2010;   6:e1000843.
doi: 10.1371/journal.pgen.1000843.

5.   Travis AJ, Norton GJ, Datta S, Sarma R, Dasgupta T, Savio FL, et al. Assessing the genetic
diversity of rice originating from Bangladesh, Assam and West Bengal. Rice. 2015; 8:35.
doi: 10.1186/s12284-015-0068-z.

6.   Mayer M, Unterseer S, Bauer E, Leon N de, Ordas B, Schön C-C. Is there an optimum
level of diversity in utilization of genetic resources. Theor Appl Genet. 2017; 130:2283–
95. doi: 10.1007/s00122-017-2959-4.

7.   Muir WM, Wong GK-S, Zhang Y, Wang J, Groenen MAM, Crooijmans RPMA, et al. Ge-
nome-wide assessment of worldwide chicken SNP genetic diversity indicates significant
absence of rare alleles in commercial breeds. Proc Natl Acad Sci. 2008; 105:17312–7.
doi: 10.1073/pnas.0806569105.

8.   Gibbs RA, Taylor JF, van Tassell CP, Barendse W, Eversole KA, Gill CA, et al. Genome-
wide survey of SNP variation uncovers the genetic structure of cattle breeds. Science.
2009; 324:528–32. doi: 10.1126/science.1167936 PMID: 19390050.

9.   Kijas JW, Townley D, Dalrymple BP, Heaton MP, Maddox JF, McGrath A, et al. A genome
wide survey of SNP variation reveals the genetic structure of sheep breeds. PLoS One.
2009; 4:e4668.

10.   Gautier M, Laloe D, Moazami-Goudarzi K. Insights into the genetic history of French
cattle from dense SNP data on 47 worldwide breeds. PLoS One. 2010; 5.
doi: 10.1371/journal.pone.0013038 PMID: 20927341.

11.   Qanbari S, Pimentel ECG, Tetens J, Thaller G, Lichtner P, Sharifi AR, et al. A ge-
nome-wide scan for signatures of recent selection in Holstein cattle. Animal genetics.
2010; 41:377–89.

12.   McTavish EJ, Decker JE, Schnabel RD, Taylor JF, Hillis DM. New World cattle show an-
cestry from multiple independent domestication events. Proc Natl Acad Sci. 2013;
110:E1398-E1406.

13.   Malomane DK, Simianer H, Weigend A, Reimer C, Schmitt AO, Weigend S. The
SYNBREED chicken diversity panel. A global resource to assess chicken diversity at high
genomic resolution. BMC Genomics. 2019; 20:345. doi: 10.1186/s12864-019-5727-9
PMID: 31064348.

14.   Perkel J. SNP genotyping. Six technologies that keyed a revolution. Nature Methods.
2008; 5:447. doi: 10.1038/nmeth0508-447.

15.   Unterseer S, Bauer E, Haberer G, Seidel M, Knaak C, Ouzunova M, et al. A powerful tool
for genome analysis in maize: development and evaluation of the high density 600 k
SNP genotyping array. BMC Genomics. 2014; 15:823.

16.   Singh N, Jayaswal PK, Panda K, Mandal P, Kumar V, Singh B, et al. Single-copy gene
based 50 K SNP chip for genetic studies and molecular breeding in rice. Sci Rep. 2015;
5:11600.

17.   Matukumalli LK, Lawley CT, Schnabel RD, Taylor JF, Allan MF, Heaton MP, et al. Devel-
opment and characterization of a high density SNP genotyping assay for cattle. PLoS
One. 2009; 4:e5350.

18.   Ramos AM, Crooijmans RPMA, Affara NA, Amaral AJ, Archibald AL, Beever JE, et al. De-
sign of a high density SNP genotyping assay in the pig using SNPs identified and charac-
terized by next generation sequencing technology. PLoS One. 2009; 4:e6524.

19. Groenen MAM, Megens H-J, Zare Y, Warren WC, Hillier LW, Crooijmans RPMA, et al.
The development and characterization of a 60K SNP chip for chicken. BMC Genomics.
2011; 12:274.

20. Boichard DA, Chung H, Dassonneville R, David X, Eggen A, Fritz S, et al. Design of a bo-
vine low-density SNP array optimized for imputation. PLoS One. 2012; 7:e34130.

21. Kranis A, Gheyas AA, Boschiero C, Turner F, Le Yu, Smith S, et al. Development of a high
density 600K SNP genotyping array for chicken. BMC Genomics. 2013; 14:59.
doi: 10.1186/1471-2164-14-59.

22. Tosser-Klopp G, Bardou P, Bouchez O, Cabau C, Crooijmans RPMA, Dong Y, et al. Design
and characterization of a 52K SNP chip for goats. PLoS One. 2014; 9:e86227.

23. Sandenbergh L, Cloete SWP, Roodt-Wilding R, Snyman MA, Bester-van der Merwe AE.
Evaluation of the OvineSNP50 chip for use in four South African sheep breeds. S Afr J
Anim Sci. 2016; 46:89–93.

24. Fan B, Du Z-Q, Gorbach DM, Rothschild MF. Development and application of high-
density SNP arrays in genomic studies of domestic animals. Asian-Australas J Anim Sci.
2010; 23:833–47.

25. Nielsen R. Population genetic analysis of ascertained SNP data. Hum Genomics. 2004;
1:1.

26. Clark AG, Hubisz MJ, Bustamante CD, Williamson SH, Nielsen R. Ascertainment bias in
studies of human genome-wide polymorphism. Genome Res. 2005; 15:1496–502.

27. Eller E. Effects of Ascertainment Bias on Recovering Human Demographic History. Hu-
man Biology. 2001; 73:411–27. doi: 10.1353/hub.2001.0034.

28. Nielsen R, Signorovitch J. Correcting for ascertainment biases when analyzing SNP data:
applications to the estimation of linkage disequilibrium. Theor Popul Biol. 2003;
63:245–55.

29. McTavish EJ, Hillis DM. How do SNP ascertainment schemes and population de-
mographics affect inferences about population history. BMC Genomics. 2015; 16:1.

30. Malomane DK, Reimer C, Weigend S, Weigend A, Sharifi AR, Simianer H. Efficiency of
different strategies to mitigate ascertainment bias when using SNP panels in diversity
studies. BMC Genomics. 2018; 19:22. doi: 10.1186/s12864-017-4416-9.

31. Lachance J, Tishkoff SA. SNP ascertainment bias in population genetic analyses: why it is
important, and how to correct it. Bioessays. 2013; 35:780–6.

32. Nielsen R, Hubisz MJ, Clark AG. Reconstituting the frequency spectrum of ascertained
single-nucleotide polymorphism data. Genetics. 2004; 168:2373–82.

33. Guillot G, Foll M. Correcting for ascertainment bias in the inference of population struc-
ture. Bioinformatics. 2009; 25:552–4. doi: 10.1093/bioinformatics/btn665 PMID:
19136550.

34. Albrechtsen A, Nielsen FC, Nielsen R. Ascertainment biases in SNP chips affect
measures of population divergence. Mol Biol Evol. 2010; 27:2534–2547.

35. West B, Zhou B-X. Did chickens go north? New evidence for domestication. Journal of
archaeological science. 1988; 15:515–33.

36. Lawal RA, Martin SH, Vanmechelen K, Vereijken A, Silva P, Al-Atiyat RM, et al. The wild
species genome ancestry of domestic chickens. BMC Biology. 2020; 18:13.
doi: 10.1186/s12915-020-0738-1.

37. Tixier-Boichard M, Bed'hom B, Rognon X. Chicken domestication. From archeology to
genomics. Comptes rendus biologies. 2011; 334:197–204.

38. Eriksson J, Larson G, Gunnarsson U, Bed'hom B, Tixier-Boichard M, Strömstedt L, et al.
Identification of the Yellow Skin Gene Reveals a Hybrid Origin of the Domestic Chicken.
PLoS Genet. 2008; 4:e1000010. doi: 10.1371/journal.pgen.1000010.

39. Crawford RD. Poultry genetic resources. evolution, diversity and conservation. In: Craw-
ford RD, editor. Poultry breeding and genetics. 2nd ed. Amsterdam: Elsevier; 1993.

40. Qanbari S, Rubin C-J, Maqbool K, Weigend S, Weigend A, Geibel J, et al. Genetics of
adaptation in modern chicken. PLoS Genet. 2019; 15:e1007989.
doi: 10.1371/journal.pgen.1007989.

41. Reference Genome Gallus gallus 5.0. UCSC 2016 [cited 25 Oct 2016]. Available from:
http://hgdownload.soe.ucsc.edu/goldenPath/galGal5/bigZips/galGal5.fa.gz.

42. Warren WC, Hillier LW, Tomlinson C, Minx P, Kremitzki M, Graves TA, et al. A New
Chicken Genome Assembly Provides Insight into Avian Genome Structure. G3. 2017;
7:109–17. doi: 10.1534/g3.116.035923.

43. DePristo MA, Banks E, Poplin R, Garimella KV, Maguire JR, Hartl C, et al. A framework
for variation discovery and genotyping using next-generation DNA sequencing data. Nat
Genet. 2011; 43:491. doi: 10.1038/ng.806.

44. van der Auwera GA, Carneiro MO, Hartl C, Poplin R, Del Angel G, Levy-Moonshine A, et
al. From FastQ data to high confidence variant calls. The Genome Analysis Toolkit best
practices pipeline. Curr Protoc Bioinformatics. 2013; 43:11.10.1-11.10.33.
doi: 10.1002/0471250953.bi1110s43 PMID: 25431634.

45. Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
2013 [updated 26 May 2013; cited 21 Mar 2019]. Available from:
https://arxiv.org/pdf/1303.3997.

46. Picard Tools 2.0.1. Broad Institute 2015. Available from:
https://broadinstitute.github.io/picard/.

47. McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, et al. The Ge-
nome Analysis Toolkit. A MapReduce framework for analyzing next-generation DNA se-
quencing data. Genome Res. 2010; 20:1297–303. doi: 10.1101/gr.107524.110.

48. ENSEMBL. Chicken Germline SNP and INDELS. 2016 [cited 6 Jan 2017]. Available from:
http://e87.ensembl.org/Gallus_gallus.

49. Broad Institute. GATK User Guide. 2018 [cited 20 Mar 2018]. Available from:
https://software.broadinstitute.org/gatk/documentation/.

50. Rocha D, Billerey C, Samson F, Boichard D, Boussaha M. Identification of the putative
ancestral allele of bovine single-nucleotide polymorphisms. J Anim Breed Genet. 2014;
131:483–6. doi: 10.1111/jbg.12095.

51. Kimura M. The neutral theory of molecular evolution. A review of recent evidence. Jpn J
Genet. 1991; 66:367–86.

52. Nei M. Genetic Distance between Populations. The American Naturalist. 1972;
106:283–92.

53. Wright S. The genetical structure of populations. Ann Eugen. 1949; 15:323–54.
doi: 10.1111/j.1469-1809.1949.tb02451.x.

54. Paradis E, Claude J, Strimmer K. APE. Analyses of phylogenetics and evolution in R lan-
guage. Bioinformatics. 2004; 20:289–90.

55. Groenen MAM, Wahlberg P, Foglio M, Cheng HH, Megens H-J, Crooijmans RPMA, et al.
A high-density SNP-based linkage map of the chicken genome reveals sequence fea-
tures correlated with recombination rate. Genome Res. 2009; 19:510–9.

56. McLaren W, Gil L, Hunt SE, Riat HS, Ritchie GRS, Thormann A, et al. The Ensembl Variant
Effect Predictor. Genome Biol. 2016; 17:122. doi: 10.1186/s13059-016-0974-4.

57. Lenth R. emmeans: Estimated Marginal Means, aka Least-Squares Means. ; 2019.

58. R Core Team. R. A Language and Environment for Statistical Computing. Vienna, Aus-
tria; 2017.

59. Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models Using
lme4. Journal of Statistical Software. 2015; 67:1–48. doi: 10.18637/jss.v067.i01.

60. Hiendleder S, Lewalski H, Janke A. Complete mitochondrial genomes of *Bos taurus* and
Bos indicus provide new insights into intra-species variation, taxonomy and domestica-
tion. Cytogenet Genome Res. 2008; 120:150–6. doi: 10.1159/000118756.

61. Heslot N, Rutkoski J, Poland JA, Jannink J-L, Sorrells ME. Impact of marker ascertain-
ment bias on genomic selection accuracy and estimates of genetic diversity. PLoS One.
2013; 8:e74612.

62. Bradbury IR, Hubert S, Higgins B, Bowman S, Paterson IG, Snelgrove PVR, et al. Evalu-
ating SNP ascertainment bias and its impact on population assignment in Atlantic cod,
Gadus morhua. Mol Ecol Res. 2011; 11:218–25.

63. Frascaroli E, Schrag TA, Melchinger AE. Genetic diversity analysis of elite European
maize (Zea mays L.) inbred lines using AFLP, SSR, and SNP markers reveals ascertain-
ment bias for a subset of SNPs. Theoretical and Applied Genetics. 2013; 126:133–41.
doi: 10.1007/s00122-012-1968-6.

64. Meuwissen THE, Hayes BJ, Goddard ME. Prediction of Total Genetic Value Using Ge-
nome-Wide Dense Marker Maps. Genetics. 2001; 157:1819–29. Available from:
https://www.genetics.org/content/157/4/1819.

65. Goddard ME, Hayes BJ. Mapping genes for complex traits in domestic animals and their
use in breeding programmes. Nat Rev Genet. 2009; 10:381–91. doi: 10.1038/nrg2575.

66. Qanbari S, Pausch H, Jansen S, Somel M, Strom T-M, Fries R, et al. Classic selective
sweeps revealed by massive sequencing in cattle. PLoS Genet. 2014; 10:e1004148.
doi: 10.1371/journal.pgen.1004148 PMID: 24586189.

67. Peripolli E, Munari DP, Silva, M. V. G. B., Lima ALF, Irgang R, Baldi F. Runs of
homozygosity: current knowledge and applications in livestock. Anim Genet. 2017;
48:255–71. doi: 10.1111/age.12526.

68. Perez-Enciso M, Rincon JC, Legarra A. Sequence- vs. chip-assisted genomic selection:
accurate   biological   information   is   advised.   Genet   Sel   Evol.   2015;   47:43.
doi: 10.1186/s12711-015-0117-5 PMID: 25956961.

69. Druet T, Macleod IM, Hayes BJ. Toward genomic prediction from whole-genome se-
quence data: impact of sequencing design on genotype imputation and accuracy of
predictions. Heredity. 2014; 112:39–47. doi: 10.1038/hdy.2013.13 PMID: 23549338.

70. Wainschtein P, Jain DP, Yengo L, Zheng Z, Cupples LA, Shadyab AH, et al. Recovery of
trait   heritability   from   whole   genome   sequence   data.   bioRxiv.   2019.
doi: 10.1101/588020.

71. Ober U, Ayroles JF, Stone EA, Richards S, Zhu D, Gibbs RA, et al. Using whole-genome
sequence data to predict quantitative trait phenotypes in Drosophila melanogaster.
PLoS Genet. 2012; 8:e1002685. doi: 10.1371/journal.pgen.1002685 PMID: 22570636.
72. Elshire RJ, Glaubitz JC, Sun Q, Poland JA, Kawamoto K, Buckler ES, et al. A robust, simple
genotyping-by-sequencing (GBS) approach for high diversity species. PLoS One. 2011;
6:e19379.
73. Andrews KR, Good JM, Miller MR, Luikart G, Hohenlohe PA. Harnessing the power of
RADseq for ecological and evolutionary genomics. Nat Rev Genet. 2016; 17:81.
787

Supporting information

S1 File. Abbreviations for breeds and accession numbers for the sequencing data.

S2 File. Supplementary methods.

S3 File. Pairwise Nei's standard genetic distances.

S4 File. Pairwise F_{ST} values.

S1 Fig. Pooled vs. ploidy two calling. Expected Heterozygosity (H_{exp}) from the calling with

assuming the correct ploidy vs. assuming ploidy two for all samples (A) and alternative allele

frequency distributions of called alleles for individually sequenced (B) and pooled sequenced

(C) populations.

S2 Fig. Alternative allele frequency spectrum for SNPs where reference allele is not ances-

tral allele (A) and alternative (B) vs. derived (C) allele frequency spectrum of all SNPs.

S3 Fig. Cumulative number of SNPs in million retained during the equal spacing step. The

red line and points represent the first remodeling according to the original array [21], while

the dashed lines and the boxplots represent the 50 random population groupings and the

black line the according median values. The algorithm starts with a very basic initial back-

bone and then adds SNPs to the backbone which are variable in either all layer lines or all

broiler lines. Separated by a vertical line, the second part of the algorithm successively fills

up potential gaps to achieve an equidistant coverage of 667 segregating SNPs/cM for each

discovery population.

S4 Fig. SNPs retained by number of populations in the discovery set.

S5 Fig. Number of SNPs retained after the equal spacing step by target density.

S6 Fig. Relative amount of SNPs shared by a specific number of runs from 50 random popu-

lation assignments.

S7 Fig. Relation of the OHE as a function of the number of discovery populations.

A - discovery, **B** – cluster removal, **C** – equal spacing, **D** – validation, **E** – downsampling. The

Boxplots are only shown for a subset of the number of discovery populations, while the

smoothing lines, which show the trend, are calculated from all observations.

S1 Table. Number of SNPs from the remodeling processes

S2 Table. Pearson correlations between the H_{exp} of the different SNP sets

A

B