Competition-by-drought interactions change phenotypic plasticity and the direction of selection on Arabidopsis traits

Claire M. Lorts[1], Jesse R. Lasky[1,2]

[1]Department of Biology
Pennsylvania State University

[2]Correspondence: lasky@psu.edu

Summary

1. Populations often exhibit genetic diversity in traits involved in responses to abiotic stressors, but what maintains this diversity is unclear. *Arabidopsis thaliana* exhibits high within-population variation in drought response. One hypothesis is that competition, varying at small scales, promotes diversity in resource use strategies. However, little is known about natural variation in competition effects on Arabidopsis physiology.

2. We imposed drought and competition treatments on diverse genotypes. We measured resource economics traits, physiology, and fitness to characterize plasticity and selection in response to treatments.

3. Plastic responses to competition differed depending on moisture availability. We observed genotype-drought-competition interactions for relative fitness: competition had little effect on relative fitness under well-watered conditions, while under drought competition caused rank changes in fitness. Early flowering was always selected. Higher $\delta^{13}C$ was selected only in the harshest treatment (drought and competition). Competitive context significantly changed the direction of selection on aboveground biomass and inflorescence height in well-watered environments.

4. Our results highlight how local biotic conditions modify abiotic selection, in some cases promoting diversity in abiotic stress response. The ability of populations to adapt to environmental change may thus depend on small-scale biotic heterogeneity.

Keywords: succession, tradeoffs, leaf economics, neighbor interactions, strigolactone

INTRODUCTION

The persistence of genetic variation in phenotypes in spite of the forces eliminating diversity is a central puzzle in biology (Mitchell-Olds *et al.*, 2007). While experiments often show strong selection under abiotic stress, populations still exhibit (perhaps surprising) diversity in response to such treatments (Hosgood & Parsons, 1968; Sultan & Bazzaz, 1993). Understanding diversity within populations is an important research goal partly because this diversity affects the adaptability of populations to environmental change (Pease *et al.*, 1989; Lynch & Lande, 1993; Jump *et al.*, 2009).

The effects of climate conditions (like precipitation) on organisms occur within the context of fine scale environmental heterogeneity, which can exacerbate or mitigate climatic conditions (Farrior *et al.*, 2013b; Urban *et al.*, 2013; Clark *et al.*, 2014; Moran & Alexander, 2014). This heterogeneity might select for phenotypic plasticity in a single optimal genotype. Alternatively, heterogeneity can promote genetic variation within populations by favoring different genotypes in different environments, *i.e.* a genotype-by-environment (G×E) interaction causing a fitness tradeoff across environmental gradients (Levene, 1953; Hamrick & Holden, 1979; Schmitt *et al.*, 2003).

Limiting resources that are influenced by climate, *e.g.* moisture, may be especially likely to exhibit interactions with competition. In plants for example, different levels of allocation to roots for water acquisition may be favored depending on the presence of neighboring individuals that compete for water (Farrior *et al.*, 2013a). Differing water use efficiencies (WUE) can be favored under moisture limitation depending on whether competitors are present (Campitelli *et al.*, 2016). Specifically, competition can favor more resource acquisitive strategies when resources are limited (Farrior *et al.*, 2013a).

Our current study system *Arabidopsis thaliana* (hereafter, Arabidopsis) is useful for studying these processes for several reasons. First, it exhibits substantial diversity in drought response within populations. Studies of genetic variation in $\delta^{13}C$, a proxy for water use efficiency (WUE), and stomata have found statistically significant but weak relationships with climate of origin (Dittberner *et al.*, 2018) as have studies of survival (Exposito-Alonso *et al.*, 2018). These weak relationships may be surprising given the tendency for local adaptation in Arabidopsis populations (Fournier-Level *et al.*, 2011) and stand in contrast with cold tolerance, which is more closely tied to local winter temperatures (Ågren & Schemske, 2012; Monroe *et*

al., 2016; Gienapp *et al.*, 2017). Second, while Arabidopsis occurs in a wide range of precipitation regimes (Hoffmann, 2002) it also occupies a wide range of microhabitats (Baron *et al.*, 2015), suggesting the opportunity for precipitation-by-microhabitat interactions. Some

70 published data suggest the potential for competition-by-climate interaction effects on performance (Baron *et al.*, 2015; Campitelli *et al.*, 2016) though the physiological basis of these interactions is largely unknown. Third, previous research has yielded insight into genetic and physiological mechanisms of drought adaptation, providing a basis for specific hypotheses about how moisture and competitive gradients interact to select for different traits (McKay *et al.*, 2003;

75 Juenger *et al.*, 2010; Des Marais *et al.*, 2012; Lovell *et al.*, 2013; Kenney *et al.*, 2014; Lasky *et al.*, 2014, 2018; El-Soda *et al.*, 2015; Bac-Molenaar *et al.*, 2016; Dittberner *et al.*, 2018; Exposito-Alonso *et al.*, 2018, 2019; DeLeo *et al.*, 2019).

Arabidopsis drought responses have been characterized into a stereotypical tradeoff between avoidance and escape (McKay *et al.* 2003; Kenney *et al.* 2014). Avoidance involves

80 phenotypes that minimize water loss (*e.g.* stomatal closure) and maximize gain (*e.g.* root growth). Escape involves phenology and rapid development that matches with time periods when moisture is abundant (Ludlow, 1989). Escape may be adaptive when low soil moisture occurs at the end of growing seasons (Kenney *et al.*, 2014) while avoidance may be adaptive when moisture is limited throughout a growing season (Campitelli *et al.*, 2016). Escape has been

85 characterized by rapid flowering, high specific leaf area (SLA), high leaf N, high carbon assimilation rate (*A*), low root:shoot mass ratio, and low water use efficiency (WUE) (Ludlow, 1989; Des Marais *et al.*, 2012; Kenney *et al.*, 2014; El-Soda *et al.*, 2015), while avoidance is characterized by the opposite values (but see Ferguson *et al.*, 2019). Arabidopsis is a rapidly maturing annual, an early successional strategy, which may be adaptive when neighbors are

90 better resource competitors. In disturbed habitats where Arabidopsis frequently occurs, competition likely increases over time as colonists grow. In this situation, moisture availability may decrease across a season due to greater uptake combined with increasing evaporative demand as temperatures warm, favoring drought escape (Cohen, 1970; McLeod *et al.*, 2012; Campitelli *et al.*, 2016).

95 Despite the importance of Arabidopsis as the model plant, there exists little information on how diverse ecotypes respond physiologically to the presence of neighboring plants. Plant responses to neighbors may use a variety of cues, including nutrient availability (Schenk, 2006),

red:far red light ratios (Aphalo *et al.*, 1999), or chemicals (Badri *et al.*, 2012). In Arabidopsis, Campitelli *et al.* (2016) found large biomass reductions and lower WUE in response to grass neighbors, even under ample water supply. One non-resource based mechanism for sensing neighbors is perception of root exudates, which are used by diverse species for belowground sensing of neighbors (Badri *et al.*, 2012; Schmid *et al.*, 2013; van Dam & Bouwmeester, 2016). Strigolactones (SLs) are exuded belowground by most plants (Al-Babili & Bouwmeester, 2015) and have been implicated in neighbor avoidance in a moss (Proust *et al.*, 2011). Furthermore, Arabidopsis plants can suppress inflorescence branching (Scaffidi *et al.*, 2014) and close stomata (Lv *et al.*, 2018) in response to SLs in growth media. To the extent that apical dominance in the presence of competitors enhances fitness (Baron *et al.*, 2015) it may be adaptive for a plant to suppress branching in response to SLs of neighbors.

We studied how competition interacts with moisture to drive plasticity and selection on life history and physiology in Arabidopsis. While additional resources vary with competition, *e.g.* nutrients, we controlled these in our experiments to focus on competition for water.

We focused on these specific questions:

1. How does Arabidopsis respond to competition and drought via changes in life history and physiology? We hypothesized that drought would induce greater WUE due to stomatal closure, reduced performance, and greater SLA and leaf N as part of a drought escape strategy. We hypothesized that competition would induce inflorescence elongation, lower performance (biomass and fitness), and greater investment in resource acquisition (greater SLA, leaf N, lower WUE), partly due to perception of SLs in soil.

2. Does competition induce rank-changing G×E in fitness, suggesting that competitive gradients could promote diversity in response to abiotic stress within populations? We hypothesized that under conditions with ample resources, the presence of neighboring plants would have little influence on the relative fitness of diverse genotypes. By contrast, under moisture limitation we expected that competition would favor genotypes that differ in multiple traits compared to when competitors are absent.

3. What traits explain genetic variation in fitness response to competition (G×E)? We hypothesized that genetic tradeoffs are associated with alternate drought response strategies, such that contrasting treatments change the sign of directional selection coefficients across treatments.

4. Do traits exhibit genetic variation in plasticity that is under selection across competitive environments? We hypothesized that changes in some traits across competitive environments would be associated with higher average fitness across competitive environments.

METHODS

Plant material

We studied 17 diverse genotypes from across the native range of Arabidopsis and the reference genotype (Columbia) for a total of 18 genotypes (Tables S1 and S2). These genotypes mostly represent diverse lines used to create a multi-parent linkage mapping population (Kover *et al.*, 2009). We obtained seeds from the Arabidopsis Biological Resource Center (ABRC). However, it was later discovered that two lines (Can-0 CS76740, and Wu-0 CS78858) were contaminants from the supplying lab, *i.e.* they were not the original labeled genotype, though they were unique unidentified genotypes (Pisupati *et al.*, 2017) and we included them in the experiments. Seeds were grown for one generation in the lab at 16ºC with a 16-hr photoperiod and the resulting next generation of seed was used in the present experiments. For competitors, we used three genotypes (Adi-10, Bd21-3, and Bd30-1) of *Brachypodium distachyon*, a widespread Mediterranean annual grass. Both Arabidopsis and *B. distachyon* occur broadly across mesic Mediterranean sites, making them potential natural competitors. However, these plant communities are diverse (Alados *et al.*, 2004) and it is likely selection on species interactions has been diffuse and not specific to this pair of species. The *B. distachyon* genotypes had similar growth form and rate and were simultaneously bulked in a greenhouse.

Experimental design and conditions

To characterize genetic variation in response to moisture differences and their interactions with competition, we grew these genotypes in a split-plot design with two crossed treatments (for a total of four conditions): (1) well-watered, no competition (*i.e.* a single Arabidopsis plant in a pot), (2) well-watered, with competition, (3) intermittent drought, no competition, and (4) intermittent drought, with competition (Fig. S1). Each Arabidopsis genotype in each treatment

had 10 total replicates per treatment, 5 for an initial harvest prior to flowering, and 5 for a post-senescence harvest (total Arabidopsis plants was 720).

We grew plants in a walk-in Conviron C1210 growth chamber (Conviron Ltd., Winnipeg, Canada). We designed a temperature and photoperiod program (Fig. S3, Table S3) to mimic a growing season starting with warm temperatures early in the local growing season, which we reasoned would be a potential site for moisture limitation, and centrally located in the species range (Online Supplement).

Competition was imposed by sowing six seeds of *B. distachyon*, along the outside perimeter of each pot. To focus on competition for water, we limited light competition by restraining *B. distachyon* tillers to the pot perimeter with a mounted stainless steel ring (Fig. S2). Drought treatments were imposed by removing bottom water at 16 days after planting (DAP), when all plants had at least the first true leaves fully expanded (Online Supplement).To limit nutrient competition pots in all treatments were fertigated at the same time of day with the same nutrition strength and volume.

Plant harvest and phenotyping during vegetative stage

To measure phenotypes during vegetative growth that required destructive harvest, we randomly selected five replicates for each genotype and treatment combination for harvest at 55-56 DAP, just prior to the initiation of the first flowering genotypes. In this harvest, *B. distachyon* shoots were cut at soil level, dried at 90 °C, and weighed. Roots of Arabidopsis plants under no competition treatments were washed, dried at 90 °C, and weighed (roots in competition were too difficult to separate from *B. distachyon* roots). Rosettes were removed from pots 1-2 cm below the soil level and were preserved in 75% ethanol at 4 °C. These rosettes were later measured for total leaf area, dry rosette biomass (excluding root), root diameter at 1 cm below the soil surface, stomatal density using a representative mid-aged leaf, specific leaf area (SLA) using one representative mid-aged leaf, leaf δ^{13}C, leaf δ^{15}N, and total leaf carbon and nitrogen composition. Leaf tissue samples for δ^{13}C, δ^{15}N, total carbon and nitrogen composition were prepared by drying the complete rosettes in coin envelopes for 12-24 hours at 65.6 °C, then grinding and homogenizing all leaves from the rosette to a fine powder using a mortar and pestle. Based on our estimated nitrogen content, we packaged a 1.5 – 2.0 mg aliquot of finely ground leaf tissue from each rosette into an 8 x 5 mm tin capsule (EA Consumables, Pennsauken, NJ). Samples

were sent to the University of California Davis Stable Isotope Facility (SIF) for $\delta^{13}C$, $\delta^{15}N$, total carbon and nitrogen analyses using an elemental analyzer interfaced to a continuous flow isotope ratio mass spectrometer (IRMS). We measured whole-plant Arabidopsis gas exchange on a separate set of plants in an experiment that replicated the temperature, soil, and photoperiod conditions in the main experiment (Online Supplement).

Response to simulated root exudate strigolactones of neighbors

We used a synthetic strigolactone applied to the soil, GR24[rac] (ChemPep, Inc., Wellington, FL), to explore how SLs from neighboring *B. distachyon,* which exude SLs from their roots (Changenet, 2018), might affect Arabidopsis performance and traits. Four of our genotypes were selected, two that were highly sensitive to competing grass neighbors (Sf-2 and Ct-1), and two that were less sensitive to neighbors (Wil-2 and Ws-0) under well-watered conditions. Plants were grown alone in pots (no competition). Growing conditions were as described in the main experiment above. Plants received bottom water until the first true leaves were expanded, at which point the water was removed and treatment applications were started. Plants were then top watered evenly throughout the top of the potting mix around the rosette using a pipette every day with 3 ml of either 7.8 uM GR24 solution, 0.78 uM GR24, or tap water, with 5 individuals per genotype per treatment. All plants were harvested 52 DAP, prior to flowering initiation. Rosette biomass and quantum yield of photosystem II using a MultispeQ V2.0 (Kuhlgert *et al.*, 2016) were measured. Leaf $\delta^{13}C$ and total leaf C and N were measured from tissue analyzed at the UC Davis SIF.

Plant harvest and phenotyping at maturity

Fecundity of those Arabidopsis plants that survived to reproduction was estimated when all siliques reached maturity (dried and brown) and the rosette leaves had senesced, using total fruit length as an estimate of fecundity (Gnan *et al.*, 2014). We measured total silique number and length of six siliques corresponding to 10, 20, 40, 60, 80, and 90th percentiles of silique positions along the inflorescence in order to represent silique length along the inflorescence (*e.g.* 10th percentile silique was higher up the inflorescence than 10% of siliques). For each plant, we also measured inflorescence height and the number of inflorescence branches. After siliques

were measured, we dried the mature inflorescence and rosette separately (in addition to rosettes that did not flower) at 90 °C for 24 hours to get aboveground dry biomass.

Data analysis

225 Broad-sense heritability within treatments was calculated as the proportion of variation in phenotypes explained by genotype identity within each treatment, using a general linear model. We used linear mixed models to study sources of phenotype variation, with drought and competition treatments as fixed effects and genotypes, genotype-by-treatment interactions, and trays as random effects. With these mixed models we tested the hypothesis that fixed effects

230 were zero and calculated variance components for random effects using the R package 'VCA' (Schuetzenmeister & Dufey, 2018). We calculated 95% confidence intervals on variance components (constrained to be non-negative) and report as significant components where the interval excludes zero. We calculated genotypes' breeding values for each trait in each treatment using similarly constructed linear mixed models, using the R package 'lme4' (Bates *et al.*, 2013).

235 To assess whether there were genetic correlations in traits often thought to represent coordinated strategies, we calculated Pearson's correlations between breeding values for flowering time, SLA, leaf C:N, $\delta^{13}C$, and *A* (assimilation rate).

We tested for evidence of selection within each treatment by fitting linear regression models of relative fitness (absolute fitness / mean fitness among genotypes within a treatment) as

240 a function of breeding values for individual unit-standardized phenotypes. Flowering time was a trait under strong selection in all treatments and thus we also tested trait correlations with fitness in multiple regression while also including a flowering time covariate, which allowed us to get better estimates of direct selection acting on a trait while accounting for flowering time selection (Lande & Arnold, 1983).

245 We tested for changing selection across competition treatments within the same moisture regime using t-tests. We focus on a soft selection scenario where individual fitness is density dependent, which may be particularly appropriate for comparisons across competitive treatments. Under this scenario, the appropriate approach is to relativize fitness within each treatment (*i.e.* combination of moisture and competitive conditions) (De Lisle & Svensson, 2017). For

250 comparison, we included tests of changing selection when fitness was relativized across

competitive treatments in the same moisture regimes, corresponding to a hard selection scenario (De Lisle & Svensson, 2017).

We also tested for selection on plasticity in response to competition, while accounting for differences among genotypes in their average traits. We fit linear models with breeding values and tested how trait in response to competition within a given moisture treatment was associated with absolute fitness in each individual competition treatment. We employed a multiple regression that included trait means averaged across competition treatments (Stinchcombe *et al.*, 2004). Greater detail on commands used in statistical software can be found in the Online Supplement.

RESULTS

Treatment effects on performance and physiology

We studied how drought and competition influenced individual Arabidopsis performance and physiology (Question 1 in Introduction). Drought and competition generally reduced plant performance, causing plants to have significantly smaller rosettes at mid-experiment harvest (rosette weight$_{vegetative}$ and total leaf area$_{vegetative}$), lower inflorescence weight, shorter inflorescences, fewer total mature fruits (siliques), shorter fruits, and less aboveground biomass (AGB$_{mature}$ *i.e.* end of experiment harvest; linear mixed-effects models LMMs, all $p < 0.05$, Table 1, Figure 1). A notable exception was that plants had larger rosette weight$_{mature}$ (at final harvest) under drought treatments compared to well-watered (LMM, drought fixed effect estimate = 55.3 mg, $p < 10^{-6}$). The number of inflorescence branches was strongly reduced under competition (LMM fixed effect, $p < 10^{-8}$), without any significant drought or interaction effects.

There was a significant interaction between competition and drought for AGB$_{mature}$, inflorescence weight, inflorescence height, fruit number, average fruit length, such that under the combination of drought and competition these traits were greater than expected from the additive contribution of the two treatments (LMM fixed effects all $p < 0.05$, Table 1). However, rosette weight$_{mature}$ and average fruit length were lower than expected under combined drought and competition (Figure 1, Table 1).

Survival to reproduction was high across treatments, with 95% of plants surviving until flowering (only one plant survived but did not flower by the end of the experiment). There was

no pre-flowering mortality in either no-competition treatment (100% survival). By contrast, with competition, survival was 98% under well-watered conditions and 84% under drought, indicating that competition combined with drought was the harshest treatment. Mean genotype absolute fitness (mean estimated total mature fruit length per initial seedling following thinning) was highest under well-watered conditions without crowding (1900 mm, standard deviation among all individuals, SD = 599), followed by drought without crowding (656 mm, SD = 264), well-watered and crowding (387 mm, SD = 181), and drought and crowding combined (131 mm, SD = 95). We did not measure fecundity on *B. distachyon*, though its total aboveground biomass was greater in well-watered treatments compared to drought treatments at the mid-experiment harvest (mean pot total biomass, well-watered = 0.36 g, SD = 0.05; drought = 0.22 g, SD = 0.04).

In contrast to the whole plant performance measures described above, traits associated with life history or water balance physiology showed more diverse responses to water and competition (Figure 1). For example, mean flowering time showed a significant competition by drought interaction, such that flowering was significantly delayed only under combined competition and drought (LMM, competition×drought fixed effect p=0.0007, Table 1). Competition resulted in significantly decreased root diameter (LMM, fixed effect $p < 10^{-8}$) and reduced $\delta^{13}C$ (a proxy for water use efficiency, $p < 10^{-5}$) across moisture treatments (Table 1). Under drought, competition resulted in even greater reductions in root diameter (LMM, competition×drought fixed effect $p < 10^{-6}$) and greater SLA (p = 0.0002), while SLA was not affected by competition in well-watered conditions. Under well-watered conditions, competition resulted in greater leaf C:N (LMM, competition×drought fixed effect $p < 10^{-6}$) and reduced $\delta^{15}N$ ($p < 10^{-8}$), while these were not affected by competition under drought (Table 1, Figure 1).

Drought treatments (apart from competition effects) resulted in reduced root weight, root:shoot ratio, greater stomatal density, lower leaf C:N, greater $\delta^{13}C$, and lower δ15N (all traits from mid-experiment vegetative harvest, LMM, all drought fixed effects p < 0.004, Table 1, Figure 1).

We also addressed whether plastic responses to competitor plants were consistent with effects of SLs in exudate. Consistent with responses to *B. distachyon* neighbors, Arabidopsis showed reductions in aboveground vegetative biomass with increased SL levels (GR24[rac]) under well-watered conditions (Figure S5, ANOVA on log proportional biomass with ranked GR24 treatment covariate, $p < 10^{-15}$, genotype-GR24 interaction p = 0.7398). Howver, the SL treatment

affected other traits in ways inconsistent with responses to *Brachypodium*, causing increased $\delta^{13}C$ (ANOVA $\delta^{13}C$ response to ranked GR24 treatment covariate, p = 0.0004, genotype-GR24 interaction p = 0.1588) and decreased leaf C:N (ANOVA, $p < 10^{-6}$, genotype-GR24 interaction p = 0.2508, Figure S5). GR24 had genotype-specific effects on the quantum yield of photosystem II, with some genotypes having greater PSII yield with increasing GR24 and one having decreased PSII yield with increasing GR24 (ANOVA quantum yield of PSII response to ranked GR24 treatment covariate, p = 0.4047, genotype-GR24 interaction p = 0.0088).

Genetic and genotype-environment interaction effects on traits

Broad-sense heritability within treatments differed widely among traits, with flowering time having the highest heritability in every treatment (ranging from 0.88 in drought and competition to 0.96 to well-watered and competition, Table S4). Heritability averaged among traits was highest in the well-watered no competition treatment, followed by well-watered competition, drought no competition, and drought competition, respectively. Heritability was higher for leaf C:N compared to leaf proportion N in most treatments and thus we below focus results on C:N.

We generally found that flowering time and $\delta^{13}C$ were negatively correlated: later flowering genotypes had lower $\delta^{13}C$ (well-watered, competition r = -0.50; drought, no competition r = -0.49, drought, competition r = -0.50) with the exception well-watered, no competition treatments where the two were unrelated (r= -0.07). As expected, SLA and leaf C:N were always negatively correlated (Figure S6). These traits were not closely related to *A* (assimilation rate) in well-watered conditions but under drought and no competition we observed that genotypes with greater *A* had greater SLA (r = 0.43), lower C:N (r = -0.56) and lower $\delta^{13}C$ (r = -0.48).

We also tested for genetic variation in environmental responses to further address Question 1. Total fitness (Figure 2) and total silique number showed significant genotype-by-competition-by-water treatment interactions but not significant genotype interactions with competition or water alone (95% CIs of variance components excluded zero, Table S5), suggesting the potential for genetic tradeoffs in performance due to drought-competition interactions. Flowering time exhibited a strong genetic variance component and also a significant genotype-by-competition-by-treatment interaction but no genotype interaction with competition or water alone. Rosette weight$_{vegetative}$ and rosette weight$_{mature}$ both showed significant genotype-

by-competition interactions but other traits did not (Figure 1, Table S5). Root:shoot, root weight, and $\delta^{15}N$ (vegetative harvest) showed significant genotype variance components but no significant genotype-by-treatment interactions. Inflorescence height, inflorescence branch number, silique length, root diameter, SLA, stomatal density, C:N, $\delta^{13}C$, total rosette leaf area (vegetative harvest) did not show significant genotype-by-treatment variance components or significant genotype variance components (Table S5).

To assess the existence of fitness tradeoffs across environments (Question 2 in the Introduction), we calculated the correlation in relative fitness between competitive treatments. Under well-watered conditions, competition did not have a strong effect on the relative fitness of genotypes (absolute fitness correlation across environments, $r = 0.82$, $p < 10^{-5}$, Figure 2). However, under drought, competition had a strong impact on relative fitness, such that absolute fitness was weakly related across competitive contexts ($r = 0.38$, $p = 0.1170$, Figure 2). Comparing moisture treatments under no competition, relative fitness was weakly correlated across environments, suggesting possible tradeoffs ($r = 0.33$, $p = 0.1813$). Under competition, fitness was more correlated across moisture treatments, indicating genotypes selected under competition have traits making them robust to moisture inputs ($r = 0.52$, $p = 0.0263$).

Selection on phenotypes within and across environments

To identify traits affecting fitness variation (Question 3 in the Introduction), we tested for univariate and multivariate selection on traits within each environment. Earlier flowering time (FT) was favored in all four environments (linear models, well-watered, no competition $p < 10^{-6}$, well-watered, competition $p = 0.0020$, drought, no competition $p = 0.0162$, drought, competition $p = 0.0231$, only well-watered significant at FDR < 0.05, Table 2). Greater inflorescence height (linear model $p = 0.0001$, significant after FDR) and inflorescence branch number ($p = 0.0337$, not significant after FDR) was linked to higher fitness in well-watered competition treatments but not in other treatments (all $p > 0.05$). Inflorescence height in well-watered competition treatments was selected even after conducting multiple regression also including a flowering time effect (inflorescence height $p = 0.0214$, flowering time $p = 0.0469$). Smaller rosettes (mature harvest) were selected under no competition treatments of both moisture regimes (well-watered, no competition $p = 0.0095$, drought, no competition $p = 0.0030$, both significant FDR < 0.05), though these patterns were not statistically significant after including flowering time in

375 multiple regression (Table 2). AGB_{mature} was positively selected under well-watered competition treatments (p = 0.0002, significant after FDR), negatively selected under drought no competition (p = 0.0223, not significant after FDR), but unrelated with fitness in well-watered no competition treatments (p = 0.5625) and drought competition (p = 0.4926). Even after including flowering time in multiple regression, AGB_{mature} was positively selected in well-watered treatments with

380 competition (AGB_{mature} p = 0.0020, FT p = 0.0159) but not so in drought treatments without competition (AGB_{mature} p = 0.4383, FT p = 0.2785). In the harshest environment, drought and competition, we found that $\delta^{13}C$ was positively selected (p = 0.0291, not significant after FDR) though this was not significant after including FT ($\delta^{13}C$ p = 0.1830, FT p = 0.1433). Stomatal density, SLA, root:shoot, root biomass, photosynthetic rate, C:N, $\delta^{15}N$, and root diameter did not

385 show significant selection in any environment (Table 2).

To determine the traits responsible for genotype-level tradeoffs across treatments (Question 3), we tested for changes in the coefficients of selection across competition treatments within a given moisture environment. When relativizing fitness within each treatment (De Lisle & Svensson, 2017), we found significant changes in selection depending on the competitive

390 environment for inflorescence height and AGB_{mature} under well-watered conditions, with plants with taller inflorescences (two-tailed t-test p = 0.007, significant with FDR < 0.05), greater inflorescence mass (p < 0.0001, significant after FDR) and greater AGB_{mature} (p = 0.0003, significant after FDR) being more fit under competitive environments but with no detectable effect on fitness under no competition (Table 3, Figure 2). These changes in selection on these

395 traits are surprising given the strong correlation (*i.e.* weak tradeoffs) between relative fitness of genotypes in well-watered conditions when comparing plants growing alone or with competitors (Figure 2). By contrast, we had observed changing ranks of relative fitness across competition treatments under drought (Figure 2). We did not find significant changes in selection on any phenotypes under drought across competition treatments (aside from the trivial case of silique

400 number), though stomatal density showed the most substantial change (t-test p = 0.0630, Table 3). To further investigate the fitness tradeoffs across competition treatments under drought, we tested change in relative fitness between treatments and found it to be most closely associated with AGB_{mature} under drought and competition (relative fitness of drought under no competition - relative fitness under competition vs. AGB_{mature} under drought and competition, general linear

405 model, p = 0.0118) while AGB_{mature} under drought no competition did not explain the fitness

changes (p = 0.8333). When relativizing fitness across competition treatments in the same moisture environment (corresponding to a scenario of hard selection) (De Lisle & Svensson, 2017), flowering time in well-watered conditions was the only trait under significant changing selection across competition treatments, with relaxed selection under competition due to lower absolute fitness in this treatment (Table S6).

When testing for evidence of selection on traits averaged across competitive environments and independently on plasticity (Question 4 in Introduction), we found significant selection under well-watered conditions for a plastic increase in inflorescence height under competition compared to no competition (multiple linear regression, p = 0.0034) while average inflorescence height across treatments was not under significant selection, although this result was not significant after FDR control. For other traits where the average value was under selection across competitive environments, we found no independent selection on plasticity (Table S7).

DISCUSSION

Organismal responses to abiotic conditions can be influenced by biotic interactions, such as interactions that affect general organismal health (*e.g.* parasitism, Price, 1991), change the balance of optimal allocation to abiotic stress tolerance (*e.g.* facilitation in arid systems, Bertness & Callaway, 1994), or directly change the availability of abiotic resources (*e.g.* competition for water, Cohen, 1970). Given rapid global climate change, we are particularly interested in limiting resources that are influenced by climate, like moisture. Such resources are likely to exhibit interactions between climate and competition in their effects on organisms. While a broad literature exists studying plant performance responses to drought combined with competition (e.g. Callaway & Walker, 1997; Young *et al.*, 2017), less is known about the underlying physiology and genetic variation. Here, we studied the model plant Arabidopsis to characterize plastic responses to drought and competition as well as to determine how competition can change selection for traits related to water balance. We found strong plastic responses to neighboring plants that differed between drought and well-watered conditions. Furthermore, we found that under drought there are major rank changes in relative fitness depending on the presence of competitors.

Plastic responses to competition and moisture

We found plastic changes in most phenotypes, with strong interactions between competition and

440   moisture. Competition and drought reduced performance in both well-watered and drought

treatments, although plants in drought and competition had higher performance (biomass and

total silique length) than expected from the additive contribution of each stressor. This

interaction may indicate multiplicative growth responses to increased resources available in the

absence of competition or with sufficient water. Additionally, plants under drought without

445   competition generally had larger rosettes at maturity than in other conditions (Figure 1), perhaps

signifying an adaptive conservative response that reduces reliance on inflorescence (cauline)

leaves for photosynthesis.

By contrast, life history and physiology showed diverse plastic responses to our

treatments. Generally, drought responses showed a strategy partly indicative of adaptive escape,

450   *i.e.* a resource acquisitive strategy: lower leaf C:N and greater stomatal density, perhaps allowing

plants to rapidly increase photosynthesis and take advantage of pulses of water. At the same

time, higher $\delta^{13}C$ under drought indicated higher water use efficiency and stomatal closure. Des

Marais *et al.*, (2012) found that Arabidopsis tends to increase SLA and decrease C:N under

drought, which the authors interpreted as drought escape. However, several non-vernalization

455   requiring genotypes in Des Marais *et al.* (2012) showed opposite patterns: greater C:N and lower

SLA under drought. Our study genotypes were non-vernalization requiring, and we found that

drought increased SLA only under competition. There was no drought effect on SLA when

plants were alone. These drought-by-competition interactions suggest the escape strategy might

be specific to certain life history strategies or combined competitive and drought conditions.

460   Responses to competition differed between drought and well-watered conditions,

demonstrating the interdependence of abiotic and biotic effects. Under well-watered conditions,

response to competition was characterized by higher leaf C:N, suggesting a decrease in a

photosynthetic rate per leaf mass, although this was coupled with a consistent but slight decrease

in $\delta^{13}C$, the latter of which suggests an acquisitive growth strategy via greater stomatal or

465   mesophyll conductance to escape competition and is consistent with findings by Campitelli *et al.*

(2016). By contrast, competition under drought increased SLA, with some genotypes also

reducing $\delta^{13}C$, perhaps indicating an acquisitive escape strategy (allowing greater whole-plant

carbon assimilation) (Des Marais *et al.*, 2012). We found that under drought, genetic variation in SLA was positively associated with whole plant carbon assimilation (Figure S6). In one of the few relevant studies, Ehleringer (1993) found that the desert shrub *Encelia farinose* (Asteraceae) showed no δ^{13}C response to competitor removal.

δ^{15}N was elevated in well-watered no-competition treatments, which we interpreted as a shift in Arabidopsis use of N pools among treatments. Synthetic fertilizers (*e.g.* Miracle Grow used here) cause lower leaf δ^{15}N relative to organic sources like peat humus (used in our media), as fertilizer has more ammonium, which requires a single assimilation event (in roots) versus multiple assimilation events (in roots and shoots) for soil nitrate sources (Evans *et al.*, 1996; Bateman *et al.*, 2005; Trandel *et al.*, 2018). We infer that well-watered Arabidopsis plants without competition used more N from the peat humus versus fertilizer. Under well-watered conditions with competitors, the fertilizer N source may have been favored by the Arabidopsis plants due to smaller root systems for exploitation of soil N, causing low leaf δ^{15}N. Under drought, N uptake may have been restricted to brief periods following fertigation, favoring the fertilizer source and causing low leaf δ^{15}N.

Even under well-watered conditions with fertilizer and direct light, Arabidopsis responded to competition with reduced performance. Such a strong response suggests Arabidopsis plants were using a chemical cue, such as root exudates (Pierik *et al.*, 2013), to sense the neighboring *Brachypodium* (Badri *et al.*, 2012; Schmid *et al.*, 2013). Previous studies have shown that SLs in growth media can suppress inflorescence branching by Arabidopsis (Scaffidi *et al.*, 2014) and cause stomatal closure (Lv *et al.*, 2018). Our treatments with a synthetic SL reduced vegetative rosette biomass, suggesting SL signaling could partly explain Arabidopsis response to neighbors. The yield of PSII was greater for two genotypes under SL treatment, consistent with findings in tomato that the synthetic SL GR24 up-regulated light-harvesting genes (Mayzlish-Gati *et al.*, 2010). However, two other genotypes showed *lower* PSII yield in response to SL; these genotypes had lower biomass response to SL, for reasons that are unclear. We found that experimental SL addition caused an increase in δ^{13}C (likely signaling stomatal closure, Lv *et al.*, 2018) and a decrease in C:N, patterns that were opposite the Arabidopsis response to *Brachypodium* in our well-watered treatment. This result indicates that other non-SL cues are required to explain the responses to *Brachypodium* that we observed under putative non-resource limited conditions (our well-watered treatment).

Genetic correlations among traits

We found that genetic variation in flowering time was negatively associated with leaf $\delta^{13}C$, suggesting that early flowering plants had higher intrinsic leaf-level WUE due to lower stomatal conductance or higher assimilation rate A (Ferguson *et al.*, 2019). This positive correlation between flowering time and $\delta^{13}C$ might be a result of maladapted late-flowering genotypes having lower A for a given stomatal conductance. Our findings contrast with some previous studies that found the early flowering plants had low leaf $\delta^{13}C$ (McKay *et al.*, 2003; Lovell *et al.*, 2013; Kenney *et al.*, 2014). Our panel of natural genotypes lacked genotypes requiring vernalization , thus these genotypes might partly be responsible for the discrepancies with these publications. Additionally, Ferguson *et al.* (2019) did not find a flowering time relationship with leaf $\delta^{13}C$ in comparisons of NILs at major flowering time QTL.

Selection under competition and changes in selection depending on competition

Researchers have more commonly investigated how competition-drought interactions affect different species in a community (Marks & Strain, 1989; Miranda-Apodaca *et al.*, 2015; Napier *et al.*, 2016), or have studied changing selection driven by the interaction between drought and herbivory (Sthultz *et al.*, 2009; Huttunen *et al.*, 2010; Grinnan *et al.*, 2013) or competition and herbivory (Tiffin, 2002). However, there are few studies of changing selection due to drought-competition interactions. In one of the few such studies, Ehleringer (1993) found that in the desert shrub *Encelia farinose*, $\delta^{13}C$ was not under selection in the presence of competitors. However, upon competitor removal, lower $\delta^{13}C$ was associated with higher growth (but not survival) possibly due to increased soil moisture.

Here, we observed significant genotype-by-competition-by-moisture treatment interactions for performance related traits (fitness components and aboveground biomass) and flowering time. The genotype-by-competition-by-moisture effects on fitness indicate that the effects of competition on relative fitness are conditional on moisture environment. This competition-by-moisture effect on selection could be seen in the substantial changes in relative fitness depending on competition under drought, but the fairly consistent relative fitness despite the presence of neighbors in well-watered conditions (Figure 2). Rank changes in relative fitness across treatments differing in competition were observed among natural Arabidopsis genotypes at a

530   moist field site (Baron *et al.*, 2015) and among genotypes segregating for a single water-use-efficiency locus under greenhouse drought (Campitelli *et al.*, 2016). Here we show selection among natural genotypes under drought. In nature, the heterogeneity of competition may select for alternate Arabidopsis traits in water limited environments, suggesting a potential mechanism promoting their high genetic diversity in drought responses (Dittberner *et al.*, 2018; Exposito-

535   Alonso *et al.*, 2018).

We identified several traits associated with fitness in specific competitive contexts. Inflorescence height and weight as well as AGB_{mature} showed the strongest evidence for changing selection depending on competition , both when well-watered. Under well-watered competition, plants with greater AGB_{mature} and larger inflorescences had greater fitness, but not in the absence

540   of competition. Selection for AGB_{mature} may indicate that the investment in leaves for resource capture (comprising much of non-reproductive of AGB, and including both rosette and cauline leaves) is adaptive when surrounded by competitors, while allocation to reproduction may be selected when competitors are absent. The lack of selection for greater AGB_{mature} and taller inflorescences in drought may indicate that aboveground investments are less useful when

545   competitors are smaller. We also found positive selection for the plastic increase in inflorescence height under well-watered competition treatments. Thus, inflorescence height may only be a good measure of fecundity under the treatment with the most competitor biomass (well-watered), where competitors lead to suppressed inflorescence branching (Schmitt *et al.*, 2003; von Wettberg *et al.*, 2008).

550   Under drought and competition, plants with greater $\delta^{13}C$ had higher fitness (though this was not significant after FDR control), indicating greater leaf-level water use efficiency was adaptive, but not in other treatments. We expected lower WUE would be favored under drought with competitors to rapidly complete development before competitors depleted resources too low (Campitelli *et al.*, 2016). Instead, higher WUE may have been favored because conditions were

555   too dry to permit success of a rapidly developing drought escape strategy, thus higher WUE may be necessary in these environments. Alternatively, despite the dramatically lower performance of plants in response to competition under drought, the intermittent watering may have been enough to favor conservative drought avoidance strategies (high $\delta^{13}C$). Our findings contrast with those of Kenney *et al.* (2014) who found terminal drought and well-watered treatments selected for

560   lower $\delta^{13}C$ and plants with greater AGB_{mature}, suggesting escape strategies were favored. Here,

early flowering was favored in competitive treatments as it was in all treatments, similar to the consistently observed selection for earlier flowering in many experiments (Austen *et al.*, 2017) including those on Arabidopsis under terminal drought (Kenney *et al.*, 2014).

565   *Conclusions*

The effects of abiotic stress on phenotypic plasticity and natural selection depend on biotic context. Arabidopsis represents an important model system for plants, including for its responses to environment. Our results highlight how natural variation in environmental responses should be interpreted in light of biotic interactions.

Acknowledgements

Jonathan Kizer, Sarah Lucas, and Crosley Kudla-Williams assisted in harvesting and phenotyping. David Des Marais provided initial seeds of *Brachypodium distachyon.*

Author contributions

CML carried out experiments, analyzed data, and wrote the manuscript. JRL planned the experiments, analyzed data, and wrote the manuscript.

References cited

Ågren J, Schemske DW. **2012**. Reciprocal transplants demonstrate strong adaptive differentiation of the model organism Arabidopsis thaliana in its native range. *New Phytologist* **194**: 1112–1122.

Alados CL, ElAich A, Papanastasis VP, Ozbek H, Navarro T, Freitas H, Vrahnakis M, Larrosi D, Cabezudo B. **2004**. Change in plant spatial patterns and diversity along the successional gradient of Mediterranean grazing ecosystems. *Ecological Modelling* **180**: 523–535.

Al-Babili S, Bouwmeester HJ. **2015**. Strigolactones, a novel carotenoid-derived plant hormone. *Annual review of plant biology* **66**: 161–186.

Aphalo PJ, Ballaré CL, Scopel AL. **1999**. Plant-plant signalling, the shade-avoidance response and competition. *Journal of Experimental Botany* **50**: 1629–1634.

Austen EJ, Rowe L, Stinchcombe JR, Forrest JRK. **2017**. Explaining the apparent paradox of persistent selection for early flowering. *New Phytologist* **215**: 929–934.

Bac-Molenaar JA, Granier C, Keurentjes JJB, Vreugdenhil D. **2016**. Genome-wide association mapping of time-dependent growth responses to moderate drought stress in Arabidopsis. *Plant, Cell & Environment* **39**: 88–102.

Badri DV, De-la-Peña C, Lei Z, Manter DK, Chaparro JM, Guimarães RL, Sumner LW, Vivanco JM. **2012**. Root Secreted Metabolites and Proteins Are Involved in the Early Events of Plant-Plant Recognition Prior to Competition. *PLOS ONE* **7**: e46640.

Baron E, Richirt J, Villoutreix R, Amsellem L, Roux F. **2015**. The genetics of intra- and interspecific competitive response and effect in a local population of an annual plant species. *Functional Ecology* **29**: 1361–1370.

Bateman AS, Kelly SD, Jickells TD. **2005**. Nitrogen Isotope Relationships between Crops and Fertilizer: Implications for Using Nitrogen Isotope Analysis as an Indicator of Agricultural Regime. *Journal of Agricultural and Food Chemistry* **53**: 5760–5765.

Bates D, Maechler M, Bolker B. **2013**. *lme4: Linear mixed-effects models using S4 classes*.

Bay RA, Rose N, Barrett R, Bernatchez L, Ghalambor CK, Lasky JR, Brem RB, Palumbi SR, Ralph P. **2017**. Predicting Responses to Contemporary Environmental Change Using Evolutionary Response Architectures. *The American Naturalist* **189**: 463–473.

Bertness MD, Callaway R. **1994**. Positive interactions in communities. *Trends in Ecology & Evolution* **9**: 191–193.

Bivand R, Lewin-Koh N. **2019**. *maptools: Tools for Handling Spatial Objects*.

Callaway RM, Walker LR. **1997**. Competition and facilitation: a synthetic approach to interactions in plant communities. *Ecology* **78**: 1958–1965.

Campitelli BE, Des Marais DL, Juenger TE. **2016**. Ecological interactions and the fitness effect of water-use efficiency: Competition and drought alter the impact of natural MPK12 alleles in Arabidopsis. *Ecology Letters* **19**: 424–434.

Changenet V. **2018**. Towards new roles for cytochrome P450s and strigolactones in Fusarium Head Blight of Brachypodium distachyon.

Clark JS, Bell DM, Kwit MC, Zhu K. **2014**. Competition-interaction landscapes for the joint response of forests to climate change. *Global Change Biology* **20**: 1979–1991.

Cohen D. **1970**. The expected efficiency of water utilization in plants under different competition and selection regimes. *Israel Journal of Botany* **19**: 50–4.

van Dam NM, Bouwmeester HJ. **2016**. Metabolomics in the Rhizosphere: Tapping into Belowground Chemical Communication. *Special Issue: Unravelling the Secrets of the*

Rhizosphere **21**: 256–265.

De Lisle SP, Svensson EI. **2017**. On the standardization of fitness and traits in comparative studies of phenotypic selection. *Evolution* **71**: 2313–2326.

DeLeo VL, Menge DNL, Hanks EM, Juenger TE, Lasky JR. **2019**. Effects of two centuries of global environmental variation on phenology and physiology of Arabidopsis thaliana. *Global Change Biology* **0**.

Des Marais DL, McKay JK, Richards JH, Sen S, Wayne T, Juenger TE. **2012**. Physiological Genomics of Response to Soil Drying in Diverse Arabidopsis Accessions. *The Plant Cell* **24**: 893–914.

Dittberner H, Korte A, Mettler-Altmann T, Weber APM, Monroe G, Meaux J de. **2018**. Natural variation in stomata size contributes to the local adaptation of water-use efficiency in Arabidopsis thaliana. *Molecular Ecology* **27**: 4052–4065.

Ehleringer JR. **1993**. Variation in leaf carbon isotope discrimination in Encelia farinosa: implications for growth, competition, and drought survival. *Oecologia* **95**: 340–346.

El-Soda M, KRUIJER W, MALOSETTI M, KOORNNEEF M, AARTS MGM. **2015**. Quantitative trait loci and candidate genes underlying genotype by environment interaction in the response of Arabidopsis thaliana to drought. *Plant, Cell & Environment* **38**: 585–599.

Evans RD, Bloom AJ, Sukrapanna SS, Ehleringer JR. **1996**. Nitrogen isotope composition of tomato (Lycopersicon esculentum Mill. cv. T-5) grown under ammonium or nitrate nutrition. *Plant, Cell & Environment* **19**: 1317–1323.

Exposito-Alonso M, Team 500 Genomes Field Experiment, Burbano HA, Bossdorf O, Nielsen R, Weigel D. **2019**. A map of climate change-driven natural selection in Arabidopsis thaliana. *bioRxiv*: 321133.

Exposito-Alonso M, Vasseur F, Ding W, Wang G, Burbano HA, Weigel D. **2018**. Genomic basis and evolutionary potential for extreme drought adaptation in Arabidopsis thaliana. *Nature Ecology & Evolution* **2**: 352.

Farrior CE, Dybzinski R, Levin SA, Pacala SW, Associate Editor: Frederick R. Adler, Editor: Mark A. McPeek. **2013a**. Competition for Water and Light in Closed-Canopy Forests: A Tractable Model of Carbon Allocation with Implications for Carbon Sinks. *The American Naturalist* **181**: 314–330.

Farrior CE, Tilman D, Dybzinski R, Reich PB, Levin SA, Pacala SW. **2013b**. Resource limitation in a competitive context determines complex plant responses to experimental resource additions. *Ecology* **94**: 2505–2517.

Ferguson JN, Meyer RC, Edwards KD, Humphry M, Brendel O, Bechtold U. **2019**. Accelerated flowering time reduces lifetime water use without penalizing reproductive performance in Arabidopsis. *Plant, Cell & Environment* **42**: 1847–1867.

Fournier-Level A, Korte A, Cooper MD, Nordborg M, Schmitt J, Wilczek AM. **2011**. A Map of Local Adaptation in Arabidopsis thaliana. *Science* **334**: 86–89.

Gienapp P, Fior S, Guillaume F, Lasky JR, Sork VL, Csilléry K. **2017**. Genomic quantitative genetics to study evolution in the wild. *Trends in Ecology & Evolution*.

Gnan S, Priest A, Kover PX. **2014**. The genetic basis of natural variation in seed size and seed number and their trade-off using Arabidopsis thaliana MAGIC lines. *Genetics* **198**: 1751–1758.

Hamrick JL, Holden LR. **1979**. Influence of Microhabitat Heterogeneity on Gene Frequency Distribution and Gametic Phase Disequilibrium in Avena barbata. *Evolution* **33**: 521–533.

Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A. **2005**. Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology* **25**:

1965–1978.

Hoffmann MH. **2002**. Biogeography of Arabidopsis thaliana (L.) Heynh. (Brassicaceae). *Journal of Biogeography* **29**: 125–134.

Hosgood SMW, Parsons PA. **1968**. Polymorphism in natural populations ofDrosophila for the ability to withstand temperature shocks. *Experientia* **24**: 727–728.

Juenger TE, SEN S, BRAY E, STAHL E, WAYNE T, MCKAY J, RICHARDS JH. **2010**. Exploring genetic and expression differences between physiologically extreme ecotypes: comparative genomic hybridization and gene expression studies of Kas-1 and Tsu-1 accessions of Arabidopsis thaliana. *Plant, Cell & Environment* **33**: 1268–1284.

Jump AS, Marchant R, Peñuelas J. **2009**. Environmental change and the option value of genetic diversity. *Trends in Plant Science* **14**: 51–58.

Kenney AM, McKay JK, Richards JH, Juenger TE. **2014**. Direct and indirect selection on flowering time, water-use efficiency (WUE, δ13C), and WUE plasticity to drought in Arabidopsis thaliana. *Ecology and Evolution*: n/a-n/a.

Kover PX, Valdar W, Trakalo J, Scarcelli N, Ehrenreich IM, Purugganan MD. **2009**. A multiparent advanced generation inter-cross to fine-map quantitative traits in Arabidopsis thaliana. *PLoS Genet.* **5**.

Kuhlgert S, Austic Greg, Zegarac Robert, Osei-Bonsu Isaac, Hoh Donghee, Chilvers Martin I., Roth Mitchell G., Bi Kevin, TerAvest Dan, Weebadde Prabode, *et al.* 2016. MultispeQ Beta: a tool for large-scale plant phenotyping connected to the open PhotosynQ network. *Royal Society Open Science* **3**: 160592.

Lande R, Arnold SJ. **1983**. The Measurement of Selection on Correlated Characters. *Evolution* **37**: 1210–1226.

Lasky JR, Des Marais DL, Lowry DB, Povolotskaya I, McKay JK, Richards JH, Keitt TH, Juenger TE. **2014**. Natural variation in abiotic stress responsive gene expression and local adaptation to climate in Arabidopsis thaliana. *Molecular Biology and Evolution* **31**: 2283–2296.

Lasky JR, Des Marais DL, McKay JK, Richards JH, Juenger TE, Keitt TH. **2012**. Characterizing genomic variation of Arabidopsis thaliana: the roles of geography and climate. *Molecular Ecology* **21**: 5512–5529.

Lasky JR, Forester BR, Reimherr M. **2018**. Coherent synthesis of genomic associations with phenotypes and home environments. *Molecular Ecology Resources* **18**: 91–106.

Levene H. **1953**. Genetic Equilibrium When More Than One Ecological Niche is Available. *The American Naturalist* **87**: 331–333.

Lovell JT, Juenger TE, Michaels SD, Lasky JR, Platt A, Richards JH, Yu X, Easlon HM, Sen S, McKay JK. **2013**. Pleiotropy of FRIGIDA enhances the potential for multivariate adaptation. *Proceedings of the Royal Society B: Biological Sciences* **280**: 20131043.

Ludlow MM. **1989**. Strategies of response to water stress. In: Structural and Functional Responses to Environmental Stresses. The Hague: SPB Academic, 269–281.

Lv S, Zhang Y, Li C, Liu Z, Yang N, Pan L, Wu J, Wang J, Yang J, Lv Y, *et al.* 2018. Strigolactone-triggered stomatal closure requires hydrogen peroxide synthesis and nitric oxide production in an abscisic acid-independent manner. *New Phytologist* **217**: 290–304.

Lynch M, Lande R. **1993**. Evolution and extinction in response to environmental change. In: Biotic Interactions and Global Change. Sunderland, MA: Sinauer Associates, Inc, 234–250.

Mayzlish-Gati E, LekKala SP, Resnick N, Wininger S, Bhattacharya C, Lemcoff JH, Kapulnik Y, Koltai H. **2010**. Strigolactones are positive regulators of light-harvesting genes in tomato. *Journal of Experimental Botany* **61**: 3129–3136.

720   **McKay JK, Richards JH, Mitchell-Olds T**. **2003**. Genetics of drought adaptation in Arabidopsis thaliana: I. Pleiotropy contributes to genetic correlations among ecological traits. *Molecular Ecology* **12**: 1137–1151.

    McLeod KA, Scascitelli M, Vellend M. **2012**. Detecting small-scale genotype–environment interactions in apomictic dandelion (Taraxacum officinale) populations. *Journal of Evolutionary*
725   *Biology* **25**: 1667–1675.

    Mitchell-Olds T, Willis JH, Goldstein DB. **2007**. Which evolutionary processes influence natural genetic variation for phenotypic traits? *Nat Rev Genet* **8**: 845–856.

    Monroe JG, McGovern C, Lasky JR, Grogan K, Beck J, McKay JK. **2016**. Adaptation to warmer climates by parallel functional evolution of CBF genes in Arabidopsis thaliana.
730   *Molecular Ecology* **25**: 3632–3644.

    Moran EV, Alexander JM. **2014**. Evolutionary responses to global change: lessons from invasive species. *Ecology Letters* **17**: 637–649.

    Msanne J, Lin J, Stone JM, Awada T. **2011**. Characterization of abiotic stress-responsive Arabidopsis thaliana RD29A and RD29B genes and evaluation of transgenes. *Planta* **234**: 97–
735   107.

    Pease CM, Lande R, Bull JJ. **1989**. A Model of Population Growth, Dispersal and Evolution in a Changing Environment. *Ecology* **70**: 1657–1664.

    Pierik R, Mommer L, Voesenek LA. **2013**. Molecular mechanisms of plant competition: neighbour detection and response strategies. *Functional Ecology* **27**: 841–853.
740   **Pisupati R, Reichardt I, Seren Ü, Korte P, Nizhynska V, Kerdaffrec E, Uzunova K, Rabanal FA, Filiault DL, Nordborg M**. **2017**. Verification of Arabidopsis stock collections using SNPmatch, a tool for genotyping high-plexed samples. *Scientific Data* **4**.

    Price PW. **1991**. The Plant Vigor Hypothesis and Herbivore Attack. *Oikos* **62**: 244–251.

    Proust H, Hoffmann B, Xie X, Yoneyama K, Schaefer DG, Yoneyama K, Nogué F, Rameau
745   **C**. **2011**. Strigolactones regulate protonema branching and act as a quorum sensing-like signal in the moss Physcomitrella patens. *Development* **138**: 1531–1539.

    Scaffidi A, Waters MT, Sun YK, Skelton BW, Dixon KW, Ghisalberti EL, Flematti GR, Smith SM. **2014**. Strigolactone Hormones and Their Stereoisomers Signal through Two Related Receptor Proteins to Induce Different Physiological Responses in Arabidopsis. *Plant Physiology*
750   **165**: 1221–1232.

    Schenk HJ. **2006**. Root competition: beyond resource depletion. *Journal of Ecology* **94**: 725–739.

    Schmid C, Bauer S, Müller B, Bartelheimer M. **2013**. Belowground neighbor perception in Arabidopsis thaliana studied by transcriptome analysis: roots of Hieracium pilosella cause biotic
755   stress. *Frontiers in Plant Science* **4**.

    Schmitt J, Stinchcombe JR, Heschel MS, Huber H. **2003**. The Adaptive Evolution of Plasticity: Phytochrome-Mediated Shade Avoidance Responses. *Integrative and Comparative Biology* **43**: 459–469.

    Schuetzenmeister A, Dufey F. **2018**. *VCA: Variance Component Analysis*.
760   **Souza ALSR de, Souza SAD, Oliveira MVVD, Ferraz TM, Figueiredo FAMMA, Silva NDD, Rangel PL, Panisset CRS, Olivares FL, Campostrini E, et al.** **2015**. Endophytic colonization of Arabidopsis thaliana by Gluconacetobacter diazotrophicus and its effect on plant growth promotion, plant physiology, and activation of plant defense. *Plant and Soil* **399**: 257–270.
765   **Stinchcombe JR, Dorn LA, Schmitt J**. **2004**. Flowering time plasticity in Arabidopsis thaliana:

a reanalysis of Westerman & Lawrence (1970). *Journal of Evolutionary Biology* **17**: 197–207.

Sultan SE, Bazzaz FA. **1993**. Phenotypic Plasticity in Polygonum Persicaria. Ii. Norms of Reaction to Soil Moisture and the Maintenance of Genetic Diversity. *Evolution* **47**: 1032–1049.

Trandel MA, Vigardt A, Walters SA, Lefticariu M, Kinsel M. **2018**. Nitrogen Isotope Composition, Nitrogen Amount, and Fruit Yield of Tomato Plants Affected by the Soil–Fertilizer Types. *ACS Omega* **3**: 6419–6426.

Urban MC, Zarnetske PL, Skelly DK. **2013**. Moving forward: dispersal and species interactions determine biotic responses to climate change. *Annals of the New York Academy of Sciences* **1297**: 44–60.

von Wettberg EJ, Remington DL, Schmitt J. **2008**. Partitioning Adaptive Differentiation across a Patchy Landscape: Shade Avoidance Traits in Impatiens Capensis. *Evolution* **62**: 654–667.

Young DJN, Stevens JT, Earles JM, Moore J, Ellis A, Jirka AL, Latimer AM. **2017**. Long-term climate and competition explain forest mortality patterns under extreme drought. *Ecology Letters* **20**: 78–86.

Table 1. Results of treatment fixed effects on phenotypes from a linear mixed-effects model where genetic and tray effects were modeled as random effects. Fixed effects are in units of the trait. Bolded p values are significant at false discovery rate < 0.05 for each contrast across traits.

	Competition - no competition (fixed effect)	p	Well watered - drought (fixed effect)	p	Competition & well watered - competition & drought (fixed effect)	p
Total silique length	-1513.893	**<0.0001**	1246.150	**<0.0001**	-974.521	**<0.0001**
AGB_m	-0.422	**<0.0001**	0.111	**<0.0001**	-0.075	**0.0006**
Rosette wt_m	-0.158	**<0.0001**	-0.055	**<0.0001**	0.061	**<0.0001**
Inflorescence wt	-0.263	**<0.0001**	0.167	**<0.0001**	-0.136	**<0.0001**
Infloresence height	-15.149	**<0.0001**	20.249	**<0.0001**	-6.960	**<0.0001**
Inflorescence branches	-5.844	**<0.0001**	0.751	0.0927	-0.654	0.2148
Flowering time	5.061	0.0572	-3.601	0.0841	-10.510	**0.0007**
Silique number	-124.911	**<0.0001**	92.501	**<0.0001**	-73.055	**<0.0001**
Average silique length	-0.643	**0.0141**	2.379	**<0.0001**	1.172	**0.0005**
Root wt_v			0.132	**0.0019**		
Root diam_v	-0.385	**<0.0001**	-0.154	**0.0178**	0.295	**<0.0001**
Root:shoot			0.018	**0.0023**		
Total leaf area_v	-37.098	**<0.0001**	6.798	**0.0054**	-2.442	0.0920
Stomatal density	19.723	0.0873	-71.873	**0.0031**	27.946	**0.0315**
Rosette wt_v	-5.543	**<0.0001**	0.470	0.2632	0.609	0.1278
log(SLA)	-0.091	0.0814	-0.004	0.9610	-0.397	**<0.0001**
Leaf C:N	4.598	**<0.0001**	4.082	**0.0006**	4.167	**<0.0001**
$\delta^{13}C$	-0.569	**<0.0001**	-1.964	**<0.0001**	-0.322	0.0533
$\delta^{15}N$	-2.761	**<0.0001**	1.916	**<0.0001**	-2.528	**<0.0001**
Brachypodium.AGB			0.135	**<0.0001**		

Table 2. Standardized selection coefficients on breeding values for traits under different treatments. Bolded p-values are shown to indicate significance with false discovery rate < 0.05. Note that root weight, roo:shoot, and *A* were measured only on plants grown in the absence of competition. All other traits had breeding values and estimated fitness compared within each of the four treatments. All selection coefficients are in units of relative fitness (individual measures divided by the mean in each treatment) per standard deviation of the trait breeding values within treatment.

	Well-watered, no comp.		Well-watered, comp.		Drought, no comp.		Drought, comp.	
	Coeff. est.	*p*	*Coeff. est.*	*p*	*Coeff. est.*	*p*	*Coeff. est.*	*p*
AGB_m	0.0461	0.5625	0.3600	**0.0002**	-0.2149	0.0230	0.0905	0.4926
Rosette wt_m	-0.1871	**0.0095**	-0.1074	0.3638	-0.2658	**0.0030**	-0.0537	0.6852
Inflorescence wt	0.2078	**0.0029**	0.4280	**0.0000**	0.1930	0.0448	0.2343	0.0622
Flowering time	-0.2728	**0.0000**	-0.3198	**0.0020**	-0.2252	0.0162	-0.2782	0.0231
Inflorescence height	0.0060	0.9401	0.3360	**0.0009**	-0.0363	0.7229	-0.0186	0.8883
Inflorescence branches	-0.0272	0.7335	0.2370	0.0337	-0.1639	0.0947	0.0182	0.8909
Silique number	0.3075	**0.0000**	0.4598	**0.0000**	0.3933	**0.0000**	0.4601	**0.0000**
Average silique length	0.0218	0.7850	0.2272	0.0431	0.1864	0.0538	0.3808	**0.0006**
Root wt_v	0.0633	0.4249	0.0693	0.5607	0.1494	0.1308	0.1602	0.2165
Root diam_v	-0.0410	0.6074	0.0887	0.4551	-0.0478	0.6397	0.0743	0.5739
Root:shoot	-0.0038	0.9620	-0.0610	0.6094	0.1307	0.1902	0.1719	0.1830
Stomatal density	-0.1140	0.1406	-0.1461	0.2111	0.0417	0.6831	-0.2333	0.0636
Rosette wt_v	0.1139	0.1410	0.1904	0.0968	-0.0749	0.4614	0.1073	0.4142
log(SLA)	-0.0991	0.2041	-0.0243	0.8390	0.1008	0.3177	-0.0174	0.8958
Leaf C:N	0.1244	0.1054	0.1269	0.2807	-0.0454	0.6568	0.0352	0.7907
A (whole plant assimilation)	0.0742	0.3478	0.1217	0.3014	-0.0693	0.4959	-0.0107	0.9361
$\delta^{13}C$	0.0406	0.6112	0.1863	0.1049	0.0406	0.6911	0.2689	0.0291
$\delta^{15}N$	0.0646	0.4148	0.0197	0.8692	0.0629	0.5372	0.1258	0.3363

Table 3. Tests for changing selection between competition treatments within the same moisture treatment. Slopes give the relationship between traits and relative fitness, where relative fitness is standardized within the treatment combination of competition and moisture conditions (corresponding to a model of soft selection), and traits are standardized across competition treatments being compared. Bold indicates those where slopes differ significantly between competition treatments with FDR < 0.05. For results when fitness is relativized across environments see Table S6.

Trait	Well-watered	Slope, w/ comp	Slope, no comp.	t-test p	Drought	Slope, w/ comp	Slope, no comp.	t-test p
AGB_m		2.77	0.15	**0.0003**		0.82	-0.57	0.1886
Rosette wt_m		-0.84	-0.32	0.4907		-0.50	-0.43	0.9487
Inflorescence wt		2.40	0.47	**<0.0001**		1.58	0.47	0.1448
Inflorescence height		0.54	0.01	**0.0070**		-0.03	-0.05	0.9340
Inflorescence branches		0.69	-0.04	0.0213		0.06	-0.22	0.4723
Flowering time		-0.25	-0.52	0.0568		-0.25	-0.40	0.4410
Silique number		2.32	0.44	**<0.0001**		2.25	0.49	**<0.0001**
Average silique length		0.24	0.02	0.1244		0.43	0.31	0.5045
Root diam_v		0.19	-0.06	0.3369		0.20	-0.15	0.4690
Stomatal density		-0.14	-0.13	0.8890		-0.27	0.04	0.0630
Rosette wt_v		1.15	0.16	0.0917		1.26	-0.25	0.2883
log(SLA)		-0.03	-0.09	0.6743		-0.03	0.14	0.4839
A (whole plant assimilation)		0.12	0.07	0.7313		-0.01	-0.07	0.7234
Leaf C:N		0.18	0.15	0.8567		0.03	-0.05	0.6264
δ13C		0.22	0.05	0.3260		0.29	0.04	0.1094
δ15N		0.07	0.20	0.7646		0.11	0.07	0.7892

Figure Legends

Figure 1. Reaction norms for traits across 18 Arabidopsis genotypes (each line represents one genotype). Lines show breeding values estimated from linear mixed models. 'NC' = no
820 competition, 'C' = competition, 'WW' = well-watered, 'D' = drought. At right in gray are three traits that were only measured in a single competition/no-competition treatment. Three traits showing significant variance components for genotype ("G") or genotype-treatment (e.g. "G×D") interactions are indicated. All traits showed significant (FDR < 0.05) fixed effects for at least one treatment or treatment interactions (Table 1 for details).

Figure 2. Reaction norms of absolute fitness (*i.e.* total silique length, first column) and relative fitness (standardized by dividing by mean fitness within each treatment, second columns) across competition treatments for the 18 studied genotypes (each line represents a genotype). Note that
830 competition treatments are on the left and no competition on the right, so as to keep consistent x-axes for traits under selection (right columns). Lines show breeding values estimated from linear mixed models. The well-watered (top row) and drought (bottom row) treatments are shown (note their changing scales). Changing selection on traits among treatments (third and fourth columns). Standardized selection coefficients are shown, with * indicating nominal significance (p<0.05)
835 and ** indicating FDR < 0.05. Horizontal bars with ** indicate slopes that differ significantly (t-test) between competition treatments in the same moisture regime with FDR < 0.05.

Figure 1.

840 **Figure 2.**

 SUPPLEMENTAL TEXT, FIGURES, AND TABLES

Growth conditions

Specifically, we chose conditions for the site of collection of the Lip-0 ecotype near Chrzanów in southern Poland. We used a growing season model (Lasky *et al.*, 2012) with WorldClim data

850 (Hijmans *et al.*, 2005) then extracted a monthly time series of maximum temperatures and minimum temperatures. We used these temperatures to program the growth chamber, though in the early growing season we were limited to 10 ºC nightly temperature because the chamber was unable to generate temperatures < 10 ºC. Local photoperiod for the growing season was obtained using the 'maptools' package (Bivand & Lewin-Koh, 2019). Weekly conditions were obtained

855 via linear interpolation of the monthly temperature values (Table S2). Our growth chamber had constant horizontal airflow to maintain homogenous conditions throughout.

Plants were grown in SC-10 Cone-tainers™ (3.81 cm diameter, 164 ml) in a 98-cell rack (RL98) leaving adjoining cells in the tray empty to prevent shading interaction between neighboring pots (Stuewe and Sons, Tangent, Oregon, USA). Each rack was placed within

860 individual flow trays. Competition treatments were spaced at every other pot location to prevent among-pot crowding, and each tray was designated a drought or well-watered treatment (Fig. S1). Genotype pot locations were randomly assigned. Potting composition within each pot consisted of 50% potting mix (Premier pro-mix PGX), 25% medium commercial grade sand, 25% calcined clay (TURFACE® ALL SPORT PRO™, Turface Athletics™, PROFILE Products

865 LLC, Buffalo Grove, IL), with 2.5 cm of the potting mix as a topsoil to assist Arabidopsis plants in germination and initial growth. Glad Press'n Seal Wrap (The Glad Products Company; Oakland, CA) was placed over pots until germination to prevent seed dehydration prior to germination.

Germination for *B. distachyon* was near 100%. The three *B. distachyon* genotypes were

870 randomly sorted into competition pots. Preliminary experiments using varying levels of *B. distachyon* in competition with *Arabidopsis* showed that greater than six plants resulted in overcrowding and very poor growth of *B. distachyon,* thus six plants were used as a competition treatment in pots.

Pots in the drought treatment were not watered during 17-19 DAP prior to fertigation

875 (combined watering and fertilizing) treatments to allow for initial drying. To simulate persistent rainfall limitation, starting at 20 DAP, each individual pot in the drought treatment was fertigated

from above with 3 ml of full strength Miracle-Gro solution (Miracle-Gro® Water Soluble All Purpose Plant Food, The Scotts Company, LLC., Marysville, OH) every other day. Throughout the entire experiment, all pots in all treatments were fertigated at the same time of day with the

880   same nutrition strength and volume to limit nutrient competition and to focus on traits involved in competition for water. At 29 DAP, fertigation was shifted to 6 ml of ½ strength Miracle Gro solution, and at 50 DAP fertigation was increased to 12 ml of ½ strength Miracle Gro solution. Well-watered pots were kept consistently moist by bottom watering with tap water. We defined the end of the plant life cycle as the time when 50% of siliques were mature (brown and brittle),

885   because at this stage the ecotypes used in this experiment stopped producing new flowers and discontinued development of existing fruits if present. Bottom water in well-watered treatments was removed when at least one plant had 50% mature (brown and brittle) siliques, and the remaining flowering plants were top watered daily with 12 ml of ½ strength Miracle Gro, to maintain the potting mix water and nutrient content of the well-watered plants, until 50% of

890   siliques were mature. Watering in drought treatments was also discontinued for a pot when 50% of siliques were mature. Potting mix volumetric water content for each treatment was monitored throughout the entire experiment with EM-50 dataloggers and EC-5 probes (METER Group Inc., Pullman, WA) within designated non-experimental pots with the reference genotype Col-0 (Fig. S4).

Photosynthesis

We measured whole-plant gas exchange on a separate set of plants in an experiment that replicated the temperature, soil, and photoperiod conditions in the main experiment. However,

900   we did not include *B. distachyon* competitors and we covered topsoil in clay. We did not use competition treatments due to the challenges of measuring whole plant gas exchange on the focal Arabidopsis plant when surrounded by *B. distachyon* competitors. We used the clay cap as suggested by the gas exchange analyzer manufacturer (LI-COR Inc., Lincoln, NE, USA) to limit soil respiration from interfering with photosynthetic gas exchange measurements. Prior to

905   planting, the top soil was covered with a 0.5 cm layer of potting clay (Msanne *et al.*, 2011; Souza *et al.*, 2015) with a 0.5 cm hole in the center for plant growth. Even with the clay and initial soil sterilization, we observed low, non-zero soil respiration from well-watered pots, and as a result

below we do not focus analysis on changes in photosynthesis between treatments, but instead use photosynthesis measurements in selection analyses within moisture treatments.

910   Five replicates per genotype per water treatment were planted in a randomized block design. Photosynthesis (*A*) was measured between 8:00 AM and 12:30 PM using a LiCor 6400XT with a LI6400-17 Whole Arabidopsis Chamber (LI-COR Inc., Lincoln, NE, USA) modified to fit SC-10 Ray Leach Cone-tainers™ (Stuewe and Sons Inc., Tangent, OR) by widening the opening that holds the Cone-tainer to fit a size -223 O-ring (McMaster Carr part

915   number 1170N95, Elmhurst, IL). This modification created an air-tight seal around the circumference of the Cone-tainer approximately 4 mm from the top of the Cone-tainer, to prevent gas leaks from the surrounding environment and to allow for continuous air flow and even concentrations of gasses throughout the chamber during measurements.

920   *Statistical analysis*

We used linear mixed models to test fixed effects of treatments on Arabidopsis traits, as well as to test for significance of variance components for genotype and genotype-treatment interaction random effects. To do so, we used the VCA package for R. Example code is shown below, where 'mydata' is a data.frame with rows as individual plants and columns that include a

925   phenotype variable, 'CompTRT' and 'WaterTRT' variables that are factors giving the competition and moisture conditions respectively, and 'Tray' giving the identity of the tray the plant was in.

```
library(VCA)
930
model1 <- anovaMM(phenotype ~ CompTRT * WaterTRT * (genotype) +(Tray),
    Data = mydata)

inf <- VCAinference(model1, VarVC=TRUE)
935
VCAinference(model1)

lc.mat <- VCA::getL(model1, c("CompTRTC-CompTRTNC", "WaterTRTWW-
    WaterTRTD", 'CompTRTC:WaterTRTWW-CompTRTC:WaterTRTD'))
940
test.fixef(m1a, lc.mat)
```

We used linear models to test for selection on traits within each treatment. Example code is shown below, where 'mydata2' is a data.frame with rows as genotypes, 'phen' are breeding

945   values for a given phenotype in a given treatment, and 'RelFit' is fitness relativized to the mean within that treatment:

```
model2 <- lm(RelFit ~ phen, data = mydata2)
summary(model2)
```
950

We also used linear models with t-tests for changes in selection among competition treatments, within the same moisture treatment. Example code is shown below, where 'mydata3' is a data.frame with rows as genotypes, each genotype having two rows, one for breeding values without competition in a given moisture environment and one for breeding values with

955   competition in a given moisture environment. 'phen' is a column giving breeding values, 'compT' is a factor giving the competition treatment, and 'RelFit' is fitness relativized to the mean within that treatment:

```
model3 <- lm(RelFit ~ phen * compT, data = mydata3)
960   summary(model3)
```

We used linear models again to test for selection on competition plasticity while accounting for average trait values. Example code is shown below, where 'mydata3' is a data.frame with rows as genotypes. 'avg_phen' is a column giving breeding values averaged across competition

965   treatments for a given moisture environment, 'plast_phen' is a column giving the difference in breeding values between competition treatments for a given moisture environment, and 'RelFit' is fitness averaged across competition treatments for a given moisture environment and then relativized to the mean.

```
970   model4 <- lm(RelFit ~ avg_phen + plast_phen, data = mydata4)
      summary(model4)
```

Figure S1. Experimental treatments, including (A) drought treatment under competition and no competition, and (B) well-watered treatment under competition and no competition. Photographs were taken 53 days after planting, prior to the first harvest.

Figure S2. Experimental plant with stainless steel wire installed to prevent *B. distachyon* shading.

1000

Figure S3. Time series of temperature (°C) conditions using a HOBO data logger (HOBO®, Onset Computer Corporation). Temporary spikes in temperature were due to brief power outages and did not result in any clear detrimental effects on experimental plants.

1005

1010

Figure S4. Percent volumetric water content (%VWC) of potting mix within pots for each treatment was monitored using EM-50 dataloggers and EC-5 probes (METER Group Inc., Pullman, WA). Seeds were planted on May 31, %VWC logging was started on June 9, and 1015 drought treatments were started on June 20.

1020

Figure S5. Response of four genotypes to increasing concentrations of GR24 applied to soil.

1025   **Figure S6**. Bivariate relationships between traits often thought to be exhibit coordinated variation. Note that gas exchange (used to caculate *A*) was only measured in the absence of competitors, these estimates are used in the comparisons with traits under competition.

 Figure S7. Principal components of trait variation within each treatment.

Table S1. The 18 parent genotypes from the Multiparent Advanced Generation Inter-Cross (MAGIC) used in this study.

1035

ABRC Stock #	Accession
76734	Bur-0
76740	Can-0*
76778	Col-0
76786	Ct-1
76831	Edi-0
76969	Kn-0
77020	Ler-0
1380	Mt-0
77128	No-0
77156	Oy-0
6839	Po-0
77222	Rsch-4
77247	Sf-2
77389	Tsu-0
78856	Wil-2
78857	Ws-0
78858	Wu-0*
78880	Zu-0

* Stock was identified as incorrect by ABRC after seed was obtained (Pisupati *et al.*, 2017).

Table S2. Climate mean conditions for non-contaminant ecotypes using WorldClim (Hijmans et al. 2005) and CGIAR CSI Aridity (Zomer et al. 2008) data.

Line	Longitude	Latitude	Alt	Ann.Mean.Tmp (ºC)	Ann.Prc (mm)	Ann.Prc/PET
Bur-0	-6.2	54.1	172	8.4	1068	1.8098
Can-0*						
Col-0						
Ct-1	15	37.3	40	17.5	503	0.4209
Edi-0	-3.16028	55.9494	149	8.3	743	1.2129
Hi-0	5	52	-2	9.4	800	1.173
Kn-0	23.8924	54.8969	30	6.7	613	0.9068
Ler-0	10.8719	47.984	621	8	962	1.2847
Mt-0	22.46	32.34	282	17.5	132	0.1044
No-0	13.2995	51.0581	276	8.1	627	0.8947
Oy-0	6.193019	60.385543	5	7.2	2190	4.4512
Po-0	7.09	50.72	67	10	739	0.97
Rsch-4	34	56.3	210	4.1	640	1.0126
Sf-2	3.03333	41.7833	146	15.4	677	0.765
Tsu-0	136.31	34.43	160	14.9	2385	2.319
Wil-2	25.3167	54.6833	106	6.4	656	0.988
Ws-0	30	52.3	130	6.7	624	0.8398
Wu-0*						
Zu-0	8.55	47.3667	424	9.3	1087	1.4135

1040

Table S3. Temperature and photoperiod program for Conviron growth chamber, to simulate early spring germination for a spring annual.

Week	Night °C	Day °C	Photoperiod (Hours:Minutes)
1	10	15	13:04
2	10	15	13:30
3	10	15	13:54
4	10	16	14:20
5	10	17	14:42
6	10	18	15:06
7	10	19	15:26
8	10	20	15:44
9	10	21	16:00
10	10	21	16:12
11	11	22	16:18
12	12	23	16:22
13	12	23	16:20
14	13	24	16:16
15	13	24	16:06
16	13	24	15:52
17	13	24	15:34
18	13	24	15:16
19	13	24	14:54
20	13	24	14:32
21	12	23	14:08
22	11	22	13:42
23	10	21	13:18
24	10	20	12:52

1045

Table S4. Broad sense heritability of traits within each treatment. 'm' suffixes indicate mature end stage harvest, while 'v' indicates the pre-flowering vegetative stage harvest.

	WW NC	WW C	D NC	D C
Absolute fitness	0.7880	0.7337	0.5693	0.2797
AGB_m	0.6026	0.5489	0.6739	0.4255
Rosette wt_m	0.8592	0.5178	0.7970	0.4030
Inflorescence wt	0.6081	0.6419	0.3673	0.4387
Flowering time	0.8932	0.9568	0.8873	0.8798
Inflorescence height	0.8378	0.7176	0.4076	0.3531
Inflorescence branches	0.6920	0.4795	0.5927	0.3241
Silique number	0.7972	0.6835	0.5705	0.3024
Average silique length	0.5640	0.6601	0.4642	0.2772
Root wt_v	0.3012		0.1627	
Root diam_v	0.2952	0.2444	0.1254	0.4691
Root:shoot	0.3087		0.2019	
Total leaf area_v	0.3101	0.3695	0.2286	0.3768
Stomatal density	0.2026	0.2080	0.3614	0.3554
Rosette wt_v	0.6003	0.4206	0.2611	0.3754
log(SLA)	0.3924	0.2513	0.3884	0.2936
Leaf prop N	0.5285	0.3096	0.2566	0.3425
Leaf prop C	0.2960	0.2139	0.2566	0.2951
Leaf C:N	0.5785	0.4204	0.3269	0.3335
$\delta^{13}C$	0.3825	0.4691	0.3838	0.2766
$\delta^{15}N$	0.2820	0.2496	0.3640	0.3437
A (whole plant assimilation)	0.1325		0.1633	
Brachypodium AGB		0.2780		0.3606

1050

Table S5. Variance components estimates for linear mixed-effects model where genotype, genotype-treatment interactions, and tray effects were modeled as random effects and treatments were fixed effects. 'est' refers to the variance component estimate and 'LCL' and 'UCL' refer to the lower and upper 95% confidence limits for the estimate. Variance components estimated and fixed at zero report 'NA' for the 95% CI.

See attached files.

Table S6. Tests for changing selection between competition treatments within the same moisture treatment. Slopes give the relationship between traits and relative fitness, where relative fitness is standardized **across** the treatment combination of competition and moisture conditions (corresponding to a model of hard selection), and traits are standardized across competition treatments being compared. Bold indicates those where slopes differ significantly between competition treatments with FDR < 0.05. For results when fitness is relativized across environments see Table 3.

Trait	Well-watered	Slope, w/ comp	Slope, no comp.	t-test p	Drought	Slope, w/ comp	Slope, no comp.	t-test p
AGB_m		0.94	0.24	0.3812		0.25	-0.97	0.2369
Rosette wt_m		-0.28	-0.53	0.6964		-0.15	-0.73	0.5193
Inflorescence wt		0.81	0.79	0.9597		0.48	0.80	0.6863
Inflorescence height		0.18	0.02	0.4455		-0.01	-0.09	0.7722
Inflorescence branches		0.23	-0.07	0.3425		0.02	-0.38	0.3158
Flowering time		-0.08	-0.86	**<0.0001**		-0.08	-0.68	0.0060
Silique number		0.78	0.73	0.6411		0.68	0.83	0.3220
Average silique length		0.08	0.04	0.7673		0.13	0.53	0.0810
Root diam_v		0.06	-0.10	0.5152		0.06	-0.26	0.5425
Stomatal density		-0.05	-0.21	0.2436		-0.08	0.06	0.4271
Rosette wt_v		0.39	0.26	0.8172		0.38	-0.42	0.5942
log(SLA)		-0.01	-0.15	0.3117		-0.01	0.23	0.3230
A (whole plant assimilation)		0.04	0.12	0.5418		0.00	-0.12	0.5139
Leaf C:N		0.06	0.25	0.2721		0.01	-0.08	0.6113
δ13C		0.07	0.09	0.9331		0.09	0.07	0.9008
δ15N		0.02	0.34	0.4730		0.03	0.12	0.6353

Table S7. Selection on mean and plasticity of trait values across competition treatments. Plasticity was calculated as no competition – competition treatments. For example, selection for an increase in inflorescence height in well-watered competition treatments is indicated by a negative coefficient. None of these tests for selection on plasticity were significant using a false discovery rate of 0.05 within each water treatment.

Trait	**Well-watered**				**Drought**			
	Trait avg. coeff.	*Trait avg. p*	*Trait plast. coeff.*	*Trait plast. p*	*Trait avg. coeff.*	*Trait avg. p*	*Trait plast. coeff.*	*Trait plast. p*
AGB_m	0.2650	0.0308	-0.2166	0.0705	-0.3588	0.0209	0.1413	0.3254
Rosette wt_m	-0.1773	0.4933	-0.0116	0.9638	-0.6082	0.0057	0.3485	0.0848
Inflorescence wt	0.3517	0.0019	-0.1323	0.1775	-0.0133	0.9222	0.2081	0.1414
Flowering time	-0.3877	0.0004	-0.1145	0.1950	-0.4004	0.0105	-0.2510	0.0870
Inflorescence height	0.1203	0.0602	-0.2056	0.0034	-0.0746	0.4506	0.0756	0.4450
Inflorescence branches	0.0818	0.4116	-0.1058	0.2915	-0.0956	0.4823	-0.0801	0.5552
Silique number	0.3852	0.0000	-0.0667	0.1681	0.3362	0.0002	0.0323	0.6357
Average silique length	0.1113	0.1363	-0.1480	0.0537	0.2355	0.0135	-0.0003	0.9970
Root diam_v	0.0491	0.5826	-0.1139	0.2125	-0.0649	0.5133	0.0386	0.6958
Stomatal density	-0.1108	0.1869	-0.0565	0.4912	-0.0602	0.5424	0.1225	0.2241
Rosette wt_v	0.1423	0.5122	-0.0109	0.9597	0.0963	0.6021	-0.1698	0.3622
log(SLA)	-0.0364	0.6589	-0.1200	0.1582	0.0799	0.4150	0.0448	0.6451
Leaf C:N	0.0995	0.2216	0.0905	0.2638	-0.0327	0.7434	0.0112	0.9108
$\delta^{13}C$	0.0964	0.2224	-0.1133	0.1553	0.1617	0.0817	-0.1151	0.2039
$\delta^{15}N$	0.0346	0.6815	0.0762	0.3714	0.0688	0.4971	0.0356	0.7241