A novel resistance pathway for calcineurin inhibitors in the human pathogenic Mucorales *Mucor circinelloides*

Sandeep Vellanki[1], R. Blake Billmyre[2, 3], Alejandra Lorenzen[1], Micaela Campbell[1], Broderick Turner[1], Eun Young Huh[1], Joseph Heitman[3], and Soo Chan Lee[1#]

[1]South Texas Center for Emerging Infectious Diseases (STCEID), Department of Biology, The University of Texas at San Antonio, San Antonio, TX.

[2]Current address: Stowers Institute for Medical Research, Kansas City, MO.

[3]Department of Molecular Genetics and Microbiology, Duke University Medical Center, Durham, NC.

#Corresponding author:

Soo Chan Lee, Ph.D.

1 UTSA Circle,

South Texas Center for Emerging Infectious Diseases (STCEID),

Department of Biology,

University of Texas at San Antonio, San Antonio, TX 78249, USA

Email: soochan.lee@utsa.edu

Phone: (210) 458-5398

(Word count: Abstract: 446; Text: 4821)

Running title – Resistance to calcineurin inhibitors in Mucorales

Abstract

Mucormycosis is an emerging lethal fungal infection in immunocompromised patients. *Mucor circinelloides* (*Mucor*) is a causal agent of mucormycosis and serves as a model system to understand genetics in Mucorales. Calcineurin is a conserved virulence factor in many pathogenic fungi and calcineurin inhibition or deletion of the calcineurin regulatory subunit (CnbR) in *Mucor* results in a shift from hyphal to yeast growth. We analyzed thirty-six calcineurin inhibitor resistant or bypass mutants that exhibited hyphal growth in the presence of calcineurin inhibitors or in the yeast-locked *cnbRΔ* mutant background without carrying any mutations in known calcineurin components. We found that a majority of the mutants had altered sequence in a gene, named here *bycA* (by̲pass of c̲alcineurin A̲). *bycA* encodes an amino acid permease. We verified that both *bycAΔ*, and the *bycAΔ cnbRΔ* double mutant are resistant to the calcineurin inhibitor FK506, thereby demonstrating a novel resistance mechanism against calcineurin inhibitors. We also found that the expression of *bycA* was significantly higher in the wild type with FK506 and in the *cnbRΔ* mutants, but significantly lower in the wild type without FK506. These findings suggest that *bycA* is a negative regulator of hyphal growth and/or a positive regulator of yeast growth in *Mucor* and calcineurin suppresses the *bycA* gene at the mRNA level to promote hyphal growth. BycA is involved in the *Mucor* hyphal-yeast transition as our data demonstrates a positive correlation between *bycA* expression, protein kinase A activity, and *Mucor* yeast-growth. Also, calcineurin activity rather than hyphal morphology primarily contributes to virulence traits including phagosome maturation blockade, host cell damages, and pro-angiogenic growth factor induction during interactions with hosts.

Importance

Mucor is intrinsically resistant to most known antifungals, which makes mucormycosis treatment challenging. Calcineurin is a serine/threonine phosphatase widely conserved across eukaryotes. When calcineurin function is inhibited in Mucor, growth shifts to a less-virulent yeast growth form which makes calcineurin an attractive target for development of new antifungal drugs. Previously we identified two distinct mechanisms through which Mucor can become resistant to calcineurin inhibitors involving Mendelian mutations in the gene for FKBP12, calcineurin A or B subunits and epimutations silencing the FKBP12 gene. Here, we identified a third novel mechanism where loss of function mutations in the amino acid permease encoding the bycA gene contribute to resistance against calcineurin inhibitors. When calcineurin activity is absent, BycA can activate PKA to promote yeast growth via a cAMP-independent pathway. Our data also shows that calcineurin activity, not morphology, primarily contributes to host - pathogen interactions in the pathogenesis of Mucor.

Introduction

Mucormycosis is a severe life-threatening infection caused by fungi belonging to the order Mucorales (1). People with weakened immune systems due to diabetes mellitus, neutropenia, or hematological and solid organ transplantation are at the highest risk of acquiring this infection (2, 3). It is the third most common invasive fungal infection in hematological and allogeneic stem transplantation patients following candidiasis and aspergillosis (1, 4). Over the past decades, there has been a global and

ongoing rise in the incidence of mucormycosis, primarily due to the increasing number

of diabetic patients and increased use of immunosuppressive drugs (5-10).

Mucormycosis is also on the rise in immunocompetent individuals (11-14). Mucorales

grow as molds in the environment and produce sporangiospores that can enter the host

via inhalation resulting in pulmonary infection, through the skin due to trauma resulting

in cutaneous infections, or through the nasal passages resulting in rhinocerebral

infections (15-17). The spores can disseminate within the host, resulting in 95-100%

mortality even with antifungal drug treatment (18). Mucorales are intrinsically resistant to

most antifungals, which makes it very difficult to treat, and surgery is often required

(19).

Calcineurin is a calcium-calmodulin-dependent phosphatase conserved widely

across eukaryotes including pathogenic fungi (20, 21). Calcineurin is a heterodimer

consisting of a catalytic and a regulatory subunit, and both subunits are required for

calcineurin function. The role of calcineurin varies depending on the fungal species, for

example in *Cryptococcus neoformans* and *Cryptococcus gattii* calcineurin is required for

growth at high temperature (37°C) and at alkaline pH (22-24), while in *Candida* spp.

calcineurin contributes to azole tolerance and is required for survival in serum, among

other functions (20, 25). In *Aspergillus fumigatus*, calcineurin mutants exhibit delayed

germination, hyphal growth with irregular branching, and abnormal septum (26). We

have previously shown that in *Mucor* spp. calcineurin regulates dimorphism, in which

the calcineurin inhibitor FK506 (tacrolimus) enforces *Mucor* to grow only as yeast (27).

Hyphal morphology is the predominant growth mode for *Mucor* spp., however by

modulating respiratory conditions *Mucor* can be forced to grow as yeast as well (28).

While aerobic conditions promote hyphal growth, low oxygen and high carbon dioxide

conditions enforce yeast growth (29-32). Targeting components involved in

mitochondrial or lipid metabolism can also promote yeast growth, even under aerobic

conditions (33-36). In addition, previous studies have shown that the addition of cyclic

AMP to *Mucor* in culture results in activation of cAMP-dependent kinase Protein Kinase

A and promotes yeast growths (37-40). Wolff et al have also shown that during

anaerobic yeast growth in *Mucor* an increased expression of PKA regulatory and

catalytic subunits are exhibited than during aerobic hyphal growth (41). *Mucor* spp have

four isoforms of PKA regulatory subunits, and each is differentially expressed

depending on the growth conditions (38, 39). Calcineurin is involved in the genetic

regulation of *Mucor* dimorphism as deletion of the gene encoding the regulatory subunit

of calcineurin (CnbR) resulted in yeast-locked growth, even under aerobic conditions

(27). *cnbRΔ* mutants are avirulent in a wax moth host model thereby making calcineurin

an attractive target for antifungal treatment in mucormycosis (27). Also, mucormycosis

incidence is low in patients receiving FK506 as an immunosuppressant (42). *cnbRΔ*

mutants are also more susceptible to antifungal drugs such as Ambisome, micafungin,

and posaconazole (43).

The cellular receptor for calcineurin is FKBP12, a member of the immunophilin

protein family with cis-trans peptide prolyl isomerase activity (44). When FK506 is

bound to FKBP12, it inhibits calcineurin phosphatase activity by binding to the interface

between the calcineurin catalytic A and regulatory B subunits, thereby preventing

substrates access to the active site (45, 46). FKBP12 also binds to rapamycin to inhibit

the Tor pathway (47), and mutations in the FKBP12 gene confer resistance to both

FK506 and rapamycin. Amino acid substitutions in the calcineurin regulatory B and catalytic A subunit surfaces that interact with the FKBP12-FK506 complex can also result in resistance to FK506 (48, 49). Another immunophilin, cyclophilin A (Cyp), serves as a cellular receptor for the drug cyclosporine A (CsA). When bound to Cyp, CsA inhibits calcineurin in a similar way to FKBP12-FK506 (50). Disruption of the gene encoding Cyp therefore confers resistance to CsA.

In our previous studies, calcineurin inhibitor resistant *Mucor* strains were found to exhibit hyphal growth instead of yeast growth were found to have mutations in the FKBP12 gene or the calcineurin catalytic A or regulatory B subunit genes (27, 43, 51, 52). In addition, Calo et al., found that *Mucor* can also silence the FKBP12 gene to become transiently resistant to FK506 and rapamycin via an RNAi-dependent epimutation pathway (51, 52). In this study, we isolated mutants that do not follow the known calcineurin inhibitor resistance mechanisms. We identified a novel mechanism through which *Mucor* can become resistant to calcineurin inhibitors. We found that mutations or deletions in a novel gene, *bycA* (bypass of calcineurin A), encoding an amino acid permease confers resistance to calcineurin inhibitors or loss of calcineurin regulatory B subunit. This gene has not been previously described in the calcineurin pathway, morphogenesis, or virulence in Mucorales. As a result, *bycA* mutation allowed us to separate the yeast – hyphal morphology switch from calcineurin function and demonstrate that calcineurin activity and not morphology *per se* contributes to *Mucor* – host interactions.

Results

Isolation of calcineurin bypass mutants in the yeast-locked *cnbRΔ* mutant background

The *cnbRΔ* mutant grows exclusively as a yeast (27). However, we isolated spontaneous mutants that exhibit hyphal growth in the *cnbRΔ* background. When the mutant (10^3 cells per spot) was grown in solid YPD medium for a prolonged incubation at 30°C exceeding 5 days, we observed that hyphal sectors emerged from the yeast colonies (Fig S1A). After two rounds of single-streak dilutions, 17 independently isolated mutants (CnSp mutants) in the *cnbRΔ* background were isolated (Fig 1, Table 1, and Fig S1B). The hyphae of the mutants continued to grow, producing aerial hyphae decorated with sporangiophores containing sporangiospores (asexual spores, hereafter denoted spores). This finding shows that these mutants can complete the entire vegetative cycle without a functional calcineurin. We previously demonstrated that calcineurin function is required for hyphal growth (27). These spontaneous mutants however do not have a functional calcineurin but filament and produce spores. Thus, the mutants carry a genetic suppressor mutation(s) of the *cnbRΔ* mutation.

Isolation of calcineurin bypass mutants in the *cnaBΔ* background

The calcineurin inhibitor CsA does not fully force *Mucor* to grow as yeast, unlike FK506. Instead, only when *Mucor* lacks the *cnaB* gene, one of the three calcineurin A catalytic subunit genes (*cnaA*, *cnaB*, and *cnaC*), does CsA fully inhibit hyphal growth and enforce a yeast-locked phenotype (43). We grew the *cnaBΔ* mutant on YPD

medium containing 2 μg/ml of CsA, on which the mutant grew exclusively as yeast

because calcineurin was inhibited by CsA. Similarly, after prolonged incubation on CsA

medium, we observed that hyphal sectors emerged from yeast colonies as the strains

became resistant to CsA (Fig S2). After two rounds of single-streak dilutions, we

isolated 19 independently derived CsA-resistant (CSR) mutants (Table 2). The mutants

exhibited hyphal growth in the presence of either CsA or FK506 (Fig 2).

Surprisingly, none of the mutants carried Mendelian mutations in genes for the

calcineurin A or B subunit, FKBP12, or two subtypes of cyclophilin A (*cypA* and *cypB*)

(data not shown). In addition, the mutants did not carry epimutations in the *cypA* and

cypB genes, in which small RNA blots did not reveal any small RNAs derived from the

cypA and *cypB* genes (51) (data not shown). Therefore, we hypothesized that the CSR

mutants were also calcineurin bypass mutants that do not require calcineurin activity for

hyphal growth.

Spontaneous mutations in the *bycA* gene result in suppression phenotypes

of the calcineurin mutation

To characterize the mutation(s) that results in bypass of the calcineurin

requirement, we sequenced the whole genomes of 6 CnSp and 6 CSR mutant strains,

along with the wild type strain MU402, which was used for transformation to obtain the

cnbR and *cnaB* mutants, using the Illumina HiSeq platform. The whole genomes of

each mutant strain were compared to that of the wild type MU402. Surprisingly, all

twelve mutants carried DNA sequence modifications in a single common locus (Table 1

and Table 2). The modifications include an insertion of short sequences, single

nucleotide polymorphisms, or deletions of a shorter or longer region in the locus. The

gene in the locus was designated *bycA* (**by**pass of **c**alcineurin). We further sequenced

the *bycA* gene in the remaining CSR and CnSp mutants and found mutations in the

bycA gene in all but 5 mutants (Table 1 and Table 2). Thirty-one out of 36 suppressor

mutants contained mutations in the *bycA* gene that could result in suppression of the

calcineurin mutation.

bycAΔ mutants are resistant to calcineurin inhibitors and bycAΔ cnbRΔ

double mutants exhibit a hyphal morphology

We further verified that *bycA* is associated with the calcineurin pathway by

generating *bycA* deletion mutants. The *bycA* gene in the wild type strain MU402 was

replaced with the *pyrG-dpl237* marker (53) and gene replacement by recombination was

confirmed by 5' and 3' junction PCR (Fig S3), ORF spanning PCR (Fig S3), and

Southern blot (not shown). We also performed RT-PCR to confirm the *bycA* gene is not

expressed in the mutants (see Fig 4A). Two independent mutants, MSL47.1 and

MSL47.2 (*bycAΔ::pyrG-dpl237*)*,* were obtained. The *bycAΔ* mutants exhibited hyphal

growth and showed no growth defects (Fig 3A). On solid agar medium containing

FK506, a *bycAΔ* mutant and CnSp4 produce a larger hyphal colony and aerial hyphae,

whereas the wild type strain formed a smaller yeast colony (Fig 3B). These results

indicate that *bycAΔ* mutants are resistant to FK506.

When grown on solid YPD media containing CsA (50 µg/ml), interestingly the

bycAΔ mutants are not fully resistant to CsA, and instead exhibiting colony size

comparable to wild type (R7B). However, CnSp4 exhibited larger colonies on media

containing CsA (Fig 3B). It is possible that the CnaB catalytic A subunit is partially

resistant to the inhibition by CsA, in which Cyp-CsA may not interfere with the function

of CnaB-CnbR-calmodulin while still inhibiting CnaA-CnbR-Calmodulin and CnaC-

CnbR-calmodulin. The *bycA∆* mutants still harbor an intact *cnaB* gene and it is possible

that partial inhibition of CnaB could result in a calcineurin independent phenotype.

Therefore, *bycA∆* mutants in the *cnaB∆* background could exhibit full resistance against

CsA, which is yet to be elucidated.

We further disrupted the *cnbR* gene in the *bycA∆* background to generate *bycA∆*

cnbR∆ double mutants. The deletion and no expression of *cnbR* were confirmed by

PCR and RT-PCR (Fig S4 and Fig S5), respectively, and Southern blot (data not

shown). Two independent mutants MSL68.1 and MSL68.2 were generated

(*bycA∆::pyrG-dpl237 cnbR∆::leuA*). As shown in Fig 3A, while *cnbR∆* mutants grow

exclusively as yeast, *bycA∆ cnbR∆* double mutants exhibit filamentous growth like wild

type and the *bycA∆* mutant. *bycA∆ cnbR∆* mutants are also completely resistant to

either FK506 or CsA (Fig 3B), as they exhibit normal hyphal growth even in the

presence of calcineurin inhibitors (Fig 3B and 3C). Taken together our results

genetically validate our hypothesis that *bycA* mutations bypass the requirement of

calcineurin for hyphal growth or suppress the lack of calcineurin.

Calcineurin regulates BycA at the mRNA level

To determine a genetic link between calcineurin and BycA, the expression of

bycA was examined under conditions in which calcineurin is either functional or non-

functional. As shown in Fig 4A, compared to WT *bycA* gene expression was significantly

higher in conditions in which calcineurin is not functional such as *cnbRΔ* mutant or the

wild type in the presence of FK506. These results suggest that calcineurin negatively

regulates the *bycA* gene at the mRNA level and BycA expression is positively correlated

with *Mucor* yeast growth.

If grown in anaerobic (or microaerobic) conditions with high CO_2, *Mucor* also

grows as a yeast (54). Interestingly, under this condition, *bycA* expression remained low

(Fig 4A). When *bycAΔ* or *bycAΔ cnbRΔ* mutants were grown anaerobically with high

CO_2 levels, they still grew as yeast (Fig 4B). These results suggest that BycA is involved

in aerobic yeast growth but not in anaerobic yeast growth. Whether calcineurin inhibits

the expression of the *bycA* gene transcriptionally or regulates the stability of the *bycA*

mRNA remains to be elucidated.

BycA serves as a link between calcineurin and protein kinase A

Previous studies by our groups and others have shown that addition of

bicarbonate ions to *Mucor* culture, or by growth in high carbon dioxide conditions, is

sufficient to induce yeast growth (27, 31), as the bicarbonate ions can activate adenylyl

cyclase which results in the generation of cAMP and in turn activates cAMP-dependent

kinase – protein kinase A (PKA) (55, 56) (Fig 8). Studies have also shown that there is

an inverse relationship between PKA and hyphal morphology as the PKA regulatory

subunit gene that inhibits the activity of PKA is more highly expressed during hyphal

growth (41). Also, high levels of cAMP promote yeast growth, while low levels of cAMP

are linked with hyphal morphology (28). Interestingly, even under aerobic conditions

when *Mucor* was forced to grow as yeast by inhibiting calcineurin activity either

genetically or by using FK506, the overall cellular PKA activity remained high (27). Therefore, PKA plays pivotal roles in yeast growth. Interestingly, in wild type when calcineurin is fully functional, the cellular PKA activity is low (27), suggesting an inverse correlation between PKA and calcineurin during morphogenesis. However, it was not clear how calcineurin and PKA are linked.

BycA is a putative amino acid permease and predicted to have ten transmembrane domains and one pectinesterase domain (Fig S6). As shown in Table 3, when *cnbR*Δ and *bycA*Δ *cnbR*Δ mutants were grown on YNB medium with either methionine, threonine, or arginine as a sole nitrogen source, the *cnbR*Δ mutants exhibited growth similar to growth on YNB complete medium, while *bycA*Δ *cnbR*Δ mutants did not produce hyphal mass. This shows that BycA is a bona fide amino acid permease. In *Saccharomyces cerevisiae* and *Candida albicans*, the general amino acid permease Gap1 is known to activate PKA in a cAMP-independent manner (57, 58). We hypothesized that in the absence of calcineurin activity, BycA may activate PKA to promote yeast-growth. To this end, we measured overall cellular PKA activity in the presence or absence of the *bycA* gene. As shown in Fig 4C, PKA activity was significantly higher in WT-yeast, *cnbR*Δ and WT + FK506 (when *bycA* expression was significantly higher) compared to WT - hyphae (when *bycA* expression is low). On the other hand, when the *bycA* gene is deleted, the *bycA*Δ or *bycA*Δ *cnbR*Δ mutants exhibited significantly lower cellular PKA activity compared to the conditions when the *bycA* gene is expressed. No significant differences were observed between WT - hyphae, *bycA*Δ, or *bycA*Δ *cnbR*Δ isolates.

Interestingly, when we measured cAMP levels, compared to WT – yeast, the

cAMP levels were significantly lower in *cnbRΔ* and WT + FK506 yeast groups despite

high PKA activity (Fig 4D). This suggests that in aerobic conditions when calcineurin is

absent, BycA increases PKA activity via a cAMP-independent pathway, which has yet to

be elucidated.

Phagosome maturation blockade upon phagocytosis by macrophages is

dependent on calcineurin, not morphology

Phagocytosis followed by phagosome maturation or acidification in macrophages

are essential innate immune pathways to control pathogens. Previously we have shown

that when the macrophage cell line J774.A1 and primary macrophages are challenged

with wild type spores or *cnbRΔ* yeast, they rapidly phagocytosed both spores and yeast,

however only the macrophages with *cnbRΔ* yeast underwent phagosome maturation

(43). This indicates *Mucor* spores escape innate immunity by blocking phagosome

maturation as spores survive better than *cnbRΔ* yeast during co-culture with

macrophages. However, it was not clear if the blockade of phagosome maturation by

Mucor is dependent on its morphology or functional calcineurin because yeast cells lack

calcineurin function. To address this question, we co-cultured WT spores, *cnbRΔ* yeast,

and *bycAΔ cnbRΔ* double mutant spores with macrophages to monitor phagosome

maturation. Macrophages containing *Mucor* spores or yeast cells were stained with

Lysotracker Green DND-26 (Thermofisher). Lysotracker only stains acidic organelles in

cells such as lysosomes or mature phagosomes and therefore can be used to

determine whether phagosomes containing *Mucor* cells are acidic, an indication of

phagosome maturation. As shown in Fig 5, only about 20% of macrophages challenged

with either wild type (n=480) or *bycAΔ* (n=290) spores underwent phagosomal

maturation, which is significantly lower when compared to macrophages challenged with

cnbRΔ (n=279; yeast-locked). Interestingly, ~80% phagosomes containing *bycAΔ*

cnbRΔ double mutants (n=413) spores underwent maturation. Our data suggest that it is

not morphology but calcineurin that is involved in inhibition of phagosome maturation.

Mucor without functional calcineurin causes reduced cell damage and

induces less FGF-2 expression in endothelial cells

Mucormycosis is an angioinvasive disease, and hence Mucorales interaction and

subsequent damage to the endothelium lining the blood vessels is an important step in

disease pathology (59). We have previously shown that *Mucor* hyphae but not yeast

(*cnbRΔ*) induces FGF-2 protein expression in lymphoblastoid cell lines and bone

marrow macrophages (43, 60). We wanted to determine the extent to which calcineurin

contributes to the endothelial damage and FGF-2 protein response by *Mucor.* To this

end, we used Human Umbilical Vein Endothelial Cells (HUVECs) and challenged with

wild type, *bycAΔ*, *cnbRΔ* or the *bycAΔ cnbRΔ* double mutant for a period of 24 hours.

When compared to HUVECs challenged with wild type or *bycAΔ* spores, the *cnbRΔ* and

bycAΔ cnbRΔ double mutant caused 70% less cytotoxicity in HUVECs as quantified by

measuring LDH levels (Fig 6A) and induced significantly less FGF-2 protein production

(Fig 6B). These data suggest that calcineurin but not morphology governs *Mucor*

interaction with endothelial cells.

bycAΔ cnbRΔ double mutants are less virulent in a _Galleria mellonella_ host model of mucormycosis

The yeast-locked _cnbRΔ_ mutant is less virulent than wild type in a _G. mellonella_ (wax moth) host (27). To test if calcineurin is involved in virulence, we injected wax moths (n=15/group) with 10,000 spores of either wild type, _bycAΔ_, _cnbRΔ_, the _bycAΔ cnbRΔ_ double mutant, or PBS (mock) and survival was monitored for 7 days. As shown in Fig 7, all of the wax moth larvae challenged with wild type or _bycAΔ_ spores succumbed to infection within a week, while _cnbRΔ_ and _bycAΔ cnbRΔ_ mutants exhibited a significantly higher survival rate. Interestingly, only about 65% of the wax moth infected with CnSp4 (spontaneous double mutant) or _bycAΔ cnbRΔ_ survived, while 100% of the wax moth infected with _cnbRΔ_ survived. We also injected wax moth larvae with 10,000 (1x), 20,000 (2x) or 30,000 spores (3x) of the _bycAΔ cnbRΔ_ double mutant; as shown in Fig S7, it was only at 3x spore inoculum when 100% mortality was achieved.

To determine virulence in an immunocompromised murine host, the wild type (R7B), _bycAΔ_, _cnbRΔ_, _bycAΔ cnbRΔ_ double mutant, or PBS (mock) were inoculated via the intratracheal or intravenous routes. However, there was no significant difference in survival between the infected groups (Fig S8). It is interesting that calcineurin mutants are significantly less virulent in the wax moth larval host but are similarly virulent as WT in the murine model.

Discussion

Calcineurin is conserved widely across pathogenic fungi and is involved in cell wall integrity, morphogenesis, and virulence (20, 21, 61). We have previously shown that when grown in the presence of a calcineurin inhibitor - FK506, *Mucor* exhibits yeast morphology even under aerobic conditions; however, when resistant to FK506 *Mucor* grows as hyphae (27). FKBP12 is the drug-receptor for FK506, and when bound, the FKBP12-FK506 complex inhibits the phosphatase activity of calcineurin by binding to the interface of the calcineurin regulatory and catalytic subunits (45). One of the common ways through which most pathogenic microbes become resistant to drugs is having mutations in the drug-receptor, in this case FKBP12 or catalytic A subunit or regulatory B subunits of calcineurin (27). Calo et al. have shown that *Mucor* can be transiently resistant to FK506 by triggering an RNA interference specific to the gene encoding FKBP12, resulting in silencing of the drug target gene (51, 52). In this study, we identified a novel resistance mechanism through which *Mucor* can also become resistant to calcineurin inhibitors. WGS and targeted sequencing of the genomes of spontaneous mutants (CnSp and CSR mutants) revealed DNA sequence alterations in the *bycA* locus in 31 out of 36 resistant isolates (Table 1 and Table 2).

A novel link between calcineurin and BycA, an amino acid permease in *Mucor*

Based on our sequencing analysis we hypothesized that deletion of *bycA* should result in stable resistance to calcineurin inhibitors. Indeed, *bycAΔ* mutants are resistant to FK506. Unlike *cnbRΔ* mutants, which grow as yeast, the *bycAΔ cnbRΔ* mutant fully

grow as hyphae like WT (Fig 3). This verifies that *Mucor* can become resistant to calcineurin inhibitors by having a second mutation in the *bycA* gene. *Mucor* has a secondary conserved mechanism that promotes hyphal growth even in the absence of a functional calcineurin. Interestingly, however, we did not observe Mendelian DNA mutations in the *bycA* locus in five of the mutants (out of 36) we performed sequencing on even though they exhibit the same phenotype as the remaining suppressors; one possibility is that *Mucor* may also produce small RNAi targeting *bycA* or there could be other mechanism(s) that confer *Mucor* resistance against calcineurin inhibitors. Future studies are required to elucidate these mechanisms further.

How is BycA linked to calcineurin and morphogenesis? It is not known if BycA is a direct post-translation modification target of calcineurin. However, our data suggest that calcineurin negatively regulates the expression of BycA at the mRNA level. It is also possible that calcineurin is involved in the stability of the *bycA* mRNA. Further investigation is ongoing to elucidate how calcineurin regulates expression of BycA. Interestingly, Chow et al revealed that, in *Cryptococcus neoformans*, a gene encoding a putative amino acid permease (CNAG_01118) is overexpressed when calcineurin is deleted and this is independent of Crz1, a well-known target of calcineurin (62). Whether deletion of the amino acid gene would result in suppression of calcineurin mutant phenotypes remains to be tested in *C. neoformans*.

BycA as a missing link between calcineurin and PKA

Studies by us and others have shown that *Mucor* morphology is primarily dependent on: a) respiratory conditions and b) calcineurin and PKA activity (27-30, 37-

39, 41). While aerobic conditions promote hyphal growth, *Mucor* grown in anaerobic

conditions with low oxygen levels undergoes yeast growth. In anaerobic conditions with

high CO_2, *Mucor* grows as yeast, and has PKA activity increased via the bicarbonate-

cAMP pathway (27, 28, 30). Previously we have shown that even under aerobic

conditions when calcineurin function is inhibited by FK506 or absent in the *cnbRΔ*

mutant, *Mucor* grows as a yeast and PKA activity remained high (27). The bicarbonate-

cAMP pathway is likely not involved in aerobic conditions as the carbon dioxide levels

are very low. It has been suggested that there is an antagonistic relationship between

calcineurin and PKA in fungal systems in *Ustilago maydis* and *S. cerevisiae* (63, 64).

In this study, we found that calcineurin and PKA are inversely related through

BycA, in which the overall PKA activity was lower in wild type hyphae with lower *bycA*

expression, but higher in yeast with higher *bycA* expression (Fig 4A and 4C). PKA

activity also remained lower in the *bycAΔ cnbRΔ* double mutant because even though

calcineurin function is suppressed, the double mutant does not have BycA function to

activate PKA and promote yeast growth. For the same reason, the *bycAΔ cnbRΔ* double

mutants exhibit hyphal morphology. We also found that the amino acid permease BycA

can activate PKA without a requirement for cAMP. Our finding that BycA can activate

PKA is congruent with previous findings in *C. albicans* and *S. cerevisiae*, where the

general amino acid permease Gap1 can activate PKA in a cAMP-independent manner

(57, 58). It is possible that this is achieved through imbalance between the catalytic and

regulatory subunits of PKA. In *S. cerevisiae*, it has been reported that kelch repeat

homolog proteins 1 and 2 (Krh1 and Krh2) promote the association between PKA

catalytic and regulatory subunits, and deletion of *krh1/2* leads to lower cAMP

requirement for PKA activation (65). Since Krh proteins are evolutionarily conserved in

eukaryotes, it is possible that such a mechanism may also exist in *Mucor.* Our future

studies will further focus on determining how BycA activates PKA in *Mucor*.

Our previous studies showed that cellular PKA activity is elevated when

calcineurin is inhibited in *C. neoformans* and *Rhizopus delemar* (27). How this link is

achieved in these fungi remains be elucidated. In *C. neoformans*, there is evidence that

higher expression of an amino acid permease is correlated with the phenotypes

regulated by PKA; treatment with urea results in a 27-fold increase of an amino acid

permease (CNAG_01118) and increased capsule production that is known to be

regulated by PKA (66-68). Our current study investigates the links between calcineurin,

amino acid permease, and PKA which could also exist in *C. neoformans* and possibly

other fungi.

Calcineurin governs virulence and plays important roles in host-pathogen

interactions in *Mucor*

Studies from several pathogenic fungi have shown that morphology is linked with

expression of virulence factors, colonization of the host, and evasion of host immune

responses, etc (69, 70). The *cnbR*Δ mutants are significantly less virulent in a

heterologous wax moth larva host model (27); however, it was not clear if the

diminished virulence potential of *cnbR*Δ mutants was due to morphology, loss of the

calcineurin, or both. In the current study using a hyphal growth strains lacking

calcineurin - *bycA*Δ *cnbR*Δ double mutant, we found that it is calcineurin and not

morphology *per se* that primarily contributes to *Mucor* - host interactions and virulence.

For example, macrophages challenged with mutants that lack calcineurin displayed

significantly higher phagosome maturation compared to wild type, irrespective of their

morphology (Fig 5). The ability of *Mucor* to cause damage to the endothelium was

dependent upon calcineurin (Fig 6A).

Angiogenesis is the process by which new blood vessels arise from pre-formed

blood vessels. Proteins such as Fibroblast Growth Factor-2 (FGF-2) and Vascular

Endothelial Growth Factor (VEGF) promote angiogenesis (71). We have previously

shown that the ability of *C. albicans* to induce FGF-2 is morphology-dependent, as non-

filamentous strains of *C. albicans* fail to induce FGF-2 (72). We have also shown that in

Mucor only WT, and not *cnbR*Δ mutants induce a host FGF-2 response (43). In this

study we further found that both *bycA*Δ *cnbR*Δ double mutant also fails to induce FGF-2

(Fig 6B). Our data suggests that the ability of *Mucor* to induce host FGF-2 is dependent

on calcineurin and not on morphology. Studies by us and others have shown that

candidalysin from *C. albicans* and gliotoxins from *Aspergillus fumigatus* regulate the

host FGF-2 response (72, 73). Hence, it is possible that toxins from *Mucor* (74, 75) also

regulate host FGF-2 response. We are currently working on identifying the *Mucor*

factor(s) that facilitates *Mucor* interactions with endothelial cells to induce FGF-2

response.

Larvae of *Galleria mellonella* have largely been used to study virulence and for

testing antifungal drugs (27, 47, 74). The wax moth larvae possess an innate immune

system that at the humoral and cellular levels is both structurally and functionally similar

to mammals (76). We have previously shown that *cnbR*Δ mutants are avirulent in this

host (27). In this study, we have shown that *bycA*Δ *cnbR*Δ double mutants are also less

virulent (Fig 7). It is congruent that more wild-type spores survived compared to *cnbRΔ*

yeast during interaction with bone marrow murine macrophages (Figure S9).

Interestingly, unlike in the wax moth host system, in the pulmonary and systemic

murine model of mucormycosis, the *cnbRΔ* mutant also caused significant mortality like

wild type. This observation may be due to the neutropenic host conditions caused by

treatment with cyclophosphamide and cortisone. Calcineurin is required to escape

innate immune cells by blocking phagosome maturation; however, lack or significant

lower numbers of phagocytic cells resulted in higher mortality by the yeast-locked

cnbRΔ mutant. Alternatively, *Mucor* yeast is more immunogenic compared to *Mucor*

spores and therefore the yeast might have caused mortality in a different manner than

spores (43). This could be an analogy to the *Cryptococcus rim101* mutants (77). The

rim101 mutant lacks virulence traits and yet is more immunogenic compared to wild

type; nevertheless, the mutants still cause mortality in a murine infection model via a

substantial host immune response. Further study is required to determine if *Mucor*

cnbRΔ yeast cause hyper immune responses in a murine lung infection model.

In summary, we identified a novel mechanism through which *Mucor* can become

resistant to calcineurin inhibitors, where loss-of-function mutations in the *bycA* gene

confer resistance against calcineurin inhibitors. This resistance mechanism is achieved

by BycA regulating the activity of protein kinase A via a cAMP-independent pathway yet

to be elucidated (Fig 8). As calcineurin is a major virulence factor in many pathogenic

fungi, it is worth investigating if this relationship between calcineurin, BycA, and PKA is

also conserved in other pathogenic fungal systems.   Calcineurin also governs key

Mucor – host interactions and is an attractive target for developing antifungals to treat

mucormycosis.

Materials and Methods

Ethics Statement

All animal experiments were conducted at the University of Texas at San Antonio

(UTSA) in accordance with the Institutional Animal Care and Use Committee (IACUC)

guidelines and in full compliance with the United States Animal Welfare Act (Public Law

98-198) and National Institute of Health (NIH) guidelines. The animal protocol - MU104

used in this study was approved by the UTSA IACUC. The experiments were conducted

in the Division of Laboratory Animal Resources (DLAR) facilities that are accredited by

the Association for Assessment and Accreditation of Laboratory Animal Care

(AAALAC).

Fungal Strains and Growth Conditions

All fungal strains and plasmids used in this study are listed in Table S1. For

spore production, *Mucor* strains were inoculated and maintained on yeast peptone

glucose (YPG, 3 g/L yeast extract, 10 g/L peptone, 20 g/L glucose, 2% agar, pH=4.5) or

yeast peptone dextrose (YPD, 10 g/L yeast extract, 20 g/L peptone, 20 g/L glucose, 2% agar,

pH=6.5) agar at 26°C or 30°C for four days. *cnbR∆* mutants were maintained on YPD

agar at 30°C. For in vitro, *Mucor* – host interaction studies or in vivo survival studies

cnbR∆ mutants were grown overnight in liquid YPD at 30°C with shaking. For high CO_2

conditions, the flasks were entirely filled with YPD broth, and the wild type (R7B) was

inoculated on the bottom of the flasks. The flasks were sealed with parafilm and were

left at room temperature overnight without disturbing. For testing with calcineurin

inhibitors, *Mucor* strains were grown in liquid YPD or on YPD agar plates supplemented

with FK506 (Astellas Pharma Inc.; 1 µg/ml) or CsA (LC laboratories; 2 or 100 µg/ml) at

30°C for 2 to 5 days.

Generation of suppressor mutants CnSp and CSR

For generation of CnSp mutants, 10^3 cells of *cnbR∆* mutants were spotted on

YPD agar and grown at 30°C for 5 or more days. Hyphal sectors emerging from yeast

colonies were propagated to fresh YPD plates. For CSR mutant generation, *cnaB∆*

mutants were grown on YPD agar containing CsA (2 µg/ml) for 5 days or longer, and

the hyphal colonies were propagated further.

Whole Genome Sequencing and targeted sequencing of CnsP and CSR

mutants

Genomic DNA of the 12 spontaneous mutant strains and MU402 strain was used

for sequencing. A single end Truseq library was constructed with the genomic DNA and

sequenced on the Ilumina HiSeq 2000 platform at the University of North Carolina at

Chapel Hill School of Medicine. The reads were mapped to the genome of CBS277.49

(78) using the short-read component of BWA (79). SNP calling was performed using the

Genome Analysis Toolkit (GATK ver. 2.4-9) pipeline and the Unified Genotyper with the

haploid setting (80).

Disruption of genes

The primers used in the study are listed in Table S2. To disrupt *bycA* gene,

nearly 1 kb upstream and downstream sequences flanking the *bycA* gene were PCR

amplified (Phusion® High-Fidelity DNA Polymerase, NEB) using primers SL3 and SL4,

and SL7 and SL8 respectively. The *pyrG-dpl237* marker (53) was amplified using SL5

and SL6 primers. All three fragments were gel excised, and 75 ng of each fragment was

amplified using overlap PCR strategy with primers SL9 and SL10 such that the marker

is placed between the flanking regions.   The product was cloned into a TOPO vector

(pSL26), which was then linearized using Sma1 enzyme and transformed into *Mucor*

strain Mu402 (*pyrG⁻ leuA⁻*) via electroporation (81). The transformants were selected on

minimal media with casamino acids pH=3.2 (MMC, 10 g casamino acids, 0.5 g yeast

nitrogen base without amino acids and ammonium sulfate, 20 g glucose, 1 mg niacin, 1

mg thiamine, and 15 g agar in 1 L dH$_2$O) containing 0.5 M sorbitol as described

previously (81). Two independent transformants (MSL47.1 and MSL47.2; Table S1A)

out of 12 were positive for disruption of *bycA.*

For disruption of the *cnbR* gene, 5' and 3' flanking regions were amplified using

SL243 and, and SL281, and SL282 and SL244 respectively. A *leuA* marker was

amplified using SCL737 and SCL738. Overlap PCR was performed using primers

SCL286 and SCL287. The pSL58 plasmid containing the disruption cassette was

linearized and used to transform MSL47.2. The transformants were selected on YNB

medium (1.5 g ammonium sulfate, 1.5 g glutamic acid, 0.5 g yeast nitrogen base without

amino acids and ammonium sulfate, 10 g glucose, 20 g agar in 1 L dH$_2$O). Two out of

18 independent transformants (MSL68.1 and MSL68.2) were positive for disruption of

cnbR.

550 Reverse Transcriptase qPCR

Based on the ORF sequences, we designed primers for *bycA* (SL346 and

SL347) and *cnbR* (SCL 578 and SCL 579) that span across two exons. *Mucor* strains

were grown overnight in YPD liquid medium overnight, and the next day total RNA was

isolated using TRIzol reagent according to the manufacturer's instructions. 1 µg of RNA

from each sample was used for cDNA synthesis (High Capacity cDNA Reverse

Transcription Kit with RNase Inhibitor, Applied Biosystems), and qPCR was performed

using SYBR Green QPCR Master mix (Thermo Scientific). Actin gene (primers SCL368

and SCL369) served as internal control. Three independent replications were performed

for each experiment with RNA obtained from two independent preparations.

561 Protein kinase A activity assay

562   *Mucor* strains were grown in appropriate conditions overnight and collection of

563 crude protein extracts, and overall PKA activity measurement was performed using The

DetectX® PKA (Protein Kinase A) Activity Kit according to the manufacturer's instructions. Briefly, crude protein extracts were obtained by subjecting the samples to activated cell lysis buffer with protease inhibitor cocktail, PMSF, and sodium orthovanadate. Each sample was vigorously shaken in a bead beater (1 minute x 5 times with 1 minute cooling) followed by centrifugation to collect supernatants containing crude extracts. Total protein was quantified using the Bradford method (Biorad). 0.5 μg of each sample was used to determine overall PKA activity according to the manufacturer's instructions.

cAMP determination

$Mucor$ strains were grown in appropriate conditions overnight and the next day cells corresponding to 60 mg (wet weight) were flash frozen and immersed in 0.1 M HCl. Crude extracts were obtained as described above. cAMP levels were measured according to the manufacturer's instructions (Direct cAMP ELISA kit, Enzo).

Phagosome maturation assay

Murine macrophage cell line J774.A1 was maintained in Dulbecco's modified Eagle Medium (DMEM) with 10% FBS, and 200 U/ml mixture of penicillin and streptomycin antibiotics at 37°C, 5% CO_2. Upon confluence, a cell scraper was used to detach cells, and 5 x 10^5 cells were plated in each well of a 24 well glass plate. After 24 hours, $Mucor$ spores or yeast were added at a multiplicity of infection of 1 along with 10

μM LysoTracker™ Green DND-26, and 1 μg/ml Hoechst 33342 stain. 30 minutes later a

Zeiss Axio Observer microscope was used to image fields containing macrophages and

Mucor cells. The experiment was performed in triplicate and was repeated on two

different occasions.

LDH assay and FGF-2 ELISA

Primary Human Umbilical Vein Endothelial cells (HUVECs) were purchased from

Lonza and were maintained in Endothelial Basal Medium (EBM) containing

hydrocortisone, ascorbic acid, Insulin Growth Factor, heparin, and FBS at 37°C, 5%

CO_2 according to the manufacturer's instructions. Confluent cells were trypsinized and 5

x 10^3 cells/well were seeded in a 96 well plate. After 24 hours, 5 x 10^4 of appropriate

fungal cells in PBS (MOI =10) or an equal volume of PBS were added to each well. 24

hours post infection, the supernatant was collected to quantify LDH levels (The CytoTox

96® Non-Radioactive Cytotoxicity Assay, Promega) or FGF-2 protein levels using

ELISA (R&D systems). The LDH release was calculated as described previously (82).

Virulence test

For production of *Mucor* spores, appropriate strains were grown on YPG agar at

26°C for 4 days under light. The *cnbRΔ* mutant was grown overnight in YPD broth at

30°C with aeration. Both the spores and yeast were washed two times with PBS, and

different inoculums ranging from 1 x 10^4 to 3 x 10^4 cells in 2 μl PBS were injected into

the wax moth host through the last left proleg. Differences between the survival curves

were evaluated for significance using the Kaplan-Meier test. The experiment was

performed on two different occasions with n = 15 animals for each group.

6 week old CD1 mice were immunocompromised with cyclophosphamide (250

mg/kg via the intraperitoneal route) and cortisone acetate (500 mg/kg via the

subcutaneous route) every 5 days starting 2 days before inoculation (83). On the day of

inoculation, the mice were anaesthetized using isoflurane and 1.5 x 10^6 spores or yeast

cells in PBS were introduced via the intratracheal route (pulmonary) were monitored

routinely. Two independent experiments were performed with n=5 for each group. Data

represented is from a single experiment.

Statistics

618   Prism (Version 7, GraphPad Software Inc) was used to perform statistical

619 analysis. A P value≤0.05 was considered significant.

Acknowledgements

622 We are indebted to Jose Lopez-Ribot, Chiung -Yu Hung, Astrid Cardona, Stephen

623 Saville, Ashraf Ibrahim, and Praveen Juvvadi for valuable discussions. We also thank

624 Anna, Averette, Gloria Adedoyin, and Alexis Garcia for technical support for this study.

625 This work was supported by NIH/NIAID R03 AI119617, a Korean Food Research

626 Institution (KFRI) grant, and UTSA Research funds to S.C.L. S.C.L. holds a Voelcker

Fund Young Investigator Pilot Award from the MAX AND MINNIE TOMERLIN VOELCKER FUND. J.H. was supported by NIH/NIAID R37 AI39115-21, R01 AI50113-15, and PO1 AI104533-05. J.H. is co-director and fellow of the Canadian Institute for Advanced Research Program Fungal Kingdom: Threats & Opportunities.

References

1.   Petrikkos G, Skiada A, Lortholary O, Roilides E, Walsh TJ, Kontoyiannis DP. 2012. Epidemiology and clinical manifestations of mucormycosis. Clin Infect Dis 54 Suppl 1:S23-34.

2.   Baldin C, Ibrahim AS. 2017. Molecular mechanisms of mucormycosis-The bitter and the sweet. PLoS Pathog 13:e1006408.

3.   Prabhu RM, Patel R. 2004. Mucormycosis and entomophthoramycosis: a review of the clinical manifestations, diagnosis and treatment. Clin Microbiol Infect 10:31-47.

4.   Eucker J, Sezer O, Graf B, Possinger K. 2001. Mucormycoses. Mycoses 44:253-260.

5.   Ambrosioni J, Bouchuiguir-Wafa K, Garbino J. 2010. Emerging invasive zygomycosis in a tertiary care center: epidemiology and associated risk factors. Int J Infect Dis 14 Suppl 3:e100-3.

6.   Prakash H, Chakrabarti A. 2019. Global epidemiology of mucormycosis. J Fungi 5(1):26.

7.   Saegeman V, Maertens J, Meersseman W, Spriet I, Verbeken E, Lagrou K. 2010. Increasing incidence of mucormycosis in University hospital, Belgium. Emerg Infect Dis 16:1456-8.

8.   Kontoyiannis DP, Yang H, Song J, Kelkar SS, Yang X, Azie N, Harrington R, Fan A, Lee E, Spalding JR. 2016. Prevalence, clinical and economic burden of mucormycosis-related hospitalizations in the United States: a retrospective study. BMC Infect Dis 16:730.

9.   Torres-Narbona M, Guinea J, Martínez-Alarcón J, Muñoz P, Gadea I, Bouza E.

2007. Impact of zygomycosis on microbiology workload: a survey study in Spain. J Clin

Microbiol 45:2051-3.

10.   Petrikkos G, Skiada A, Drogari-Apiranthitou M. 2014. Epidemiology of

mucormycosis in Europe. Clin Microbiol Infect 20 Suppl 6:67-73.

11.   Mignogna MD, Fortuna G, Leuci S, Adamo D, Ruoppo E, Siano M, Mariani U.

2011. Mucormycosis in immunocompetent patients: a case-series of patients with

maxillary sinus involvement and a critical review of the literature. Int J Infect Dis

15:e533-40.

12.   Venkatesh D, Dandagi S, Chandrappa PR, Hema KN. 2018. Mucormycosis in

immunocompetent patient resulting in extensive maxillary sequestration. J Oral

Maxillofac Pathol 22:S112-s116.

13.   Shatriah I, Mohd-Amin N, Tuan-Jaafar TN, Khanna RK, Yunus R, Madhavan M.

2012. Rhino-orbito-cerebral mucormycosis in an immunocompetent patient: case report

and review of literature. Middle East Afr J Ophthalmol 19:258-61.

14.   Pozo Laderas JC, Pontes Moreno A, Pozo Salido C, Robles Arista JC, Linares

Sicilia MJ. 2015. Mucormicosis diseminadas en pacientes sin inmunodeficiencias: una

enfermedad que también existe. Revista Iberoamericana de Micología 32:63-70.

15.   Fanfair R, Benedict K, Bos J, D Bennett S, Lo Y-C, Adebanjo T, Etienne K, Deak

E, Derado G, Shieh W-J, Drew C, Zaki S, Sugerman D, Gade L, H Thompson E, Sutton

D, Engelthaler D, Schupp J, E Brandt M, J Park B. 2012. Necrotizing cutaneous

mucormycosis after a tornado in Joplin, Missouri, in 2011. N Engl J Med 367:2214-

2225.

16. Muqeetadnan M, Rahman A, Amer S, Nusrat S, Hassan S, Hashmi S. 2012.

Pulmonary mucormycosis: an emerging infection. Case Rep Pulmonol 2012:3.

17. Tribble DR, Rodriguez CJ. 2014. Combat-related invasive fungal wound

infections. Curr Fungal Infect Rep 8:277-286.

18. Katragkou A, Walsh TJ, Roilides E. 2014. Why is mucormycosis more difficult to

cure than more common mycoses? Clin Microbiol Infect 20:74-81.

19. Spellberg B, Edwards J, Jr., Ibrahim A. 2005. Novel perspectives on

mucormycosis: pathophysiology, presentation, and management. Clin Microbiol Rev

18:556-69.

20. Juvvadi PR, Lee SC, Heitman J, Steinbach WJ. 2017. Calcineurin in fungal

virulence and drug resistance: prospects for harnessing targeted inhibition of calcineurin

for an antifungal therapeutic approach. Virulence 8:186-197.

21. Park H, Lee SC, Cardenas ME, Heitman J. 2019. Calcium-calmodulin-calcineurin

signalling: a globally conserved virulence cascade in eukaryotic microbial pathogens.

Cell Host Microbe 26:453-462.

22. Odom A, Muir S, Lim E, Toffaletti DL, Perfect J, Heitman J. 1997. Calcineurin is

required for virulence of *Cryptococcus neoformans*. Embo J 16:2576-89.

23. Chen YL, Lehman VN, Lewit Y, Averette AF, Heitman J. 2013. Calcineurin

governs thermotolerance and virulence of *Cryptococcus gattii*. G3: Genes, Genomes,

Genetics 3:527-39.

24. Del Poeta M, Cruz MC, Cardenas ME, Perfect JR, Heitman J. 2000. Synergistic

antifungal activities of bafilomycin A(1), fluconazole, and the pneumocandin MK-

0991/caspofungin acetate (L-743,873) with calcineurin inhibitors FK506 and L-685,818

against *Cryptococcus neoformans*. Antimicrob Agents Chemother 44:739-46.

25.   Blankenship JR, Wormley FL, Boyce MK, Schell WA, Filler SG, Perfect JR,

Heitman J. 2003. Calcineurin is essential for *Candida albicans* survival in serum and

virulence. Eukaryot Cell 2:422-30.

26.   Juvvadi PR, Lamoth F, Steinbach WJ. 2014. Calcineurin-mediated regulation of

hyphal growth, septation, and virulence in *Aspergillus fumigatus*. Mycopathologia

178:341-348.

27.   Lee SC, Li A, Calo S, Heitman J. 2013. Calcineurin plays key roles in the

dimorphic transition and virulence of the human pathogenic zygomycete *Mucor*

circinelloides. PLoS Pathog 9:e1003625.

28.   Orlowski M. 1991. *Mucor* dimorphism. Microbiol Rev 55:234-58.

29.   Bartnicki Garcia S. 1963. Symposium on biochemical bases of morphogenesis in

fungi. III. Mold-yeast dimorphism of *Mucor*. Bacteriol Rev 27:293-304.

30.   Bartnicki-Garcia S, Nickerson WJ. 1962. Induction of yeast-like development in

Mucor by carbon dioxide. J Bacteriol 84:829-40.

31.   Bartnicki-Garcia S, Nickerson WJ. 1962. Assimilation of carbon dioxide and

morphogenesis *Mucor rouxii*. Biochim Biophys Acta 64:548-51.

32.   Bartnicki-Garcia S, Nickerson WJ. 1962. Nutrition, growth, and morphogenesis of

Mucor rouxii. J Bacteriol 84:841-58.

33.   Zorzopulos J, Jobbagy AJ, Terenzi HF. 1973. Effects of

ethylenediaminetetraacetate and chloramphenicol on mitochondrial activity and

morphogenesis in *Mucor rouxii*. J Bacteriol 115:1198-204.

34. Clark-Walker GD. 1973. Relationship between dimorphology and respiration in

Mucor genevensis studied with chloramphenicol. J Bacteriol 116:972-980.

35. Ito ET, Cihlar RL, Inderlied CB. 1982. Lipid synthesis during morphogenesis of

Mucor racemosus. J Bacteriol 152:880-7.

36. Garcia JR, Hiatt WR, Peters J, Sypherd PS. 1980. S-adenosylmethionine levels

and protein methylation during morphogenesis of *Mucor racemosus*. J Bacteriol

142:196-201.

37. Lubbehusen T, Gonzalez Polo V, Rossi S, Nielsen J, Moreno S, McIntyre M,

Arnau J. 2004. Protein kinase A is involved in the control of morphology and branching

during aerobic growth of *Mucor circinelloides*. Microbiol 150:143-50.

38. Ocampo J, McCormack B, Navarro E, Moreno S, Garre V, Rossi S. 2012. Protein

kinase A regulatory subunit isoforms regulate growth and differentiation in *Mucor*

circinelloides: essential role of PKAR4. Eukaryot Cell 11:989-1002.

39. Ocampo J, Fernandez Nuñez L, Silva F, Pereyra E, Moreno S, Garre V, Rossi S.

2009. A subunit of protein kinase A regulates growth and differentiation in the fungus

Mucor circinelloides. Eukaryot Cell 8:933-944.

40. Zaremberg V, Donella-Deana A, Moreno S. 2000. Mechanism of activation of

cAMP-dependent protein kinase: in *Mucor rouxii* the apparent specific activity of the

cAMP-activated holoenzyme is different than that of its free catalytic subunit. Arch

Biochem Biophys 381:74-82.

41. Wolff AM, Appel KF, Petersen JB, Poulsen U, Arnau J. 2002. Identification and

analysis of genes involved in the control of dimorphism in *Mucor circinelloides* (syn.

racemosus). FEMS Yeast Res 2:203-13.

42.   Singh N, Aguado JM, Bonatti H, Forrest G, Gupta KL, Safdar N, John GT, Pursell

KJ, Munoz P, Patel R, Fortun J, Martin-Davila P, Philippe B, Philit F, Tabah A, Terzi N,

Chatelet V, Kusne S, Clark N, Blumberg E, Julia MB, Humar A, Houston S, Lass-Florl

C, Johnson L, Dubberke ER, Barron MA, Lortholary O. 2009. Zygomycosis in solid

organ transplant recipients: a prospective, matched case-control study to assess risks

for disease and outcome. J Infect Dis 200:1002-11.

43.   Lee SC, Li A, Calo S, Inoue M, Tonthat NK, Bain JM, Louw J, Shinohara ML,

Erwig LP, Schumacher MA, Ko DC, Heitman J. 2015. Calcineurin orchestrates

dimorphic transitions, antifungal drug responses and host-pathogen interactions of the

pathogenic mucoralean fungus *Mucor circinelloides*. Mol Microbiol 97:844-65.

44.   Siekierka JJ, Hung SH, Poe M, Lin CS, Sigal NH. 1989. A cytosolic binding

protein for the immunosuppressant FK506 has peptidyl-prolyl isomerase activity but is

distinct from cyclophilin. Nature 341:755-7.

45.   Liu J, Farmer JD, Jr., Lane WS, Friedman J, Weissman I, Schreiber SL. 1991.

Calcineurin is a common target of cyclophilin-cyclosporin A and FKBP-FK506

complexes. Cell 66:807-15.

46.   Juvvadi PR, Fox D, Bobay BG, Hoy MJ, Gobiel SM, Venters RA, Chang Z, Lin

JJ, Averette AF, Cole DC, Barrington BC, Wheaton JD, Ciofani M, Trzoss M, Li X, Lee

SC, Ying-Lien C, Mutz M, Spicer LD, Schumacher MA, Heitman J, Steinbach WJ. 2019.

Harnessing calcineurin-FK506-FKBP12 crystal structures from invasive fungal

pathogens to develop antifungal agents. Nat Commun 10:4275.

47.   Bastidas RJ, Shertz CA, Lee SC, Heitman J, Cardenas ME. 2012. Rapamycin

exerts antifungal activity in vitro and in vivo against *Mucor circinelloides* via FKBP12-

dependent inhibition of Tor. Eukaryot Cell 11:270-81.

48.   Cardenas ME, Muir RS, Breuder T, Heitman J. 1995. Targets of immunophilin-

immunosuppressant complexes are distinct highly conserved regions of calcineurin A.

Embo J 14:2772-83.

49.   Fox DS, Cruz MC, Sia RA, Ke H, Cox GM, Cardenas ME, Heitman J. 2001.

Calcineurin regulatory subunit is essential for virulence and mediates interactions with

FKBP12-FK506 in *Cryptococcus neoformans*. Mol Microbiol 39:835-49.

50.   Jin L, Harrison SC. 2002. Crystal structure of human calcineurin complexed with

cyclosporin A and human cyclophilin. Proc Natl Acad Sci U S A 99:13522-6.

51.   Calo S, Nicolas FE, Lee SC, Vila A, Cervantes M, Torres-Martinez S, Ruiz-

Vazquez RM, Cardenas ME, Heitman J. 2017. A non-canonical RNA degradation

pathway suppresses RNAi-dependent epimutations in the human fungal pathogen

Mucor circinelloides. PLoS Genet 13:e1006686.

52.   Calo S, Shertz-Wall C, Lee SC, Bastidas RJ, Nicolás FE, Granek JA,

Mieczkowski P, Torres-Martínez S, Ruiz-Vázquez RM, Cardenas ME, Heitman J. 2014.

Antifungal drug resistance evoked via RNAi-dependent epimutations. Nature 513:555.

53.   Garcia A, Adedoyin G, Heitman J, Lee SC. 2017. Construction of a recyclable

genetic marker and serial gene deletions in the human pathogenic Mucorales *Mucor*

circinelloides. G3: Genes, Genomes, Genetics 7:2047-2054.

54.   Lubbehusen TL, Nielsen J, McIntyre M. 2003. Morphology and physiology of the

dimorphic fungus *Mucor circinelloides* (syn. *M. racemosus*) during anaerobic growth.

Mycol Res 107:223-30.

55.   Chen Y, Cann MJ, Litvin TN, Iourgenko V, Sinclair ML, Levin LR, Buck J. 2000.

Soluble adenylyl cyclase as an evolutionarily conserved bicarbonate sensor. Science

289:625-8.

56.   Turnham RE, Scott JD. 2016. Protein kinase A catalytic subunit isoform

PRKACA; history, function and physiology. Gene 577:101-8.

57.   Donaton MC, Holsbeeks I, Lagatie O, Van Zeebroeck G, Crauwels M, Winderickx

J, Thevelein JM. 2003. The Gap1 general amino acid permease acts as an amino acid

sensor for activation of protein kinase A targets in the yeast *Saccharomyces cerevisiae*.

Mol Microbiol 50:911-29.

58.   Kraidlova L, Van Zeebroeck G, Van Dijck P, Sychrová H. 2011. The *Candida*

albicans GAP gene family encodes permeases involved in general and specific amino

acid uptake and sensing. Eukaryot Cell 10:1219-29.

59.   Ibrahim AS, Spellberg B, Walsh TJ, Kontoyiannis DP. 2012. Pathogenesis of

mucormycosis. Clin Infect Dis 54:S16-S22.

60.   Wang L, Pittman KJ, Barker JR, Salinas RE, Stanaway IB, Williams GD, Carroll

RJ, Balmat T, Ingham A, Gopalakrishnan AM, Gibbs KD, Antonia AL, Heitman J, Lee

SC, Jarvik GP, Denny JC, Horner SM, DeLong MR, Valdivia RH, Crosslin DR, Ko DC.

2018. An atlas of genetic variation linking pathogen-induced cellular traits to human

disease. Cell Host Microbe 24:308-323.e6.

61.   Juvvadi PR, Lamoth F, Steinbach WJ. 2014. Calcineurin as a multifunctional

regulator: unraveling novel functions in fungal stress responses, hyphal growth, drug

resistance, and pathogenesis. Fungal Biol Rev 28:56-69.

62.   Chow EW, Clancey SA, Billmyre RB, Averette AF, Granek JA, Mieczkowski P,

Cardenas ME, Heitman J. 2017. Elucidation of the calcineurin-Crz1 stress response

transcriptional network in the human fungal pathogen *Cryptococcus neoformans*. PLoS

Genet 13:e1006667.

63.   Egan JD, Garcia-Pedrajas MD, Andrews DL, Gold SE. 2009. Calcineurin is an

antagonist to PKA protein phosphorylation required for postmating filamentation and

virulence, while PP2A is required for viability in *Ustilago maydis*. Mol Plant Microbe

Interact 22:1293-301.

64.   Kafadar KA, Cyert MS. 2004. Integration of stress responses: modulation of

calcineurin signaling in *Saccharomyces cerevisiae* by protein kinase A. Eukaryot Cell

3:1147-53.

65.   Peeters T, Louwet W, Geladé R, Nauwelaers D, Thevelein JM, Versele M. 2006.

Kelch-repeat proteins interacting with the Gα protein Gpa2 bypass adenylate cyclase for

direct regulation of protein kinase A in yeast. PNAS 103:13034-13039.

66.   Frazzitta AE, Vora H, Price MS, Tenor JL, Betancourt-Quiroz M, Toffaletti DL,

Cheng N, Perfect JR. 2013. Nitrogen source-dependent capsule induction in human-

pathogenic *Cryptococcus* species. Eukaryot Cell 12:1439-1450.

67.   O'Meara TR, Norton D, Price MS, Hay C, Clements MF, Nichols CB, Alspaugh

JA. 2010. Interaction of *Cryptococcus neoformans* Rim101 and protein kinase A

regulates capsule. PLoS Pathog 6:e1000776.

68.   Alspaugh JA. 2016. Unveiling protein kinase A targets in *Cryptococcus*

neoformans capsule formation. mBio 7:e00021-16.

69.   Boyce KJ, Andrianopoulos A. 2015. Fungal dimorphism: the switch from hyphae

to yeast is a specialized morphogenetic adaptation allowing colonization of a host.

FEMS Microbiol Rev 39:797-811.

70.   Gauthier GM. 2017. Fungal dimorphism and virulence: molecular mechanisms

for temperature adaptation, immune evasion, and in vivo survival. Mediators Inflamm

2017:8491383.

71.   Yoo SY, Kwon SM. 2013. Angiogenesis and its therapeutic opportunities.

Mediators Inflamm 2013:127170.

72.   Vellanki S, Huh EY, Saville SP, Lee SC. 2019. *Candida albicans* morphology-

dependent host FGF-2 response as a potential therapeutic target. J Fungi 5:22.

73.   Ben-Ami R, Lewis RE, Leventakos K, Kontoyiannis DP. 2009. *Aspergillus*

fumigatus inhibits angiogenesis through the production of gliotoxin and other secondary

metabolites. Blood 114:5393-9.

74.   Lee SC, Billmyre RB, Li A, Carson S, Sykes SM, Huh EY, Mieczkowski P, Ko

DC, Cuomo CA, Heitman J. 2014. Analysis of a food-borne fungal pathogen outbreak:

Virulence and genome of a *Mucor circinelloides* isolate from yogurt. mBio 5:e01390-14.

75.   Mueller KD, Zhang H, Serrano CR, Billmyre RB, Huh EY, Wiemann P, Keller NP,

Wang Y, Heitman J, Lee SC. 2019. Gastrointestinal microbiota alteration induced by

Mucor circinelloides in a murine model. J Microbiol 57:509-520.

76.   Desbois AP, Coote PJ. 2012. Utility of greater wax moth larva (*Galleria*

mellonella) for evaluating the toxicity and efficacy of new antimicrobial agents. Adv Appl

Microbiol 78:25-53.

77.   O'Meara TR, Holmer SM, Selvig K, Dietrich F, Alspaugh JA. 2013. *Cryptococcus*

neoformans Rim101 is associated with cell wall remodeling and evasion of the host

immune responses. mBio 4:e00522-12.

78.   Corrochano LM, Kuo A, Marcet-Houben M, Polaino S, Salamov A, Villalobos-

Escobedo JM, Grimwood J, Alvarez MI, Avalos J, Bauer D, Benito EP, Benoit I, Burger

G, Camino LP, Canovas D, Cerda-Olmedo E, Cheng JF, Dominguez A, Elias M, Eslava

AP, Glaser F, Gutierrez G, Heitman J, Henrissat B, Iturriaga EA, Lang BF, Lavin JL, Lee

SC, Li W, Lindquist E, Lopez-Garcia S, Luque EM, Marcos AT, Martin J, McCluskey K,

Medina HR, Miralles-Duran A, Miyazaki A, Munoz-Torres E, Oguiza JA, Ohm RA,

Olmedo M, Orejas M, Ortiz-Castellanos L, Pisabarro AG, Rodriguez-Romero J, Ruiz-

Herrera J, Ruiz-Vazquez R, Sanz C, Schackwitz W, et al. 2016. Expansion of signal

transduction pathways in fungi by extensive genome duplication. Curr Biol 26:1577-

1584.

79.   Lopman B, Vennema H, Kohli E, Pothier P, Sanchez A, Negredo A, Buesa J,

Schreier E, Reacher M, Brown D, Gray J, Iturriza M, Gallimore C, Bottiger B, Hedlund

KO, Torven M, von Bonsdorff CH, Maunula L, Poljsak-Prijatelj M, Zimsek J, Reuter G,

Szucs G, Melegh B, Svennson L, van Duijnhoven Y, Koopmans M. 2004. Increase in

viral gastroenteritis outbreaks in Europe and epidemic spread of new norovirus variant.

Lancet 363:682-8.

80.   DePristo MA, Banks E, Poplin R, Garimella KV, Maguire JR, Hartl C, Philippakis

AA, del Angel G, Rivas MA, Hanna M, McKenna A, Fennell TJ, Kernytsky AM,

Sivachenko AY, Cibulskis K, Gabriel SB, Altshuler D, Daly MJ. 2011. A framework for

variation discovery and genotyping using next-generation DNA sequencing data. Nature

Genetics 43:491.

81.   Vellanki S, Navarro-Mendoza MI, Garcia A, Murcia L, Perez-Arques C, Garre V,

Nicolas FE, Lee SC. 2018. *Mucor circinelloides*: growth, maintenance, and genetic

manipulation. Curr Protoc Microbiol 49:e53.

82.   Watkins TN, Gebremariam T, Swidergall M, Shetty AC, Graf KT, Alqarihi A,

Alkhazraji S, Alsaadi AI, Edwards VL, Filler SG, Ibrahim AS, Bruno VM. 2018. Inhibition

of EGFR Signaling protects from mucormycosis. mBio 9:e01384-18.

83.   Luo G, Gebremariam T, Lee H, French SW, Wiederhold NP, Patterson TF, Filler

SG, Ibrahim AS. 2013. Efficacy of liposomal Amphotericin B and Posaconazole in

intratracheal models of murine mucormycosis. Antimicrob Agents Chemother 57:3340.

Figure Legends

Figure 1. Calcineurin suppressor mutations (CnSp) in the *cnbRΔ* mutant background restore hyphal growth. A) Growth of calcineurin suppressor mutants (CnSp1 to CnSp4) and *cnbRΔ* mutant on a YPD agar plate at 30°C for four days post inoculation. While the *cnbRΔ* mutant shows smaller yeast colonies, the CnSp mutants form larger hyphal colonies. B) *Mucor* WT (R7B), *cnbRΔ* mutant, or CnSp mutants were grown overnight in liquid YPD medium with shaking at 30°C. Micrographs show that the CnSp mutants exhibit hyphal growth like WT (scale bar = 10 μm).

Figure 2. Calcineurin suppressor mutations (CSR) in the *cnaBΔ* mutant background confer resistance to calcineurin inhibitors. A) Growth of cyclosporine resistant mutants (CSR) exhibiting hyphal growth and *cnaBΔ* mutant exhibiting yeast growth when incubated on YPD agar with CsA (100 μg/ml) or FK506 (1 μg/ml) for four days at 30°C. CSR1, CSR2, CSR3, and CSR4 are shown. The other CSR mutants exhibited a similar CsA resistance phenotype (data not shown).

Figure 3. *bycAΔ* and *bycAΔ cnbRΔ* double mutant is resistant to calcineurin inhibitors. A) After three days of growth on solid YPD agar the *cnbRΔ* mutant grows as yeast; however, despite no calcineurin function, the *bycAΔ cnbRΔ* double mutant exhibits hyphal growth like WT and the *bycAΔ* mutant. B) In the presence of CsA (upper; 100 μg/ml) the *bycAΔ cnbRΔ* double mutant and CnSp4 formed larger hyphal colonies compared to WT, indicating resistance to CsA. No major difference in colony

size was noted between *bycAΔ* mutant and WT. In the presence of FK506 (lower; 1

μg/ml) the WT forms a smaller yeast colony whereas the *bycAΔ* mutant, *bycAΔ cnbRΔ*

double mutant and CnSp4 each form a larger hyphal colony. C) When *Mucor* was

grown overnight in YPD medium containing FK506 (1 μg/ml) at 30°C with shaking, the

bycAΔ mutant, *bycAΔ cnbRΔ* double mutant, and CnSp4 mutant exhibited resistance to

FK506 as evident by larger biomass and hyphal morphology, whereas the WT is

sensitive to FK506 as they not only form less biomass but also grow as a yeast. As

expected, the *cnbRΔ* mutant remains in its yeast-locked form (scale bar = 20 μm).

Figure 4. BycA is involved in the *Mucor* hyphal-yeast transition under aerobic

conditions. A) Reverse transcriptase quantitative PCR showed that in the absence of

calcineurin function (*cnbRΔ* and WT + FK506; yeast morphology) *bycA* expression was

six-fold higher compared to WT (hyphal morphology) with calcineurin function

suggesting that calcineurin regulates *bycA* expression at the mRNA level. When *Mucor*

WT was grown anaerobically under high CO_2 conditions (see methods) it grows as a

yeast; however, there is no significant difference in *bycA* expression compared to WT-

hyphae group. One-way ANOVA was significant (P = 0.0001). Dunnett's post-hoc test

was used to compare *cnbRΔ*, WT + FK506, WT-yeast with WT-hyphae group (*P

<0.05). As expected, no expression of the *bycA* gene was detected in both of the *bycAΔ*

mutants. B) When WT, *cnbRΔ*, *bycAΔ*, and *bycAΔ cnbRΔ* mutants were grown

anaerobically overnight in high CO_2 conditions, they all exhibited a yeast morphology

thereby suggesting that neither calcineurin nor BycA has a role in anerobic

morphological pathways (scale bar = 20 μm). C) Crude protein extracts (0.5 µg) were

used to measure overall PKA activity, compared to WT-yeast (grown anaerobically) the

WT-hyphae and *bycA*Δ have significantly lower PKA activity. In WT + FK506 and *cnbR*Δ

mutant PKA activity remained higher, however *bycA*Δ *cnbR*Δ mutant has lower PKA

activity despite no calcineurin function. One-way ANOVA was significant (P <0.0001).

Dunnett's multiple comparison test was used to compare each group with WT-yeast (*P

<0.05). D) Crude extracts from 60 mg biomass were used to measure overall cAMP

activity, compared to WT-yeast, the WT + FK506 and *cnbR*Δ had significantly lower

cAMP activity thereby suggesting that under aerobic conditions, the PKA activity is

elevated in *Mucor* yeast in a cAMP-independent manner. One-way ANOVA was

significant (P <0.0001). Dunnett's multiple comparison test was used to compare each

group with WT-yeast (*P <0.05).

Figure 5. Phagosome maturation in macrophages containing *Mucor* is dependent

on pathogen calcineurin function and not morphology. A) 5 x 10^5 J774.A1 were

challenged with *Mucor* spores or yeast at a MOI of 1 along with LysoTracker™ Green

DND-26, and Hoechst 33342 stain (blue). White arrows indicate where phagosome

maturation should be observed in the field (scale bar = 5 μm). B) Compared to WT, the

macrophages containing *cnbR*Δ and *bycA*Δ *cnbR*Δ underwent significantly higher

phagosome maturation. Data is shown as percentage maturation. The number (N) of

macrophages containing *Mucor* counted for each group is: WT = 480; *bycA*Δ = 290;

*cnbR*Δ = 279; *bycA*Δ *cnbR*Δ = 413. One-way ANOVA was significant (P <0.0001).

Dunnett's multiple comparison test was used to compare each group with WT (*P

<0.05).

Figure 6. Calcineurin mutants cause less endothelial cell damage and FGF-2

protein secretion. 5 x 10^3 HUVECs were challenged with *Mucor* spores or yeast at a

MOI of 10. After 24 hours, the supernatant was collected. A) LDH levels were quantified

which indicate cytotoxicity. Compared to WT, calcineurin mutants cause less damage.

One-way ANOVA was significant (P <0.0001). Dunnett's multiple comparison test was

used to compare each group with WT (*P <0.05). B) FGF-2 levels were quantified using

ELISA. The WT induce significantly higher FGF-2 protein secretion when compared to

calcineurin mutants. One-way ANOVA was significant (P <0.0001). Dunnett's multiple

comparison test was used to compare each group with WT (*P <0.05).

Figure 7. Calcineurin mutants are less virulent in a *Galleria mellanolla* (wax moth)

model of mucormycosis. The wax moth larvae (n=15/group) were inoculated with 1 x

10^4 *Mucor* spores or yeast in 2 μl PBS via injection into the last left proleg and were

monitored for survival. All wax moth challenged with WT or *bycA*Δ succumbed to

mortality within a week. The *cnbR*Δ was avirulent, whereas about 65% of wax moth

inoculated with *bycA*Δ *cnbR*Δ or CnSp4 survived. A log-rank (Mantel-Cox) test was

statistically significant (p <0.0001). A pair-wise comparison was also performed to

compare the following groups: WT vs *bycA*Δ (p = 0.82); WT vs *cnbR*Δ or *bycA*Δ *cnbR*Δ

or CnSp4 (*p <0.0001).

Figure 8. Calcineurin, BycA, and PKA in the morphogenesis of *Mucor* in aerobic conditions. Calcineurin is the master regulator of *Mucor* morphology. Active calcineurin positively regulates hyphal growth and negatively regulates yeast growth. This is achieved by suppressing the expression of the *bycA* gene thereby preventing an increase in PKA activity. Calcium-calmodulin activates calcineurin to promote hyphal growth under aerobic conditions. However, when calcineurin is not functional, *bycA* gene expression is significantly elevated. BycA then activates PKA in a cAMP-independent manner to promote yeast growth. In anaerobic conditions with high CO_2 levels, PKA is activated through the CO_2 – cAMP pathway. Neither BycA nor calcineurin has a defined role in the regulation of *Mucor* morphology under anaerobic conditions. Can2: carbonyl anhydrase, AC: adenylyl cyclase, and CaM: calmodulin.

Supporting information

Figure S1. Emergence of hyphal sectors from *cnbR*Δ mutants, and isolation of calcineurin suppressor mutants. *cnbR*Δ mutants were plated on YPD agar for 5 or more days at 30°C. The hyphae emerging from the yeast colonies were transferred to fresh YPD agar to propagate further.

Figure S2. Emergence of hyphal sectors from *cnaB*Δ mutant grown on CsA containing medium. *cnaB*Δ mutants were grown in the presence of CsA (2 µg/ml) for 5 or more days at 30°C. This resulted in emergence of hyphal sectors from yeast colonies.

1020

Figure S3. Confirmation of disruption of *bycA* gene by junction PCR and ORF

spanning PCR. A) Illustration of the *bycAΔ::pyrG-dpl237* and *bycA* alleles with ~ 1 kb

upstream and downstream flanking sequence. P1 (SL3) and P4 (SL8) recognize

sequences outside the disruption cassette. P2 (SCL566) and P3 (SCL567) recognize

pyrG-dpl237. P5 (SL182) and P6 (SL183) recognize *bycA*. B) 5' end: P1 and P2 amplify

a 2,203 bp region, and 3' end: P3 and P4 amplify a 1,994 bp region in the *bycAΔ::pyrG-*

dpl237, the primer pairs does not produce a fragment in the WT. P5 and P6 amplify 690

bp region within the *bycA* gene, so no amplification is noted in the mutants. Image not to

scale.

1030

Figure S4. Confirmation of disruption of *cnbR* gene in the *bycAΔ* background by

junction PCR and ORF spanning PCR. A) Illustration of the *cnbRΔ::leuA* deletion in

the *bycAΔ::pyrG-dpl237* strain and *cnbR* alleles with ~ 1 kb upstream and downstream

flanking sequence. P1 (SL243) and P4 (SL244) recognize sequences outside the

disruption cassette. P2 (SL391) and P3 (SL392) are specific for *leuA*. P5 (SCL578) and

P6 (SCL579) are specific for *cnbR*. B) 5' end: P1 and P2 amplify a 2,388 bp region, and

3' end: P3 and P4 amplify a 1,620 bp region in the *cnbRΔ::leuA* region, the primer pair

does not produce a fragment in the WT. P1 and P4 produce a 5,711 bp product in the

mutant and 3,342 bp product in the WT. P5 and P6 amplify 170 bp region within the

cnbR gene, so no amplification is noted in the mutants. Image not to scale.

1041

Figure S5. Confirmation of disruption of *cnbR* gene in the *bycA∆* background by RT-qPCR. Primers SCL578 and SCL579 were used to perform qPCR on the cDNA obtained from WT and mutants (see methods). Actin (SCL368 and SCL369) served as a control. No expression for *cnbR* was detected in the *bycA∆ cnbR∆* double mutant.

Figure S6. Predicted structure of BycA showing ten transmembrane domains, and one cytosolic domain. BycA is predicted to have 10 transmembrane domains, and one cytosolic domain. The transmembrane domains were predicted with the TMHMM 2.0 software package (http://www.cbs.dtu.dk/services/TMHMM/). The predicted cytosolic region contains a pectinesterase domain.

Figure S7. *bycA∆ cnbR∆* double mutant is lethal in a wax moth larva host only at higher inoculum. The wax moth larvae (n=15/group) were inoculated with 1×10^4 (1X) to 3×10^4 (3X) *bycA∆ cnbR∆* double mutant or WT spores (1X) in 2 µl PBS via the last left proleg and were monitored for survival. All wax moth challenged with WT or *bycA∆ cnbR∆* (3X) succumbed to mortality within a week. 65% of wax moths inoculated with *bycA∆ cnbR∆* (1X) and 45% of the wax moths inoculated with *bycA∆ cnbR∆* (2X) survived. A log-rank (Mantel-Cox) test was statistically significant ($p < 0.0001$). A pair-wise comparison was also performed to compare the following groups: WT vs *bycA∆ cnbR∆* (1X) (*$p < 0.0001$); WT vs *bycA∆ cnbR∆* (2X) (*$p = 0.0014$); WT vs *bycA∆ cnbR∆* (3X) ($p = 0.9460$).

Figure S8. *Mucor* **virulence in the murine model of systemic and pulmonary**

mucormycosis. A) Cyclophosphamide and cortisone acetate treated CF1 mice were

challenged with 1.5 x 10^6 spores or yeast cells in PBS via lateral tail vein (systemic

mucormycosis model). The animals (n = 5/group) were monitored for survival. A log-

rank (Mantel-Cox) test was not statistically significant (p = 0.0585) suggesting no major

difference in survival between WT and calcineurin mutant groups. B) For pulmonary

mucormycosis model (n = 5/group), the same inocula were introduced via intratracheal

route and monitored for survival. A log-rank (Mantel-Cox) test was not statistically

significant (p = 0.1188).

Figure S9. Survival of wild-spores and *cnbRΔ* yeast during co-cultures with bone

marrow macrophages. The spores survive the macrophages significantly higher

compared to the yeast-locked mutant. The experiment was performed as described

previously (Jung et al. 2019).

1.   **Jung, K.-W., Y. Lee, E. Y. Huh, S. C. Lee, S. Lim, and Y.-S. Bahn.** 2019.

     Rad53- and Chk1-dependent DNA damage response pathways cooperatively

     promote fungal pathogenesis and modulate antifungal drug susceptibility. mBio

     10:e01726-18.

1084   **Table S1** A) List of strains used in the study and B) List of primers used in the study.

1085

1086

1087

1088

1089

1090

1091

1092

1093

1094

1095

1096

1097

1098

1099

1100

1101

1102   **Table 1.** Characterization of the bypass mutants isolated in the *cnbRΔ* mutant

1103   background.

Strain	Genotype in the *bycA* allele	Remark	Type of mutation
CnSp1	1350G>A		nonsense
CnSp2	long deletion around *bycA*	whole genome sequenced	
CnSp3	2030C>A	whole genome sequenced	nonsense
CnSp4	361_1579del	whole genome sequenced	frameshift
CnSp5	potential deletion	no PCR product of *bycA*	
CnSp6	unknown		
CnSp7	995C>T	whole genome sequenced	missense
CnSp8	1921_1922insTGACATTGCTTCAG CAG	whole genome sequenced	frameshift
CnSp9	320T>C		missense
CnSp10	2319_2320insC	whole genome sequenced	frameshift
CnSp11	2129_2130insTCACC		frameshift
CnSp12	potential deletion	no PCR product of *bycA*	
CnSp13	1784_1793delTCAATTTCAT		frameshift
CnSp14	566T>A		nonsense
CnSp15	2238_2239insTC		frameshift
CnSp16	unknown		
CnSp17	1732C>T		nonsense

1104

1105

1106

1107

1108

1109

1110   **Table 2.** Characterization of the calcineurin bypass mutants isolated in the *cnaBΔ*

1111   mutant background.

1112

Strain	Genotype in the *bycA* allele	Remark	Type of mutation
CSR1	C1240del		frameshift
CSR2	potential deletion	no PCR product of *bycA*	
CSR3	733 C>A		missense
CSR4	633delG	whole genome sequenced	frameshift
CSR5	407C>T	whole genome sequenced	missense
CSR6	2188delC	whole genome sequenced	frameshift
CSR7	806_825delCCTACCCGCCAGTGGAAGCA		frameshift
CSR8	1711delG		frameshift
CSR9	2188delC	whole genome sequenced	frameshift
CSR10	1649delT	whole genome sequenced	frameshift
CSR11	578G>A		nonsense
CSR12	621delT		frameshift
CSR13	2296_2297insAAT		frameshift
CSR14	218G>A		nonsense
CSR16	2263delC	whole genome sequenced	frameshift
CSR28	215G>A		missense
CSR29	unknown		
CSR35	unknown		
CSR38	unknown		

1113

1114

1115

1116

1117

1118

1119

1120

1121 **Table 3**. Summary for *cnbRΔ* and *bycAΔ cnbRΔ* double mutant growth in the presence

1122 of Methionine, Threonine, and Arginine as sole nitrogen source.

Ammonium source	cnbRΔ mutant growth [a]	bycAΔ cnbRΔ mutant growth [a]
None	-	-
Methionine	+	-
Threonine	+	-
Arginine	+	-
Casamino acids	++	++

1123   [a] *+ or ++ represent the extent of growth.*

1124

1125

Figure 1

Figure 2

YPD + CsA

YPD + FK506

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8