1 Running title: A cargo-regulated transition point in endocytosis
2
3

4 **Cargo licenses maturation of stalled endocytic sites through a**
5 **regulatory checkpoint**
6
7

8 Ross TA Pedersen[1,3], Julian E Hassinger[1,2,3], Paul Marchando[1], David G Drubin[1*]
9
10 [1]Department of Molecular and Cell Biology and
11 [2]Biophysics Graduate Group, University of California, Berkeley, Berkeley, CA 94720
12 [3]These authors contributed equally to this work
13 *Correspondence: drubin@berkeley.edu

Abstract

During Clathrin-mediated endocytosis (CME), over 50 different proteins assemble on the

plasma membrane to reshape it into a cargo-laden vesicle. Studies of *Saccharomyces*

cerevisiae have traced a detailed timeline of assembly of the CME machinery. Assembly of

this machinery is subject to regulation to ensure cargo capture; however, which CME steps

are regulated, and which steps are deterministic, is unknown. In this study, we

quantitatively and spatially describe progression through the CME pathway. We pinpoint a

regulatory transition point that governs progression from the initiation phase of CME to

the internalization phase. Availability of endocytic cargo influences maturation through

this transition point, indicating that it is a cargo checkpoint. While previous studies in yeast

and mammalian cells suggested that cargo ensures its own internalization through

regulating CME initiation rates or frequency of abortive events, our data instead identify

maturation through a checkpoint in the pathway as the cargo-sensitive step.

1 Introduction

Membrane trafficking processes require coordinated cargo capture and membrane

reshaping. One such process is clathrin-mediated endocytosis (CME), which is the major

trafficking route into the cell (Bitsikas et al., 2014). Numerous studies have suggested that

cargo influences the rate of CME by modulating the proportion of endocytic sites that

continue to completion as opposed to aborting precociously (Ehrlich et al., 2004; Liu et al.,

2010; Loerke et al., 2009; Mettlen et al., 2010). However, this proposed mode of

coordinating cargo capture with endocytosis was called into question when studies of

genome-edited mammalian cells did not reveal widespread abortive CME events (Doyon et

al., 2011; Hong et al., 2015). Thus, how the CME pathway is regulated and which steps are

subject to regulation remain unresolved.

The budding yeast *Saccharomyces cerevisiae* is an ideal organism for systematic

studies of the CME pathway. Aside from practical advantages such as simplicity, highly

tractable genetics, and accessible genome editing, *S. cerevisiae* has a remarkably regular

CME pathway that invariably proceeds to completion (Kaksonen et al., 2005, 2003). Over

50 proteins are recruited to every CME site in budding yeast with a nearly invariant order

and with fairly regular kinetics (Lu et al., 2016).

Budding yeast CME proteins have been organized conceptually into functional

modules based on shared timing of recruitment and dynamic behavior, common functions,

and spatial organization at CME sites (Figure 1A, Carroll et al., 2012; Kaksonen et al., 2005;

Lu et al., 2016). Briefly, proteins of the early and early coat modules initiate CME sites

(Godlee and Kaksonen, 2013; Lu and Drubin, 2017; Stimpson et al., 2009). Next,

intermediate coat proteins, which bind to actin filaments and couple actin growth to

membrane invagination, are recruited (Carroll et al., 2012; Skruzny et al., 2012). Late coat

proteins arrive and recruit actin assembly machinery consisting of proteins of the

WASP/Myosin complex (Bradford et al., 2015; Sun et al., 2015). The WASP/myosin

complex orchestrates a critical burst of actin assembly and myosin activity to bend the

plasma membrane into a pit (Lewellyn et al., 2015; Manenschijn et al., 2019; Pedersen and

Drubin, 2019; Sun et al., 2006). While the early coat, intermediate coat, and late coat

proteins internalize with the plasma membrane, the early module proteins dissociate from

CME sites at the time of actin assembly (Carroll et al., 2012). Actin filaments recruit a host

of actin binding proteins and disassembly factors collectively referred to as the actin

module. At the same time, proteins of the scission module arrive to release the nascent

vesicle. The timing and modular organization of this process are conserved from yeast to

mammals (Taylor et al., 2011).

Despite this detailed knowledge of the CME machinery, how key transitions in the

process are triggered remains poorly understood (Figure 1A). The budding yeast early and

early coat proteins persist at the CME site for a variable period of time before recruitment

of the late arriving coat components, which are reported to have more regular lifetimes

(Carroll et al., 2012; Newpher et al., 2005; Peng et al., 2015a; Stimpson et al., 2009). A

previous study provided preliminary evidence consistent with cargo regulating the

transition from the "variable phase" to the "regular phase", but this model was not

thoroughly tested, the point in the pathway that is regulated was not identified, and the

physiological implications of the model were not examined (Carroll et al., 2012).

We set out to determine in budding yeast by systematic live-cell imaging analyses

(1) which steps in the CME pathway are subject to regulation and which steps are

deterministic, and (2) how cargo influences progression through the CME pathway to ensure its own timely internalization. Our data quantitatively separate the CME pathway into an early phase and a late phase. The behavior of proteins in each phase predicts the behavior of others in the same phase, but not the behavior of proteins in the other phase. Further experiments lead us to propose that the presence of cargo accelerates the transition from the early phase to the late phase, so this regulatory transition is likely a cargo checkpoint. We propose that endocytic sites are initiated widely over the yeast cell surface but mature to the late, internalization phase more readily in regions rich in cargo.

 Results

2 **Systematic live-cell imaging of clathrin-mediated endocytosis reveals inherent**
3 **variability in timing and abundance of protein recruitment**
4
We took a systematic imaging approach to quantitatively describe progression though the

CME pathway. While each of the proteins of the CME pathway we examined had been

imaged previously to determine a recruitment order, we aimed to perform a deeper

quantitative analysis to uncover previously undetected features and relationships among

the proteins. Recruitment order and timing in the budding yeast CME pathway were

previously determined by many pair-wise imaging iterations using two-color fluorescence

microscopy. Here we imaged CME protein recruitment timing relative to three reference

proteins, providing an orthogonal approach to determining the recruitment order and

relative timing.

We acquired live-cell two-color total internal reflection fluorescence microscopy

(TIRFM) movies of budding yeast strains expressing nearly two-dozen different fluorescent

fusion proteins (Supplementary Table 1). Each yeast strain encoded an endocytic query

protein tagged with GFP and a reference protein for various stages of progression through

CME tagged with a red fluorophore. All proteins were endogenously tagged and expressed

from their own promoters in the genome to avoid overexpression artifacts. We chose the

Eps15 homologue Ede1, a key component of the early endocytic module (Carroll et al.,

2012; Lu and Drubin, 2017; Stimpson et al., 2009), as a reference protein to report on CME

site initiation. Sla1, a late-arriving endocytic coat protein (Kaksonen et al., 2003), was used

as a reference protein to report on the transition from the variable stage of CME to the

regular stage. Finally, we chose the actin binding protein Abp1, a reporter of endocytic

actin networks in budding yeast (Kaksonen et al., 2003), to report on initiation of actin

assembly. All of these reference proteins are relatively abundant at endocytic sites (and

hence bright). We chose TIRFM in lieu of the more standard imaging modality of widefield

microscopy in a medial focal plane of yeast cells (Kaksonen et al., 2005, 2003). TIRFM has a

superior signal-to-noise ratio as well as minimal photobleaching, although this comes at

the cost of losing the ability to track the endocytic proteins once they move away from the

plasma membrane during internalization. The benefits of TIRFM were especially important

for faithfully detecting and tracking the early endocytic proteins, because they produce a

dim, long-lived signal at CME sites. By imaging different GFP-tagged "query" proteins in

two-color movies with common reference markers, we were able to compare the dynamics

of a large subset of the presently known CME proteins at endocytic sites.

To chronicle progression through the CME pathway, we chose GFP-tagged query

proteins from each endocytic module. We imaged the early module proteins Syp1 (an FCho

homologue) and Hrr25 (a casein kinase homologue) and the early coat proteins Pal1 and

Apl1 (a component of the AP-2 complex) with reference to the early protein Ede1 to

capture the earliest stages of CME. We also imaged the early module proteins Ede1, Syp1,

and Hrr25 and the early coat proteins Pal1 and Apl1 with reference to the late coat protein

Sla1 to monitor the transition from the early stage of CME to the late stage. Finally, we

imaged a large sample of CME proteins with reference to Abp1 to monitor the late stage of

CME and the initiation of actin assembly in particular. We used Ede1 and Syp1 as

representative early proteins; Sla2 (human homologue: Hip1R) as a representative

intermediate coat protein; Pan1 (intersectin) and Sla1 as representative late coat proteins;

Las17 (Wiscott-aldrich syndrome protein, WASP), Bzz1 (toca1), Vrp1 (WASP interacting

protein, WIP), and Myo5 (myosin 1e) as representative WASP/Myosin module proteins;

Arc15 (an Arp2/3 complex subunit), Sac6 (fimbrin), and Ark1 (GAK/AAK) as

representative actin module proteins; and Rvs167 (amphiphysin) as a representative

scission module protein. Movies of these query proteins yielded new information about

every stage of the CME pathway.

A combination of established and custom MATLAB scripts was used to monitor

fluorescently tagged proteins at CME sites. CME sites are smaller than the diffraction limit

of fluorescence microscopy, so they appear as ~200 nm diameter spots in our movies.

Figure 1B depicts a representative montage of an endocytic site. In this case, the GFP-

tagged query protein Las17, the yeast WASP, arrives before the reference protein Abp1-

RFP. Both proteins then disappear as the endocytic pit internalizes and leaves the TIRFM

field (See also Video 1). We used the cmeAnalysis MATLAB package to automatically track

all of the endocytic sites in many fields of yeast cells (Aguet et al., 2013). We then used

custom MATLAB scripts to: identify and associate colocalized green and red spots, reject

CME sites that were too close to one another to resolve independently, and extract

background-subtracted fluorescence intensities for each fluorescence channel at each site

(See Materials and Methods for full details). Using our software, we tracked individual

endocytic sites from their initiation to their disappearance and extracted time-resolved

fluorescence intensity data for each channel (Figure 1B-C).

Our systematic imaging data set independently supports the previously proposed

recruitment order for CME proteins while providing important additional mechanistic

insights. We generated plots of fluorescence intensity vs. time for each fluorescently-tagged

protein at each tracked CME site and recorded lifetimes, maximum fluorescence intensities,

and the delay between query protein arrival and reference protein arrival (Figure 1C).

When we ordered the proteins using our time to reference protein arrival data, we recapitulated the previously deduced recruitment order (Figure 1D – F, Supplementary Tables 2, 3, and 4). First, our plots show that all early arriving proteins (Ede1, Syp1, Pal1, Hrr25, and Apl1) arrive at CME sites at about the same time when imaged with reference to Ede1-RFP (Figure 1D, Carroll et al., 2012). When imaged with reference to Sla1-mCherry, Ede1 appears to be the earliest arriving protein, although this is likely due to Ede1 being the brightest of the group (Figure 1E, also see Figure 1 Supplement 1A). Our dataset using Abp1 as a reference protein recapitulates the canonical timeline of the CME pathway, an ordering which was previously established only through painstaking sequential rounds of two-color imaging by several groups across many studies (Figure 1F, see Lu et al., 2016 for review). Even fine-grained details are borne out in these plots. For example, the actin assembly factor Arc15-GFP (a component of the Arp2/3 complex) precedes the actin-binding protein Sac6-GFP at CME sites, and Sac6-GFP is followed closely by Ark1-GFP, a kinase recruited to CME sites by actin binding proteins (Figure 1F, Cope et al., 1999). Our ability to recapitulate fine-grained differences in timing gave us confidence to draw new conclusions from our dataset.

Maximum intensity and lifetime measurements also recapitulate trends from the literature. While we did not calibrate fluorescence intensity to count molecules, relative fluorescence intensities we measured are in agreement with those reported previously (Figure 1 Supplement 1A, Supplementary Tables 5, 6, and 7; Picco et al., 2015; Sun et al., 2019). The median lifetimes we measured are also in general agreement with published results (Figure 1 Supplement 1B, Supplementary Tables 8, 9, and 10, Carroll et al., 2012; Kaksonen et al., 2005, 2003; Lu et al., 2016; Peng et al., 2015a; Stimpson et al., 2009). The

instances in which our data deviate slightly from the literature are likely due to the

exponential decay of TIRFM illumination with distance from the coverslip (Axelrod, 1989).

We undercounted Arc15 slightly, most likely due to the actin network extending further

from the coverslip than other endocytic components we imaged. Our recorded lifetimes are

also systematically slightly lower than those previously reported, likely due to a failure to

record the final moments of each CME event when the invagination/vesicle is moving out of

the TIRFM field. Despite these minor artifacts, trends within our systematic TIRFM dataset

agree with the published literature, empowering our deeper analysis, described below.

One unexpected result from our initial analysis of CME protein dynamics is the

amount of variability in our measurements (Figure 1D-F, Figure 1 Supplement 1A-B).

Budding yeast CME has canonically been described as being highly regular, particularly

during the late stages (Kaksonen et al., 2006; Stimpson et al., 2009; Sun et al., 2015).

However, our dataset made us question this characterization. One measure of variability is

the coefficient of variation (CV), the ratio of the standard deviation to the mean (Reed et al.,

2002). Our data confirm that the earlier stages of CME (CV: ~0.5) are more variable then

the later stages, but we nevertheless detected considerable variability during the later

stages of the process (CV: ~0.3, See Supplementary Tables 8, 9, and 10). Despite repeated

characterization of CME as being "highly regular" in the literature, our CV values appear

consistent with previously reported data (Kaksonen et al., 2005, 2003; Sun et al., 2006,

where CV values can be estimated from plots of mean with standard deviation). We

exploited this previously underappreciated variability to gain new systems-level insights

into the mechanisms of CME.

23
24

Quantitative evidence for a regulatory transition point in the clathrin-mediated endocytosis pathway

Analysis of paired lifetime and intensity data for individual CME events (as opposed to ensemble averages from many events) revealed at least two distinct behaviors for proteins in the CME pathway. Given the highly interconnected nature of protein interactions at CME sites (Holland and Johnson, 2018; Tonikian et al., 2009), one might predict that longer assembly times at endocytic sites would lead to greater abundances. To test this prediction, we plotted the maximum intensity from each tracked endocytic event for each query protein as a function of its lifetime (Figure 2, Figure 2 Supplement 1), and performed linear regression analyses. The R^2 value of each fit can be interpreted as the percentage of the variability in maximum intensity that can be explained by the variation in lifetime (Glantz et al., 2001). For early endocytic proteins, exemplified by Ede1, lifetime is poorly correlated with abundance, with linear fits yielding low slopes (0.29) and R^2 values (0.05) (Figure 2A, Supplementary Table 11). Late arriving coat proteins and actin associated proteins, exemplified by Las17 and Sac6, show moderate correlations between lifetime and abundance, with higher slopes (0.97 and 0.91 respectively) and better fits (R^2 = 0.21 and 0.1, respectively) (Figure 2B-C, Supplementary Table 11). Finally, proteins involved in scission and disassembly of the endocytic machinery once again have abundances that are poorly predicted by residence times (Figure 2D, Supplementary Table 11). To summarize the results from our analysis of each CME protein imaged, we generated bar graphs with the height of the bars representing the slope of the fit and the color of the bars representing the R^2 value (Figure 2D, see Figure 2 Supplement 1 for individual plots, Supplementary Table 11 for fit statistics). From these plots, two distinct behaviors for CME proteins emerge. Early arriving proteins and very late arriving proteins have abundances that are

largely independent of their lifetimes, while components of the endocytic coat and WASP/Myosin module show a stronger correlation between these parameters. Such temporally resolved differences in behavior hint that a shift in CME protein behavior may occur sometime after endocytic site initiation but prior to actin assembly.

We next investigated interrelationships between the behaviors of different CME proteins. Many previous studies have presented two-color imaging of CME proteins (Kaksonen et al., 2005, 2003). However, previous analyses have typically relied on separating and averaging measurements of the two protein species imaged, so correlations between the behaviors of the two proteins at individual CME events imaged were lost (e.g., Kaksonen et al., 2005).

Since some CME proteins have lifetimes that are correlated with their own maximum intensities, and because CME proteins are extensively interconnected through protein-protein interactions, we wondered whether the lifetimes of any of our query proteins would be predictive of the abundances of our reference proteins. When we plotted the maximum intensity of our reference proteins at each CME event as a function of the lifetime of the corresponding query protein, we once again detected a quantitative transition point within the CME pathway (Figure 3, Figure 3 Supplement 1). Early proteins such as Ede1 have lifetimes that are poorly correlated with the maximum abundance of the CME actin marker Abp1 (Slope = -0.06, R^2 = 0.00, Figure 3A, Supplementary table 12). In contrast, lifetimes of late arriving coat proteins, WASP/Myosin module proteins, and actin binding proteins, exemplified by Las17 and Sac6, are moderately predictive of maximum Abp1 levels (Slopes = 0.96 and 0.70 respectively, R^2 = 0.13 and 0.07 respectively, Figure 3B-C, Supplementary Table 12). Finally, lifetimes of proteins involved in scission and

disassembly are once again relatively less correlated with maximum Abp1 intensity (Figure

3D, Supplementary Table 12). We again summarized the data from our linear fits by

making bar graphs as in Figure 2D (Figure 3D, see Figure 3 Supplement 1 for individual

plots, Supplementary Table 12 for fit statistics). While the lifetimes of the earliest and latest

arriving CME proteins are poorly correlated with the abundance of the actin marker Abp1,

all proteins of the late coat and WASP/Myosin CME modules have lifetimes that are

comparatively more positively correlated with Abp1 abundance (Figure 3D). We

interpreted a correlation between query protein behavior and reference protein behavior

to indicate shared protein-protein interactions. These data therefore show that a large

fraction of the proteins in the CME pathway can be thought of as an extensively

interconnected network, but the earliest and latest arriving CME proteins may be less well

connected into this network.

To determine how behaviors of individual CME proteins are correlated with

progression through the pathway as a whole, we tested whether lifetimes of any of our

query proteins are predictive of the lifetimes of our reference proteins. Early proteins, once

again exemplified by Ede1, have lifetimes that are well correlated with other early protein

lifetimes (slopes ranging from 0.45 – 0.71, R^2 values ranging from 0.21 – 0.55), but poorly

correlated with the lifetimes of the late arriving proteins Sla1 and Abp1 (slopes of 0.17 and

0.08 respectively, R^2 values of 0.06 and 0.03 respectively, Figure 4A, Supplementary Tables

13-15). Late arriving coat proteins and proteins of the WASP/Myosin module have

lifetimes that are moderately positively correlated to the lifetime of actin assembly as

reported by Abp1 (For Las17, a slope of 0.4 and an R^2 value of 0.24, Figure 4B,

Supplementary Table 15). Actin module proteins, such as Sac6, have lifetimes that are well

correlated with the lifetime of the actin marker Abp1 (slope = 0.83, R^2 = 0.71, Figure 4C,

Supplementary Table 15). All queried early arriving proteins have lifetimes that are well

correlated with the lifetime of the early arriving reference protein Ede1 (Figure 4D, Figure

4 Supplement 1, Supplementary Table 13), suggesting that a signal or a stochastic event,

the nature of which is currently unknown, may be required to trigger concerted maturation

of the early site. However, long lifetimes of early arriving CME proteins are not predictive

of the lifetime of any of the later-arriving reference proteins that we tested; our queried

early CME proteins have lifetimes that are poorly correlated with both Sla1 and Abp1

lifetimes (Figure 4E-F, Figure 4 Supplements 2-3, Supplementary Tables 14-15). Thus,

while behavior of any one early arriving CME protein is predictive of the behavior of the

others, the early proteins are molecularly insulated from the proteins in the late part of the

CME pathway.

Together, these data indicate that the CME pathway in budding yeast can be thought

of as occurring in two, quantitatively recognizable phases. Early arriving proteins initiate

CME sites and persist for a variable period of time, but the length of this variable phase is

not predictive of events that occur downstream in the pathway. Around the time of Sla2

recruitment, the CME site transitions into a more regular phase. The behavior of any one

protein predicts the behavior of other proteins in this phase, although there is still

variability in the behavior of proteins. Sla2 has been classified as an intermediate coat

protein (Carroll et al., 2012), and it is indeed "intermediate" by nearly every measure in our

data set. Its lifetime is less variable than those of the early proteins, but more variable than

the lifetimes of the late phase proteins (Early protein CV ~0.5, Late protein CV ~ 0.3, Sla2

CV ~0.4. Supplemental Tables 8, 9, and 10). The lifetime of Sla2 is also intermediately

correlated with its own abundance, the abundance of Abp1, and the lifetime of Abp1 (Figure 2F, Figure 3F, and Figure 4F). Sla2 may therefore be involved in transitioning from the early phase of CME into the late phase. Having substantiated the existence of this transition point based on an unbiased quantitative imaging approach, we decided to investigate molecular determinants that dictate the rate of maturation from the early phase into the late phase.

Maturation through the CME transition point is faster in cellular regions with concentrated endocytic cargo

Cargo has previously been suggested to influence the maturation rate of individual CME sites (Carroll et al., 2012; Layton et al., 2011). We set out to determine whether cargo influences maturation through tuning the rate of transition from the early phase of CME to the late phase.

As an initial strategy for varying the presence of cargo at CME sites, we took advantage of polarized secretion during the budding yeast cell cycle. When mitotically replicating budding yeast are in the small to medium-budded cell cycle stages, exocytic events are localized primarily to the growing bud, whereas exocytic events are redirected toward either side of the bud neck at cytokinesis (Lew and Reed, 1995). Endocytic cargos such as vesicle-associated SNARE proteins must be retrieved from the plasma membrane by CME following exocytosis (Lewis et al., 2000). Because septin filaments at the bud neck prevent lateral diffusion between the mother and the bud (Takizawa et al., 2000), the polarized pattern of secretion in small-budded cells concentrates CME cargo in the bud relative to the mother.

To determine whether the presence of endocytic cargo influences the rate of

transition from the early phase of CME to the late phase, we imaged a representative

protein from each respective endocytic phase through the cell cycle. We chose Ede1-GFP as

a representative marker of the early phase of CME and Sla1-mCherry as a representative

marker of the late phase. During the small- and medium-budded stages of the cell cycle,

when cargo is plentiful in the bud but sparse in the mother, we observed considerable

polarization of Sla1-mCherry-marked late CME sites without coincident polarization of

Ede1-GFP-marked early CME sites (Figure 5A, Video 2). The disparity in polarization

diminishes as cells near cytokinesis, when cargoes are more evenly distributed between

the mother and bud due to exocytic events occurring on either side of the contractile ring

(Figure 5A, Video 2). As cells begin the next cell cycle, Sla1-marked late phase CME sites

again become polarized (Figure 5A, Video 2). Based on the observation that early CME sites

are depolarized while late CME sites are polarized within the same cell, we concluded that

the presence of cargo does not affect the rate of CME site initiation appreciably, but it does

increase the proportion of initiated CME sites that mature.

The abundance of early CME sites relative to late CME sites in the mother cells of

polarized budding yeast could be explained by the presence of either abortive or stalled

CME events in the mother cells. The "extra" Ede1-GFP-marked early sites that we observed

could either abort, leading to the dearth of Sla1-marked late CME sites in the mother cells,

or they could mature at a slower rate, also resulting in fewer late phase sites. To distinguish

between these possibilities, we used a two-step imaging regime to first identify highly

polarized and depolarized cells, and to then track the fate of early CME sites in each cell

type. First, we collected 5 μm z-stacks (11 slices, 500 nm each) to identify highly polarized

cells, defined as small-budded cells with 3 or fewer Sla1-mCherry-marked late CME sites in

the mother, and depolarized cells, defined as medium- or large-budded cells with greater

than 3 Sla1-mCherry-marked late CME sites in the mother (Figure 5B, left photos). Next, we

collected time series in the medial focal plane of the same cells for 4 minutes with an

interval of 2 seconds between frames and generated circumferential kymographs along the

mother cortex (Figure 5B, right photos, Video 3). Our imaging protocol allowed us to

confidently identify highly polarized and depolarized cells and to visualize CME at high

temporal resolution in the cells identified.

When we analyzed the fates of individual CME sites in polarized and depolarized

cells, we determined that the rate of maturation through the transition point is accelerated

by proximity to exocytic sites and/or the cell polarity machinery. Quantification of the ratio

of the number of late CME sites to early sites in mothers of highly polarized and

depolarized cells confirms the observation that there are significantly more Ede1-GFP-

marked early CME sites per Sla1-mCherry-marked late site in mothers of small-budded

cells (median ratio of 0.08 vs. 0.44, Figure 5B left photos, Figure 5C). Tracking the fate of

every early CME site in the medial focal plane by generating circumferential kymographs

failed to reveal any abortive CME events, but it revealed many persistent early sites (Figure

5B right photos, Figure 5D, Video 3). While we could not reliably extract Ede1 lifetimes due

to a large fraction of Ede1 puncta being present at the start and end of our movies, we

observed significantly more long-lived Ede1-marked early CME sites in kymographs from

mothers in highly polarized cells than from mothers in depolarized cells (21.3% persistent

sites compared to 0.3%, 58.7% "partial" sites compared to 34.9%, Figure 5D, see figure

legend). This observation suggests that the maturation rate from the early phase of CME to

the late phase is slower when a CME site is further from cargo and the cell polarity machinery. When we conducted the same kinds of experiments with two late-phase CME markers, Sla1-GFP and Abp1-RFP, neither the ratio of numbers of puncta nor the lifetimes of these CME markers were sensitive to polarization state (median ratios 0.56 compared to 0.63, Figure 5 Supplement 1A, B, and C). While Abp1 lifetimes differ slightly (by 0.5 s, $p = 0.0247$), the small magnitude of this difference makes it unlikely to be biologically relevant. Based on these observations, we concluded that proximity to cargo and/or polarity machinery accelerates the rate of maturation from the early phase of CME to the late phase, but does not affect the kinetics of late events in CME.

The availability of cargo, rather than cell cycle stage, dictates maturation rate through the regulatory transition point

To determine whether the differences we observed in CME site maturation rate were due to polarization state or cell cycle stage, we used osmotic shock to reposition CME cargos and cell polarity proteins independent of cell cycle stage. A previous study from our lab used a temperature-sensitive secretion mutant to halt accumulation of endocytic cargo on the plasma membrane and reported increased numbers of long-lived Ede1-GFP-marked CME sites (Carroll et al., 2012). We conducted the converse experiment by redistributing endocytic cargo from the plasma membrane of buds in highly polarized cells to the plasma membrane of mother cells, testing whether long-lived early phase CME sites could be induced to mature simply though loss of cell polarity and redirection of cargo. Diluting *S. cerevisiae* cells out of media into water has previously been reported to trigger depolarization (Pringle et al., 1989). We found that osmotically shocking highly polarized

cells by rapidly diluting them with water caused dispersal of the polarity marker Bem1-

GFP (Figure 6A-B). Depolarization also results in redistribution of GFP-Sec4-marked

secretory vesicles from a bud-localized distribution to a more uniform one, suggesting that

exocytic traffic (and therefore endocytic cargo) is repositioned by this manipulation

(Figure 6 Supplement 1A). We therefore used osmotic shock to acutely depolarize cells

without influencing the cell cycle.

Forced depolarization through osmotic shock accelerates maturation through the

transition point in mothers of small-budded cells. When we examined the CME markers

Ede1-GFP and Sla1-mCherry in small-budded mothers before and after osmotic shock, we

found that the treatment caused Sla1-mCherry sites to depolarize. Ede1-GFP sites were not

noticeably affected (Figure 6C). Quantification of the ratio of the number of late phase CME

sites to early phase CME sites in these cells reveals a change in ratio reminiscent of the one

that occurs naturally during the cell cycle (median ratio changes from 0.09 to 0.43, Figure

6D, see also Figure 5C). To determine whether this transition was caused by a change in the

maturation rate of CME sites, we examined Ede1-GFP tracks in circumferential kymographs

made from cells before and after osmotic shock (Figure 6E). We observed a significant

decrease in the proportion of long-lived early CME sites upon osmotic shock, suggesting

that early phase CME sites mature to the late, regular phase more quickly (from 22.9%

persistent sites to 0.2%, 54.3% partial sites compared to 38.6%, Figure 6F). Interestingly,

the apparent shortening of Ede1-GFP lifetimes by osmotic shock does not extend to late

phase CME proteins. While the ratio of the number of Abp1-RFP to Sla1-GFP puncta was

unchanged, both Sla1-GFP and Abp1-RFP lifetimes were significantly increased upon

osmotic shock (median lifetimes of 22 and 10 s, respectively, before osmotic shock and 34

and 13 s, respectively, after. Figure 6 Supplement 1B-E). Lengthening of Sla1-GFP and Abp1-RFP lifetimes is likely due to a mechanical burden on membrane invagination/internalization caused by increased turgor pressure in hypotonic solution (Aghamohammadzadeh and Ayscough, 2009; Boulant et al., 2011; Hassinger et al., 2017). The results of our osmotic shock experiments demonstrate that moving endocytic cargo from places where it is concentrated to places where it is scarce can activate stalled endocytic sites, even as vesicle internalization itself is slowed.

In contrast to forced depolarization, elimination of endocytic cargo causes endocytic sites to stall. We acutely blocked accumulation of endocytic cargo on the plasma membrane through treating cells with the drug Brefeldin A (BFA), a secretion pathway inhibitor (Graham et al., 1993). BFA inhibits secretion in cells lacking the sterol synthesis gene *ERG6*, but it does not affect secretion in wild-type cells (Graham et al., 1993). Large-budded *erg6Δ* cells resemble large-budded wild-type cells, with both early and late CME sites visible in the mother (Figure 6G, Figure 6 Supplement 1F). Treating *erg6Δ* cells with BFA significantly reduced the ratio of late endocytic sites to early endocytic sites in large-budded mothers (Figure 6G-H). Kymographs along the mother cortex and analysis of early endocytic site turnover indicate that slowed maturation rate and increased frequency of persistent, early endocytic sites accounts for this change in ratio of late endocytic sites to early ones (Figure 6I-J). Interestingly, treatment of wild-type cells with BFA caused a slight increase in the ratio of late endocytic sites to early ones, accounted for by a slight drop in the proportion of endocytic sites that turnover, although this effect was very small (Figure 6 Supplement 1F-I). These data demonstrate that availability of cargo at endocytic sites

licenses CME site maturation from the early phase to the late phase of the pathway. The

transition point we identified can therefore be considered a cargo checkpoint.

The mechanism of cargo sensing within the CME pathway remains unclear. We

considered one candidate cargo-sensing molecule, the casein kinase Hrr25. A previous

study from our lab identified several proteins from the early phase of CME as Hrr25

substrates (Peng et al., 2015a). In the same study, cells lacking Hrr25 were found to have

fewer Sla1 sites, while cells overexpressing Hrr25 had more Sla1 sites. We wondered

whether this effect could be due to modulation of the rate of transit through the cargo

checkpoint. While depletion of Hrr25 using an auxin inducible degron (AID, Nishimura et

al., 2009) caused a growth defect detectable at low temperatures (Figure 6 Supplement

2A), acutely-induced and nearly complete degradation of the kinase caused only a

moderate increase in the fraction of long-lived Ede1-GFP sites in small-budded mothers

(Figure 6 Supplement 2B-D). Conversely, overexpression of Hrr25 using a β-estradiol

induced overexpression system (McIsaac et al., 2011) caused only a moderate increase in

the fraction of short-lived Ede1-GFP sites in mothers of small-budded cells (Figure 6

Supplement 2E-F). While the mechanism of cargo sensing in the CME pathway remains

unclear, these findings suggest that the casein kinase Hrr25 may contribute to the

transition.

19
20
21   **The regulatory transition point occurs between recruitment of the intermediate**
22   **coat proteins and late coat proteins**
23
24   Since a representative CME protein from before the regulatory transition point (Ede1) was

25   not significantly polarized in small-budded cells but a late phase CME protein (Sla1) was

(Figure 5), we used this difference in CME protein distribution as an assay to pinpoint the time of the regulatory transition point in the CME pathway. We recorded Z-stacks of proteins from the early, early coat, intermediate coat, and late coat modules using either epifluorescence microscopy or spinning disc microscopy (depending on how well we could distinguish CME puncta above background noise). The early proteins Syp1 and Hrr25, the early coat proteins Apl1 and Pal1, and the intermediate coat proteins Sla2, Ent1, and Ent2 all display distributions similar to that of Ede1: in highly polarized cells with few Sla1-mCherry-marked late CME sites, we nevertheless observed many puncta of these query proteins, placing them upstream of the transition point (median ratios of 0.14 vs. 0.59 (Syp1), 0.09 vs. 0.49 (Sla2), 0.16 vs. 0.79 (Ent2), Figure 7A-B, Figure 7 Supplement 1). The late coat marker Pan1, on the other hand, polarizes with Sla1-mCherry-marked late CME sites (median ratio of 0.62 vs. 0.64, Figure 7A-B). These data indicate that the transition point in the CME pathway occurs after recruitment of the intermediate coat module but before recruitment of the late coat proteins Pan1 and Sla1.

Since we concluded that polarized cargo distribution drives polarization of late endocytic sites during the cell cycle through controlling maturation rate, we predicted that mutant cells that could not sense cargo at endocytic sites would also fail to undergo cycles of late endocytic site polarization and depolarization. A mutant with genes encoding seven early arriving endocytic proteins deleted ("*7Δ*") has previously been reported to complete endocytosis without internalizing cargo (Brach et al., 2014). When we examined these mutants, we found that they were no longer polarized during the small-budded stage of the cell cycle (Figure 7C). We compared the ratio of late CME sites (marked by Sla1-mCherry) to early sites (marked by Sla2-GFP) in small- and large-budded *7Δ* mothers and found that

1   the ratio was no longer significantly different in the *7Δ* mutants (Figure 7 C-D). In contrast,

2   small- and large-budded mothers were insignificantly different with respect to the ratio of

3   the number of puncta of two late endocytic markers to one another (Figure 7 Supplement

4   1B-C). Thus the cargo-sensitive maturation rate of individual endocytic sites accounts for

5   polarization of late endocytic sites during specific phases of the cell cycle.

Discussion

Despite identification of dozens of proteins involved in CME through decades of research,

how progression through this pathway is regulated is poorly understood. In this study, we

conducted systematic imaging experiments of CME in budding yeast with the goal of

revealing new regulatory principles. Previous studies used two-color imaging to describe a

protein recruitment cascade at CME sites; however, these studies typically analyzed the

recruitment dynamics of each protein independently, therefore losing information about

how the behavior of one protein is correlated with the behavior of another at individual

endocytic sites. To uncover inter-relationships in the behavior of individual components of

the CME machinery, we performed quantitative analysis of the pair-wise recruitment and

abundance behavior of CME proteins at >17,000 endocytic sites.

Our data set revealed more variation in the behavior of individual proteins in the

budding yeast CME pathway than had previously been appreciated. CME has been

described as a highly regular molecular pathway. The ordered arrival of CME proteins at

endocytic sites is roughly conserved from yeasts to humans and is nearly invariant (Doyon

et al., 2011; Kaksonen et al., 2005, 2003; Sun et al., 2019; Taylor et al., 2011). While the

initial steps of CME were known to occur with variable timing, the timing of the late steps

of the pathway had been considered to be remarkably regular (Kaksonen et al., 2006).

Previous studies even used lifetimes of late phase CME proteins as readouts for endocytic

efficiency (Kaksonen et al., 2005). While our data confirm that the early stages of CME

occur with more variable timing than the late stages, we were surprised to nevertheless

observe variability even in what has previously been considered the "regular phase" of the

process. We took advantage of this previously underappreciated variability in timing to

gain new insights into the CME pathway by looking for covariation in the pair-wise

behavior of endocytic proteins.

Our analysis suggests that a full complement of early arriving CME proteins is

present at each endocytic site at initiation. Each early arriving endocytic protein has been

reported to have a variable lifetime (Carroll et al., 2012, 2009; Newpher et al., 2005; Peng

et al., 2015a; Stimpson et al., 2009). However, the implication of this variability was

previously unknown. One possible interpretation was that long-lived early CME sites

represent sites that are incompletely assembled, potentially lacking some component of the

early or early coat module. Another possibility was that the early proteins assemble fully

but that maturation to the next phase of the pathway is stalled due to an inhibitory

checkpoint signal. Our pair-wise analysis of the lifetimes of each early arriving protein and

Ede1 shows that they are well correlated. This correlation is consistent with the possibility

that CME proteins assemble at the nascent sites in a coordinated manner, but that

advancement to the next phase awaits release of a checkpoint.

Analyses of correlations between the behaviors of different pairs of proteins at

individual CME sites also reveal two quantitatively separable stages, substantiating the

existence of a regulatory transition point in the CME pathway (Carroll et al., 2012, Figure

8A). This conclusion is based on the loss of correlation of lifetimes and intensities for

proteins spanning a specific step in the pathway. While lifetimes of early arriving proteins

are predictive of one another (for example, a plot of Ede1 lifetime vs. Syp1 lifetime yields a

linear fit with a slope of 0.65 and an R^2 of 0.35), in no case does the behavior of an early

arriving protein strongly correlate with the behavior of a late arriving protein (for example,

a plot of Ede1 lifetime vs. Abp1 lifetime yields a linear fit with a slope of 0.08 and an R^2 of

0.03). Conversely, lifetimes of late arriving proteins are fairly good predictors of the maximum intensity of the endocytic actin reporter Abp1 (Slopes ranging from 0.46 to 0.96). The striking correlations within groups of proteins recruited early or late, but not between these groups, is consistent with the notion that robust protein interaction networks are at play during CME: an early complex of proteins establishes endocytic sites and a late complex mediates membrane invagination and vesicle scission (Figure 8A). The poor correlation between the lifetimes and intensities of these two networks likely indicates that there are fewer molecular links between the two networks than within each respective network. Since the behavior of proteins in the early and late phases have behaviors that are predictive of behaviors of other proteins in the same phase, but not the other phase, CME initiation and internalization can be thought of as two separable processes.

Through observation of CME site composition and dynamics while directionality of polarized yeast growth changed naturally in a programmed manner through the cell cycle, and through genetic and chemical perturbation of secretory traffic, we determined that progression from the early to the late phase of CME is controlled by cargo (Figure 8A). Although it is formally possible that the cell polarity machinery composed of proteins such as Cdc42 and Cdc24 might directly trigger the transition from the early to the late phase of CME, we favor the possibility that cargo plays a direct role. This is in part because CME would be more efficient with a cargo sensing mechanism. Our studies also suggest that a portion of the variability observed for early arriving endocytic proteins might be explained by differences in the polarization state of the cells analyzed. Thus, in *S. cerevisiae*, cargo-

mediated regulation of CME progression occurs through endocytic stalling, rather than

abortion, in the absence of cargo.

While the specific molecular mechanism of this proposed cargo checkpoint is

incompletely understood, several intriguing possibilities provide exciting avenues for

further research. Importantly, several studies have suggested that a cargo checkpoint might

regulate progress through the mammalian CME pathway (Liu et al., 2010; Loerke et al.,

2009; Mettlen et al., 2010; Puthenveedu and von Zastrow, 2006). A recent reconstitution

study of mammalian CME suggests that clathrin continually assembles and disassembles

during the early stage of the process, with cargo binding stabilizing membrane-associated

clathrin structures (Chen et al., 2019). Fluorescently-tagged early arriving CME proteins in

budding yeast are also known to fluctuate in intensity, suggesting that such a dynamic

proofreading mechanism may be evolutionarily conserved (Carroll et al., 2012). Cargo

sensing might involve post-translational modification. We provided evidence that the

casein kinase Hrr25 may play a role in progression of CME from the cargo collection phase

to the internalization phase. In addition, ubiquitination has also been suggested to play a

role in CME (Weinberg and Drubin, 2014). One attractive possibility is that early arriving

CME proteins dynamically bind and unbind nascent endocytic sites until cargo binding

induces a conformational change, facilitating post-translational modification and

maturation.

Because representative early phase CME proteins do not appear polarized in

budding yeast while representative late phase CME proteins do, we assessed polarization

state of more proteins to precisely define which CME proteins are present on either side of

the transition point in the CME pathway. We found that sites containing Ede1, Syp1, Hrr25,

the AP-2 complex, Pal1, Sla2, and the epsins all behaved similarly in that they are found both proximal and distal to sites of polarized growth. In contrast, sites containing Pan1 and Sla1, late coat proteins, are highly polarized (Figure 8A). Importantly, it is interesting to consider implications of our new observations on certain previous observations. Acute depletion of late coat proteins Pan1 and End3, the first proteins to arrive at CME sites after Sla2 and the epsins, leads to arrest of early endocytic sites on the plasma membrane and aberrant assembly of the late endocytic proteins including the actin cytoskeleton in the cytoplasm (Sun et al., 2015). Thus recruitment of these late coat proteins is likely the crucial step in transitioning from the early phase to the late phase of CME. Conversely, deletion of genes encoding seven early arriving proteins (Syp1, Ede1, Apl1, Pal1, Pal2, Yap1801, and Yap1802) leads to defects in cargo collection but does not halt endocytosis itself (Brach et al., 2014). Loss of a "checkpoint" protein that halts progress in endocytosis in the absence of cargo may be permissive for progress in the CME pathway if that protein does not perform an additional, essential CME function. One of these seven early arriving proteins might therefore be responsible for detecting cargo in a nascent endocytic pit and releasing the block on Pan1/End3 recruitment.

Together, the data we collected and analyzed in this study suggest a simple conceptual model to explain where and when endocytic internalization is triggered (Figure 8B). Endocytic sites are initiated widely across the plasma membrane. However, in the absence of cargo, they mature to the internalization phase slowly due to an unsatisfied cargo checkpoint, rather than aborting in the absence of cargo, as has been proposed previously in several studies of mammalian cells (Ehrlich et al., 2004; Liu et al., 2010; Loerke et al., 2009; Mettlen et al., 2010). Localized delivery of endocytic cargos via

exocytosis triggers localized maturation of CME sites through release of the cargo

checkpoint. This interpretation represents an attractive explanation for the decades-old

observation that "actin patches" (late CME sites) are polarized in small and medium

budded *S. cerevisiae* cells (Adams and Pringle, 1984). Because secretion is directed

primarily to the bud while it is growing (Field and Schekman, 1980), concentrated

endocytic cargos in the bud trigger local maturation of endocytic sites through the cargo

checkpoint, after which actin assembles.

2   **Strains and yeast husbandry**
3
4   The strains used in this study are listed in Supplementary Table 1. All budding yeast

5   strains were derived from the wild-type diploid DDY1102 and propagated using standard

6   techniques (Amberg et al., 2005). C-terminal fluorescent protein fusions were constructed

7   specifically for this study or for earlier studies, as indicated in Table 1, as described

8   previously (Longtine et al., 1998).

9
10
11   **Live-cell imaging**
12
Cells were grown to mid log phase in imaging media (synthetic minimal media

supplemented with Adenine, L-Histidine, L-Leucine, L-Lysine, L-Methionine, Uracil, and 2%

glucose), then adhered to coverslips coated with 0.2 mg/ml Concanavalin A.

TIRFM imaging was performed on a Nikon Eclipse Ti2 inverted microscope with a

Nikon CFI60 60× 1.49 numerical aperture (NA) Apo oil immersion TIRFM objective and a

Hamamatsu Orca-Flash 4.0 V2 sCMOS camera. GFP and RFP/mCherry were excited using

488- and 561-nm lasers and detected using a Chroma HC TIRFM Quad Dichroic (C-FL TIRF

Ultra Hi S/N 405/488/561/638) and Chroma HC Quad emission filters BP 525/550 and BP

600/650, respectively. Channels were acquired sequentially. The system was controlled

with Nikon Elements software and maintained at 25°C by an OkoLab environmental

chamber. Frames were separated by 1 sec.

Epifluorescence imaging was carried out on a Nikon Eclipse Ti inverted microscope

with a Nikon 100x 1.4 NA Plan Apo VC oil immersion objective and an Andor Neo 5.5

sCMOS camera. GFP and RFP/mCherry fluorescence were excited using a Lumencore

Spectra X LED light source with 550/15 nm and 470/22 nm excitation filters. For two-color

imaging, channels were acquired sequentially using an FF-493/574-Di01 dual pass dichroic

mirror and FF01-524/628-25 dual pass emission filters (Semrock, Rochester, NY). The

system was controlled with Nikon Elements software and maintained at 25°C by an

environmental chamber (In Vivo Scientific, St. Louis, MO).

Spinning disc confocal microscopy was carried on a Nikon Eclipse Ti inverted

microscope with a Nikon 100x 1.45 NA Plan Apo λ oil immersion objective, an Andor IXon

X3 EM-CCD camera and Andor CSU-X spinning disc confocal equipment. GFP fluorescence

was excited using a 488 nm laser and detected with a Chroma 535/20 nm emission filter

(Bellows Falls, VT). mCherry fluorescence was excited using a 565 nm laser and detected

with a Chroma 605/52 nm emission filter. The system was controlled with Nikon Elements

software. Imaging was conducted at room temperature (~23°C).

13
14
15 **Osmotic shock and BFA experiments**
16
For osmotic shock and BFA experiments, cells were adhered to Concanavalin A coated

coverslips for live cell imaging. For osmotic shot experiments, the cells were overlaid with

only 250 μL imaging media. Osmotic shock was achieved through rapid addition of 4 mL of

sterile water or 4 mL imaging media as a control. For BFA experiments, the cells were

overlaid with 500 μL imaging media and BFA treatment was initiated by rapidly adding

imaging media supplemented with BFA such that the final concentration was 75 μg/mL

(Graham et al., 1993).

24
25

Auxin depletion experiments

For auxin depletion experiments, 500 µM 3-indoleacetic acid (auxin, Sigma) was prepared

freshly in DMSO and added to standard *S. cerevisiae* growth media. As depletion is most

efficient in slightly acidic media, YPD media used in growth assays was supplemented with

0.1 M potassium phosphate buffer (pH 6.2). Liquid imaging media was sufficiently acidic

without buffering.

      For growth comparisons in the presence of auxin, mid log phase starter cultures

were diluted to an OD_{600} of 0.1, 0.02, and 0.004, spotted on plates with and without added

auxin, and grown for 48 hours at 18° C and 25° C.

      For immunoblot and live cell imaging analysis, auxin was added directly to mid log

phase cultures in imaging media. The cultures were then incubated at room temperature

(~23° C) for the indicated time interval, after which they were either imaged by

epifluorescence microscopy or processed for immunoblot analysis.

β-estradiol-induced overexpression experiments

Hormone-induced overexpression of Hrr25 was carried out essentially as in Peng et al.

2015a. Briefly, 50 mM β-estradiol (Sigma) was prepared freshly in DMSO and added to mid

log phase cultures in imaging media. The cultures were then incubated at 25° C for the

indicated time interval, after which they were either imaged by epifluorescence microscopy

or processed for immunoblot analysis.

Immunoblot analysis

Cells were grown to mid log phase in the appropriately supplemented imaging media, harvested by centrifugation, washed with water, and lysed by bead beating in a lysis buffer with 5 mM Tris-HCl pH 8, 5 mM EDTA pH 8, and 15% Trichloroacetic acid (TCA). The TCA precipitate was then washed with ice-cold acetone and resuspended in Tris/urea sample buffer (62.5 mM Tris pH 6.8, 3 M urea, 1% sodium dodecyl sulfate, 5% beta mercaptoethanol, 0.05% bromophenol blue). Extracts were separated electrophoretically on 10% polyacrylamide gels and transferred to nitrocellulose. Membranes were blocked with 5% milk and probed with the one or more of the following antibodies (indicated): a mouse monoclonal antibody against the V5 epitope tag (Invitrogen), mouse monoclonal antibody clone 9E10 against a 13-myc tag (prepared from hybridoma supernatant (Lewellyn et al., 2015)), a mouse antibody raised against Hrr25 (Abmart, Peng et al., 2015b), or a mouse monoclonal antibody against Pgk1 (459250; Novex, Winston-Salem, NC). Membranes were then incubated with secondary antibodies conjugated to infrared fluorophores (Li-Cor Biosciences inc., Lincoln, NE). Bands were visualized using an Odyssey infrared imaging system (Li-Cor Biosciences).

Image and data analysis

Tracking of endocytic sites was performed using the MATLAB package cmeAnalysis (Aguet et al., 2013). The red and green channels for each movie were tracked independently, and the centroid position over time for each site determined. These data were used as input in custom MATLAB scripts that associated colocalized tracks in the red and green channels while rejecting sites that appeared within 350 nm of one another (Hong et al., 2015). The

fluorescence intensity for each spot was calculated as an integrated intensity within a circular region with a diameter of 7 pixels (756 nm), centered at the spot position determined by cmeAnalysis. The background fluorescence was calculated as the average fluorescence intensity of an annulus 2 pixels (216 nm) wide surrounding the circular region used to calculate the spot intensity.

Custom MATLAB scripts were then used to clean our dataset by rejecting tracks that fall into several categories: sites with a low signal-to-noise (brightness not significantly above background, Hong et al., 2015) were excluded from further analysis. Tracks that began or terminated within 5 frames of the beginning or end of the movie were excluded so that only complete tracks were kept for analysis. Sites whose calculated background-subtracted intensity went below $-0.25\times$ the maximum intensity were excluded to prevent abnormally high background or presence of nearby sites from affecting intensity measurements. Finally, outliers in spot lifetime, fluorescence intensity, time to arrival, and time to disappearance (defined as more than $1.5\times$ the interquartile distance from the median value) were also excluded. This was done to exclude erroneous traces in which two distinct tracks had been inadvertently linked, where two sites overlapped and thus were substantially brighter, as well as erroneous putative colocalizations. While these analyses were blind to cell cycle stage, long-lived early sites in polarized mothers represented a small fraction of the final data set owing to the requirement for association of query protein tracks with reference protein tracks and exclusion of incomplete tracks.

GFP maximum fluorescence intensity and the lifetimes and maximum intensities of the red reference proteins were fit to the lifetimes of the GFP-tagged query proteins via simple linear regression. To compare the slopes of the fit across GFP-tagged proteins with

differing brightnesses and lifetimes, the data were first normalized by dividing by their

respective medians, which does not change the R^2 of the fit.

      Analysis of the influence of cargo on CME site maturation was carried out using Fiji

software (National Institutes of Health, Bethesda, MD). For quantification of Ede1-GFP

turnover, circumferential kymographs were generated by drawing a line of width 5 pixels

around the circumference of each cell. For quantification of Sla1-GFP and Abp1-RFP patch

lifetimes, radial kymographs were generated using a custom Fiji macro that generates a

kymograph at every radius around a circle in 2-degree increments. The lifetimes of the first

5 kymographs generated were measured to avoid bias. For intensity profile analysis of

Bem1-GFP, intensity profiles were normalized to cell length and averaged using a Fiji

plugin provided by C. Brownlee (Brownlee and Heald, 2019). Exact numbers of cells and

CME sites measured are described in the figure legends.

Figure preparation

For figure panels and movies, individual cells were cropped, background fluorescence was

uniformly subtracted from the image stack, and photo bleaching was fit to a linear decay

function and corrected in Fiji. Look up tables used for display are linear. Plots were

generated in MATLAB or Prism 8 (GraphPadSoftware, San Diego, CA). Figures were

assembled in Adobe illustrator Creative Cloud 2019.

Reproducibility of experiments

All experimental results presented were replicated in at least three distinct experiments to

ensure reproducibility.

For TRIFM imaging experiments, endocytic sites from each of the dozens of cells in

several fields of view were tracked simultaneously. As the results from each dataset were

indistinguishable, they were pooled for further analysis. The number of tracked spots for

each protein pair is presented in Supplementary Tables 2-10. Statistical analysis was

conducted in MATLAB.

For epifluorescence and confocal fluorescence imaging experiments, multiple cells

from   each   replicate   were   analyzed.   As   data   from   different   replicates   were

indistinguishable, they were pooled for statistical analysis. The specific number of cells

analyzed is indicated in each figure legend. Statistical analysis was conducted in Prism 8.

10
11
12   **Data and code availability**

13   The data and code described can be found at the following links:
14   https://github.com/DrubinBarnes/Pedersen_Hassinger_Marchando_Drubin_CME_Manuscr
15   ipt_2019
16   https://drive.google.com/drive/folders/1I-v1BiOq0la4_07y74EOlG6_6JexlqW4
17
18

Acknowledgements

We thank Yui Iwamoto for critically reading the manuscript. We are grateful to Tony

Bretscher for providing the GFP-Sec4 yeast strain, Michelle Lu for providing the Bem1-GFP

yeast strain, and Marko Kaksonen for providing the *7Δ* strains. Spinning disc confocal

microscopy was conducted at the University of California, Berkeley Cancer Research

Laboratory Molecular Imaging center, supported by the Gordon and Betty Moore

foundation. We would like to thank H. Aaron and F. Ives for their microscopy training and

assistance. This research was conducted with US Government support, under and awarded

by Department of Defense, Air Force Office of Scientific Research, National Defense Science

and Engineering Graduate Fellowship 32 CFR 168a (to J.E.H.); National Institutes of Health

Grant R35GM118149 (to D.G.D.).

1 **Author Contributions**

2 R.T.A. Pedersen, J.E. Hassinger, and D.G. Drubin conceived of the experiments. R.T.A.

3 Pedersen and P. Marchando generated the reagents. R.T.A. Pedersen, J.E. Hassinger, and P.

4 Marchando performed the experiments, and analyzed the data. R.T.A. Pedersen, J.E.

5 Hassinger, and D.G. Drubin wrote the manuscript. D.G. Drubin secured funding.

1 **Competing interests**

2 The authors declare no competing interests.

References

Adams AE, Pringle JR. 1984. Relationship of actin and tubulin distribution to bud growth in
wild- type and morphogenetic-mutant *Saccharomyces cerevisiae. J Cell Biol* **98**:934–
945. doi:10.1083/jcb.98.3.934
Aghamohammadzadeh S, Ayscough KR. 2009. Differential requirements for actin during
yeast and mammalian endocytosis. *Nat Cell Biol* **11**:1039–1042. doi:10.1038/ncb1918
Aguet F, Antonescu CN, Mettlen M, Schmid SL, Danuser G. 2013. Advances in analysis of low
signal-to-noise images link dynamin and AP2 to the functions of an endocytic
checkpoint. *Dev Cell* **26**:279–291. doi:10.1016/j.devcel.2013.06.019
Amberg DC, Burke DJ, Strathern JN. 2005. Methods in Yeast Genetics: A Cold Spring Harbor
Laboratory Course Manual, 2005 Edition, A Cold Spring Harbor Laboratory Course
Manual.
Axelrod D. 1989. Total Internal Reflection Fluorescence Microscopy. *Methods Cell Biol*
30:245–270.
Bitsikas V, Corrêa IR, Nichols BJ. 2014. Clathrin-independent pathways do not contribute
significantly to endocytic flux. *Elife* **3**:1–26. doi:10.7554/elife.03970
Boulant S, Kural C, Zeeh J-C, Ubelmann F, Kirchhausen T. 2011. Actin dynamics counteract
membrane tension during clathrin-mediated endocytosis. *Nat Cell Biol* **13**:1124–31.
doi:10.1038/ncb2307
Brach T, Godlee C, Moeller-Hansen I, Boeke D, Kaksonen M. 2014. The initiation of clathrin-
mediated endocytosis is mechanistically highly flexible. *Curr Biol* **24**:548–54.
doi:10.1016/j.cub.2014.01.048
Bradford MK, Whitworth K, Wendland B. 2015. Pan1 regulates transitions between stages
of clathrin-mediated endocytosis. *Mol Biol Cell* **26**:1371–1385. doi:10.1091/mbc.E14-
11-1510
Brownlee C, Heald R. 2019. Importin α Partitioning to the Plasma Membrane Regulates
Intracellular Scaling. *Cell* **176**:805-815.e8. doi:10.1016/j.cell.2018.12.001
Carroll SY, Stimpson HEM, Weinberg J, Toret CP, Sun Y, Drubin DG. 2012. Analysis of yeast
endocytic site formation and maturation through a regulatory transition point. *Mol*
Biol Cell **23**:657–668. doi:10.1091/mbc.E11-02-0108
Carroll SY, Stirling PC, Stimpson HEM, Gießelmann E, Schmitt MJ, Drubin DG. 2009. A Yeast
Killer Toxin Screen Provides Insights into A/B Toxin Entry, Trafficking, and Killing
Mechanisms. *Dev Cell* **17**:552–560. doi:10.1016/j.devcel.2009.08.006
Chen Y, Yong J, Martinez-sanchez A, Yang Y, Wu Y, De Camilli P, Fernández-busnadiego R,
Wu M. 2019. Dynamic Instability of Clathrin Assembly Provides Proofreading Control
for Endocytosis. *J Cell Biol*. doi:10.1083/jcb.201804136
Cope MJ, Yang S, Shang C, Drubin DG. 1999. Novel protein kinases Ark1p and Prk1p
associate with and regulate the cortical actin cytoskeleton in budding yeast. *J Cell Biol*
144:1203–18.
Doyon JB, Zeitler B, Cheng J, Cheng AT, Cherone JM, Santiago Y, Lee AH, Vo TD, Doyon Y,
Miller JC, Paschon DE, Zhang L, Rebar EJ, Gregory PD, Urnov FD, Drubin DG. 2011.
Rapid and efficient clathrin-mediated endocytosis revealed in genome-edited
mammalian cells. *Nat Cell Biol* **13**:331–337. doi:10.1038/ncb2175
Ehrlich M, Boll W, Van Oijen A, Hariharan R, Chandran K, Nibert ML, Kirchhausen T. 2004.

Endocytosis by random initiation and stabilization of clathrin-coated pits. *Cell* **118**:591–605. doi:10.1016/j.cell.2004.08.017

Field C, Schekman R. 1980. Localized secretion of acid phosphatase reflects the pattern of cell surface growth in *Saccharomyces cerevisiae*. *J Cell Biol* **86**:123–128. doi:10.1083/jcb.86.1.123

Glantz SA, Slinker BK, Neilands TB. 2001. Primer of Applied Regression and Analysis of Variance, 3rd ed. New York, NY: McGraw-Hill.

Godlee C, Kaksonen M. 2013. From uncertain beginnings: Initiation mechanisms of clathrin-mediated endocytosis. *J Cell Biol* **203**:717–725. doi:10.1083/jcb.201307100

Graham TR, Scott PA, Emr SD. 1993. Brefeldin A reversibly blocks early but not late protein transport steps in the yeast secretory pathway. *EMBO J* **12**:869–877. doi:10.1002/j.1460-2075.1993.tb05727.x

Hassinger JE, Oster G, Drubin DG, Rangamani P. 2017. Design principles for robust vesiculation in clathrin-mediated endocytosis. *Proc Natl Acad Sci U S A* **114**:E1118–E1127. doi:10.1073/pnas.1617705114

Holland DO, Johnson ME. 2018. Stoichiometric balance of protein copy numbers is measurable and functionally significant in a protein-protein interaction network for yeast endocytosis. *PLoS Comput Biol* **14**:1–34. doi:10.1371/journal.pcbi.1006022

Hong SH, Cortesio CL, Drubin DG. 2015. Machine-Learning-Based Analysis in Genome-Edited Cells Reveals the Efficiency of Clathrin-Mediated Endocytosis. *Cell Rep* **12**:2121–2130. doi:10.1016/j.celrep.2015.08.048

Kaksonen M, Sun Y, Drubin DG. 2003. A pathway for association of receptors, adaptors, and actin during endocytic internalization. *Cell* **115**:475–87.

Kaksonen M, Toret CP, Drubin DG. 2006. Harnessing actin dynamics for clathrin-mediated endocytosis. *Nat Rev Mol Cell Biol* **7**:404–414. doi:10.1038/nrm1940

Kaksonen M, Toret CP, Drubin DG. 2005. A modular design for the clathrin- and actin-mediated endocytosis machinery. *Cell* **123**:305–20. doi:10.1016/j.cell.2005.09.024

Layton AT, Savage NS, Howell AS, Carroll SY, Drubin DG, Lew DJ. 2011. Modeling vesicle traffic reveals unexpected consequences for Cdc42p-mediated polarity establishment. *Curr Biol* **21**:184–194. doi:10.1016/j.cub.2011.01.012

Lew DJ, Reed SI. 1995. Cell cycle control of morphogenesis in budding yeast. *Curr Opin Genet Dev* **5**:17–23. doi:10.1016/S0959-437X(95)90048-9

Lewellyn EB, Pedersen RTA, Hong J, Lu R, Huntly M, Drubin DG. 2015. An Engineered Minimal WASP-Myosin Fusion Protein Reveals Essential Functions for Endocytosis. *Dev Cell* **35**:281–294. doi:10.1016/j.devcel.2015.10.007

Lewis MJ, Nichols BJ, Prescianotto-Baschong C, Riezman H, Pelham HRB. 2000. Specific Retrieval of the Exocytic SNARE Snc1p from Early Yeast Endosomes. *Mol Biol Cell* **11**:23–38. doi:10.1091/mbc.11.1.23

Liu AP, Aguet F, Danuser G, Schmid SL. 2010. Local clustering of transferrin receptors promotes clathrin-coated pit initiation. *J Cell Biol* **191**:1381–1393. doi:10.1083/jcb.201008117

Loerke D, Mettlen M, Yarar D, Jaqaman K, Jaqaman H, Danuser G, Schmid SL. 2009. Cargo and Dynamin Regulate Clathrin-Coated Pit Maturation. *PLoS Biol* **7**:e1000057. doi:10.1371/journal.pbio.1000057

Longtine MS, McKenzie a, Demarini DJ, Shah NG, Wach a, Brachat a, Philippsen P, Pringle JR. 1998. Additional modules for versatile and economical PCR-based gene deletion

and modification in *Saccharomyces cerevisiae. Yeast* **14**:953–61. doi:10.1002/(SICI)1097-0061(199807)14:10<953::AID-YEA293>3.0.CO;2-U

Lu R, Drubin DG. 2017. Selection and stabilization of endocytic sites by Ede1, a yeast functional homologue of human Eps15. *Mol Biol Cell* **28**:567–575. doi:10.1091/mbc.E16-06-0391

Lu R, Drubin DG, Sun Y. 2016. Clathrin-mediated endocytosis in budding yeast at a glance. *J Cell Sci* **129**:1531–1536. doi:10.1242/jcs.182303

Manenschijn HE, Picco A, Mund M, Ries J, Kaksonen M. 2019. Type-I myosins promote actin polymerization to drive membrane bending in endocytosis. *Elife* **17**:490011. doi:10.1101/490011

McIsaac RS, Silverman SJ, McClean MN, Gibney PA, Macinskas J, Hickman MJ, Petti AA, Botstein D. 2011. Fast-acting and nearly gratuitous induction of gene expression and protein depletion in *Saccharomyces cerevisiae. Mol Biol Cell* **22**:4447–4459. doi:10.1091/mbc.E11-05-0466

Mettlen M, Loerke D, Yarar D, Danuser G, Schmid SL. 2010. Cargo- and adaptor-specific mechanisms regulate clathrin-mediated endocytosis. *J Cell Biol* **188**:919–933. doi:10.1083/jcb.200908078

Newpher TM, Smith RP, Lemmon V, Lemmon SK. 2005. In vivo dynamics of clathrin and its adaptor-dependent recruitment to the actin-based endocytic machinery in yeast. *Dev Cell* **9**:87–98. doi:10.1016/j.devcel.2005.04.014

Nishimura K, Fukagawa T, Takisawa H, Kakimoto T, Kanemaki M. 2009. An auxin-based degron system for the rapid depletion of proteins in nonplant cells. *Nat Methods* **6**:917–922. doi:10.1038/nmeth.1401

Pedersen RT, Drubin DG. 2019. Type I myosins anchor actin assembly to the plasma membrane during clathrin-mediated endocytosis. *J Cell Biol* **218**:1138–1147. doi:10.1083/jcb.201810005

Peng Y, Grassart A, Lu R, Wong CCL, Yates J, Barnes G, Drubin DG. 2015a. Casein Kinase 1 Promotes Initiation of Clathrin-Mediated Endocytosis. *Dev Cell* **32**:231–240. doi:10.1016/j.devcel.2014.11.014

Peng Y, Moritz M, Han X, Giddings TH, Lyon A, Kollman J, Winey M, Yates III J, Agard DA, Drubin DG, Barnes G. 2015b. Interaction of CK1δ with TuSC ensures proper microtubule assembly and spindle positioning. *Mol Biol Cell* **26**:1–37. doi:10.1091/mbc.E14-12-1627

Picco A, Mund M, Ries J, Nedelec F, Kaksonen M. 2015. Visualizing the functional architecture of the endocytic machinery. *Elife* **4**:1–29. doi:10.7554/eLife.04535

Pringle JR, Preston RA, Adams AEM, Stearns T, Drubin DG, Haarer BK, Jones EW. 1989. Fluorescence Microscopy Methods for Yeast. *Methods Cell Biol* **31**:357–435. doi:10.1016/S0091-679X(08)61620-9

Puthenveedu MA, von Zastrow M. 2006. Cargo Regulates Clathrin-Coated Pit Dynamics. *Cell* **127**:113–124. doi:10.1016/j.cell.2006.08.035

Reed GF, Lynn F, Meade BD. 2002. Use of Coefficient of Variation in Assessing Variability of Quantitative Assays. *Clin Diagn Lab Immunol* **9**:1235–1239. doi:10.1128/CDLI.9.6.1235

Skruzny M, Brach T, Ciuffa R, Rybina S, Wachsmuth M, Kaksonen M. 2012. Molecular basis for coupling the plasma membrane to the actin cytoskeleton during clathrin-mediated endocytosis. *Proc Natl Acad Sci U S A* **109**:E2533–E2542.

doi:10.1073/pnas.1207011109

Stimpson HEM, Toret CP, Cheng AT, Pauly BS, Drubin DG. 2009. Early-Arriving Syp1p and Ede1p Function in Endocytic Site Placement and Formation in Budding Yeast. *Mol Biol Cell* **20**:4640–4651. doi:10.1091/mbc.E09

Sun Y, Leong NT, Wong T, Drubin DG. 2015. A Pan1/End3/Sla1 complex links Arp2/3-mediated actin assembly to sites of clathrin-mediated endocytosis. *Mol Biol Cell* **26**:3841–3856. doi:10.1091/mbc.E15-04-0252

Sun Y, Martin AC, Drubin DG. 2006. Endocytic Internalization in Budding Yeast Requires Coordinated Actin Nucleation and Myosin Motor Activity. *Dev Cell* **11**:33–46. doi:10.1016/j.devcel.2006.05.008

Sun Y, Schöneberg J, Chen X, Jiang T, Kaplan C, Xu K, Pollard TD, Drubin DG. 2019. Direct comparison of clathrin-mediated endocytosis in budding and fission yeast reveals conserved and evolvable features. *Elife* **8**. doi:10.7554/eLife.50749

Takizawa PA, Derisi JL, Wilhelm JE, Vale RD. 2000. Plasma Membrane Compartmentalization in Yeast by Messenger RNA Transport and a Septin Diffusion Barrier. *Science* **290**:341–344. doi:10.1126/science.290.5490.341

Taylor MJ, Perrais D, Merrifield CJ. 2011. A high precision survey of the molecular dynamics of mammalian clathrin-mediated endocytosis. *PLoS Biol* **9**. doi:10.1371/journal.pbio.1000604

Tonikian R, Xin X, Toret CP, Gfeller D, Landgraf C, Panni S, Paoluzi S, Castagnoli L, Currell B, Seshagiri S, Yu H, Winsor B, Vidal M, Gerstein MB, Bader GD, Volkmer R, Cesareni G, Drubin DG, Kim PM, Sidhu SS, Boone C. 2009. Bayesian modeling of the yeast SH3 domain interactome predicts spatiotemporal dynamics of endocytosis proteins. *PLoS Biol* **7**. doi:10.1371/journal.pbio.1000218

Weinberg JS, Drubin DG. 2014. Regulation of Clathrin-mediated endocytosis by dynamic ubiquitination and deubiquitination. *Curr Biol* **24**:951–959. doi:10.1016/j.cub.2014.03.038

Figure 1: Quantitatively probing the budding yeast clathrin-mediated endocytosis

pathway through systematic imaging

(A) Illustration of key steps in the budding yeast CME pathway. Mechanisms of site

initiation, transition from the early, variable lifetime stages to the later, more regular

lifetime stages, and initiation of actin assembly remain poorly understood. Diagram

adapted from Tonikian et al., 2009. (B) Top: montage of Las17-GFP and Abp1-RFP

recruitment to an endocytic site viewed *en face* by live two-color TIRF microscopy of

budding yeast. Bottom: centroid position tracks in the x and y dimensions for the Las17-

GFP (green) and Abp1-RFP (magenta) spots depicted in the montage. • - beginning, x – end.

See Video 1. (C) Fluorescence intensity vs. time for Las17-GFP and Abp1-RFP spots

depicted in panel B. Timepoints of note and key measurements recorded are indicated by

dotted lines and arrows, respectively. (D – F) Box and whisker plots of the time from the

appearance of the indicated GFP-tagged query protein to the appearance of (D) Ede1-RFP,

(E) Sla1-mCherry, and (F) Abp1-RFP. Red lines are median values, boxes indicate

interquartile range, and whiskers indicate full range of the data.

Figure 1 – Figure Supplement 1: Lifetimes and relative fluorescence intensities of

GFP-tagged query proteins agree with prior results but reveal inherent variability

(A) Box and whisker plots of lifetimes of the indicated GFP-tagged query protein when

colocalized with Ede1-RFP (left), Sla1-mCherry (center), and Abp1-RFP (right). (B) Box and

whisker plots of maximum fluorescence intensity of the indicated GFP-tagged query

protein imaged with reference to Ede1-RFP (left), Sla1-mCherry (center), and Abp1-RFP

(right). Red lines are median values, boxes indicate interquartile range, and whiskers

indicate full range of the data.

Figure 2: Lifetime and maximum intensity are positively correlated for individual

later arriving CME proteins, but are less correlated for earlier arriving proteins

(A – C) Scatterplots of maximum intensity vs. lifetime for (A) Ede1-GFP, (B) Las17-GFP, and

(C) Sac6-GFP. Color indicates the normalized data density in the neighborhood around each

point. Red lines are linear fits to the data with the indicated R^2 values. (D) Summary of

linear fits to plots of maximum intensity vs. lifetime for the indicated GFP-tagged query

proteins. The bar height and color indicate the normalized slope and the R^2 of the fit,

respectively. The grayed area highlights proteins displaying similar behavior. Association

of query protein signal with signal of the reference protein Abp1-RFP was used to eliminate

spurious events. See Figure 2 Supplement 1 for all related scatter plots.

Figure 2 – Figure Supplement 1: Maximum intensity vs. lifetime plots for GFP-tagged

query proteins

Scatter plots of query protein maximum intensity vs. lifetime for GFP-tagged (A) Ede1, (B)

Syp1, (C) Sla2, (D) Pan1, (E) Las17, (F) Sla1, (G) Vrp1, (H) Bzz1, (I) Arc15, (J) Myo5, (K)

Sac6, (L) Ark1, and (M) Rvs167. Color indicates the normalized data density in the

neighborhood around each point. Red lines are linear fits to the data with the indicated R^2

value. Association of query protein signal with signal of the reference protein Abp1-RFP

was used to eliminate spurious events.

Figure 3: Correlations between reference protein abundance and query protein

lifetime provide further evidence for a regulatory transition point

(A – C) Scatterplots of the maximum intensities of Abp1-RFP vs. the lifetimes of (A) Ede1-

GFP, (B) Las17-GFP, and (C) Sac6-GFP. Colors indicate the normalized data density in the

neighborhood around each point. Red lines are linear fits to the data with the indicated R^2

value. (D) Summary of linear fits to plots of maximum Abp1-RFP intensity vs. lifetime for

the indicated GFP-tagged query proteins. The bar height and color indicate the normalized

slope and the R^2 of the fit, respectively. The grayed area highlights proteins displaying

similar behavior. See Figure 3 Supplement 1 for all related scatter plots.

Figure 3 – Figure Supplement 1: Abp1-RFP maximum intensity vs. query protein

lifetime plots for GFP-tagged query proteins

Scatter plots of Abp1-RFP maximum intensity vs. lifetime for GFP-tagged (A) Ede1, (B)

Syp1, (C) Sla2, (D) Pan1, (E) Las17, (F) Sla1, (G) Vrp1, (H) Bzz1, (I) Arc15, (J) Myo5, (K)

Sac6, (L) Ark1, and (M) Rvs167. Color indicates the normalized data density in the

neighborhood around each point. Red lines are linear fits to the data with the indicated R^2

value.

Figure 4: CME protein lifetimes are correlated amongst earlier arriving proteins, but

early protein lifetimes are not correlated with late protein lifetimes

(A – C) Scatterplots of the lifetimes of Abp1-RFP vs. the lifetimes of (A) Ede1-GFP, (B)

Las17-GFP, and (C) Sac6-GFP. Colors indicate the normalized data density in the

neighborhood around each point. Red lines are linear fits to the data with the indicated R^2

value. (D – F) Summaries of linear fits to plots of (D) Ede1-RFP, (E) Sla1-mCherry, and (F)

Abp1-RFP lifetime vs. lifetime for the indicated GFP-tagged query proteins. The bar height

and color indicate the normalized slope and the R^2 of the fit, respectively. See Figure

Supplements 1 - 3 for all related scatter plots.

Figure 4 – Figure Supplement 1: Ede1-RFP lifetime vs. query protein lifetime plots for

GFP-tagged query proteins.

Scatter plots of Ede1-RFP lifetime vs. query protein lifetime for GFP-tagged (A) Syp1, (B)

Hrr25 (3X-GFP tagged), (C) Pal1, and (D) Apl1. Color indicates the normalized data density

in the neighborhood around each point. Red lines are linear fits to the data with the

indicated R^2 value.

Figure 4 – Figure Supplement 2: Sla1-mCherry lifetime vs. query protein lifetime

plots for GFP-tagged query proteins.

Scatter plots of Sla1-mCherry lifetime vs. query protein lifetime for GFP-tagged (A) Ede1,

(B) Syp1, (C) Hrr25 (3X-GFP tagged), (D) Pal1, and (E) Apl1. Color indicates the normalized

data density in the neighborhood around each point. Red lines are linear fits to the data

with the indicated R^2 value.

Figure 4 – Figure Supplement 3: Abp1-RFP lifetime vs. query protein lifetime plots

for GFP-tagged query proteins.

Scatter plots of Abp1-RFP lifetime vs. query protein lifetime for GFP-tagged (A) Ede1, (B)

Syp1, (C) Sla2, (D) Pan1, (E) Las17, (F) Sla1, (G) Vrp1, (H) Bzz1, (I) Arc15, (J) Myo5, (K)

Sac6, (L) Ark1, and (M) Rvs167. Color indicates the normalized data density in the

neighborhood around each point. Red lines are linear fits to the data with the indicated R^2

value.

Figure 5: Proximity to sites of exocytosis and cell polarity signaling influences the

rate of endocytic site maturation.

(A) Montage from a maximum intensity-projected epifluorescence video of a cell

endogenously expressing Sla1-mCherry (magenta) and Ede1-GFP (Green). Times are

hours:minutes. See Video 2. (B) Maximum intensity projections of z-stacks (left) of

polarized and depolarized cells paired with circumferential kymographs around the

mother cortex from medial focal plane videos of the same cells (right) endogenously

expressing Sla1-mCherry (magenta) and Ede1-GFP (Green). See Video 3. (C) Quantification

of the ratio of the number of Sla1-mCherry sites to Ede1-GFP sites in mother cells from

maximum intensity projections of z-stacks of 60 polarized and 60 depolarized cells. A two-

tailed p value from a Mann-Whitney U test with the null hypothesis that the two ratios are

identical (U = 1) is displayed. The median and interquartile ranges are denoted with error

bars. (D) Percentage of 596 Ede1 patches from 59 polarized mother cells and 1034 Ede1

patches from 56 depolarized mother cells that persist throughout the duration of a 4-

minute video (persistent), that are present at the start or end of the video (partial), or that

assemble and disassemble within the interval of the video (turnover). Numbers indicate

the percentage of patches observed in each category. A two-tailed p value from a Chi-

Square test with the null hypothesis that the proportion distributions are identical (chi-

square = 1349, 2 degrees of freedom) is displayed.

Figure 5 – Figure Supplement 1: Proximity to sites of exocytosis and cell polarity signaling does not influence the rate of late steps in the endocytic pathway

(A) Maximum intensity projections of z-stacks (top) of polarized and depolarized cells paired with radial kymographs of individual endocytic events from medial focal plane epifluorescence videos of the same cells (bottom) endogenously expressing Sla1-GFP (green) and Abp1-RFP (magenta). Kymograph generation and analysis were carried out in mother cells. (B) Quantification of the ratio of the number of Abp1-RFP sites to Sla1-GFP sites from maximum intensity projections of z-stacks of 50 polarized and 50 depolarized cells. Analysis was carried out in mother cells. A two-tailed p value from a Mann-Whitney U test with the null hypothesis that the two ratios are identical (U = 1051) is displayed. The median and interquartile ranges are denoted with error bars. (C) Sla1-GFP and Abp1-RFP lifetimes measured in mother cells of 40 polarized cells and 40 depolarized cells (5 representative sites per cell). Two-tailed p values from Mann-Whitney U tests (U_{Sla1} = 19732, U_{Abp1} = 17429) with the null hypothesis that the lifetimes are identical are displayed. The median and interquartile ranges are denoted with error bars.

Figure 6: Endocytic site maturation rate is controlled by the presence of cargo.

(A) Maximum intensity projections of cells endogenously expressing polarity marker Bem1-GFP before and 5 minutes after 17-fold dilution into isotonic imaging media (control, left) or water (right). (B) Average ± standard deviation for Bem1-GFP intensity profiles from 20 (media, left) and 30 (water, right) cells before and after dilution into the indicated

media. Individual intensity profiles generated from 25 pixel-wide lines were normalized to the maximum value before dilution and to cell length. (C) Maximum intensity projections of a cell endogenously expressing Sla1-mCherry (magenta) and Ede1-GFP (green) before and 5 minutes after 17-fold dilution into water. (D) Quantification of the ratio of the number of Sla1-mCherry sites to Ede1-GFP sites from maximum intensity projections of z-stacks of 30 small-budded mother cells before and after osmotic shock. A two-tailed p value from a Mann-Whitney U test with the null hypothesis that the two ratios are identical (U = 7) is displayed. The median and interquartile ranges are denoted with error bars. (E) Circumferential kymographs around the mother cortex from medial focal plane videos of cells endogenously expressing Sla1-mCherry (magenta) and Ede1-GFP (Green) before and 5 minutes after 17-fold dilution into water. (F) Percentage of 280 Ede1 patches from 30 small-budded mother cells before and 594 Ede1 patches from 30 small-budded mother cells after osmotic shock that persist throughout the duration of a 4-minute video (persistent), are present at the start or end of the video (partial), or assemble and disassemble within the interval of the video (turnover). Numbers indicate the percentage of patches observed in each category. A two-tailed p value from a Chi-Square test with the null hypothesis that the proportions distributions are identical (chi-square = 543.1, 2 degrees of freedom) is displayed. (G) Maximum intensity projections of an *erg6Δ* cell endogenously expressing Sla1-mCherry (magenta) and Ede1-GFP (green) before and 10 minutes after treatment with Brefeldin A. (H) Quantification of the ratio of the number of Sla1-mCherry sites to Ede1-GFP sites from maximum intensity projections of z-stacks of 24 large-budded, *erg6Δ* mother cells before and after BFA treatment. A two-tailed p value from a Mann-Whitney U test with the null hypothesis that the two ratios are identical (U =

25.5) is displayed. The median and interquartile ranges are denoted with error bars. (I)

Circumferential kymographs around the mother cortex from medial focal plane videos of

erg6Δ cells endogenously expressing Sla1-mCherry (magenta) and Ede1-GFP (Green)

before and 10 minutes after treatment with BFA. (F) Percentage of 277 Ede1 patches from

18 large-budded, *erg6Δ* mother cells before and 266 Ede1 patches from 22 large-budded,

erg6Δ mother cells after BFA treatment that persist throughout the duration of a 4-minute

video (persistent), are present at the start or end of the video (partial), or assemble and

disassemble within the interval of the video (turnover). Numbers indicate the percentage

of patches observed in each category. A two-tailed p value from a Chi-Square test with the

null hypothesis that the proportions distributions are identical (chi-square = 1122, 2

degrees of freedom) is displayed.

Figure 6, figure supplement 1: Acute loss of cell polarity does not accelerate late

steps of the endocytic pathway; Brefeldin A does not affect endocytosis in insensitive

cells

(A) Maximum intensity projections of cells endogenously expressing GFP-Sec4 before and 5

minutes after 17-fold dilution into isotonic imaging media (control, left) or water (right).

(B) Maximum intensity projections of a cell endogenously expressing Sla1-GFP (Green) and

Abp1-RFP (magenta) before and 5 minutes after 17-fold dilution into water. (C) Radial

kymographs of individual endocytic events in small-budded mother cells from medial focal

plane videos of cells endogenously expressing Sla1-GFP (green) and Abp1-RFP (magenta)

before and 5 minutes after 17-fold dilution into water. (D) Quantification of the ratio of the

number of Abp1-RFP sites to Sla1-GFP sites from maximum intensity projections of z-stacks of 30 small-budded mother cells before and after osmotic shock. A two-tailed p value from a Mann-Whitney U test with the null hypothesis that the two ratios are identical (U = 441.5) is displayed. The median and interquartile ranges are denoted with error bars. (E) Sla1-GFP and Abp1-RFP lifetimes measured in 30 small-budded mother cells prior to osmotic shock and 30 cells after osmotic shock (5 representative sites per cell). Two-tailed p values from Mann-Whiney U tests with the null hypothesis that the lifetimes are identical (U_{Sla1} = 1678, U_{Abp1} = 2998) are displayed. The median and interquartile ranges are denoted with error bars. (F) Maximum intensity projections of a wild-type cell endogenously expressing Sla1-mCherry (magenta) and Ede1-GFP (green) before and 10 minutes after treatment with Brefeldin A. (G) Quantification of the ratio of the number of Sla1-mCherry sites to Ede1-GFP sites from maximum intensity projections of z-stacks of 20 large-budded mother cells before and after BFA treatment. A two-tailed p value from a Mann-Whitney U test with the null hypothesis that the two ratios are identical (U = 132.5) is displayed. The median and interquartile ranges are denoted with error bars. (H) Circumferential kymographs around the mother cortex from medial focal plane videos of cells endogenously expressing Sla1-mCherry (magenta) and Ede1-GFP (Green) before and 10 minutes after treatment with BFA. (I) Percentage of 375 Ede1 patches from 22 large-budded mother cells before and 477 Ede1 patches from 24 large-budded mother cells after BFA treatment that persist throughout the duration of a 4-minute video (persistent), are present at the start or end of the video (partial), or assemble and disassemble within the interval of the video (turnover). Numbers indicate the percentage of patches observed in each category. A two-tailed p value from a Chi-Square test with the null hypothesis that the

proportions distributions are identical (chi-square = 10.77, 2 degrees of freedom) is

displayed.

Figure 6, figure supplement 2: Testing whether Hrr25 regulates endocytic

maturation

(A) Growth of cells of the indicated genotypes on YPD and auxin-containing plates at 18°C

and 25°C. Photos are representative of results from three separate experiments. (B)

Immunoblot illustrating degradation kinetics of auxin inducible degron (AID)-tagged Hrr25

upon treatment with auxin. Hrr25-AID was detected via a V5 epitope tag. Pgk1 was probed

as a loading control. (C) Quantification of relative Hrr25 levels as a function of treatment

time in auxin from three independent experiments. Hrr25 band intensities were

normalized to the Pgk1 loading control and expressed as a fraction of the preAuxin signal.

(D) Percentage of Ede1 patches in 30 small-budded mother cells of the indicated genotype

treated as indicated with auxin for 30 min. that persist throughout the duration of a 4-

minute video (persistent), are present at the start or end of the video (partial), or assemble

and disassemble within the interval of the video (turnover). Numbers indicate the

percentage of patches observed in each category. Two-tailed p values from Chi-Square tests

with the null hypothesis that the proportions distributions are identical are displayed. (E)

Immunoblot illustrating overexpression of a 13myc-tagged copy of Hrr25 upon treatment

with β-estradiol for 2 hours. Overexpressed Hrr25 was detected via the 13myc epitope tag.

Endogenous Hrr25 and Pgk1 were probed directly as loading controls. (F) Percentage of

Ede1 patches in 30 small-budded mother cells of the indicated genotype treated as

indicated with β-estradiol for 2 hrs. analyzed as in panel D. Numbers indicate the

percentage of patches observed in each category. Two-tailed p values from Chi-Square tests with the null hypothesis that the proportions distributions are identical are displayed.

Figure 7: Defining molecular components of the early and late steps of the endocytic pathway

(A) Maximum intensity projections of epifluorescence z-stacks of clusters of cells endogenously expressing Sla1-mCherry (magenta) and Syp1-GFP (green, first row), Sla2-GFP (green, second row), Ent2-GFP (green, third row), and Pan1-GFP (green, last row). Individual channels are shown in gray scale at left. (B) Quantification of the ratio of the number of Sla1-mCherry sites to the number of Syp1-GFP (top row), Sla2-GFP (second row), Ent2-GFP (third row), and Pan1-GFP (last row) sites from maximum intensity projections of z-stacks of 30 small-budded mother cells before and after osmotic shock. Two-tailed p values from Mann-Whitney U test with the null hypothesis that the two ratios are identical are displayed ($U_{Syp1} = 0$, $U_{Sla2} = 0$, $U_{Ent2} = 0$, $U_{Pan1} = 448$). The median and interquartile ranges are denoted with error bars. (C) Maximum intensity projections of epifluorescence z-stacks of clusters of "7Δ" cells endogenously expressing Sla1-mCherry (magenta) and Sla2-GFP (green). Individual channels are shown in gray scale at left. Yellow arrowheads indicate highlight low concentration of Sla1-mCherry sites in a small bud, where late endocytic sites would normally be plentiful. (D) Quantification of the ratio of the number of Sla1-mCherry sites to the number of Sla2-GFP sites in small budded and large budded wild-type and 7Δ cells (n = 30 cells per category). Numbers are p values from Kruskal-Wallis tests followed by Dunn's multiple comparisons test. The median and interquartile ranges are denoted with error bars.

Figure 7, figure supplement 1: Defining molecular components of the early and late steps of the endocytic pathway

(A) Maximum intensity projections of spinning disc confocal z-stacks of clusters of cells endogenously expressing Sla1-mCherry (magenta) and Hrr25-3xGFP (green, top), Apl1-GFP (green, second row), Pal1-GFP (green, third row), or Ent1-GFP (green, last row). Individual channels are shown in gray scale at left. Images are representative of at least three separate experiments. (B) Maximum intensity projections of epifluorescence z-stacks of clusters of 7Δ cells endogenously expressing Sla1-GFP (green) and Abp1-mCherry (magenta). Individual channels are shown in gray scale at left. (C) Quantification of the ratio of the number of Abp1-mCherry sites to the number of Sla1-GFP sites in small budded and large budded wild-type and 7Δ cells (n = 30 cells per category). Numbers are p values from Kruskal-Wallis tests followed by Dunn's multiple comparisons test. The median and interquartile ranges are denoted with error bars.

Figure 8: Cargo regulates the transition between two quantitatively distinguishable phases of endocytosis.

(A) Revised timeline for the clathrin-mediated endocytosis pathway indicating the proposed temporal location of the cargo checkpoint. Proteins upstream and downstream of the checkpoint are listed under each respective phase. (B) Schematic cartoon illustrating the proposed mechanism leading to polarized endocytosis in budding yeast. In highly polarized cells, endocytic cargos (black arrows) are delivered primarily to the growing bud by exocytosis (gray arrows), locally accelerating the transition from the initiation and

cargo-collection phase of CME (green spots) into the internalization phase (magenta spots). As exocytosis becomes depolarized later in the cell cycle, cargos are more plentiful across the plasma membrane, leading to global acceleration of maturation through the cargo checkpoint.

Video 1: Related to Figure 1B

TIRFM movie of several yeast cells endogenously expressing Las17-GFP (green) and Abp1-RFP (magenta). Each Las17 punctum is punctuated by a burst of Abp1 signal. Frames are separated by 1 second and played back at 15 frames per second (fps).

Video 2: Related to Figure 5A

Time-lapse of maximum intensity projections from epifliorescence z-stacks of a cell endogenously expressing Sla1-mCherry (magenta) and Ede1-GFP (green). A dynamic cycle of polarization/depolarization is apparent for the magenta Sla1 sites, but less apparent for the green Ede1 sites. Frames are separated by 10 minutes and played back at 10 fps.

Video 3: Related to Figure 5B

Epifluorescence movies in the media focal plane of the polarized and depolarized cells displayed in Figure 5B. The cells endogenously express Sla1-mCherry (magenta) and Ede1-GFP (green). Persistent Ede1 sites are present in the mother of the polarized cell (left), but not in the mother of the depolarized cell (right). Frames are separated by 2 seconds and played back at 15 fps. Circumferential kymographs in Figure 5B were generated from these movies.

Figure 1

Figure 1 - Figure Supplement 1

A

B

Figure 2

Figure 2 - Figure Supplement 1

Figure 3

Figure 3 - Figure Supplement 1

Figure 4

Figure 4 - Figure Supplement 1

Figure 4 - Figure Supplement 2

Figure 5

A

B Sla1-mCherry Ede1-GFP

C

Sla1/Ede1 site ratio

p < 0.0001

Polarized   Depolarized

D

Percent of Ede1 sites (%)

p < 0.0001

0.3%

	Persistant
21.3	
34.9	Partial
58.7	
20	Turnover
64.8	

Polarized   Depolarized

Figure 5 - Figure Supplement 1

Figure 6

A Control: Media Wash-in   Water Wash-in
Bem1-GFP
Before After   Before After

B Before Wash-in   After Wash-in

C Sla1-mCherry Ede1-GFP
Before   After
Osmotic Shock

D p < 0.0001
Before Water   After Water

E Before Water   After Water
Time   4 Minutes

F p < 0.0001   0.2%
Persistant / Partial / Turnover
Before Water   After Water

G Sla1-mCherry Ede1-GFP
erg6Δ
Before   After
Brefeldin A Treatment

H p < 0.0001
Before BFA   After BFA

I Before BFA   After BFA
Time   4 Minutes

J p < 0.0001   1.41%
Persistant / Partial / Turnover
Before BFA   After BFA

Figure 6 - Figure Supplement 1

A Control: Media Wash-in   Water Wash-in
GFP-Sec4
Before   After   Before   After   2μm

B   Water Wash-in
Sla1-GFP Abp1-mRFP
2μm
Before   After

C   Water Wash-in
Sla1-GFP
Abp1-mRFP
Merge   1.5 μm   1.5 μm
1 Minute   1 Minute
Before   After

D
ns
p = 0.9036
Abp1/Sla1 site ratio
Before Water   After Water

E   Sla1-GFP Lifetimes   Abp1-mRFP Lifetimes
p < 0.0001   p < 0.0001
Lifetime (s)   Lifetime (s)
Before Water   After Water   Before Water   After Water

F   Sla1-mCherry Ede1-GFP
2μm
Before   After
Brefeldin A Treatment

G
p = 0.0261
Sla1/Ede1 site ratio
Before BFA   After BFA

H
Distance along Mother cortex
2μm
Time   4 Minutes
Before BFA   After BFA

I   p = 0.0046
0.3%   1.1%
Percent of Ede1 sites (%)
43.3   46.6
56.5   52.3
Before BFA   After BFA
Persistant
Partial
Turnover

Figure 6 - Figure Supplement 2

Figure 7

A

Syp1-GFP   Sla1-mCherry   Merge

5μm

Sla2-GFP

Ent2-GFP

Pan1-GFP

B

p < 0.0001

Sla1/Syp1 site ratio

Polarized   Depolarized

p < 0.0001

Sla1/Sla2 site ratio

Polarized   Depolarized

p < 0.0001

Sla1/Ent2 site ratio

Polarized   Depolarized

p = 0.9795

Sla1/Pan1 site ratio

Polarized   Depolarized

C

Sla2-GFP   Sla1-mCherry   Merge

5μm

pal1Δ, pal2Δ, syp1Δ, ede1Δ, apl1Δ, yap1801Δ, yap1802Δ (7Δ)

D

p < 0.0001   p = 0.3246

Sla1/Sla2 site ratio

WT, Sm Bud   WT, Lg Bud   7Δ, Sm Bud   7Δ, Lg Bud

Figure 7 - Figure Supplement 1

A

Hrr25-3xGFP Sla1-mCherry Merge

5µm

Apl1-GFP

Pal1-GFP

Ent1-GFP

B

Sla1-GFP Abp1-mCherry Merge

5µm

pal1Δ, pal2Δ, syp1Δ, ede1Δ, apl1Δ, yap1801Δ, yap1802Δ (7Δ)

C

p > 0.9999

p = 0.6037

Abp1/Sla1 site ratio

WT, Sm Bud WT, Lg Bud 7Δ, Sm Bud 7Δ, Lg Bud

Figure 8

A

B

● Initiated endocytic sites awaiting cargo	● Internalizing endocytic sites

Supplementary table 1: strains used in this study

Name	Genotype	Source
DDY1102	*MATa/MATα his3-Δ200/his3-Δ200, leu2-3, 112/leu2-3, 112, ura3-52/ura3-52, ade2-1/ADE2, lys2-801/LYS2*	Drubin Lab
DDY5713	*MATa his3-Δ200, leu2-3, 112, ura3-52, ABP1-mRFP::HIS3, SLA1-GFP::KanMX*	This study
DDY5714	*MATa his3-Δ200, leu2-3, 112, ura3-52, ABP1-mRFP::HIS3, Bzz1-GFP::HIS3*	This study
DDY5715	*MATa his3-Δ200, leu2-3, 112, ura3-52, ABP1-mRFP::HIS3, ARC15-GFP::KanMX*	This study
DDY5716	*MATa his3-Δ200, leu2-3, 112, ura3-52, ABP1-mRFP::HIS3, RVS167-GFP::HIS3*	This study
DDY5717	*MATa his3-Δ200, leu2-3, 112, ura3-52, ABP1-mRFP::HIS3, SLA2-GFP::HIS3*	This study
DDY5718	*MATa his3-Δ200, leu2-3, 112, ura3-52, ABP1-mRFP::HIS3, ARK1-GFP::KanMX*	This study
DDY5719	*MATa his3-Δ200, leu2-3, 112, ura3-52, ABP1-mRFP::HIS3, SAC6-GFP::HIS3*	This study
DDY5720	*MATa his3-Δ200, leu2-3, 112, ura3-52, bar1Δ::NatR, EDE1-mRFP::KanMX, PAL1-GFP::HIS3*	This study
DDY5721	*MATa his3-Δ200, leu2-3, 112, ura3-52, bar1Δ::NatR, EDE1-mRFP::HIS3, SYP1-GFP::KanMX*	This study
DDY5722	*MATa his3-Δ200, leu2-3, 112, ura3-52, bar1Δ::NatR, EDE1-mRFP::KanMX, Hrr25-3xGFP::HIS3*	This study
DDY5723	*MATa his3-Δ200, leu2-3, 112, ura3-52, bar1Δ::NatR, SLA1-mCherry::HIS3, PAL1-GFP::HIS3*	This study
DDY5724	*MATa his3-Δ200, leu2-3, 112, ura3-52, bar1Δ::NatR, SLA1-mCherry::HIS3, APL1-GFP::HIS3*	This study
DDY5725	*MATa his3-Δ200, leu2-3, 112, ura3-52, bar1Δ::NatR, SLA1-mCherry::HIS3, EDE1-GFP::HIS3*	This study
DDY5726	*MATa his3-Δ200, leu2-3, 112, ura3-52, bar1Δ::NatR, SLA1-mCherry::HIS3, SYP1-GFP::KanMX*	This study
DDY5727	*MATa his3-Δ200, leu2-3, 112, ura3-52, bar1Δ::NatR, SLA1-mCherry::HIS3, Hrr25-3xGFP::HIS3*	This study
DDY3293	*MATα his3-Δ200, leu2-3, 112, ura3-52, ABP1-mRFP::HIS3, LAS17-GFP::HIS3*	Drubin lab
DDY5617	*MATa his3-Δ200, leu2-3, 112, ura3-52, ABP1-mRFP::HIS3, LAS17-GFP::HIS3*	Drubin lab (Sun et al., 2017)
DDY3288	*MATa his3-Δ200, leu2-3, 112, ura3-52, ABP1-mRFP::HIS3, MYO5-GFP::URA3::KanMX*	Drubin lab
DDY5640	*MATa his3-Δ200, leu2-3, 112, ura3-52, ABP1-mRFP::HygMX, MYO5-GFP::HIS3*	Drubin lab (Sun et al., 2017)
DDY3063	*MATα his3-Δ200, leu2-3, 112, ura3-52, ABP1-mRFP::HIS3, Pan1-GFP::HIS3*	Drubin lab

DDY3314	*MATa his3-Δ200, leu2-3, 112, ura3-52, ABP1-mRFP::HIS3, Pan1-GFP::KanMX*	Drubin lab
DDY3867	*MATα his3-Δ200, leu2-3, 112, ura3-52, ABP1-mRFP::HIS3, SYP1-GFP::KanMX*	Drubin lab (Stimpson et al., 2009)
DDY3868	*MATa his3-Δ200, leu2-3, 112, ura3-52, ABP1-mRFP::HIS3, Ede1-GFP::HIS3*	Drubin lab (Stimpson et al., 2009)
DDY3061	*MATα his3-Δ200, leu2-3, 112, ura3-52, ABP1-mRFP::HIS3, CLC1-GFP::HIS3*	Drubin lab
DDY5635	*MATa his3-Δ200, leu2-3, 112, ura3-52, ABP1-mRFP::HygMX, VRP1-GFP::HIS3*	Drubin lab (Sun et al., 2017)
DDY3837	*MATα, his3-Δ200, leu2-3, 112, ura3-52, bar1Δ::LEU2, EDE1-mRFP::HIS3, APL1-GFP::HIS3*	Drubin lab
DDY5728	*MATα, his3Δ1, leu2Δ0, lys2Δ0, ura3Δ0, GFP-SEC4::URA3*	Bretscher lab (Donovan and Bretscher, 2015)
DDY4884	*MATα, his3-Δ200, leu2-3, 112, ura3-52, BEM1-GFP::HIS3*	Drubin Lab
DDY4083	*MATα, his3-Δ200, leu2-3, 112, ura3-52, SLA1-mCherry::HIS3, SLA2-GFP::HIS3*	Drubin lab (Carroll et al., 2012)
DDY4084	*MATα, his3-Δ200, leu2-3, 112, ura3-52, SLA1-mCherry::HIS3, ENT1-GFP::HIS3*	Drubin lab (Carroll et al., 2012)
DDY4085	*MATα, his3-Δ200, leu2-3, 112, ura3-52, SLA1-mCherry::HIS3, ENT2-GFP::HIS3*	Drubin lab (Carroll et al., 2012)
DDY4834	*MATα, his3-Δ200, leu2-3, 112, ura3-52, SLA1-mCherry::HIS3, PAN1-GFP::HIS3*	Drubin lab (Sun et al., 2015)
DDY5734	*MATα, his3-Δ200, leu2-3, 112, ura3-52, EDE1-GFP::HIS3, SLA1-mCherry::HIS3, hrr25-AID::KanMX, TIR1::LEU2*	This study
DDY5735	*MATα, his3-Δ200, leu2-3, 112, ura3-52, EDE1-GFP::HIS3, SLA1-mCherry::HIS3, hrr25-AID::KanMX*	This study
DDY5736	*MATα, his3-Δ200, leu2-3, 112, ura3-52, EDE1-GFP::HIS3, SLA1-mCherry::HIS3, TIR1::LEU2*	This study

DDY5737	*MATa, his3-Δ200, leu2-3, 112, ura3-52, GEV::URA3, pGAL-HRR25-13Myc::LEU2, EDE1-GFP::HIS3, SLA1-mCherry::HIS3, bar1Δ::NatR*	This study
DDY5738	*MATa, his3-Δ200, leu2-3, 112, ura3-52, pGAL-HRR25-13Myc::LEU2, EDE1-GFP::HIS3, SLA1-mCherry::HIS3, bar1Δ::NatR*	This study
DDY5739	*MATa, his3-Δ200, leu2-3, 112, ura3-52, GEV::URA3, EDE1-GFP::HIS3, SLA1-mCherry::HIS3, bar1Δ::NatR*	This study
DDY5730	*MATa, his3Δ200, leu2-3,112, ura3-52, lys2-801, SLA1-EGFP::HIS3MX6, ABP1-mCherry::kanMX4*	Kaksonen lab (Brach et al. 2014)
DDY5731	*MATa, his3Δ200, leu2-3,112, ura3-52, lys2-801, SLA2-EGFP::natNT2, Sla1-mCherry::kanMX4*	Kaksonen lab (Brach et al. 2014)
DDY5732	*MATa, his3Δ200, leu2-3,112, ura3-52, lys2-801, pal1::natNT2, pal2::natNT2, syp1::hphNT1, ede1::natNT2, apl1::natNT2, yap1801::natNT2, yap1802::natNT2, SLA1-EGFP::HIS3MX6, ABP1-mCherry::kanMX4*	Kaksonen lab (Brach et al. 2014)
DDY5733	*MATa, his3Δ200, leu2-3,112, ura3-52, lys2-801, pal1::natNT2, pal2::natNT2, syp1::hphNT1, ede1::natNT2, apl1::natNT2, yap1801::natNT2, yap1802::natNT2, SLA2-EGFP::HIS3MX6, SLA1-mCherry::kanMX4*	Kaksonen lab (Brach et al. 2014)

Supplementary table 2: Time to arrival for GFP-tagged query proteins colocalized with Ede1-RFP. Related to Figure 1D

Query protein	N	Median time to arrival (s)	Mean (s)	St. Dev (s)
Apl1	329	-7.0	-7.7	14.3
Hrr25	291	-1.0	-1.9	9.3
Pal1	207	-1.0	-2.6	12.0
Syp1	351	2.0	1.7	9.9

Supplementary table 3: Time to arrival for GFP-tagged query proteins colocalized with Sla1-mCherry. Related to Figure 1E

Query protein	N	Median time to arrival (s)	Mean (s)	St. Dev (s)
Apl1	728	3.0	3.8	12.1
Ede1	260	29.0	29.6	26.1
Hrr25	444	11.5	11.3	16.5
Pal1	229	3.0	6.0	15.7
Syp1	211	15.0	17.4	15.3

Supplementary table 4: Time to arrival for GFP-tagged query proteins colocalized with Abp1-RFP. Related to Figure 1F

Query protein	N	Median time to arrival (s)	Mean (s)	St. Dev (s)
Arc15	1878	2.0	2.3	2.5
Ark1	260	-2.0	-2.3	2.2
Bzz1	2257	5.0	5.7	3.9
Ede1	252	42.5	48.0	27.9
Las17	582	14.0	14.1	6.9
Myo5	1202	1.0	1.5	1.8
Pan1	158	15.0	16.3	6.7
Rvs167	1802	-3.0	-3.1	2.0
Sac6	2512	0.0	0.0	1.1
Sla1	316	13.0	14.0	5.9
Sla2	274	18.0	18.9	9.9
Syp1	315	21.0	24.3	17.6
Vrp1	1604	9.0	9.0	4.4

Supplementary table 5: Maximum fluorescence intensity of GFP-tagged query proteins colocalized with Ede1-RFP. Related to Figure 1, Figure supplement 1A

Query protein	N	Median maximum intensity (au)	Mean (au)	St. Dev (au)
Apl1	329	517	565	186
Hrr25	291	1083	1199	542
Pal1	207	462	483	160
Syp1	351	694	868	508

Supplementary table 6: Maximum fluorescence intensity of GFP-tagged query proteins colocalized with Sla1-mCherry. Related to Figure 1, Figure supplement 1A

Query protein	N	Median maximum intensity (au)	Mean (au)	St. Dev (au)
Apl1	728	466	486	140
Ede1	260	1843	2183	1434
Hrr25	444	984	1052	441
Pal1	229	349	363	97
Syp1	211	419	461	183

Supplementary table 7: Maximum fluorescence intensity of GFP-tagged query proteins colocalized with Abp1-RFP. Related to Figure 1, Figure supplement 1A

Query protein	N	Median maximum intensity (au)	Mean (au)	St. Dev (au)
Arc15	1878	3358	4291	3116
Ark1	260	457	465	142
Bzz1	2257	1241	1342	539
Ede1	252	1972	2292	1364
Las17	582	851	1012	615
Myo5	1202	3168	3743	2457
Pan1	158	1196	1479	896
Rvs167	1802	3378	3620	1558
Sac6	2512	10954	11916	7004
Sla1	316	2356	2749	1717
Sla2	274	1144	1399	687
Syp1	315	899	979	405
Vrp1	1604	1183	1329	681

Supplementary table 8: Lifetimes of GFP-tagged query proteins colocalized with Ede1-RFP. Related to Figure 1, Figure supplement 1B

Query protein	N	Median lifetime (s)	Mean lifetime (s)	St. Dev (s)	Coeff. of variation
Apl1	329	23.0	25.7	13.9	0.54
Hrr25	291	25.0	27.9	16.0	0.57
Pal1	207	24.0	27.6	16.5	0.60
Syp1	351	26.0	28.8	16.2	0.56

Supplementary table 9: Lifetimes of GFP-tagged query proteins colocalized with Sla1-mCherry. Related to Figure 1, Figure supplement 1B

Query protein	N	Median lifetime (s)	Mean lifetime (s)	St. Dev (s)	Coeff. of variation
Apl1	728	29.0	30.0	13.7	0.46
Ede1	260	48.5	49.5	27.4	0.55
Hrr25	444	28.0	30.5	15.8	0.52
Pal1	229	27.0	29.9	15.5	0.52
Syp1	211	30.0	33.3	17.2	0.52

Supplementary table 10: Lifetimes of GFP-tagged query proteins colocalized with Sla1-mCherry. Related to Figure 1, Figure supplement 1B

Query protein	N	Median lifetime (s)	Mean lifetime (s)	St. Dev (s)	Coeff. of variation
Arc15	1878	13.0	14.0	4.2	0.3
Ark1	260	9.0	9.9	3.0	0.3
Bzz1	2257	15.0	15.3	4.8	0.32
Ede1	252	49.0	54.2	26.7	0.49
Las17	582	24.0	24.4	8.3	0.34
Myo5	1202	12.0	12.3	3.6	0.29
Pan1	158	21.0	22.0	7.9	0.36
Rvs167	1802	8.0	9.1	2.9	0.32
Sac6	2512	11.0	11.3	2.5	0.22
Sla1	316	21.0	22.3	7.2	0.32
Sla2	274	26.0	28.0	11.2	0.40
Syp1	315	28.0	31.6	16.2	0.51
Vrp1	1604	18.0	18.6	5.3	0.29

Supplementary table 11: Fit statistics from linear fits of intensity vs. lifetime plots for GFP-tagged query proteins. Upper and lower bounds indicate the 95% confidence interval. Related to Figure 2D

Query protein	N	Slope	Slope upper bound	Slope lower bound	R^2	R^2 upper bound	R^2 lower bound
Arc15	1878	0.86	0.73	0.98	0.09	0.06	0.11
Ark1	260	0.32	0.21	0.42	0.12	0.05	0.20
Bzz1	2257	0.59	0.54	0.64	0.19	0.16	0.22
Ede1	252	0.29	0.14	0.45	0.05	0.01	0.12
Las17	582	0.97	0.82	1.12	0.21	0.16	0.28
Myo5	1202	1.13	1.00	1.26	0.19	0.15	0.23
Pan1	158	1.14	0.88	1.39	0.33	0.21	0.45
Rvs167	1802	0.37	0.31	0.43	0.08	0.06	0.11
Sac6	2512	0.91	0.81	1.02	0.10	0.08	0.13
Sla1	316	1.07	0.86	1.27	0.25	0.17	0.33
Sla2	274	0.73	0.59	0.88	0.28	0.19	0.37
Syp1	315	0.35	0.27	0.43	0.20	0.13	0.28
Vrp1	1604	0.96	0.88	1.04	0.24	0.21	0.28

Supplementary table 12: Fit statistics from linear fits of Abp1-RFP intensity vs. lifetime plots for GFP-tagged query proteins. Upper and lower bounds indicate the 95% confidence interval. Related to Figure 3D

Query protein	N	Slope	Slope upper bound	Slope lower bound	R^2	R^2 upper bound	R^2 lower bound
Arc15	1878	0.65	0.54	0.75	0.07	0.05	0.10
Ark1	260	0.19	0.04	0.34	0.02	0.00	0.07
Bzz1	2257	0.46	0.39	0.52	0.08	0.06	0.10
Ede1	252	-0.06	-0.24	0.12	0.00	0.03	0.01
Las17	582	0.96	0.76	1.16	0.13	0.08	0.18
Myo5	1202	0.95	0.79	1.11	0.10	0.07	0.14
Pan1	158	0.73	0.47	1.00	0.16	0.07	0.27
Rvs167	1802	0.28	0.22	0.35	0.04	0.02	0.06
Sac6	2512	0.70	0.60	0.80	0.07	0.05	0.09
Sla1	316	0.74	0.49	0.98	0.10	0.05	0.17
Sla2	274	0.50	0.31	0.70	0.09	0.03	0.16
Syp1	315	0.16	0.01	0.30	0.01	0.00	0.05
Vrp1	1604	0.59	0.49	0.70	0.07	0.05	0.09

Supplementary table 13: Fit statistics from linear fits of Ede1-RFP lifetime vs. lifetime plots for GFP-tagged query proteins. Upper and lower bounds indicate the 95% confidence interval. Related to Figure 4D

Query protein	N	Slope	Slope upper bound	Slope lower bound	R^2	R^2 upper bound	R^2 lower bound
Apl1	329	0.45	0.35	0.54	0.21	0.14	0.30
Hrr25	291	0.71	0.64	0.79	0.55	0.47	0.62
Pal1	207	0.64	0.53	0.75	0.39	0.29	0.49
Syp1	351	0.65	0.55	0.74	0.35	0.27	0.43

Supplementary table 14: Fit statistics from linear fits of Sla1-mCherry lifetime vs. lifetime plots for GFP-tagged query proteins. Upper and lower bounds indicate the 95% confidence interval. Related to Figure 4E

Query protein	N	Slope	Slope upper bound	Slope lower bound	R^2	R^2 upper bound	R^2 lower bound
Apl1	728	0.31	0.27	0.36	0.19	0.14	0.24
Ede1	260	0.17	0.09	0.25	0.06	0.02	0.13
Hrr25	444	0.09	0.04	0.14	0.03	0.00	0.06
Pal1	229	0.14	0.05	0.23	0.04	0.01	0.10
Syp1	211	0.20	0.13	0.28	0.12	0.05	0.21

Supplementary table 15: Fit statistics from linear fits of Abp1-RFP lifetime vs. lifetime plots for GFP-tagged query proteins. Upper and lower bounds indicate the 95% confidence interval. Related to Figure 4F

Query protein	N	Slope	Slope upper bound	Slope lower bound	R^2	R^2 upper bound	R^2 lower bound
Arc15	1878	0.66	0.63	0.69	0.59	0.56	0.61
Ark1	260	0.42	0.34	0.51	0.27	0.18	0.36
Bzz1	2257	0.23	0.20	0.26	0.11	0.09	0.14
Ede1	252	0.08	0.02	0.14	0.03	0.00	0.08
Las17	582	0.40	0.34	0.46	0.24	0.19	0.31
Myo5	1202	0.63	0.59	0.67	0.42	0.38	0.46
Pan1	158	0.24	0.13	0.35	0.10	0.03	0.21
Rvs167	1802	0.21	0.19	0.24	0.15	0.12	0.18
Sac6	2512	0.83	0.81	0.85	0.71	0.69	0.73
Sla1	316	0.24	0.16	0.33	0.09	0.04	0.16
Sla2	274	0.19	0.11	0.27	0.08	0.03	0.15
Syp1	315	-0.01	-0.07	0.05	0.00	0.02	0.01
Vrp1	1604	0.36	0.32	0.40	0.18	0.14	0.21

Supplemental References

Brach T, Godlee C, Moeller-Hansen I, Boeke D, Kaksonen M. 2014. The initiation of clathrin-mediated endocytosis is mechanistically highly flexible. *Curr Biol* **24**:548–54. doi:10.1016/j.cub.2014.01.048

Carroll SY, Stimpson HEM, Weinberg J, Toret CP, Sun Y, Drubin DG. 2012. Analysis of yeast endocytic site formation and maturation through a regulatory transition point. *Mol Biol Cell* **23**:657–668. doi:10.1091/mbc.E11-02-0108

Donovan KW, Bretscher a. 2015. Tracking individual secretory vesicles during exocytosis reveals an ordered and regulated process. *J Cell Biol* 1–14. doi:10.1083/jcb.201501118

Stimpson HEM, Toret CP, Cheng AT, Pauly BS, Drubin DG. 2009. Early-Arriving Syp1p and Ede1p Function in Endocytic Site Placement and Formation in Budding Yeast. *Mol Biol Cell* **20**:4640–4651. doi:10.1091/mbc.E09

Sun Y, Leong NT, Jiang T, Tangara A, Darzacq X, Drubin DG. 2017. Switch-like Arp2/3 activation upon WASP and WIP recruitment to an apparent threshold level by multivalent linker proteins in vivo. *Elife* **6**:1–25. doi:10.7554/elife.29140

Sun Y, Leong NT, Wong T, Drubin DG. 2015. A Pan1/End3/Sla1 complex links Arp2/3-mediated actin assembly to sites of clathrin-mediated endocytosis. *Mol Biol Cell* **26**:3841–3856. doi:10.1091/mbc.E15-04-0252