Boosting Brain Signal Variability Underlies Liberal Shifts in Decision Bias

Niels A. Kloosterman[1,2*], Julian Q. Kosciessa[1,2], Ulman Lindenberger[1,2], Johannes Jacobus Fahrenfort[3], Douglas D. Garrett[1,2]

[1]Max Planck UCL Centre for Computational Psychiatry and Ageing Research and [2]Center for Lifespan Psychology, Max Planck Institute for Human Development, Lentzeallee 94, 14195 Berlin, Germany

[3]Department of Experimental and Applied Psychology, Vrije Universiteit, van der Boechorststraat 1, 1081 BT Amsterdam, The Netherlands

[4]Department of Psychology, University of Amsterdam, The Netherlands;

*Correspondence: Niels Kloosterman

Email: kloosterman@mpib-berlin.mpg.de

Author ORCIDs

Niels A Kloosterman http://orcid.org/0000-0002-1134-7996

Julian Kosciessa https://orcid.org/0000-0002-4553-2794

Ulman Lindenberger https://orcid.org/0000-0001-8428-6453

Johannes Jacobus Fahrenfort http://orcid.org/0000-0002-9025-3436

Douglas D Garrett https://orcid.org/0000-0002-0629-7672

Classification

Biological Sciences / Neuroscience

Keywords

Brain signal variability, decision bias, perceptual decision making, signal detection theory, cognitive flexibility

Author Contributions

Niels Kloosterman, Conceptualization, Data curation, Software, Formal analysis, Investigation, Visualization, Methodology, Writing—original draft, Project administration, Writing—review and editing; Julian Kosciessa, Software, Formal analysis, Writing—review and editing; Ulman Lindenberger, Resources, Funding acquisition, Writing—review and editing; Johannes Jacobus Fahrenfort, Conceptualization, Data curation, Software, Formal analysis, Supervision, Visualization, Methodology, Writing—original draft, Project administration, Writing—review and editing; Douglas Garrett, Conceptualization, Resources, Formal analysis, Supervision, Funding acquisition, Investigation, Methodology, Writing—review and editing.

Abstract

Strategically adopting decision biases allows organisms to tailor their choices to environmental demands. For example, a liberal response strategy pays off when target detection is crucial, whereas a conservative strategy is optimal for avoiding false alarms. Using conventional time-frequency analysis of human electroencephalographic (EEG) activity, we previously showed that bias setting entails adjustment of evidence accumulation in sensory regions (Kloosterman et al., 2019), but the presumed prefrontal signature of a strategic conservative-to-liberal bias shift has remained elusive. Here, we show that a liberal bias shift relies on frontal regions adopting a more unconstrained neural regime (boosted entropy) that is suited to the detection of unpredictable events. Overall EEG variation, spectral power and event-related potentials could not explain this relationship, highlighting the unique contribution of moment-to-moment neural variability to bias shifts. Neural variability modulation through prefrontal cortex appears instrumental for permitting an organism to tailor its decision bias to environmental demands.

Impact statement

Moment-to-moment variability is a prominent feature of neural activity. Rather than representing mere noise, this variability might enable us to flexibly adapt our decision biases to the environment.

Introduction

We often reach decisions not only by objectively weighing different alternatives, but also by allowing subjective biases to influence our choices. Ideally, such biases should be under internal control, allowing us to flexibly adapt to changes in task context while performing a challenging task. Specifically, contexts which prioritize

target detection benefit from a liberal response strategy, whereas a conservative strategy should be used at times when it is important to avoid errors of commission (e.g., false alarms). Strategic shifts in decision bias are presumed to rely on prefrontal cortex (Rahnev et al., 2016a), but despite growing interest (Chen et al., 2015a; Reckless et al., 2014a; Windmann et al., 2002a), the spatio-temporal neural signature of such within-person bias shifts is currently unknown.

One candidate neural signature of decision bias shifts that has not been considered thus far is the variability of brain activity, as reflected in its moment-to-moment irregularity. Temporal neural variability is a prominent feature in all types of neural recordings (single-cell, local field potentials, EEG/MEG, fMRI) and has traditionally been considered noise that corrupts neural computation (Dinstein et al., 2015; Faisal et al., 2008). In contrast, increased neural variability is increasingly proposed to support cognitive flexibility by allowing the brain to continuously explore its dynamical repertoire, helping it to quickly adapt to and process a novel stimulus (Ghosh et al., 2008; Mišić et al., 2010). Indeed, a growing body of evidence suggests that neural variability can prove optimal for neural systems, allowing individuals to perform better, respond faster, and adapt quicker to their environment (Garrett et al., 2015, 2013a, 2011).

One tangible possibility is that cognitive flexibility emerges when a neural system avoids locking into a stereotypical, rhythmic pattern of activity, while instead continuously exploring its full dynamic range to better prepare for unpredictably occurring events. Consistent with this notion of exploration, influential attractor models of neural population activity (Chaudhuri et al., 2019; Inagaki et al., 2019; Wimmer et al., 2014) typically contain a "noise" component that drives a dynamical system from attractor state to attractor state within a high-dimensional state space

(Deco et al., 2009; Deco and Romo, 2008). This noise element might indeed

translate into modulations of moment-to-moment neural variability during cognitive

operations that can be empirically observed. Here, we perform a crucial test of the

utility of temporal neural variability in the context of adaptive human decision making.

In line with recent ideas that a high-fidelity, more variable neural encoding regime

may be particularly required in more complex, non-deterministic situations (Garrett et

al., 2020; Marzen and DeDeo, 2017; Młynarski and Hermundstad, 2018), we

hypothesized that increased neural variability might underlie a state of higher

receptiveness to and preparedness for events of interest that are not predictable in

time, permitting the adoption of a more liberal bias toward confirming that such an

event has indeed occurred.

Interestingly, improved cognitive function has also recently been linked to

reduced neural variability, in line with the presumed corruptive role of noise for

cognitive operations (Faisal et al., 2008). In particular, transient variability decreases

after stimulus onset – called "quenching" (Churchland et al., 2010) – have been

proposed to reflect the settlement of an attractor into a stable state (Churchland et

al., 2010; Schurger et al., 2015; Wang, 2002), with quenching reportedly being

stronger during conscious perception relative to when a stimulus passes unseen

(Schurger et al., 2015). Stronger quenching has also been reported in observers with

higher perceptual sensitivity (Arazi et al., 2017), in line with a central assumption of

signal detection theory (SDT) that internal noise is detrimental for sensitivity and thus

should be suppressed (Green and Swets, 1966). To attend to this conceptual

discrepancy, we also asked whether a quenching effect can also be observed in

moment-to-variability, and if so, whether it reflects strategic decision bias shifts and

perceptual sensitivity.

We investigated these issues using previously published data from humans

performing a continuous target detection task under two different decision bias

manipulations, while non-invasively recording their electroencephalogram (EEG)

(Kloosterman et al., 2019). Sixteen participants (three experimental sessions each)

were asked to detect orientation-defined squares within a continuous stream of line

textures of various orientations and report targets via a button press (Figure 1A). In

alternating nine-minute blocks of trials, we actively biased participants' perceptual

decisions by instructing them either to report as many targets as possible (liberal

condition), or to only report high-certainty targets (conservative condition). We

played auditory feedback after errors and imposed monetary penalties to enforce

instructions.

Figure 1 | Experimental paradigm and behavioral results A. Top, target and non-target stimuli.

Subjects detected targets (left panel) within a continuous stream of diagonal and cardinal line stimuli

(middle panel), and reported targets via a button press. In different blocks of trials, subjects were

instructed to actively avoid either target misses (liberal condition) or false alarms (conservative

condition). Auditory feedback was played directly after the respective error in both conditions (right

panel). Bottom, time course of an experimental session. The two conditions were alternatingly

administered in blocks of nine minutes. In between blocks participants were informed about current

task performance and received instructions for the next block. Subsequent liberal and conservative

blocks were paired for within-participant analyses (see panel D, and Figure 3C). **B.** Distributions of

participants' criterion in both conditions. A positive criterion indicates a more conservative bias,

whereas a negative criterion indicates a more liberal bias. **C.** Lines indicating the criteria used by each

participant in the two conditions, highlighting individual differences both in overall criterion (line

intercepts), and in the size of the criterion shift between conditions (slopes). **D.** Within-person bias

shifts for liberal–conservative block pairs (see panel A, bottom). Participants were sorted based on

average criterion shift before plotting.

The following figure supplement is available for Figure 1:

Figure supplement 1 | Perceptual sensitivity and relationship between decision bias and sensitivity.

In our previous paper on these data, we reported within-participant evidence that

decision bias in each condition separately is implemented by modulating the

accumulation of sensory evidence in posterior brain regions through oscillatory EEG

activity in the 8-12 Hz (alpha) and gamma (60-100 Hz) frequency ranges

(Kloosterman et al., 2019). In no brain region, however, did we find a change-change

relationship between participants' liberal–conservative shifts in decision bias and in

spectral power, despite substantial available data per participant (on average 1733

trials) and considerable individual differences in the bias shift. Reasoning that

moment-to-moment variability of neural activity may instead better capture the

strategic bias shift from person to person and possibly reveal its hypothesized

prefrontal signature, we here measured temporal variability in the EEG data using a

novel extension of multi-scale entropy (MSE)(Costa et al., 2002). We then tested for

a change-change relationship by correlating within-person liberal–conservative shifts

in decision bias with those estimated via our modified MSE (mMSE) measure. We

indeed found that those participants who strategically shifted more towards a liberal

bias, also showed a stronger boost in mMSE. This relationship could not be

explained by overall EEG signal variation, band-specific spectral power, and event-

related potentials, highlighting the unique contribution of moment-to-moment neural

variability to the bias shift. Finally, we show that interactions between spectral power

and phase in ultra-low frequencies (1-3 Hz) may underlie the observed effects.

Results

Large individual differences in size of strategic decision bias shift

Participants differentially adopted the intended decision biases in the respective

conditions, as quantified by the criterion SDT measure of bias (Green and Swets,

1966). Subjects assumed a lower criterion (more liberal bias) when target detection

was emphasized (c = –0.13, standard deviation (SD) 0.4) and adopted a higher

criterion (more conservative bias) when instructed to avoid false alarms (c = 0.73,

SD 0.36; liberal vs. conservative, p = 0.001, two-sided permutation test, 1000

permutations) (Figure 1B). Participants varied substantially not only in the average

criterion they used across the two conditions (range of c = –0.24 to 0.89), but also in

the size of the criterion shift between conditions (range of Δc = –1.54 to –0.23).

Highlighting the extent of individual differences, participant's biases in the two

conditions were only weakly correlated (Spearman's rho = 0.24, p = 0.36), as can be

seen from the subjects' large variation in criterion intercept and slope between the

two conditions (Figure 1C). Moreover, each participant's bias shift also fluctuated

spontaneously over the course of the experiment, as indicated by variation in

criterion differences between successive, nine-minute liberal and conservative

blocks (participant-average SD 0.37, Figure 1D). Participants also varied widely in

their ability to detect targets (range in SDT d´ 0.26 to 3.97), but achieved similar d´ in

both bias conditions (rho = 0.97, p < 0.001, Figure 1, figure supplement 1).

Moreover, the liberal–conservative bias shift was only weakly correlated with a shift

in sensitivity across participants (rho = 0.44, p = 0.09), indicating that the bias

manipulation largely left perceptual sensitivity unaffected. In our previous paper on

these data (Kloosterman et al., 2019), we also quantified decision bias in these data

in terms of the "drift bias" parameter of the drift diffusion model (Ratcliff and McKoon,

2008). We chose to focus on SDT criterion in the current paper due to its

predominant use in the literature and its comparably simpler computation, while

noting the substantial overlap between the two measures as indicated by their high

correlation (rho = –0.89, as reported in Kloosterman et al., 2019). Taken together,

we observed considerable variability in strategic decision bias shifts as a result of our

bias manipulation, both at the group level and within single individuals.

Measuring neural variability with modified multi-scale entropy

We exploited the between- and within-participant variations in liberal–conservative

criterion differences to test our hypothesis that a boost in brain signal variability

underlies a liberal bias shift. To this end, we developed a novel algorithm based on

multi-scale entropy (MSE) that directly quantifies the temporal irregularity of the EEG

signal at shorter and longer timescales by counting how often temporal patterns in

the signal reoccur during the signal's time course (Costa et al., 2002)(Figure 2A,

bottom). In general, signals that tend to repeat over time, such as neural oscillations,

are assigned lower entropy, whereas more irregular, non-repeating signals yield

higher entropy. Please see the Materials and Methods section for a step-by-step

description of the MSE computation in our EEG data.

We developed time-resolved, modified MSE (mMSE), that differs from

traditional MSE in two ways. First, slower timescales are usually assessed in

conventional entropy analysis by "coarsening" the data through averaging of data

samples close to each other in time, and repeating the pattern counting operation

(see Figure 2A). Although this method can remove faster dynamics from the data in

a straightforward way, it is prone to aliasing artifacts and thereby possibly obscures

genuine entropy effects in the data. Therefore, we instead coarsened the data by

applying a Butterworth low-pass filter followed by skipping of data points (Figure 2B),

thereby retaining better control over the frequencies present in the coarsened signal

(Semmlow, 2014; Valencia et al., 2009). Second, conventional entropy analysis

requires substantial continuous data (in the order of minutes) for robust estimation,

which makes the standard method unsuitable for studying brief, transient cognitive

operations such as perceptual decision making. To investigate entropy dynamics

over time, we calculated entropy across discontinuous data segments aggregated

across trials via a sliding window approach (Grandy et al., 2016) (Figure 2A, top),

allowing us to examine entropy fluctuations from moment to moment. Please see

Materials and Methods for details on the various analysis steps and our

modifications of the MSE algorithm.

Figure 2 | mMSE estimation procedure. A. Discontinuous entropy computation procedure. Data

segments of 0.5 s duration centered on a specific time point from each trial's onset (top row) are

selected and concatenated (middle row). Entropy is then computed on this concatenated time series

while excluding discontinuous segment borders by counting repeats of both m (here, m = 1 for

illustration purposes) and m+1 (thus 2) sample patterns and taking the log ratio of the two pattern

counts (bottom row). We used m = 2 in our actual analyses. The pattern similarity parameter r

determines how lenient the algorithm is towards counting a pattern as a repeat by taking a proportion

of the signal's standard deviation (SD), indicated by the width of the horizontal gray bars. The

pattern counting procedure is repeated at each step of the sliding window, resulting in a time course

of entropy estimates computed across trials. **B.** "Filt-skip" coarsegraining procedure used to estimate

entropy on longer timescales, consisting of low-pass filtering followed by point-skipping. Filter cutoff

frequency is determined by dividing the data sampling rate (here, 256 Hz i.e. 1 sample per 3.9 ms) by

the index of the timescale of interest (top row). The signal is then coarsened by intermittently skipping

samples (bottom row). In this example, every second sample is skipped at timescale 2, resulting in

two different time courses depending on the starting point. Patterns are counted independently for

both starting points and summed before computing entropy.

Boosted entropy in frontal electrodes is linked to strategic decision bias shift

We tested for a relationship between shifts in decision bias and neural variability

between the conservative and liberal conditions by Spearman-correlating joint

modulations of mMSE and criterion across participants (averaged over the three

sessions), for all electrodes, time points, and timescales. Strikingly, we found a

negative cluster of correlations in mid- and left-frontal electrodes ($p = 0.015$, cluster-

corrected for multiple comparisons (Maris and Oostenveld, 2007)) indicating that

participants who showed a larger bias shift from the conservative to the liberal

condition were those who also exhibited a larger boost in frontal entropy (Figure 3A).

The cluster ranged across timescales from ~20-164 ms, with most of the cluster

located after trial initialization (solid vertical line in Figure 3A). To illustrate this

correlation, we averaged liberal–conservative mMSE within the significant cluster

and plotted the across-participant change-change correlation (rho = –0.87) with

criterion in a scatter plot (Figure 3B). Statistically controlling for the participants'

perceptual ability to detect targets, quantified as the liberal–conservative shift in SDT

sensitivity measure d´ (Green and Swets, 1966) did not affect the relationship (partial

rho = –0.88, $p < 0.0001$), indicating that perceptual sensitivity could not explain our

results.

Figure 3 | Change-change correlation between liberal–conservative shifts in mMSE and bias.
A. Significant negative electrode-time-timescale cluster observed via Spearman correlation between liberal–conservative mMSE and liberal–conservative SDT criterion, identified using a cluster-based permutation test across participants to control for multiple comparisons. Correlations outside the significant cluster are masked out. Left panel, time-timescale representation showing the correlation cluster integrated over the electrodes included in the cluster, indicated by the black dots in the topographical scalp map in the right panel. Dot size indicates how many time-timescale bins contribute to the cluster at each electrode. Color borders are imprecise due to interpolation used while plotting. The solid vertical line indicates the time of trial onset. The dotted vertical line indicates time of (non)target onset. Right panel, scalp map of mMSE integrated across time-timescale bins belonging to the cluster. P-value above scalp map indicates statistical significance of the cluster. The black triangle indicates participants' median reaction time, averaged across participants and conditions. **B.** Scatter plot of the correlation after averaging mMSE within time-timescale-electrode bins that are part of the three-dimensional cluster. Since the cluster was defined based on significant change-change correlations, averaging mMSE across the significant time-timescale-electrode bins before correlating represents no new information and serves only to illustrate the negative relationship identified in panel A. Both Pearson's r and Spearman's rho are indicated. **C.** Within-participant mMSE vs. criterion

slopes across liberal–conservative block pairs completed across the experiment. r_{rm}, repeated

measures correlation across all block pairs performed after centering each participant's shifts in

mMSE and criterion around zero by removing the intercept. Gray lines, individual participant slopes fit

across liberal–conservative mMSE vs criterion block pairs. Black line, slope averaged across

participants.

The following source data and figure supplements are available for Figure 3:

Source data 1. This MATLAB file contains the data for Figure 3.

Figure supplement 1. Correlation between liberal–conservative shifts in mMSE and bias in split data.

Figure supplement 2. Correlations between liberal–conservative shifts in mMSE, criterion, EEG

signal SD and spectral power.

Figure supplement 3. Control analyses investigating signal SD and point averaging method.

Figure supplement 4. EEG spectral power normalized by subtracting the condition-average pre-trial

baseline.

Figure supplement 5. Event-related responses in both conditions.

Entropy-bias correlation is also present within participants and in split data

Correlating brain and behavior across a relatively modest number of participants can

be unreliable (Yarkoni, 2009), depending on the amount of data underlying each

observation. Therefore, we next employed two complementary approaches to

strengthen evidence for the observed link between shifts in neural variability and

decision bias. We first asked whether mMSE and bias were also linked *within*

participants across the nine liberal–conservative block pairs that each participant

performed throughout the three sessions (see Figure 1A, bottom for task structure

and Figure 1D for criterion shifts in single block pairs). Critically, we observed a

negative repeated measures correlation (Bakdash and Marusich, 2017) between

within-participant shifts in criterion and mMSE ($r_{rm} = -0.19$, p = 0.046, Figure 3C),

providing convergent within-person evidence for a link between shifts in decision

bias and neural variability. Second, we tested whether the observed across-

participant correlation was present within two separate halves of the data after an

arbitrary split based on odd and even trials. We found significant correlations in both

data halves, indicating reliable between-subject associations (odd, rho = −0.77, p =

0.001; even, rho = −0.75, p = 0.001)(Figure 3, figure supplement 1).

Entropy-bias relationship is not explained by total signal variation or spectral power

Next, we investigated whether the entropy-behavior correlation could alternatively be

explained by total signal variation (quantified via the signal SD), or spectral power.

Specifically, the variance structure of a signal can influence entropy estimates

through the pattern similarity (r) parameter (width of gray bars in Figure 2), even

when this parameter is recomputed for each timescale after coarsening, as we did

(Kosciessa et al., 2020). In addition, E/MEG data is often quantified in terms of

oscillatory spectral power in canonical delta (1-2 Hz), theta (3-7 Hz), alpha (8-12 Hz),

beta (13-30 Hz) and gamma (60-100 Hz) bands, which might be able to explain the

entropy results through a similar dependency. (See Kloosterman et al., 2019 for

detailed spectral analysis of the current dataset). Therefore, we tested whether the

Δbias-Δentropy correlation could be explained by broadband signal SD and band-

specific spectral power. To make the computation of spectral power and entropy as

similar as possible, we used the same 0.5 s sliding window and 50 ms step size for

spectral analysis (1 s window to allow delta power estimation, see methods), and

selected spectral power within the same electrodes and time points in which the

mMSE effect was indicated.

Strikingly, we found that the Δbias-Δentropy correlation remained strong and

significant both when controlling for signal SD (partial rho = −0.82, p < 0.0001), and

even when controlling for all major power bands simultaneously (delta, theta, alpha,

beta, gamma; partial rho = −0.75, p = 0.005). See Figure 3, figure supplement 2 for

correlations between mMSE and various potentially confounding factors. We also

found similar results when separately controlling for signal SD within each time-

timescale bin while correlating modulations of mMSE and criterion in all electrodes,

time points, and timescales (Figure 3, figure supplement 3A). Importantly, the results

did depend on our modified entropy estimation method, since the frontal correlation

cluster was smaller and non-significant when performing the Δbias-Δentropy

correlation using conventional MSE combined with our novel sliding window

approach (cluster p = 0.37)(Costa et al., 2002)(Figure 3, figure supplement 3B). In

contrast to mMSE, spectral power was not linked to the strategic bias shift. We found

no significant clusters when correlating the liberal–conservative shifts in bias versus

raw spectral power and versus percent signal change power modulation using either

the condition-specific pre-stimulus baseline, or a condition-average baseline

subtraction (Figure 3, figure supplement 4). Taken together, neither overall signal

variation, nor spectral power could account for this observed correlation between

shifts in mMSE and the strategic decision bias.

Entropy-bias relationship is not explained by event-related potentials

Next, we investigated whether event-related potentials (ERPs), as a relatively simple

and widely used EEG metric, could explain the observed link between shifts in

criterion and entropy (Luck et al., 2000). We computed ERPs for the liberal and

conservative conditions by averaging trial time courses and tested them against

each other for all electrodes and time points using the cluster-based permutation

test. We observed one significant positive cluster (p = 0.001, cluster-corrected,

indicating a stronger ERP for the liberal condition) over central and parietal

electrodes, and one negative cluster (p = 0.001) in midfrontal, central and parietal

electrodes (Figure 3, figure supplement 4). The timing and topography of the positive

ERP cluster closely corresponded to the centroparietal positivity (CPP) signal

thought to reflect sensory evidence accumulation (O'Connell et al., 2012). This is in

line with our previous report of increased evidence accumulation in the liberal

condition (Kloosterman et al., 2019). Importantly, when repeating the change-change

correlation between liberal–conservative ERPs and criterion as performed for

mMSE, we found no significant clusters (lowest p-value positive cluster, p = 0.69;

negative cluster, p = 0.23). Furthermore, we repeated the mMSE analysis after

removing the ERP from the overall EEG activity by subtracting the event-related

potential (computed by averaging all trials within a condition, session, and

participant) from each single trial. ERP subtraction from single trials is typically

performed to remove stimulus-evoked activity and focus on ongoing or "induced"

neural activity (Klimesch et al., 1998). The bias-entropy correlation remained virtually

unchanged after removing the ERP (rho = –0.90 with removal versus rho = –0.87

without removal). See for a similar finding Mišić et al. (2010). Taken together, these

results suggest that liberal–conservative ERPs cannot account for the brain-behavior

link observed between shifts in bias and entropy.

Entropy-bias relationship possibly mediated by power-phase interactions in

the delta range

Given that overall signal variation, spectral power, and ERPs were not able to

explain our entropy findings, an intriguing question remains exactly which aspect of

the EEG signal underlies the observed link between entropy and decision bias. Since

we previously found stronger midfrontal oscillatory activity in these data in the

delta/theta (2-6 Hz) frequency range (Kloosterman et al., 2019), we next examined

the impact of systematically removing the lowest frequencies in the data on the

strength of the observed brain-behavior relationship. To this end, we performed

entropy analysis after applying a high-pass filter with 1, 2, or 3 Hz cutoff frequencies

to the time-domain data (filter order of 4, trials mirror-padded to 4 s to allow robust

estimation (Cohen, 2014)). Note that we applied a 0.5 Hz high-pass filter during data

preprocessing to remove slow drift in all cases. To quantify the strength of the brain-

behavior correlation at each filtering step, we averaged mMSE within the time-space-

timescale cluster that showed the strong negative correlation in our principal (non-

filtered) analysis (see Figure 3B) and plotted scatterplots of the correlation.

Figure 4 shows the results of this analysis for non-filtered (Figure 4A, copied

from Figure 3B), 1, 2, and 3 Hz high-pass filters (Figure 4B-D). Interestingly, we

found that the brain-behavior relationship progressively weakened with higher cutoff

frequencies, such that the correlation was non-significant after applying a 3 Hz high-

pass filter before entropy analysis. Whereas this finding suggests that these low

frequencies contribute to our entropy effects, our control analysis in Figure 3, figure

supplement 2G indicates that statistically controlling for 1-2 Hz (delta) power does

not affect the brain-behavior relationship. One explanation for these seemingly

incongruent results could be the different ways in which oscillatory phase is treated

in these two analyses: whereas statistically controlling for delta power does not take

delta phase into account, the high-pass filter removes both power and phase

information from the signal before entropy is computed. Taken together, these

analyses reveal that the lowest frequencies present in the data might play a role in

the entropy-behavior relationship, possibly through non-linear interactions between

spectral power and phase of these frequencies.

Figure 4 | Liberal–conservative change-change correlation between mMSE and decision bias for non-filtered data (**A.**) and after 1,2, and 3 Hz high-pass filtering (**B. C.** and **D.**)

Entropy quenching is not related to behavior

Finally, we tested whether variability quenching was related to behavior in our data. Specifically, improved perceptual sensitivity has been linked to a transient, post-stimulus decrease in neural variability (Arazi et al., 2017; Churchland et al., 2010; Schurger et al., 2015). Quenching is directly predicted by attractor models of brain organization (Wang, 2002), and is consistent with the main principle of signal detection theory that suppression of neural noise enhances perception (Green and Swets, 1966). Quenching has also been reported in the human EEG in terms of a variance reduction across trials in visual cortex following stimulus onset (Arazi et al., 2017), although this type of quenching can be attributed to the well-known suppression of low-frequency (alpha and beta) spectral power following stimulus onset (Daniel et al., 2019). To our knowledge only Arazi et al. (2017) report an across-participant correlation between perceptual sensitivity and variance quenching. However, this correlation could be explained by the absolute variability in the pre-stimulus period, suggesting that pre-stimulus variability was the relevant factor for behavior, rather than the post-stimulus quench. Therefore, we tested the

link between entropy quenching and behavior in our data without any strong prior hypothesis.

To investigate this issue, we computed mMSE quenching by converting the raw mMSE values into percentage modulation from the pre-stimulus baseline and testing this modulation against zero. Besides a lateral occipital enhancement of mMSE modulation (Figure 5A) that could not be explained by spectral power modulation (Figure 5B), we also found a suppression of mMSE with a focal, mid-occipital topography, in line with quenching (Figure 5C). This focal topography was highly similar to that of the SSVEP evoked by the strong, visual stimulation frequency at 25 Hz (see Figure 3A of (Kloosterman et al., 2019). In addition, spectral analysis of the ERP-subtracted data also revealed involvement of the subharmonic of this stimulation frequency at 12.5 Hz (data not shown). The highly periodic nature of this boosted SSVEP is bound to decrease the temporal irregularity of the signal, which could explain the observed mMSE suppression. In addition, the effect is strongest in shorter time scales below 40 ms because of the progressive low-pass filter implemented for longer timescales in the coarse graining procedure, which removes these SSVEP-related frequencies from timescales slower than ca. 40 ms. mMSE quenching was indeed strongly positively correlated with low-frequency power encompassing 12.5 and 25 Hz (Figure 5D). Thus, the strongly periodic SSVEP boost after stimulus onset likely increased the temporal regularity of the EEG signal, which in turn suppressed post-stimulus entropy and manifested as quenching.

Contrasting transient mMSE percentage modulation between the two conditions, we found a significant positive cluster in midfrontal electrodes, indicating a stronger transient mMSE increase following trial onset in the liberal condition

(Figure 5E). Furthermore, when change-change correlating liberal–conservative

mMSE modulation and criterion, we observed a left-lateralized negative cluster in

temporal electrodes, but no cluster in occipital electrodes (Figure 5F). Finally, we

found no significant cluster when correlating liberal–conservative mMSE quenching

with shifts in perceptual sensitivity (d′). Thus, although we did find occipital, transient

entropy quenching likely due to the strengthened SSVEP response, we found no

convincing link between entropy quenching and behavior. Taken together, the

various analyses reported in the Results section suggest a unique contribution of

moment-to-moment neural entropy to bias shifts in human decision making that

cannot be accounted for by overall brain signal variation, oscillatory neural dynamics,

ERPs, variability quenching, or perceptual sensitivity.

Figure 5 | mMSE modulation with respect to pre-trial baseline. A. Significant positive cluster

observed in longer timescales after normalizing mMSE values to percent signal change (psc) units

with respect to the pre-trial baseline (–0.2 to 0 s) and averaging across conditions. **B.** Correlation

between mMSE modulation in the positive cluster depicted in A. and spectral power modulation in midfrontal electrodes. Left panel, 3-7 Hz; right panel, 12-30 Hz. **C. D.** As B. but for the posterior negative cluster. **E.** Significant positive cluster observed in mid-frontal electrodes in the liberal–conservative contrast of mMSE modulation. **F.** Significant cluster resulting from the correlation between liberal–conservative mMSE modulation with liberal–conservative SDT criterion. Conventions as in Figure 3.

Discussion

Strategic decision biases allow organisms to adapt their choices to the context in which decisions are made. Frontal cortex has previously been shown to be involved in strategic bias shifts in humans (Chen et al., 2015b; Rahnev et al., 2016b; Reckless et al., 2014b; Windmann et al., 2002b) and monkeys (Ferrera et al., 2009), but its spatiotemporal neural signature has to date remained elusive. Here, we provide first evidence that flexible adjustment of moment-to-moment variability (entropy) in frontal regions may underlie such strategic shifts in decision bias, independent of total brain signal variation, oscillatory neural dynamics, and ERPs. This entropy-behavior relation was observed at longer time scales, for which estimation was biased in a large amount of previous work (Kosciessa et al., 2020). Since the results reported here were exclusively observed with principled extensions of the mMSE algorithm, our finding provides initial evidence for the unique value of brain signal irregularity at longer time scales.

The observed relationship between shifts in bias and neural variability in anterior brain regions complements our previous findings in the frequency domain that humans can intentionally control prestimulus 8–12 Hz (alpha) oscillatory power in posterior regions to strategically bias decision making (Kloosterman et al., 2019). Notably, we previously observed increased oscillatory 2—6 Hz (theta) power in the

liberal condition in the same midfrontal electrodes implicated here in the Δbias-

Δentropy correlation, but this theta power difference was not correlated with the bias

shift. This suggests that the strategic bias shift may be reflected both in low-

frequency spectral power and entropy in midfrontal regions, but that only entropy is

linked to the magnitude of the decision-maker's bias shift. One possible explanation

for such a dissociation is that spectral power exclusively reflects the amplitude of

oscillatory signal contributions while discarding their phase information. In contrast,

entropy is sensitive to both variations in the magnitude as well as the phase of signal

fluctuations. This notion is also in line with our finding that low-frequency spectral

power is insufficient to explain our observed brain-behavior relationship, while the

presence of these frequencies (including narrowband non-linear phenomena such as

phase resets or temporal dependencies in the amplitude of signals (Linkenkaer-

Hansen et al., 2001)) during entropy estimation is sufficient and necessary for the

relationship to emerge (Figure 4). Moreover, whereas spectral analysis strictly

assumes a sinusoidal waveform of EEG signal fluctuations (Cole and Voytek, 2017;

Jones, 2016), entropy is agnostic to the shape of the waveforms present in the

signal. Entropy thus provides a more unrestricted description of moment-to-moment

fluctuations in neural activity that is highly predictive of strategic decision bias shifts

across participants in our data. Our results suggest that entropy taps into information

in the EEG signal that is not available in ERPs and spectral power – the most

popular analysis methods used in the field. Intriguingly, this suggests that many

previous E/MEG studies analyzing ERPs and/or spectral power only might have

structurally overlooked a crucial property of the EEG signal that is in fact strongly

linked to behavior. It could thus be that many interesting brain-behavior links are still

hidden in existing EEG datasets, which can now be uncovered using mMSE.

Despite the consistent liberal bias shift that participants exhibited between the

two conditions and the strong link between entropy and behavior, mMSE was not

significantly higher on average in the liberal condition. This is in contrast to the raw

alpha and theta power differences that we reported previously, which did show

significant differences (see Kloosterman et al., 2019). Paradoxically, we show here

that the shift in mMSE did correlate with the liberal–conservative bias shift, whereas

the shifts in alpha and theta did not – this is indeed the research advance of the

current paper. This indicates that inter-individual differences in mMSE may be

sensitive to behavioral adjustments even in the absence of a group-wise shift. A

possible explanation for such a dissociation is that a main effect and a correlation

address somewhat divergent research questions that may conflict with each other.

On the one hand, a main effect tends to occur when subjects respond similarly to an

experimental condition, which typically requires that individual differences be

relatively small. On the other hand, chances of detecting a correlation with behavior

typically increase when individual differences are larger, (e.g., (Lindenberger et al.,

2008)). Thus, the common presumption that a main effect of condition is a

prerequisite for detecting behavioral effects is unfounded in our view. In our study,

certain participants may have used the correct and others the incorrect strategy,

leading to a lack of average across- participant condition effect, yet revealing why

their behaviors differed on task. Therefore, we did not *a priori* hypothesize a main

effect of condition in the liberal–conservative contrast, but rather focused our

hypotheses on inter-individual adjustments in mMSE that tracked the magnitude of

the individual bias shift. Finally, we note that we did indeed observe a significantly

stronger mMSE transient increase after trial onset in the liberal condition (after

baseline-correcting the data, as is common in time-frequency EEG analysis).

Perhaps mediated by the logic outlined above, the change-change correlation with

behavior is weaker using baseline corrected mMSE.

In apparent contrast to the view that neural variability facilitates cognition,

previous work has suggested that a temporary stabilization of neural activity after

stimulus onset (quenching, quantified as a transient suppression of time-domain

variance) is beneficial for perception (Arazi et al., 2017; Schurger et al., 2015). We

also observed a suppression in baseline-corrected mMSE, which is likely due to

increased regularity of the time-domain signal due to the boosted power at the

SSVEP frequency (Kloosterman et al., 2019). Previous work has linked variance

quenching to post-stimulus suppression of rhythmic low-frequency (alpha and beta)

power (Daniel et al., 2019). Future work could investigate whether entropy indeed

increases after suppression of these temporally regular signals. Crucially however –

and divergent from our finding that boosting variability facilitates a strategic bias shift

– we found no evidence for a change-change relationship between entropy

quenching and decision bias or perceptual sensitivity. Since the relations between

quenching observed in neural spiking (Churchland et al., 2010), trial-by-trial variance

of E/MEG (Arazi et al., 2017), and mMSE are currently unclear, further investigation

on this issue is needed (Garrett et al., 2013b). Future studies could also explore how

neural variability quenching and boosting in different timescales are related to

various aspects of decision making such as perceptual sensitivity and different kinds

of biases (Fleming et al., 2010; Talluri et al., 2018; Urai et al., 2019), but also to

confidence and metacognitive processes (Fleming and Dolan, 2012; Yeung and

Summerfield, 2012). Furthermore, individual decision bias has also been linked to

the magnitude of transient dilations of the pupil (Gee et al., 2017, 2014) and to

entropy of EEG (Waschke et al., 2019), suggesting that pupil-linked neuromodulation

(Joshi and Gold, 2020) could be related to decision bias through adjustments to

moment-to-moment neural variability. Further investigation of such relationships

could yield fruitful insights about the neurochemical mechanisms underlying

associations between neural variability and higher-order cognitive function (Alavash

et al., 2018; Garrett et al., 2015).

Our findings may have important implications for dynamical attractor models

of neural population activity, which have become increasingly influential in recent

years (Chaudhuri et al., 2019; Inagaki et al., 2019; Wimmer et al., 2014). Attractor

models cast cognitive outcomes (e.g. decisions, perceptual experiences, or retention

of an item in working memory) as low-dimensional, stable states ("attractors") within

a high-dimensional energy landscape (Deco et al., 2009; Deco and Romo, 2008).

These models typically contain a "noise" component that enables probabilistic

exploration of the energy landscape, increasing chances of attraction to the most

optimal state (e.g. the correct decision) and limiting the likelihood of settling too

rigidly into any particular attractor state. In multi-stable visual perception, for

example, noise is thought to underlie the spontaneous, flexible switching between

perceptual states reported by observers while viewing bi-stable visual illusions

(Kloosterman et al., 2015; Moreno-Bote et al., 2007). Our results suggest that an

element of noise facilitating cognitive flexibility might translate into modulations of

moment-to-moment neural variability that can be measured in cortical population

activity. Future work on dynamical attractor modelling could investigate exactly which

characteristics of this noise component underlie effective exploration of the state

space within these models, for example by modulating noise amplitude as well as

the relative contribution of different noise frequencies (noise color). Modeling insights

could then guide the search for signatures of noise supporting cognitive operations in

moment-to-moment neural variability.

Our results suggest that dynamic adjustment of neural variability in frontal

regions is crucial for adaptive behavior. Based on our findings, we propose that

heightened frontal entropy results from a more dynamic, irregular neural regime that

enables an individual to be more prepared to process and act upon uncertain, yet

task-relevant information. We believe that quantifying shifts in neural entropy could

help elucidate the mechanisms allowing organisms to adapt to their environment and

ultimately increase their chances of survival.

Materials and Methods

Key resources table

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
biological sample (Humans)	Participants	(Kloosterman et al., 2019)	https://doi.org10.7554/elife.37321	See Subjects section in Materials and Methods
software, algorithm	MATLAB	Mathworks	MATLAB_R2016b, RRID:SCR_001622	
software, algorithm	Presentation	NeuroBS	Presentation_v9.9, RRID:SCR_002521	

software, algorithm	Custom analysis code	Kloosterman (2018)	https://github.com/ nkloost1/critEEG	
other	EEG data experimental task	(Kloosterman et al., 2018)	https://doi.org/10.6084/ m9.figshare.6142940	

We report a novel analysis of a previously published dataset involving a target detection task during two different decision bias manipulations (Kloosterman et al., 2019).

Subjects Sixteen participants (eight females, mean age 24.1 years, ± 1.64) took part in the experiment, either for financial compensation (EUR 10,– per hour) or in partial fulfillment of first year psychology course requirements. Each participant completed three experimental sessions on different days, each session lasting ca. 2 hours, including preparation and breaks. One participant completed only two sessions, yielding a total number of sessions across subjects of 47. Due to technical issues, for one session only data for the liberal condition was available. One participant was an author. All participants had normal or corrected-to-normal vision and were right handed. Participants provided written informed consent before the start of the experiment. All procedures were approved by the ethics committee of the University of Amsterdam.

Stimuli Stimuli consisted of a continuous semi-random rapid serial visual presentation (rsvp) of full screen texture patterns. The texture patterns consisted of line elements approx. 0.07° thick and 0.4° long in visual angle. Each texture in the rsvp was presented for 40 ms (i.e. stimulation frequency 25 Hz), and was oriented in one of four possible directions: 0°, 45°, 90° or 135°. Participants were instructed to

fixate a red dot in the center of the screen. At random inter trial intervals (ITI's)

sampled from a uniform distribution (ITI range 0.3 – 2.2 s), the rsvp contained a fixed

sequence of 25 texture patterns, which in total lasted one second. This fixed

sequence consisted of four stimuli preceding a (non-)target stimulus (orientations of

45°, 90°, 0°, 90° respectively) and twenty stimuli following the (non)-target

(orientations of 0°, 90°, 0°, 90°, 0°, 45°, 0°, 135°, 90°, 45°, 0°, 135°, 0°, 45°, 90°, 45°,

90°, 135°, 0°, 135° respectively) (see Figure 1A). The fifth texture pattern within the

sequence (occurring from 0.16 s after sequence onset) was either a target or a

nontarget stimulus. Nontargets consisted of either a 45° or a 135° homogenous

texture, whereas targets contained a central orientation-defined square of 2.42°

visual angle, thereby consisting of both a 45° and a 135° texture. 50% of all targets

consisted of a 45° square and 50% of a 135° square. Of all trials, 75% contained a

target and 25% a nontarget. Target and nontarget trials were presented in random

order. To avoid specific influences on target stimulus visibility due to presentation of

similarly or orthogonally oriented texture patterns temporally close in the cascade, no

45° and 135° oriented stimuli were presented directly before or after presentation of

the target stimulus. All stimuli had an isoluminance of 72.2 cd/m^2. Stimuli were

created using MATLAB (The Mathworks, Inc., Natick, MA, USA) and presented using

Presentation version 9.9 (Neurobehavioral systems, Inc., Albany, CA, USA).

Experimental design The participants' task was to detect and actively report targets

by pressing a button using their right hand. Targets occasionally went unreported,

presumably due to constant forward and backward masking by the continuous

cascade of stimuli and unpredictability of target timing (Fahrenfort et al., 2007). The

onset of the fixed order of texture patterns preceding and following (non-)target

stimuli was neither signaled nor apparent. At the beginning of the experiment,

participants were informed they could earn a total bonus of EUR 30, -, on top of their regular pay of EUR 10, - per hour or course credit. In two separate conditions within each session of testing, we encouraged participants to use either a conservative or a liberal bias for reporting targets using both aversive sounds as well as reducing their bonus after errors. In the conservative condition, participants were instructed to only press the button when they were relatively sure they had seen the target. The instruction on screen before block onset read as follows: "Try to detect as many targets as possible. Only press when you are relatively sure you just saw a target." To maximize effectiveness of this instruction, participants were told the bonus would be diminished by 10 cents after a false alarm. During the experiment, a loud aversive sound was played after a false alarm to inform the participant about an error. During the liberal condition, participants were instructed to miss as few targets as possible. The instruction on screen before block onset read as follows: "Try to detect as many targets as possible. If you sometimes press when there was nothing this is not so bad." In this condition, the loud aversive sound was played twice in close succession whenever they failed to report a target, and three cents were subsequently deducted from their bonus. The difference in auditory feedback between both conditions was included to inform the participant about the type of error (miss or false alarm), to facilitate the desired bias in both conditions. After every block, the participant's score (number of missed targets in the liberal condition and number of false alarms in the conservative condition) was displayed on the screen, as well as the remainder of the bonus. After completing the last session of the experiment, every participant was paid the full bonus as required by the ethical committee.

Participants performed six blocks per session lasting ca. nine minutes each. During a block, participants continuously monitored the screen and were free to

respond by button press whenever they thought they saw a target. Each block

contained 240 trials, of which 180 target and 60 nontarget trials. The task instruction

was presented on the screen before the block started. The condition of the first block

of a session was counterbalanced across participants. Prior to EEG recording in the

first session, participants performed a 10-min practice run of both conditions, in

which visual feedback directly after a miss (liberal condition) or false alarm

(conservative) informed participants about their mistake, allowing them to adjust their

decision bias accordingly. There were short breaks between blocks, in which

participants indicated when they were ready to begin the next block.

Behavioral analysis We defined decision bias as the criterion measure from SDT

(Green and Swets, 1966). We calculated the criterion *c* across the trials in each

condition as follows:

$$c = -\frac{1}{2}\left[Z(Hit\text{-}rate) + Z(FA\text{-}rate)\right]$$

676   where hit-rate is the proportion target-present responses of all target-present trials,

677   false alarm (FA)-rate is the proportion target-present responses of all target-absent

678   trials, and Z(...) is the inverse standard normal distribution. Furthermore, we

679   calculated perceptual sensitivity using the SDT measure d´:

$$d' = Z(Hit\text{-}rate) - Z(FA\text{-}rate)$$

EEG recording Continuous EEG data were recorded at 256 Hz using a 48-channel

BioSemi Active-Two system (BioSemi, Amsterdam, the Netherlands), connected to a

standard EEG cap according to the international 10-20 system. Electrooculography

(EOG) was recorded using two electrodes at the outer canthi of the left and right

eyes and two electrodes placed above and below the right eye. Horizontal and

vertical EOG electrodes were referenced against each other, two for horizontal and

two for vertical eye movements (blinks). We used the FieldTrip toolbox (Oostenveld

et al., 2011) and custom software in MATLAB R2016b (The Mathworks Inc., Natick,

MA, USA; RRID:SCR_001622) to process the data. Data were re-referenced to the

average voltage of two electrodes attached to the earlobes. We applied a

Butterworth high-pass filter (fourth order, cutoff 0.5 Hz) to remove slow drifts from the

data.

Trial extraction We extracted trials of variable duration from 1 s before target

sequence onset until 1.25 after button press for trials that included a button press

(hits and false alarms), and until 1.25 s after stimulus onset for trials without a button

press (misses and correct rejects). The following constraints were used to classify

(non-)targets as detected (hits and false alarms), while avoiding the occurrence of

button presses in close succession to target reports and button presses occurring

outside of trials: 1) A trial was marked as detected if a response occurred within 0.84

s after target onset; 2) when the onset of the next target stimulus sequence started

before trial end, the trial was terminated at the next trial's onset; 3) when a button

press occurred in the 1.5 s before trial onset, the trial was extracted from 1.5 s after

this button press; 4) when a button press occurred between 0.5 s before until 0.2 s

after sequence onset, the trial was discarded. After trial extraction the mean of every

channel was removed per trial.

Artifact rejection Trials containing muscle artifacts were rejected from further

analysis using a standard semi-automatic preprocessing method in Fieldtrip. This

procedure consists of bandpass-filtering the trials of a condition block in the 110–125

Hz frequency range, which typically contains most of the muscle artifact activity,

followed by a Z-transformation. Trials exceeding a threshold Z-score were removed

completely from analysis. We used as the threshold the absolute value of the

minimum Z-score within the block, + 1. To remove eye blink artifacts from the time

courses, the EEG data from a complete session were transformed using

independent component analysis (ICA), and components due to blinks (typically one

or two) were removed from the data. In addition, to remove microsaccade-related

artifacts we included two virtual channels in the ICA based on channels Fp1 and

Fp2, which included transient spike potentials as identified using the saccadic

artefact detection algorithm from (Hassler et al., 2011). This yielded a total number of

channels submitted to ICA of 48 + 2 = 50. The two components loading high on

these virtual electrodes (typically with a frontal topography) were also removed.

Blinks and eye movements were then semi-automatically detected from the

horizontal and vertical EOG (frequency range 1–15 Hz; z-value cut-off 4 for vertical;

6 for horizontal) and trials containing eye artefacts within 0.1 s around target onset

were discarded. This step was done to remove trials in which the target was not

seen because the eyes were closed. Finally, trials exceeding a threshold voltage

range of 200 mV were discarded. To attenuate volume conduction effects and

suppress any remaining microsaccade-related activity, the scalp current density

(SCD) was computed using the second-order derivative (the surface Laplacian) of

the EEG potential distribution (Perrin et al., 1989).

ERP removal In a control analysis, we removed stimulus-evoked EEG activity

related to external events by computing the event-related potential (ERP) and

subtracting the ERP from each single trial prior to entropy or spectral analysis. This

was done to focus on ongoing (termed "induced", (Klimesch et al., 1998)) activity. To

eliminate differences in evoked responses between sessions and conditions, we

performed this procedure separately for ERPs computed in each condition, session,

and participant.

Entropy computation We measured temporal neural variability in the EEG using a

form of multiscale entropy (MSE)(Costa et al., 2002), which we modified in several

ways. MSE characterizes signal irregularity at multiple time scales by estimating

sample entropy (SampEn) of a signal's time series at various sampling rates. The

estimation of SampEn involves counting how often specific temporal patterns

reoccur over time, and effectively measures how unpredictably the signal develops

from moment to moment. At a given time scale, the estimation of SampEn consists

of the following steps:

1) A to-be-counted temporal 'template' pattern consisting of m samples is

selected, starting at the beginning of the time series.

2) The data is discretized to allow comparing patterns of samples rather than

exact sample values (which are rarely exactly equal in physiological

timeseries). A boundary parameter r is used to determine whether other

patterns in the time series match the template. r denotes the proportion of the

time series standard deviation (SD, see also Figure 2A), within which a

pattern is a match, as follows:

$$\text{Boundary parameter} = r \times \text{SD} \tag{1}$$

3) Template pattern repeats throughout the time series are counted, yielding

pattern count $N(m)$.

4) Steps 1 to 3 are repeated for patterns consisting of $m + 1$ samples, yielding

pattern count $N(m + 1)$.

5) Steps 1 to 4 are iterated to assess each temporal pattern as a template once.

Counts for each template pattern are then summed, yielding total counts

across templates for $N(m)$ and $N(m+1)$.

6) Finally, SampEn is computed as the logarithm of the ratio of the counts for m

and $m+1$:

$$\mathrm{SampEn} = \ln \frac{N(m)}{N(m+1)} \qquad (2)$$

Thus, SampEn estimates the proportion of similar sequences of m samples that are

still similar when the next sample, i.e., $m+1$, is added to the sequence. Here, we

use $m = 2$ and $r = 0.5$, as typically done in neurophysiological settings (Courtiol et

al., 2016; Grandy et al., 2016; Richman and Moorman, 2000).

We have implemented three modifications of regular MSE that we outline in the next

sections. We refer to our own measure as modified MSE (mMSE) throughout the

manuscript.

MSE modification #1: multi-scale implementation through filtering and point

skipping In multiscale entropy, the computation of SampEn is repeated for multiple

time scales after progressively lowering the time series sampling rate by a process

called "coarsening" (Costa et al., 2002). By default, SampEn quantifies entropy at the

time scale that corresponds to the sampling rate of the time series, which is typically

in the order of milliseconds or lower in (non-downsampled) neurophysiological data.

To enable estimation of entropy at longer time scales, the time series is typically

coarsening by averaging groups of adjacent samples ("point averaging") and

repeating the entropy computation (Costa et al., 2002). However, despite being

straightforward, this method is suboptimal for eliminating short temporal scales from

the time series. Point averaging is equivalent to low-pass filtering using a finite-

impulse response filter, which does not effectively eliminate high frequencies

(Semmlow, 2014; Valencia et al., 2009). For this reason, we used an improved

coarsening procedure involving replacement of the multi-point average by a low-pass

Butterworth filter, which has a well-defined frequency cutoff and precludes aliasing

(Valencia et al., 2009)(Figure 2B, top). The filter cutoff frequency $CutoffFreq$ is

determined as:

$$CutoffFreq = NyquistFreq \times \frac{1}{\text{scale number}} \qquad\qquad (3)$$

where $NyquistFreq$ is the highest estimable frequency given the signal's sampling

rate. This filtering ensures that an increasingly larger portion of the higher

frequencies is removed for slower time scales. Note that low-pass filtering affects the

temporal structure of the time-domain signal, which could hamper the interpretation

of the EEG dynamics due to smearing of responses (VanRullen, 2011). This issue is

largely mitigated, however, due to the liberal–conservative subtraction that we

perform before correlating with behavior, since this issue presumably affects both

conditions similarly. Low-pass filtering is followed by a point-skipping procedure to

reduce the sampling rate of the time series (Figure 2B, bottom). Since point-skipping

omits increasingly large portions of the filtered time series depending on the starting

point of the point-skipping procedure, we counted patterns separately for each

starting point within a scale (see section Entropy computation above), summed their

counts for $N(m)$ and $N(m + 1)$ and computed entropy as described above.

MSE modification #2: Pattern similarity recomputed at each time scale By

increasingly smoothing the time series, coarse-graining affects not only on the

signal's entropy, but also its overall variation, as reflected in the decreasing standard

deviation as a function of time scale (Nikulin and Brismar, 2004). In the original

implementation of the MSE calculation, the similarity parameter r was set as a

proportion of the original (scale 1) time series' standard deviation and applied to all

the scales (Costa et al., 2002). Because of the decreasing variation in the time series

due to coarse graining, the similarity parameter therefore becomes increasingly

tolerant at slower time scales, resulting in more similar patterns and decreased

entropy. This decreasing entropy can be attributed both to changes in signal

complexity, but also in overall variation (Kosciessa et al., 2020; Nikulin and Brismar,

2004). To overcome this limitation, we recomputed the similarity parameter for each

scale, thereby normalizing mMSE with respect to changes in overall time series

variation at each scale.

MSE modification #3: Time-resolved computation An important limitation of MSE

is the need for substantial continuous data for robust estimation. Heuristically, the

recommended number of successive data points for estimation at each scale is 100

(minimum) to 900 (preferred) points using typical MSE parameter settings (Grandy et

al., 2016). This limitation precludes the application of MSE to neuroimaging data

recorded during cognitive processes that unfold over brief periods of time, such as

perceptual decisions. Grandy et al. (Grandy et al., 2016) has shown that the pattern

counting process can be extended to discontinuous data segments that are

concatenated across time, as long as the counting of artificial patterns across

segment borders is avoided (as these patterns are a product of the concatenation

and do not occur in the data itself). We applied the mMSE computation across

discontinuous segments of 0.5 s duration (window size). To track the evolution of

mMSE over the trial, we slid this window across the trials in steps of 50 milliseconds

from -0.2 s until 0.6 s, each time recomputing mMSE across segments taken from

the time window in each trial.

Given our segments of 0.5 s window length sampled at 256 Hz, we computed

mMSE for scales 1 (129 samples within the window) until 42 (three or four samples

within the window, depending on the starting point). Note that using a pattern

parameter $m = 2$, a minimum of three samples within a segment is required to

estimate entropy across the segments of continuous data, yielding a maximum

possible scale of 42. In line with the MSE literature (Courtiol et al., 2016), we

converted the time scale units to milliseconds by taking the duration between

adjacent data points after each coarsegraining step. For example, time scale 1

corresponds to 1000 ms / 256 Hz = 3.9 ms, and time scale 42 to 1000 / (256/42) =

164 ms.

Spectral analysis We used a sliding window Fourier transform; step size, 50 ms;

window size, 500 ms; frequency resolution, 2 Hz) to calculate time-frequency

representations (spectrograms) of the EEG power for each electrode and each trial.

We used a single Hann taper for the frequency range of 3–35 Hz (spectral

smoothing, 4.5 Hz, bin size, 1 Hz) and the multitaper technique for the 36 – 100 Hz

frequency range (spectral smoothing, 8 Hz; bin size, 2 Hz; five tapers)(Mitra and

Bokil, 2007). See (Kloosterman et al., 2019) for similar settings. Finally, to

investigate spectral power between 1-3 Hz (delta band), we performed an additional

time-frequency analysis with a window size of 1 s (i.e. frequency resolution 1 Hz)

without spectral smoothing (bin size 0.5 Hz). Spectrograms were aligned to the onset

of the stimulus sequence containing the (non)target. Power modulations during the

trials were quantified as the percentage of power change at a given time point and

frequency bin, relative to a baseline power value for each frequency bin. We used as

a baseline the mean EEG power in the interval 0.4 to 0 s before trial onset,

computed separately for each condition. If this interval was not completely present in

the trial due to preceding events (see Trial extraction), this period was shortened

accordingly. We normalized the data by subtracting the baseline from each time-

frequency bin and dividing this difference by the baseline (x 100 %). In an additional

analysis, we performed a baseline correction by subtracting the condition-averaged

pre-stimulus power, without converting into percent signal change.

Statistical significance testing of mMSE and spectral power and correlations

across space, time, and timescales/frequencies. To determine clusters of

significant modulation with respect to the pre-stimulus baseline without any a priori

selection, we ran statistics across space-time-frequency bins using paired t-tests

across subjects performed at each bin. Single bins were subsequently thresholded at

$p < 0.05$ and clusters of contiguous time-space-frequency bins were determined. For

the correlation versions of this analysis, we correlated the brain measure at each bin

with the criterion and converted the r-values to a t-statistic using the Fisher-

transformation (Fisher, 1915). We used a cluster-based procedure (Maris and

Oostenveld, 2007) to correct for multiple comparisons using a cluster-formation

alpha of $p < 0.05$ and a cluster-corrected alpha of $p = 0.05$, two-tailed (10.000

permutations). For visualization purposes, we integrated (using MATLAB's trapz

function) power or entropy values in the time-frequency/entropy representations

(TFR/TTR) across the highlighted electrodes in the topographies. For the

topographical scalp maps, modulation was integrated across the saturated time-

frequency bins in the TFRs/TTRs. See (Kloosterman et al., 2019) for a similar

procedure in the time-frequency domain.

High-pass filtering analysis To examine the effect of systematically removing lower

frequencies from the data before computing mMSE, we high-pass filtered the data

using 1,2 and 3Hz highpass filters (filter order of 4). We mirror-padded trials to 4 s to

allow robust estimation (Cohen, 2014). After high-pass filtering, we performed mMSE

analysis as reported above.

Correlation analysis We used both Pearson correlation and robust Spearman

correlation across participants to test the relationships between the behavioral

variables as well as with the EEG entropy and power (modulation). To test whether

behavior and EEG activity were linked within participants, we used repeated

measures correlation. Repeated measures correlation determines the common

within-individual association for paired measures assessed on two or more

occasions for multiple individuals by controlling for the specific range in which

individuals' measurements operate, and correcting the correlation degrees of

freedom for non-independence of repeated measurements obtained from each

individual (Bakdash and Marusich, 2017; Bland and Altman, 1995). To test whether

spectral power could account for the observed correlation between criterion and

mMSE, we used partial Spearman and Pearson correlation controlling for other

variables.

Data and code sharing The data analyzed in this study are publicly available on

Figshare (Kloosterman et al., 2018). We programmed mMSE analysis in a MATLAB

function within the format of the FieldTrip toolbox (Oostenveld et al., 2011). Our

ft_entropyanalysis.m function takes as input data produced by Fieldtrip's

ft_preprocessing.m function. In our function, we employed matrix computation of

mMSE for increased speed, which is desirable due to the increased computational

demand with multi-channel data analyzed with a sliding window. The function

supports GPU functionality to further speed up computations. The software can be

found on https://github.com/LNDG/mMSE. A tutorial for computing mMSE within the

FieldTrip   toolbox   can   be   found   on   the   FieldTrip   website

(http://www.fieldtriptoolbox.org/example/entropy_analysis/). Analysis scripts for the

current paper can be found on https://github.com/kloosterman/critEEGentropy.

Acknowledgments: Funding: Emmy Noether Grant to Douglas D Garrett, Max

Planck UCL Centre for Computational Psychiatry and Ageing Research to Douglas

Garrett, Niels Kloosterman, and Max Planck Society. **Author contributions:** Niels

Kloosterman, Conceptualization, Data curation, Software, Formal analysis,

Investigation, Visualization, Methodology, Writing—original draft, Project

administration, Writing—review and editing; Julian Kosciessa, Software, Formal

analysis, Writing—review and editing; Ulman Lindenberger, Resources, Funding

acquisition, Writing—review and editing; Johannes Jacobus Fahrenfort,

Conceptualization, Data curation, Software, Formal analysis, Supervision,

Visualization, Methodology, Writing—original draft, Project administration, Writing—

review and editing; Douglas Garrett, Conceptualization, Resources, Formal analysis,

Supervision, Funding acquisition, Investigation, Methodology, Writing—review and

editing. **Competing interests:** Authors declare no competing interests. **Data and

materials availability:** All data analyzed during this study are publicly available

(Kloosterman et al., 2018).

Source data and figure supplements

The following source data and figure supplements are included in this article:

Figure 1 – Figure supplement 1.

Figure 3 – Figure supplements 1, 2, 3, 4 and 5.

Figure 3 – Source data 1 (sourceFigure3.zip).

Figure supplement legends

Figure 1—figure supplement 1 | Perceptual sensitivity and relationship between decision bias and sensitivity. A. Distributions of participants' d' in both conditions. A higher d' indicates higher sensitivity. B. Scatter plot depicting the correlation between participant's d' in both conditions. C. Corresponding individual participant slopes. D. Correlation between d' and criterion averaged across conditions. E. Correlation between liberal – conservative d' and liberal – conservative criterion.

Figure 3—figure supplement 1 | Correlation between liberal – conservative mMSE and bias shift is reliable in split data. Spearman correlations across participants in both data halves after arbitrary split into even trials (left panel) and odd trials (right panel).

Figure 3—figure supplement 2 | Change-change correlations between liberal– conservative mMSE, criterion, EEG signal SD and spectral power. A. – H. Scatter plots of change-change correlations. See titles and axes labels of each plot for details. Least squares lines (gray) were plotted when the Pearson correlation reached significance ($p<0.05$). Both the signal SD (panel A.) and delta (panel B.) were negatively correlated with mMSE, whereas the other canonical frequency bands correlated only weakly (theta, rho = –0.31; alpha, rho = 0.18; beta, rho = 0.26; gamma, rho = 0.29, all $p > 0.24$). Signal SD and delta were also highly correlated with each other (rho = 0.93, $p < 0.001$, panel C.), indicating that delta contributed strongly to overall signal SD, as expected based on the known 1/f characteristic of

neural data. Neither liberal–conservative SD (panel D.) nor delta (panel E.)

correlated as strongly with the criterion shift as liberal–conservative mMSE (see

Figure 3B and panels F. and G.). In contrast, the correlation between SD and

criterion was not significant when controlling for mMSE (panel H.), highlighting the

unique ability of mMSE to explain the bias shift.

Figure 3—figure supplement 3 | Control analyses investigating signal SD and

point averaging method. A. Liberal – conservative mMSE vs. criterion correlation

when statistically controlling for the r parameter (signal SD) across participants. The

cluster remains significant and the topography is similar, but the effect is more

widespread across electrodes, and less widespread across timescales. **B.** As A. but

using traditional MSE, including coarse graining through point averaging to asses

longer timescales and a fixed r parameter across timescales. The cluster does not

reach significance.

Figure 3—figure supplement 4 | EEG spectral power normalized by subtracting

the condition-average pre-trial baseline. In addition to the condition-specific

baseline used in our previous paper (Kloosterman et al., 2019), we also computed

spectral power modulation using a condition-averaged baseline correction based to

investigate the link between spectral power and our experimental manipulation.

Power modulation were similar to those reported in our previous paper. **A.** Significant

clusters found within the liberal (left panels) and conservative (right panels)

conditions. **B.** Liberal – conservative contrast, showing stronger broadband power

modulation in the liberal condition, except in the alpha range. Please see our

previous paper for detailed analysis of the alpha effect. **C.** Change-change

correlations between liberal–conservative shifts in power modulation and strategic

decision bias. We observed two clusters (rows), none of which were significant after

multiple comparison correction.

Figure 3—figure supplement 5 | Event-related responses in both conditions. A.

Positive (liberal > conservative) cluster in parietal and central electrodes between

0.37 – 0.8 s after trial onset. **B.** Negative (conservative > liberal) cluster in parietal

and central electrodes between 0.08 – 0.59 s after trial onset. Conventions as in

Figure 3.

References

Alavash M, Lim S-J, Thiel C, Sehm B, Deserno L, Obleser J. 2018. Dopaminergic
modulation of hemodynamic signal variability and the functional connectome
during cognitive performance. *Neuroimage* **172**:341–356.
doi:10.1016/j.neuroimage.2018.01.048
Arazi A, Censor N, Dinstein I. 2017. Neural Variability Quenching Predicts Individual
Perceptual Abilities. *J Neurosci* **37**:97 109. doi:10.1523/jneurosci.1671-16.2017
Bakdash JZ, Marusich LR. 2017. Repeated Measures Correlation. *Front Psychol*
8:491. doi:10.3389/fpsyg.2017.00456
Bland JM, Altman DG. 1995. Statistics notes: Calculating correlation coefficients with
repeated observations: Part 1—correlation within subjects. *Bmj* **310**:446 446.
doi:10.1136/bmj.310.6977.446
Chaudhuri R, Gerçek B, Pandey B, Peyrache A, Fiete I. 2019. The intrinsic attractor
manifold and population dynamics of a canonical cognitive circuit across waking
and sleep. *Nat Neurosci* **22**:1512–1520. doi:10.1038/s41593-019-0460-x
Chen M-Y, Jimura K, White CN, Maddox WT, Poldrack RA. 2015a. Multiple brain
networks contribute to the acquisition of bias in perceptual decision-making. *Front*
Neurosci-switz **9**:63. doi:10.3389/fnins.2015.00063
Chen M-Y, Jimura K, White CN, Maddox WT, Poldrack RA. 2015b. Multiple brain
networks contribute to the acquisition of bias in perceptual decision-making. *Front*
Neurosci-switz **9**:63. doi:10.3389/fnins.2015.00063
Churchland MM, Yu BM, Cunningham JP, Sugrue LP, Cohen MR, Corrado GS,
Newsome WT, Clark AM, Hosseini P, Scott BB, Bradley DC, Smith MA, Kohn A,
Movshon JA, Armstrong KM, Moore T, Chang SW, Snyder LH, Lisberger SG,
Priebe NJ, Finn IM, Ferster D, Ryu SI, Santhanam G, Sahani M, Shenoy KV.
2010. Stimulus onset quenches neural variability: a widespread cortical
phenomenon. *Nat Neurosci* **13**:369 378. doi:10.1038/nn.2501
Cohen MX. 2014. Analyzing neural time series data: theory and practice.
Cole SR, Voytek B. 2017. Brain Oscillations and the Importance of Waveform
Shape. *Trends Cogn Sci* **21**:137–149. doi:10.1016/j.tics.2016.12.008
Costa M, Goldberger AL, Peng C-K. 2002. Multiscale entropy analysis of complex
physiologic time series. *Phys Rev Lett* **89**:068102.
doi:10.1103/physrevlett.89.068102

Courtiol J, Perdikis D, Petkoski S, Müller V, Huys R, Sleimen-Malkoun R, Jirsa VK.
2016. The multiscale entropy: Guidelines for use and interpretation in brain signal
analysis. *J Neurosci Meth* **273**:175 190. doi:10.1016/j.jneumeth.2016.09.004
Daniel E, Meindertsma T, Arazi A, Donner TH, Dinstein I. 2019. The Relationship
between Trial-by-Trial Variability and Oscillations of Cortical Population Activity.
Sci Rep-uk **9**:16901. doi:10.1038/s41598-019-53270-7
Deco G, Rolls ET, Romo R. 2009. Stochastic dynamics as a principle of brain
function. *Prog Neurobiol* **88**:1–16. doi:10.1016/j.pneurobio.2009.01.006
Deco G, Romo R. 2008. The role of fluctuations in perception. *Trends Neurosci*
31:591 598. doi:10.1016/j.tins.2008.08.007
Dinstein I, Heeger DJ, Behrmann M. 2015. Neural variability: friend or foe? *Trends*
Cogn Sci **19**:322–328. doi:10.1016/j.tics.2015.04.005
Fahrenfort JJ, Scholte HS, Lamme VAF. 2007. Masking disrupts reentrant
processing in human visual cortex. *Journal of Cognitive Neuroscience* **19**:1488
1497.
doi:10.1162/jocn.2007.19.9.1488&url_ctx_fmt=info:ofi/fmt:kev:mtx:ctx&rft_val_fmt
=info:ofi/fmt:kev:mtx:journal&rft.atitle=masking
Faisal AA, Selen LPJ, Wolpert DM. 2008. Noise in the nervous system. *Nat Rev*
Neurosci **9**:292 303. doi:10.1038/nrn2258
Ferrera VP, Yanike M, Cassanello C. 2009. Frontal eye field neurons signal changes
in decision criteria. *Nat Neurosci* **12**:1458 1462. doi:10.1038/nn.2434
Fisher RA. 1915. Frequency Distribution of the Values of the Correlation Coefficient
in Samples from an Indefinitely Large Population. *Biometrika* **10**:507.
doi:10.2307/2331838
Fleming SM, Dolan RJ. 2012. The neural basis of metacognitive ability. *Philosophical*
Transactions Royal Soc B Biological Sci **367**:1338–1349.
doi:10.1098/rstb.2011.0417
Fleming SM, Thomas CL, Dolan RJ. 2010. Overcoming status quo bias in the human
brain. *Proc National Acad Sci* **107**:6005–6009. doi:10.1073/pnas.0910380107
Garrett DD, Epp S, Kleemeyer M, Lindenberger U, Polk TA. 2020. Higher performers
upregulate brain signal variability in response to more feature-rich visual input.
Neuroimage **217**:116836. doi:10.1016/j.neuroimage.2020.116836
Garrett DD, Kovacevic N, McIntosh AR, Grady CL. 2013a. The Modulation of BOLD
Variability between Cognitive States Varies by Age and Processing Speed. *Cereb*
Cortex **23**:684–693. doi:10.1093/cercor/bhs055
Garrett DD, Kovacevic N, McIntosh AR, Grady CL. 2011. The importance of being
variable. *J Neurosci* **31**:4496 4503. doi:10.1523/jneurosci.5641-10.2011
Garrett DD, Nagel IE, Preuschhof C, Burzynska AZ, Marchner J, Wiegert S,
Jungehülsing GJ, Nyberg L, Villringer A, Li S-C, Heekeren HR, Bäckman L,
Lindenberger U. 2015. Amphetamine modulates brain signal variability and
working memory in younger and older adults. *Proc National Acad Sci* **112**:7593–
7598. doi:10.1073/pnas.1504090112
Garrett DD, Samanez-Larkin GR, MacDonald SWS, Lindenberger U, McIntosh AR,
Grady CL. 2013b. Moment-to-moment brain signal variability: A next frontier in
human brain mapping? *Neurosci Biobehav Rev* **37**:610–624.
doi:10.1016/j.neubiorev.2013.02.015
Gee JW de, Colizoli O, Kloosterman NA, Knapen T, Nieuwenhuis S, Donner TH.
2017. Dynamic modulation of decision biases by brainstem arousal systems. *Elife*
6:309. doi:10.7554/elife.23232

Gee JW de, Knapen T, Donner TH. 2014. Decision-related pupil dilation reflects
upcoming choice and individual bias. *Proc National Acad Sci* **111**:E618 25.
doi:10.1073/pnas.1317557111
Ghosh A, Rho Y, McIntosh AR, Kötter R, Jirsa VK. 2008. Noise during rest enables
the exploration of the brain's dynamic repertoire. *Plos Comput Biol* **4**:e1000196.
doi:10.1371/journal.pcbi.1000196
Grandy TH, Garrett DD, Schmiedek F, Werkle-Bergner M. 2016. On the estimation
of brain signal entropy from sparse neuroimaging data. *Sci Rep-uk* **6**:23073.
doi:10.1038/srep23073
Green DM, Swets JA. 1966. Signal detection theory and psychophysics. *New York:*
Wiley **1**:521.
Hassler U, Trujillo-Barreto N, Gruber T. 2011. Induced gamma band responses in
human EEG after the control of miniature saccadic artifacts. *Neuroimage* **57**:1411
1421. doi:10.1016/j.neuroimage.2011.05.062
Inagaki HK, Fontolan L, Romani S, Svoboda K. 2019. Discrete attractor dynamics
underlies persistent activity in the frontal cortex. *Nature* **566**:212–217.
doi:10.1038/s41586-019-0919-7
Jones SR. 2016. When brain rhythms aren't 'rhythmic': implication for their
mechanisms and meaning. *Curr Opin Neurobiol* **40**:72–80.
doi:10.1016/j.conb.2016.06.010
Joshi S, Gold JI. 2020. Pupil Size as a Window on Neural Substrates of Cognition.
Trends Cogn Sci. doi:10.1016/j.tics.2020.03.005
Klimesch W, Russegger H, Doppelmayr M, Pachinger T. 1998. A method for the
calculation of induced band power: implications for the significance of brain
oscillations. *Electroencephalogr Clin Neurophysiology Evoked Potentials Sect*
108:123 130. doi:10.1016/s0168-5597(97)00078-6
Kloosterman N, Meindertsma T, Hillebrand A, Dijk Bw, Lamme V, Donner T. 2015.
Top-down modulation in human visual cortex predicts the stability of a perceptual
illusion. *J Neurophysiol* **113**:1063–76. doi:10.1152/jn.00338.2014
Kloosterman NA, Gee JW de, Werkle-Bergner M, Lindenberger U, Garrett DD,
Fahrenfort JJ. 2019. Humans strategically shift decision bias by flexibly adjusting
sensory evidence accumulation. *Elife* **8**:e37321. doi:10.7554/elife.37321
Kloosterman NA, Gee JW de, Werkle-Bergner M, Lindenberger U, Garrett DD,
Fahrenfort JJ. 2018. Data from: Humans strategically shift decision bias by flexibly
adjusting sensory evidence accumulation in visual cortex.
doi:https://doi.org/10.6084/m9.figshare.6142940
Kosciessa JQ, Kloosterman NA, Garrett DD. 2020. Standard multiscale entropy
reflects neural dynamics at mismatched temporal scales: What's signal irregularity
got to do with it? *Plos Comput Biol* **16**:e1007885.
doi:10.1371/journal.pcbi.1007885
Lindenberger U, Nagel IE, Chicherio C, Li S-C, Heekeren HR, Bäckman L. 2008.
Age-related decline in brain resources magnifies genetic effects on cognitive
functioning. *Front Neurosci-switz* **2**:234–244. doi:10.3389/neuro.01.039.2008
Linkenkaer-Hansen K, Nikouline VV, Palva JM, Ilmoniemi RJ. 2001. Long-Range
Temporal Correlations and Scaling Behavior in Human Brain Oscillations. *J*
Neurosci **21**:1370–1377. doi:10.1523/jneurosci.21-04-01370.2001
Luck SJ, Woodman GF, Vogel EK. 2000. Event-related potential studies of attention.
Trends Cogn Sci **4**:432–440. doi:10.1016/s1364-6613(00)01545-x
Maris E, Oostenveld R. 2007. Nonparametric statistical testing of EEG- and MEG-
data. *J Neurosci Meth* **164**:177–190. doi:10.1016/j.jneumeth.2007.03.024

Marzen SE, DeDeo S. 2017. The evolution of lossy compression. *J Roy Soc Interface* **14**:20170166. doi:10.1098/rsif.2017.0166

Mišić B, Mills T, Taylor MJ, McIntosh AR. 2010. Brain Noise Is Task Dependent and Region Specific. *J Neurophysiol* **104**:2667–2676. doi:10.1152/jn.00648.2010

Mitra P, Bokil H. 2007. Observed Brain Dynamics - Partha Mitra, Hemant Bokil - Google Books.

Młynarski WF, Hermundstad AM. 2018. Adaptive coding for dynamic sensory inference. *Elife* **7**:e32055. doi:10.7554/elife.32055

Moreno-Bote R, Rinzel J, Rubin N. 2007. Noise-Induced Alternations in an Attractor Network Model of Perceptual Bistability. *J Neurophysiol* **98**:1125 1139. doi:10.1152/jn.00116.2007

Nikulin VV, Brismar T. 2004. Comment on "Multiscale entropy analysis of complex physiologic time series". *Phys Rev Lett* **92**:089803 author reply 089804. doi:10.1103/physrevlett.92.089803

Oostenveld R, Fries P, Maris E, Schoffelen J-M. 2011. FieldTrip: open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. *Comput Intel Neurosc* **2011**:1 9. doi:10.1155/2011/156869

Perrin F, Pernier J, Bertrand O, Echallier JF. 1989. Spherical splines for scalp potential and current density mapping. *Electroen Clin Neuro* **72**:184 187. doi:10.1016/0013-4694(89)90180-6

Rahnev D, Nee DE, Riddle J, Larson AS, D'Esposito M. 2016a. Causal evidence for frontal cortex organization for perceptual decision making. *Proc National Acad Sci* **113**:6059–6064. doi:10.1073/pnas.1522551113

Rahnev D, Nee DE, Riddle J, Larson AS, D'Esposito M. 2016b. Causal evidence for frontal cortex organization for perceptual decision making. *Proc National Acad Sci* **113**:6059–6064. doi:10.1073/pnas.1522551113

Ratcliff R, McKoon G. 2008. The Diffusion Decision Model: Theory and Data for Two-Choice Decision Tasks. *Neural Comput* **20**:873–922. doi:10.1162/neco.2008.12-06-420

Reckless GE, Ousdal OT, Server A, Walter H, Andreassen OA, Jensen J. 2014a. The left inferior frontal gyrus is involved in adjusting response bias during a perceptual decision making task. *Brain Behav* **4**:398–407. doi:10.1002/brb3.223

Reckless GE, Ousdal OT, Server A, Walter H, Andreassen OA, Jensen J. 2014b. The left inferior frontal gyrus is involved in adjusting response bias during a perceptual decision making task. *Brain Behav* **4**:398–407. doi:10.1002/brb3.223

Richman JS, Moorman JR. 2000. Physiological time-series analysis using approximate entropy and sample entropy. *Phys Rev A* **278**:H2039 H2049. doi:10.1103/physreva.29.975

Schurger A, Sarigiannidis I, Naccache L, Sitt JD, Dehaene S. 2015. Cortical activity is more stable when sensory stimuli are consciously perceived. *Proc National Acad Sci* **112**:E2083–E2092. doi:10.1073/pnas.1418730112

Semmlow J. 2014. Signal Processing and Communications. doi:10.1201/9780203024058

Talluri BC, Urai AE, Tsetsos K, Usher M, Donner TH. 2018. Confirmation Bias through Selective Overweighting of Choice-Consistent Evidence. https://www.cell.com/current-biology/fulltext/S0960-9822(18)30982-5

Urai AE, Gee JW de, Tsetsos K, Donner TH. 2019. Choice history biases subsequent evidence accumulation. *Elife* **8**:e46331. doi:10.7554/elife.46331

Valencia JF, Porta A, Vallverdu M, Claria F, Baranowski R, Orłowska-Baranowska E, Caminal P. 2009. Refined multiscale entropy: application to 24-h Holter

recordings of heart period variability in healthy and aortic stenosis subjects. *Ieee*
T Bio-med Eng **56**:2202 2213. doi:10.1109/tbme.2009.2021986
VanRullen R. 2011. Four Common Conceptual Fallacies in Mapping the Time
Course of Recognition. *Front Psychol* **2**:365. doi:10.3389/fpsyg.2011.00365
Wang X-J. 2002. Probabilistic Decision Making by Slow Reverberation in Cortical
Circuits. *Neuron* **36**:955–968. doi:10.1016/s0896-6273(02)01092-9
Waschke L, Tune S, Obleser J. 2019. Local cortical desynchronization and pupil-
linked arousal differentially shape brain states for optimal sensory performance.
Elife **8**:e51501. doi:10.7554/elife.51501
Wimmer K, Nykamp DQ, Constantinidis C, Compte A. 2014. Bump attractor
dynamics in prefrontal cortex explains behavioral precision in spatial working
memory. *Nat Neurosci* **17**:431–439. doi:10.1038/nn.3645
Windmann S, Urbach TP, Kutas M. 2002a. Cognitive and Neural Mechanisms of
Decision Biases in Recognition Memory. *Cereb Cortex* **12**:808–817.
doi:10.1093/cercor/12.8.808
Windmann S, Urbach TP, Kutas M. 2002b. Cognitive and Neural Mechanisms of
Decision Biases in Recognition Memory. *Cereb Cortex* **12**:808–817.
doi:10.1093/cercor/12.8.808
Yarkoni T. 2009. Big Correlations in Little Studies: Inflated fMRI Correlations Reflect
Low Statistical Power-Commentary on Vul et al. (2009). *Perspect Psychol Sci*
4:294 298. doi:10.1111/j.1745-6924.2009.01127.x
Yeung N, Summerfield C. 2012. Metacognition in human decision-making:
confidence and error monitoring. *Philosophical Transactions Royal Soc B*
Biological Sci **367**:1310–1321. doi:10.1098/rstb.2011.0416

1191

1192