Leaderless short ORFs in mycobacteria comprise a translational regulon

Jill G. Canestrari[1], Erica Lasek-Nesselquist[1], Ashutosh Upadhyay[1], Martina Rofaeil[2],

Matthew M. Champion[2], Joseph T. Wade[1], Keith M. Derbyshire[1], Todd A. Gray[1]

[1] Division of Genetics, Wadsworth Center, New York State Department of Health,

Albany, NY 12208

[2] Department of Chemistry and Biochemistry, University of Notre Dame, Notre Dame, IN

46556

Key words:

Operon

Regulon

Attenuation

Cysteine

Mycobacteria

Leaderless mRNAs

Short ORFs

ABSTRACT

Genome-wide transcriptomic analyses have revealed abundant expressed short open

reading frames (ORFs) in bacteria. Whether these short ORFs, or the small proteins

they encode, are functional remains an open question. One quarter of mycobacterial

mRNAs are leaderless, meaning the RNAs begin with a 5'-AUG or GUG initiation

codon. Leaderless mRNAs often encode an unannotated short ORF as the first gene

of a polycistronic transcript. Consecutive cysteine codons are highly

overrepresented in mycobacterial leaderless short ORFs. Here we show that

polycysteine-encoding leaderless short ORFs function as cysteine-responsive

attenuators of operonic gene expression. Through detailed mutational analysis, we

show that one such polycysteine-encoding short ORF controls expression of the

downstream genes by causing ribosome stalling under conditions of low cysteine.

Ribosome stalling in turn blocks mRNA secondary structures that otherwise

sequester the Shine-Dalgarno ribosome-binding site of the 3'gene. This translational

attenuation does not require competing transcriptional terminator formation, a

mechanism that underlies traditional amino acid attenuation systems. We further

assessed cysteine attenuation in *Mycobacterium smegmatis* using mass spectrometry

to evaluate endogenous proteomic responses. Notably, six cysteine metabolic loci

that have unannotated polycisteine-encoding leaderless short ORF architectures

responded to cysteine supplementation/limitation, indicating that cysteine-

responsive attenuation is widespread in mycobacteria. Individual leaderless short

ORFs confer independent operon-level control, while their shared dependence on

cysteine ensures a collective response. Bottom-up regulon coordination is the

antithesis of traditional top-down master regulator regulons and illustrates one

utility of the many unnanotated short ORFs expressed in bacterial genomes.

INTRODUCTION

Short open reading frames (sORFs) are extremely difficult to computationally identify in genome sequences; their shortened gene length approaches statistical random ORF background frequencies and their amino acid sequences have limited bioinformatic value (Frith et al. 2006; Hemm et al. 2008; Hobbs et al. 2011; Crappe et al. 2013). Conventional mass spectrometry proteomic studies also systematically underrepresent the small proteins (sproteins) encoded by sORFs, as they can be lost during sample preparation or provide too few detectable peptides (Hemm et al. 2010). These limitations have contributed to both a lag in (i) recognition of sORFs, and (ii) assessment of their functional potential. This knowledge gap has been underscored by recent descriptions of many potential novel sORFS in *Escherichia coli* and in bacteria found in the human microbiome (Meydan et al. 2019; Sberro et al. 2019; Weaver et al. 2019). Previous work applying complementary transcriptomic approaches to mycobacteria, in both slow-growing *Mycobacterium tuberculosis*, and fast-growing *Mycobacterium smegmatis* and *Mycobacterium abscessus* suggested that their genomes contained hundreds of sORFs actively producing sproteins (Shell et al. 2015; Miranda-CasoLuengo et al. 2016). Ribosome profiling (Ribo-seq), together with RNA-seq and transcription start site (TSS) mapping, provided genome-wide empirical evidence for the location and ribosome occupancy of mycobacterial mRNAs that express sproteins (Cortes et al. 2013; Shell et al. 2015; Miranda-CasoLuengo et al. 2016).

Many ORFs initiated by leaderless mRNAs (LL-mRNAs) in mycobacteria are short

(defined here as less than 150 nt) and unannotated. Importantly, the ORFs initiated at the

combined transcription/translation initiation start sites of LL-mRNAs are readily

identifiable from transcriptomic data sets and thus provide a high-confidence list of

expressed novel mycobacterial sORFs. Insights into the mechanistic and functional

attributes of LL-mRNAs have lagged, as they are rare or poorly expressed in *E. coli* but

are abundant in archaea, Actinobacteria and extremophiles (Beck and Moll 2018). LL-

sORFs represent the first (5'-most) ORF of a transcript, which positions them for a role in

cis-regulation of the downstream operonic genes. There is ample precedent in eukaryotes

for regulation of downstream genes by such "upstream ORFs" (uORFs) (Hinnebusch et

al. 2016; Couso and Patraquim 2017). In prokaryotes, mechanisms have also been

previously described in which uORFs have been shown to regulate expression of

downstream genes through a process known as attenuation (Oppenheim and Yanofsky

1980; Bechhofer 1990; Henkin and Yanofsky 2002).

Attenuation is a *cis*-regulatory mechanism often mediated by short uORFs enriched in

codons for the amino acid product of that biosynthetic operon. Attenuation occurs when

abundant charged tRNA levels allow translating ribosomes to quickly clear the

modulating uORF, promoting the formation of an intrinsic terminator that aborts

transcription of the operon. Low levels of charged tRNA cause ribosome stalling in the

uORF at codons for the end-product amino acid, facilitating the formation of a competing

anti-terminator structure, thereby releasing attenuation to allow transcription to extend

into the biosynthetic operon (Turnbough 2019). uORF-mediated attenuation mechanisms

for cysteine have not been described. A subset of predicted mycobacterial LL-sORFs

conspicuously encode consecutive cysteine residues, and these were found upstream of

genes annotated to be involved in cysteine biosynthesis (Shell et al. 2015). We

hypothesized that these LL-sORFs function as cysteine-sensitive attenuators.

RESULTS

We identified 304 putative LL-sORFs in *Mycobacterium smegmatis* (Supplementary

Information Table 1(Shell et al. 2015; Martini et al. 2019)). We compared amino acid

content of the encoded sproteins and found that consecutive cysteines were

overrepresented relative to cysteine content (chi-square $p<.01$, Extended Data Table 1)

and relative to consecutive cysteine frequency in annotated genes. A subset of predicted

mycobacterial LL-sORFs conspicuously encode consecutive cysteine residues, and these

were often found upstream of genes annotated to be involved in cysteine biosynthesis

(Shell et al. 2015). We hypothesized that these LL-sORFs function as cysteine-sensitive

attenuators.

***Ms5788* is regulated in response to cysteine abundance**

One predicted LL-sORF (here denoted Ms5788A) encodes eight consecutive cysteines in

its C-terminus, and is followed by operonic genes, including a putative thiosulfate

sulfurtransferase, *cysA2* (Fig 1A). RNA-seq and Ribo-seq profiles indicate that this

unannotated LL-sORF is abundantly transcribed and translated in *M. smegmatis*

(Extended Data Fig 1 A). Moreover, transcription start site mapping strongly suggests

that a homologous LL-sORF is expressed in *M. tuberculosis* (Extended Data Fig 1B). To

determine whether the genes located immediately downstream of *Ms5788A* are regulated

in response to changes in cysteine levels, we generated a luciferase translational reporter

in which a constitutive promoter drives a leaderless transcript that begins at the native

GUG initiation codon of *Ms5788A* and continues to the initiation codon of the annotated

gene downstream, *Ms5788* (Fig 1B). We then measured expression of the reporter in *M.*

smegmatis cells in the presence or absence of cysteine in the growth medium. Luciferase

activity decreased for cells grown with cysteine supplementation (Fig 1B i). Hence, we

hypothesized that expression of *Ms5788* is regulated in response to cysteine levels by an

attenuation mechanism involving the upstream LL-sORF *Ms5788A*.

We next tested whether cysteine-dependent regulation of *Ms5788* requires translation of

the upstream *Ms5788A* LL-sORF. We mutated the GUG translation initiation codon of

Ms5788A to ACC in the context of the luciferase reporter, to prevent ribosome loading.

This non-start mutation reduced luciferase expression, and abolished attenuation (Fig 1B

compare i vs ii). These data indicate that translation of *Ms5788A* is required for cysteine-

dependent regulation of *Ms5788*, and that in the absence of *Ms5788A* translation,

expression of *Ms5788* is locked in an attenuated state. We speculated that ribosome

occupancy of the C-terminal polycysteine tract of *Ms5788A* is particularly important for

this regulation. We created a nonsense mutant, Ser8Stop, to truncate *Ms5788A* ten amino

acids before the first Cys codon. This truncating mutation also dramatically reduced

luciferase expression and attenuation by cysteine (Fig 1B iii). The residual cysteine

response of this nonsense mutant may result from translation reinitiation after the stop

codon.

Since ribosome occupancy of the LL-sORF appeared to be required for cysteine-

dependent regulation of the luciferase reporter, we postulated that ribosomes stalled in

the *Ms5788A* polycysteine tract due to limiting levels of charged tRNA^{cys} would increase

luciferase expression by relieving attenuation. We hypothesized that recoding the

polycysteine tract should impair the observed cysteine response. We created an out-of-

frame (OoF) *Ms5788A* mutant luciferase reporter to replace the eight consecutive

cysteine codons with eight consecutive leucine codons, leaving a single Cys18 codon

(Fig 1B iv). This OoF mutant was not affected by cysteine supplementation, indicating

the importance of the polycysteine tract in relieving attenuation. Interestingly, expression

of this reporter appears to be in the active state, suggesting that limiting leucine may

functionally substitute for limiting cysteine.

Ribosome stalling in the *Ms5788A* sORF modulates RNA structure in the *Ms5788* 5'

UTR

Ribosome occupancy of *Ms5788A* could affect the formation of mRNA secondary

structures in the *Ms5788* mRNA leader, as occurs in previously described attenuation

mechanisms (Turnbough 2019). The predicted RNA secondary structure for the mRNA

through the *Ms5788A* LL-sORF and up to *Ms5788*, indicates the potential for an

energetically stable structure over most of its length (Fig 2). Importantly, nucleotides in

the polycysteine tract of *Ms5788A* are predicted to base pair with complementary

sequences near the Shine-Dalgarno sequence of *Ms5788*. This structure suggests a

mechanism in which stalled ribosomes in *Ms5788A* free the Shine-Dalgarno sequence to

recruit and position ribosomes for canonical translation initiation of *Ms5788*.

Whereas the *Ms5788A* OoF mutant was predicted to have no effect on mRNA structure,

we created mutants intended to selectively disrupt predicted duplex pairing near the

Ms5788 Shine-Dalgarno sequence (Fig 2). We first changed the invariant guanine in the

2^{nd} position of cysteine codons (U<u>G</u>Y) to cytosine (U<u>C</u>Y) in the last five codons of

Ms5788A (Fig 2, recoded red series bottom strand). The nucleotide changes should

reduce the stability of base pair interactions with the Shine-Dalgarno region, while

recoding polyserine for the final five codons of *Ms5788A*: Cys(26-30)Ser. This reporter

exhibited constitutively elevated expression that was insensitive to cysteine (Fig 2 ii),

consistent with full Shine-Dalgarno sequence accessibility.

To differentiate the effect of RNA duplex formation from the effect of *Ms5788A* amino

acid recoding, we created a mutant that only disrupted the base pairing, by changing the

predicted five cognate nucleotides near the Shine-Dalgarno sequence (Fig 2, recoded red

top strand). Even though the polycysteine tract remained intact, luciferase expression

from this reporter was insensitive to cysteine supplementation (Fig 2 iii), consistent with

the base-paired structure corresponding to the attenuated state. We next constructed a

mutant that combined the recoded *Ms5788A* and cognate non-coding mutants, such that

base pairing should be restored. As expected, combining the complementary mutations

reduced expression of the luciferase reporter for cells grown in the presence of cysteine,

consistent with restored base pairing (Fig 2 iv). Interestingly, the restored response to

cysteine indicated that the residual four-cysteine codon content of the LL-sORF was

sufficient to confer sensitivity to cysteine.

We constructed a silent *Ms5788A* mutant that switched the nucleotide sequence of six

cysteine codons to retain polycysteine coding, and yet are predicted to disrupt base

pairing with the Shine-Dalgarno region (Fig 2 polycysteine purple bottom strand). The

elevated luciferase activity of this mutant reporter was not attenuated under cysteine-

replete conditions, separating the roles of nucleotide and encoded amino acid sequence of

Ms5788A (Fig 2 v). We also created a mutant of the predicted complementary bases near

the Shine-Dalgarno sequence (Fig 2, polycysteine purple top strand). Surprisingly, this

mutant exhibited an attenuation response similar to wild type (Fig 2, compare i and vi).

We reassessed the base pairing potential of the mRNA produced from this mutant and

found that it was predicted to fold into a stable, wild-type-like structure that would also

reduce Shine-Dalgarno sequence availability (Extended Data Fig 2). Combining the

Ms5788A and peri-Shine-Dalgarno nucleotide changes in this series resulted in an

expression pattern similar to that of the wild-type construct, consistent with the model

(Fig 2 vii).

Cysteine-dependent regulation involving *Ms5788A* affects expression of downstream

operonic genes

Classical models of ribosome-mediated attenuation invoke competing mRNA stem loops

that form an intrinsic terminator structure when ribosomes rapidly translate the sORF,

resulting in regulation at the level of transcription (Yanofsky 1981; Turnbough 2019). In

the case of *Ms5788* attenuation, regulation appears to occur at the level of translation. To

test this, we created a reporter to assess mRNA extension beyond the start of *Ms5788*. To

maintain the predicted RNA structure of the 5' leader region, an independent Shine-

Dalgarno sequence was added to efficiently initiate translation of transcripts that extend

into the luciferase gene (Fig 3 ii). Luciferase activity of this reporter was insensitive to

cysteine, and expression levels were consistently high. These data indicate that *Ms5788*

attenuation does not result from transcription termination. Translational repression can

indirectly affect transcription over longer distances due to polarity that is caused by Rho-

dependent transcription termination and/or by enhanced RNase processing of the

untranslated RNA (Deana and Belasco 2005; Martini et al. 2019). To determine if

translational repression of *Ms5788* by attenuation leads to polar effects on downstream

genes, we constructed a translational fusion that extends to *Ms5789* (*cysA2*). This more

distal site, 516 nt 3' of the *Ms5788* start, exhibited cysteine responsiveness (Fig 3 iii).

Taken together, our data support a model in which *Ms5788* is regulated by translational

attenuation through *Ms5788A*-controlled Shine-Dalgarno availability, while *Ms5789* is

likely regulated by polarity, due to the absence of elongating ribosomes in *Ms5788*.

Cysteine-dependent regulation of *Ms5789* and *Ms5790* by *Ms5788A* occurs in a

chromosomal context

To assess whether our reporter-supported model is valid in the native locus context, we

performed quantitative mass spectrometry-based proteomics (LFQ) to determine

differences in protein expression for cells grown with or without cysteine

supplementation. Whole cell extracts were prepared from cultures of *M. smegmatis*

grown in minimal media +/- cysteine supplementation. Tryptic digests of whole cell

lysates were subjected to nanoUHPLC-MS/MS identified and quantitated using label-free

based peak integration (Cox and Mann 2008; Bosserman et al. 2017). As expected, the

abundance of most proteins is unchanged between the two conditions (+/- cysteine),

reflected in the linear correlation on the diagonal of the scatter plot (Fig 4A). Proteins

below the diagonal were more abundant in cells grown without cysteine supplementation.

The small (159 AA) and hydrophobic Ms5788 was not detected in these experiments.

However, *Ms5789* and *Ms5790* are predicted to be co-transcribed in an operon with

Ms5788 (Martini et al. 2019), and expression of the encoded proteins Ms5789 and

Ms5790 was higher in cells grown without cysteine (Fig 4A, yellow diamonds),

consistent the attenuation model.

To test the hypothesis that polycysteines in *Ms5788A* control Ms5789 and Ms5790

expression, we generated an out-of-frame (OoF) mutation in *Ms5788A* by deletion of a

single nucleotide from the chromosomal locus. This deletion shifts the reading frame to

encode valines and alanines in place of the polycysteine tract, and adds six additional

amino acids (ERSRAL) prior to encountering a stop codon (Fig 4B), while preserving the

potential for nearly wild-type RNA duplex formation. This mutant exhibited low Ms5789

and Ms5790 expression levels consistent with a stable mRNA leader structure

sequestering the Shine-Dalgarno of *Ms5788*. Expression of Ms5789 and Ms5790 was

unaffected by cysteine supplementation (Fig 4B, yellow diamonds in diagonal),

highlighting the role of cysteine codons in relieving attenuation.

We reasoned that if *Ms5788A* directs attenuation, its deletion would elevate operon

expression of the downstream genes. We created a precise deletion of *Ms5788A* and

subjected this mutant to LFQ proteomics. As predicted, Ms5789 and Ms5790 were no

longer responsive to ambient cysteine supplementation, appearing in the diagonal of

unresponsive genes (Fig 4C, yellow diamonds in diagonal). Moreover, absolute levels of

Ms5789 and Ms5790 were higher in the Δ*5788A* strain than in either the wild-type or the

OoF mutant, regardless of cysteine supplementation. This indicates that the *Ms5788A*

deletion leaves the *Ms5788* Shine-Dalgarno sequence fully available for canonical

translation initiation, and results in an elevated expression of the operonic Ms5789 and

Ms5790 (Fig 4B, C yellow diamonds). Thus, data from wild-type and targeted

chromosomal mutant *M. smegmatis* are in agreement with our reporter-generated data,

and strengthen the conclusion that the *Ms5788A* LL-sORF modulates operonic gene

expression through an attenuation mechanism.

Widespread cysteine-dependent regulation in *M. smegmatis* associated with cysteine-

rich LL-sORFs

Given that LL-sORFs in *M. smegmatis* are enriched for polycysteine, we hypothesized

that additional cysteine-responsive genes are regulated by an associated polycysteine LL-

sORF. We identified six more LL-sORFs that contain at least two consecutive cysteine

codons and are located upstream of annotated genes. We then examined our proteomic

data for the protein levels encoded by putative operonic genes downstream of these LL-

sORFs for cells grown with/without cysteine supplementation. Remarkably, all of the

detectable proteins that were encoded by polycysteine LL-sORF-led operons were

upregulated, falling below the diagonal (Fig 4A, black circles), consistent with

attenuation release. In cysteine-replete medium, expression of some of these proteins

(Ms0113, Ms0934, Ms4527, Ms5279, and Ms5280) was below the threshold of detection

for reliable quantification [$<10^5$ LFQ intensity (a.u.)], indicative of tight attenuation. The

polycysteine LL-sORFs exhibit RNA-seq and Ribo-seq expression profiles consistent

with their robust expression during cysteine replete growth (Extended Data Figs 1 and 3).

None of these expressed LL-sORFs were identified by genome annotation pipelines.

Each of these cysteine-responsive operons contains genes annotated for cysteine

associated activities (Table 1). Collectively, these data reveal a cysteine-metabolic

regulon, whose concerted response is controlled independently at each locus by an

expressed LL-sORF that includes consecutive cysteine codons. This bottom-up

coordinated regulation contrasts with the conventional top-down, master regulator-driven

mechanism of transcriptional regulons.

LFQ proteomics does not comprehensively identify all proteins in a proteome. We looked

for additional loci with the same hallmarks of responsive attenuation: an expressed

polycysteine LL-sORF upstream of annotated operonic genes. *Ms4536A* was identified

upstream of a single gene (Table 1, Extended Data Fig 3 E). Without corroborating

indicators of cysteine response or function of the encoded operon protein, we only

speculate at its membership in the *M. smegmatis* cysteine LL-sORF regulon. The

independent evolution of LL-sORFs makes it unlikely that our experimental reference

species, *M. smegmatis*, contains all of the LL-sORF operons in the mycobacterial pan-

genome. In one example, transcriptomic profile data for *M. tuberculosis* clearly identified

Rv2334A as an unannotated gene meeting all of the criteria demonstrated in the *M.*

smegmatis regulon: an expressed LL-sORF with a C-terminal polycysteine tract,

followed by genes annotated as *cysK1* and *cysE* (Table 1, Extended Data Fig 3 G).

The availability of complete genome sequence information on diverse mycobacteria

provided an opportunity to track the evolution of these polycysteine sORFs. We searched

the genomes of 41 species using complementary approaches predicated on the sequence

of each *M. smegmatis* or *M. tuberculosis* LL-sORF, or on the position of the flanking

annotated orthologous genes as landmarks. We identified genomic sequences consistent

with conservation of the LL-sORFs expressed in *M. smegmatis* and *M. tuberculosis*

(Extended Data Table 2). The distribution of the conserved LL-sORFs is summarized as

barcodes adjacent to each species on the *Mycobacterium* genus phylogenetic tree (Fig 5).

Ms4527A, Ms4533A, and Ms5788A are deeply rooted, indicating both an emergence that

predates mycobacterial radiation and a selective advantage to retaining these sequences.

The presence/absence of others (e.g., *Ms0932A* or *Ms5280A*) is consistent with horizontal

gene transfer or sporadic gene loss by deletion or degradation. Sequence logos derived

from multiple alignments of the amino acid sequences encoded by the LL-sORFs further

support the evolutionary selection of their consecutive cysteine tracts (Fig 5). The

similarity between Ms4536A and Rv2334A strongly suggests homology by common

origin, yet the context, including operon genes, is not homologous. The occurrence of

Ms4536A—TQXA or Rv2334A—cysK1-cysE is mutually exclusive, suggesting origin

by rearrangement rather than a merodiploid duplication or horizontal gene transfer event.

Non-cysteine amino acids are also conserved in some LL-sORFs, suggesting that they

provide function through ribosomal interaction, or support activities as an independent

small protein product, or are encoded by codons that are constrained at the nucleotide

level.

DISCUSSION

An attenuation mechanism that controls translation in response to amino acid

availability

In the work presented here, we demonstrate attenuation of a cysteine biosynthesis pathway

locus. Attenuation is a recurring theme in biosynthetic pathways for nucleosides and amino

acids, in which the end product of the pathway interacts with the 5' leader of the

biosynthesis-encoding transcript to reduce (attenuate) expression of the operonic ORFs

downstream (Turnbough 2019). However, these models typically involve the formation of

competing alternate hairpin structures that function as an intrinsic terminator when the end

product is plentiful. By contrast, the *Ms5788A* LL-sORF featured here defines a class of

translational attenuator that indirectly assesses charged tRNA^{cys} availability to modulate

expression of the downstream operonic genes.

The need for cysteine regulation in mycobacteria

Why might mycobacteria need a multi-locus cysteine regulon? Mycobacteria do not

produce glutathione, which modulates redox balance in most bacteria. Instead, they rely on

mycothiol (MSH), a cysteine derivative (Xu et al. 2011; Loi et al. 2015). The *mshA* gene

(*Ms0933*) encodes the first enzyme in MSH biosynthesis, and it resides in an operon with

the hallmarks of cysteine attenuation (Extended Data Fig 3 B). The multiple roles of

cysteine in mycobacterial protein synthesis, redox and sulfur metabolism likely require

subtle, independent fine-tuning of the enzymes involved in these respective pathways. The

slight differences in cysteine composition and architectures of the polycysteine LL-sORFs

could impart varying mechanisms as well as customized levels of operon gene expression.

Attenuating sORF requirements

What are the cysteine codon requirements of an effective small ORF attenuator? Our

criterion that LL-sORFs encode two consecutive cysteine codons may seem to be a low

threshold, yet the *cis*-encoded proteins only detectable in cysteine-limiting conditions

(baseline of Fig 4A) demonstrate the effectiveness of two consecutive cysteines in LL-

sORFs in relieving attenuation. Our Ms5788A LL-sORF analysis demonstrated that base

pairing was needed to impose attenuation, but it is premature to speculate that it is required

at all responsive loci. Transcription activation was not a regulatory factor at the *Ms5788*

locus; the promoter remained intact in the *Ms5788A* mutants in Fig 4 B, C, indicating that

all of the cysteine response at this locus is directed by the attenuation mechanism we

detailed and not via transcriptional activation. RNA-seq profiles of the LL-sORFs

presented here are also consistent with robust constitutive transcription in cysteine-replete

medium.

Evolution of a cysteine attenuation regulon

The independent evolution of similar polycysteine LL-sORF architecture associated with cysteine attenuation at multiple loci in mycobacteria indicates that coordinated expression is important, and that LL-sORF directed attenuation is effective. The co-regulation of individual operons functionally defines a regulon, akin to regulons controlled by dedicated DNA-binding transcription factors. We speculate that the evolution of regulation by attenuation is simplified in mycobacteria by the robust nature of leaderless translation, and that the evolution of a dedicated transcription factor and its cognate binding sites in the promoters of target operons is more problematic than exploiting LL-sORFs in a genus that exhibits frequent and robust LL-mRNA expression. As the first genes in their transcripts, LL-mRNAs are ideally positioned to *cis*-regulate expression of downstream genes. Additionally, transcription start sites are preferentially associated with purines (R), and the +2 transcript position is preferentially associated with pyrimidines (Y) (Martini et al. 2019). Thus, transcription often begins at RYN trinucleotide sequences. Our previous study showed that a 5' RUG trinucleotide is both necessary and sufficient for robust leaderless translation initiation (Shell et al. 2015), so many transcription start sites are predicted to already initiate leaderless translation, and others are only one or two changes from initiating translation. It is not yet clear whether leaderless architecture *per se* offers advantages for attenuation and may have been selected over canonical Shine-Dalgarno translation initiation, or whether the sole criterion of an RUG sequence at the transcription start site is simply a relatively modest requirement. It is clear, however, that in mycobacteria, polycysteine LL-sORFs integrate two levels of coordination: locally by

modulating polycistronic operon gene expression, and globally by synchronizing operon

response.

Concluding remarks

Given the prevalence of LL-sORFs in mycobacteria, we speculate that other translational

regulons have evolved as an alternative to transcriptional regulons controlled by DNA-

binding transcription factors. LL translation is considered to be the ancestral form of

ribosome delivery (Nakamoto 2009; Zheng et al. 2011; Duval et al. 2013), indicating that

LL-sORF regulons may be ancient and widespread. LL-sORFs define a functional

subclass of small *de novo* genes that effectively decode translational stress into broad

regulatory effects.

METHODS

Bacterial strains and culture

M. smegmatis wild-type mc^{2}155 and its derivatives were grown in tryptic soy broth +

0.05% Tween 80 (TSBT) or on TSA plates, and cultured at 37°C. Antibiotic selection for

reporter maintenance or mutation selection strategies included apramycin (12.5 µg/ml on

agar, 10 µg/ml in broth), hygromycin (100 µg/ml and 25 µg/ml), kanamycin (50 µg/ml

and 10 µg/ml), and zeocin (50 µg/ml and 25 µg/ml).

For the cysteine attenuation study, bacteria were cultured in minimal media. Base

medium per liter: 6g Na$_2$HPO4 (anhydrous), 3g KH$_2$PO4, 0.5g NaCl, 1g NH$_4$Cl, 0.05%

(v/v) Tween-80. After autoclaving, 0.2 % glucose and micronutrients were added to final

concentrations: $MgSO_4$ to 1 mM, $CaCl_2$ to 100 µM, H_3BO_3 to $4x10^{-7}$ M, $CoCl_2.6H_2O$ to

$3x10^{-8}$ M $CuSO_4.5H_2O$ to $1x10^{-8}$ M, $MnCl_2.4H_2O$ to $8x10^{-8}$ M, $ZnSO_4.7H_2O$ to $1x10^{-8}$

M, $FeSO_4.7H_2O$ to $1x10^{-6}$ M. L-cysteine or L-cystine (exogenously stable dimeric

cysteine) was supplemented at 200 µg/ml as noted.

Luciferase assays

Reporters were generated by long-primer-dimer PCR to recreate the LL-sORF and leader

sequence of *M. smegmatis msmeg_5788*. The products were cloned by Infusion and

verified by DNA sequence analysis. The NanoLuc (Promega) luciferase ORF is carried

on a plasmid that confers apramycin resistance and integrates at the L5 *attB* site in

mycobacteria. Mycobacterial cultures were grown in minimal media for luciferase assays.

Luciferase activity in a culture was assessed by the addition of NanoGlo (Promega)

substrate and then measuring luminescence normalized to culture density.

Chromosomal mutants of *M. smegmatis*

A precise deletion of *Ms5788A* was created using a targeting plasmid that integrates via a

single cross-over allowing selection of a hyg^r intermediate and, then, is resolved by a

second homology-driven event in which a *sacB/galK* counter-selection allows enrichment

for the deleted recombinant (Barkan et al. 2011). The OoF point mutant was created by a

recombineering approach that used a single-stranded oligonucleotide template to

introduce an additional adenine in *Ms5788A* (van Kessel and Hatfull 2008). Co-

electroporation of 2×10^{-10} mol of 60-mer oligo with 500 ng of an episomal zeo^r plasmid

(pGE324, Zeo-sacB) allowed zeocin selection and isolation of the electrocompetent

population of *M. smegmatis*. Mismatch-sensitized PCR (MAMA-PCR) screening (Cha et

al. 1992; Swaminathan et al. 2001) of isolates identified clones that integrated the

additional adenine. Deletion and point mutants were verified by genomic DNA PCR and

sequencing.

Mass spectrometry

Wild-type and mutant derivative *M. smegmatis* were cultured in minimal media with or

without cysteine supplementation, harvested by centrifugation and cryo-milled (Retsch

MM400, Haan, Germany) for mass spectrometry analysis. Reagents were of LC-MS

quality or higher and obtained from Sigma Aldrich unless indicated. Milled cell pellets

were digested with trypsin using commercial S-Traps (Protifi, NY). Briefly 50 μg of

protein from each milled cell pellet was re-suspended in 6% SDS 10mM Tris-2-Carboxy

ethyl phosphine in 100 mM tri ethyl ammonium bicarbonate (TEAB), heated at 95 ºC for

3minutes, then alkylated with 10 mM iodoacetamide in the dark for 15 min. Samples

were acidified by addition of H_3PO4_4 to 1.2% Final (v/v) and flocculated by 7-fold

addition of 95/5 MeOH:100 mM TEAB prior to collection on the S-trap column

(Zougman et al. 2014). Washing and conditioning was preformed three times by 150 μl

addition of MeOH buffer as above and 1 μg of sequencing grade trypsin (Promega, WI)

was added to each sample and digested at 37 ºC for 8 hours. Peptides were isolated,

acidified and desalted using Stage tips packed into P100 pipette tips, and dried using a

MiVac (Genvac UK) prior to LC-MS/MS analysis (Rappsilber et al. 2003).

NanoUHPLC-MS/MS was performed essentially as described (Bosserman et al. 2017;

Bosserman et al. 2019). 1μg of each digest was analyzed in technical triplicate and

biological duplicate on an Orbitrap instrument running a TOP15 data-dependent

acquisition (Q-Exactive Thermo San Jose, CA). Protein spectral matching and Label Free

quantification were performed using MaxQuant (Cox and Mann 2008) against the *M.*

smegmatis FASTA combined with contaminants from the Uniprot database, LFQ param

(UP000000757 6,595 entries). LFQ parameters were set to default, quantification was

restricted to proteins >2 peptides (missing peaks enabled). Target-decoy was used to

determine False Discovery Rates (Elias and Gygi 2007) and proteins at a global 1% FDR

were used for quantification. Data reduction and significance testing were performed

using a modified LIMMA methodology (Efstathiou et al. 2017). Protein search and RAW

data files are accessible at the Center for Computational Mass Spectrometry via

ftp://MSV000084381@massive.ucsd.edu (Deutsch et al. 2017). Proteins quantitatively

replicated in one condition but absent in another, were given an arbitrary intensity of

5×10^4 (below detection threshold) for ease in visualization.

Bioinformatics

Phylogeny

The genomes and proteomes for *Mycobacterium abscessus* subsp. massiliense

(NC_018150.2), *M. africanum* strain 25 (CP010334,1), *M. litorale* strain F4

(CP019882.1), *M. avium* 104 (NC008595.1), *M. ulcerans* Agy99 (NC_008611.1), *M.*

vanbaalenii PYR-1 (NC_008726.1), *M. marinum* M (NC_010612.1), *M. liflanddii*

128FXT (NC_020133.1), *M. kansasii* ATCC 12478 (NC_022663.1), *M. gilvum* Spyr1

(NC_014814.1), *M. bovis* AF2122/97 (NC_002945.4), *M. tuberculosis* H37Rv

(NC_000963.3), *M. sinense* strain JDM601 (NC_015576.1), *M. canettii* CIPT 140010059

(NC_015848.1), *M. chubuense* NBB4 (NC_018027.1), *M. intracellulare* MOTT-64

(NC_016948.1), *M. intracellulare* ATCC 13950 (NC_016946.1), *M. neoaurum* VKM ac-

1815D (NC_023036.2), *M. haemophilum* ATCC 29548 (NZ_CP0118883.2), *M. simiae*

ATCC 25275 (NZ_HG315953.1), *M. goodii* strain X7B (NZ_CP012150.1), *M. fortuitum*

strain CT6 (NZ_CP011269.1), *M. phlei* strain CCUG 21000 (NZ_CP014475.1), *M.*

immunogenum strain CCUG 47286 (NZ_CP011530.1), *M. chelonae* CCUG 47445

(NZ_CP007220.1), *M. vaccae* 95051 (NZ_CP011491.1), *M. chimaera* strain AH16

(NZ_CP012885.2), *M. caprae* strain Allgaeu (NZ_CP016401.1), *M. colombiense* CECT

3035 (NZ_CP020821.1), *M. dioxanotrophicus* strain PH-06 (NZ_CP020809.1), *M.*

marseillense strain FLAC0026 (NZ_CP023147.1), *M. lepraemurium* strain Hawaii

(NZ_CP021238.1), *M. shigaense* strain UN-152 (NZ_AP018164.1), *M. stephanolepidis*

(NZ_AP018165.1), *M. pseudoshottsii* JCM 15466 (NZ_AP018410.1), *M. paragordonae*

49061 (NZ_CP025546.1), *M. rutilum* strain DSM 45405 (NZ_LT629971.1), *M.*

thermoresistibile strain NCTC10409 (NZ_LT906483.1), *M. hassiacum* DSM 44199

(NZ_LR026975.1), and *M. microti* strain 12 (CP010333.1) were downloaded from NCBI.

Orthologs of *M. smegmatis* proteins were identified by a reciprocal best blast hit

approach and aligned in MAFFT v.7.058b (Katoh and Standley 2013). A supermatrix

was created from the concatenation of 1560 of these protein alignments with an in-house

Python script. Alignments included in the supermatrix were required to have 40-41 of

the 41 mycobacterial species present. IQ-TREE v.1.6.9 (Nguyen et al., 2015) generated a

maximum likelihood (ML) phylogeny under an LG+R4 model of evolution (Soubrier et

al., 2012; Yang 1995; Le and Gascuel 2008). Support values were generated by the

ultrafast bootstrap method with 1000 replicates (Minh et al., 2013). The ML tree was

visualized in FigTree v.1.4.4 (http://tree.bio.ed.ac.uk/software/figtree/).

Identification of ssORFs in additional mycobacterial genomes

The orthologs to all *M. smegmatis* ORFs with an upstream ssORF were identified in other

Mycobacteria via a reciprocal best blast hit approach and the regions 1000 nucleotides

upstream of these genes were extracted. *M. smegmatis* ssORF proteins were queried

against these upstream regions via TBLASTN searches to identify orthologous

sequences. The coordinates for these sequences were obtained from the blast results and

extended to the match the length of the corresponding *M. smegmatis* ssORF. To ensure

that we captured the start and stop codon, we extended these coordinates by an additional

5-10 amino acids on both 5' and 3' ends. These coordinates and the strand information

captured from the blast results were used to extract and translate ssORFs from 1 Kb

upstream regions with an in-house Python script. All ssORFs were visually inspected

and initially aligned in MEGA v. 70.26 (Kumar et al., 2016). In some cases, additional

ssORF orthologs were identified by predicting proteins (anything between two stop

codons) in upstream 1 Kb regions with the getorf subroutine from EMBOSS v. 6.3.1

(Rice et al., 2000).

Multiple alignments of deduced LL-sORF amino acid sequences were performed using a

web-based tool and default clustal algorithm (Madeira et al. 2019). The aligned

sequences were compiled into a sequence logo infographic (weblogo.berkeley.edu).

Supplementary Information Table 1. LL-sORFs predicted by transcriptomic data: Ribo-

seq, RNA-seq, and RUG transcription start sites (Shell et al. 2015; Martini et al. 2019).

REFERENCES

Barkan D, Stallings CL, Glickman MS. 2011. An improved counterselectable
marker system for mycobacterial recombination using galK and 2-deoxy-
galactose. *Gene* **470**: 31-36.
Bechhofer DH. 1990. Triple post-transcriptional control. *Mol Microbiol* **4**: 1419-
1423.
Beck HJ, Moll I. 2018. Leaderless mRNAs in the Spotlight: Ancient but Not
Outdated! *Microbiol Spectr* **6**.
Bosserman RE, Nguyen TT, Sanchez KG, Chirakos AE, Ferrell MJ, Thompson CR,
Champion MM, Abramovitch RB, Champion PA. 2017. WhiB6 regulation of
ESX-1 gene expression is controlled by a negative feedback loop in
Mycobacterium marinum. *Proc Natl Acad Sci U S A* **114**: E10772-E10781.
Bosserman RE, Nicholson KR, Champion MM, Champion PA. 2019. A New ESX-1
Substrate in Mycobacterium marinum That Is Required for Hemolysis but
Not Host Cell Lysis. *J Bacteriol* **201**.
Cha RS, Zarbl H, Keohavong P, Thilly WG. 1992. Mismatch amplification
mutation assay (MAMA): application to the c-H-ras gene. *PCR Methods*
Appl **2**: 14-20.
Cortes T, Schubert OT, Rose G, Arnvig KB, Comas I, Aebersold R, Young DB.
2013. Genome-wide mapping of transcriptional start sites defines an
extensive leaderless transcriptome in Mycobacterium tuberculosis. *Cell*
Rep **5**: 1121-1131.
Couso JP, Patraquim P. 2017. Classification and function of small open reading
frames. *Nat Rev Mol Cell Biol* **18**: 575-589.
Cox J, Mann M. 2008. MaxQuant enables high peptide identification rates,
individualized p.p.b.-range mass accuracies and proteome-wide protein
quantification. *Nat Biotechnol* **26**: 1367-1372.
Crappe J, Van Criekinge W, Trooskens G, Hayakawa E, Luyten W, Baggerman G,
Menschaert G. 2013. Combining in silico prediction and ribosome
profiling in a genome-wide search for novel putatively coding sORFs.
BMC Genomics **14**: 648.
Deana A, Belasco JG. 2005. Lost in translation: the influence of ribosomes on
bacterial mRNA decay. *Genes Dev* **19**: 2526-2533.
Deutsch EW, Csordas A, Sun Z, Jarnuczak A, Perez-Riverol Y, Ternent T,
Campbell DS, Bernal-Llinares M, Okuda S, Kawano S et al. 2017. The
ProteomeXchange consortium in 2017: supporting the cultural change in
proteomics public data deposition. *Nucleic Acids Res* **45**: D1100-D1106.
Duval M, Korepanov A, Fuchsbauer O, Fechter P, Haller A, Fabbretti A, Choulier
L, Micura R, Klaholz BP, Romby P et al. 2013. Escherichia coli ribosomal
protein S1 unfolds structured mRNAs onto the ribosome for active
translation initiation. *PLoS biology* **11**: e1001731.
Efstathiou G, Antonakis AN, Pavlopoulos GA, Theodosiou T, Divanach P,
Trudgian DC, Thomas B, Papanikolaou N, Aivaliotis M, Acuto O et al.

2017. ProteoSign: an end-user online differential proteomics statistical
analysis platform. *Nucleic Acids Res* **45**: W300-W306.
Elias JE, Gygi SP. 2007. Target-decoy search strategy for increased confidence
in large-scale protein identifications by mass spectrometry. *Nat Methods*
4: 207-214.
Frith MC, Forrest AR, Nourbakhsh E, Pang KC, Kai C, Kawai J, Carninci P,
Hayashizaki Y, Bailey TL, Grimmond SM. 2006. The abundance of short
proteins in the mammalian proteome. *PLoS Genet* **2**: e52.
Hemm MR, Paul BJ, Miranda-Rios J, Zhang A, Soltanzad N, Storz G. 2010. Small
stress response proteins in Escherichia coli: proteins missed by classical
proteomic studies. *J Bacteriol* **192**: 46-58.
Hemm MR, Paul BJ, Schneider TD, Storz G, Rudd KE. 2008. Small membrane
proteins found by comparative genomics and ribosome binding site
models. *Mol Microbiol* **70**: 1487-1501.
Henkin TM, Yanofsky C. 2002. Regulation by transcription attenuation in
bacteria: how RNA provides instructions for transcription
termination/antitermination decisions. *Bioessays* **24**: 700-707.
Hinnebusch AG, Ivanov IP, Sonenberg N. 2016. Translational control by 5'-
untranslated regions of eukaryotic mRNAs. *Science* **352**: 1413-1416.
Hobbs EC, Fontaine F, Yin X, Storz G. 2011. An expanding universe of small
proteins. *Curr Opin Microbiol* **14**: 167-173.
Loi VV, Rossius M, Antelmann H. 2015. Redox regulation by reversible protein S-
thiolation in bacteria. *Front Microbiol* **6**: 187.
Madeira F, Park YM, Lee J, Buso N, Gur T, Madhusoodanan N, Basutkar P, Tivey
ARN, Potter SC, Finn RD et al. 2019. The EMBL-EBI search and sequence
analysis tools APIs in 2019. *Nucleic Acids Res* **47**: W636-W641.
Martini MC, Zhou Y, Sun H, Shell SS. 2019. Defining the Transcriptional and
Post-transcriptional Landscapes of Mycobacterium smegmatis in Aerobic
Growth and Hypoxia. *Front Microbiol* **10**: 591.
Meydan S, Marks J, Klepacki D, Sharma V, Baranov PV, Firth AE, Margus T, Kefi
A, Vazquez-Laslop N, Mankin AS. 2019. Retapamulin-Assisted Ribosome
Profiling Reveals the Alternative Bacterial Proteome. *Mol Cell* **74**: 481-
493 e486.
Miranda-CasoLuengo AA, Staunton PM, Dinan AM, Lohan AJ, Loftus BJ. 2016.
Functional characterization of the Mycobacterium abscessus genome
coupled with condition specific transcriptomics reveals conserved
molecular strategies for host adaptation and persistence. *BMC Genomics*
17: 553.
Nakamoto T. 2009. Evolution and the universality of the mechanism of
initiation of protein synthesis. *Gene* **432**: 1-6.
Oppenheim DS, Yanofsky C. 1980. Translational coupling during expression of
the tryptophan operon of Escherichia coli. *Genetics* **95**: 785-795.
Rappsilber J, Ishihama Y, Mann M. 2003. Stop and go extraction tips for matrix-
assisted laser desorption/ionization, nanoelectrospray, and LC/MS
sample pretreatment in proteomics. *Anal Chem* **75**: 663-670.

Sberro H, Fremin BJ, Zlitni S, Edfors F, Greenfield N, Snyder MP, Pavlopoulos
GA, Kyrpides NC, Bhatt AS. 2019. Large-Scale Analyses of Human
Microbiomes Reveal Thousands of Small, Novel Genes. *Cell* **178**: 1245-
1259 e1214.
Shell SS, Wang J, Lapierre P, Mir M, Chase MR, Pyle MM, Gawande R, Ahmad R,
Sarracino DA, Ioerger TR et al. 2015. Leaderless Transcripts and Small
Proteins Are Common Features of the Mycobacterial Translational
Landscape. *PLoS Genet* **11**: e1005641.
Swaminathan S, Ellis HM, Waters LS, Yu D, Lee EC, Court DL, Sharan SK. 2001.
Rapid engineering of bacterial artificial chromosomes using
oligonucleotides. *Genesis* **29**: 14-21.
Turnbough CL, Jr. 2019. Regulation of Bacterial Gene Expression by
Transcription Attenuation. *Microbiol Mol Biol Rev* **83**.
van Kessel JC, Hatfull GF. 2008. Efficient point mutagenesis in mycobacteria
using single-stranded DNA recombineering: characterization of
antimycobacterial drug targets. *Mol Microbiol* **67**: 1094-1107.
Weaver J, Mohammad F, Buskirk AR, Storz G. 2019. Identifying Small Proteins
by Ribosome Profiling with Stalled Initiation Complexes. *mBio* **10**.
Xu X, Vilcheze C, Av-Gay Y, Gomez-Velasco A, Jacobs WR, Jr. 2011. Precise null
deletion mutations of the mycothiol synthesis genes reveal their role in
isoniazid and ethionamide resistance in Mycobacterium smegmatis.
Antimicrob Agents Chemother **55**: 3133-3139.
Yanofsky C. 1981. Attenuation in the control of expression of bacterial operons.
Nature **289**: 751-758.
Zheng X, Hu GQ, She ZS, Zhu H. 2011. Leaderless genes in bacteria: clue to the
evolution of translation initiation mechanisms in prokaryotes. *BMC*
Genomics **12**: 361.
Zougman A, Selby PJ, Banks RE. 2014. Suspension trapping (STrap) sample
preparation method for bottom-up proteomics analysis. *Proteomics* **14**:
1006-1000.
654
655

Figure 1. Polycysteine-encoding LL-sORF attenuates expression of operonic genes.

A. Locus schematic of *Ms5788A* upstream of *Ms5788-5790*. The conceptual amino acid

sequence of the wild-type Ms5788A shows 9 cysteine codons (bold C), with eight

consecutive codons at the 3' end. **B.** Ribosome occupancy of *Ms5788A* relieves

attenuation of luciferase reporter gene expression in the absence (-) of cysteine. The same

wild-type (i) reporter data are shown in Figs 2-3 as a comparative reference for the

mutant derivatives. Non-start (ii), nonsense (iii), and frameshift (iv) mutations decrease

basal or attenuated expression (or both) of the luciferase gene. Lower case letters below

665   clones indicated nucleotide substitutions and insertion/deletions (+/-) used to create

666   mutants. Asterisks indicate significance $p <.01$ for cysteine supplementation, by two-

667   tailed t-test in Figs 1-3.

668
669

670
671
Figure 2. Predicted duplex mRNA structures regulate Shine-Dalgarno availability for

expression of *Ms5788* luciferase reporter gene. A stable mRNA structure model

(RNAWebSuite/RNAfold) shows extensive base pairing forming a stem between the

polycysteine coding region of the LL-sORF and the peri-Shine-Dalgarno region. The

color of each nucleotide indicates the confidence in the predicted structure (as indicated

in the heatmap gradient key). Clustered point mutations were introduced into the

Ms5788A LL-sORF or the predicted pairing nucleotides to disrupt base pairing but, when

combined, should restore the modeled stable mRNA structure when combined. Two

series of mutants were generated (red, ii – iv and purple, v – vii). The effect of structure-

destabilizing mutations and re-stabilizing combination were tested for their effect on the

luciferase reporter fused to *Ms5788.*

Figure 3. Attenuation regulates translation initiation of Ms5788. Nluc translated via an

independent Shine-Dalgarno and initiation codon (M in white box) gages mRNA

extension beyond the structured leader (ii). Placement of the luciferase reporter at the

initiation codon of Ms5789 (iii) indicates that translation attenuation in Ms5788 has polar

effects, propagating the effects of *Ms5788A*.

Figure 4. Cysteine attenuation is observed in the native context. a. Label-Free

Quantitative proteomics (LFQ) was used to identify and quantitate changes in the

abundance of proteins from whole-cell extracts of wild-type *M. smegmatis* subjected to

trypsin digestion and nanoUHPLC-MS/MS. *M. smegmatis* was cultured in minimal

media (X-axis) or supplemented with cysteine (Y-axis). Units are normalized LFQ peak

area (protein) in arbitrary units. The baselines for both X and Y-axes were artificially set

at 5×10^4 counts to allow depiction of proteins not expressed in one of the two conditions.

Peptides from Ms5789 and Ms5790 (yellow diamond 1 and 2, respectively) increased in

abundance (shift right along the X-axis) under cysteine limitation. Peptides of annotated

proteins from loci similarly encoded on polycysteine-encoding LL-sORF mRNAs are

indicated by black filled circles (Fig S3; 3 = Ms0113; 4 = Ms0114; 5 = Ms0934; 6 =

Ms4527; 7 = Ms4533; 8 = Ms5279; 9 = Ms5280). b. A single nucleotide deletion caused

a frameshift mutation (OoF) in *Ms5788A* and changes the CCCCCCC* to

VAVVVVAVersral*. This OoF LL-sORF mutant is insensitive to cysteine and does not

release Ms5789 and Ms5790 from attenuation. c. Deleting *Ms5788A* elevates expression

of Ms5789 and Ms5790 and is insensitive to cysteine, indicating a completely

unattenuated state.

710
711

Figure 5. Phylogenetic distribution of polycysteine LL-sORFs in mycobacteria. A robust

reference phylogenetic tree was constructed from complete genome sequences of 41

diverse *Mycobacterium* spp. All nodes had bootstrap values of 100, except where

indicated. Species of the slow grower clade appear in blue text, and fast growers in

magenta. The presence (black square) or absence (white square) of each LL-sORF forms

a binary barcode for each species. Barcode key and fixed-length sequence logo for each

LL-sORF is shown.

720
721

gene #	gene name	encoded protein predicted function	HMM Pfam	uniprot
Ms0113	tauC	taurine transport system permease protein TauC; sulfur starvation	PF00528	A0QNP1
Ms0114		hnD/SsuA/transferrin family substrate-binding protein	PF04069	A0QNP2
Ms0116		taurine import ATP-binding protein TauB	PF00005	A0QNP3
Ms0932		ROK family protein	PF00480	A0QQZ7
Ms0933	mshA	MshA glycosyltransferase	PF00534	A0QQZ8
Ms0934		YbjN domain-containing protein		A0QQZ9
Ms4527		ferredoxin sulfite reductase	PF01077	A0R0W1
Ms4528	cysH	phosphoadenosine phosphosulfate reductase	PF01507	A0R0W2
Ms4529		secreted protein, sirohydrochlorin chelatase	PF01903	A0R0W3
Ms4530	cysA	sulfate ABC transporter, ATP-binding protein	PF00005	A0R0W4
Ms4531	cysW	sulfate ABC transporter, permease protein	PF00528	A0R0W5
Ms4532	cysT	sulfate ABC transporter, permease protein CysT	PF00528	A0R0W6
Ms4533		sulfate ABC transporter substrate-binding protein	TIGR00971	A0R0W7
Ms4536		TQXA domain-containing protein; surface-expressed		A0R0X0
Ms5280		cysteine dioxygenase type I superfamily protein	PF05995	A0R2Y8
Ms5279		sulfurtransferase		A0R2Y7
Ms5788		DUF4395 domain-containing protein		A0R4C8
Ms5789	cysA2	putative thiosulfate sulfurtransferase	PF00581	A0R4C9
Ms5790		DUF1416 domain-containing protein, SseC protein	PF07210	A0R4D0
Rv2334	cysK1	O-acetylserine sulfhydrylase; Interacts with CysE		P9WP55
Rv2335	cysE	serine acetyltransferase; acetylates serine to make cysteine		P95231

722
723
724   Table 1. Annotated genes of the cysteine translation attenuation regulon. Gene identifiers

725   are listed for *M. smegmatis* (Ms) or *M. tuberculosis* (Rv). The predicted functions and

726   features of the encoded proteins are shown, as annotated in Pfam or uniport record.

727
728

 EXTENDED DATA FILES

a

b

Extended Data Fig 1. a. Schematic and expression profile of the *Ms5788A* locus. RNA-seq and Ribo-seq profiles show robust occupancy of the positive strand (maximum displayed read depth = 500), corresponding to the LL-sORF (shaded yellow). The end of the transcriptionally independent *thiX* gene is shown, and *cysA2* (Ms5789) represents operon genes downstream. Reads from a single replicate are shown for space efficiency. Note that *Ms5787* (gold line shown as an annotated gene reading from right to left) is likely a misannotation, as we detect no transcriptional or translational activity for this anti-sense gene. b. Schematic and expression profile for the *M. tuberculosis* orthologous locus, *Rv0815A*.

Extended Data Fig 2. RNA-folding model of the polycysteine *Ms5788A* mutant. The *Ms5788A* ORF mutations (indicated by filled triangles) retain polycysteine coding, but disrupt the stability and folding of the wild-type mRNA as indicated in the modeled structures. Mutations of the complementary nucleotides (indicated by open triangles) near the *Ms5788* Shine-Dalgarno sequence, are predicted to form G-U wobble base pairs; the resulting overall structure and stability are comparable to that of the wild-type mRNA. The comparable wild-type predicted folding of the complementary stem mutant is consistent with the comparable wild-type expression of the reporter (Fig 2 vi). Combining each mutated stem restores the predicted mRNA 2° structure and cysteine attenuation (Fig 2 vii). Color scheme is as for Fig 2.

732
733

3a

3b

Extended Data Fig 3. Additional loci that feature expressed polycysteine LL-sORFs upstream of annotated genes. JBrowse images of each LL-sORF locus show the gene context and sequence and transcriptomic profiles. Yellow arrow boxes indicate the LL-sORF in each locus, and the gene encoding the cysteine-responsive protein detected by mass spectrometry is indicated by the black circle with white number as in the mass spectrometry scatter plot (Fig 5a). Panels are shown for (a) *Ms0113A/0013/Ms0114*, (b) *Ms0932A/0932/0933/0934.* Note that genes (B) *Ms0931* and (C) *Ms4528* are likely misannotated based on our transcriptional and translational data. Pipeline prediction algorithms annotated none of these LL-sORFs. Screen shots captured from http://www.wadsworth.org/research/scientific-resources/interactive-genomics.

3c

3d

(C) *Ms4527A/4527/4528/4529*, (D) *Ms4533A/4533/4532/4531/4530*

3e

3f

(e) *Ms4536A/4536*, (f) *Ms5280/5279*

3g

(g) *Rv2334A*

ANNOTATED

Amino Acid	x1	x2	x3	x4	x5	x6	x7	x8	x9	x10	x11	x12	x13	x14	x15	x16	x17	x18	x19	freq	exp x2	obs >1x en	>1x inst	total AA
A, alanine	197695	29314	4605	729	186	33	1	1	1											0.118932	27659.338	1.2606954	34870	232565
C, cysteine	16898	197	1	1																0.008743	149.48337	1.3312517	199	17097
D, asparate	115764	7902	627	26	4		6													0.063581	7904.9284	1.0835013	8565	124329
E, glutatmate	103393	5090	167	6																0.055566	6037.5465	0.8717117	5263	108656
F, phenylalanine	61161	2052	73	1		1														0.032365	2048.3106	1.0384167	2127	63288
G, glycine	155845	13140	1222	123	25	5	2			1			1							0.087123	14842.557	0.9782007	14519	170364
H, histidine	45231	1405	54	2	1															0.023878	1114.9531	1.3112659	1462	46693
I, isoleucine	85428	3059	82	1																0.045294	4011.6806	0.7832129	3142	88570
K, lysine	43100	1377	37	3																0.022766	1013.4559	1.3981862	1417	44517
L, leucine	168666	16936	1363	158	9	1	1													0.095699	17908.469	1.0312439	18468	187134
M, methioine	41595	881	16	3																0.021732	923.48263	0.9745717	900	42495
N, asparagine	44477	1187	33																	0.023369	1067.8947	1.1424347	1220	45697
P, proline	107885	5629	544	103	33	19	11	2	3	1			1							0.058417	6672.9984	0.9509968	6346	114231
Q, glutamine	56801	2368	115	5	3		1													0.030322	1797.8767	1.3860795	2492	59293
R, arginine	127562	11444	1220	138	15	6	3		1											0.071794	10079.041	1.2726409	12827	140389
S, serine	99161	5508	365	38	8	1	3		1											0.05374	5647.2176	1.0490122	5924	105085
T, threonine	114519	7025	569	42	10	4	4	1	2		1								1	0.062481	7633.7703	1.003305	7659	122178
V, valine	151768	14766	1590	156	10	1														0.086062	14483.544	1.1408119	16523	168291
W, tryptophan	29995	632	10																	0.015667	480.00475	1.3374868	642	30637
Y, tyrosine	42884	1038	19	1																0.022472	987.44451	1.0714526	1058	43942
TOTALS	1809828	130950	12712	1536	304	71	32	4	8	2	1	0	1	1	0	0	0	0	1	1	118632.23	21.84775	145623	1955451

LL-sORFs

Amino Acid	x1	x2	x3	x4	x5	x6	x7	x8	x9	x10	x11	x12	x13	x14	x15	x16	x17	x18	x19	freq	exp x2	obs >1x en	>1x inst	total AA
A, alanine	437	40	3																	0.09548	38.070312	1.129489	43	526
C, cysteine	142	14	2	2				1												0.034852	5.0724457	3.7457277	19	192
D, asparate	204	15																		0.042476	7.5343651	1.9908778	15	234
E, glutatmate	212	11	1																	0.043021	7.7287923	1.5526358	12	237
F, phenylalanine	103	4	1																	0.020693	1.7882352	2.7960527	5	114
G, glycine	407	31																		0.085133	30.26639	1.0242384	31	469
H, histidine	159	2																		0.029588	3.655865	0.5470661	2	163
I, isoleucine	158	3																		0.029769	3.7008599	0.8106224	3	164
K, lysine	88	2	1																	0.017245	1.24183	2.4157896	3	95
L, leucine	285	27	3																	0.063169	16.663777	1.8003122	30	348
M, methioine	90	4																		0.017789	1.3214998	3.0268639	4	98
N, asparagine	101																			0.018334	1.4036463	0	0	101
P, proline	331	17	3																	0.067889	19.246783	1.0391347	20	374
Q, glutamine	149	6																		0.029225	3.5667009	1.6822268	6	161
R, arginine	571	95	12	4																0.147577	90.948604	1.2204695	111	813
S, serine	419	35	5	3																0.093665	36.636531	1.1736919	43	516
T, threonine	318	25	2																	0.067889	19.246783	1.4028318	27	374
V, valine	283	27	4																	0.063351	16.759683	1.849677	31	349
W, tryptophan	114	2																		0.021419	1.915927	1.0438811	2	118
Y, tyrosine	63																			0.011436	0.54613	0	0	63
TOTALS	4634	360	37	9	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	176.92714	19.825789	407	5509

Extended Data Table 1. Amino acid content and clustering in proteins encoded by annotated and LL-sORF genes. Single or consecutive instances are totaled. Freq = specific AA/total AA, exp x2 = (Freq)^2*total AA, obs >1x en = obs/exp, >1x inst = total of polyAA occurrences.

LL-sORF	0113A	0932A	4527A	4533A	4536A	5280A	5788A	Rv2334A
Mycobacterium_abscessus	Cys x 4		Cys x 1X2	Cys x 2		Cys x 2	Cys x 6	
Mycobacterium_africanum		Cys x 3	Cys x 1X1X1	Cys x 2		Cys x 2	Cys x 5X1	Cys x 4
Mycobacterium_avium_104			Cys x 1X1X1	Cys x 2	Cys x 6		Cys x 7	
Mycobacterium_bovis		Cys x 3	Cys x 1X1X1	Cys x 2		Cys x 2	Cys x 5 x 1	Cys x 4
Mycobacterium_canettii		Cys x 3	Cys x 1X1X1	Cys x 2		Cys x 2		Cys x 4
Mycobacterium_caprae		Cys x 3	Cys x 1X1X1	Cys x 2		Cys x 2	Cys x 5X1	Cys x 4
Mycobacterium_chelonae	Cys x 3		Cys x 1X2	Cys x 2		Cys x 2	Cys x 6	
Mycobacterium_chimaera			Cys x 1X1X1	Cys x 2			Cys x 7	Cys x 1 x 2
Mycobacterium_chubuense			Cys x 1X3	Cys x 4			Cys x 5X1	
Mycobacterium_colombiense			Cys x 1X1X1	Cys x 2	Cys x 7		Cys x 7	
Mycobacterium_dioxanotrophicus		Cys x 3	Cys x 1X3	Cys x 2	Cys x 4		Cys x 7	
Mycobacterium_fortuitum		Cys x 4	Cys x 1X3	Cys x 2		Cys x 2	Cys x 7	
Mycobacterium_gilvum			Cys x 1X3	Cys x 3			Cys x 7	
Mycobacterium_goodii	Cys x 4	Cys x 4	Cys x 1X3	Cys x 2		Cys x 2	Cys x 7	
Mycobacterium_haemophilum			Cys x 1X3		Cys x 3 x 2		Cys x 5X1	
Mycobacterium_hassiacum			Cys x 1X3	Cys x 3			Cys x 6	
Mycobacterium_immunogenum	Cys x 3		Cys x 1X2			Cys x 2	Cys x 7	
Mycobacterium_intracellulare			Cys x 1X1X1	Cys x 2			Cys x 7	Cys x 1 x 2
Mycobacterium_kansasii		Cys x 3		Cys x 2	Cys x 5			
Mycobacterium_leprae					Cys x 1 x 2		Cys x 1 x 1 x 1	
Mycobacterium_lepraemurium			Cys x 1X3		Cys x 6		Cys x 5	
Mycobacterium_liflandii		Cys x 3	Cys x 1X3				Cys x 5X1	Cys x 5
Mycobacterium_litorale			Cys x 1X3	Cys x 2			Cys x 6	
Mycobacterium_marinum_M		Cys x 3	Cys x 1X3				Cys x 5 x 1	Cys x 5
Mycobacterium_marseillense			Cys x 1X1X1	Cys x 2			Cys x 7	Cys x 3
Mycobacterium_microti		Cys x 3	Cys x 1X1X1	Cys x 2		Cys x 2	Cys x 5X1	Cys x 4
Mycobacterium_neoaurum	Cys x 4		Cys x 1X3	Cys x 3		Cys x 2	Cys x 5	
Mycobacterium_paragordonae		Cys x 3	Cys x 1X3	Cys x 2	Cys x 5 x 1			
Mycobacterium_phlei	Cys x 3	Cys x 4	Cys x 1X3				Cys x 6	
Mycobacterium_pseudoshottsii		Cys x 3	Cys x 1X3				Cys x 5X1	Cys x 5
Mycobacterium_rutilum		Cys x 3	Cys x 1X3	Cys x 4			Cys x 6	
Mycobacterium_shigaense			Cys x 1X1X1	Cys x 2	Cys x 2 x 2	Cys x 2	Cys x 2X3	
Mycobacterium_simiae			Cys x 1X1X1	Cys x 2	Cys x 2 x 2		Cys x 7 x 1	
Mycobacterium_sinense			Cys x 1X1X1	Cys x 2		Cys x 3	Cys x 4X1	
Mycobacterium_smegmatis	Cys x 4	Cys x 4	Cys x 1X3	Cys x 2	Cys x 3	Cys x 2	Cys x 8	
Mycobacterium_stephanolepidis	Cys x 3		Cys x 1X2	Cys x 2		Cys x 2	Cys x 6	
Mycobacterium_thermoresistibile		Cys x 3	Cys x 1X3					
Mycobacterium_tuberculosis		Cys x 3	Cys x 1X1X1	Cys x 2		Cys x 2	Cys x 5X1	Cys x 4
Mycobacterium_ulcerans		Cys x 3	Cys x 1X3					Cys x 5
Mycobacterium_vaccae			Cys x 1X3	Cys x 3			Cys x 6	
Mycobacterium_vanbaalenii		Cys x 4	Cys x 1X3	Cys x 3			Cys x 6	

747
748
749 Extended Data Table 2. Polycysteine LL-sORF gene conservation in mycobacteria. Each
750 column designates the query LL-sORF. Open cells did not return an orthologous gene.
751 Cys x describes the array of cysteines in the polycysteine cluster; arrays that are
752 interrupted by non-cysteine codons represent that codon by X.
753