1 **Comparative genomic analyses and a novel linkage map for cisco (*Coregonus artedi*)**
2 **provide insights into chromosomal evolution and rediploidization across salmonids**
3

4 Danielle M. Blumstein[1], Matthew A. Campbell[2], Matthew C. Hale[3], Ben J. G. Sutherland[4],
5 Garrett J. McKinney[5], Wendylee Stott[6], Wesley A. Larson[7, 1]
6
[1] College of Natural Resources, University of Wisconsin-Stevens Point
[2] University of Alaska Museum of the North, University of Alaska Fairbanks, Fairbanks, AK,
99775, USA
[3] Department of Biology, Texas Christian University, Fort Worth, TX, 76122
[4] Pacific Biological Station, Fisheries and Oceans Canada, Nanaimo, V9T 6N7, British
Columbia, Canada
[5] National Marine Fisheries Service, Northwest Fisheries Science Center
[6] U.S. Geological Survey, Great Lakes Science Center, Ann Arbor, MI
[7] U.S. Geological Survey, Wisconsin Cooperative Fishery Research Unit, College of Natural
Resources, University of Wisconsin-Stevens Point
17
18

22 **Corresponding author:** Wesley Larson (e-mail: wlarson@uwsp.edu, phone: 715-346-3150,
23 office mailing address: U.S. Geological Survey, Wisconsin Cooperative Fishery Research Unit,
24 College of Natural Resources, University of Wisconsin)
25
26
27

Abstract

Whole-genome duplication (WGD) is hypothesized to be an important evolutionary mechanism that can facilitate adaptation and speciation. Genomes that exist in states of both diploidy and residual tetraploidy are of particular interest, as mechanisms that maintain the ploidy mosaic after WGD may provide important insights into evolutionary processes. The Salmonidae family exhibit residual tetraploidy within chromosomes, approximately 25% of the genome. This combined with the evolution diversity after an ancestral autotetraploidization event, makes this group a useful evolutionary system. In this study, we generate a novel linkage map for cisco (*Coregonus artedi*) and conduct comparative genomic analyses to refine our understanding of chromosomal fusion/fission history across salmonids. To facilitate this comparative approach, we propose the adoption of the naming strategy of protokaryotype identifiers (PKs), a series of identifiers that associate duplicated chromosomes to their putative ancestral state. The female linkage map for cisco contains 20,450 loci, 3,383 of which are likely contained within residually tetraploid regions of the genome. Comparative genomic analysis revealed that patterns of residual tetrasomy are generally conserved across species although interspecific variation persists in terms of relative extent of rediploidization. To determine the broad-scale retention of residual tetrasomy across the salmonids, we analyze sequence similarity of the currently available reference genomes using known state training sets and find evidence of residual tetrasomy in seven of the eight chromosomes that have been previously hypothesized to show this pattern. This interspecific variation that exists may have important implications for understanding salmonid evolutionary histories and informing future conservation efforts.

Keywords: coregonines; comparative genomics; linkage mapping; residual tetrasomy; Salmonidae; whole genome duplication

Introduction

The evolutionary significance of whole-genome duplications (WGDs) has been intensively debated for decades (e.g; Ohno 1970; Taylor et al. 2003; Santini et al. 2009; Wood et al. 2009; Zhan et al. 2014; Mayrose et al. 2015; Van de Peer et al. 2017). Multiple studies have hypothesized that WGD is an important evolutionary mechanism that can facilitate adaptation on short- and long-term evolutionary timescales (Ohta 1989, Selmecki et al. 2015; Van de Peer et al. 2017). For example, genes found in polyploid regions are able to gain new function (i.e., neofunctionalization) without the consequences of deleterious mutations affecting the main function of the second gene copy. This may facilitate adaptive molecular divergence and the evolution of new phenotypes (Wittbrodt et al 1998; Wendel 2000; Rastogi and Liberles 2005). However, other studies have hypothesized that WGD presents significant challenges for meiosis and mitosis (Hollister 2015) and may not have as much of an effect on evolution as originally considered (Mayrose et al. 2011; Arrigo and Barker 2012; Vanneste et al. 2014; Clarke et al. 2016). A consensus is therefore yet to be reached on the evolutionary impact of WGD relative to other evolutionary forces.

Conducting genetic studies on organisms with relatively recent WGD has been challenging due to the inability to differentiate alleles and sequences from the same chromosome (homologs) from those on the duplicated chromosome (homeologs) (Limborg et al. 2016b). Fortunately, approaches leveraging gamete manipulation, high sequencing coverage, and long read sequencing have improved our ability to characterize duplicated regions. Linkage mapping with haploids and doubled haploids has facilitated analysis of duplicated regions in salmonids (Brieuc et al. 2014; Kodama et al. 2014; Lien et al. 2016; Waples et al. 2016). Further, long-read sequencing technologies have made it possible to assemble complex genomes with convoluted duplication histories (Kyriakidou et al. 2018). These technological advances have revolutionized our ability to understand genomic architecture in species that have adaptively radiated into dozens of species following an ancestral WGD in lineages, such as salmonids (Lien et al. 2016; Robertson et al. 2017; Campbell et al. 2019) and many plant species (Alix et al. 2017).

Salmonids are derived from an ancestral species that underwent a WGD ~100 million years ago (Ss4R, Allendorf and Thorgaard 1984; Berthelot et al. 2014; Macqueen and Johnston 2014; Lien et al. 2016) and have since diversified into a broad array of ecologically and genetically distinct taxa. The Salmonidae family is comprised of three subfamilies: Salmoninae (salmon, trout, and char), Thymallinae (graylings), and Coregoninae (whitefish and ciscoes) (Norden 1961), and diversification of these subfamilies post-dates the Ss4R, occurring 40-50 million years ago (Campbell et al. 2013; Macqueen and Johnston 2014). Phylogenetic analysis has revealed that the majority of the salmonid genome returned to a diploid inheritance state prior to the basal split of each of the three subfamilies (Robertson et al. 2017). However, the rediploidization process is still incomplete and approximately 20-25% of each salmonid genome still shows signals of tetrasomic inheritance (i.e., residual tetrasomy, or the recombination between homeologs that results in the exchange of alleles between homeologous chromosomes (Allendorf et al. 2015; Lien et al. 2016; Robertson et al. 2017)).

Early evidence for residual tetrasomy in the salmonids was identified using allozyme
studies in experimental crosses (Allendorf and Danzmann 1997) and, more recently, by linkage
maps, sequenced genomes, and sequence capture (Kodama et al. 2014; McKinney et al. 2017;
Robertson et al. 2017; Christensen et al. 2018b; Pearse et al. 2019). High-density linkage maps
that include both duplicated and non-duplicated markers have revealed that eight pairs of
homeologous chromosomes repeatedly display evidence of residual tetraploidy despite
independent fusion and fission events (Brieuc et al. 2014; Kodama et al. 2014; Sutherland et al.
2016). These chromosomes, often referred to as the "magic eight," have been observed in
linkage mapping studies of coho salmon *Oncorhynchus kisutch* (Kodama et al. 2014), Chinook
salmon *O. tshawytscha* (Brieuc et al. 2014; McKinney et al. 2016; McKinney et al. 2019), pink
salmon *O. gorbuscha* (Tarpey et al. 2017), chum salmon *O. keta* (Waples et al. 2016), and
sockeye salmon *O. nerka* (Larson et al. 2015). Mapping studies have also revealed that at least
one of the two homeologs exhibiting residual tetraploidy is within a chromosomal fusion,
suggesting their role in tetrasomy persistence (Brieuc et al. 2014; Kodama et al. 2014; Sutherland
et al. 2016). Analysis of sequenced genomes for Atlantic salmon (*Salmo salar*) and rainbow trout
(*O. mykiss*) also suggest evidence of residual tetraploidy, although in these genomic studies only
seven of these pairs were identified as displaying clear signals of conserved residual tetraploidy
(Lien et al. 2016; Campbell et al. 2019). Sequence capture analysis also identified seven pairs
displaying conserved signals of residual tetrasomy across species (Robertson et al. 2017). This
led to the definition of two types of homeologous regions: 1) ancestral ohnologue resolution
(AORe) regions with relatively low sequence similarity with ancestral homeologs that likely
rediploidized prior to species diversification; and 2) lineage-specific ohnologue resolution
(LORe) regions with high sequence similarity among homeologs, likely maintained by residual
tetraploidy (Robertson et al. 2017).
Over the past decade, an extensive proliferation of genomic resources has occurred
within salmonids. Currently, linkage maps that include duplicated regions are available for five
Oncorhynchus species, and genome assemblies are available for grayling *Thymallus thymallus*
(Savilammi et al. 2019), Atlantic salmon (Lien et al. 2016), Arctic char *Salvelinus alpinus*
(Christensen et al. 2018b), rainbow trout (Pearse et al. 2019), and Chinook salmon (Christensen
et al. 2018a). These resources permit investigation into the processes of rediploidization and
residual tetrasomy across the salmonid family. There is however an underrepresentation of other
lineages within the salmonid family, such as the Coregoninae subfamily (but see Gagnaire et al.
2013, and recently De-Kayne et al. 2019). There has not been any characterization in the
duplicated regions to date in Coregoninae. In the current study, we develop a high-density
linkage map for North American cisco (*Coregonus artedi*), the first haploid linkage map for the
Coregoninae subfamily, and analyze existing genomic resources for other salmonids with the
goal of investigating patterns of residual tetrasomy and chromosomal fusion and fission history
across the salmonid family, with particular focus on the coregonines. Our results suggest that (1)
interspecific variation in residual tetrasomy is greater than previously observed; (2) binary
definitions of chromosome ploidy status may not adequately capture variation within and among

species; (3) linkage maps and sequenced genomes identify slightly different patterns regarding
residual tetrasomy; and (4) a large number of fissions and fusions are specific to the base of the
Coregoninae subfamily with species-specific fusions within Coregoninae being rare. This study
uses new and existing resources to conduct the most comprehensive analysis of residual
tetrasomy across the salmonid phylogeny to date, revealing important patterns that likely
impacted the evolutionary history of this diverse and important taxonomic group.
138
Methods
Experimental crosses for linkage mapping
Genotypes from diploid and haploid families were used to build sex-specific linkage
maps. Diploid crosses were constructed from cisco collected in northern Lake Huron (45º
58'51.6" N -84 º19'40.8" W, USA) during spawning season (November 2015) by U. S. Fish and
Wildlife Service crews using standardized gill net assessment methods. Gametes for haploid
crosses were collected following the same methods, from the same location and month but in
2017. Gametes were extracted from mature fish and eggs were combined directly with sperm to
produce diploid crosses or with sperm that had been irradiated with 300,000 µJ/cm2 UV light for
two minutes to break down the DNA to produce haploid crosses. UV irradiation leaves the sperm
intact so that the egg can be activated but no paternal genetic material is contributed (i.e.,
gynogenesis, Chourrout 1982), resulting in haploid embryos developing with maternal genetic
material only. Crosses were made in the field and transported to the U. S. Geological Survey-
Great Lakes Science Center, Ann Arbor, Michigan (USA) for rearing. Tissue samples (fin clips)
were taken from adult parents, from offspring of the diploid crosses at age two, and from
haploids approximately 50 days post fertilization. All samples were preserved in a combination
of 95% ethanol and 5% EDTA and sent to the University of Wisconsin, Stevens Point Molecular
Conservation Genetics Lab for processing. Laboratory and field collections were conducted
under the auspices of the U.S. Fish and Wildlife Service and U.S. Geological Survey-Great
Lakes Science Center and all necessary animal care and use protocols were filed by these
agencies.
160
DNA extraction and RAD-sequencing library preparation
DNA was extracted using DNeasy 96 Blood and Tissue Kits (Qiagen, Valencia,
California) per the manufacturer's instructions. Quality and quantity of the extracted genomic
DNA was measured using the Quant-iT PicoGreen double-stranded DNA Assay (Life
Technologies) with a plate reader (BioTek). To confirm ploidy of haploid samples, parents and
offspring were genotyped using six polymorphic microsatellite loci known to occur at diploid
sites developed by Angers et al. (1995), Patton et al. (1997), and Rogers et al. (2004), and
individuals were classified as haploids if only a single allele was present at all loci. The
probability of not detecting a diploid if a diploid was present is ~1.09% based on microsatellite
heterozygosity in the parental population (unpublished data, Wendylee Stott).

Genomic DNA from diploids and confirmed haploids was prepared for RAD sequencing
using the SbfI restriction enzyme following the methods outlined in Ali et al. (2016) except
shearing restriction digested DNA was done with NEBNext® dsDNA Fragmentase® (New
England Biolabs, Inc) instead of sonication. DNA was then purified and indexed using
NEBNext® Ultra™ DNA Library Prep Kit for Illumina® per the manufacturer's instructions
(New England Biolabs, Inc). Libraries were sequenced on a HiSeq4000 with paired end 150bp
chemistry at the Michigan State Genomics Core Facility (East Lansing, MI).
178
SNP discovery and genotyping
Quality filtering, SNP identification, and genotyping was conducted using *STACKS* v.2.2
(Rochette and Catchen 2017). First, samples were demultiplexed with *process_radtags* with
flags -c, -q, -r, -t 140, --*bestrad*. Markers were discovered *de novo* and genotyped within
individuals with *ustacks* (flags = -m 3, -M 5, -H --*max_locus_stacks* 4, --*model_type* bounded, --
bound_high 0.05, --*disable-gapped*). A catalog of loci was created using a subset of the
individuals (diploid parents = 8, haploid parents = 5, wild fish = 38, total cisco = 51) with *cstacks*
(-n of 3, --*disable-gapped*). The 38 wild fish used in the catalog were collected from the same
geographic area using the same collection methods as listed above and were included to search
for a sex identification marker, which was unsuccessful (*data not shown*).
Putative loci within each individual fish were matched against the catalog with *sstacks*
(flag = --*disable-gapped*), *tsv2bam* was used with only the forward reads to orient the data by
SNP, and *gstacks* was used to combine genotypes across individuals. Only the forward reads
from the paired-end data were used in *gstacks* due to variable read depth in reverse reads and
thus less reliable genotyping. *gstacks* was also run separately with the forward and reverse reads
using *tsv2bam* to assemble longer contigs for sequence alignment and annotation. Final genotype
calls were output as VCF files with *populations* (flags = -r 0.75), with each family grouped as a
separate population in the *popmap* sample interpretation file. VCFtools (Danecek et al. 2011)
was used to identify and remove individuals from the study that were missing more than 30% of
data.
Maximum likelihood-based methods developed by Waples et al. (2016) were used to
identify loci that could be mapped in haploid crosses and to identify potentially duplicated loci.
Custom Python scripts available on GitHub (Python Software Foundation version 2.7) (see *Code
Availability*), were used to filter the haplotype VCF file output from the *populations* module to
identify loci that could be mapped in the diploid families. Loci missing more than 25% of data
and loci that were genotyped as heterozygous in both parents of diploid families (and therefore
could not be reliably mapped) were removed (as in Larson et al. 2015). Individual genotypes
were exported with the custom Python scripts as *LepMap3* input files. As a final step before
linkage mapping, genotypes from all seven families (haploid and diploid) were combined into a
single dataset to form the final female *LepMap3* input file and the four diploid families were
combined into a single dataset to form the final male *LepMap3* input file.
210

Linkage mapping
The program *LepMap3* (Rastas 2017) was used to construct linkage maps following the
methods of McKinney et al. (2016). Due to heterochiasmy (i.e., recombination rate differences
between males and females) that occurs in the Salmonidae family (Sakamoto et al. 2000), a
separate map was constructed for each sex. Loci were filtered and clustered into linkage groups
(LGs) based on recombination rates by calculating logarithm of the odds (LOD) scores between
all pairs of loci with the *SeparateChromosomes2* module. The LOD scores were chosen by
increasing the LOD value by one until the number of LGs stabilized and was similar to that
expected based on the haploid karyotype of cisco (N=40, Phillips et al. 1996). The final LOD
scores used to generate the map were LOD = 15 and 5 for the female and the male maps,
respectively. Loci were then ordered within LGs by utilizing paternal and maternal haplotypes as
inheritance vectors with the *OrderMarkers2* module. We used a minimum marker number per
LG of 100 for the female map and 40 for the male map. LGs were reordered and markers
removed until no large gaps remained (Rastas 2017).
225
Comparative analysis of syntenic regions of linkage maps via MAPCOMP
MAPCOMP can be used to compare syntenic relationships among markers between
linkage maps of any related species using a genome intermediate from another related species
(Sutherland et al. 2016). Here, MAPCOMP was used to compare the cisco map with other
Coregonus spp.; lake whitefish *C. clupeaformis* (Gagnaire et al. 2013), European whitefish *C.*
lavaretus "Albock" (De-Kayne and Feulner 2018), as well as other representative species from
other salmonid genera (i.e., Atlantic salmon (Lien et al. 2016), brook trout *S. fontinalis*
(Sutherland et al. 2016), and Chinook salmon (Brieuc et al. 2014) and a representative of the
outgroup to the salmonid WGD, northern pike *Esox lucius,* that does not have a duplicated
genome (Rondeau et al. 2014). All code to collect and prepare maps, and run the analysis are
available on GitHub (see *Code Availability*).
MAPCOMP pairs loci between the two compared linkage maps if they align at the same
locus or close to each other on the same contig or scaffold on the intermediate reference genome
(Sutherland et al. 2016). Due to the large phylogenetic distance covered in this analysis, two
reference genomes; grayling (Savilammi et al. 2019) and Atlantic salmon (Lien et al. 2016) were
used for comparisons. As salmonid chromosomal evolution is typified by Robertsonian fusions
(Phillips and Rab 2001), fused chromosome arms in cisco, lake whitefish, and European
whitefish were identified by aligning cisco markers to multiple salmonid genomes to identify
cases where one cisco LG corresponded with at least two chromosome arms in another species.
The fusion and fission phylogenetic history was plotted based on the most parsimonious
explanation of common fusions among *Coregonus* spp. basing the approximate timings of
fusions on shared fusions among species. Fusion history shared at the base of the Salmoninae
lineage was taken from earlier work (Sutherland et al. 2016).
249
Homeolog identification, similarity and inheritance mode

Homeologous chromosome arms can be identified in haploid crosses by mapping
multiple alleles of duplicated markers as described in Brieuc et al. (2014). Duplicated markers in
cisco were mapped using this method, and duplicated markers from previously constructed
linkage maps for coho salmon (Kodama et al. 2014), Chinook salmon (McKinney et al. 2019),
pink salmon (Tarpey et al. 2017), chum salmon (Waples et al. 2016), and sockeye salmon
(Larson et al. 2015) were obtained. Homeologs were then ranked based on the number of
markers supporting each known homeologous relationship and the number of duplicated markers
was used to determine patterns of inheritance (i.e., disomy vs. tetrasomy) of the homeologous
pair.
Homeology was assessed by comparing DNA sequence similarity between homeologous
arms from chromosome-level genome assemblies. Genomes included in this analysis were
grayling (GCA_004348285.1, Savilammi et al. 2019) Atlantic salmon (GCF_000233375.1, Lien
et al. 2016), Arctic char (GCF_002910315.2, Christensen et al. 2018b), rainbow trout
(GCA_002163495.1, Pearse et al. 2019), and Chinook salmon (GCF_002872995.1, Christensen
et al. 2018a). Homeologous arms were inferred either as identified in the original genome paper
(references above) or through MAPCOMP comparisons. Homeologous arms were then aligned to
determine sequence similarity using LASTZ v1.02 (Harris 2007) following methods outlined in
Lien et al. (2016). Options specified with LASTZ included --chain --gapped --gfextend --
identity=75.0..100. --ambiguous=iupac --exact=20. The analysis was restricted to alignments
with minimum percent match values of 75%, and a minimum length of 1,000 base pairs to
minimize the likelihood of spurious alignments that might be due to gene family duplication
rather than WGD. Overall similarity of a homeologous pair was represented by the median
percent similarity of all alignments, weighted by alignment length, and summarized with
boxplots for each homeologous pair in each species ordered based on descending median
percentage sequence similarity.
Each homeologous pair was classified into one of two categories, tetrasomic or disomic,
using a machine learning approach. Previous research indicates that salmonids are undergoing
rediploidization with tetrasomic homeologous pairs, disomic homeologous pairs, and
intermediate homeologous pairs of uncertain affinity (Campbell et al. 2019; Lien et al. 2016). To
objectively classify protokaryotypes into tetrasomic homeologous and disomic homeologous
pairs in each species, a training set was constructed containing the four highest and four lowest
sequence similarity homeolog pairs. A k – nearest neighbor classification (knn) approach was
then applied to the dataset using this training set. The k - nearest neighbor method uses votes
from the training set to classify, therefore it supplies an objective method to identify
intermediates and towards which class they are more similar based on the number and kind of
votes received from the training set. The k nearest-neighbors was determined by using a repeated
10-fold cross validation of 100 iterations with the trainControl function in the R package caret
(v6.0-84, Kuhn 2019). The k for each species with the largest number of neighbors exhibiting the
highest accuracy with the training set was selected. This k was then used to classify homeologous
pairs as disomic or tetrasomic, along with the predefined training set (knn function, Ripley and

Venables 2019). Overall, similarity between the two predicted categories from the *knn*
classification was tested with a Wilcoxon test (R Core Team 2018) incorporating percent
similarity from all alignments to determine whether the categories displayed significantly
different sequence similarity between homeologs (alpha = 0.01). All scripts used in this analysis
are available on GitHub (see *Code Availability*).
296
Results
RADseq, SNP discovery, and data filtering
RADseq data were obtained from 746 cisco across seven families, with an average of 4.1
M reads per individual (range: 1.1 – 30.8 M reads per individual). Individuals that were
genotyped at more than 30% of loci and loci that were genotyped in more than 75% of the total
individuals were retained, resulting in a dataset of 676 individuals (n = 333 diploid offspring;
330 haploid offspring; and 13 parents) and 49,998 unique loci (Supplementary file S1).
304
Linkage Mapping
A total of 22,459 unique loci were identified in the female (Figure 1) and male linkage
maps (Supplementary file S2). The female map included 20,450 loci distributed across 38 LGs
(Table 1), the male map included 6,340 loci distributed across 40 LGs, and 4,012 loci were
present on both maps (Supplementary file S3). A total of 40 chromosomes was expected from
karyotyping of coregonine fishes from the Great Lakes (Phillips et al. 1996), which matches the
number of LGs mapped in males. However, male LGs 39 and 40 contained relatively few
markers and may be fragments of other linkage groups rather than the two linkage groups that
were not mapped in females. Eight LGs (i.e., Cart01 – Cart08) were identified as metacentric
based on homology to two chromosome arms in other salmonids using MAPCOMP (*see below*). In
the female map, metacentric LGs were on average 85.44 cM (57.72 – 101.35 cM) and contained
739 loci (range: 685 - 867). Putative acrocentric LGs in the female map were on average 59.12
cM (50.97 - 64.53 cM) and contained 484 loci (range: 296 - 585). The total length of the female
map was 2,456.98 cM. The average lengths of metacentric LGs on the male map were 66.76 cM
(51.62 – 87.14 cM) and they contained 212 loci on average (range: 159 – 278). Putative
acrocentric LGs in the male map were on average 57.00 cM (40.54 – 83.66 cM) and contained
145 loci (range: 41 – 224). The total length of the male map was 2,357.97 cM. We identified
3,383 putatively duplicated loci on the female linkage map, and of these, 2,671 loci mapped to
one paralog and 709 loci mapped to both paralogs.
324
Comparative analysis of syntenic regions of linkage maps via MAPCOMP
The main focus of our comparative analysis was to define the homologous and
homeologous relationships among the linkage groups available for the coregonines, specifically
in cisco (current study), lake whitefish (Gagnaire et al. 2013), and European whitefish (De-
Kayne and Feulner 2018), and bring these species into the context of the broader chromosomal
correspondence by identifying the homologous chromosome arms in brook trout, Atlantic

salmon, and Chinook salmon, as well as the non-duplicated northern pike (Table 2, Supplementary file S4). To facilitate these comparisons, we have defined the term protokaryotype identifier (PK) for use in chromosome naming here. PKs correspond to hypothetical ancestral salmonid chromosomes pre-duplication inferred from observing a representative genome of the non-duplicated sister lineage of salmonids from the Esociformes (Ishiguro et al. 2003; Lopez et al. 2004) and are ordered as PK 01-25. Protokaryotypes correspond 1:1 with the northern pike genome but have two descendant homeologous regions within salmonid genomes. For example, PK 01 corresponds to northern pike chromosome 01 and was an ancestral pre-duplication salmonid chromosome which gave rise to homeologous Atlantic salmon chromosomes Ssa09c (PK 01.2) and Ssa20b (PK 01.1) and to homeologous rainbow trout Omy27 (PK 01.1) and Omy24 (PK 01.2) (Supplemental file S7). PKs in the previously hypothesized "magic eight" from linkage mapping studies are PKs 02, 06, 09, 11, 20, 22, 23, 25. PKs defined as LORes by Robertson et al. (2017) and those that displayed residual tetraploidy in previous genome-based studies (Lien et al. 2016; Campbell et al. 2019) are the same as these with the exception of PK 06, which is not identified as residually tetraploid. The use of this naming system corresponds to the identifiers used in Sutherland et al. (2016), maintaining the correspondence of PK .1 or .2.

Most PKs were identifiable in the MAPCOMP analysis, with some notable exceptions for each coregonine species. In cisco, chromosome arms PK 22.2, 25.1 and 25.2 were unidentified; two of these arms (PK 25.1 and 25.2) were also unidentified in the European whitefish linkage map (De-Kayne and Feulner 2018). Additionally, it was difficult to determine correspondences for PK 09 and 23. In European whitefish, five chromosome arms were unidentifiable (PK 09.1, 09.2, 23.2, 25.1 and 25.2), and there were homeology ambiguities for PK 02, as well as homology ambiguities for PK arm 05.2 (Table 2). In lake whitefish, five arms were unidentifiable (i.e., PK 09.2, 11.2, 20.2, 22.1, and 25.2), and there were homeology ambiguities for PK 02.2 and 06.2. In multiple species, arms where it was difficult to determine homologous relationships often had a high proportion of duplicated loci, presumably making distinguishing homologs and homeologs challenging. Nonetheless, most homologs and homeologs (42/50) were identified in all three coregonine species. This information was then leveraged to characterize the fusion/fission history within the Coregoninae lineage using the methods outlined in Sutherland et al. (2016).

The fusion/fission analysis indicated far fewer species-specific fusions than identified for salmonids in Sutherland et al. (2016), with most fusions occurring prior to the divergence of the coregonines (Figure 2). This difference in species-specific fusions may also be related to the general lower number of fusions in coregonines relative to *Salmo* and *Oncorhynchus* (Supplementary file S5); although, in the coregonines the majority of fusions were observed in more than one species, which was not observed in most other species previously characterized. Two strongly supported fusions were observed in all three coregonine species: fusions PK 05.1-06.1 and 10.2-24.1. PK 11.1-21.1 was fused in both cisco and European whitefish, which presumably underwent a fission in lake whitefish (Figure 2). However, evidence for the

correspondence for lake whitefish for PK 11.1 and 21.1 was not highly conclusive, and a more
recent analyses of lake whitefish by regenerating the linkage map suggest that PK 11.1-21.1 may
have not underwent a fission in this species and is indeed still fused (Claire Merot, *pers. comm.*).
Therefore, more work is needed to determine whether this fusion is conserved in all three
species. The full characterization of fissions will require the resolution of the ambiguous arms
that are considered as probable in the current analysis, and this may be further clarified in future
work.
In summary, five fusions were likely shared among all three species, and one was shared
between cisco and European whitefish, and possibly all three species (PK 11.1-21.1). Cisco had
two species-specific fusions (PK 10.1-20.1 and 22.1-20.2), bringing the total count of observed
fusions to eight. European whitefish had one species-specific fusion (PK 20.2-10.1), bringing the
total count of observed fusions to seven. Lake whitefish also had one species-specific fusion (PK
20.1-23.2), bringing the total count to six. Interestingly, the PK 09.2-17.1 fusion that was
originally proposed to be shared among all known salmonids (Sutherland et al. 2016), was found
not to be fused in any of the species here, suggesting either that this fusion occurred after the
divergence of *Coregonus* from the ancestor of the rest of the salmonids, or that a fission occurred
at the base of the coregonines (Figure 2). The observation that this fusion was not present in
grayling Varadharajan et al. 2018) also suggests the former.
389
Homeolog identification, similarity, and inheritance mode
A second major goal of this study was to compare homeologous relationships and modes
of inheritance within and among species. We identified 17 of the 25 homeologous chromosome
pairs (PK) in cisco using the markers that could be mapped to both homeologs in the linkage
map, and each homeologous pair shared between one and 86 duplicated loci (Fig. 3,
Supplementary file S6). Of the 17 homeolog pairs, six (PK 02, 06, 09, 11, 20, and 23) had many
loci (42-86) supporting homeology; these are six of the "magic eight" discussed above. The other
11 had few markers supporting homeology (i.e., 1-6) and are not members of the "magic eight".
The other two arms found in the magic eight were not identifiable in cisco, potentially because
they contained a large number of duplicated loci. All of the previously constructed linkage maps
for salmonids that included duplicated regions had a large number of markers supporting
homeology for the "magic eight" with the exception of pink salmon, where seven of the eight
PKs had high support (34 – 68 loci) but one pair (PK25) displayed substantially lower support
(nine loci) (Tarpey et al. 2017) (Figure 3, Supplementary file S6).
To better understand the genetic similarity between homeologs and infer inheritance
mechanisms (i.e., residual tetrasomy or disomy), all 25 known homeologous relationships were
compared in reference genomes for grayling, Atlantic salmon, Arctic char, rainbow trout, and
Chinook salmon (Figures 3 and 4, Supplementary file S7). Using the machine learning algorithm
(see Methods), the optimal k – nearest neighbor for each species was identified as five. Those
five nearest neighbors from the training sets voted on the assignment of a particular PK to either
putatively tetrasomic or disomic classes (Figure 4), and the proportion of votes supporting each

assignment are reported in Supplemental Document S8. The highest observed vote proportion for
assignment to a class is 4 of 5 as a result of the limit on training set size to four of each class and
the five optimal k – nearest neighbors indicated for accuracy.
For Atlantic salmon and all *Oncorhynchus* spp. (i.e., rainbow trout and Chinook salmon),
the same eight PKs (i.e., PK 01, 02, 09, 11, 20, 22, 23, 25) were classified as tetrasomic using the
machine learning approach. This list of PKs includes all of those defined as LORes by Robertson
et al. (2017), and one additional (i.e., PK 01), but does not include PK 06, which is considered to
be part of the "magic eight" using linkage map evidence. Arctic char showed evidence for
residual tetrasomy in seven of these eight PKs, with the exception of PK 11 (see below for
details regarding this discrepancy due to other chromosome arms in this fusion). Grayling also
shared seven of the eight residually tetraploid homeolog pairs, with the exception of PK 01. Most
PKs received the highest possible vote proportions for their classifications (0.8), however, PK01
in Atlantic salmon and rainbow trout demonstrated a lower vote proportion (0.6) (Figure 4,
Supplemental Document S8), suggesting reduced support (i.e., lower sequence similarity) for
this homeologous pair being tetrasomic. Additionally, PK 19 in grayling, did not have the
highest vote proportion and was assigned as diploid but had the highest sequence similarity in
that class (Figure 4, Supplemental Document S8). Sequence similarity was significantly higher
for the tetrasomic PKs across all species ($P < 0.0001$).
Although the group of tetrasomic PKs was largely conserved across species, there was
substantial variation in the relative sequence similarity between these homeolog pairs (i.e., order
of highest to lowest similarity) among species. PK 01 consistently displayed the lowest sequence
similarity of all the PKs in all five species where it was classified as tetrasomic and did not
always receive the highest observed vote proportion (see above). However, there were a number
of other homeolog pairs that displayed highly variable sequence similarity rankings across
species (Figure 4). For example, PK 09 had the highest sequence similarity in the grayling
genome, the sixth highest in the rainbow trout genome, and the fourth or fifth highest in the other
genomes. This variation suggests that the frequency of tetravalent meiosis for each PK may
differ across species and that the process of diploidization has occurred in a species-specific
manner post WGD as suggested in the mechanisms proposed by Robertson et al (2017).
440
Discussion
The amount of genomic resources available for salmonids has increased drastically over
the last decade. However, many previous studies investigating genome evolution in salmonids
focus on one or a few species, with fewer studies considering broader subsets of available taxa to
understand patterns of genome evolution across the Salmonidae family (but see Sutherland et al.
2016; Robertson et al. 2017). Here, we utilize genomic resources along with a newly generated
high-density linkage map for cisco to compare patterns of homology, fusion/fission events,
homeology, and residual tetrasomy across species. The cisco linkage map incorporates
duplicated regions and contains 20,450 loci, making it denser than most salmonid RAD-based
haploid linkage (typically built from 3,000 to 5,000 loci). Higher marker density helped identify

orthologous relationships between coregonines and other salmonids as well as to identify
homeologous chromosomes in cisco. We also demonstrate the use of the protokaryotype ID
(PK), defined here but first used in Sutherland et al. (2016), for comparative analyses in
salmonids in order to unify and facilitate comparative approaches in salmonid linkage maps and
chromosome-level assemblies. Comparisons across Salmonidae revealed that patterns of
rediploidization are relatively similar across genera and loosely correspond with phylogeny.
However, we did identify substantial variation in sequence similarity between homeologs both
within species across homeolog pairs, and among species, suggesting that frequently used binary
classifications such as AORe/LORe and "magic eight" may be oversimplified.
460
Protokaryotype identifiers to facilitate comparative genomics in salmonids and other fishes
Comparative genomics within Salmonidae is important for the interpretation of the
effects of rediploidization after WGD on genome evolution (e.g., Berthelot et al. 2014; Kodama
et al. 2014; Lien et al. 2016). However, most studies use independent naming schemes for each
species, making it difficult to directly compare studies without complicated lookup tables or
alignments to confirm homology (e.g. Brieuc et al. 2014; Kodama et al. 2014). Recently
developed methods for connecting linkage maps through reference genomes (Sutherland et al.
2016) facilitated description of homologous relationships for all linkage group arms across
salmonids (with a few exceptions in coregonines). Additionally, Sutherland et al. (2016) and
Savilammi et al. (2019) have explored the utility of naming chromosomes based on homology to
northern pike, which as a representative of the non-duplicated sister lineage of salmonids, can be
used to represent the ancestral (pre-duplication) state of salmonid genomes. Each northern pike
chromosome is represented in salmonids by two homeologous (i.e., arose from the WGD event)
chromosome arms, which are designated with a .1 or .2 following the ancestral chromosome
number. This naming system, if applied broadly, will facilitate comparative genomics in
salmonids by creating a "Mueller element"-like system (reviewed in Schaeffer 2018), where
each chromosome arm has a universal identifier. However, there also remains value in species-
specific identifiers; for example, Cart03 is the third named linkage group in the Cisco linkage
map (Table 2). By comparison, Cart03 named via the PK system could be Cart03 (PK 08.2-09.1)
or Cart03$_{PK08.2-09.1}$ as Cart03 represents the fusion of two ancestral salmonid chromosome arms
08.2 and 09.1 (Figure 1, Table 2).
The PK system can facilitating comparisons of chromosomes containing genes for
adaptive potential in sockeye salmon (So13$_{PK18.2}$, TULP4, Larson et al. 2017), for run timing in
Chinook (Ots28$_{PK03.2}$, GREB1L, Prince et al. 2017), and for age-at-maturity in Atlantic salmon
(Ssa25$_{PK16.2}$, VGLL3, Barson et al. 2015). While there may be some sections of the PK that are
not always retained (e.g., some transposition of parts of chromosomes), as long as the majority of
the chromosome is preserved, then the PK system enables general comparisons. The PK system
proposed here will facilitate quick and accurate comparisons across taxa, adding significant
value to the myriad studies searching for adaptively important genes and regions in salmonids by
leveraging comparative approaches. A case study of the utility of this system is the comparison

of the chromosome containing the sex determining gene in salmonids (sdY), which is known to
move between chromosomes (see Table 1 in Sutherland et al. 2017). This comparison
demonstrated that some chromosome arms more frequently contain or are fused to the
chromosome that contains the sex determining gene than would be expected by chance or
explainable by phylogenetic conservation (i.e., PK 01.2 (AC04q), PK 03.1 (Cclu25, Co30,
So09), PK 19.1 (So09.5, AC04q.1), PK 15.1 (AC04q.2, BC35), Sutherland et al. 2017). Even
more intriguing is that the northern pike naming was based on the threespine stickleback
Gasterosteus aculeatus (Rondeau et al. 2014), and PK 19 is the sex determining chromosome in
three spine stickleback (Peichel et al. 2004). As observed above, this chromosome is often fused
with sex chromosomes in salmon (Sutherland et al. 2017). By comparison, using the naming
system, it is easy to observe that LG24 in northern pike (i.e., PK 24 in salmonids), recently
identified to hold the sex determining gene in northern pike (Pan et al. 2019), does not appear to
contain the sex-determining locus in any tested salmonid. Deriving this information would be
more difficult without the PK system and would require extensive cross-referencing.
The example of sex chromosomes from the PK system indicates a broad phylogenetic
utility of this nomenclature as it applies to threespine stickleback (a neoteleost) as well as
Esociformes and Salmoniformes. The protokaryotypes as defined here can represent the ancestral
karyotype of the five major euteleost lineages and be applicable of comparative genomic studies
among and within (1) Esociformes and Salmoniformes, (2) Stomiatii, (3) Argentiniformes, (4)
Galaxiiformes, and (5) Neoteleostei (Betancur-R et al. 2013). Exploration of the PK system as
defined here and its applicability across euteleosts should be conducted to determine the
suitability of the PK system for comparative genomics in the Euteleostei.
513
Homology and fission/fusion history in coregonines
Comparisons using linkage maps for three coregonine species (i.e., cisco, lake whitefish
(Gagnaire et al. 2013), and European whitefish (De-Kayne and Feulner 2018)), allowed us to
assess homology and variation in karyotypes across the genus. Within lake whitefish, aneuploidy
has been documented in diverged populations and historical contingency (Dion-Côté et al. 2015,
2017) and our results show ambiguity in homologous relationships remained for at least five
chromosome arms in all three coregonines; this degree of uncertainty was much higher than
documented in *Salmo*, *Oncorhynchus*, and *Salvelinus* by Sutherland et al. (2016), where there
were only two ambiguities across these groups. Coregonines appear to have a number of
relatively small acrocentric chromosomes (Phillips and Rab 2001), some of which contain a high
degree of duplicated loci, making constructing linkage maps more difficult than for other
salmonids (Gagnaire et al. 2013; De-Kayne and Feulner 2018). For example, PK 25 has never
been successfully mapped in coregonines, likely because it is small, submetacentric or
acrocentric, and contains many duplicates. In other salmonids, where PK 25 is part of larger
and/or metacentric chromosome, mapping becomes easier as there are many disomically
inherited markers on the chromosome. Interestingly, the fact that PK 25 likely residually
tetrasomic in cisco, even though it is likely an acrocentric or submetacentric chromosome,

suggests that metacentric chromosomes are not required for homeologous recombination. This
contradicts previous theory which suggests that homeologous recombination requires at least one
chromosome arm to be metacentric (Kodama et al. 2014). Additionally, PK 25.2 in grayling is a
submetacentric chromosome and also displays signals of residual tetrasomy (Savilammi et al.
2019), providing further evidence that a small secondary arm may be sufficient to facilitate
tetrasomic meiosis.
The fusion history in coregonines differs substantially from many other members of the
salmonid family. Members of the *Coregonus*, *Salvelinus*, and *Thymallus* genera possess the "A
karyotype," with a diploid chromosome number (2N) ~80 and many acrocentric chromosomes,
whereas *Oncorhynchus* and *Salmo*, possess the "B karyotype," with 2N ~60 and many
metacentric chromosomes (Phillips and Rab 2001). Given that these both come from an ancestral
type of n = 50 chromosome arms, species with the "A karyotype" have undergone fewer fusions
than lineages with the "B karyotype". Interestingly, it appears that "A karyotype" species also
generally contain a lower proportion of species-specific fusions compared to "B karyotype"
species, suggesting that the reduction in chromosome number and the higher frequency of
metacentric chromosomes characteristic of the "B karyotype", comes from species-specific
fusions. Sutherland et al. (2016) investigated fusion history within many species from the
Oncorhynchus genus and found that most species had many species-specific fusions (e.g., 17
species-specific fusions in pink salmon). However, Sutherland et al. (2016) only investigated one
species from the *Coregonus* and *Salvelinus* genera as this was all that was available at the time of
publication, and no species from *Thymallus*. Our current study is the first to investigate fusion
history across multiple coregonines and illustrates that most fusions are shared among species in
the *Coregonus* genus, contrasting the pattern observed in *Oncorhynchus* (Sutherland et al. 2016).
The functional effect of differing fusion histories is yet to be determined, and remains an
important question differentiating species within the *Coregonus*, *Salvelinus*, and *Thymallus*
genera from other salmonids. Further information from genome sequencing projects, for example
the European whitefish genome (De-Kayne et al. 2019) should facilitate important future studies
contrasting genomic processes and structure in species with differing fusion histories.
559
Patterns of homeology and residual tetrasomy across salmonids
Although patterns of residual tetrasomy were generally conserved, variation within and
among species was observed when examining results from linkage maps versus reference
genomes. Sequence similarity analyses using reference genomes suggested that all species
showed evidence for residual tetrasomic inheritance in seven homeologous pairs (PK 02, 09, 11,
20, 22, 23, and 25) with the exception of PK11 in Arctic char (see below). Using linkage maps,
these same seven homeologous pairs have been found to be tetrasomic in *Oncorhynchus*
(Kodama et al. 2014; McKinney et al. 2019), *Salvelinus* (Sutherland et al. 2016; Nugent et al.
2017), *Salmo* (Robertson et al. 2017), and likely *Coregonus* (results reported herein), strongly
suggesting that tetravalent meioses can and do form between these homeologs in all investigated
species to date. However, evidence for residual tetrasomy differed between linkage map and

genome methods for multiple PKs, most notably PK 06, which was classified as tetrasomic in
linkage mapping studies but not in genome analyses, and PK 01 which was classified as
tetrasomic in genome analyses but not linkage maps. It is likely that some of these differences
are the result of methodological limitations of the current approach and point to future analysis
approaches that may be able to improve upon the framework presented here. This is further
described below.
The observation that PK11 did not display high sequence similarity in Arctic char might
suggest a difference in diploidization rates in *Salvelinus* compared to other salmonids for this
homeologous pair, but it is more likely that methodological limitations prevented us from
detecting residual tetrasomy, as a linkage map study in Arctic char found a high number of
duplicated markers on this PK (Nugent et al. 2017). The percentage similarity analysis applied in
the present study uses complete chromosome alignments and requires post-filtering to remove
non-homeologous alignments. This method appears to be robust when chromosome arms are
well defined but, PK 11 in Arctic char appears to be composed of four chromosome arms that
have come together in a series of species-specific fusions (inferred from Christensen et al.
2018b). Since arm boundaries were not well defined, alignments in this PK produced a wide
interquartile range, suggesting that, while some regions of the PK are likely undergoing residual
tetrasomy, the alignments may have masked these regions by integrating over multiple
chromosome arms. This would be particularly problematic if the chromosomes being compared
both contained non-target chromosomes that were homeologous. To improve upon the method
applied here, better definition of the breaks between chromosome fusions could be applied and
this could prevent such ambiguities or noise in the sequence similarity calculated. We therefore
conclude that PK 11 is likely tetrasomic in Arctic char, but that we were unable to classify it as
such due to methodological limitations. The sequence similarity method applied here is generally
robust, but the fusion history of the species being analyzed needs to be considered to avoid
unexpected and erroneous similarity values. Ideally, only the section containing the ancestral
chromosome of interest would be being compared between the homeologs. This is an avenue of
method development that will be valuable for future work.
Contrastingly, the finding that PK 06 is not tetrasomic does not appear to be due to
methodological limitations of our genome analysis but may be due to differences in estimating
extent of residual tetraploidy between linkage mapping and genome assembly approaches.
Linkage mapping in *Oncorhynchus* and *Salvelinus* consistently finds support for tetrasomic
inheritance at PK 06 (Larson et al. 2017; Nugent et al. 2017), but the genome analysis conducted
here and that conducted for rainbow trout (Campbell et al. 2019) found that this PK displayed
intermediate sequence similarity consistent with disomic homeologs. One of the ways the two
approaches differ is the length of the sequence used during each analysis. The genome analysis
conducted here calculated similarity by using alignments of at least 1,000 bp, whereas linkage
maps compare alleles within ~100-150 bp RADtags. The short sequences analyzed by software
such as STACKs (Rochette and Catchen 2017; Rochette et al. 2019) make it possible to collapse
sequences into a single locus that can be mapped at both paralogs, even when sequence

divergence in a given region is relatively large. This therefore makes linkage maps a less
conservative characterization method for determining residual tetrasomy. In addition, many
genome assemblers applied to salmonid genomes (e.g. Chin et al. 2016; Koren et al. 2017; Ruan
and Li 2019) are not optimized for paralogous regions in polyploid genomes. This could be
especially problematic for genomes that combine both disomic and tetrasomic regions, such as in
salmonids. The end results are that duplicated regions may be detected as single copies as a
result of sequence collapse during the assembly process (Alkan et al. 2011; Varadharajan et al.
2018). If sequences do not collapse during assembly, contigs might be fragmented and
misassembled in the genome, making it difficult to differentiate between homologs and
homeologs (Kyriakidou et al. 2018). This could lead to homeologous regions being missed
altogether in genome sequences, particularly in comparisons that require chromosome-level
assemblies. However, the fact that support for tetrasomic inheritance in other PKs identified as
tetrasomic through linkage mapping was consistent with that observed in genome analysis
strongly suggests that there is something unique with PK 06 rather than a fault with the genome
analyses conducted here. Perhaps, as suggested by Campbell et al (2019), the PK 06
chromosome arms are returning to a diploid state faster than the other seven tetrasomic
homeolog pairs or the tetrasomically inherited portion of PK 06 is smaller than other tetrasomic
PKs.
Another notable difference between linkage mapping and genome analysis was the
consistent classification of PK 01 as tetrasomic in the genome analysis (five of six species) but
not in any linkage map. PK 01 uniformly exhibited the least similarity between tetrasomic
homeologous pairs and was assigned to the putatively tetrasomic class of PKs with less certainty
by the machine learning algorithm. This suggests that PK 01 may have low levels of tetrasomy.
We also observed some consistent patterns of variation in sequence similarity within disomic
markers. For example, homeolog pairs for PK 24 and 21 generally displayed the lowest sequence
similarity, and homeolog pairs for PK 07 and 19 displayed higher similarity. Our study therefore
presents additional nuances into the rediploidization process by identifying a core group of
conserved tetrasomic homeologs, potentially intermediate homeologs (PK 01, 06) and
consistently diverged homeologs (PK 21, 24). Future investigations can be refined to examine
four well-defined categories across PKs: tetrasomic, intermediate, disomic, and most diverged.
This enhanced refinement should reduce noise from the incorrect pooling of homeologs and aid
in understanding the rediploidization process in salmonids.
Interestingly, more variation in sequence similarity was observed within tetrasomic
homeologs than was observed in disomic homeologs. For example, PK 23 has the second highest
sequence similarity in Arctic char, the fourth highest in rainbow trout, the sixth highest in
Atlantic salmon, and the seventh highest in grayling. While this may be in part due to differences
in genome assembly method and assembly quality, the fact that variation exists even among the
highest quality genomes (Atlantic salmon and rainbow trout) suggests that rediploidization rates
at tetrasomic PKs may vary among species, even though the same seven PKs are consistently
classified as tetrasomic. In other words, although there appears to be a large amount of

conservation of tetrasomic inheritance between species, our genome analyses also suggest some independence in the return to disomy since the three subfamilies of salmonid split ~ 50MYA.

Conclusions

Here we provide the most complete analysis of chromosomal rearrangements in coregonines using the currently available genomic resources as well and a newly developed haploid linkage map for cisco. We also suggest a naming system using a protokaryotype ID (PK) to facilitate a comparative analysis of all currently available salmonid genomes and linkage maps. Our study revealed that patterns of tetrasomic inheritance are largely conserved across the salmonids, but that there is substantial variation in these patterns both within and among species. For example, while the same seven PKs appear to be tetrasomically inherited across all species examined, their relative rates of sequence similarity differ within species, suggesting the potential of independent evolutionary trajectories following speciation. Additionally, we documented that analyses based on linkage maps do not identify the same tetrasomically inherited PKs as genome analyses and postulate that this may be due to inconsistencies with genome assemblies or due to differences the length of sequence used in comparisons. When inferring genomic patterns, more robust results can be obtained by using comparative approaches as opposed to a single resource. In conclusion, our study presents important insights about the WGD in salmon and also provides a framework that can be built upon to improve our understanding of WGDs both within and beyond salmonids.

Acknowledgements

This project was funded by the Great Lakes Restoration Initiative (GLRI). Special thanks to the USFWS crew members Chris Olds, Paul Haver, Kaley Genther, Steve Nimcheski, and Matt McLean for assistance in field sampling, USGS Great Lakes Science Center Aquatic Research Wet Lab for egg rearing, the University of Wisconsin- Stevens Point Molecular Conservation Genetics Lab for assistance in lab work, and the support of the Turing High Performance Computing cluster at Old Dominion University. Thanks to Kris Christensen, Claire Mérot, and Eric Rondeau for valuable and constructive comments on the manuscript, including the discussion of non-salmonid sex chromosomes. Any use of trade, product, or company name is for descriptive purposes only and does not imply endorsement by the U.S. Government.

Code Availability

Linkage mapping: https://github.com/DaniBlumstein/Cisco-Linkage-Map

Collecting *Coregonus* maps and running MAPCOMP:
https://github.com/bensutherland/coregonus_mapcomp

Genome homeology relationships: https://github.com/MacCampbell/residual-tetrasomy

References

Ali OA, et al. 2016. RAD Capture (Rapture): Flexible and Efficient Sequence-Based
Genotyping. Genetics 202: 389-400. doi: 10.1534/genetics.115.183665

Alkan C, Sajjadian S, Eichler EE 2011. Limitations of next-generation genome sequence
assembly. Nat Methods 8: 61-65. doi: 10.1038/nmeth.1527

Allendorf FW, et al. 2015. Effects of crossovers between homeologs on inheritance and
population genomics in polyploid-derived salmonid fishes. J Hered 106: 217-227. doi:
10.1093/jhered/esv015

Allendorf FW, Danzmann RG 1997. Secondary Tetrasomic Segregation of MDH-B and
Preferential Pairing of Homeologues in Rainbow Trout. Genetics 145: 1083-1092.

Allendorf, FW, Thorgaard GH 1984. Tetraploidy and the evolution of salmonid
fishes. Evolutionary genetics of fishes. Springer, Boston, MA. 1-53.

Angers B, Bernatchez L, Angers A, Desgroseillers L 1995. Specific microsatellite loci for brook
charr reveal strong population subdivision on a microgeographic scale. Journal of Fish Biology
47: 177-185.

Arrigo N, Barker MS 2012. Rarely successful polyploids and their legacy in plant genomes. Curr
Opin Plant Biol 15: 140-146. doi: 10.1016/j.pbi.2012.03.010

Barson NJ, et al. 2015. Sex-dependent dominance at a single locus maintains variation in age at
maturity in salmon. Nature 528: 405-408. doi: 10.1038/nature16062

Berthelot C, et al. 2014. The rainbow trout genome provides novel insights into evolution after
whole-genome duplication in vertebrates. Nature communications 5: 3657. doi:
10.1038/ncomms4657

Betancur-R, R et al. 2013. The tree of life and a new classification of bony fishes. PLoS
currents, 5.

Brieuc MS, Waters CD, Seeb JE, Naish KA 2014. A dense linkage map for Chinook salmon
(Oncorhynchus tshawytscha) reveals variable chromosomal divergence after an ancestral whole
genome duplication event. G3 (Bethesda) 4: 447-460. doi: 10.1534/g3.113.009316

Campbell MA, Hale MC, McKinney GJ, Nichols KM, Pearse DE 2019. Long-Term
Conservation of Ohnologs Through Partial Tetrasomy Following Whole-Genome Duplication in
Salmonidae. G3 (Bethesda) 9: 2017-2028. doi: 10.1534/g3.119.400070

Campbell MA, Lopez JA, Sado T, Miya M 2013. Pike and salmon as sister taxa: detailed
intraclade resolution and divergence time estimation of Esociformes + Salmoniformes based on
whole mitochondrial genome sequences. Gene 530: 57-65. doi: 10.1016/j.gene.2013.07.068

Chin CS, et al. 2016. Phased diploid genome assembly with single-molecule real-time
sequencing. Nat Methods 13: 1050-1054. doi: 10.1038/nmeth.4035

Chourrout D 1982. Gynogenesis caused by ultraviolet irradiation of salmonid sperm. Journal of
Experimental Zoology 223: 175-181.

Christensen KA, et al. 2018a. Chinook salmon (Oncorhynchus tshawytscha) genome and
transcriptome. PLoS One 13: e0195461. doi: 10.1371/journal.pone.0195461

Christensen KA, et al. 2018b. The Arctic charr (Salvelinus alpinus) genome and transcriptome
assembly. PLoS One 13: e0204076. doi: 10.1371/journal.pone.0204076

Clarke JT, Lloyd GT, Friedman M 2016. Little evidence for enhanced phenotypic evolution in
early teleosts relative to their living fossil sister group. Proc Natl Acad Sci U S A 113: 11531-
11536. doi: 10.1073/pnas.1607237113

Crete-Lafreniere A, Weir LK, Bernatchez L 2012. Framing the Salmonidae family phylogenetic
portrait: a more complete picture from increased taxon sampling. PLoS One 7: e46662. doi:
10.1371/journal.pone.0046662

Danecek P, et al. 2011. The variant call format and VCFtools. Bioinformatics 27: 2156-2158.
doi: 10.1093/bioinformatics/btr330

De-Kayne R, Feulner PGD 2018. A European Whitefish Linkage Map and Its Implications for
Understanding Genome-Wide Synteny Between Salmonids Following Whole Genome
Duplication. G3 (Bethesda) 8: 3745-3755. doi: 10.1534/g3.118.200552

De-Kayne R, Zoller S, Feulner, PG 2019 A de novo chromosome-level genome assembly
of Coregonus sp. "Balchen": one representative of the Swiss Alpine whitefish radiation. bioRxiv
doi: 10.1101/771600

Dion-Côté A-M, Symonová R, Ráb P, Bernatchez L. 2015. Reproductive isolation in a nascent
speciespair is associated with aneuploidy in hybrid offspring. Proceedings of the Royal Society
B: Biological Sciences. 282: 20142862.

Dion-Côté A-M, Symonová R, Lamaze FC, Pelikánová Š, Ráb P, Bernatchez L.
2017. Standingchromosomal variation associated with rapid divergence in Lake Whitefish

species pairs: the role of historical contingency and relevance for speciation. Molecular Ecology.
26:178–192"

Gagnaire PA, Normandeau E, Pavey SA, Bernatchez L 2013. Mapping phenotypic, expression
and transmission ratio distortion QTL using RAD markers in the Lake Whitefish (Coregonus
clupeaformis). Mol Ecol 22: 3036-3048. doi: 10.1111/mec.12127

Gerstein AC, Otto SP 2009. Ploidy and the causes of genomic evolution. Journal of Heredity
100: 571-581. doi:10.1093/jhered/esp057
Harris RS 2007. Improved pairwise alignment of genomic DNA. The Pennsylvania State
University.

Hollister JD 2015. Polyploidy: adaptation to the genomic environment. New Phytologist 205:
1034-1039. doi: 10.1111/nph.12939

Ishiguro NB, Miya M, Nishida M 2003. Basal euteleostean relationships: a mitogenomic
perspective on the phylogenetic reality of the "Protacanthopterygii". Molecular phylogenetics
and evolution, 27(3).

Karine Alix et al. 2017. Polyploidy and interspecific hybridization: partners for adaptation,
speciation and evolution in plants, Annals of Botany, 120: 183-194,
https://doi.org/101093/aob/mcx079

Kodama M et al. 2014. Comparative mapping between Coho Salmon (Oncorhynchus kisutch)
and three other salmonids suggests a role for chromosomal rearrangements in the retention of
duplicated regions following a whole genome duplication event. G3 (Bethesda) 4: 1717-1730.
doi: 10.1534/g3.114.012294

Koren S et al. 2017. Canu: scalable and accurate long-read assembly via adaptive k-mer
weighting and repeat separation. Genome Res 27: 722-736. doi: 10.1101/gr.215087.116

Kuhn M 2019. caret: Classification and Regression Training. R package version 6.0-84.

Kyriakidou M, Tai HH, Anglin NL, Ellis D, Stromvik MV 2018. Current Strategies of Polyploid
Plant Genome Sequence Assembly. Front Plant Sci 9: 1660. doi: 10.3389/fpls.2018.01660

Larson WA, et al. 2017. Genomic islands of divergence linked to ecotypic variation in sockeye
salmon. Mol Ecol 26: 554-570. doi: 10.1111/mec.13933

Larson WA, et al. 2015. Identification of Multiple QTL Hotspots in Sockeye Salmon
(Oncorhynchus nerka) Using Genotyping-by-Sequencing and a Dense Linkage Map. J Hered
107: 122-133. doi: 10.1093/jhered/esv099

Lien S, et al. 2016. The Atlantic salmon genome provides insights into rediploidization. Nature
533: 200-205. doi: 10.1038/nature17164

Limborg MT, McKinney GJ, Seeb LW, Seeb JE 2016a. Recombination patterns reveal
information about centromere location on linkage maps. Mol Ecol Resour 16: 655-661. doi:
10.1111/1755-0998.12484

Limborg MT, Seeb LW, Seeb JE 2016b. Sorting duplicated loci disentangles complexities of
polyploid genome masked by genotyping by sequencing. Molecular ecology 25: 2117–2129.

López JA, Chen WJ, & Ortí G 2004. Esociform phylogeny. Copeia, 2004(3), 449-464.

Macqueen DJ, Johnston IA 2014. A well-constrained estimate for the timing of the salmonid
whole genome duplication reveals major decoupling from species diversification. Proc Biol Sci
281: 20132881. doi: 10.1098/rspb.2013.2881

Mayrose I, et al. 2015. Methods for studying polyploid diversification and the dead end
hypothesis: a reply to Soltis et al. (2014). New Phytologist 206: 27-35. doi: 10.1111/nph.13192

Mayrose I, et al. 2011. Recently formed polyploid plants diversify at lower rates. Science 333:
1257. doi: 10.1126/science.1207205

McKinney GJ, et al. 2016. An integrated linkage map reveals candidate genes underlying
adaptive variation in Chinook salmon (Oncorhynchus tshawytscha). Mol Ecol Resour 16: 769-
783. doi: 10.1111/1755-0998.12479

McKinney G et al. 2017. Paralogs are revealed by proportion of heterozygotes and deviations in
read ratios in genotyping-by-sequencing data from natural populations. Mol Ecol Resour, 17:
656-669. doi:10.1111/1755-0998.12613

McKinney GJ, et al. 2019. Dense SNP panels resolve closely related Chinook salmon
populations. Canadian Journal of Fisheries and Aquatic Sciences. doi: 10.1139/cjfas-2019-0067

Norden CR 1961. Comparative osteology of representative salmonid fishes, with particular
reference to the grayling (Thymallus arcticus) and its phylogeny. Journal of the Fisheries Board
of Canada 18: 679-791.

Nugent CM, Easton AA, Norman JD, Ferguson MM, Danzmann RG 2017. A SNP Based
Linkage Map of the Arctic Charr (Salvelinus alpinus) Genome Provides Insights into the
Diploidization Process After Whole Genome Duplication. G3 (Bethesda) 7: 543-556. doi:
10.1534/g3.116.038026

Ohno S 1970. Evolution by gene duplication: Springer Science & Business Media.

Ohta T 1989. Role of gene duplication in evolution. Genome, 31(1), 304-310. doi:10.1139/g89-
048

Pan Q et al. 2019. Identification of the master sex determining gene in Northern pike (Esox
lucius) reveals restricted sex chromosome differentiation. PLoS Genet 15: e1008013. doi:
10.1371/journal.pgen.1008013

Patton JC, Gallaway BJ, Fechhelm RG, Cronin MA 1997. Genetic variation of microsatellite and
mitochondrial DNA markers in broad whitefish (Coregonus nasus) in the Colville and
Sagavanirktok rivers in northern Alaska. Canadian Journal of Fisheries and Aquatic Sciences 54:
1548-1556. doi: 10.1139/f97-062

Pearse DE, et al. 2019. Sex-dependent dominance maintains migration supergene in rainbow
trout. Nat Ecol Evol 3, 1731–1742 doi:10.1038/s41559-019-1044-6

Peichel CL, et al. 2004. The master sex-determination locus in threespine sticklebacks is on a
nascent Y chromosome. Current biology 16: 1416-1424. doi: 10.1016/j.cub.2004.08.030

Phillips R, Rab P 2001. Chromosome evolution in the Salmonidae (Pisces): an update. Biological
Reviews 76: 1-25. doi: 10.1017/S1464793100005613

Phillips RB, Reed KM, Ráb P 1996. Revised karyotypes and chromosome banding of coregonid
fishes from the Laurentian Great Lakes. Canadian Journal of Zoology 74: 323-329. doi:
10.1139/z96-040

Prince DJ, et al. 2017. The evolutionary basis of premature migration in Pacific salmon
highlights the utility of genomics for informing conservation. Science advances 3: e1603198.
doi: 10.1126/sciadv.1603198

Python Software Foundation version 2.7. Python Language Reference, Available at
http://www.python.org.

R Core Team 2018. R: A language and environment for statistical computing. Vienna, Austria: R
Foundation for Statistical Computing.

Rastas P 2017. Lep-MAP3: robust linkage mapping even for low-coverage whole genome
sequencing data. Bioinformatics 33: 3726-3732. doi: 10.1093/bioinformatics/btx494

Rastogi S, Liberles DA 2005. Subfunctionalization of duplicated genes as a transition state to
neofunctionalization. BMC Evolutionary Biology 5: 28. doi:10.1186/1471-2148-5-28

Ripley B, Venables W 2019. class: Functions for Classification. R package version 7.3-15.

Robertson FM, et al. 2017. Lineage-specific rediploidization is a mechanism to explain time-lags
between genome duplication and evolutionary diversification. Genome Biol 18: 111. doi:
10.1186/s13059-017-1241-z

Rochette NC, Catchen JM 2017. Deriving genotypes from RAD-seq short-read data using
Stacks. Nat Protoc 12: 2640-2659. doi: 10.1038/nprot.2017.123

Rochette NC, Rivera-Colón AG, Catchen JM 2019. Stacks 2: Analytical Methods for Paired-end
Sequencing Improve RADseq-based Population Genomics. bioRxiv: 615385. doi:
10.1101/615385

Rogers SM, Marchand M-H, Bernatchez L 2004. Isolation, characterization and cross-salmonid
amplification of 31 microsatellite loci in the lake whitefish (Coregonus clupeaformis, Mitchill).
Molecular Ecology Notes 4: 89-92. doi: 10.1046/j.1471-8286.2003.00578.x

Rondeau EB, et al. 2014. The genome and linkage map of the northern pike (Esox lucius):
conserved synteny revealed between the salmonid sister group and the Neoteleostei. PLoS One
9: e102089. doi: 10.1371/journal.pone.0102089

Ruan J, Li H 2019. Fast and accurate long-read assembly with wtdbg2. bioRxiv: 530972. doi:
10.1101/530972

Sakamoto T, et al. 2000. A Microsatellite Linkage Map of Rainbow Trout (Oncorhynchus
mykiss) Characterized by Large Sex-Specific Differences in Recombination Rates. Genetics 155:
1331–1345.

Santini F, Harmon LJ, Carnevale G, Alfaro ME 2009. Did genome duplication drive the origin of
teleosts? A comparative study of diversification in ray-finned fishes. BMC Evol Biol 9: 194. doi:
10.1186/1471-2148-9-194

Savilammi T, et al. 2019. The Chromosome-Level Genome Assembly of European Grayling
Reveals Aspects of a Unique Genome Evolution Process Within Salmonids. G3 (Bethesda) 9:
1283-1294. doi: 10.1534/g3.118.200919

Schaeffer SW 2018. Muller "Elements" in Drosophila: How the Search for the Genetic Basis for
Speciation Led to the Birth of Comparative Genomics. Genetics 210: 3-13. doi:
10.1534/genetics.118.301084

Schwager EE, et al. 2017. The house spider genome reveals an ancient whole-genome
duplication during arachnid evolution. BMC Biol 15: 62. doi: 10.1186/s12915-017-0399-x

Selmecki AM, et al. 2015. Polyploidy can drive rapid adaptation in yeast. Nature 519: 349-352.
doi: 10.1038/nature14187

Sutherland BJG, et al. 2016. Salmonid Chromosome Evolution as Revealed by a Novel Method
for Comparing RADseq Linkage Maps. Genome Biol Evol 8: 3600-3617. doi:
10.1093/gbe/evw262

Sutherland BJG, Rico C, Audet C, Bernatchez L 2017. Sex Chromosome Evolution,
Heterochiasmy, and Physiological QTL in the Salmonid Brook Charr Salvelinus fontinalis. G3
(Bethesda) 7: 2749-2762. doi: 10.1534/g3.117.040915

Tarpey CM, Seeb JE, McKinney GJ, Seeb LW 2017. A dense linkage map for odd-year lineage
pink salmon incorporating duplicated loci. In. Alaska Department of Fish and Game: School of
Aquatic and Fishery Sciences, University of Washington. p. 1-50.

Taylor JS, Braasch I, Frickey T, Meyer A, Van de Peer Y 2003. Genome duplication, a trait
shared by 22000 species of ray-finned fish. Genome Res 13: 382-390. doi: 10.1101/gr.640303

Van de Peer Y, Mizrachi E, Marchal K 2017. The evolutionary significance of polyploidy. Nat
Rev Genet 18: 411-424. doi: 10.1038/nrg.2017.26

Vanneste K, Maere S, Van de Peer Y 2014. Tangled up in two: a burst of genome duplications at
the end of the Cretaceous and the consequences for plant evolution. Philos Trans R Soc Lond B
Biol Sci 369. doi: 10.1098/rstb.2013.0353

Varadharajan S, et al. 2018. The Grayling Genome Reveals Selection on Gene Expression
Regulation after Whole-Genome Duplication. Genome Biol Evol 10: 2785-2800. doi:
10.1093/gbe/evy201

Waples RK, Seeb LW, Seeb JE 2016. Linkage mapping with paralogs exposes regions of
residual tetrasomic inheritance in chum salmon (Oncorhynchus keta). Mol Ecol Resour 16: 17-
28. doi: 10.1111/1755-0998.12394

Wendel JF. 2000. Genome evolution in polyploids. In. Plant molecular evolution: Springer. p.
225-249.

Wittbrodt K, Meyer A, Schartl M 1998. More genes in fish? BioEssays 20: 511-515. doi:
10.1002/(sici)1521-1878(199806)20:6<511::Aid-bies10>3.0.Co;2-3

Wolfe KH 2001. Yesterday's polyploids and the mystery of diploidization. Nature reviews
genetics 2: 333.

Wood TE, et al. 2009. The frequency of polyploid speciation in vascular plants. Proc Natl Acad
Sci U S A 106: 13875-13879. doi: 10.1073/pnas.0811575106

Zhan SH, Glick L, Tsigenopoulos CS, Otto SP, Mayrose I 2014. Comparative analysis reveals
that polyploidy does not decelerate diversification in fish. J Evol Biol 27: 391-403. doi:
10.1111/jeb.12308

Table 1. Results for the female and male cisco linkage maps. Duplicated and singleton loci are from the female linkage map, and % duplicated is the percentage of duplicated loci on each female linkage group (LG). LG type is denoted with acrocentric (A) and metacentric (M).

Cisco LG	Duplicated Loci	Singleton Loci	% Duplicated	Length Female (cM)	Male (cM)	Total Loci Female	Male	Loci/cM Female	Male	LG Type
Cart01	212	510	0.42	101.35	87.14	722	224	7.12	2.57	M
Cart02	394	384	1.03	97.77	61.11	778	159	7.96	2.6	M
Cart03	226	477	0.47	93.56	78.98	703	248	7.51	3.14	M
Cart04	48	547	0.09	92.45	67.80	595	194	6.44	2.86	M
Cart05	283	584	0.48	91.73	58.07	867	219	9.45	3.77	M
Cart06	108	729	0.15	91.36	68.95	837	278	9.16	4.03	M
Cart07	189	496	0.38	57.72	60.40	685	204	11.87	3.38	M
Cart08	189	532	0.36	57.56	51.62	721	170	12.53	3.29	M
Cart09	238	271	0.88	64.53	44.47	509	92	7.89	2.07	A
Cart10	21	488	0.04	63.98	49.68	509	152	7.96	3.06	A
Cart11	38	510	0.07	63.72	49.41	548	165	8.6	3.34	A
Cart12	393	177	2.22	63.18	74.69	570	80	9.02	1.07	A
Cart13	27	558	0.05	62.90	53.67	585	181	9.3	3.37	A
Cart14	73	454	0.16	62.66	59.74	527	140	8.41	2.34	A
Cart15	96	318	0.30	62.45	48.26	414	121	6.63	2.51	A
Cart16	24	456	0.05	62.25	45.80	480	153	7.71	3.34	A
Cart17	27	478	0.06	62.25	53.19	505	160	8.11	3.01	A
Cart18	25	446	0.06	62.16	42.08	471	136	7.58	3.23	A
Cart19	29	541	0.05	61.85	75.08	570	213	9.22	2.84	A
Cart20	31	536	0.06	60.19	48.36	567	224	9.42	4.63	A
Cart21	31	388	0.08	59.04	61.23	419	142	7.1	2.32	A

Cart22	20	544	0.04	58.96	49.49	564	204	9.57	4.12	A
Cart23	52	473	0.11	58.57	55.39	525	162	8.96	2.92	A
Cart24	13	556	0.02	58.38	57.23	569	195	9.75	3.41	A
Cart25	26	458	0.06	58.28	51.08	484	161	8.3	3.15	A
Cart26	30	445	0.07	58.09	50.91	475	147	8.18	2.89	A
Cart27	32	499	0.06	58.00	60.69	531	213	9.16	3.51	A
Cart28	27	444	0.06	57.79	54.44	471	141	8.15	2.59	A
Cart29	15	404	0.04	57.78	62.64	419	149	7.25	2.38	A
Cart30	29	384	0.08	57.76	71.50	413	141	7.15	1.97	A
Cart31	25	541	0.05	56.93	77.82	566	184	9.94	2.36	A
Cart32	19	478	0.04	56.56	75.17	497	177	8.79	2.35	A
Cart33	20	460	0.04	56.37	49.57	480	146	8.52	2.95	A
Cart34	262	82	3.20	55.54	83.66	344	62	6.19	0.74	A
Cart35	42	254	0.17	54.74	40.54	296	81	5.41	2	A
Cart36	19	332	0.06	54.45	45.50	351	117	6.45	2.57	A
Cart37	25	402	0.06	53.16	55.82	427	122	8.03	2.19	A
Cart38	25	431	0.06	50.97	73.21	456	171	8.95	2.34	A
Cart39	0	0	0.00	0.00	45.13	0	41	0	0.91	A
Cart40	0	0	0.00	0.00	58.46	0	71	0	1.21	A
Averages	89.03	449.13	0.20	64.66	58.95	538.16	158.5	8.36		
Total	3383	17067	0.20	2456.98	2357.98	20450	6340	317.71		

Table 2. MAPCOMP results documenting homologous chromosomes for the three coregonines; cisco, lake whitefish (Gagnaire et al. 2013), and European whitefish (De-Kayne and Feulner 2018), integrated with Atlantic salmon (Lien et al. 2016), brook trout (Sutherland et al. 2016), and Chinook salmon (Brieuc et al. 2014). Homologous chromosomes for all species are named according to the corresponding Northern Pike linkage group as a reference (Rondeau et al. 2014), as per Sutherland et al. (2016), here termed protokaryotype ID (PK). Letters after linkage group (LG) names indicate the first (a) or second (b) arm of the LG, ^ indicates weak evidence and * indicates uncertainty between homeologs from MAPCOMP analysis.

Northern Pike (PK)	Cisco	European Whitefish	Lake Whitefish	Brook Trout	Atlantic Salmon	Chinook Salmon
01.1	Cart23	Calb16	Cclu28	Sf25	Ssa20b	Ots13q
01.2	Cart14	Calb33	Cclu35	Sf38	Ssa09c	Ots14q
02.1	Cart01a	Calb02b^*	Cclu04a	Sf06a	Ssa26	Ots04q
02.2	Cart12	Calb02b^*	Cclu04a^*	Sf28	Ssa11a	Ots12q
03.1	Cart25	Calb19	Cclu25	Sf22	Ssa14a	Ots10q
03.2	Cart26	Calb22	Cclu26	Sf11	Ssa03a	Ots28
04.1	Cart30	Calb29	Cclu16	Sf33	Ssa09b	Ots08q
04.2	Cart21	Calb30	Cclu29	Sf07b	Ssa05a	Ots21
05.1	Cart06b	Calb01a	Cclu05a	Sf01a	Ssa19b	Ots24
05.2	Cart18	Calb35 or Calb40	Cclu15	Sf27	Ssa28	Ots25
06.1	Cart06a	Calb01b	Cclu05b	Sf01b	Ssa01b	Ots01q
06.2	Cart15	Calb27	Cclu05b^*	Sf36	Ssa18a	Ots06q
07.1	Cart20	Calb06	Cclu13	Sf08b	Ssa13b	Ots09p
07.2	Cart19	Calb07	Cclu08	Sf09	Ssa04b	Ots30
08.1	Cart27	Calb17	Cclu36	Sf04a	Ssa23	Ots01p
08.2	Cart03a	Calb08	Cclu06a	Sf17	Ssa10a	Ots05q
09.1	Cart03b*	missing	Cclu06b	Sf42	Ssa02b	Ots32
09.2	Cart34*	missing	missing	Sf03b	Ssa12a	Ots02q
10.1	Cart08a	Calb20b	Cclu10	Sf23	Ssa27	Ots13p
10.2	Cart04b	Calb09a	Cclu24a	Sf34	Ssa14b	Ots31
11.1	Cart05a	Calb13b	Cclu18	Sf14	Ssa06a	Ots27
11.2	Cart09	Calb34	missing	Sf08a	Ssa03b	Ots09q
12.1	Cart33	Calb14	Cclu27	Sf18	Ssa13a	Ots22
12.2	Cart16	Calb28	Cclu14	Sf30	Ssa15b	Ots16q
13.1	Cart17	Calb25	Cclu34	Sf06b	Ssa24	Ots04p
13.2	Cart32	Calb31	Cclu37	Sf40	Ssa20a	Ots12p
14.1	Cart01b	Calb02a	Cclu04b	Sf13	Ssa01c	Ots20
14.2	Cart38	Calb11	Cclu33	Sf10	Ssa11b	Ots33

15.1	Cart10	Calb18	Cclu31	Sf35	Ssa09a	Ots08p
15.2	Cart31	Calb10	Cclu22	Sf12	Ssa01a	Ots11q
16.1	Cart07a	Calb03b	Cclu02b or Cclu03	Sf26	Ssa21	Ots26
16.2	Cart28	Calb21	Cclu32	Sf24	Ssa25	Ots03q
17.1	Cart24	Calb05	Cclu38	Sf03a	Ssa12b	Ots02p
17.2	Cart22	Calb12	Cclu21	Sf21	Ssa22	Ots07q
18.1	Cart29	Calb24	Cclu40	Sf19	Ssa15a	Ots05p
18.2	Cart37	Calb23	Cclu17	Sf31	Ssa06b	Ots18
19.1	Cart13	Calb04	Cclu30	Sf15	Ssa10b	Ots19
19.2	Cart11	Calb15b	Cclu11	Sf20	Ssa16a	Ots06p
20.1	Cart08b^	Calb36	Cclu01a	Sf07a	Ssa05b	Ots23
20.2	Cart02b	Calb20a	missing	Sf29	Ssa02a	Ots03p
21.1	Cart05b	Calb13a	Cclu12	Sf05b	Ssa29	Ots29
21.2	Cart36	Calb26	Cclu39	Sf16	Ssa19a	Ots16p
22.1	Cart02a	Calb39^	missing	Sf39	Ssa17a	Ots07p
22.2	missing	Calb15a	Cclu19^	Sf05a	Ssa16b	Ots14p
23.1	Cart07b^*	Calb03a	Cclu02a	Sf02b	Ssa07b	Ots15p
23.2	Cart07b^*	missing	Cclu01b^	Sf37	Ssa17b	Ots17
24.1	Cart04a	Calb09b	Cclu24b	Sf02a	Ssa07a	Ots15q
24.2	Cart35	Calb32	Cclu23	Sf32	Ssa18b	Ots10p
25.1	missing	missing	Cclu09^	Sf04b	Ssa04a	Ots34
25.2	missing	missing	missing	Sf41	Ssa08a	Ots11p

Figure 1. A) Female linkage map for cisco containing 20,450 loci. Each dot represents a locus, duplicated loci are blue and singleton loci are gray. Lengths are in centimorgans (cM). Approximate location of centromeres for metacentric LGs are denoted in red. Metacentric LGs were identified through homologous relationships of chromosome arms with other Salmonids via MAPCOMP. B) Circos plot of cisco LGs highlighting 17 supported homeologous regions within the linkage map. Included in the 17 homeologus regions are six of the eight regions that are likely still residually tetrasomic across the Salmonids. Colors represent the number of markers supporting relationship, with darker colors representing higher marker numbers (maximum support = 86 markers) and theoretical links inferred via MAPCOMP.

Figure 2. Fusions and fissions in the Coregoninae and Salmoninae lineages. This is an extension of Figure 4 from Sutherland et al. (2016). White boxes display the fusion events, where the homologous chromosomes for all species are named according to the protokaryotype ID. Bold and underlined chromosome numbers are the homeologous pairs that exhibit residual tetraploidy (i.e "magic eight"), * indicate uncertainty in one species, and ** indicates uncertain in two species (i.e., *C. artedi* is ambiguous for homeolog 09.1 or 09.2 while *C. lavaretus* is missing 09.1 and 09.2). Above the species names are conserved fusions, whereas below are the species-specific fusions. The phylogeny is adapted from (Crete-Lafreniere et al. 2012). Branch lengths do not represent phylogenetic distance, only relative phylogenetic position.
[1]Arms 11.1-21.1 were fused in both cisco and European whitefish, but likely underwent fission in lake whitefish.

Figure 3. Ranking of homeologous chromosome pairs based on putative residual tetrasomic inheritance as measured by the number of markers shared among homeologs for linkage maps or percent sequence similarity for genomes. A lower rank represents more marker pairs supporting a homeolog and or a higher sequence similarity. Chromosomes for all species are named according to the protokaryotype ID (PK). PKs are ordered in the figure by averaging the ranks across all species and then sorting the averages from smallest to largest (i.e., ordered from highest support for residual tetrasomy to lowest). Grey indicates that no duplicated loci could be mapped to both homeologs. Species abbreviations are grayling (T. thy), Atlantic salmon (S. sal), Arctic char (S. alp), rainbow trout (O. myk), Chinook salmon (O. tsh), cisco (C. art), and coho salmon (O. kis).

Figure 4. Distribution of protokaryotypes (PK) similarity in aligned sections between the homeolog pairs across salmonids based on genome assemblies. For each species with a genome sequence, the percent similarity (y – axis) of the 25 PK pairs as shown as box plots. PK pairs are ranked from highest to lowest median similarity for each species (x – axis), with the average similarity of protokaryotypes presented as a dashed line. The classification of protokaryotypes by the machine learning approach described in the main text into putatively tetrasomic and disomic pairs is shown through coloring of the boxplots into purple (putatively tetrasomic) and yellow (putatively disomic). The number of alignments used in computing similarity is presented at the top of each bar. Those protokaryotypes that did not receive the highest observed voting proportion for the assigned class are indicated with an asterisk (*). PKs with high variance (e.g. PK 11 in S. alp) may be due to methodological limitations that have caused additional non-homeologous chromosome arms to be included in the comparisons (see discussion). Species abbreviations are grayling (T. thy), Atlantic salmon (S. sal), Arctic char (S. alp), rainbow trout (O. myk), and Chinook salmon (O. tsh)

Supplementary tables and figures:

S1. Sampling information for cisco (Coregonus artedi) families collected from Lake Huron the number of individuals used. The number of offspring shown only includes those used for mapping after the removal of individuals with low sequencing coverage. The numbers of SNPs shown include those retained following quality control filtering but before inclusion in linkage mapping.

S2. Male linkage map for cisco (Coregonus artedi) containing 6340 loci. Each dot represents a locus. Lengths are in centimorgans (cM).

S3. Information for each marker on the female and male cisco (Coregonus artedi) linkage maps. Tag is the RAD tag, and marker name is the tag followed by a designation used to differentiate duplicated loci. The "Sequence P1 column" is the sequence from the single end read for each RAD tag, and the "Sequence PE" is the sequence obtained from paired-end assemblies.

S4. MAPCOMP determination of homologous chromosome arms.

S5: Probable metacentric chromosomes from the MAPCOMP analysis for coregonines.

S6. Homeologous chromosome pairs for currently available haploid linkage maps and the number of maker pairs supporting homeology (Kodama et al. 2014; Larson et al. 2015; Waples et al. 2016; Tarpey et al. 2017). Homologous chromosomes are named according to the corresponding Northern Pike linkage group Protokaryotype ID (PK).

S7. Homeologous chromosome pairs for all available salmonid genomic resources.

S8. Support for classifications from k – nearest neighbor machine learning algorithm. Cross validation indicated five nearest neighbors should be used for classification across species. The vote proportion for assignment to either putatively tetrasomic or putatively disomic categories are presented as the proportion of votes out of five at the top of each bar. Median percent similarity for each protokaryotype pair is presented as the y – axis and protokaryotype pairs are ranked from most similar to least (x – axis).