Genomic characterization of the non-O1/non-O139 *Vibrio cholerae* strain that caused a gastroenteritis outbreak in Santiago, Chile, 2018

Mónica Arteaga[1]*, Juliana Velasco[1]*, Shelly Rodriguez[1], Maricel Vidal[2], Carolina Arellano[3], Francisco Silva[4], Leandro J. Carreño[5,6], Roberto Vidal[3,6], David A. Montero[3,5] **

[1]Servicio de Urgencia Infantil, Hospital Clínico de la Universidad de Chile "Dr. José Joaquín Aguirre", Santiago, Chile. [2]Laboratorio de Salud Pública Ambiental y Laboral, Secretaría Regional Ministerial de Salud Región Metropolitana, Santiago, Chile. [3]Programa de Microbiología y Micología, Facultad de Medicina, Universidad de Chile, Santiago, Chile. [4]Servicio de Laboratorio Clínico, Hospital Clínico de la Universidad de Chile "Dr. José Joaquín Aguirre", Santiago, Chile. [5]Programa de Inmunología, Instituto de Ciencias Biomédicas, Facultad de Medicina, Universidad de Chile, Santiago, Chile. [6]Instituto Milenio de Inmunología e Inmunoterapia, Facultad de Medicina, Universidad de Chile, Santiago, Chile.

* Co-first authors. ** Corresponding author. Correspondence should be addressed to Montero DA. (Email: davidmontero@med.uchile.cl)

Keywords: Non-O1/non-O139 Vibrio cholerae, Gastroenteritis outbreak, Type III secretion system (T3SS), Type VI secretion system (T6SS), Multidrug resistance genomic island

Abstract

We investigated the virulence determinants and phylogeny of a non-O1/non-O139 *Vibrio cholerae* strain that caused an outbreak in Santiago, Chile, 2018. Our results suggest that the high virulence of this strain, as well as other non-O1/non-O139 strains, may be the result of the acquisition of genomic islands carrying antibiotic resistance genes, T3SS and T6SS.

V. cholerae strains belonging to serogroups O1 and O139 are known to cause cholera
outbreaks and epidemics. These serogroups generally produce the cholera toxin (CT) and the
toxin coregulated pilus (TCP), which are responsible for secretory diarrhea and intestinal
colonization, respectively (1). Serogroups other than O1 and O139, called non-O1/non-O139,
typically lack the CT and TCP-encoding genes (2). However, several non-O1/non-O139 *V.*
cholerae strains harbor additional virulence factors that contribute to pathogenicity (3). In fact,
non-O1/non-O139 *V. cholerae* strains have been isolated from sporadic cases of gastroenteritis,
bacteremia and wound infections (1,2,4). To date, the pathogenic mechanisms of non-O1/non-
O139 *V. cholerae* strains have not been fully investigated.
94
There were several cholera outbreaks in South America in the 1990s (5). In Chile, the last
reported cases of cholera corresponded to the 1997-1998 outbreak of San Pedro de Atacama.
Until 2017, only sporadic cases of gastroenteritis or bacteremia caused by non-O1/non-O139 *V.*
cholerae strains had been reported in the country (6). On August 2, 2018, an outbreak of acute
gastroenteritis started in Santiago, Chile. As of February 16, 2019, 55 gastroenteritis cases were
reported, of which 13 required hospitalization, with diarrhea, nausea and vomiting being the main
symptoms. The epidemiological study of the outbreak performed at the Chilean National Health
Institute (Instituto de Salud Pública de Chile, ISP) identified a clonal strain of non-toxigenic and
non-O1/non-O139 *V. cholerae* as the etiology agent in 44/55 cases, although the source of
contamination could not be determined (7). Here, we performed comparative genomic and
phylogenetic analyses to decipher the virulence determinants and phylogenetic origin of this
outbreak strain.
107

The Study

The *V. cholerae* strain characterized in this study, which we named the Santiago strain, was
isolated from a 12-year-old boy hospitalized with bloody diarrhea and abdominal pain. This was
one of the first gastroenteritis cases of the outbreak. The microbiological study of the strain was
approved by written consent of the parents and the Ethics Committee of the Clinical Hospital
University of Chile. Moreover, a Pulsed-field gel electrophoresis analysis performed at the ISP
indicated that this and another 43 strains isolated during the outbreak were clonal (personal
communication of Dr. Juan Carlos Hormazabal, Sub-department of Infectious Diseases, ISP).
116

Initially, an antimicrobial drug susceptibility test showed that the Santiago strain is resistant to
trimethoprim-sulfamethoxazole, erythromycin and nalidixic acid (**Table**). Next, the phylogeny of
this outbreak strain was investigated. For this, its genomic DNA was sequenced (**Supplementary**
information) and the draft genome deposited in GenBank under the accession number
SRLP00000000. In addition, a set of genomes of *V. cholerae* strains isolated worldwide that are
available in GenBank were included in the phylogenetic analysis (**Supplementary Table 1**). As
shown in the maximum likelihood phylogenetic tree (**Figure 1A**), the strains were clustered into
twelve lineages. While lineages 4 and 5 clustered the O1, O139 and O65 strains, the rest of the
lineages clustered only non-O1/non-O139 strains. In particular, the Santiago strain was clustered

in lineage 2 along with strains isolated from India, Bangladesh and Haiti. Moreover, MLST
sequence types were consistent with the topology of the tree.

This outbreak strain caused several hospitalizations but lacks the CT and, consequently other
virulence genes must be contributing to its pathogenicity. Therefore, we analyzed its genome
searching for other virulence genes (**Supplementary Table 2**). As expected, the strain lacks the
Phage CTXφ and the genomic islands (GIs) (VPI-I, VPI-II, VSP-I and VSP-II) that are generally
carried by O1 and O139 strains (**Supplementary Figure 1**). However, this strain carries genes
that encode toxins such HlyA and MARTX, and proteins of the Type III and Type VI secretion
systems (T3SS/T6SS) (**Figure 1B**). Similarly, most of the non-O1/non-O139 strains lack the
Phage CTXφ and the GIs mentioned; rather, they carry genes encoding proteins of T3SS and
T6SS.

We analyzed the genetic context of the T3SS and T6SS genes in the Santiago strain and
found that they are located in GIs. The GIVch-T3SS identified was previously reported in the *V.*
cholerae str. AM-19226 and promotes colonization and infection (8). We found that this GI of ~64-
kb is inserted in the tRNA-*ser* gene and located next to a nan-nag region involved in the sialic
acid catabolism (**Figure 2A)**, which is also harbored by the VPI-II (9). The second GI, which we
named GIVch-T6SS$_{Santiago}$, was partially identified and contains a CRISPR-Cas region and genes
that encode Hcp and VgrG alleles, which are structural T6SS components (**Figure 2B**). The
GIVch-T6SS$_{Santiago}$ was also identified in the *V. cholerae* str. HC36A1. The Hcp from GIVch-
T6SS$_{Santiago}$ has 96.5% amino acid identity with the Hcp1 and Hcp2 alleles reported in *V. cholerae*.
In contrast, the VgrG from GIVch-T6SS$_{Santiago}$ has 66.7%, 66.6% and 59.4% amino acid identity
with the VgrG-1, VgrG-2 and VgrG-3 alleles reported in V. *cholerae*, respectively (10). Therefore,
we named this uncharacterized allele VgrG-4. Moreover, we found that Hcp and VgrG-4 alleles
from GIVch-T6SS$_{Santiago}$ are also harbored by the GIVchS12 (11). Thus, the identification of T3SS
genes and the VgrG-4 allele in non-O1/non-O139 strains of several lineages suggests these GIs
are widely distributed (**Figure 1**).

Finally, since the Santiago strain is multidrug-resistant (**Table**), we analyzed its genome
searching for genes or mutations that mediate its antimicrobial resistance profile. As a result, we
identified a multidrug resistance (MDR) genomic island of ~31-kb that we named GIVch-
MDR$_{Santiago}$ (**Figure 2C**). This GI harbors the *sul1* and *dfrA15* genes, which confer resistance to
sulfonamide and trimethoprim, respectively. The genetic structure of the GIVch-MDR$_{Santiago}$ is also
relatively similar to the GIVchHai6 (12). On the other hand, the Santiago strain has the mutations
S83I and A171S in the GyrA protein, which confer resistance to nalidixic acid (13)
(**Supplementary information**). Genes or mutations conferring resistance to erythromycin were
not identified.

Conclusions

The fact that a *V. cholerae* strain like the one characterized in the present study, which lacks the classical virulence factors (i.e. CT and CTP) yet causes a gastroenteritis outbreak with several hospitalized patients, represents a public health concern that justifies the permanent surveillance of these pathogens. Moreover, the multidrug-resistant phenotype of the Santiago strain, as well as the spread of MDR islands (12), are also epidemiological factors that need to be considered by public health authorities. In particular, the Santiago strain was isolated from a case of bloody diarrhea. Although bloody diarrhea is not a common symptom caused by *V. cholerae*, there are some reports in which non-O1/non-O139 strains have caused it (14). The above is of major importance for the diagnosis and treatment of these infections.

We show that the population structure of *V. cholerae* and particularly of the non-O1/non-O139 strains is heterogeneous, having a clear phylogenetic diversity and where known virulence factors are widely distributed among lineages. Indeed, we found that in the absence of the classic virulence factors, the non-O1/non-O139 strains have acquired several GIs encoding T3SS and T6SS, which may enhance their virulence. T3SS has a key role in the pathogenesis of *Vibrio parahaemolyticus* (15). Although less studied, the role of T3SS in the pathogenesis of *V. cholerae* and other pathogenic *Vibrio* species has begun to be understood (8,16). These genes could be considered molecular risk markers for these pathogens and may be useful in epidemiological monitoring studies. We also identified new Hcp and VgrG alleles that in future studies will be functionally characterized. In conclusion, our results highlight the pathogenic potential of the Santiago strain as well as other non-O1/non-O139 *V. cholerae* strains.

About the authors

Mónica Arteaga and Juliana Velasco are medical doctors interested in infectious diseases and the epidemiology of enteropathogens.

Acknowledgments

This work was supported by Postdoctoral FONDECYT grant 3190524, awarded to D. Montero, and FONDECYT grant 1161161, awarded to R. Vidal. We wish to thank Dr. Juan Carlos Hormazabal for the critical reading of this manuscript and the epidemiological information of the Santiago strain.

210 **Table 1.** Antimicrobial resistance profile of the *V. cholerae* str. Santiago [1]

Antimicrobial agent	Disk diffusion	Zone Diameter (mm)	MIC (µg/ml)	Interpretative criteria
Ampicillin	10 µg	20	-	Susceptible
Azithromycin	-	-	2	Susceptible
Erythromycin [2]	-	-	8	Resistant
Gentamicin	10 µg	16	-	Susceptible
Tetracycline	30 µg	25	-	Susceptible
Ciprofloxacin	5 µg	22	-	Susceptible
Trimethoprim-Sulfamethoxazole	1.25/23.75 µg	6	-	Resistant
Chloramphenicol	30 µg	31	-	Susceptible
Nalidixic acid [2]	30 µg	6	-	Resistant

211 [1] The disk diffusion method or broth dilution was performed according to Clinical and Laboratory Standards
212 Institute guidelines. [2] Interpretive criteria adapted from Enterobacteriaceae

Figure 1. **Phylogenetic relationship between 70 *V. cholerae* strains**. **A)** Maximum likelihood phylogenetic tree (midpoint rooted) based on whole genome SNPs (146,534 SNPs within 2,483,145 positions, which were found in all analyzed genomes). The genome of the *V. cholerae* str. N16961 was used as the reference. Bayesian analysis of population structure grouped the strains into five sequence clusters (SC; SC1 to SC5), which were further divided into 12 lineages. The epidemiological data of each strain is shown, including country and year of isolation. MLST sequence types (STs) are shown. **B)** Heat map showing the presence, absence and variation of major virulence-associated genes distributed among the strains. Presence and variation (nucleotide identity levels, ranging from 80 to 100%) for each gene are indicated by color intensity (red to yellow), as shown in the legend. The analysis was performed using BLASTn. Absence was defined as an identity and/or gene coverage of less than 80% and 60%, respectively and is indicated in gray.

Figure 2. Comparison of the genetic structures of genomic islands carried by the _V. cholerae_ str. Santiago. Predicted genes and the direction of transcription are represented as block arrows. Genes are color-coded according to gene function, as indicated in the legends at the bottom. The names of some genes are indicated. Conserved regions are shaded in gray and the intensity of the color indicates nucleotide identity levels, as indicated in the legends at the right. Contig boundaries are shown as red lines. **A)** Genomic islands encoding a T3SS. GenBank accessions: GI_Vch_-T3SS_AM-19226_ (AATY02000004 and AATY02000003), GI_Vch_-T3SS_10432-62_ (CP010812), GI_Vch_-T3SS_Santiago_ (SRLP00000000). **B)** Genomic islands encoding CRISPR-Cas and T6SS genes. GenBank accessions: GI_Vch_-T6SS_HC-36A1_ (AXDR01000008.1), GI_Vch_-T6SS_Santiago_ (SRLP00000000), GI_Vch_S12 (KU722393). Open reading frames (ORFs) of the GI_Vch_-T6SS_Santiago_ are listed in Supplementary Table 3. **C)** Multidrug resistance (MDR) islands. The GI_Vch_-MDR_Santiago_ (GenBank accession: SRLP00000000) harbors the _sul1_ and _dfrA15_ genes, which confer resistance to sulfonamide and trimethoprim, respectively. ORFs of the GI_Vch_-MDR_Santiago_ are listed in Supplementary Table 4. GI_Vch_Hai6 (GenBank accession: AXDR01000001).

References

1.	Dutta D, Chowdhury G, Pazhani GP, Guin S, Dutta S, Ghosh S, et al. Vibrio cholerae non-O1, non-O139 serogroups and cholera-like diarrhea, Kolkata, India. Emerg Infect Dis. 2013 Mar;19(3):464–7.

2.	Octavia S, Salim A, Kurniawan J, Lam C, Leung Q, Ahsan S, et al. Population Structure and Evolution of Non-O1/Non-O139 Vibrio cholerae by Multilocus Sequence Typing. Zhou D, editor. PLoS One. 2013 Jun 11;8(6):e65342.

3.	Chatterjee S, Ghosh K, Raychoudhuri A, Chowdhury G, Bhattacharya MK, Mukhopadhyay AK, et al. Incidence, Virulence Factors, and Clonality among Clinical Strains of Non-O1, Non-O139 Vibrio cholerae Isolates from Hospitalized Diarrheal Patients in Kolkata, India. J Clin Microbiol. 2009 Apr 1;47(4):1087–95.

4.	Deshayes S, Daurel C, Cattoir V, Parienti J-J, Quilici M-L, de La Blanchardière A. Non-O1, non-O139 Vibrio cholerae bacteraemia: case report and literature review. Springerplus. 2015 Dec 5;4(1):575.

5.	Domman D, Quilici M, Dorman MJ, Njamkepo E, Mutreja A, Mather AE, et al. Integrated view of Vibrio cholerae in the Americas. Science. 2017 Nov 10;358(6364):789–93.

6.	Olivares F, Domínguez I, Dabanch J, Porte L, Ulloa MT, Osorio G. Bacteriemia por Vibrio cholerae no-O1/no-O139 que porta una región homóloga a la isla de patogenicidad VpaI-7. Rev Chil infectología. 2019 Jun;36(3):392–5.

7.	Ministerio de Salud de Chile. Departamento de Epidemiología. Minuta: Situación epidemiológica de brote de diarrea aguda por Vibrio cholerae no toxigénico. Chile, año 2018-2019 [Internet]. 2019. Available from: Not publuished

8.	Chaand M, Miller KA, Sofia MK, Schlesener C, Weaver JWA, Sood V, et al. Type Three Secretion System Island-Encoded Proteins Required for Colonization by Non-O1/Non-O139 Serogroup Vibrio cholerae. Camilli A, editor. Infect Immun. 2015 Jul;83(7):2862–9.

9.	Almagro-Moreno S, Boyd EF. Sialic Acid Catabolism Confers a Competitive Advantage to Pathogenic Vibrio cholerae in the Mouse Intestine. Infect Immun. 2009 Sep 1;77(9):3807–16.

10.	Zheng J, Ho B, Mekalanos JJ. Genetic Analysis of Anti-Amoebae and Anti-Bacterial Activities of the Type VI Secretion System in Vibrio cholerae. Riggs PD, editor. PLoS One. 2011 Aug 31;6(8):e23876.

11.	Labbate M, Orata FD, Petty NK, Jayatilleke ND, King WL, Kirchberger PC, et al. A genomic island in Vibrio cholerae with VPI-1 site-specific recombination characteristics contains CRISPR-Cas and type VI secretion modules. Sci Rep. 2016 Dec 15;6(1):36891.

12.	Carraro N, Rivard N, Ceccarelli D, Colwell RR, Burrus V. IncA/C Conjugative Plasmids Mobilize a New Family of Multidrug Resistance Islands in Clinical Vibrio cholerae Non-

O1/Non-O139 Isolates from Haiti. MBio. 2016 Sep 7;7(4).

13.   Zhou Y, Yu L, Li J, Zhang L, Tong Y, Kan B. Accumulation of mutations in DNA gyrase
and topoisomerase IV genes contributes to fluoroquinolone resistance in Vibrio cholerae
O139 strains. Int J Antimicrob Agents. 2013 Jul;42(1):72–5.

14.   Fraga SG, De Trejo AV, Pichel M, Figueroa S, Merletti G, Caffer MI, et al.
Caracterización de aislamientos de Vibrio cholerae no-O1, no-O139 asociados a
cuadros de diarrea. Rev Argent Microbiol. 2009;41(1):11–9.

15.   Broberg CA, Calder TJ, Orth K. Vibrio parahaemolyticus cell biology and pathogenicity
determinants. Microbes Infect [Internet]. 2011;13(12–13):992–1001. Available from:
http://dx.doi.org/10.1016/j.micinf.2011.06.013

16.   Zhao Z, Liu J, Deng Y, Huang W, Ren C, Call DR, et al. The Vibrio alginolyticus T3SS
effectors, Val1686 and Val1680, induce cell rounding, apoptosis and lysis of fish
epithelial cells. Virulence. 2018 Dec 31;9(1):318–30.

A.

Type 3 Secretion System (T3SS)

nan-nag region

str. AM-19226

str. 10432-62

str. Santiago

- ■ tRNA-*ser* gene
- ■ Genes involved in DNA mobility
- ■ Transcriptional regulator genes
- ■ genes coding for T3SS-translocated proteins (Vops)
- ■ Genes coding for T3SS structural proteins
- ■ Genes coding for predicted virulence-related proteins (*acfs*) and thermostable direct hemolysin (*tdh*)
- ■ Genes coding for sialic acid catabolism
- ■ Core genome genes
- □ Genes coding for uncharacterized "hypothetical" proteins

identity 100% — 76%

B.

DR1-DR2

CRISPR-Cas region

Type VI secretion system region

DR1-DR2

str. HC36A1

str. Santiago

GI*Vch*S12 (28 kb)

- ■ tmRNA gene
- ■ Genes involved in DNA mobility
- □ Genes coding for uncharacterized "hypothetical" proteins
- ■ CRISPR-Cas genes
- ■ Type VI secretion system genes
- ■ Core genome genes

identity 100% — 66%

C.

str. Santiago

In36A1 ACSSuTmT

Mercury resistance transposon

GI*Vch*Hai6

- ■ Genes involved in DNA mobility
- ■ Genes coding for resistence determinants
- ■ Toxin-antitoxin gene
- ■ Transcriptional regulator genes
- ■ Genes coding for Type I restriction-modification enzymes
- ■ Core genome genes

identity 100% — 65%