Assembly of the *Drosophila* mushroom body circuit and its regulation by Semaphorin 1a

Chen-Han Lin[1,2] and Suewei Lin[1,2,#]

[1]Department of Life Sciences and Institute of Genome Sciences, National Yang-Ming University, Taipei 112, Taiwan
[2]Institute of Molecular Biology, Academia Sinica, Taipei 115, Taiwan

[#]Correspondence: sueweilin@gate.sinica.edu.tw

Summary

The *Drosophila* mushroom body (MB) is a learning and memory center in the fly brain. It is the most extensively studied brain structure in insects, but we know little about the molecular and cellular mechanisms underlying assembly of its neural circuit. The MB is composed of around 2200 intrinsic Kenyon cells (KCs), whose axons are bundled to form multiple MB lobes. The MB lobes are innervated by a large number of extrinsic neurons. Twenty types of dopaminergic neurons (DANs) and 21 types of MB output neurons (MBONs) have been identified. Each type of these extrinsic neurons innervates specific compartments or zones in the MB lobes. Here, we characterize the assembly of the MB circuit and reveal several intriguing features of the process. The DANs and MBONs innervate zones in the MB vertical lobes in specific sequential orders. Innervation of DAN axons in some zones precedes that of MBON dendrites, and *vice versa* in other zones. MBON and DAN innervations are largely independent of each other. Removing one type of extrinsic neuron during early development has a limited effect on the MB lobe innervations of the other type of extrinsic neurons. However, KC axons are essential for zonal elaboration of DAN axons and MBON dendrites. Competition also exists between MB zones for some MBONs, so when the cognate zones for these MBONs are missing, their dendrites are misdirected to other zones. Finally, we identify Semaphorin 1a (Sema1a) as a crucial guidance molecule for MBON dendrites to innervate specific MB lobe zones. Ectopic expression of Sema1a in some DANs is sufficient to re-direct their dendrites to those zones, demonstrating a potential to rewire the MB circuit. Taken together, our work provides an initial characterization of the cellular and molecular mechanisms underlying MB circuit assembly.

Introduction

During brain development, large numbers of neurons are precisely assembled in three-dimensional space to form functional networks that ultimately give rise to complex behaviors. The assembly process is regulated by intricate cellular and molecular mechanisms. Disruption of these mechanisms can lead to severe behavioral disorders or even death. Since Roger Sperry proposed his famous chemoaffinity hypothesis (Sperry, 1963), postulating that the precise wiring of the nervous system is established by specific chemical affinities generated from individual neurons, a large number of guidance molecules and principles governing neural circuit development has been discovered (Chédotal and Richards, 2010; Dickson and Zou, 2010; Hong and Luo, 2014; Kolodkin and Tessier-Lavigne, 2011; Sanes and Yamagata, 2009). However, our understanding of neural circuit assembly remains profoundly insufficient and we still cannot explain fully the mechanisms underpinning the wiring process of any neural circuit. Furthermore, given the remarkably diverse cell types and circuit configurations in the brain, additional guiding strategies and molecules surely await discovery.

The genetically tractable yet sufficiently complex nervous system of the fruit fly *Drosophila* has proven to be an excellent model for studying neural development. Many important principles of molecular guidance and neural circuit organization were first discovered in the fly (Evans, 2016; Hong and Luo, 2014; Kolodkin and Tessier-Lavigne, 2011; Plazaola-Sasieta et al., 2017; Tessier-Lavigne and Goodman, 1996). Most fly studies have focused on the ventral nerve cord, neuromuscular systems, antennal lobes, and optical lobes, but assembly of the central brain circuits has only begun to be explored (Oliva et al., 2016; Xie et al., 2017, 2019).

The *Drosophila* mushroom body (MB) is a multi-lobe structure at the center of the fly brain. It is a computational center mediating learning and memory (de Belle and Heisenberg, 1994; Heisenberg, 2003), sleep (Joiner et al., 2006; Pitman et al., 2006), and motivated behavior (Krashes et al., 2009; Senapati et al., 2019; Tsao et al., 2018). The MB circuit is composed of three major types of neurons—Kenyon cells (KCs), MB output neurons (MBONs), and dopaminergic neurons (DANs) (Aso et al., 2014). There are around 2200 KCs in each fly brain hemisphere. They are the intrinsic MB neurons whose axonal bundles form five MB lobes, with the α and α' lobes extending vertically and the β, β' and γ lobes projecting horizontally. The MB lobes are innervated by the dendrites of ~34 MBONs and the axons of ~130 DANs. Based on their morphology and molecular markers, 21 types of MBONs and 20 types of DANs have been identified. Each type of MBON or DAN extends its dendrites or axons, respectively, into specific compartments or zones of the MB lobes. Together, MBONs and DANs subdivide the MB lobes into 15 distinct zones. Most MBONs and DANs innervate only one or two zones. Given that the MB lobes are bundles of continuous axonal processes, the way in which each type of MBON and DAN synapses with specific domains of the MB lobes represents an elaborated form of subcellular targeting, through a process that is not yet well understood.

MB development has been studied extensively, but mainly focused on the KCs. KCs can be classified into three major types, i.e., γ, α'β', and αβ. They are sequentially produced from four neuroblasts during development (Lee et al., 1999). Several molecular mechanisms have been identified as governing the specification of KC cell fates (Liu et al., 2019, 2015; Marchetti and Tavosanis, 2017; Zhu et al., 2006). Targeting and pruning of KC axons have also been well characterized (Awasaki and Ito, 2004; Boulanger et al., 2011; Boyle et al., 2006; Lee et al., 2000; Lin et al., 2009; Ng et al., 2002; Rabinovich et al., 2016; Shimizu et al., 2011; Shin and DiAntonio, 2011; Smith-Trunova et al., 2015; Wang et al., 2002; Yu et al., 2013; Zwarts et al., 2016). In contrast, much less is known about the development of MBONs or DANs and how they are assembled into the MB circuit. To fill this knowledge gap, we first characterized the targeting of MBON dendrites and DAN axons to the MB vertical lobes during the pupal stage when the adult MB circuit is being assembled. We then used ablation experiments to probe the requirement of MBONs, DANs, and KCs for the assembly of the

87 MB circuit. Finally, we identified that *semaphorin 1a* (*sema1a*), a classical guidance molecule
88 (Kolodkin et al., 1993), is expressed in some MBONs, which is critical for their dendritic targeting to
89 specific MB lobe zones. Ectopic expression of *sema1a* in some DANs is sufficient to redirect their
90 dendrites to those zones. Our work provides the first characterization of the assembly of the MB
91 circuit and reveals a role for *sema1a* in the dendritic targeting of MBONs.
92

Results

The innervation timings of DAN axons and MOBN dendrites in the MB vertical lobes

To investigate how the MB circuits are assembled, we first endeavored to identify GAL4 lines in which individual types of DANs and MBONs are labeled throughout their morphogenesis. We focused on the MBONs and DANs that innervate the MB vertical lobes because they are fewer in type and number than those innervating the horizontal lobes (Aso et al., 2014). Among all of the GAL4 lines we assessed (**Table S1**), *MB058B-splitGAL4* and *MB091C-splitGAL4* specifically label PPL1-α'2α2 DANs and MBON-α'2, respectively, throughout their development (**Fig. 1A-L'**). Since PPL1-α'2α2 DANs and MBON-α'2 are the only DANs and MBONs that innervate the α'2 zone (Aso et al., 2014), we could study the innervation order of DAN axons and MBON dendrites in the same MB lobe zone by following the development of these neurons. We used *MB058B* and *MB091C* to drive the expression of mCD8::GFP in PPL1-α'2α2 DANs and MBON-α'2 and checked their morphologies at various times after puparium formation (APF). The PPL1-α'2α2 DANs begin to touch the MB lobes at 18 h APF, starting from the lateral surface of the α'2 zone (**Fig. 1A-A'**). At 24 h APF, innervation became more obvious, with axons extending ventrally along the interfaces between the α' and α lobes (**Fig. 1B-B' and M-M''**). Dense zonal innervation of the PPL1-α'2α2 axons was observed at 36 h APF (**Fig. 1D-D'**). Dendritic targeting of MBON-α'2 to the α'2 zone occurred at a later time-point. At 42 h APF, MBON-α'2 dendrites only began to touch the medial surface of the α'2 zone (**Fig. 1K-K'**), and dense zonal elaboration of the MBON-α'2 dendrites was only seen at 48 h APF (**Fig. 1L-L'**). Therefore, in the α'2 zone, DAN axons innervate the MB lobes earlier than MBON dendrites.

Interestingly, it seems that DAN axons do not always precede MBON dendrites in all MB lobe zones. For instance, in the α'3 zone, MBON-α'3 dendrites likely innervate the MB lobes earlier than the PPL1-α'3 axons. Innervation by MBON-α'3 dendrites (labeled with *MB027B-splitGAL4*) was observed at 6 h APF, i.e., much earlier than observed for neighboring MBON-α'2 (**Fig. S1**). However, when we labeled the DANs with antibodies against tyrosine hydroxylase (TH), an enzyme required for the synthesis of dopamine (Neckameyer and Quinn, 1989), we found that TH expression is sequentially turned on, beginning at the base of the vertical lobes and finishing at the tips of the α and α' lobes (**Fig. S2**). Expression of TH in the α'3 zone only became obvious at 54 h APF, representing the latest time-point among all vertical compartments. Since anti-TH signal was first seen in the α'2 and α2 zones at 24 h APF, TH expression appears to correlate with the innervations by DANs (**Fig. 1B-B' and Fig. S2**). These data suggest that MBON dendrites precede DAN axons in the α'3 zone.

DANs and MBONs independently innervate MB zones

Given that the axons of PPL1-α'2α2 DANs precede the MBON-α'2 dendrites in the α'2 zone, next we examined the possibility that DAN axons provide guidance cues for MBON dendrites. We genetically ablated PPL1-α'2α2 DANs by ectopically expressing the apoptotic genes *head involution defective* (*hid*) and *reaper* (*rpr*) (Grether et al., 1995; White et al., 1994) in these neurons using *MB058B-splitGAL4*. No *MB058B*-positive cells were seen in PPL1-α'2α2 DAN-ablated adult brains, and the α'2 and α2 MB lobe zones lacked anti-TH staining signal (**Fig. 2A-A' and B-B'**). Ablation of PPL1-α'2α2 DANs occurred quite early during development and all *MB058B*-positive neurites were eliminated before 24 h APF, i.e., by the time PPL1-α'2α2 DANs start to innervate MB lobes in wild-type brains (**Fig. S3**). TH-negative zones in the PPL1-α'2α2 DAN-ablated brains suggest that axon-axon contacts between neighboring DANs have a minimal effect in determining the borders of DAN axon zonal networks in MB lobes. Ablation of PPL1-α'2α2 DANs did not affect the dendritic targeting of MBON-α'2 (labeled with *R20G03-LexA::P65*-driven *rCD2-RFP*), suggesting that PPL1-α'2α2 DANs do not

provide guidance cues for MBON-α'2 dendrites (**Fig. 2A''-A''' and B''-B'''**). Ablation of PPL1-α'2α2
DANs also had no effect on the dendritic elaboration of MBON-α'3 (labeled with *R53F03-LexA::P65*-
driven *rCD2-RFP*) or MBON-γ2α'1 (labeled with *R25D01-LexA::P65*-driven *rCD2-RFP*) in the
neighboring zones (**Fig. S4**). Similarly, ablation of MBON-α'2 before its dendrites innervate the MB
lobes did not influence zonal elaboration of MBON-α'3 and MBON-γ2α'1 dendrites (**Figs. 2C-F'' and
S5**), indicating that dendritic tiling is not the mechanism by which the borders between MBON
dendrites in neighboring zones are determined. Therefore, zonal elaborations of DAN axons and
MBON dendrites are largely independent of each other.
152
Interestingly, when we ablated MBON-α'3, we found that the morphology of the α' lobe (labeled by
the KC-specific driver *MB247-LexA::p65* and *LexAop-rCD2::RFP*) was altered (**Fig. S3**) in that it
became shorter and the tip of the α' lobe (i.e. the α'3 zone) became smaller in adult brains lacking
MBON-α'3 (**Fig. S3A-E**). We followed the development of the MB lobes at different pupal stages in
MBON-α'3-ablated brains and found that growth of the α' lobe was slower than for wild-type
controls (**Figs. S7 and S3F**). This result suggests that the shorter α'3 tip was caused by a
developmental defect, not neurodegeneration. The α'3 zone did not completely disappear when
MBON-α'3 was ablated because when we labeled MBON-α'2 with *R20G03-LexA::P65*, its dendrites
remained clear of the reduced tip of the α' lobe (**Fig. S3G-H''**). These data suggest that the MBON-
α'3 dendrites might provide a positive signal to promote growth of the KC axons. Alternatively,
innervation by the MBON-α'3 dendrites might simply inflate the α'3 zone in wild-type brains.
However, this latter explanation seems less likely because we did not observe a similar reduction of
the α'2 zone when MBON-α'2 dendrites were ablated (**Fig. S3I**).
166
167
168 **KC axons are essential for zonal elaborations of DAN axons**
169
To investigate the role of KC axons in axonal and dendritic targeting of DANs and MBONs,
respectively, we took advantage of the *alpha lobe absent* (*ala*) fly mutant. Brains of the *ala* mutant
have been shown to randomly lose vertical or horizontal lobes of the MB (Pascual and Préat, 2001;
Pascual et al., 2005). We found that in *ala* mutants that lose MB vertical lobes on one side of the
brain, both contralateral and ipsilateral PPL1-α'2α2 axons wandered around the areas near their
missing targets without forming normal zonal elaboration patterns (**Fig. 3A-B''**). Mistargeting of
PPL1-α'2α2 axons was completely correlated with the lobe-missing phenotype (n=9), so it was not
caused by loss of the *ala* gene *per se* but was a consequence of the missing lobe. We also examined
PPL1-α'2α2 axons in the *ala* mutant brains at 30 h APF, when obvious zonal innervation by the axons
is first observed (**Fig. 1C-C'**). We found that when vertical lobes were absent, PPL1-α'2α2 axons
failed to form a zonal network from the very beginning of the elaboration process, and some
mistargeted neurites could be found slightly above and posterior to where the α'2 and α2 zones are
supposed to be located (**Fig. 3C-D'**). Since these mistargeted axons could still project to sites close to
their normal targets (**Fig. 3**), KC axons do not appear to provide long-range guidance cues for these
axons. Instead, the KC axons may provide local positional cues for the DAN axons to form zonal
networks. This latter scenario was further evidenced when we used a multi-color flip-out technique
to label PPL1-α'2α2 and PPL1-α'3 DANs in different colors. We found that the primary neurites of
these two DANs entered the MB lobes at a similar location (**Fig. S8**). However, PPL1-α'2α2 axons
exclusively occupied the α'2 and α2 zones below the entry point, whereas PPL1-α'3 axons specifically
covered the α'3 zone above the entry point (**Fig. S8**), suggesting local cues exist that direct zonal
elaboration of DAN axons.
191
192
Competition between MB lobe zones for MBON-α'2 dendrites

Surprisingly, when the MB vertical lobes were missing, MBON-α'2 dendrites targeted the β'2mp and
α'1 zones instead (**Fig. 4A-B''**). A previous study showed that in *ala* mutants lacking vertical MB lobes,
approximately half of the KCs lose their vertical branches and the vertical branches of the remaining
half project horizontally (Pascual et al., 2005). However, the mistargeting of MBON-α'2 we report
here is unlikely caused by the horizontally-projecting KC vertical branches for two reasons. First, the
zones innervated by the mistargeted MBON-α'2 dendrites are located at either end, but not the
middle, of the β' lobe. Second, the guidance molecules required for MBON-α'2 to innervate the α'2
zone are different from those required for it to innervate the β'2 and α'1 zones (see below). When
we followed the development of the MBON-α'2 dendrites in the absence of their normal target, we
found that they never formed a zonal network near the missing target, but started to innervate the
β'2mp and α'1 zones at around 72 h APF (**Fig. 4C-E'**). Taken together, these results indicate that
there are competing guidance cues from multiple MB lobe zones for the MBON-α'2 dendrites. When
the stronger guidance cues from the α'2 zones are missing, the MBON-α'2 dendrites are attracted to
other zones.
208
209
210   **Overexpression of *sema1a* in DANs mistargets their dendrites to specific MB lobe zones**
211
During the course of identifying genes that control MB circuit assembly, we found that when *sema1a*
was overexpressed in PPL1-α'2α2 DANs, their dendrites that normally elaborate loosely behind the
MB vertical lobes (**Fig. 5A-A''**) instead innervated the β'2p, α'1, and α'3 zones (**Fig. 5B-B''**). We
confirmed that these misguided neurites are dendrites by labeling PPL1-α'2α2 with DenMark and
dSyd1-GFP (**Fig. 5E-E''**). The dSyd-1-GFP-labeled axons were not affected by overexpression of
sema1a. Sema1a can function as a ligand or receptor in different developmental contexts (Cafferty
et al., 2006; Godenschwege et al., 2002; Komiyama et al., 2007; Winberg et al., 1998). We found that
overexpression in PPL1-α'2α2 DANs of *sema1a* lacking its intracellular domain (Sema1a-mEC-5xmyc
and Sema1a-mEC.Fc-5xmyc) (Jeong et al., 2012) had no effect on their dendrites (**Fig. 5C-D''**),
suggesting that when overexpressed, Sema1a works as a receptor to guide PPL1-α'2α2 dendrites to
the MB lobe zones where the ligands for Sema1a are likely produced. Overexpression of *sema1a* in
PPL1-γ2α'1 DANs using *MB296B-splitGAL4* also caused mistargeted dendrites, but in this instance
they innervated the β'2mp and β'2a zones (**Fig.5F-I''**). It is striking that overexpression of *sema1a* in
different DANs targets their dendrites to distinct β'2 zones, suggesting that Sema1a may work with
other guidance molecules to determine zonal specificity for different dendrites.
227
Overexpression of *sema1a* cannot always outcompete the endogenous dendritic targeting program,
as is evident for MBON-α'2 in which overexpression of *sema1a* did not lead to any dendritic
targeting phenotype (**Fig. S9**). However, ectopic innervation by *sema1a*-overexpressed PPL1-α'2α2
and PPL1-γ2α'1 dendrites in the β'2mp and α'1 zones is reminiscent of the mistargeting of MBON-
α'2 dendrites in the absence of MB vertical lobes (**Fig. 4**). Therefore, we tested if *sema1a* is required
for the mistargeting of MBON-α'2 dendrites in *ala* mutant brains. We knocked down *sema1a* from
MBON-α'2 in *ala* mutant flies and found that the MBON-α'2 dendrites with reduced *sema1a*
expression no longer innervated the β'2mp and α'1 zones when their normal target was missing (**Fig.
6**). However, knockdown of *sema1a* did not affect MBON-α'2 dendritic targeting to the α'2 zone in
the normal vertical lobes (**Fig. 6**). These results suggest that MBON-α'2 may express *sema1a* as well
as a yet unidentified guidance molecule. In wild-type brains, this unknown guidance molecule
outcompetes *sema1a* to target MBON-α'2 dendrites to the α'2 zone. However, when the α'2 zone is
missing, a *sema1a*-depedent guidance mechanism takes precedence, directing the MBON-α'2
dendrites to the β'2mp and α'1 zones.
242
243
244   ***Sema1a* is required for dendritic targeting of several MBONs**

245
The ectopic innervation of *sema1a*-overexpressed PPL1-α'2α2 and PPL1-γ2α'1 dendrites in the β'2,
α'1, and α'3 zones prompted us to examine the role of *sema1a* in dendritic targeting of the MBONs
innervating these three zones. We found that RNAi knockdown of *sema1a* in MBON-β2β'2a using
MB399C-splitGAL4 resulted in mistargeting of the dendrites that normally innervate the β'2a zone
(**Fig. 7A-D**). Instead of innervating the β'2a zone, these dendrites projected upwards to regions near
the central complex. The MBON-β2β'2a dendrites innervating the β2 zone were minimally affected
by *sema1a* knockdown (**Fig. 7A-D**), suggesting that *sema1a* regulates dendritic targeting of MBON-
β2β'2a in a zone-specific manner. Similarly, when *sema1a* was knocked down in MBON-α'3 using
MB027B-splitGAL4, the MBON-α'3 dendrites failed to concentrate in the α'3 zone but elaborated
loosely around the α'3 zone (**Fig. 7E-H**). We confirmed these findings by creating *sema1a* MARCM
(mosaic analysis with a repressible cell marker; (Lee and Luo, 1999)) clones for MBON-β2β'2a (**Fig.**
8A-B') and MBON-α'3 (**Fig. 8C-D'**). The dendritic targeting defects observed in these *sema1a*-mutant
MBON-β2β'2a and MBON-α'3 were consistent with the phenotypes caused by *sema1a* RNAi
knockdown. To confirm that MBON-β2β'2a and MBON-α'3 express *sema1a*, we used *sema1a*
Artificial exon that produces V5-tagged Sema1a and LexA::p65 under the control of the endogenous
sema1a enhancer when an FRT-stop-FRT cassette in the exon is removed by Flipase (FLP) (Pecot et
al., 2013). Consistently, V5-tagged Sema1a and LexA::P65 driven by the *sema1a* enhancer were
detected in MBON-β2β'2a and MBON-α'3 when *MB399C-splitGAL4* and *MB027B-splitGAL4* were
used, respectively, to drive *UAS-FLP* (**Fig. S10**).
265
MARCM also allowed us to test the role of *sema1a* in MBONs for which we do not have GAL4 lines to
drive RNAi sufficiently early during their development. Apart from innervation by MBON-β2β'2a
dendrites, the β'2 zone is also innervated by MBON-β'2mp and MBON-γ5β'2a dendrites, and the α'1
zone is further innervated by MBON-γ2α'1 and MBON-α'1 dendrites. We found that dendritic
targeting of MBON- γ2α'1 and MBON-γ5β'2a remained normal in *sema1a* MARCM clones (**Fig. S11**).
However, *sema1a* MARCM clones for MBON-α'1 exhibited a severe dendritic targeting defect (**Fig.**
8E-F'). We induced clones at 4-8 h after egg laying (AEL) and labeled the MBON-α'1 dendrites using
R24H08-GAL4, which also labels MBON-α'3 dendrites. Interestingly, all of the MBON-α'1 clones were
accompanied by MBON-α'3 clones, suggesting that these two neuron types are from the same
neuronal lineage. We also observed a dendritic targeting defect in *sema1a* MARCM clones for
MBON-β'2mp (**Fig. 8G-H'**). Taken together, these results indicate that *sema1a* is required for
dendritic targeting in most MBONs that innervate the β'2, α'1, and α'3 zones.
278
279
280 **Canonical partners of Sema1a are not essential for the targeting of MBON dendrites**
281
Sema2a, Sema2b, and PlexA have been identified as ligands for Sema1a (Sweeney et al., 2011; Yu et
al., 2010). To test if these molecules play a role in the targeting of MBON dendrites, we first
examined the dendritic innervation patterns of MBON-β2β'2a and MBON-α'3 in flies carrying
homozygous mutations in both *sema2a* and *sema2b* genes (*Sema2b^{f02042}, sema2a^{p2}*) (Sweeney et al.,
2011). Although MB lobe structure is slightly abnormal in these flies, morphology of MBON-β2β'2a
and MBON-α'3 dendrites and their zonal innervations were essentially normal (**Fig. S12A-B'**).
Therefore, Sema2a and Sema2b are unlikely to be the guidance molecules that direct *sema1a*-
positive MBON dendrites to specific MB lobe zones. Flies lacking PlexA activity fail to survive to
adulthood. To examine the potential role of PlexA in guiding the MBON dendrites, we took
advantage of a weaker *sema1a-RNAi-2* line whose overexpression resulted in mild targeting defects
of MBON-α'3 and MBON-β2β'2a dendrites (**Fig. S12C-C' and E-E'**) and then checked if loss of one
plexA allele exacerbated the defects. However, the targeting defects of MBON-β2β'2a and MBON-
α'3 dendrites were not exacerbated in flies heterozygous for a *plexA* mutant allele, *plexA^{EY16548}* (**Fig.**
S12C-F'). We also expressed an *UAS-plexA-RNAi* construct that has been shown to knock down *plexA*

effectively in all neurons (*nSyb-GAL4*) or in all glia (*repo-GAL4*) (Pecot et al., 2013). We then
examined the innervation by MBON-β2β'2a and MBON-β'2mp (*R12C11-LexA*-driven rCD2-RFP) and
MBON-α'3 (*R53F03-LexA*-driven rCD2-RFP) dendrites in the MB lobes. However, we did not observe
any dendritic targeting phenotype in all cases (**Fig. S12G-N'**). Therefore, there is no evidence that the
three known Sema1a ligands are involved in guiding MBON dendrites during assembly of the MB
circuit.
302

Discussion

The MB is the most studied structure in the fly brain. It is conventionally considered as an olfactory learning and memory center (Aso et al., 2014; de Belle and Heisenberg, 1994; Heisenberg, 2003). Recently, it has also been shown to receive other sensory inputs and to mediate motivational control of innate behavior (Krashes et al., 2009; Masek and Scott, 2010; Masek et al., 2015; Senapati et al., 2019; Tsao et al., 2018; Vogt et al., 2014, 2016). Almost all the neurons that form the MB circuit have been identified, and genetic means of labeling individual neuronal types have been developed (Aso et al., 2014). However, our understanding of the assembly process of this important structure is essentially lacking. In this study, we provide an initial characterization of this process and reveal several interesting principles.

The most intriguing feature of the organization of the MB circuit is the zonal innervation of the MB lobes by DAN axons and MBON dendrites. Electron microscopy reconstruction of the MB lobes shows that in each zone the DAN axons and MBON dendrites form synapses with each other and the KC axons (Takemura et al., 2017). The borders of the zones are clear, and there is minimal overlap between DAN axons or MBON dendrites in the neighboring zones. With such a highly organized neural network, elaborate interactions among the extrinsic neurons might be expected. For example, dendritic tiling, as observed between dendritic arborization (da) neurons in fly larvae, might be required for the formation of zonal borders (Grueber et al., 2002), and matching between DAN axons and MBON dendrites in the same zone might be important for these neurites to establish proper zonal innervation patterns. However, our results suggest that the targeting and elaboration networks of DAN axons and MBON dendrites are largely independent. In the α'2 zone, where innervation by DAN axons precedes that by MBON dendrites, ablation of DANs does not affect zonal elaboration of the MBON dendrites. Moreover, upon ablation of one type of DAN or MBON, morphologies of the neighboring neurites appear to be normal. Therefore, the extent and location of zonal network elaboration by DAN axons and MBON dendrites might be primarily determined by the interactions between KC axons and these extrinsic neurons.

The KC axons are apparently essential for zonal arborization of the DAN axons and MBON dendrites. When their target lobes are absent, DAN axons wander around the missing target and generate less extensive neurite arbors. Therefore, the KC axons appear to promote growth and arborization of DAN axons. Do the KC axons also determine the extent and borders of the zonal arbors? It is still possible that the KC axons simply provide an anchoring point on which DAN axons and MBON dendrites grow, and that the extent of their arborization is determined cell-autonomously as an intrinsic property. However, the axonal innervation pattern of PPL1 DANs argues against this latter possibility. The axonal projections of PPL1-α'3 and PPL1-α'2α2 DANs enter the MB vertical lobes at almost the same location, but specifically innervate distinct zones on opposite sides of the entry point. Therefore, different segments in the KC axons might provide positional cues for the DANs and MBONs, although direct evidence for this possibility awaits investigation. If they exist, what could be the nature of these positional cues? One possibility is that the KC axonal segment in each zone expresses a unique guidance molecule that binds to a receptor on the neurites of the DANs and MBONs projecting to that segment. However, when the target zone of MBON-α'2 is lost, its dendrites instead innervate the β'2mp and α'1 zones. This result suggests that an MBON can respond to multiple positional cues. Furthermore, since *sema1a* is required for MBONs to innervate multiple specific MB lobe zones, the positional cues are likely combinatorial.

Sema1a can function as both a ligand and a receptor (Cafferty et al., 2006; Godenschwege et al., 2002; Komiyama et al., 2007; Winberg et al., 1998). Using *sema1a* transgenes that lack the intracellular domain, we demonstrated that it works as a receptor in MBONs to guide their dendrites. PlexA and Sema2a/2b are known ligands for Sema1a (Sweeney et al., 2011; Yu et al., 2010). However,

dendritic targeting by MBONs was minimally affected in flies homozygous for *sema2a/2b* double
mutants or when we knocked down PlexA either pan-neuronally or in glia. Moreover, loss of one
copy of *plexA* did not exacerbate the dendritic targeting defect caused by RNAi knockdown of
sema1a in MBONs. Therefore, these canonical Sema1a ligands do not seem to play a role in guiding
MBON dendrite targeting. Thus, the molecular mechanisms that direct *sema1a*-positive MBON
dendrites to specific MB lobe zones remain unknown.

Experimental procedures

Fly strains and husbandry

Fly lines used in this study are: (1) *MB077C-splitGAL4* (Bloomington Drosophila Stock Center, BDSC:68284) (Aso et al., 2014); (2) *MB091C-splitGAL4* (gift from Y. Aso) (Aso et al., 2014). *MB027B-splitGAL4* (BDSC:68301) (Aso et al., 2014); (3) *MB058B-splitGAL4* (BDSC:68278) (Aso et al., 2014); (4) *MB296B-splitGAL4* (BDSC:68308) (Aso et al., 2014); (5) *MB399C-splitGAL4* (gift from Y. Aso) (Aso et al., 2014); (6) *R53F03-GAL4* (BDSC:48197) (Jenett et al., 2012); (7) *R21D02-GAL4* (BDSC:48939) (Jenett et al., 2012); (8) *R24H08-GAL4* (BDSC:49100) (Jenett et al., 2012); (9) *R25D01-GAL4* (BDSC:49122) (Jenett et al., 2012); (10) *repo-GAL4* (gift from Y-H. Sun); (11) *VT41043-GAL4* (gift from C-L. Wu); (12) *nSyb-GAL4* (Lin et al., 2012); (13) *UAS-DenMark,UAS-Dsyd-1::GFP* (Owald et al., 2015); (14) *UAS-mCD8::GFP*(LL4) (Lee and Luo, 1999); (15) *UAS-mCD8::GFP*(II) (Lee and Luo, 1999); (16) *UAS-mCD8::GFP*(III) (Lee and Luo, 1999); (17) *UAS-(FRT stop)myr-SmGdp-HA, UAS-(FRT stop)myr-SmGdp-V5, UAS-(FRT stop)myr-SmGdp-FLAG* (BDSC:64085) (Nern et al., 2015); (18) *MB247-LexA::p65, LexAop-rCD2::RFP* (Awasaki et al., 2011); (19) *LexAop-rCD2::RFP*(II) (Awasaki et al., 2011); (20) *UAS-hid, UAS-rpr* (gift from S. Waddell); (21) UAS-Dicer (Vienna Drosophila RNAi Center, VDRC:36148); (22) *20XUAS-FLPG5* (BDSC:55807) (Nern et al., 2011); (23) *R53F03-LexA::p65* (BDSC:53502) (Pfeiffer et al., 2010); (24) *R20G03-LexA::p65* (BDSC:52570) (Pfeiffer et al., 2010); (25) *R25D01-LexA::p65* (BDSC:53519) (Pfeiffer et al., 2010); (26) *R12C11-LexA::p65* (BDSC:52444) (Pfeiffer et al., 2010); (27) *hs-FLPG5* (BDSC:77140) (Nern et al., 2011); (28) *UAS-sema1a* (BDSC:65734) (Jeong et al., 2012); (29) *UAS-sema1.mEC-5xmyc* (BDSC:65739) (Jeong et al., 2012); (30) *UAS-sema1.mEC.Fc-5xmyc* (BDSC:65740) (Jeong et al., 2012); (31) *UAS-sema1a-RNAi-1* (BDSC:34320) (Perkins et al., 2015); (32) *UAS-sema1a-RNAi-2* (VDRC:36148) (Dietzl et al., 2007); (33) *UAS-plexA-RNAi* (BDSC:30483) (Pecot et al., 2013; Perkins et al., 2015); (34) *w, ala^{E13}* (BDSC:9463) (Pascual and Préat, 2001); (35) *plexA^{EY16548}* (BDSC:23097) (Bellen et al., 2011; Shin and DiAntonio, 2011); (36) *40A, sema1a^{k13702}* (Kyoto Stock Center:111328) (Yu et al., 1998); (37) *Sema2b^{f02042}, sema2a^{p2}, UAS-mCD8::GFP/CyO* (gift from L. Luo) (Sweeney et al., 2011); (38) *hs-FLP122, 40A, tubp-GAL80* (gift from H-H. Yu); and (39) *w; sema1a Artificial exon/CyO; LexAop-tdTomato* (gift from L. Zipursky) (Pecot et al., 2013). The genotypes of the flies used in each figure are shown in **Table S2**. All flies were reared at 23 °C with 60% relative humidity, under a 12 h light: 12 h dark cycle. All our data were derived from male flies.

Mosaic analysis with a repressible marker (MARCM) and Flip-out

Flies with proper genotypes were heat-shocked at 37 °C at different developmental stages for different durations (see **Table S2** for each of these parameters used in each figure). For flip-out clones induced at the adult stage, we waited for at least 3 days after heat-shock to ensure proper expression of the marker proteins. Brains were then dissected before performing immunofluorescence staining.

Immunofluorescence staining

Fly brains were dissected in phosphate buffered saline (PBS; Sigma, P4417) and fixed in PBS containing 4% formaldehyde (Sigma, F8775) at room temperature for 20 min. The brains were then washed three times with PBS containing 0.5% Triton X-100 (PBST, Sigma-Aldrich, T8787); each time, the brains were incubated in PBST at room temperature for at least 20 min. After that, the brains were incubated in PBST containing 5% normal goat serum (blocking solution; normal goat serum, Jackson ImmunoResearch, 005-000-121) for 30 min at room temperature. After blocking, the brains were incubated overnight in blocking solution containing primary antibodies at 4 °C. The next day, the brains were washed three times, for 20 min each time, with PBST at room temperature before they were incubated overnight in blocking solution containing secondary antibodies at 4 °C. The next

day, the brains were washed again three times, for 20 min each time, with PBST at room
temperature. Then the brains were mounted with Gold Antifade reagent (Thermo Fisher Scientific,
S36937). Antibodies used in each figure are listed in **Table S2**. In some cases, we performed
sequential staining in which the brains were stained with the first set of primary and secondary
antibodies before being incubated with the second set of primary and secondary antibodies. The
staining procedures (single vs. sequential) for each figure are provided in **Table S2**. The primary
antibodies used in this study are Rat anti-mCD8 (1:100; Invitrogen, MCD0800), Chicken anti-GFP
(1:1000; Invitrogen, A10262), Rabbit anti-Dsred (1:100; Takara, 632496), Mouse anti-Trio (1:100;
Developmental Studies Hybridoma Bank, DSHB, 9.4A), Mouse anti-FasII (1:100; DSHB, 1D4), Mouse
anti-DLG (1:100; DSHB, 4F3), Mouse anti-TH (1:1000; Immunostar, 22941), Rat anti-FLAG (1:200;
Novus Biologicals, NBP1-06721), Mouse anti-V5 (1:500; Bio-RAD, MCA1360), and Rabbit anti-V5
(1:500; Abcam, AB9116). The secondary antibodies used in this study are Goat anti-rat-488 (1:400;
Jackson ImmunoResearch, 112-545-167), Goat anti-rabbit-Cy3 (1:400; Jackson ImmunoResearch,
111-165-144), Donkey anti-chicken-488 (1:400; Jackson ImmunoResearch, 703-545-155), Goat anti-
mouse-Cy5 (1:400; Jackson ImmunoResearch, 115-175-166), Goat anti-mouse-Cy3 (1:400; Jackson
ImmunoResearch, 115-165-166), Goat anti-mouse-488 (1:400; Invitrogen, A11029), and Goat anti-
rabbit-Cy5 (1:400; Jackson ImmunoResearch, 111-175-144). The stained brains were imaged using a
confocal microscope (Zeiss LSM880) and analyzed using Fiji/ImageJ(Schindelin et al., 2012) or Imaris
x64 (v9.2.1, Oxford Instruments).
431
432 **Statistics**
433
434 Statistical analysis was performed in Prism 8. All data were tested for normality using the Shapiro-
435 Wilk normality test. No data points were excluded from the analysis. Since all our data are normally
436 distributed, we used two-tailed unpaired Student's *t*-test with Welch's correction.
437
438

Author contributions

C-H.L. and S.L. conceived the project and designed the experiments. C-H.L. performed the experiments and analyzed the data. S.L. wrote the paper and directed the research.

Acknowledgments

We thank Y. Aso (Janelia Farm Research Campus, USA), G. Rubin (Janelia Farm Research Campus, USA), T. Lee (Janelia Farm Research Campus, USA), L. Luo (Stanford University, USA), Y-H. Sun (Academia Sinica, Taiwan), S. Waddell (University of Oxford, UK), H-H. Yu (Academia Sinica, Taiwan), C-L. Wu (Chang Gung University, Taiwan), L. Zipurski (University of California, Los Angeles, USA), and FlyLight (Janelia Farm Research Campus, USA), the Bloomington Drosophila Stock Center, Vienna Drosophila RNAi Center, Harvard TRiP RNAi Stock Center, Kyoto Stock Center, and Taiwan Fly Core for fly stocks. S. L. is funded by the Ministry of Science and Technology, Taiwan (107-2311-B-001-042-MY3) and Academia Sinica, Taiwan.

References

Aso, Y., Hattori, D., Yu, Y., Johnston, R.M., Iyer, N.A., Ngo, T.-T.B., Dionne, H., Abbott, L.F., Axel, R., Tanimoto, H., et al. (2014). The neuronal architecture of the mushroom body provides a logic for associative learning. Elife *3*, e04577.

Awasaki, T., and Ito, K. (2004). Engulfing action of glial cells is required for programmed axon pruning during Drosophila metamorphosis. Curr. Biol. *14*, 668–677.

Awasaki, T., Huang, Y., O'Connor, M.B., and Lee, T. (2011). Glia instruct developmental neuronal remodeling through TGF-β signaling. Nat. Neurosci. *14*, 821–823.

de Belle, J.S., and Heisenberg, M. (1994). Associative odor learning in Drosophila abolished by chemical ablation of mushroom bodies. Science *263*, 692–695.

Bellen, H.J., Levis, R.W., He, Y., Carlson, J.W., Evans-Holm, M., Bae, E., Kim, J., Metaxakis, A., Savakis, C., Schulze, K.L., et al. (2011). The Drosophila gene disruption project: progress using transposons with distinctive site specificities. Genetics *188*, 731–743.

Boulanger, A., Clouet-Redt, C., Farge, M., Flandre, A., Guignard, T., Fernando, C., Juge, F., and Dura, J.-M. (2011). ftz-f1 and Hr39 opposing roles on EcR expression during Drosophila mushroom body neuron remodeling. Nat. Neurosci. *14*, 37–44.

Boyle, M., Nighorn, A., and Thomas, J.B. (2006). Drosophila Eph receptor guides specific axon branches of mushroom body neurons. Development *133*, 1845–1854.

Cafferty, P., Yu, L., Long, H., and Rao, Y. (2006). Semaphorin-1a functions as a guidance receptor in the Drosophila visual system. J. Neurosci. *26*, 3999–4003.

Chédotal, A., and Richards, L.J. (2010). Wiring the brain: the biology of neuronal guidance. Cold Spring Harb. Perspect. Biol. *2*, 1–18.

Dickson, B.J., and Zou, Y. (2010). Navigating intermediate targets: the nervous system midline. Cold Spring Harb. Perspect. Biol. *2*, a002055.

Dietzl, G., Chen, D., Schnorrer, F., Su, K.-C., Barinova, Y., Fellner, M., Gasser, B., Kinsey, K., Oppel, S., Scheiblauer, S., et al. (2007). A genome-wide transgenic RNAi library for conditional gene inactivation in Drosophila. Nature *448*, 151–156.

Evans, T.A. (2016). Embryonic axon guidance: insights from Drosophila and other insects. Curr. Opin. Insect Sci. *18*, 11–16.

Godenschwege, T.A., Hu, H., Shan-Crofts, X., Goodman, C.S., and Murphey, R.K. (2002). Bi-directional signaling by Semaphorin 1a during central synapse formation in Drosophila. Nat. Neurosci. *5*, 1294–1301.

Grether, M.E., Abrams, J.M., Agapite, J., White, K., and Steller, H. (1995). The head involution defective gene of Drosophila melanogaster functions in programmed cell death. Genes Dev. *9*, 1694–1708.

Grueber, W.B., Jan, L.Y., and Jan, Y.N. (2002). Tiling of the Drosophila epidermis by multidendritic sensory neurons. Development *129*, 2867–2878.

Heisenberg, M. (2003). Mushroom body memoir: from maps to models. Nat. Rev. Neurosci. *4*, 266–275.

Hong, W., and Luo, L. (2014). Genetic control of wiring specificity in the fly olfactory system. Genetics *196*, 17–29.

Jenett, A., Rubin, G.M., Ngo, T.-T.B., Shepherd, D., Murphy, C., Dionne, H., Pfeiffer, B.D., Cavallaro, A., Hall, D., Jeter, J., et al. (2012). A GAL4-driver line resource for Drosophila neurobiology. Cell Rep. *2*, 991–1001.

Jeong, S., Juhaszova, K., and Kolodkin, A.L. (2012). The Control of semaphorin-1a-mediated reverse signaling by opposing pebble and RhoGAPp190 functions in drosophila. Neuron *76*, 721–734.

Joiner, W.J., Crocker, A., White, B.H., and Sehgal, A. (2006). Sleep in Drosophila is regulated by adult mushroom bodies. Nature *441*, 757–760.

Kolodkin, A.L., and Tessier-Lavigne, M. (2011). Mechanisms and molecules of neuronal wiring: A primer. Cold Spring Harb. Perspect. Biol. *3*, 1–14.

Kolodkin, A.L., Matthes, D.J., and Goodman, C.S. (1993). The semaphorin genes encode a family of
transmembrane and secreted growth cone guidance molecules. Cell *75*, 1389–1399.
Komiyama, T., Sweeney, L.B., Schuldiner, O., Garcia, K.C., and Luo, L. (2007). Graded expression of
semaphorin-1a cell-autonomously directs dendritic targeting of olfactory projection neurons. Cell
128, 399–410.
Krashes, M.J., DasGupta, S., Vreede, A., White, B., Armstrong, J.D., and Waddell, S. (2009). A neural
circuit mechanism integrating motivational state with memory expression in Drosophila. Cell *139*,
416–427.
Lee, T., and Luo, L. (1999). Mosaic analysis with a repressible cell marker for studies of gene function
in neuronal morphogenesis. Neuron *22*, 451–461.
Lee, T., Lee, A., and Luo, L. (1999). Development of the Drosophila mushroom bodies: sequential
generation of three distinct types of neurons from a neuroblast. Development *126*, 4065–4076.
Lee, T., Marticke, S., Sung, C., Robinow, S., and Luo, L. (2000). Cell-autonomous requirement of the
USP/EcR-B ecdysone receptor for mushroom body neuronal remodeling in Drosophila. Neuron *28*,
807–818.
Lin, S., Huang, Y., and Lee, T. (2009). Nuclear receptor unfulfilled regulates axonal guidance and cell
identity of Drosophila mushroom body neurons. PLoS One *4*, e8392.
Lin, S., Kao, C.-F., Yu, H.-H., Huang, Y., and Lee, T. (2012). Lineage analysis of Drosophila lateral
antennal lobe neurons reveals notch-dependent binary temporal fate decisions. PLoS Biol. *10*,
e1001425.
Liu, L.-Y., Long, X., Yang, C.-P., Miyares, R.L., Sugino, K., Singer, R.H., and Lee, T. (2019). Mamo
decodes hierarchical temporal gradients into terminal neuronal fate. Elife *8*, 1–28.
Liu, Z., Yang, C.-P., Sugino, K., Fu, C.-C., Liu, L.-Y., Yao, X., Lee, L.P., and Lee, T. (2015). Opposing
intrinsic temporal gradients guide neural stem cell production of varied neuronal fates. Science *350*,
317–320.
Marchetti, G., and Tavosanis, G. (2017). Steroid Hormone Ecdysone Signaling Specifies Mushroom
Body Neuron Sequential Fate via Chinmo. Curr. Biol. *27*, 3017-3024.e4.
Masek, P., and Scott, K. (2010). Limited taste discrimination in Drosophila. Proc. Natl. Acad. Sci. U. S.
A. *107*, 14833–14838.
Masek, P., Worden, K., Aso, Y., Rubin, G.M., and Keene, A.C. (2015). A dopamine-modulated neural
circuit regulating aversive taste memory in Drosophila. Curr. Biol. *25*, 1535–1541.
Neckameyer, W.S., and Quinn, W.G. (1989). Isolation and characterization of the gene for Drosophila
tyrosine hydroxylase. Neuron *2*, 1167–1175.
Nern, A., Pfeiffer, B.D., Svoboda, K., and Rubin, G.M. (2011). Multiple new site-specific recombinases
for use in manipulating animal genomes. Proc. Natl. Acad. Sci. *108*, 14198–14203.
Nern, A., Pfeiffer, B.D., and Rubin, G.M. (2015). Optimized tools for multicolor stochastic labeling
reveal diverse stereotyped cell arrangements in the fly visual system. Proc. Natl. Acad. Sci. U. S. A.
112, E2967–E2976.
Ng, J., Nardine, T., Harms, M., Tzu, J., Goldstein, A., Sun, Y., Dietzl, G., Dickson, B.J., and Luo, L. (2002).
Rac GTPases control axon growth, guidance and branching. Nature *416*, 442–447.
Oliva, C., Soldano, A., Mora, N., De Geest, N., Claeys, A., Erfurth, M.-L., Sierralta, J., Ramaekers, A.,
Dascenco, D., Ejsmont, R.K., et al. (2016). Regulation of Drosophila Brain Wiring by Neuropil
Interactions via a Slit-Robo-RPTP Signaling Complex. Dev. Cell *39*, 267–278.
Owald, D., Felsenberg, J., Talbot, C.B., Das, G., Perisse, E., Huetteroth, W., and Waddell, S. (2015).
Activity of defined mushroom body output neurons underlies learned olfactory behavior in
Drosophila. Neuron *86*, 417–427.
Pascual, A., and Préat, T. (2001). Localization of long-term memory within the Drosophila mushroom
body. Science *294*, 1115–1117.
Pascual, A., Chaminade, M., and Préat, T. (2005). Ethanolamine kinase controls neuroblast divisions
in Drosophila mushroom bodies. Dev. Biol. *280*, 177–186.
Pecot, M.Y., Tadros, W., Nern, A., Bader, M., Chen, Y., and Zipursky, S.L. (2013). Multiple interactions

control synaptic layer specificity in the Drosophila visual system. Neuron *77*, 299–310.

Perkins, L.A., Holderbaum, L., Tao, R., Hu, Y., Sopko, R., McCall, K., Yang-Zhou, D., Flockhart, I., Binari, R., Shim, H.-S., et al. (2015). The Transgenic RNAi Project at Harvard Medical School: Resources and Validation. Genetics *201*, 843–852.

Pfeiffer, B.D., Ngo, T.T.B., Hibbard, K.L., Murphy, C., Jenett, A., Truman, J.W., and Rubin, G.M. (2010). Refinement of tools for targeted gene expression in Drosophila. Genetics *186*, 735–755.

Pitman, J.L., McGill, J.J., Keegan, K.P., and Allada, R. (2006). A dynamic role for the mushroom bodies in promoting sleep in Drosophila. Nature *441*, 753–756.

Plazaola-Sasieta, H., Fernández-Pineda, A., Zhu, Q., and Morey, M. (2017). Untangling the wiring of the Drosophila visual system: developmental principles and molecular strategies. J. Neurogenet. *31*, 231–249.

Rabinovich, D., Yaniv, S.P., Alyagor, I., and Schuldiner, O. (2016). Nitric Oxide as a Switching Mechanism between Axon Degeneration and Regrowth during Developmental Remodeling. Cell *164*, 170–182.

Sanes, J.R., and Yamagata, M. (2009). Many paths to synaptic specificity. Annu. Rev. Cell Dev. Biol. *25*, 161–195.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source platform for biological-image analysis. Nat. Methods *9*, 676–682.

Senapati, B., Tsao, C.-H., Juan, Y.-A., Chiu, T.-H., Wu, C.-L., Waddell, S., and Lin, S. (2019). A neural mechanism for deprivation state-specific expression of relevant memories in Drosophila. Nat. Neurosci. *in press*.

Shimizu, K., Sato, M., and Tabata, T. (2011). The Wnt5/planar cell polarity pathway regulates axonal development of the Drosophila mushroom body neuron. J. Neurosci. *31*, 4944–4954.

Shin, J.E., and DiAntonio, A. (2011). Highwire regulates guidance of sister axons in the Drosophila mushroom body. J. Neurosci. *31*, 17689–17700.

Smith-Trunova, S., Prithviraj, R., Spurrier, J., Kuzina, I., Gu, Q., and Giniger, E. (2015). Cdk5 regulates developmental remodeling of mushroom body neurons in Drosophila. Dev. Dyn. *244*, 1550–1563.

Sperry, R.W. (1963). Chemoaffinity in the orderly growth of nerve fiber patterns and connections. Proc. Natl. Acad. Sci. U. S. A. *50*, 703–710.

Sweeney, L.B., Chou, Y.-H., Wu, Z., Joo, W., Komiyama, T., Potter, C.J., Kolodkin, A.L., Garcia, K.C., and Luo, L. (2011). Secreted semaphorins from degenerating larval ORN axons direct adult projection neuron dendrite targeting. Neuron *72*, 734–747.

Takemura, S.-Y., Aso, Y., Hige, T., Wong, A., Lu, Z., Xu, C.S., Rivlin, P.K., Hess, H., Zhao, T., Parag, T., et al. (2017). A connectome of a learning and memory center in the adult Drosophila brain. Elife *6*, 1–43.

Tessier-Lavigne, M., and Goodman, C.S. (1996). The molecular biology of axon guidance. Science *274*, 1123–1133.

Tsao, C.-H., Chen, C.-C., Lin, C.-H., Yang, H.-Y., and Lin, S. (2018). Drosophila mushroom bodies integrate hunger and satiety signals to control innate food-seeking behavior. Elife *7*, e35264.

Vogt, K., Schnaitmann, C., Dylla, K. V, Knapek, S., Aso, Y., Rubin, G.M., and Tanimoto, H. (2014). Shared mushroom body circuits underlie visual and olfactory memories in Drosophila. Elife *3*, e02395.

Vogt, K., Aso, Y., Hige, T., Knapek, S., Ichinose, T., Friedrich, A.B., Turner, G.C., Rubin, G.M., and Tanimoto, H. (2016). Direct neural pathways convey distinct visual information to Drosophila mushroom bodies. Elife *5*, e14009.

Wang, J., Zugates, C.T., Liang, I.H., Lee, C.-H.J., and Lee, T. (2002). Drosophila Dscam is required for divergent segregation of sister branches and suppresses ectopic bifurcation of axons. Neuron *33*, 559–571.

White, K., Grether, M.E., Abrams, J.M., Young, L., Farrell, K., and Steller, H. (1994). Genetic control of programmed cell death in Drosophila. Science *264*, 677–683.

Winberg, M.L., Noordermeer, J.N., Tamagnone, L., Comoglio, P.M., Spriggs, M.K., Tessier-Lavigne, M.,
and Goodman, C.S. (1998). Plexin A is a neuronal semaphorin receptor that controls axon guidance.
Cell *95*, 903–916.

Xie, X., Tabuchi, M., Brown, M.P., Mitchell, S.P., Wu, M.N., and Kolodkin, A.L. (2017). The laminar
organization of the Drosophila ellipsoid body is semaphorin-dependent and prevents the formation
of ectopic synaptic connections. Elife *6*, e25328.

Xie, X., Tabuchi, M., Corver, A., Duan, G., Wu, M.N., Kolodkin, A.L., Xie, X., Tabuchi, M., Corver, A.,
Duan, G., et al. (2019). Semaphorin 2b Regulates Sleep-Circuit Formation in the Drosophila Central
Brain Article Semaphorin 2b Regulates Sleep-Circuit Formation in the Drosophila Central Brain.
Neuron 1–16.

Yu, H.H., Araj, H.H., Ralls, S.A., and Kolodkin, A.L. (1998). The transmembrane Semaphorin Sema I is
required in Drosophila for embryonic motor and CNS axon guidance. Neuron *20*, 207–220.

Yu, L., Zhou, Y., Cheng, S., and Rao, Y. (2010). Plexin a-semaphorin-1a reverse signaling regulates
photoreceptor axon guidance in Drosophila. J. Neurosci. *30*, 12151–12156.

Yu, X.M., Gutman, I., Mosca, T.J., Iram, T., Ozkan, E., Garcia, K.C., Luo, L., and Schuldiner, O. (2013).
Plum, an immunoglobulin superfamily protein, regulates axon pruning by facilitating TGF-β signaling.
Neuron *78*, 456–468.

Zhu, S., Lin, S., Kao, C.-F., Awasaki, T., Chiang, A.-S., and Lee, T. (2006). Gradients of the Drosophila
Chinmo BTB-zinc finger protein govern neuronal temporal identity. Cell *127*, 409–422.

Zwarts, L., Goossens, T., Clements, J., Kang, Y.Y., and Callaerts, P. (2016). Axon Branch-Specific
Semaphorin-1a Signaling in Drosophila Mushroom Body Development. Front. Cell. Neurosci. *10*, 210.

Figure legends

Figure 1. Timing of innervations by DANs and MBONs targeting the α'2 zone. (A-F') Axonal innervation in the MB vertical lobes by PPL1-α'2α2 DANs, labeled by *MB058B-splitGAL4* (green, A-F'), at different hours after puparium formation (APF). The brains were counterstained with *LexAop-rCD2::RFP* driven by *MB247-LexA::p65* (magenta, A'-F'). The white arrowheads in (A-B') indicate where the axons start to innervate the MB lobes. **(G-L')** Dendritic innervation in the MB vertical lobes by MBON-α'2, labeled by *MB091C-splitGAL4* (green, G-L'), at different hours APF. The brains were counterstained with *LexAop-rCD2::RFP* driven by *MB247-LexA::p65* (magenta, G'-L'). The white arrowheads in (K-K') indicate where the dendrites start to innervate the MB lobes. **(M-M')** 3-D images of the PPL1-α'2α2 axons in the MB lobes. Axons were labeled with *UAS-mCD8::GFP* driven by *MB058B-splitGAL4* (green, M-M') and all MB lobes were labeled with *LexAop-rCD2::RFP* driven by *MB247-LexA::p65* (magenta, M'-M'), and the α and β lobes were labeled with anti-FasII antibody (blue, M-M'). The white arrowheads in indicate the axonal processes extending at the junctions between the α and α' lobes. **(M'')** A magnified view of the area outlined by a white dashed square in (M'). Scale bar in (A-L'): 50 µm. Scale bar in (M-M''): 15 µm. Detailed genotypes are listed in Table S2.

Figure 2. Ablation of PPL1-α'2α2 DANs has no effect on MB targeting by neighboring DANs or MBON-α'2. (A-A''') A wild-type brain whose PPL1-α'2α2 DANs were labeled with *UAS-mCD8::GFP* driven by *MB058B-splitGAL4* (green, A and A''') and whose MBON-α'2 was labeled with *LexAop-rCD2::RFP* driven by *R20G03-LexA* (magenta, A'' and A'''). The brain was also counterstained with anti-TH antibody (blue, A' and A'''). A single focal plane is shown. The white dashed circle in (A') and the white arrowhead in (A''') indicates the α'2 zone. **(B-B''')** As for (A-A'''), except that PPL1-α'2α2 DANs were ablated with *UAS-rpr,hid* driven by *MB058B-splitGAL4*. **(C-C'')** A wild-type brain whose MBON-α'2 was labeled with *UAS-mCD8::GFP* driven by *MB091C-splitGAL4* (green, C and C'') and its MBON-α'3 labeled with *LexAop-rCD2::RFP* driven by *R53F03-LexA* (magenta, C' and C''). The brain was also counterstained with anti-Trio antibody to label the α', β' and γ lobes (blue, C''). The yellow and white arrowheads indicate the α'2 and α'3 zones, respectively. **(D-D'')** As for (C-C''), except that MBON-α'2 was ablated using *UAS-rpr,hid* driven by *MB091C-splitGAL4*. **(E-E'')** A wild-type brain whose MBON-α'2 was labeled with *UAS-mCD8::GFP* driven by *MB091C-splitGAL4* (green, E and E'') and its MBON-γ2α'1 labeled with *LexAop-rCD2::RFP* driven by *R25D01-LexA* (magenta, E' and E''). The brain was also counterstained with anti-Trio antibody to label the α', β' and γ lobes (blue, E''). A single focal plane is shown. The yellow and white arrowheads indicate the α'2 and α'1 zones, respectively. **(F-F'')** As for (E-E''), except that MBON-α'2 was ablated using *UAS-rpr,hid* driven by *MB091C-splitGAL4*. Scale bars: 50 µm. Detailed genotypes are listed in Table S2.

Figure 3. Loss of vertical lobes results in mistargeted PPL1-α'2α2 axons. (A-B'') Single-cell flip-out clones of PPL1-α'2α2 DANs (green; A, A'', B, and B'') in *ala* mutant brains from which the MB has lost its vertical lobes on the ipsilateral (A-A'') or contralateral (B-B'') side to the clones. The brains were counterstained with anti-Trio and anti-FasII antibodies to label all MB lobes (magenta; A'-A'' and B'-B''). The white arrowheads in (A, A'', B and B'') indicate the mistargeted axons. **(C-C')** Axonal innervation patterns of PPL1-α'2α2 DANs in an *ala* mutant brain at 30 h APF. The PPL1-α'2α2 DANs were labeled with *UAS-mCD8::GFP* driven by *MB058B-splitGAL4* (green), and all KCs were labeled with *LexAop-rCD2::RFP* driven by *MB247-LexA::p65* (magenta, C'). The vertical lobes were missing in one hemibrain. The white arrowheads indicate the mistargeted axons near the missing lobes. **(D-D')** A single focal plane from (C-C') showing the α'2 zones. The white arrowheads indicate the position of the α'2 zone. **(E-E')** A single focal plane from (C-C') showing the layer immediately posterior to the vertical lobes. The white arrowheads indicate the mistargeted axons locate laterally and dorsally to the putative missing α'2/α2 zones. White dashed lines outline the putative position of the missing lobes. Scale bars: 50 µm. Detailed genotypes are listed in Table S2.

Figure 4. MBON-α'2 dendrites innervate other MB lobe zones when their normal target zone is missing. (A-A'') Innervation pattern of MBON-α'2 dendrites labeled with *UAS-mCD8::GFP* driven by *MB091C-splitGAL4* (green, A and A'') in an *ala* mutant brain from which the vertical lobes are missing in one hemibrain. The KCs were labeled with *LexAop-rCD2::RFP* driven by *MB247-LexA::p65* (magenta, A' and A''). White arrowheads in (A and A'') indicate the α'1 and β'2mp zones. Yellow arrowheads in (A and A'') indicate the normal α'2 zone. **(B-B'')** As for (A-A''), except that the MBON-α'2 was labeled with *UAS-dSyd-1-GFP* (green, presynaptic marker, B and B'') and *UAS-DenMark* (magenta, postsynaptic marker, B' and B''). White arrowheads in (B' and B'') indicate the α'1 and β'2mp zones. **(C-E')** Targeting of MBON-α'2 dendrites labeled with *UAS-mCD8::GFP* driven by *MB091C-splitGAL4* (green) in *ala* mutant brains at 36 (C-C'), 48 (D-D'), and 72 (E-E') h APF. The KCs were labeled with *LexAop-rCD2::RFP* driven by *MB247-LexA::p65* (magenta, C', D' and E'). Vertical lobes were missing from one hemibrain. The white arrowheads in (C-D') indicate the α'2 zone, and the arrowheads in (E-E') indicate ectopic innervations of the MBON-α'2 dendrites. Scale bars: 50 μm. Detailed genotypes are listed in Table S2.

Figure 5. Overexpression of *sema1a* redirects PPL1-α'2α2 and PPL1-γ2α'1 dendrites to distinct MB lobe zones. (A-A'') PPL1-α'2α2 DANs labeled with *UAS-mCD8::GFP* driven by *MB058B-splitGAL4* (green, A and A'') in a wild-type brain. The brain was counterstained with anti-Trio and anti-FasII antibodies to label all MB lobes (magenta, A' and A''). **(B-B'')** As for (A-A''), except that Sema1a was additionally overexpressed in PPL1-α'2α2 DANs using *UAS-sema1a* driven by *MB058B-splitGAL4*. Arrowheads in (B and B'') indicate the β'2p, α'1, and α3 zones. **(C-C'')** As for (B-B''), but *UAS-sema1a* was replaced with *UAS-sema1.mEC-5xmyc*. **(D-D'')** As for (B-B''), but *UAS-sema1a* was replaced with *UAS-sema1.mEC.Fc-5xmyc*. **(E-E'')** As for (B-B''), except that the PPL1-α'2α2 DANs were labeled with *UAS-dSyd-1-GFP* (green, presynaptic marker, E and E'') and *UAS-DenMark* (magenta, postsynaptic marker, E' and E''). Arrowheads in (E' and E'') indicate the β'2p, α'1, and α3 zones. **(F-F'')** PPL1-γ2α'1 DANs labeled with *UAS-mCD8::GFP* driven by *MB296B-splitGAL4* (green, F and F'') in a wild-type brain. The brain was counterstained with anti-Trio and anti-FasII antibodies to label all MB lobes (magenta, F' and F''). **(G-G'')** As for (F-F''), except that Sema1a was additionally overexpressed in PPL1-γ2α'1 DANs using *UAS-sema1a* driven by *MB296B-splitGAL4*. Arrowheads in (G and G'') indicate the β'2mp and β'2a zones. Note that this manipulation resulted in loss of MB vertical lobes due to an unknown mechanism. **(H-H'')** A single focal plane from (G-G'') showing the innervation of ectopic PPL1-γ2α'1 neurites in the β'2a zone (outlined by white dashed lines). **(I-I'')** As for (G-G''), except that the PPL1-γ2α'1 DANs were labeled with *UAS-dSyd-1-GFP* (green, I and I'') and *UAS-DenMark* (magenta, I' and I''). Arrowheads in (I' and I'') indicate the β'2mp and β'2a zones. Scale bars: 50 μm. Detailed genotypes are listed in Table S2.

Figure 6. *sema1a* is required for the mistargeting of MBON-α'2 dendrites due to missing vertical lobes. (A-A'') The innervation pattern of MBON-α'2 dendrites labeled with *UAS-mCD8::GFP* driven by *MB091C-splitGAL4* (green, A and A'') in an *ala* mutant brain lacking MB vertical lobes in one hemibrain. All MB lobes were labeled with *LexAop-rCD2::RFP* driven by *MB247-LexA::p65* (magenta, A' and A''). **(B-B'')** As for (A-A''), except that *UAS-sema1a-RNAi* was additionally expressed in MBON-α'2. The white arrowheads indicate the mistargeted dendrites in the β'2mp and α'1 zones and the yellow arrowheads indicate the normal innervation in the α'2 zones. Note the greatly reduced ectopic dendritic innervations and that normal α'2 innervation is not affected by *sema1a* knockdown. Scale bars: 50 μm. Detailed genotypes are listed in Table S2.

Figure 7. *sema1a* is required for the dendritic targeting of MBON-β2β'2a and MBON-α3. (A-A'') The innervation pattern of MBON-β2β'2a dendrites labeled with *UAS-mCD8::GFP* driven by *MB399C-splitGAL4* (green, A-A''). The α', β' and γ lobes were labeled with anti-Trio antibody (magenta, A' and A''). The β'2a zones are outlined with white dashed circles, and the β2 zones are indicated by yellow arrowheads. (A'') is a high magnification view of a single focal plane showing the β'2a and β2 zones.

(B-B'') As for (A-A''), except that *UAS-sema1a-RNAi-1* was additionally expressed by *MB399C-*
splitGAL4 in MBON-β2β'2a. The white arrowheads in (B and B') indicate the mistargeted neurites. **(C-**
D) Wild-type (C) and *UAS-sema1a-RNAi-1*-expressing (D) MBON-β2β'2a labeled with dSyd-1-GFP
(green, presynaptic marker) and DenMark (magenta, postsynaptic marker) driven by *MB399C-*
splitGAL4. The brains were counterstained with anti-Trio and anti-FasII antibodies to label all MB
lobes (blue). The white arrowhead in (D) indicates the mistargeted dendrites. **(E-F')** Wild-type (E-E')
and *UAS-sema1a-RNAi-*1-expressing (F-F') MBON-α'3 labeled with *UAS-mCD8::GFP* driven by
MB027B-splitGAL4 (green). The brains were counterstained with anti-Trio antibody to label the α', β'
and γ lobes (magenta, E' and F'). The white arrowheads in (F and F') indicate the mistargeted
dendrites. **(G-H)** Wild-type (G) and *UAS-sema1a-RNAi-1*-expressing (H) MBON-α'3 labeled with dSyd-
1-GFP (green, presynaptic marker) and DenMark (magenta, postsynaptic marker) driven by *MB027B-*
splitGAL4. The brains were counterstained with anti-Trio and anti-FasII antibodies to label all MB
lobes (blue). The white arrowhead in (H) indicates the mistargeted dendrites.
746　
Figure 8. *sema1a* mutant MARCM clones for MBON-β2β'2a, MBON-α'3, MBON-α'1, and MBON-
β'2mp exhibit defective dendritic targeting. (A-B') Wild-type (A-A') and *sema1a^{k13702}* (B-B') MARCM
clones for MBON-β2β'2a labeled with *UAS-mCD8::GFP* driven by *R21D02-GAL4* (green). The brains
were counterstained with anti-Trio antibody to label the α', β' and γ lobes (magenta, A' and B'). The
white arrowheads in (B-B') indicate the mistargeted dendrites. **(C-D')** Wild-type (C-C') and
sema1a^{k13702} (D-D') MARCM clones for MBON-α'3 labeled with *UAS-mCD8::GFP* driven by *R53F03-*
GAL4 (green). The brains were counterstained with anti-Trio antibody to label the α', β' and γ lobes
(magenta, C' and D'). The white arrowhead in (D-D') indicates the mistargeted dendrites. **(E-F')** Wild-
type (E-E') and *sema1a^{k13702}* (F-F') MARCM clones for MBON-α'3 and MBON-α'1 labeled with *UAS-*
mCD8::GFP driven by *R24H08-GAL4* (green). The brains were counterstained with anti-Trio antibody
to label the α', β' and γ lobes (magenta, E' and F'). The yellow arrowheads in (E-E') indicate the
innervation of MBON-α'3 and the white arrowheads in the same images indicate the innervation of
MBON-α'1. Note that MBON-α'3 was always accompanied by MBON-α'1 in the wild-type MARCM
clones (n = 17), indicating that they are from the same lineage. MBON-α'3 and MBON-α'1 dendritic
innervations were both defective in *sema1a^{k13702}* MARCM clones (F-F'). **(G-H')** Wild-type (G-G') and
sema1a^{k13702} (H-H') MARCM clones for MBON-β'2mp labeled with *UAS-mCD8::GFP* driven by
VT41043-GAL4 (green). The brains were counterstained with anti-Trio antibody to label the α', β'
and γ lobes (magenta, G' and H'). The white arrowheads in (H-H') indicate the mistargeted dendrites.
Scale bars: 50 μm. Detailed genotypes are listed in Table S2.
766　
767　

Supplementary materials

Figure S1. Innervation timings for MBON-α'3. (A-D') Dendritic targeting of MBON-α'3 flip-out clones labeled with *UAS-mCD8::GFP* driven by *MB027B-splitGAL4* (green) at different hours after puparium formation (APF). All KC axons were labeled with *LexAop-rCD2::RFP* driven by *MB247-LexA::p65* (magenta, A'-D'). Scale bars: 50 μm. Detailed genotypes are listed in Table S2.

Figure S2. Anti-TH signals in the MB lobe during metamorphosis. Brains at different hours APF were stained with anti-TH antibody (green). All KC axons were labeled with *LexAop-rCD2::RFP* driven by *MB247-LexA::p65* (magenta). Note that TH signals in zones α'3 and α3 only start to appear at 48 h APF. Scale bars: 50 μm. Detailed genotypes are listed in Table S2.

Figure S3. PPL1-α'2α2 DANs are ablated early in development. (A-B') Axonal innervation patterns for PPL1-α'2α2 DANs labeled with *UAS-mCD8::GFP* driven by *MB058B-GAL4* (green) at 24 (A-A') and 30 (B-B') h APF. The brains were counterstained with anti-FasII antibody to label the α, β, and γ lobes (magenta, A' and B'). Arrowheads indicate the innervations in the MB lobes (A and B). Insets in (A' and B') are the cell bodies of PPL1-α'2α2 DANs. **(C-D')** As for (A-B'), except that *UAS-hid,rpr* was additionally expressed by *MB058B-GAL4* to ablate PPL1-α'2α2 DANs. Note that although the cell bodies of PPL1-α'2α2 DANs were still observed at 24 h APF, no axonal innervation in the lobes was apparent (C-C'). Cell bodies were completely ablated at 30 h APF (D-D'). Scale bars: 50 μm. Detailed genotypes are listed in Table S2.

Figure S4. Ablation of PPL1-α'2α2 DANs has no effect on the dendritic targeting of neighboring MBONs. (A-A'') A wild-type adult brain whose PPL1-α'2α2 DANs and MBON-α'3 were labeled, respectively, with *UAS-mCD8::GFP* driven by *MB058B-splitGAL4* (green, A and A'') and *LexAop-rCD2::RFP* driven by *R53F03-LexA* (magenta, A' and A''). The brain was counterstained with anti-Trio antibody to label the α', β', and γ lobes (blue, A''). The white arrowheads indicate the α'2α2 and α'3 innervations. **(B-B'')** As for (A-A''), except that *UAS-rpr,hid* was additionally expressed using *MB058B-splitGAL4* to ablate PPL1-α'2α2 DANs. The white arrowheads indicate the innervation by MBON-α'3 dendrites (B' and B''). **(C-C'')** As for (A-A''), except that *R53F03-LexA* was replaced with *R25D01-LexA* to label MBON-γ2α'1 (magenta, C'-C''). The white arrowheads indicate the α'2α2 and α'1 innervations. **(D-D'')** As for (C-C''), but *UAS-rpr,hid* was additionally expressed using *MB058B-splitGAL4* to ablate PPL1-α'2α2 DANs. The white arrowheads indicate the innervation by MBON-γ2α'1 dendrites of the α'1 zone (D' and D''). Scale bars: 50 μm. Detailed genotypes are listed in Table S2.

Figure S5. MBON-α'2 are ablated early in development. (A-A') MBON-α'2 labeled with *UAS-mCD8::GFP* driven by *MB091C-splitGAL4* (green) at 24 h APF. The brains were counterstained with anti-FasII antibody to label the α, β, and γ lobes (magenta, A'). The white arrowheads in (A) indicate the cell bodies of MBON-α'2. **(B-B')** As for (A-A'), except that *UAS-hid,rpr* was additionally expressed by *MB091C-splitGAL4* to ablate MBON-α'2. **(C-D')** As for (A-B'), but the brain was dissected at 36 h APF. The arrowheads in (C) indicate the targeting of MBON-α'2 dendrites to the α'2 zones. Note that the cell bodies of MBON-α'2 were completely ablated as early as 24 h APF. Scale bars: 50 μm. Detailed genotypes are listed in Table S2.

Figure S6. Ablation of MBON-α'3 impairs development of the α' lobe. (A-A''') A wild-type brain whose MBON-α'3 was labeled with *UAS-mCD8::GFP* driven by *MB027B-splitGAL4* (green, A and A''') and its KCs labeled with *LexAop-rCD2::RFP* driven by *MB247-LexA::p65* (magenta, A'' and A'''). The brain was also counterstained with anti-Trio antibody to label the α', β' and γ lobes (blue, A' and A'''). The white arrowheads in (A'-A''') indicate the α'3 zone. **(B-B'')** As for (A-A'''), except that MBON-α'3 was ablated with *UAS-rpr,hid* driven by *MB027B-splitGAL4*. **(C)** The parameters used to measure the

position of the α'3 zone relative to the α lobe tip (*h* and *d*) and the width of the α'2 zone (*w*). *d* was
measured as the horizontal distance between the most lateral position of the α'3 zone and the
middle of the α lobe tip. *h* was measured as the vertical distance between the α and α' lobe tips. *w*
was measured as the horizontal distance between two points, one on the inner border and the other
on the outer border of the α'2 zone. The points were selected as the position closest to the brain
midline on each border. **(D)** The lateral position of the α'3 zone (*d*) was reduced when MBON-α'3
was ablated (p<0.0001, n = 10, two-tailed unpaired *t*-test with Welch's correction). **(E)** The relative
height of the α' lobe was reduced (increased in *h*) when MBON-α'3 was ablated (p=0.0009, n = 10,
two-tailed unpaired *t*-test with Welch's correction). **(F)** Growth of the α' lobe was impaired when
MBON-α'3 was ablated. Growth was measured as the relative positions of the tips of the α and α'
lobes (*h*) at 24, 48, and 72 h APF (24 h APF: p=0.7224, n = 6; 48 h APF: p=0.0263, n = 6; 72 h APF:
p=0.002, n = 6; two-tailed unpaired *t*-test with Welch's correction). **(G-G'')** A wild-type brain whose
MBON-α'3 was labeled with *UAS-mCD8::GFP* driven by *MB027B-splitGAL4* (green, G and G'') and its
MBON-α'2 labeled with *LexAop-rCD2::RFP* driven by *R20G03-LexA* (magenta, G' and G''). The brain
was also counterstained with anti-Trio antibody to label the α', β' and γ lobes (blue, G''). The white
arrowheads in (G'') indicate the α'2 zone. **(H-H'')** As for (G-G''), except that the MBON-α'3 was
ablated using *UAS-rpr,hid* driven by *MB027B-splitGAL4*. **(I)** The width of the α'2 zone (*w*) was not
altered upon MBON-α'2 dendrite ablation (p=0.3051, n = 10, two-tailed unpaired *t*-test with Welch's
correction). Scale bars: 50 μm. Detailed genotypes are listed in Table S2.
838
Figure S7. Development of the α' lobe without MBON-α'3. (A-C'') MB lobes labeled with *LexAop-*
rCD2::RFP driven by *MB247-LexA::p65* (magenta, A'-A'', B'-B'', and C'-C'') at 24 (A-A''), 48 (B-B''), and
72 (C-C'') h APF. MBON-α'3 was simultaneously labeled with *UAS-mCD8::GFP* driven by *MB027B-*
splitGAL4 (green; A, A', B, B', C and C'). The brains were counterstained with anti-FasII antibody to
label the α, β, and γ lobes (blue; A'', B'' and C''). Yellow and white arrowheads indicate the tips of
the α and α' lobes, respectively. **(D-F'')** As for (A-C''), except that *UAS-hid,rpr* was additionally
expressed by *MB027B-splitGAL4* to ablate MBON-α'3. Scale bars: 50 μm. Detailed genotypes are
listed in Table S2.
847
848 **Figure S8. Multi-color flip-out clones of PPL1-α'2α2 and MBON-α'3 DANs in the same adult brain.**
849 **(A-A'')** Multi-color flip-out clones of a PPL1-α'2α2 DAN (labeled by both V5-tag and FLAG-tag, white;
850 A and A''), and a PPL1-α'3 DAN (labeled by V5-tag, magenta; A'-A'') in the same adult brain. Images
851 were processed in Imaris. The yellow arrowheads indicate the common entry point in the MB lobes
852 for both neuron types. Scale bars: 20 μm. Detailed genotypes are listed in Table S2.
853
Figure S9. Targeting of MBON-α'2 dendrites is not affected by *sema1a* overexpression. (A-A'')
Targeting of MBON-α'2 dendrites labeled by *UAS-mCD8::GFP* driven by *MB091C-splitGAL4* (green, A
and A''). All KCs in (A-A'') were labeled with *LexAop-rCD2::RFP* driven by *MB247-LexA::p65* (magenta,
A' and A''). The brain in (B-B'') was counterstained with anti-Trio and anti-FasII antibodies to label all
MB lobes (magenta, B' and B''). **(B-B'')** As for (A-A''), except that *UAS-sema1a* was additionally
expressed in MBON-α'2 by *MB091C-splitGAL4*. Scale bars: 50 μm. Detailed genotypes are listed in
Table S2.
861
Figure S10. MBON-β2β'2a and MBON-α'3 express *sema1a*. (A-B'') Expression of V5-tagged
endogenous Sema1a (A and B) and tdTomato driven by the endogenous *sema1a* enhancer (A' and B')
in MBON-β2β'2a (A-A'') and MBON-α'3 (B-B'') dendrites. Arrowheads in (A-A') indicate the
innervations by MBON-β2β'2a dendrites in the β2 and β'2a zones. Arrowheads in (B-B') indicate the
innervations of MBON-α'3 dendrites in the α'3 zones. *MB399C-splitGAL4* and *MB027B-splitGAL4*
were used to remove the stop cassette from *sema1a* Artificial exon in MBON-β2β'2a and MBON-α'3,
respectively. Scale bars: 50 μm. Detailed genotypes are listed in Table S2.
869

Figure S11. *sema1a* **is not required for dendritic targeting of MBON-γ2α'1 and MBON-γ5β'2a. (A-B'')** Wild-type (A-A'') or *sema1a^k13702* (B-B'') MARCM clones for MBON-γ2α'1 labeled with *UAS-mCD8::GFP* driven by *R25D01-GAL4* (green; A, A'', B and B''). The brains were counterstained with anti-Trio to label the α', β' and γ lobes (magenta; A'-A'' and B'-B''). Arrowheads in (A'' and B'') indicate the α'1 zone. **(C-D'')** As for (A-B''), except that the clones are of MBON-γ5β'2a labeled by *R21D02-GAL4*. Arrowheads in (C'' and D'') indicate the β'2a and γ5 zones. Scale bars: 50 μm. Detailed genotypes are listed in Table S2.

Figure S12. Canonical Sema1a ligands are not required for the dendritic targeting of MBONs. (A-A') MBONs innervating the β'2mp, β'2a and β2 zones (indicated by white arrowheads and a dashed circle) were labeled with *UAS-mCD8::GFP* driven by *R21D02-GAL4* (green) in a *Sema2b^f02042, sema2a^p2* double mutant brain. The brain was counterstained with anti-Trio antibody to label the α', β', and γ lobes (magenta, A'). A signal focal plane is shown. Note that dendritic innervations in the MB lobe zones were largely unaffected. **(B-B')** Dendritic innervation of MBON-α3 labeled with *UAS-mCD8::GFP* driven by *R53F03-GAL4* (green) in a *Sema2b^f02042, sema2a^p2* double mutant brain. The brain was counterstained with anti-Trio antibody to label the α', β', and γ lobes (magenta, B'). Note that although the tip of the α' lobe is enlarged, the MBON-α3 dendrites are largely confined to the α'3 zone (indicated by white arrowheads). **(C-D')** MBON-α3 dendrites expressing *UAS-sema1a-RNAi-2* and *UAS-mCD8::GFP* driven by *MB027B-splitGAL4* (green) in a wild-type (C-C') or a *plexA^EY16548* heterozygous mutant brain (D-D'). The brains were counterstained with anti-Trio and anti-FasII antibodies to label all MB lobes (magenta, C' and D'). White arrowheads indicate the mistargeted dendrites. Note that the dendritic mistargeting phenotype was not exacerbated in the *plexA^EY16548* heterozygous mutant brain (compare C and D as well as Fig. 8E). **(E-F')** MBON-β2β'2a dendrites expressing *UAS-sema1a-RNAi-2* and *UAS-mCD8::GFP* driven by *MB0399C-splitGAL4* (green) in a wild-type (E-E') or a *plexA^EY16548* heterozygous mutant brain (F-F'). The brains were counterstained with anti-Trio and anti-FasII antibodies to label all MB lobes (magenta, E' and F'). White arrowheads indicate the mistargeted dendrites. Note that the dendritic mistargeting phenotype was not exacerbated in the *plexA^EY16548* heterozygous mutant brain (compare E and F as well as Fig. 8A). **(G-H')** MBONs innervating the β'2mp, β'2a and β2 zones (indicated by white arrowheads and a dashed circle) were labeled with *LexAop-rCD2::RFP* driven by *R12C11-LexA* (magenta) in a wild-type brain (G-G') or a brain in which *UAS-plexA-RNAi* was expressed pan-neuronally by *nSyb-GAL4* (H-H'). The brains were counterstained with anti-Trio and anti-FasII antibodies to label all MB lobes (green, G'and H'). **(I-J')** MBON-α'3 waslabeled with *LexAop-rCD2::RFP* driven by *R53F03-LexA* (magenta) in a wild-type brain (I-I') or a brain in which *UAS-plexA-RNAi* was expressed pan-neuronally by *nSyb-GAL4* (J-J'). The brains were counterstained with anti-Trio and anti-FasII antibodies to label all MB lobes (green, I'and J'). The white arrowheads indicate the α'3 zones. **(K-N')** As for (G-J'), except that *UAS-plexA-RNAi* was expressed in all glia using *repo-GAL4* (L-L' and N-N'). Scale bars: 50 μm. Detailed genotypes are listed in Table S2.

Table S1. GAL4 lines examined for their potential to follow MBONs or DANs during their morphogenesis.

Table S2. Fly genotypes, immunostaining procedures, and heat-shock conditions for each figures.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

MB399C (MBON-β2β'2a)

Single focal plane

MB399C > *dSyd-1-GFP*, *DenMark*

WT

MB399C > sema1a-i-1

A A' β'2a β2

A'' β'2a β2

C

B B'

B''

10 μm

D

MB027B (MBON-α'3)

MB027B > *dSyd-1-GFP*, *DenMark*

WT

MB027B > sema1a-i-1

E E'

G

F F'

H

Figure 8

Figure S1

	6h APF	12h APF	18h APF	24h APF
MB027B (MBON-α'3)	A / A'	B / B'	C / C'	D / D'

Figure S2

Figure S3

Figure S4

Figure S5

Figure S6

Figure S7

Figure S8

| PPL1-α'2α2 | PPL1-α'3 | Merged |

Figure S9

MB091C (MBON-α'2)	*MB247-LexA* (KCs)	Merged
A	A'	A''

WT

MB091C (MBON-α'2)	Anti-Trio, anti-FasII	Merged
B	B'	B''

MB091C > sema1a

Figure S10

Figure S11

Figure 12