On species delimitation, hybridization and population structure of cassava whitefly in

Africa

Elfekih S[1,2a], Tay WT[3a†], Polaszek A[4], Gordon KHJ[3], Kunz D[3], Macfadyen S[3], Walsh TK[3],

Vyskočilová S[5], Colvin J[5], De Barro PJ[6†]

(1) CSIRO, Australian Animal Health Laboratory, Geelong, VIC, Australia

(2) University of Melbourne, Bio21 Institute, Parkville, VIC, Australia

(3) CSIRO, Black Mountain Laboratories, ACT, Australia

(4) Department of Life Sciences, Natural History Museum, London, UK

(5) Natural Resources Institute, University of Greenwich, Kent, UK

(6) CSIRO, Ecosciences Precinct, Brisbane, Australia

(a) These authors contributed equally

[†] Corresponding authors

Key words: gene flow, speciation, hybridization, pest, genomics

Abstract

The *Bemisia* cassava whitefly complex includes species that cause severe crop damage through vectoring cassava viruses in eastern Africa. Currently, the cassava whitefly complex is divided into species and subgroups based on very limited molecular markers that did not allow clear definition of species and population structure. Based on 14,358 genome-wide SNPs from 63 cassava whitefly individuals belonging to sub-Saharan African (SSA1, SSA2 and SSA4) species, and using a well-curated mtCOI gene database, we show clear incongruities in previous taxonomic approaches underpinned by effects from pseudogenes. We show that the SSA4 species is part of the SSA2 species, and that populations of the SSA1 species comprise of south-eastern (Madagascar, Tanzania) and north-western (Nigeria, Democratic Republic of Congo and Burundi) sub-species that show signatures of allopatric incipient speciation, with a hybrid zone separating these adjacent sub-species. These findings provide the first genomic insights into the evolution and molecular ecology of a highly cryptic hemipteran insect complex in African, and allow the systematic use of genomic data in management and control strategies for this important cassava pest.

Introduction

Confidence in species identification underpins strategies for biological conservation and management of natural resources, providing insight into the evolutionary forces that drive species diversification. While morphological differences have enabled many species to be identified, cryptic species complexes present a particular challenge due to their general lack of readily distinguishable morphological characters. Collecting ecological and life history data, including host plant use and mating behaviour, can be used effectively to mitigate and address these challenges (Liu, et al. 2007; Vyskočilová, et al. 2019; Vyskočilová, et al. 2018), but are often time-consuming and impractical, especially with large numbers of samples. Incorporating DNA markers, i.e., diagnostic mitochondrial DNA (mtDNA) markers (e.g., Behere, et al. 2008; Elfekih, et al. 2018b; Walsh, et al. 2019), genome-wide sequencing (Anderson, et al. 2016; Elfekih, et al. 2018a), and whole-genome resequencing (Anderson, et al. 2018), can increase confidence in differentiating between cryptic species in a complex and identifying inter-species hybrids.

The challenges can become critical when individual cryptic species in a complex have a serious impact on health or agricultural productivity. An example of the latter is presented by insect vectors of virus diseases affecting cassava, the most important food crop grown in Africa (55.2% global production share; average from 1994-2011; FAOSTAT 2017). Since the 1990's, cassava production has been affected by a pandemic of cassava mosaic disease (CMD) caused by a number of cassava mosaic geminiviruses (CMGs) (Legg, et al. 2014; Patil and Fauquet 2009). Although cassava is also widely grown in Latin America (its original home) and South East Asia, CMD has been reported in Africa since the early 1970's (Macfadyen, et al. 2018) and more recently in South-East Asia (Minato, et al. 2019; Wang, et al. 2016). While the vector responsible for the spread of the CMD in South-East Asia is not fully understood, the epidemic in east and central Africa was associated with the distribution

of invasive Sub-Saharan African (SSA) species of the *Bemisia* whitefly complex (Legg, et al.

2014). Invasive whitefly species have become major pests of global agriculture, spreading

plant virus diseases and becoming resistant to chemical control agents (De Barro, et al. 2011).

In particular, the invasive cassava whitefly lineage SSA2 (previously 'Ug2'; Boykin, et al.

2017) has been found in regions affected by the CMD pandemic, and since 2007 has also

successfully expanded into southern Europe (Hadjistylli, et al. 2015), while another lineage,

the SSA1 (previously 'Ug1'; Boykin, et al. 2017), was found beyond the pandemic-affected

geographical regions (i.e., widespread throughout the African continent).

Defining the actual CMD vectors in Africa remains a challenge due to the cryptic

nature of the many species in the Aleyrodidae whitefly *Bemisia* genus (De Barro, et al. 2011;

Lee, et al. 2013; Mugerwa, et al. 2018; Vyskočilová, et al. 2018). At least 38 cryptic species

have now been recognised within what was previously considered a single *Bemisia tabaci*

(Gennadius) species (Martin and Mound 2007) through the characterisation of a partial

mitochondrial DNA cytochrome oxidase subunit I (mtCOI) 3' gene region of 657bp (De

Barro, et al. 2011; Dinsdale, et al. 2010; Kunz, et al. 2019b). However, re-analyses of partial

mtCOI sequences in GenBank relating to the Mediterranean (MED) species by Vyskočilová

et al. (2018), and of the *B. tabaci* standard partial mtCOI dataset (Boykin, et al. 2017) by

Kunz et al. (2019b), demonstrated that a substantial portion (*ca*. 60-65%) were artefacts of

nuclear mitochondrial DNA sequences (NUMTs) or other PCR artefacts (Elfekih, et al.

2018a; Kunz, et al. 2019b; Tay, et al. 2017a). This analysis, together with molecular

characterisation of the original voucher specimens, is beginning to disentangle the species

complex, identifying the Mediterranean (MED) species as the true *B. tabaci* (Tay, et al.

2012), the *B. tabaci* AsiaII_7 species as *B. emiliae* Corbett (Tay, et al. 2017b), assisting with

identifying historical species ranges (Kunz, et al. 2019a; Tay, et al. 2017b; Tay, et al. 2012),

and provided phylogenetic evidence that the cassava whitefly (i.e., SSA1, SSA2, SSA3,

SSA6, SSA9) were ancestral to the 'B. tabaci cryptic species complex' (Kunz, et al. 2019b).

The advent of high-throughput sequencing (HTS) technologies has driven the availability of

genome-wide data that is further transforming the Bemisia taxonomy. HTS has confirmed

that the B. tabaci Middle East Asia Minor 2 (MEAM2), previously recognised as a distinct

species, is in fact the result of an artefact of sequencing nuclear mitochondrial DNA

sequences (NUMTs) (Tay, et al. 2017a ; Elfekih, et al. 2018a).

The taxonomy, population structure and evolutionary genetics of the cassava whitefly

Bemisia species complex (i.e., SSA1, SSA2, SSA3, SSA4, SSA6, and SSA9; see Kunz, et al.

2019b) remain poorly understood, despite the importance of defining and managing the

vectors of CMD. In a recent study of taxonomic structure and gene flow among genetically

diverse SSA1, SSA2, SSA3 and SSA4 species, Wosula et al. (2017) described serious

discrepancies between their results obtained either from mtCOI or by nextRAD sequencing,

with possible implications for understanding the dynamics of the CMD-transmitting species.

The findings of Wosula et al. (2017) have wider implications to the future usage of this

partial mtCOI gene as the current standard DNA marker for species identification of B. tabaci

and related Bemisia cryptic species. This is especially so in the sub-Saharan African Bemisia

species context, where species identified to-date would need to be reanalysed using a

genome-wide sequencing approach. The diversity within genome-wide SNP data, and

complex gene flow patterns existed between these newly defined SSA genetic groups

(Wosula, et al. 2017) would appeared to also have a different operational taxonomic unit to

challenge the traditional 'species' unit.

In this study, we seek to understand why the African cassava whitefly SSA1 (which is

also further refined into sub-groups 1, 2, and 3 in some studies; e.g., Legg, et al. (2014)),

SSA2, and SSA4 species, should have such conflicting species status and admixture patterns

depending on whether the mtCOI or nuclear DNA markers were used. To reconcile these

contrasting findings, we re-analysed the mtCOI data of Wosula et al. (2017) and compared it

to a strictly filtered set of mtCOI sequences for species status delimitation. Identification and

removal of NUMT-derived mtCOI resolved conflict of results from mtCOI discrepancies as

reported by Wosula et al. (2017). Reanalyses of genome-wide SNP data enabled insights into

SSA1-species-wide population substructure to identify the geographic contact zone and

signature of intra- and inter-species hybridization. Our findings provide powerful insights

into the evolutionary forces shaping incipient speciation and vector competence of this

agriculturally invasive species complex with the capacity to impact on food security in sub-

Saharan Africa.

Material and Methods

mtCOI data handling. The mtCOI sequences from the study by Wosula et al. (2017) were

downloaded from GenBank (accession numbers MF417578-MF417602), including all their

published sequences and published reference sequences to identify genetic groups within the

SSA species. These partial mtCOI sequences were imported into Geneious R11.1 and aligned

with reference sequences (Suppl. Fig. 1) using the MAFFT alignment program (Katoh and

Standley 2013) with default settings. We examined all sequences for insertion/deletions

(indels) as signatures of pseudogenes (NUMTs). Sequences that did not contain indels were

assessed also for amino acid substitution patterns at evolutionary conserved regions in the

partial COI gene region to assist with further identification of potential NUMTs.

Identification of amino acid residue conservation sites across the partial mtCOI gene regions

was as described in Kunz et al. (2019b).

nextRAD data processing. The raw data from nextRAD sequencing of Wosula et al. (2017)

was accessed from GenBank (SRP103541). The fastq sequences were screened using FastQC

for quality control. The retained reads were trimmed by quality to a total length of 151 bp

using Trimmomatic (Bolger, et al. 2014).

Mapping quality. We mapped the reads from each individual sample to the *B. tabaci*

MEAM1 genome (Chen, et al. 2016) using Burrows-Wheeler Aligner (BWA) v.0.7.12 (Li

and Durbin 2010). The alignments used the BWA-MEM algorithm with default settings. The

SAM files were converted to BAM output using Samtools (Li, et al. 2009). The BAM files

were subsequently sorted and indexed, and checked for quality and mapping percentages per

scaffold.

SNP genotyping. SNPs were first called using a *de novo* approach in PyRAD (Eaton 2014).

PyRAD is a pipeline designed for RADseq datasets that aims to capture variation at the

species/clade level. It allows clustering of highly divergent sequences and takes into

consideration indel variation. We also performed SNP calling in PyRAD, using reads mapped

to the MEAM1 reference genome (Chen, et al. 2016). Given the low-depth nature of the data,

we analysed the data using an additional pipeline. We used the BAM files generated in BWA

as input for the program ANGSD (Korneliussen, et al. 2014), which relies on a statistical

approach that takes into consideration genotype uncertainty (Fumagalli, et al. 2013).

Genetic clusters and tests for introgression. Genetic clusters within the dataset that consists

of 63 SSA samples were investigated by Principal Component Analysis (PCA) using the

SNPRelate R package (Zheng, et al. 2012). We used ADMIXTURE v.1.3.0 (Alexander, et al.

2009) on the 63-sample dataset to identify genetic clusters and estimate the genetic ancestry

of the SSA samples. We ran the program with K ancestral clusters varying from 1 to 10 with

each 'K' value repeated 100 times. A cross-validation test was performed to determine the

optimal K value. The program ANGSD v.0.911 (Korneliussen, et al. 2014) was used to run

an ABBA-BABA test (D-statistics) to investigate introgression patterns between the 63

samples (SSA1 and SSA2 species) and with the MEAM1 genome as an outgroup. The

ABBA-BABA test is based on the comparison of the number of tree topologies of the ABBA

and BABA patterns. In this study, the SSA species group patterns we investigated were as

defined   by   Wosula   et   al.   (2017)   and   consisted   of

[ESA+CA/SSA2+SSA4/ECA+WA/MEAM1], where 'ESA', 'CA', 'ECA', and 'WA'

represented East South Africa, Central Africa, East Central Africa, and Western Africa,

respectively. We used the MEAM1 species as outgroup for comparison and to infer

introgression patterns. We tested the significance of these ABBA/BABA patterns using the

corresponding Z-scores, obtained in ANGSD using a jack-knife procedure. An absolute Z-

score value of ≥ 3 is used as a cut-off value (Reich, et al. 2009).

Phylogenetic analyses. The topology of the phylogenetic relationships between individuals

/populations/subspecies and species within the dataset (63 SSA samples and three *B. afer*

individuals (TZ_CHA1, TZ_CHA2, TZ_KIL1) as the outgroup) was examined using a

maximum likelihood (ML) approach. The SSA species were first identified based on the

mtCOI marker, then, a phylogeny was reconstructed using the nextRAD genome-wide SNPs

excluding samples with low genotype quality to minimize biases that could potentially be

introduced by missing data. The phylogenetic reconstruction was carried out in RAxML

v.7.2.8 (Stamatakis 2006) using the GTR substitution model and GTRGAMMA as the

GAMMA model of rate heterogeneity, with 1,000 bootstrap replications. The phylogenetic

clustering obtained in RAxML was used to infer a population ML tree in TreeMix (Pickrell

and Pritchard 2012), in order to identify genetic mixing, the history of population splits and

admixture patterns. We ran the TreeMix simulations using 0 to 5 migrations events with 100

bootstrap replications.

Results

Integrity of African cassava *Bemisia* SSA mtCOI sequences. Of the 11 SSA4 sequences (originally labelled as 'Sub-Saharan III: Western Africa-Cameroon/Cassava'; see Berry et al. (2004), Table 1), only two (Cam Ayos 1 WO2 and Cam Ayos 2 WO3) did not exhibit any premature stop codons at the amino acid level or have indels at the nucleotide level (Suppl. Fig. 1). Both of these sequences were also much shorter than the remaining nine sequences within the Sub-Saharan III: Western Africa-Cameroon/Cassava clade. Comparison of amino acid composition against 21 high-throughput sequencing-generated mtCOI genes representing 12 evolutionary divergent *Bemisia* cryptic species (Kunz, et al. 2019b), enabled Cam Ayos 2 WO3 to be identified as a potential NUMT sequence. Indels, stop codons, non-conservation of amino acid residues in highly conserved mtCOI region therefore ruled out 10 of the 11 SSA4 sequences as potential NUMTs. The remaining single, short but 'functional' Cam Ayos 1 WO2 sequence (AF344246) that clustered with the 10 NUMT-related SSA4 sequences suggested that as a whole, these SSA4 sequences were likely of nuclear genomic origin. The clustering of SSA4 sequences to SSA2 indicates it likely to be of SSA2 nuclear origin, similar to that reported for *B. tabaci* MEAM2 NUMT that originated from the nuclear genome of MEAM1 (Elfekih, et al. 2018a; Tay, et al. 2017a).

The examination of nucleotide substitution patterns identified indels-affected sequences and inconsistencies with respect to number of base substitutions for distinguishing 'subgroups' (SG) within the SSA1 species. For example, the SSA1-SG1/SG2 sequences (accession numbers: MF417578-MF417602) were affected by indels that should not be present in a mitochondrial DNA COI protein coding gene, suggesting that these sequences were associated with sequencing errors and/or derived from non-coding NUMT DNA sequences. The re-assessment of the two SSA1-SG5 sequences (MF417585, MF417586) showed that they differed from the SSA1-SG3 sequences by only three C/T substitutions that resulted in 0.56% nucleotide difference. The basis for suggesting MF417585 and MF417586

belong to different 'sub-groups' was therefore unclear, given that similar numbers of

nucleotide substitutions between sequences (e.g., MF417602 and MF417587: SSA1-SG2;

two transitions (i.e., 2xC/T), 0.37% nucleotide difference), and sequences with greater

numbers of nucleotide substitutions (e.g., MF417590, MF417582: SSA1-SG1; three

transitions (2xG/A, 1xT/C) and one transversion (A/T); 0.75% nucleotide difference) were

placed within the same subgroups.

Species delimitation and introgression. The genome-wide SNP-based Principal Component

Analysis (PCA) discriminated between SSA1 and SSA2, suggesting that they are genetically

distinct and likely to be different species, whereas samples from SSA4 clustered with SSA2

(Fig. 1). Three groupings were identified within SSA1, suggesting a level of intra-species

substructure. These observations were further explored using ML phylogenies generated with

RAxML. An initial such analysis based on the mtCOI sequences (Fig. 2A) broadly divided

these 63 SSA individuals into three clusters containing the populations of (Wosula, et al.

2017): (i) SSA2 cluster containing individuals previously identified as either SSA4 or SSA2

by mtCOI gene sequences with 100% bootstrap confidence, (ii) SSA1 cluster of the

populations defined as 'SSA-CA/SSA-ESA' (node confidence 89%), and (iii) SSA1 cluster

that defined as 'SSA-ECA/SSA-WA' populations (node confidence 96%).

Fig. 1. Principal component analysis of SSA *Bemisia* samples. Principal component

analysis (PCA) was based on 14,358 genome-wide SNPs across the 63 sub-Saharan African

cassava *Bemisia* samples. Clustering of SSA2 (blue circle) from SSA1 (red circles) supported

their genetic distinctness as separate species. Three groupings are identified within SSA1,

suggesting a level of intra-species substructure including the potential gene flow contact

zone. Country of origins for these SSA1 and SSA2 *Bemisia* species are shown by unique

colour codes.

A comprehensive RAxML phylogenomic analysis based on the 14,358 genome-wide

SNPs (Figs. 2B, 2C) confirmed SSA1 and SSA2 species status and is consistent with that

earlier identified using partial mtCOI gene sequence identity. This analysis clearly showed

SSA1 to comprise two clusters, denoted SSA1-SE (South-Eastern) and SSA1-NW (North-

Western) (Fig. 2C), respectively, reflecting the geographic distribution of the populations

identified within them (SSA1-SE comprising samples from Democratic Republic of Congo

(DRC), Tanzania (TZ), and Madagascar (MAD); and SSA1-NW comprising samples from

Cameroon (CMR), Nigeria (NGA), Democratic Republic of Congo (DRC), Rwanda (RWA),

Burundi (BUR) and Tanzania (TZ)). Within SSA1-NW, the 'SSA-ECA' and 'SSA-WA'

populations as defined by Wosula et al. (2017) are clustered separately with 99% and 96%

confidence values. Within SSA1-SE, the 'SSA-ESA' and 'SSA-CA' populations as defined

by Wosula et al. (2017) are partially resolved, sharing only 34% confidence and suggesting

insufficient genomic evidence for this phylogenetic analysis to confirm them as separate

genetic groups.

Fig. 2. Phylogeny of SSA populations. A: Phylogeny based on mtCOI of 63 SSA *Bemisia*

individuals. The phylogeny was constructed using RAxML with 1,000 bootstrap replications.

Sample names are as in Fig. S1, and GenBank accession numbers for the sequences used, and

corresponding mtCOI SSA1-subgroups (SG) of either SG1, SG2 and SG3, are listed on the

right. The *Bemisia* SSA2-SSA4 clade is indicated by yellow branches, and the lineages

corresponding to the SSA1 populations analysed are indicated as follows: CA (pink), ESA

(black), WA (green) and ECA (red). **B:** Phylogeny based on 14,358 genome-wide SNPs of

the individuals analysed in A. The phylogeny was constructed using RAxML with 1,000

bootstrap replications. Sample names and lineage colours are as for **A** above. **C:** Detail of the

RAxML phylogeny in **B** showing only the SSA1 and SSA2-SSA4 lineages, with population

names added. Bootstrap confidence percentages are visible for the main branches.

Fig. 2A

Fig. 2B

Fig. 2C

Admixture analysis identified three genetic clusters (Fig. 3) that separated the SSA2

species (highlighted in green) from the SSA1 species, which was again divided into two

distinct genetic backgrounds corresponding to SSA1-NW (containing SSA-WA and SSA-

ECA (highlighted in blue)) and SSA1-SE (containing SSA-CA and SSA-ESA (highlighted in

red)). Some SSA2 and all the SSA1-CA samples showed evidence for low levels of

admixture. Interestingly, F_{st} analysis (Table 1) supported the status of the two sub-groups

SSA1-NW and SSA-SE as intermediate at the species-level differentiation between any of

the SSA1 populations and the pooled SSA2 populations (the latter ranging from 0.28582 –

0.33406). This was particularly evident for SSA1-NW versus the SSA-ESA population of

SSA1-SE, with F_{st} of 0.22605 - 0.24304. F_{st} differentiation was less clear for the more

weakly defined SSA-CA population, whose F_{st} scores versus SSA1-NW were not

significantly different to those between populations within the two major SSA1 groups.

While the overall differentiation between SSA1-NW and SSA1-SE is consistent with them

being sub-species of SSA1, the observations concerning the SSA-CA population suggest

gene flow between CA and neighbouring populations from both sub-species.

Fig. 3. Populations and species of *Bemisia* across sub-Saharan Africa. Population

structure analysis using Admixture of 63 individuals from the sub-Saharan *Bemisia* cryptic

species clade based on 14,358 SNPs (K=3). Individuals were clustered into the SSA2 species

group, as well as two genetic clusters (red, blue) that belonged to SSA1 based on partial

mtCOI gene signature. The samples are shown on the map: green squares represent SSA-

WA, red squares represent SSA-ECA, pink squares represent SSA-CA and black squares

represent SSA-ECA individuals. Yellow and blue squares represent SSA2 and 'SSA4'

individuals, respectively (the latter being NUMT artefacts of SSA2 – see text). One SSA1

genetic cluster in the Admixture panel included the SSA-WA and SSA-ECA groups (35 blue

individuals and one red-blue individual), and the other SSA1 genetic cluster included five

SSA-CA (blue-red/blue-red-green) and 12 SSA-ESA (red) individuals. Evidence of gene

flow between SSA1 genetic clusters were predominantly detected in SSA-CA individuals

(four red-blue individuals in the Admixture analysis). Three inter-specific hybrids (two from

SSA2; one from SSA-CA with SSA1 (red-blue) and SSA2 (green) genetic backgrounds)

were also detected in the SSA *Bemisia* individuals analysed. The gene flow contact zone

between populations on the NW-SE transect is indicated by the thick blue dashed line. SSA1-

NW and SSA1-SE sub-species boundary lines are indicative only. Landscape elevations

across the sampling sites are shown in insert.

Fig. 3A

319 **Fig. 3B**

330 **Table 1.** Population substructure based on the F-statistics (F_{st}) of (Weir and Cockerham

331 1984). Treatment of SSA1 as single species against the SSA2/SSA4 species showed

332 significant population substructure (F_{st} range 0.28 – 0.33). This limited gene flow between

333 SSA2/SSA4 and the SSA1 (i.e., SSA-WA, SSA-ECA, SSA-CA. SSA-ESA) supported their

separate species status. Treatment of the four SSA1 populations as two well-defined groups, the proposed sub-species, showed that that comprising SSA-WA and SSA-ECA overall shared intermediate levels of gene flow with the SSA-ESA population. Gene flow at the contact zone (SSA-CA) is highest with the SSA-WA, SSA-ECA, and SSA-ESA genetic groups as evident by the low F_{st} values.

Species	SSA2	SSSA1			
Sub-species		SSA1-NW		SSA1-SE	
Populations	SSA2/SSA4	SSA-WA	SSA-ECA	SSA-CA	SSA-ESA
SSA2/4					
SSA-WA	0.29068				
SSA-ECA	0.33406	0.17048			
SSA-CA	0.28582	0.1658	0.1575		
SSA-ESA	0.33173	0.24304	0.22605	0.16026	

Within sub-species SSA1-NW, the SSA-WA population is predominantly found in the lowlands (*ca*. 300m - 1500m), whereas the SSA-ECA/SSA-CA populations were found at higher altitudes (from *ca*. 600m - ≥1500m). Individuals belonging to the SSA-ESA group were predominantly from coastal landscape of Tanzania and Madagascar (Fig. 3). A gene flow contact zone was detected at the SSA-CA region represented by individuals from DRC (n = 5), and one TZA individual from SSA-ECA group. Hybridization was detected in two SSA2 individuals (green/blue genetic backgrounds) and in one DRC individual from SSA-CA (red/green/blue in Fig. 3).

Results from our D-statistics test (ABBA-BABA test) between three groups

[ESA+CA], [SSA2+SSA4] and [ECA+WA] using MEAM1 as an outgroup identified strong

signals of introgression between the [ESA+CA] and [ECA+WA] groups which belonged to

the same SSA1 species based on mtCOI gene analysis, where the strong introgression signals

between [ESA+CA] and [ECA+WA] could potentially represent well-defined sub-

populations/sub-species occupying different geographical landscapes, with gene flow

occurring between these sub-populations/sub-species at the contact zone (Fig. 4).

Fig. 4. Introgression site patterns using a four-Taxon ABBA/BABA test. A. The ABBA

and BABA sites used in the test are derived from ESA+CA and ECA+WA belonging to

SSA1 species, compared to the SSA2+ SSA4 (**Fig. 4B**) group which are previously reported

as two distinct species but were identified as belonging to one species (SSA2) in this study.

Fig. 4A

Fig. 4B

The test also detects signals of introgression within the same geographical region between the SSA2 and SSA1 (i.e., [ECA+WA]) species. The same signatures of interspecies admixture were also evident from the Treemix results (Fig. 5). The ML tree generated in Treemix also supported the SSA4 species as part of SSA2. The populations/subspecies SSA-ECA and SSA-WA cluster together which is consistent with their geographical origin (i.e., SSA1-NW), whereas the SSA-CA population clusters with the SSA-ECA population (i.e., representing the south-eastern geographic origin). Gene flow between these SSA1 populations is especially evident at the contact zone (i.e., at the SSA-CA geographical location). The migration edges (m = 2) revealed gene flow between SSA2 and the SSA1 species from central Africa (i.e., SSA-CA) and supports detection of hybrid individuals identified in the admixture analysis (Fig. 3).

381 **Fig. 5. Gene flow between SSA species.** TreeMix analysis showing gene flow from SSA1 to

382 SSA2/SSA4 (red arrow). Clustering between ECA and WA, and between ESA and CA

383 indicated close evolutionary relationship. SSA2 and SSA4 shared a genetic clade with no

384 branch depth indicating these two species were identical.

Discussion

We combined genome-wide SNPs and mtCOI markers to gain unprecedented

evolutionary ecological and population genomic insights into the sub-Saharan African

cassava whitefly species complex. By taking into consideration the impact of NUMTs, we

showed that there is no conflict between nuclear and mitochondrial DNA markers in terms of

defining species status, contrary to the study of Wosula et al. (2017) based on the same data

set. A thorough filtering of genome-wide SNP markers also discerned the evolutionary forces

shaping incipient speciation and admixture in the *Bemisia* SSA1 whitefly species. Various

evolutionary genetic studies based on the partial mtCOI gene have proposed the presence of

population subgroups (e.g., Ghosh, et al. 2018; Legg, et al. 2014) in the SSA1 species. Based

on genome-wide SNP markers, we found no evidence to support the subgroup classification,

although for the first time, the SSA1 species as a whole, could be broadly divided into two

well-defined north western (NW) and south eastern (SE) populations/sub-species, with a

contact zone separated by the eastern highland and central/western lowland regions. Evidence

for interspecies hybridisation was also detected between the SSA1 and SSA2 species, where

bidirectional genetic contributions were observed between SSA2 and SSA1, and will have

important implications to understanding the evolutionary genetics associated between plant

virus transmission capacity and the socioeconomically important SSA *Bemisia* whitefly pest

complex.

In the study of Elfekih et al. (2018a), mtCOI sequences from the MEAM1, MED,

Indian Ocean (IO) and Australia (AUS) *B. tabaci* cryptic species were interrogated prior to

generating nextRAD genome-wide SNP data to investigate the history of global introduction

pathways of the MED and MEAM1 species. Elfekih et al. (2018a), Elfekih et al. (2015),

Elfekih et al. (2016a), and Elfekih et al. (2016b) showed that species status of these cryptic *B.*

tabaci species as ascertained by the partial mtCOI gene was corroborated by genome-wide

SNP data. Similar to this study, genome-wide SNP markers also detected genomic signatures

of hybridization between the MED, MEAM1 and IO, while also revealing the role played by

NUMTs in confounding our effort to define species status such as between the MEAM1 and

MEAM2 species (Delatte, et al. 2005; Elfekih, et al. 2018a; Tay, et al. 2017a).

Wosula et al. (2017) adopted the nextRAD genome-wide sequencing method outlined

by Elfekih et al. (2015), Elfekih et al. (2016a), and Elfekih et al. (2016b) to investigate gene

flow patterns and diversity in the African cassava whitefly *Bemisia* SSA1, SSA2, SSA4

species complex, but overlooked the impact of NUMTs associated with the *B. tabaci* mtCOI

partial sequences. This oversight led to the major conclusion that the widely used mtCOI

gene was an unsuitable marker for differentiating between African *Bemisia* cryptic species.

Increasingly, reanalyses of the *B. tabaci* complex cryptic species partial mtCOI gene dataset

have detected the presence of significant NUMTs and pseudogenes (Kunz, et al. 2019b; Tay,

et al. 2017a; Vyskočilová, et al. 2018). These studies highlighted the challenges faced by

researchers to delimit species in the cryptic *B. tabaci* species complex, despite the promises

of molecular systematics and species delimitation via mitochondrial barcoding approaches.

Similar to MEAM2 (Boykin, et al. 2013; Dinsdale, et al. 2010), the *Bemisia* cryptic species

SSA4 (Berry, et al. 2004) was shown to be the product of NUMTs. This species

misidentification underpinned all subsequent interpretations of gene flow and evolutionary

ecology in the SSA *Bemisia* cassava whitefly species (Wosula, et al. 2017) based on genome-

wide SNP markers.

Significant genetic diversity in the *B. tabaci* species complex can obscure species and

sub-species identification. For instance, Elfekih et al. (2018a) detected significant diversity in

the MED species based on nextRAD SNP data; however, the MED-ASL species was not

identified until confirmed by complete mitogenome study, mating experiments (Vyskočilová,

et al. 2018), and host preference studies (Vyskočilová, et al. 2019). Similar studies are needed

to provide additional biological support for subgroup/sub-species status within SSA1.

In developing the DNA barcoding approach, Hebert et al. (2003) showed that the COI

gene is especially effective at differentiating at the species, order and phylum levels due to

the associated nucleotide substitutional rates for this gene, while for fine-scale identification

such as within species level, the use of genes with greater substitutional rates were

recommended. Subgroup classification within the SSA1 African cassava whitefly species

based on the mtDNA gene might be possible, provided that a gene region with an appropriate

substitutional rate is identified. Nevertheless, there is currently no evidence for maternal

lineage-based patterns to population dispersal behaviour, host adaptation, and fitness cost

differentiation in the *Bemisia tabaci* complex and the African cassava *Bemisia* species

complex, and therefore the subgroup delimitation based on the partial mtCOI gene is not

justified.

Previously, Legg et al. (2014) reported that the SSA1-SG1, representing the major

SSA1 mtCOI subgroup, was implicated in the severe CMD pandemic in East Africa. Based

on the subgroup classification, Wosula et al. (2017) reclassified these SSA1-SG1 individuals

into three SNP-based groups of SSA-ECA, SSA-CA, and SSA-WA due to perceived

inconsistencies (e.g., SSA1-SG3, SSA1-SG5; see Ghosh et al. (2015)) based solely on the

utilization of the short (i.e., *ca*. 657bp) mtCOI partial gene marker. Based on the same

dataset, our reanalysis found that while phylogenetic signals were detected for the separate

clustering of the 'WA' and 'ECA' populations, there was not a strong genetic difference

between the 'CA' and 'ESA' populations, and we therefore caution against the adoption of

the nomenclature as proposed by Wosula et al. (2017) in future studies. Furthermore,

admixture analysis also showed a substantial gene flow limitation between the 'WA/ECA'

groups and the 'CA/ESA' groups.

Although adopting the subgroup status could help resolve virus transmission capacity

and provide insights into understanding pest introduction pathways (Anderson, et al. 2018;

Tabachnick 1996), there is currently insufficient evidence to link either the SSA1-NW (i.e.,

'WA/ECA') or SSA1-SW (i.e., 'CA/ESA') populations/subgroups to historical CMD

outbreaks. Legg et al. (2002) described a distinct genetic cluster (*B. tabaci* Ug2) assumed to

be associated with the CMD epidemic in the 1990's, mainly due to its geographic distribution

in relation to the front of the epidemic. However, transmission leading to outbreaks of CMD

in the sub-Saharan Africa region is complex, with one factor being the sharing and

propagation of diseased cassava cuttings between farmers. There was a recombination event

between two geminiviruses that produced a more virulent hybrid (Colvin, et al. 2004),

however we have little historical information about how the viruses interacted with the

different *Bemisia* cryptic species in terms of disease dynamics (Polston, et al. 2014). The role

of the SSA2 species in the CMD epidemic is also unclear given that the SSA2's habitat range

(east and west Africa) overlaps the SSA1 species (Legg, et al. 2002; Mugerwa, et al. 2018;

Wosula, et al. 2017). The exact relationship between SSA1/SSA2 and outbreaks of CMD

(Manani, et al. 2017) should be re-investigated, taking into consideration the current

nextRAD SNP data as genomic evidence that supports the presence of hybrid SSA1/SSA2

individuals. Furthermore, the vector competencies of other common cassava whitefly species

(such as *Bemisia afer*) to different cassava viruses also needs to be fully understood.

Stringent delimitation of population groups across the full habitat range of the SSA1

species will be needed to ascertain whether the SSA1 as a whole is represented by genetically

distinct groups where limited gene flow could be further identified to external (e.g.,

geographic, climatic) and/or metagenomic (e.g., secondary symbionts) factors. The genetic

differentiation between the two SSA1 genetic groups suggests they can be considered as

subspecies of SSA1. Recognition of such biological classifications and their distribution in

sub-Saharan Africa will be important for management of these agriculturally and biosecurity

significant species, as for others posing a threat to global health and biosecurity (Paini, et al.

2016).

The South-Eastern SSA1 sub-populations belonging to Madagascar (SSA-ESA) and

Tanzania (SSA-CA) are geographically isolated from the North-Western sub-populations

sampled from Nigeria (SSA-WA), Democratic Republic of Congo and Burundi (SSA-ECA).

The mountain range separating these two SSA1 categories (SE and NW sub-populations)

poses a geographical barrier to gene flow, and representing a potential factor that could

promote incipient allopatric speciation. For instance, the East African Rift Valley is a major

geographical barrier providing the right environment for incipient species and recent

radiations to thrive and evolve (Gottelli, et al. 2004). Specifically, the western branch of the

Rift Valley, i.e., the Albertine Rift covering part of Tanzania, Burundi, Rwanda, Uganda and

Democratic Republic of Congo, is a chain of mountains situated at 1500–3500 meters above

sea level, and formed by volcanic activity that started during the plio-pleistocene. In addition

to the volcanic activity, climate change during this glacial period led to habitat fragmentation

and therefore species diversification in the newly formed biogeographical zones (Huhndorf,

et al. 2007). In addition to allopatric forces, species diversification in the whitefly cryptic

species system has been explained by host plant associations and the ability of this system to

feed on a wide range of host plants including acquiring new ones (Matsubayashi, et al. 2010).

Using RNAseq, Malka et al. (2018) provided insights into how the variation in level of

expression of detoxification genes might have conferred plasticity to the complex and that

likely also contributed to its diversification.

B. tabaci sensu lato on cassava in sub-Saharan Africa is a species complex going

through a continuous diversification process. The examination of the genetic diversity via

mitochondrial and genome-wide markers validated discrimination between SSA1 and SSA2,

and the latter being the same as SSA4. This was mainly facilitated by careful data analysis,

and taking into consideration artefacts caused by pseudogenes (NUMTs). The SSA1 species

shows signatures of incipient speciation with sub-populations divided into two

biogeographical niches; North Western (NW) and South Eastern (SE) regions with a hybrid

zone between Tanzania, Uganda and Democratic Republic of Congo. We conclude that

accurate species identification that takes into consideration all the evolutionary forces driving

the diversification in this system will be critical in the implementation of any future

management strategy for this invasive and economically important insect pest complex.

Further comparative analyses on the endosymbiont metacommunities and host association

can shed more light onto the major causes underlying the gradual process of species

diversification and adaptive radiations in the cassava whitefly system.

Acknowledgements

SE was funded by the CSIRO Office of the Chief Executive (OCE) post-doctoral fellowship

(R-4546-1) and a European Molecular Biology Organization (EMBO) fellowship ASTF-

6889. WTT, SE, KG DK and TW were supported by CSIRO H&B Genes of Biosecurity

Importance fund (R-8681-1). SM, AP, WTT, SV, JC, PDB were supported by the Natural

Resources Institute, University of Greenwich from a grant provided by the Bill & Melinda

Gates Foundation (Grant Agreement OPP1058938).

Literature cited

Alexander DH, Novembre J, Lange K 2009. Fast model-based estimation of ancestry in

unrelated individuals. Genome Research 19: 1655-1664. doi: 10.1101/gr.094052.109

Anderson CJ, et al. 2018. Hybridization and gene flow in the mega-pest lineage of moth,

Helicoverpa. Proc Natl Acad Sci U S A 115: 5034-5039. doi: 10.1073/pnas.1718831115

Anderson CJ, Tay WT, McGaughran A, Gordon K, Walsh TK 2016. Population structure and

gene flow in the global pest, *Helicoverpa armigera*. Mol Ecol 25: 5296-5311. doi:

10.1111/mec.13841

Behere GT, Tay WT, Russell DA, Batterham P 2008. Molecular markers to discriminate

among four pest species of *Helicoverpa* (Lepidoptera: Noctuidae). Bull Entomol Res 98:

599-603. doi: 10.1017/S0007485308005956

Berry SD, et al. 2004. Molecular evidence for five distinct *Bemisia tabaci* (Homoptera :

Aleyrodidae) geographic haplotypes associated with cassava plants in sub-Saharan

Africa. Annals of the Entomological Society of America 97: 852-859. doi: Doi

10.1603/0013-8746(2004)097[0852:Meffdb]2.0.Co;2

Bolger AM, Lohse M, Usadel B 2014. Trimmomatic: a flexible trimmer for Illumina

sequence data. Bioinformatics 30: 2114-2120. doi: 10.1093/bioinformatics/btu170

Boykin LM, Bell CD, Evans G, Small I, De Barro PJ 2013. Is agriculture driving the

diversification of the *Bemisia tabaci* species complex (Hemiptera: Sternorrhyncha:

Aleyrodidae)?: Dating, diversification and biogeographic evidence revealed. BMC Evol

Biol 13: 228. doi: 10.1186/1471-2148-13-228

Boykin LM, Savill A, De Barro P 2017. Updated mtCOI reference dataset for the *Bemisia*

tabaci species complex F1000Res 6: 1835. doi: 10.12688/f1000research.12858.1

Chen WB, et al. 2016. The draft genome of whitefly *Bemisia tabaci* MEAM1, a global crop

pest, provides novel insights into virus transmission, host adaptation, and insecticide

resistance. BMC Biology 14: 110. doi: 10.1186/s12915-016-0321-y

Colvin J, Omongo CA, Maruthi MN, Otim-Nape GW, Thresh JM 2004. Dual begomovirus

infections and high *Bemisia tabaci* populations: two factors driving the spread of a

cassava mosaic disease pandemic. Plant Pathology 53: 577-584. doi: 10.1111/j.1365-

3059.2004.01062.x

De Barro PJ, Liu SS, Boykin LM, Dinsdale AB 2011. *Bemisia tabaci*: a statement of species

status. Annu Rev Entomol 56: 1-19. doi: 10.1146/annurev-ento-112408-085504

Delatte H, et al. 2005. A new silverleaf-inducing biotype Ms of *Bemisia tabaci* (Hemiptera :

Aleyrodidae) indigenous to the islands of the south-west Indian Ocean. Bulletin of

Entomological Research 95: 29-35. doi: 10.1079/Ber2004337

Dinsdale A, Cook L, Riginos C, Buckley YM, De Barro P 2010. Refined Global Analysis of

Bemisia tabaci (Hemiptera: Sternorrhyncha: Aleyrodoidea: Aleyrodidae) Mitochondrial

Cytochrome Oxidase 1 to Identify Species Level Genetic Boundaries. Annals of the

Entomological Society of America 103: 196-208. doi: 10.1603/An09061

Eaton DAR 2014. PyRAD: assembly of de novo RADseq loci for phylogenetic analyses.

Bioinformatics 30: 1844-1849. doi: 10.1093/bioinformatics/btu121

Elfekih S, et al. 2018a. Genome-wide analyses of the *Bemisia tabaci* species complex reveal

contrasting patterns of admixture and complex demographic histories. PLoS One 13:

e0190555. doi: 10.1371/journal.pone.0190555

Elfekih S, et al. 2015. Evolutionary genomics of *Bemisia tabaci* and characterization of its

endosymbiont metacommunities using nextRAD sequencing. International Plant and

Animal Genome Asia 2015 July 12-15th, 2015: Singapore.

Elfekih S, et al. 2016a. Genome-Wide SNPs Decipher Global Incursion Pathways in the

Bemisia tabaci Species Complex. International Plant and Animal Genome Conferences.

2016a 9-13 January 2016: San Diego, CA, U. S. A.

Elfekih S, et al. 2016b. Genome-wide scans unravel fine-scale invasion routes in the *Bemisia*

tabaci species complex. 2nd International Whitefly Symposium, . 2016b 14-19 February

2016 Arusha, Tanzania. p38.

Elfekih S, Tay WT, Gordon K, Court LN, De Barro PJ 2018b. Standardized molecular

diagnostic tool for the identification of cryptic species within the *Bemisia tabaci*

complex. Pest Manag Sci 74: 170-173. doi: 10.1002/ps.4676

FAOSTAT 2017 [cited 2018 Jan-2018]. Available from: http://www.fao.org/faostat/en/ -

data/QC/visualize

Fumagalli M, et al. 2013. Quantifying Population Genetic Differentiation from Next-

Generation Sequencing Data. Genetics 195: 979-+. doi: 10.1534/genetics.113.154740

Ghosh S, Bouvaine S, Maruthi MN 2015. Prevalence and genetic diversity of endosymbiotic

bacteria infecting cassava whiteflies in Africa. BMC Microbiology 15: 93. doi:

10.1186/s12866-015-0425-5

Ghosh S, Bouvaine S, Richardson SCW, Ghanim M, Maruthi MN 2018. Fitness costs

associated with infections of secondary endosymbionts in the cassava whitefly species

Bemisia tabaci. Journal of Pest Science 91: 17-28. doi: 10.1007/s10340-017-0910-8

Gottelli D, Marino J, Sillero-Zubiri C, Funk SM 2004. The effect of the last glacial age on

speciation and population genetic structure of the endangered Ethiopian wolf (*Canis*

simensis). Molecular Ecology 13: 2275-2286. doi: 10.1111/j.1365-294X.2004.02226.x

Hadjistylli M, Roderick GK, Gauthier N 2015. First report of the Sub-Saharan Africa 2

species of the *Bemisia tabaci* complex in the Southern France. Phytoparasitica 43: 679-

687. doi: 10.1007/s12600-015-0480-3

Hebert PDN, Cywinska A, Ball SL, DeWaard JR 2003. Biological identifications through

DNA barcodes. Proceedings of the Royal Society B-Biological Sciences 270: 313-321.

doi: 10.1098/rspb.2002.2218

Huhndorf MH, Peterhans JCK, Loew SS 2007. Comparative phylogeography of three

endemic rodents from the Albertine Rift, east central Africa. Molecular Ecology 16: 663-

674. doi: 10.1111/j.1365-294X.2007.03153.x

Katoh K, Standley DM 2013. MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. Mol Biol Evol 30: 772-780. doi:

10.1093/molbev/mst010

Korneliussen TS, Albrechtsen A, Nielsen R 2014. ANGSD: Analysis of Next Generation

Sequencing Data. Bmc Bioinformatics 15: 356. doi: 10.1186/s12859-014-0356-4

Kunz D, et al. 2019a. Draft mitochondrial DNA genome of a 1920 Barbados cryptic *Bemisia*

tabaci 'New World' species (Hemiptera: Aleyrodidae). Mitochondrial DNA Part B-

Resources 4: 1183-1184. doi: 10.1080/23802359.2019.1591197

Kunz D, Tay WT, Elfekih S, Gordon KHJ, De Barro PJ 2019b. Take out the rubbish -

Removing NUMTs and pseudogenes from the *Bemisia tabaci* cryptic species mtCOI

database. Sci Rep Submitted.

Lee W, Park J, Lee GS, Lee S, Akimoto S 2013. Taxonomic status of the *Bemisia tabaci*

complex (Hemiptera: Aleyrodidae) and reassessment of the number of its constituent

species. PLoS One 8: e63817. doi: 10.1371/journal.pone.0063817

Legg JP, French R, Rogan D, Okao-Okuja G, Brown JK 2002. A distinct *Bemisia tabaci*

(Gennadius) (Hemiptera : Sternorrhyncha : Aleyrodidae) genotype cluster is associated

with the epidemic of severe cassava mosaic virus disease in Uganda. Molecular Ecology

11: 1219-1229. doi: DOI 10.1046/j.1365-294X.2002.01514.x

Legg JP, et al. 2014. Spatio-temporal patterns of genetic change amongst populations of

cassava *Bemisia tabaci* whiteflies driving virus pandemics in East and Central Africa.

Virus Res 186: 61-75. doi: 10.1016/j.virusres.2013.11.018

Li H, Durbin R 2010. Fast and accurate long-read alignment with Burrows-Wheeler

transform. Bioinformatics 26: 589-595. doi: 10.1093/bioinformatics/btp698

Li H, et al. 2009. The Sequence Alignment/Map format and SAMtools. Bioinformatics 25:

2078-2079. doi: 10.1093/bioinformatics/btp352

Liu SS, et al. 2007. Asymmetric mating interactions drive widespread invasion and

displacement in a whitefly. Science 318: 1769-1772. doi: 10.1126/science.1149887

Macfadyen S, et al. 2018. Cassava whitefly, *Bemisia tabaci* (Gennadius) (Hemiptera:

Aleyrodidae) in East African farming landscapes: a review of the factors determining

abundance. Bull Entomol Res 108: 565-582. doi: 10.1017/S0007485318000032

Malka O, et al. 2018. Species-complex diversification and host-plant associations in *Bemisia*

tabaci: A plant-defence, detoxification perspective revealed by RNA-Seq analyses.

Molecular Ecology 27: 4241-4256. doi: 10.1111/mec.14865

Manani DM, Ateka EM, Nyanjom SRG, Boykin LM 2017. Phylogenetic Relationships

among Whiteflies in the *Bemisia tabaci* (Gennadius) Species Complex from Major

Cassava Growing Areas in Kenya. Insects 8: 25. doi: 10.3390/insects8010025

Martin JH, Mound LA 2007. An annotated check list of the world's whiteflies (Insecta :

Hemiptera : Aleyrodidae). Zootaxa: 1-84.

Matsubayashi KW, Ohshima I, Nosil P 2010. Ecological speciation in phytophagous insects.

Entomologia Experimentalis Et Applicata 134: 1-27. doi: 10.1111/j.1570-

7458.2009.00916.x

Minato N, et al. 2019. Surveillance for Sri Lankan cassava mosaic virus (SLCMV) in

Cambodia and Vietnam one year after its initial detection in a single plantation in 2015.

PLoS One 14: e0212780. doi: 10.1371/journal.pone.0212780

Mugerwa H, et al. 2018. African ancestry of New World, *Bemisia tabaci*-whitefly species.

Sci Rep 8: 2734. doi: 10.1038/s41598-018-20956-3

Paini DR, et al. 2016. Global threat to agriculture from invasive species. Proceedings of the

National Academy of Sciences of the United States of America 113: 7575-7579. doi:

10.1073/pnas.1602205113

Patil BL, Fauquet CM 2009. Cassava mosaic geminiviruses: actual knowledge and

perspectives. Mol Plant Pathol 10: 685-701. doi: 10.1111/j.1364-3703.2009.00559.x

Pickrell JK, Pritchard JK 2012. Inference of Population Splits and Mixtures from Genome-

Wide Allele Frequency Data. PLoS Genetics 8: e1002967. doi:

10.1371/journal.pgen.1002967

Polston JE, De Barro P, Boykin LM 2014. Transmission specificities of plant viruses with the

newly identified species of the *Bemisia tabaci* species complex. Pest Management

Science 70: 1547-1552. doi: 10.1002/ps.3738

Reich D, Thangaraj K, Patterson N, Price AL, Singh L 2009. Reconstructing Indian

population history. Nature 461: 489-U450. doi: 10.1038/nature08365

Stamatakis A 2006. RAxML-VI-HPC: Maximum likelihood-based phylogenetic analyses

with thousands of taxa and mixed models. Bioinformatics 22: 2688-2690. doi:

10.1093/bioinformatics/btl446

Tabachnick WJ 1996. *Culicoides variipennis* and bluetongue-virus epidemiology in the

United States. Annual Review of Entomology 41: 23-43. doi: DOI

10.1146/annurev.en.41.010196.000323

Tay WT, et al. 2017a. The Trouble with MEAM2: Implications of Pseudogenes on Species

Delimitation in the Globally Invasive *Bemisia tabaci* (Hemiptera: Aleyrodidae) Cryptic

Species Complex. Genome Biology and Evolution 9: 2732-2738. doi:

10.1093/gbe/evx173

Tay WT, et al. 2017b. Novel molecular approach to define pest species status and tritrophic

interactions from historical *Bemisia* specimens. Scientific Reports 7: 429. Doi:

10.1038/s41598-017-00528-7

Tay WT, Evans GA, Boykin LM, De Barro PJ 2012. Will the Real *Bemisia tabaci* Please

Stand Up? PLoS One 7: e50550. DOI: 10.1371/journal.pone.0050550

Vyskočilová S, Seal S, Colvin J 2019. Relative polyphagy of "Mediterranean" cryptic

Bemisia tabaci whitefly species and global pest status implications. Journal of Pest

Science 92: 1071-1088. doi: 10.1007/s10340-019-01113-9

Vyskočilová S, Tay WT, van Brunschot S, Seal S, Colvin J 2018. An integrative approach to

discovering cryptic species within the *Bemisia tabaci* whitefly species complex. Sci Rep

8: 10886. doi: 10.1038/s41598-018-29305-w

Walsh TK, et al. 2019. Mitochondrial DNA genomes of five major *Helicoverpa* pest species

from the Old and New Worlds (Lepidoptera: Noctuidae). Ecol Evol 9: 2933-2944. doi:

10.1002/ece3.4971

Wang HL, et al. 2016. First Report of Sri Lankan cassava mosaic virus Infecting Cassava in

Cambodia. Plant Disease 100: 1029-1029. doi: 10.1094/Pdis-10-15-1228-Pdn

Weir BS, Cockerham CC 1984. Estimating F-Statistics for the Analysis of Population-

Structure. Evol. 38: 1358-1370. doi: Doi 10.2307/2408641

Wosula EN, Chen WB, Fei ZJ, Legg JP 2017. Unravelling the Genetic Diversity among

Cassava *Bemisia tabaci* Whiteflies Using NextRAD Sequencing. Genome Biology and

Evolution 9: 2958-2973. doi: 10.1093/gbe/evx219

Zheng XW, et al. 2012. A high-performance computing toolset for relatedness and principal

component analysis of SNP data. Bioinformatics 28: 3326-3328. doi:

10.1093/bioinformatics/bts606