1

Genome-wide association study of musical beat synchronization demonstrates high polygenicity

Maria Niarchou[1,2*], Daniel E. Gustavson[1,2], J. Fah Sathirapongsasuti[3], Manuel Anglada-Tort[4], Else Eising[5],

Eamonn Bell[6,7], Evonne McArthur[1], Peter Straub[1], The 23andMe Research Team[3^], J. Devin

McAuley[8], John A. Capra[9], Fredrik Ullén[10,11], Nicole Creanza[12], Miriam A. Mosing[13,10,11], David Hinds[3], Lea

K. Davis[2,1,14,15,16*†], Nori Jacoby[4†], Reyna L. Gordon[17,1,18,19*†]

[1]Vanderbilt Genetics Institute, Vanderbilt University Medical Center, TN, USA
[2]Division of Genetic Medicine, Department of Medicine, Vanderbilt University Medical Center, TN, USA
[3]23andMe, Inc., Sunnyvale, CA, USA
[4]Computational Auditory Perception Research Group, Max Planck Institute for Empirical Aesthetics, Frankfurt,
Germany
[5]Department of Language and Genetics, Max Planck Institute for Psycholinguistics, Nijmegen, Netherlands
[6]Department of Music, Columbia University, New York, NY, USA
[7]Department of Computer Science, Durham University, Durham, UK
[8]Department of Psychology, Michigan State University, MI, USA
[9]Bakar Computational Health Sciences Institute and Department of Epidemiology & Biostatistics, University of
California, San Francisco, CA, USA
[10]Department of Neuroscience, Karolinska Institutet, Sweden
[11]Department of Cognitive Neuropsychology, Max Planck Institute for Empirical Aesthetics, Frankfurt, Germany
[12]Department of Biological Sciences, Vanderbilt University, Nashville, TN, USA
[13]Melbourne School of Psychological Sciences, University of Melbourne, Australia
[14]Department of Biomedical Informatics, Vanderbilt University Medical Center, TN, USA
[15]Department of Psychiatry and Behavioral Sciences, Vanderbilt University Medical Center, TN, USA
[16]Department of Molecular Physiology and Biophysics, Vanderbilt University, TN, USA
[17]Department of Otolaryngology – Head & Neck Surgery, Vanderbilt University Medical Center, TN, USA
[18]Department of Psychology, Vanderbilt University, TN, USA
[19]Vanderbilt Brain Institute, Vanderbilt University, Nashville, TN, USA
^full list of authors in Acknowledgements
31

32   *Corresponding authors: Reyna L. Gordon (reyna.gordon@alumni.usc.edu https://orcid.org/0000-

33 0003-1643-6979), Lea K. Davis (lea.k.davis@gmail.com https://orcid.org/0000-0001-5143-2282) and

34 Maria Niarchou (maria.niarchou@vumc.org https://orcid.org/0000-0002-8855-7393)

35   †Co-senior authors

Abstract

Moving in synchrony to the beat is a fundamental component of musicality. Here, we conducted a genome-wide association study (GWAS) to identify common genetic variants associated with beat synchronization in 606,825 individuals. Beat synchronization exhibited a highly polygenic architecture, with sixty-nine loci reaching genome-wide significance (p<5x10^{-8}) and SNP-based heritability (on the liability scale) of 13%-16%. Heritability was enriched for genes expressed in brain tissues, and for fetal and adult brain-specific gene regulatory elements, underscoring the role of central nervous system-expressed genes linked to the genetic basis of the trait. We performed validations of the self-report phenotype (through internet-based experiments) and of the GWAS (polygenic scores for beat synchronization were associated with patients algorithmically classified as musicians in medical records of a separate biobank). Genetic correlations with breathing function, motor function, processing speed, and chronotype suggest shared genetic architecture with beat synchronization and provide avenues for new phenotypic and genetic explorations.

Introduction

Our tendency to perceive, create, and appreciate rhythms in a variety of contexts (e.g., speech, music, movement) is a key feature of the human experience[1–3]. Rhythmic patterns provide predictable and robust sensorimotor structure to every-day interactions[4,5], helping guide our attention to communicatively important moments in time[6,7]. Even young children are sensitive to the social and linguistic signals carried by rhythm[8–10] and parents use rhythmic vocalizations and synchronous movement (e.g., lullabies and rocking) to interact with their infants from birth[11,12]. Rhythmic musical interactions are structured around the percept of a stable periodic pulse (termed the "beat" in Western music and present in music of most cultures[1,13], though its precise instantiation in musical structure varies cross-culturally[14,15]). While music in general and rhythmic structures in particular vary globally[15–17], there is evidence that hierarchical beat structure of most music is robust to cultural transmission[2] and indeed common in many types of music[1].

Beat perception and synchronization (i.e. perceiving, predicting, and moving predictively in synchrony to a musical beat[18]) is an important feature of musical experiences across many human cultures and musical genres[1,19]. The predictive temporal mechanisms afforded by beat structure enhance general perceptual and learning processes in music, including melody perception and production, singing, and joint music-making[3,6]. While some features of rhythm perception and production vary across listeners from different cultures[13,19–21], the same studies showed considerable consistencies across cultures for other features (e.g., preference for beat-based isochrony). Musicality (broadly encompassing musical behavior, music engagement and musical skill[22]) impacts society by supporting pro-social behavior[11,23] and well-being[24]. Many have proposed that beat perception and synchronization evolved in humans to support communication and group cohesion[18,22,25,26]. In modern humans, beat perception and synchronization are predictive of language and literacy skills[27,28] and are related to cognition, motor function, and social coordination[29]. Thus, the biology of beat synchronization has general importance for understanding human ability to perceive and predict natural rhythms, may have relevance for characterizing phenotypes such as developmental speech-language disorders which demonstrate associations with atypical rhythm[30], and may further elucidate mechanisms of rhythm-based rehabilitation (e.g., for stroke and Parkinson's[31]).

Neuroimaging findings highlight auditory-motor networks in the brain underlying beat perception and
production[32], during which there is precise entrainment of neural oscillatory activity to musical signals,
primarily involving motor planning areas and auditory regions of the brain, even during passive listening
to music[33,34]. Neural mechanisms of entrainment, prediction, and reward work in concert to coordinate
the timing of beat-related expectancies to musical signals during listening, playing, singing, and
dance[26,34]. The significant inter-individual variation of beat synchronization[35] are thought to be
influenced, in part, by common genetic variation; thus genetic approaches can be used to gain a
foothold on the biological basis of musicality and human rhythm traits.
89
Indeed, twin-modelling and other family-based studies point to moderate heritability of rhythm-related
traits such as duration discrimination[36,37], rhythm discrimination[38], isochronous sensori-motor
production[39], and off-beat detection[40]. Much less is known at the molecular level about human genome
variation underlying rhythm, and more generally musicality[41] which to date has been investigated in
relatively small samples[37], due to the challenge of assessing such phenotypes in samples large enough to
provide sufficient power to detect common variants with small effects (as expected for complex traits[42]).
Large-scale genome-wide association studies (GWASs) of rhythm-based traits (e.g. beat synchronization)
are thus needed to advance this field. Our understanding of the biological underpinnings of beat
synchronization, from its genetic architecture to its neural instantiation, behavioral manifestation, and
relationship to health, requires mechanistic multi-methodological approaches. Post-GWAS approaches
(i.e., enrichment of gene expression in central nervous system tissues and genetic correlations) can be
deployed to illuminate the relationship between the genetic and neural architecture of music-related
traits, and shared underlying biology with other health traits.
103
Here, we report a large-scale genome-wide interrogation of beat synchronization. Our approach was as
follows (Supplementary Figure 1): 1) We validated a subjective self-reported beat synchronization item
("Can you clap in time with a musical beat?", referred to in this paper as the "target question"), in
relation to measured beat synchronization and rhythm perception task performance. 2) We performed a
GWAS in N=606,825 to identify genomic loci associated with beat synchronization. 3) We further
investigated the genetic architecture of beat synchronization by estimating SNP-based heritability,
partitioned heritability, and conducting gene property and gene set enrichment analyses. Lastly, we
evaluated the contribution of genomic regions that have experienced significant human-specific
evolutionary shifts. 4) We then validated GWAS results by testing whether a cumulative sum of the
genetic effects for beat synchronization detected in our GWAS (i.e., polygenic score or PGS) was
significantly associated with algorithmically identified musical engagement in a separate sample. 5) We
explored shared genetic effects (pleiotropy) on beat synchronization and other traits through genetic
correlation and genomic structural equation analyses.

Results

Overview. Validating the self-reported beat synchronization phenotype

In light of prior work suggesting that musicality and rhythm skills are complex traits that can be
quantified with both objective (experiment-derived) assessment and subjective self-reported data[43,44],
we performed a series of validations of the GWAS target question (i.e., the self-report "Can you clap in
time with a musical beat?"), in relation to rhythm perception and beat production tasks. Both studies
were administered in English for consistency. We also explored the relationship between task-based
beat synchronization ability, a self-reported rhythm scale, and musicality. Study overviews and key
results are summarized in Figure 1.

126
Figure 1. Phenotype validation studies overview and results. A) Schema of internet-based phenotype validation studies. In phenotype
experiment #1, participants performed a musical rhythm perception test and provided self-report of the same target question in the GWAS
study ("Can you clap in time with a musical beat?"). In phenotype experiment #2, participants performed beat synchronization tasks (which
involved tapping to the beat of musical excerpts) as well as responding to the same target question, in addition to a series of other
questionnaires about their musical engagement/ability and health traits. B) Phenotype Experiment 1 results in N=724 show rhythm perception
task performance is correlated with Yes vs. No responses to GWAS target question, Odds Ratio (OR)=1.92, McFadden's R^2 = 0.06, p=0.002. C-H):
Phenotype Experiment 2 results. C) Beat synchronization task performance (n=542) is highly correlated with Yes vs. No responses to the target
question in OR=0.28, McFadden's R^2 =0.24, p<0.001; note that lower values of SD of the asynchrony correspond to more accurate tapping in
time to the musical beat. D) Beat synchronization task performance is correlated with responses to a similar self-report question asked on a
Likert scale, in n=542, r=−0.40, p<0.001. E) Self-reported rhythm questionnaire (seven-item scale in N=1,412) is correlated with responses to the
target question, McFadden's R^2 =0.34, p<0.001. F) Beat synchronization task performance is correlated with Self-reported rhythm questionnaire
in n=542, r=0.41, p<0.001. G) Gold-MSI (self-reported musical sophistication questionnaire) is correlated with responses to the target question
in N=1,412, OR=4.16, McFadden's R^2 = 0.18, p<0.001. H) Beat synchronization task performance is correlated with Gold-MSI in n=542, r=-0.36,
p<0.001. Within each plot for panels B,C,E and G, distributions are displayed using violin plots (mirrored density plot showing probability
density on the left), jittered individual data plots (right), and box plots in the center (horizontal line at median, lower and upper hinges
correspond to the first and third quartiles. The upper and lower whisker extends from the hinges to the value no further than 1.5 * interquartile
range from the hinge). Data beyond the end of the whiskers are called "outlying" points and are plotted individually. Panels D, F, and H
scatterplots are shown with dots colored by density to illustrate distribution. Taken together, these results show that self-reported beat
synchronization is a reasonable proxy of the trait.
146

Phenotype Experiment 1: Rhythm perception task performance.
In this experiment N=724 (see Table 1 for demographics) were asked the target question and performed
a musical rhythm perception test (Supplementary Figure 2). In each of the 32 trials of the task,
participants judged whether two rhythms were the same or different (see Figure 1A), following a
standard procedure for assessing musical perception ability[45] and utilizing rhythm sequences with
simple (highly metrical) and complex (syncopated) rhythms[46]. The rhythm perception task yielded
quantitative scores (*d'*). Individuals with better performance in the rhythm perception test (higher total
d') were more likely to answer Yes (vs. No) to the target question (OR=1.92, p=0.002, McFadden's
R^2=0.06, 95% CI=1.27,2.95; Figure 1B). All tests in both phenotype experiments were two-tailed.
156
Phenotype Experiment 2: Beat synchronization task performance
We then validated self-reported beat synchronization phenotype (N=1,412) as a proxy for directly-
measured beat synchronization ability. Participants (Table 1) completed a questionnaire on musicality,
health, and personality, and were asked to tap in real time to the beat of 4 different musical excerpts
(Supplementary Figure 3). Beat synchronization tapping accuracy was assessed similarly to lab-based
studies[35], but with a recently developed online-based technology that precisely measures asynchrony of
participants' taps along to music clips - i.e., REPP (Rhythm ExPeriment Platform[47]) for additional details
and pre-registered hypotheses (H1-H6), see Methods and Supplementary Notes. Key results of this study
are summarized in Figure 1 and Supplementary Table 1. Note that more accurate tapping is reflected in
lower tapping asynchrony scores, i.e., more accurate timing of taps in relation to the beat.
167
As predicted (OSF pre-registered H1), individuals who responded Yes to the target question ("Can you
clap in time with a musical beat") had lower tapping asynchrony, OR=0.28, p<.001, McFadden's R^2=.24,
95% CI=0.18,0.42 (Figure 1C). Tapping asynchrony was also negatively correlated with responses to a
highly similar item ("I can tap in time to a musical beat") when asked on a seven-point Likert agreement
scale (1= disagree; 7 = agree), *r*= -.40, p<.001, 95% CI=-0.47,-0.33] (H1a; Figure 1D). Similarly, individuals
with higher self-reported rhythmic ability (from another multi-item questionnaire) were much more
likely to respond "Yes" to the target question, OR=7.34, p<.001, McFadden's R^2=.34, 95% CI=4.90,11.52],
(Figure 1E), and demonstrate lower tapping asynchrony, r = -.41, p <.001, 95% CI=-0.47,-0.33] (Figure 1F)
(H2). Controlling for confidence judgments or confidence as a personality trait did not diminish the
associations between self-report and tapping asynchrony (H3; Supplementary Notes). Musical
Sophistication[43] was positively associated with the target question, OR=4.16, p <.001, McFadden's
R^2=.18, 95% CI=2.90,6.12 (Figure 1G) and negatively correlated with tapping asynchrony r= -.36, p <.001,
95% CI =-0.43,-0.28 (Figure 1H; H5). There was no credible evidence that Musical Sophistication or
prior/current musician status interacted with the tapping asynchrony to predict responses to the target
question (H6). All associations reported here were maintained when controlling for age, sex, and
education (Supplementary Table 1). Key analyses were repeated using vector length (variability of
relative phase of participants' tapping) as an outcome, and showed the same pattern of results as SD of
the asynchrony (Methods and Supplementary Notes). Taken together, results show that the self-
reported target question is a valid phenotype, and that other similar self-reported rhythm measures are
also valid proxies of beat synchronization.
188
189 **Beat Synchronization GWAS**.
190 *Genomic study population.* The discovery GWAS sample consisted of N=606,825 unrelated participants
191 of European ancestry (see Table 1 for demographics), who consented to participate in research with
192 23andMe, Inc. and answered Yes (91.57%) or No (8.43%) to the target question "Can you clap in time
193 with a musical beat?"
194

GWAS results and SNP-based heritability estimation. GWAS was conducted using logistic regression
under an additive genetic model, while adjusting for age, sex, the first five principal components from
genetic data, and genotype platforms (Methods). Seventy "sentinel" SNPs (after two rounds of LD
pruning, first at r^2=0.6 and then at r^2=0.1, kb = 250) at 69 genomic loci reached genome-wide
significance (p<5x10^{-8}; two-tailed; Figure 2, Table 2, and Supplementary Table 2), with a total of 6,160
SNPs passing the genome-wide significance threshold. Sixty-seven loci were autosomal and two were on
the X chromosome; locus 28 contains two independent sentinel SNPs. QQ-plot is provided in
Supplementary Figure 4, and local association plots at each locus are in the Regional Plots Supplemental
document. The LD score regression intercept was 1.02 (se=0.01) the ratio was 0.03, indicating that the
majority of inflation in test statistics was due to true polygenicity instead of population stratification.
205

206
Figure 2. Manhattan plot of GWAS results of beat synchronization. Results of GWAS in N=606,825 with 23andMe. The GWAS phenotype is
participants' responses of Yes (N=555,660) vs. No (N=51,165) to the question "Can you clap in time with a musical beat?". The GWAS controlled
for age, sex, top 5 PC's for ancestry, and genotype platform. The x-axis shows chromosomal position and the y-axis shows -log10 p-values).
Sixty-nine loci (70 sentinel SNPs, with one locus containing two independent sentinel SNPs) surpassed the threshold for genome-wide
significance of p<5x10^{-8} (dotted horizontal line). For illustration purposes, only 500,000 SNPs with p<0.1 are shown; gene symbols for sentinel
SNPs are notated when FUMA provided a gene mapped to nearest sentinel SNP.

213
214
The top-associated locus (rs848293) was mapped at chromosome 2 close to *VRK2 (*Vaccinia
Serine/Threonine Kinase 2 which codes for a protein kinase with multiple spliced isoforms expressed in
the brain) and *FANCL*, within a region previously linked to multiple neurological phenotypes[48,49]. Another
strongly associated locus at chromosome 17 (rs4792891) included the Microtubule Associated Protein
Tau (*MAPT*) gene, a Parkinson's disease[50] associated locus. The Mitogen-Activated Protein Kinase 3
(*MAPK3*) gene at 16p11.2, a region known to harbor rare variants which influence neurodevelopmental
disorders [51] and language-related phenotypes[52], was also strongly implicated. We also identified a locus
at Glycoprotein M6A (*GPM6A*), whose gene promoter contains a transcription factor binding site for
GATA2, a gene previously related to music phenotypes[37].

224
225 SNP-based heritability estimates on the liability scale[53] ranged from 13% to 16% when adjusted for a
226 range of estimated population prevalence for atypical beat synchronization (3.0% to 6.5%;

227   Supplementary Table 3; see Supplementary Notes for explanation of prevalence estimates). The
228   observed (unadjusted) genetic variance explained 5% (se=0.002) of the phenotypic variance.
229
Gene based GWAS. Gene-based genome-wide association analyses performed with MAGMA yielded 129
genes surpassing the threshold of $p<2.56x10^{-6}$ (two-tailed; Supplementary Table 4), with top two hits at:
CCSER1, in the 4q22 region in proximity to genes previously associated with multiple musicality
phenotypes[54], and *VRK2* (converging with the top locus in our SNP-based GWAS). Within these
associations, we examined potential replication of 29 genetic associations with musicality in humans
from prior reports[37,54,55]; none reached significance after genome-wide correction (Supplementary Table
5, Supplementary Notes), neither independently, nor as a gene-set (p=0.297).
237
In silico functional analyses.
Gene property and gene set enrichment analyses. To understand the biological functions and gene
expression associations of beat synchronization, we performed gene set analysis (GSA) and gene
property enrichment analyses[56] on the gene-based p-values, using MAGMA[57] implemented in FUMA[58].
Results of conditional gene property analysis (based on GTEx data tissue types[59] and controlling for
average expression) demonstrated that the genetic architecture of beat synchronization was
significantly enriched in genes expressed in brain tissues (Figure 3A), including cortex, cerebellum, and
basal ganglia (putamen, caudate and nucleus accumbens), converging with subcortical and cortical
regions supporting beat perception and synchronization[34].
247
248   To further examine potential biological functions associated with beat synchronization, we performed
249   exploratory GSA[57] (Supplementary Table 6). The genetic architecture of beat synchronization was
250   enriched for two gene sets associated with nervous system function: gene sets for synaptic membrane
251   adhesion ($p=1.01 \times 10^{-7}$) and synaptic adhesion-like molecules ($p=8.35x\ 10^{-7}$).
252
Partitioned Heritability. Complementing these gene-based enrichment analyses, we also performed
stratified LDSC[60] on the GWAS results to partition heritability according to genomic properties, using
specific functional categories to gain insight into the types of variation that contribute most to beat
synchronization. Among broad SNP annotation categories[61] (Supplementary Table 7), we found
enrichment (all $p<9.6x10^{-4}$) of: regions conserved in mammals (considered under purifying selection[62]),
regulatory regions marked by acetylation of histone H3 at lysine 9 (H3K9ac; a marker for active
chromatin, and monomethylation of histone H3 at lysine 4 (H3K4me1; a marker for enhancers),
supporting the hypothesis that identified associations may affect gene regulation. We next used LDSC-
Specifically Enriched Genes (LDSC-SEG[63]) to determine whether genes expressed in specific cell- or
tissue-types (conditional to the other annotations) are enriched for beat synchronization-associated
variants. For tissue-specific annotations of active chromatin and enhancers (marked by H3K9ac,
H3K27ac, DNase hypersensitivity sites and H3K4me1), heritability was enriched in central-nervous-
system- and skeletal muscle-specific regulatory regions (Supplementary Table 8). Cell-type specific,
multi-tissue chromatin, and multi-tissue gene expression results are shown in Supplementary Figures 5,
6 and 7 respectively. Enrichment in brain-specific regulatory elements, in multiple fetal and adult tissue-
specific elements as well as CNS-specific cell cultures, are shown in Figure 3B.
269
270
271

Figure 3. Genetic architecture of beat synchronization is enriched for brain-related expression. <u>A. Genes associated with beat synchronization are enriched for expression in brain tissue.</u> Results of MAGMA gene-property analysis are based on gene expression levels from GTEx v8, in 54 tissues, conditioned on average expression across tissues. Associations with beat synchronization were significantly enriched in brain-expressed genes (-log-10 p-values are on the y-axis, with tissue type on the x-axis). Dotted line shows p-value threshold for significant enrichment after Bonferroni correction. <u>B. Partitioned heritability shows enrichment in brain-specific regulatory regions of the genome.</u> Partitioned heritability analysis was performed with LDSC-SEG. Tissue-specific regulatory elements are marked by histone 3 acetylation or DNase hypersensitivity (for open chromatin) and H3K4me1 (for enhancers). Regulatory regions in adult brain tissues are shown in yellow, with regulatory elements in in

280   cell cultures in teal, and in fetal brain tissue shown in dark purple. The graph shows -log-10 p-values are on y-axis, with tissue and marker type
281   on x-axis. The dotted line shows p-value threshold for significant enrichment after Bonferroni correction for number of gene sets tested.
282
283

Evolutionary Analyses

Given evolutionary hypotheses about the origins of rhythm[4,18,64], we evaluated the contribution of regions of the human genome that have experienced significant human-specific shifts in evolutionary pressure, using stratified LDSC[53,60]. In particular, we analyzed the contribution to beat synchronization heritability from variants in genomic loci that are conserved across non-human species, but have elevated substitution rate on the human lineage[65]. Many of these human accelerated regions (HARs) play roles in human-enriched traits[66], including cognition[67]. Two variants significantly ($p < 5 \times 10^{-8}$) associated with beat synchronization (rs14316 at locus 66, rs1464791 at locus 20) fall within HARs. This is 11.2 times more overlap than expected by chance ($\mu = 0.178$ overlaps; p=0.017, based on 10,000 permutations). The rs1464791 variant is near *GBE1*, a gene associated with neuromuscular disease[68], reaction time[69] and cognitive impairments[70]. Applying LDSC to consider the full set of association statistics, we find that genetic variants in HARs contribute 2.26 times more to the observed heritability of beat synchronization than would be expected if heritability were distributed uniformly across variants ($p = 0.14$). Given the small number of common variants within HARs, this stratified heritability analysis is substantially underpowered (0.17% of variants considered are in HARs). The general agreement of these two approaches supports the enrichment of functional variation relevant to beat synchronization in HARs. We also evaluated the contribution of genetic variants detected in the Neanderthal genome to the heritability of beat synchronization (Supplementary Notes and Supplementary Table 9).

Polygenic scores (PGS) for beat synchronization are related to musicality reported in a health care context

We investigated whether the polygenic model of beat synchronization from the GWAS would differentiate self-identified musicians from non-musicians in a separate sample. Musicians (n=1,259) and matched controls (n=4,893) were drawn from a study[71] that algorithmically identified musically active patients (Methods and Supplementary Notes). PGS for beat synchronization were significantly associated with musical engagement (OR=1.33 per SD increase in PGS, p<2 x 10^{-16}, Nagelkerke's R^2=2%, 95% CI=1.25,1.42) consistent with beat synchronization capturing a dimension of musicality.

Cross-trait analyses.

Genetic correlations. To determine if beat synchronization shares genetic architecture with other traits, we tested genetic correlations[72] between beat synchronization GWAS and available GWAS of 64 traits classified into seven domains (Supplementary Table 10 and Supplementary Notes for details). There were 15 statistically significant genetic correlations (p<7.8 x 10^{-4}) (Figure 4, Supplementary Table 11). Results included positive correlations with motor function (grip strength and usual walking pace) and heaviness of smoking, and negative correlations with risk-taking and smoking initiation. There were two correlations with hearing traits (positive correlation with tinnitus and negative correlation with hearing difficulty). From the cognitive traits, processing speed (faster perceptual motor speed) was genetically correlated with beat synchronization, in addition to executive function - shifting (from a GWAS of trail-making, a task that involves complex processing speed). There were multiple associations with other biological rhythms*:* breathing function traits (positive associations with peak expiratory flow, forced expiratory volume, forced vital capacity, and a negative correlation with shortness of breath) and negative associations with sleep-related traits (insomnia and morning chronotype).

Figure 4. Cross-trait genetic correlations with beat synchronization. Results of exploratory genetic correlation analyses between beat synchronization and 64 traits from eight domains, conducted with LDSC. The x-axis is magnitude of genetic correlation (r_g) with standard error visualized, and the (uncorrected) p-values for each trait's correlation with beat synchronization are shown next to each trait label. Significant genetic correlations (after adjusting for multiple comparisons with a threshold of $p<7.8\times10^{-4}$) are shown with filled-in circles; empty circles are results that did not pass this threshold. See Supplementary Notes for detail on source studies. There are significant positive associations between beat synchronization and two of the cognitive domain GWASs; associations with smoking and risk-taking; two associations with hearing traits; two positive associations with motor function; and multiple associations with other biological rhythms (morning/evening chronotype, insomnia, and four breathing-related traits).

Genomic Structural Equation Modeling (SEM). We conducted Genomic SEM[73] to examine whether specific associations between beat synchronization and a subset of associated traits (e.g., musculoskeletal strength, walking pace, breathing function, and processing speed[74–76]) that are known to be related among each other in prior research[74–76] represent distinct genetic relationships or share a common set of genetic influences with beat synchronization. The best fitting model, displayed in Figure 5, included a common genetic factor that accounted for genetic correlations among beat synchronization, grip strength, processing speed, usual walking pace, and expiratory flow. This common factor explained 11.6% of total variance in the beat synchronization GWAS and 9.6-25.0% of the variance in the other GWASs (see Supplementary Notes).

347
Figure 5. Genomic SEM model of beat synchronization and rhythm-related traits. The best-fitting genomic structural equation model of beat
synchronization with GWASs of processing speed, grip strength, usual walking pace, and peak expiratory flow. 95% confidence intervals of
factor loadings and correlations are displayed in brackets. Results suggest that beat synchronization was associated with the other traits
through a set of common genetic influences. Model fit: χ^2 (4) = 10.85, p = .028, CFI = .983, SRMR = .017.

Common Factor GWAS: Rhythm-Related Traits. Using genomic SEM, we conducted a multivariate GWAS
(Supplementary Notes) on the latent genetic factor from the model presented above and portrayed in
Figure 5. The heritability of this latent genetic factor was 7.27% (s.e.=0.25%) and there were 130
genome-wide significant loci (Supplementary Table 12; Supplementary Figure 8). Heritability was
enriched for genes expressed in cerebellum (Supplementary Figure 9).

Cross-trait phenotypic extension of genetic correlations. Data from Phenotype Experiment 2 was
analyzed to examine whether a subset of genetic correlations would be reflected in true phenotypic
associations (pre-registered H4). Less accurate beat synchronization was weakly associated with a
morningness preference (r=-.10, p=.015), more shortness of breath (r=-.16, p<.001), and smoking 20 or
more (lifetime) cigarettes (r=-.11, p<.001) (Supplementary Table 13). In other words, accuracy in beat
synchronization was correlated with eveningness chronotype, reduced shortness of breath when
walking on level ground, and smoking abstinence (these associations go in the same direction as the
genetic correlations; moreover, these associations remained significant after controlling for age, sex and
education, and/or removing professional musicians from the sample.

367
368 *Additional sensitivity analyses*. Our results are robust to several potential biases (Supplementary Notes):
369 the GWAS beat synchronization results are not driven by 1) shared genetic effects with general cognitive
370 ability or educational attainment or 2) subtle residual population substructure, and 3) the *MAPT*
371 association is not confounded with Parkinson's Disease.

372
373
374

Discussion

Our GWAS revealed highly polygenic architecture of the human capacity to synchronize to a musical beat, representing a significant advancement of our understanding of the genomic basis of a musicality phenotype. Heritability of beat synchronization is enriched for functions of the central nervous system on a number of dimensions: SNPs involved in neural development and brain-specific regulatory regions of the genome; genes involved in synaptic function; and gene expression in cortical and subcortical brain tissues aligned with auditory-motor regions previously linked beat perception and synchronization[34]. Polygenic scores for beat synchronization were associated with self-identified musicians in a separate cohort, showing that the GWAS taps into the larger construct of musicality. Genetic correlations pointed to pleiotropy between beat synchronization and biological rhythms (including breathing function, walking pace, and chronotype), paving the way to a better understanding of the biological underpinnings of musicality and its health relevance.

In a series of phenotypic experiments, we also demonstrate that self-reported beat synchronization/rhythm measures can be used in large-scale population-based studies as suitable proxies for measuring individual differences in beat synchronization. Our findings indicate that the GWAS phenotype beat synchronization question was highly related to beat synchronization task performance (i.e., accuracy in tapping along to musical excerpts). Clearly the self-report is an imperfect correlate of beat synchronization; nevertheless, we demonstrate that it is a suitable proxy for very large-scale studies in which task administration is impractical. Furthermore, the GWAS phenotype is also significantly associated with: rhythm perception task performance[46], a multi-item Rhythm questionnaire, and a well-established assessment of musical sophistication[43]. These results also converge with earlier work showing shared variance among task performance of beat synchronization, rhythm perception, and musical engagement/training[44,77–80]. The phenotypic associations were robust to demographic factors and self-confidence, and were not driven by the presence of professional musicians in the sample. These phenotype validation studies represent critical groundwork (see[81]) enabling brief rhythm self-report questionnaires to be deployed online in large-scale population genetic cohorts.

With sixty-nine loci surpassing the threshold for genome-wide significance, the polygenic architecture of the beat synchronization GWAS aligns with expectations for complex traits[82,83]. The top-associated locus mapped to *VRK2*, a gene previously associated with behavioral and psychiatric phenotypes (i.e. depression[84], schizophrenia[85] and developmental delay[86]), suggesting a biological connection between beat synchronization and neurodevelopment. The SNP-based heritability of beat synchronization on the liability scale was moderate, ranging from 13 to 16%, similar to heritability estimates of other complex traits (e.g., chronotype GWAS[87]) and consistent with moderate heritability estimates of musical rhythm abilities reported in twin studies[38–40]. Still, the limitation of the heritability adjustment on the liability scale is that the exact population prevalence of atypical beat synchronization is unknown and had to be estimated based on other indices of rhythm (see Supplementary Notes); this limitation should be addressed in future work.

We examined potential mechanisms linking genetic variation to neural architecture of the beat synchronization trait using multiple in-silico enrichment methods. Results showed enrichment of the heritability of beat synchronization in many brain tissues including cerebellum, dorso-lateral prefrontal cortex, inferior temporal lobe, and basal ganglia nuclei (i.e., putamen, caudate, and nucleus accumbens). This pattern of results likely reflects a genetic contribution to subcortical-cortical networks underlying musical rhythm perception and production[32,34]; furthermore, enrichment of brain-tissue-specific enhancers and active-regulatory regions in tandem with gene expression enrichments in brain tissue

suggest that regions of the genome involved in regulation of gene expression within the beat perception
and synchronization network contribute to phenotypic variance. Moreover, the partitioning heritability
chromatin results showed an enrichment in both fetal and adult brain tissues, suggesting that beat
synchronization may be the result of neurodevelopmental or basic brain processes. Gene set
enrichments were also observed for synaptic function in the nervous system. Taken together, these
results are a building block towards understanding how genes influence neural processes during beat
perception and production, complementing results obtained with neuroimaging methods[88–93].
429
Insights about the evolution of rhythm traits are suggested by the occurrence of two of the beat-
synchronization-associated loci in human-accelerated regions (HARS) of the genome. In particular,
rs1464791 is an expression quantitative trait locus (eQTL) that regulates expression of *GBE1* in multiple
tissues, including muscle[59]; *GBE1* is also linked to neuromuscular disease[68] and reaction time[69]. HARs are
involved in many functions, so it is difficult to explicitly link their accelerated evolution to beat
synchronization. It is too early to tell whether the overlap between beat synchronization-associated loci
and those two HARS supports evolutionary theories about music (e.g., joint synchronous music-making
has been posited to exert selective pressures in early humans by enhancing group social cohesion and
family bonding[26,94]). The contribution of the genetic architecture of motor function to beat
synchronization is further suggested by enriched heritability of SNPs that are enhancers located in
musculoskeletal-tissue-specific regulatory regions of the genome, as well as our findings of genetic
correlations between walking pace, musculoskeletal strength, and beat synchronization.
442
Moreover, our findings are promising for future large-scale genomic interrogations using comprehensive
music phenotyping yielding continuous musicality variables (whether questionnaire-based[43,95] or
measured aptitude-based variables[38]), providing a path to examine potential genetic correlations
between beat synchronization and other musical traits, such as music training or pitch discrimination, in
line with family-based findings[36,37,41,96]. While the current data show a clear connection between the
beat synchronization and broader musicality at the phenotypic and genetic levels, further genomic
investigation in well-powered samples is needed to disentangle the *specificity* of genetic influences on
beat synchronization from other genetic influences on musical traits, or motor or auditory function.
Finally, although our GWAS was based on self-report, the magnitude of the sample size bolstered
statistical power. This is important because previous GWAS of other health traits based on self-report
have effectively replicated associations from other GWAS of deeper phenotypes[83], and it is generally
acknowledged that powerful sample size can overcome some of the limitations arising from modest
measurement error[97].
456
Moving in synchrony to a musical beat encompasses beat perception and extraction, motor periodicity,
meter perception, and auditory-motor entrainment (see [4,32,98] and Glossary in Supplementary Notes).
Despite this complexity, beat is a highly frequent feature of many musical systems[1,3,26]. Indeed, we
found that the heritability of beat synchronization is enriched in auditory-motor regions known to be
active during rhythm tasks.[34] It should be noted that beat perception and production does not depend
on musical training or music genre, and atypical beat synchronization is not linked to lack of music
exposure[99]. A limitation of the current work is the restriction of the genetic sample to a European
ancestry (due to GWAS methodology constraints); investigating beat synchronization, musicality, and
cross-trait correlations in populations of non-European ancestry should be a future priority for capturing
the spectra of musicality traits in a wider range of ethnic, cultural and socio-economic contexts (see[100]).
Regrettably, early research on individual differences in musicality in the early 1900's was pursued not
only using what we now recognize as highly culturally biased assessments, but also explicitly through the
lens of eugenics (see[101]), similar to early research on individual differences in cognition. We strongly

condemn the intent and design of those studies, and emphasize that the value of this work arises not
from the hypothetical ranking of interindividual differences in beat synchronization (indeed, genomic
associations with beat synchronization cannot be used to make deterministic predictions about
individual abilities or aptitudes[102, 103]). Rather the value arises from discovering that the shared
experience of rhythm, different though it is across cultures, is, in part, hardwired into our human
genome. Furthermore, new knowledge on the genetic basis of musicality must be used ethically and
fairly for research discovery and never for harm (e.g., discouraging a child from accessing musical
activities).
478
We replicated previous findings implicating location 4q22.1 in musicality-related traits[36,55] (*CCSER1* was
the top-associated gene in our MAGMA analysis), but did not find support for previous gene associations
from a set of genes that was drawn from prior candidate-gene, linkage, and GWAS studies with
relatively small samples[54]. This is potentially due to well-known methodological problems with these
methods particularly when applied to complex traits in small samples[104]. Without a second comparably
sized GWAS available within which to conduct replication of the loci discovered in the primary GWAS,
we were still able to demonstrate generalizability of these results by showing that PGS for beat
synchronization predicts a musical trait in a separate biobank sample. The GWAS results of beat
synchronization were nearly identical even after conditioning the results on GWASs of educational
attainment and general cognition (g-factor); these results align with twin findings of specific genetic
effects of rhythmic aptitude over and above any common genetic influences between rhythm and non-
verbal cognitive variables[39,105]. Moreover, given both the likely capturing of genetic variation related to
SES[106] by the educational attainment GWAS summary stats, and the observation that our beat
synchronization GWAS loci are robust to educational attainment, SES is unlikely to play a major role in
our findings.
494
Our cross-trait explorations revealed pleiotropic effects between beat synchronization and several
breathing-related phenotypes (peak expiratory flow, forced vital capacity, forced expiratory volume, and
shortness of breath). We demonstrated phenotypically that more accurate beat synchronization task
performance was related to lower likelihood of shortness of breath, mirroring the genetic correlations
between beat synchronization and breathing function. In light of our genetic correlation between beat
synchronization and three categories of traits (breathing, motor, and cognitive functions) previously
shown to be genetically interrelated during the aging process[74,75], we used genomic SEM to uncover
shared genetic variance among beat synchronization and enhanced breathing function, greater grip
strength, faster walking pace, and faster processing speed. Poor beat synchronization could be tied to
certain health risks during aging, in light of other genetic and epidemiological work showing that lung
function decline predicts later declines in motor function and psychomotor speed in older adults[107–110].
506
The genetic correlation results suggest that beat synchronization shares common biology with a
constellation of health traits, converging with the growing literature on the overlapping biomechanical
and perceptual mechanisms of rhythms harnessed during synchronization, communication, muscle
tensioning, and breathing; these relationships start very early in development[111,112]. The cerebellum in
particular plays important roles in the control of coordinated movement, balance, respiration, dance,
and even rhythm perception during passive listening to music[33]. Indeed, our rhythm-related traits multi-
variate GWAS demonstrated enriched heritability of genes expressed in Cerebellar tissue, potentially of
note in relation to experimental findings of functional synchronization of respiratory and upper limb
movements during vocalization[5]. Moreover, "beat gestures" in speech involve the cerebellum[113] and are
inextricably linked to respiration, upper limb movement, and postural control, all of which may be
biomechanically related to tapping or clapping to music.

518
Another dimension of biological chronometry was captured in the genetic correlation between
chronotype and beat synchronization, which we replicated phenotypically (individuals who self-
identified as 'evening people' tended to tap more accurately to music, even after removing professional
musicians from the analysis). These results complement recent evidence of the increased prevalence of
eveningness in musicians[114], indicating that the relationship between chronotype and musicianship
cannot solely be explained by environment (i.e., nocturnal job demands of professional musicians), but
that also other shared biological factors may play a role. Given the genetic correlation between beat
synchronization and lowered incidence of insomnia, the relationship between regulation of sleep,
musicality, and rhythm represents an area for further exploration.
528
Our GWAS effectively identified alleles at 69 separate loci differentially associated with typical vs.
atypical beat synchronization, complementing existing evidence of underlying neural
mechanisms[77,79,80,99]. Future genetic studies could study beat synchronization as a continuous trait,
either through self-report or internet-based task paradigms (i.e., REPP[47]). Prior literature on liability
threshold models has shown that case-control GWAS of complex traits yield similar results to those
obtained through continuous phenotypic measures (e.g., the genetic architecture of continuous
measures of psychiatric symptoms is highly similar to the genetic architecture of cases versus
controls[115]). Moreover, the use here of a population-based control group that is not "super-normal"
increases the likelihood that the genetic correlations that we uncovered are reliable and not biased
upward[116].
539
Taken together, our results advance knowledge of the biological basis of beat synchronization by
identifying genomic regions associated with individual differences in beat synchronization, estimating its
cumulative SNP-based heritability, successfully applying a polygenic score model in a separate genetic
sample, and exploring the enrichment of heritability in genes tied to central nervous system function.
Movement in synchrony with a musical beat is a fundamental feature of music, and sensitivity to the
beat emerges early in development, supporting childhood development in numerous ways[3,11,27,30] and
with importance over the lifespan[117]. We have elucidated the genetic architecture of beat
synchronization and revealed its health relevance through cross-trait analyses. This study also provides a
solid foundation for future exploration of how specific genetic variants contribute to neural mechanisms
of entrainment, prediction, and reward harnessed during musical interactions[118].
550

Methods
Phenotype validation studies
Phenotype Validation Experiment 1.
Overview. Phenotype Validation Experiment 1 was designed to determine if self-reported rhythm
abilities measured with the question used in the GWAS (i.e., 'Can you clap in time with a musical beat?')
would be associated with task-based rhythm perception performance. The study was conducted in
Amazon's Mechanical Turk and received ethical approval from the Columbia University Institutional
Review Board; participants gave their written informed consent, and the research complied with all
relevant ethical regulations. We selected the Beat-based Advantage paradigm as a rhythm
discrimination (perception) test due to its design of stimuli with simple and complex meter[119] and prior
history investigating individual differences in rhythm perception in a variety of brain and behavioural
studies in adults and children with typical and atypical development[46,120–122] as well as feasibility for

internet-based adaptation. A questionnaire (self-report questions) was administered prior to the
perception task, to avoid biasing participant self-report responses by how they perceived their own task
performance. See Supplementary Notes for additional details on procedure, compensation, and self-
report questionnaire.
567
Participants. The study sample was N=724 participants recruited anonymously in Amazon's Mechanical
Turk. All consented and passed a common headphone check[123] that guarantees good listening
conditions and the ability to follow basic instructions; this test also effectively filters out bots.
Participants (333 females; 387 males; 4 self-reported "other") were 18-73 years old (mean = 36.1 years,
SD=10.9) with 0-45 years of self-reported musical experience (mean 3.7 years, SD=5.7), representing an
average degree of musical experience (see norms in[43]); demographics are reported in Table 1 (note that
n=3 did not report their age).
575
Rhythm Perception Task. Stimuli for the rhythm perception task consisted of 32 rhythms drawn from
prior work[46,119]. For each participant, we randomized with probability of one half the occurrence of
"simple" rhythms (strong beat-based metrical structure and generally easier to discriminate) and
"complex" rhythms (weaker metrical structure due to syncopation and generally more challenging to
discriminate). Each rhythm was presented using pure tone stimuli in one of 6 frequencies (294, 353, 411,
470, 528, and 587 Hz, selected at random), and one of 4 durations (ISI of 220, 230, 240, and 250 ms).
Each trial consisted of 3 rhythms separated by 1500 ms of silence; there were 32 trials of the task. The
two first presentations were always identical, and in half of the trials (counterbalanced) the third rhythm
was also identical (standard condition); in the other half of the trials, the rhythm differed by having one
interval swapped (deviant condition). The pairings and structure of standard and deviant trials were
taken from[46]. Participants were instructed that in each trial, they would listen to the series of three
rhythms (the first two were always identical, and the third could be the same or different), and they had
to indicate if the third rhythm was the same or different (see Supplementary Figure 2). Additional
technical details are provided in the Supplementary Notes.
590
Data analysis.
Self-report. Responses to the target question were as follows: n=654 (90.3%) participants answered
'Yes', n=25 (3.5%) answered 'No' and n=45 (6.2%) answered "I'm not sure." Regarding an additional self-
report question 'Do you have a good sense of rhythm?', n=503(67%) answered 'Yes', 102(14%)
answered 'No' and n=117(16%) answered 'I don't know'. n=488 answered 'Yes' to both questions; the
tetrachoric correlation between these two self-report questions was *r*=0.73.
597
Rhythm perception test. Responses to the rhythm perception test were analysed using signal detection
theory[46,124]; this method is appropriate for discrimination tasks where the participant has to categorize
stimuli along some dimension with the resulting *d'* values the strength of detection of the signal relative
to noise. *d'* values were calculated on the 32 test trials. As expected from prior work[46,125], individuals
performed better at discriminating simple rhythms (mean d'= 1.98, SD =0.91) than complex rhythms
(mean d'=1.43, SD =0.97) (t(724)=11.11, *p*<0.001, Cohen's d=0.58).
604
To examine whether the target question was related to the objective (experimenter-measured)
performance on the rhythm perception test, we performed a logistic regression analysis in which the
clap-beat target question (Yes vs. No) was the outcome and quantitative scores on the rhythm
perception test (*d'* scores) were the predictor. Covariates included age, education, and sex. McFadden's
R^2 was also computed. We did not include 'I'm not sure' in the regressions, because this response was
not available for data analysis in the GWAS. Given that the simple rhythms have a strong metrical

structure that is known to facilitate detection and synchronization of the beat[46], we also tested whether
performance on the simple rhythm trials predicted self-reported beat synchronization (i.e., those who
responded Yes to the clap-beat question). See Supplementary Notes for additional analyses.
614
Phenotype Experiment 2.
Overview.
The aims of Phenotype Experiment 2 were two-fold: 1) to validate self-reported beat synchronization
phenotype as a proxy for objectively measured beat synchronization ability, and 2) to explore
phenotypic associations between rhythm/beat synchronization and assorted traits found to be
genetically correlated with beat synchronization. Phenotype Experiment 2 was pre-registered with Open
Science framework (https://osf.io/exr2t) on July 8, 2020, prior to data collection. This internet-based
study consisted of a beat synchronization task to assess the accuracy of participants' tapping in time
with musical excerpts, and a series of questionnaires assessing self-reported rhythm, musicality/music
engagement, selected health traits, confidence as a personality trait, and demographics. We used
REPP[47] to measure participants' tapping responses online with high temporal fidelity. The item from the
GWAS study, "Can you clap in time with a musical beat?" with possible responses: Yes/No/I'm not sure,
is referred to as the "target question."
628
We tested the following hypotheses: *H1*: Self-report responses to the target question will be correlated
with beat synchronization task performance (i.e., accuracy of tapping to the beat of music), such that
individuals who respond Yes to the "target question" are predicted to tap more accurately to the beat of
musical excerpts (i.e., they will have lower standard deviation of asynchrony than individuals who
respond No to the target question). *H1*a: Self-report on a highly similar self-report question ("I can tap in
time with a musical beat") with responses on a 7-point agreement Likert scale are predicted to be
correlated with tapping accuracy. *H2a:* The target question will be associated with broader rhythm
ability/engagement (measured with a rhythm scale from seven other self-report questions). *H2b:* Beat
synchronization task performance reflects broader self-reported rhythm ability/engagement. *H3:* To
examine whether confidence (either as a personality trait or sureness in one's own task performance)
affects the reliability of self-reported beat synchronization. *H4:* Selected traits found to be genetically
correlated with beat synchronization in the GWAS will be phenotypically correlated with beat
synchronization task performance and the Rhythm Scale. Specifically: better beat/rhythm is correlated
with evening chronotype (H4a), less shortness of breath (H4b), more tinnitus and loud music exposure
(H4c), and more smoking (H4d); and that these associations would survive controlling for age, sex, and
education (H4e). *H5.* Responses to the target question will be positively correlated with musical
engagement measured with the Gold-MSI. *H6*. The associations in H4 would interact with being a
musician, or more generally, with musical engagement.
647
Participants. A total of N=1,412 individuals met participation criteria outlined in the pre-registration
(including passing the attention check item and not abandoning the study before completion). The study
took place in Amazon Mechanical Turk and all participants provided informed consent in accordance
with the Max Planck Society Ethics Council's approved protocol; the research complied with all relevant
ethical regulations. Participants (728 females; 678 males; 6 prefer not answer) were 18-77 years old
(mean=36.3 years, SD=11.9) and had of 1-2 years of self-reported musical experience (Table 1). To
ensure that the tapping technology measured beat synchronization with high temporal fidelity, it was
crucial that participants complied with instructions to perform the tapping task (e.g., using the laptop
speakers instead of headphones, with minimal background noise, etc.), and also used hardware and
software without any technical issues that would preclude the recording signal (e.g., malfunctioning
speakers or microphones, or the use of strong noise cancellation technology; see[47]). Thus, several

precautions, including calibration tests and practice trials, were taken to make sure the tapping
technology would work effectively, excluding cases that did not meet the requirements (see
Supplementary Notes for details). A subset of n=542 had appropriate hardware to complete all parts of
the study (including the tapping tests). Questionnaires were administered in the full sample of
participants. Sample demographics are reported in Table 1. Demographics of the participants that
completed the tapping experiment was highly similar to the full sample, as shown in the table;
furthermore, 65.3% of the full sample and 64.9% of tapping sample had a Bachelor's degree or higher.
666
Data collection for Phenotype Experiment 2.
The first questionnaire included self-report items, including the "target question," and also covering a
variety of musical, health, and interest phenotypes. The health phenotype questions were chosen from
phenotypes (chronotype, smoking, shortness of breath, and tinnitus) found to be genetically correlated
with beat synchronization in our genetic analyses. Rhythm questions were selected for their particular
relevance to various aspects of interacting/engaging with musical rhythm. The order of the questions
was fixed for all participants. In addition, we used an attention check item[126] between item 10 and 11, in
order to exclude fraudulent responders, such as computer bots or disengaged participants responding
randomly to the experiments. The end-questionnaire consisted of items covering the following
additional self-report topics: another question about being a musician, a task confidence rating
question, a Confidence scale, a 16-item short version of the Gold-MSI[43] (items were chosen due to their
high reliability scores: reliability omega = 0.92), and a Demographic questionnaire. Questionnaire items
for Phenotype Experiment 2 are listed in the Appendix of the Supplementary Notes.
680
Tapping technology. Beat synchronization is particularly challenging to study with online research,
where variability in participants' hardware and software can introduce delay in latency and jitter into
the recorded time stamps[127,128]. Here we used REPP (see[47] for full details and a validation study of the
technology), a robust cross-platform solution for measuring sensorimotor synchronization in online
experiments that has high temporal fidelity and can work efficiently using hardware and software
available to most participants online. To address core issues related to latency and jitter, REPP uses a
free-field recording approach: specifically, the audio stimulus is played through the laptop speakers and
the original signal is simultaneously recorded with participants' tapping responses using the built-in
microphone. The resulting recording is then analyzed using signal processing techniques to extract and
align timing cues with high temporal accuracy.
691
Beat synchronization task. The beat synchronization task procedure consisted of three parts: calibration
tests, practice phase, and main tapping phase. Participants started with the calibration tests, including a
volume test to calibrate the volume of the laptop speakers to a level sufficient for detection by the
microphone, a background noise test to make sure participants were in a quiet environment, and a
tapping test to help participants practice how to tap on the surface of their laptop in the right level and
location to be detected by the microphone. Participants were then presented with the practice phase,
which consisted of four 15-second trials of isochronous tapping to a metronome beat (two with inter-
onset interval of 500 msec and two with inter-onset interval of 600 msec). Following the practice phase,
participants were presented with the main tapping task consisting of eight trials (4 musical excerpts,
each played twice), with each trial 30 seconds long. The order of presentation of the practice trials and
test trials was randomized for each participant.
703
The musical excerpts were drawn from the MIREX 2006 Audio Beat Tracking database in which musical
excerpts had been annotated for beat locations by 30 listeners who tapped along to the music[129]. We
chose these four MIREX clips that represent different music genres with different tempos and tapping

difficulty: track 1 ("You're the First, the Last, My Everything" by Barry White), track 3 ("El Contrapunto"
by Los Mensajeros de La Libertad), track 7 ("Le Sacre du Printemps" by Stravinsky), and track 19
("Possessed to Skate" by Suicidal Tendencies) of the MIREX training set (respectively). Based on the
annotations in[129], we identified the target beat locations from those consistently produced by the
annotators. Additional technical details are provided in the Supplementary Notes, and Supplementary
Figure 2 illustrates the instructions for participants.
713
Data Analysis.
Beat synchronization task performance: Tapping accuracy analysis
Let S_t and R_t be the stimulus and response onsets, respectively. In case of the metronome S_t are the
metronome onset (practice phase) and for music clips S_t is the target beat location based on the
annotations. We define the asynchrony as $a_t = R_t - R_t$. Based on prior work[130], we chose the standard
deviation of the asynchrony ($std(a_t)$) as our main target interest variable, as this appears to be a robust
measure of individual performance and tightly linked to musical abilities[131]. We used metronome onsets
to mark the beat metric level in an unambiguous way[132]. We emphasize that the metronome onsets
were only physically present during the beginning and end of each clip. We used only the participant-
produced asynchronies during the epoch at beats in which the guiding metronome was *not* present, in
order to test the ability of the participants to synchronize to music without the metronome sounds
(results were nearly identical when we included all onsets including the ones where physical metronome
onsets were present). For the main test scores, we used the asynchronies computed relative to the
virtual beat locations computed from prior human annotators in MIREX. We also computed vector
length in order to confirm key associations of interest between the target question and beat
synchronization accuracy (See Supplementary Notes).
730
Regression analyses
In accordance with the OSF preregistration, we examined whether responses to self-reported beat
synchronization phenotype were associated with objectively-measured tapping accuracy, other self-
reported measures of rhythm ability, confidence, and/or musical sophistication using logistic regression
and McFadden's R^2 (for H1, H2a, H3, and H5) and linear regression (for H1a and H2b). Likewise, we used
linear regression to examine potential replication of cross-trait associations uncovered by genetic
analyses (H4a-d), to examine whether musical background interacted with the above associations (H6).
Analyses were conducted in R version 3.5.1[133]. As described in our preregistration, individuals were
recruited using MTurk and were included unless they failed and attention check item or abandoned the
experiment before completing the study (N=1,412). Usable tapping data was available for n=542
individuals. The majority of exclusions were due to technical reasons detected by REPP's signal
processing pipeline during the practice trials (e.g., poor signal, noisy environment, wearing headphone,
issues with laptop microphone, or people not tapping at all), but some additional subjects (n=19) were
excluded for not having enough usable trials during data analysis. Missing covariates were handled using
pair-wise deletion. Exclusion criteria are detailed in the Supplementary Notes.
746
GWAS of beat synchronization.
Genome-wide association study summary statistics were generated from data acquired by personal
genetics company 23andMe, Inc. Phenotypic status was based on responses to an English-language
online questionnaire in which individuals self-reported "Yes" (cases) or "No" (controls) to the question
'Can you clap in time with a musical beat?'. Individuals who responded "I'm not sure" were excluded
from the genomic dataset as their data was not available. The GWAS included a total of 555,660 cases
and 51,165 controls (total N=606,825, mean age(SD)=52.09(18.5), prevalence=92%), unrelated
individuals of European ancestry; age range breakdown is provided in Table 1. All individuals provided

informed consent according to 23andMe's human subject protocol, which is reviewed and approved by
Ethical & Independent Review Services, a private institutional review board
(http://www.eandireview.com); the study complied with all relevant ethical regulations.
758
GWAS was conducted using logistic regression under an additive genetic model, while adjusting for age,
sex, the top five principal components estimated from genetic data in order to control for population
stratification, and indicators for genotype platforms to account for batch effects. We excluded SNPs with
Minor Allele Frequency (MAF) <0.01, low imputation quality (R^2<0.3) and indels, resulting in 8,288,850
SNPs in the GWAS summary statistics. Genotyping and QC details are provided in the Supplementary
Notes.
765
Post GWAS enrichment analyses
FUMA-based analyses. The FUMA[58] web application was used on the Genome-Wide Association
summary statistics to identify genomic loci along with the "sentinel" SNPs that were independent in our
analysis with a genome-wide significant P-value (<5 x 10^{-8}) that are in approximate linkage
disequilibrium (LD) with each other at r^2<0.1 and to generate Manhattan plots and Quantile-Quantile
plots. GWAS Catalogue associations for top loci were performed in FUMA (Supplementary Table 16).
772
Next, using the GWAS summary statistics as input for MAGMA (v1.08), we conducted a gene-based test
of association, a gene property enrichment test, and a gene-set enrichment analysis. Gene property
analysis[56] utilized GTEx v8 data integrated in FUMA, with gene expression values log2 transformed
average TPM per tissue type after winsorization at 50 based on GTEx RNA-seq data; this analysis was
performed for 54 tissue types where the result of gene analysis was tested for one side while
conditioning on average expression across all tissue types. We also performed exploratory GSA[57] in
FUMA using 15,556 Gene Ontology gene sets from the MsigDB database[134,135]; a Bonferroni threshold of
3.2x10^{-6} was used.
781
SNP-based heritability and partitioned heritability.
SNP-heritability was computed with LD Score regression software[60], and heritability estimates were
adjusted to the liability scale based on population prevalence of atypical beat synchronization of 3.0%-
6.5% (Supplementary Table 3, Supplementary Notes). We partitioned heritability of beat
synchronization by 52 broad functional categories (Supplementary Table 7), using stratified LD score
regression[60,63] (Bonferroni-corrected significance level of p=9.6x10^{-4}). We hypothesized that SNPs falling
into open chromatin regulatory regions (i.e., accessible to transcriptional machinery), and regions with
human-specific variation, would be enriched for beat synchronization-associated variation.
790
We further investigated (SNP-based) cell-type-specific and tissue-specific enrichments with LDSC-SEG
(LDSC Specifically Expressed Genes)[67], using a total of 697 gene sets (3 Cahoy gene sets, 205 Multi-tissue
gene expression sets and 489 Multi-tissue chromatin sets from the RoadMap Epigenomics and ENCODE
datasets); the Bonferroni-corrected significance level for this analysis was 7.1x10^{-5} (Supplementary Table
8). The X chromosome was not included in these analyses or any subsequent analyses using LDSC, given
that the file that is required for LDSC analysis (w_hm3_snplist) does not include chromosome X SNPs.
797
Evolutionary analyses.
The set of human accelerated regions (HARs) was taken from[65]. All variants in perfect LD (r^2 = 1.0 in 1000
Genomes European participants) with variants in HARs were considered in the analysis. Similarly,
variants tagging Neanderthal introgressed haplotypes were defined as in[136]. All variants in perfect LD
with a Neanderthal tag SNP were considered Neanderthal variants. For each set, we performed

stratified LDSC (v1.0.0) with European LD scores and the baseline LD-score annotations v2.1. The
heritability enrichment is defined as the proportion of heritability explained by SNPs in the annotation
divided by the proportion of SNPs in the annotation. Standard effect size (), which quantifies the effects
unique to the annotation, is the proportionate change in per-SNP heritability associated with a one
standard deviation increase in the value of the annotation, conditional on other annotations in the
baseline v2.1 model[62]. To determine the expected number of overlaps between the N loci significantly
associated with beat synchronization and HARs, we computed all overlaps between these sets of
genomic regions (in hg19 coordinates) using bedtools2[137]. We then randomly shuffled the locations of
HARs around the genome choosing segments of equal lengths and avoiding gaps in the genome
assembly. We repeated this process 10,000 times and for each iteration computed the number of
overlaps observed with the significantly associated loci. Based on this empirical distribution created with
no association between the region sets, we computed the enrichment and p-value for the observed
number of overlaps.
816
Genetic correlations. The genetic correlation method is designed to show whether there is shared
genetic variation linked to a particular trait (here, our beat synchronization trait) and traits measured in
other GWAS studies. We curated GWAS summary statistics for 64 complex traits representing a broad
range of phenotypic categories: cognitive, psychiatric, neuro-imaging/other neurological, motor, other
biological rhythms (circadian, heart, and breathing), overall health/well-being, and hearing (see
Supplementary Table 10 and Supplementary Notes for details of the source studies). We estimated
genetic correlations between beat synchronization and each of these traits using LDSC[72], with a
Bonferroni threshold of 7.5 x 10^{-4} (Supplementary Table 11).
825
Beat synchronization Polygenic Score (PGS) prediction of music engagement reported in health records
<u>Overview.</u> We examined whether beat synchronization polygenic scores (PGS) would be associated with
music engagement reported in health records. Individuals who disclosed music engagement to their
care providers (which was subsequently recorded by their provider) were drawn from a recent
phenome-wide study of 9,803 musicians[71] identified from keyword searches of patient electronic health
records (EHRs) in Vanderbilt University Medical Center's de-identified research database (Synthetic
Derivative). The phenotyping method was based on mining of clinical notes, utilizing 4 keywords and 449
regular expressions (i.e., "musician", "plays the piano"); see Supplementary Notes and[71] for details. The
method was then validated with manually conducted chart review, with a positive predictive value (PPV)
of 93%. Here we accessed the subset of n=1,259 musicians and 4,893 controls (matched for sex, median
age (across the patients' medical record), ethnicity, race, and length of record) that were also part of the
BioVU database and had genotyped data on file, to test the hypothesis that higher PGS for beat
synchronization would be associated with musical engagement operationalized as a having musician-
related keywords/regular expressions recorded in an individual's electronic health record.
840
We only selected individuals of European ancestry with genetic data that met standard quality control
thresholds due to the poor performance of PGS trained in individuals of one ancestry and applied to
individuals of another. This resulted in n=1,259 individuals (553 (44%) females, mean median age of
record (SD)=53.1(16.5)) as musician "cases" and 4,893 controls (1,963(40%) females, mean median age
of record (SD)=53.2(16.3)). See Supplementary Notes for details on the phenotyping, the samples,
genotyping, and QC.
847
<u>Polygenic scores.</u> We used an IBD filter of 0.2 in order to include only unrelated European samples of
BioVU. PGS were generated using the beat synchronization GWAS summary statistics, using software
PRS_CS[138]. Briefly, this method uses a Bayesian regression framework and places continuous shrinkage

(CS) prior on SNP effect sizes; this method outperforms previous methods in terms of prediction
accuracy especially when the training sample size is large[138], as is the case with the beat synchronization
GWAS. The 1000genomes European reference set was used. The PGS was standardized to have a mean
of 0 and SD of 1. Chromosome X was not included in the BioVU sample.
Data analysis. We conducted a logistic multivariable regression where the outcome variable was
musician vs. control, the predictor variable was PGS for beat synchronization, and covariates included
median age across the patients' medical record, sex, top 10 principal components estimated from BioVU
genetic data.
859
860 **Study FAQ:** A live FAQ for the study is at: https://www.vumc.org/music-cognition-lab/news/faq-about-
861 beat-synchronization-gwas-study

References

1. Savage, P. E., Brown, S., Sakai, E. & Currie, T. E. Statistical universals reveal the structures and functions of human
music. *Proc Natl Acad Sci U S A* **112**, 8987–8992 (2015).
2. Ravignani, A., Delgado, T. & Kirby, S. Musical evolution in the lab exhibits rhythmic universals. *Nat. Hum. Behav.* (2017)
doi:10.1038/s41562-016-0007.
3. Mehr, S. A. *et al.* Universality and diversity in human song. *Science (80-.).* (2019) doi:10.1126/science.aax0868.
4. Kotz, S. A., Ravignani, A. & Fitch, W. T. The Evolution of Rhythm Processing. *Trends in Cognitive Sciences* (2018)
doi:10.1016/j.tics.2018.08.002.
5. Pouw, W., Paxton, A., Harrison, S. J. & Dixon, J. A. Acoustic information about upper limb movement in voicing. *Proc.*
Natl. Acad. Sci. U. S. A. (2020) doi:10.1073/pnas.2004163117.
6. Large, E. W. & Jones, M. R. The dynamics of attending: How we track time varying events. *Psychol. Rev.* **106**, 119–159
(1999).
7. Nobre, A. C. & Van Ede, F. Anticipated moments: Temporal structure in attention. *Nature Reviews Neuroscience* (2018)
doi:10.1038/nrn.2017.141.
8. Hannon, E. E. & Trehub, S. E. Tuning in to musical rhythms: Infants learn more readily than adults. *Proc. Natl. Acad. Sci.*
U. S. A. (2005) doi:10.1073/pnas.0504254102.
9. Winkler, I., Haden, G. P., Ladinig, O., Sziller, I. & Honing, H. Newborn infants detect the beat in music. *Proc. Natl. Acad.*
Sci. U. S. A. **106**, 2468–2471 (2009).
10. Zentner, M. & Eerola, T. Rhythmic engagement with music in infancy. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 5768–5773
(2010).
11. Cirelli, L. K., Trehub, S. E. & Trainor, L. J. Rhythm and melody as social signals for infants. *Ann. N. Y. Acad. Sci.* **1423**, 66–
72 (2018).
12. Nazzi, T., Bertoncini, J. & Mehler, J. Language discrimination by newborns: toward an understanding of the role of
rhythm. *J Exp Psychol Hum Percept Perform* **24**, 756–766 (1998).
13. Polak, R. *et al.* Rhythmic Prototypes Across Cultures. *Music Percept. An Interdiscip. J.* (2018)
doi:10.1525/mp.2018.36.1.1.
14. London, J., Polak, R. & Jacoby, N. Rhythm histograms and musical meter: A corpus study of Malian percussion music.
Psychon. Bull. Rev. (2017) doi:10.3758/s13423-016-1093-7.
15. Clayton, M., Sager, R. & Will, U. In time with the music : the concept of entrainment and its significance for
ethnomusicology. *Eur. Meet. Ethnomusicol.* (2005).
16. Polak, R. & London, J. Timing and Meter in Mande Drumming from Mali. *Music Theory Online* (2014)
doi:10.30535/mto.20.1.1.
17. Polak, R., London, J. & Jacoby, N. Both isochronous and non-isochronous metrical subdivision afford precise and stable
ensemble entrainment: A corpus study of malian jembe drumming. *Front. Neurosci.* (2016)
doi:10.3389/fnins.2016.00285.
18. Patel, A. D. & Iversen, J. R. The evolutionary neuroscience of musical beat perception: the Action Simulation for
Auditory Prediction (ASAP) hypothesis. *Front. Syst. Neurosci.* **8**, 57 (2014).
19. Jacoby, N. & McDermott, J. H. Integer Ratio Priors on Musical Rhythm Revealed Cross-culturally by Iterated
Reproduction. *Curr. Biol.* (2017) doi:10.1016/j.cub.2016.12.031.
20. Cameron, D. J., Bentley, J. & Grahn, J. A. Cross-cultural influences on rhythm processing: Reproduction, discrimination,
and beat tapping. *Front. Psychol.* (2015) doi:10.3389/fpsyg.2015.00366.
21. Neuhoff, H., Polak, R. & Fischinger, T. Perception and evaluation of timing patterns in drum ensemble music from Mali.

904   *Music Perception* (2017) doi:10.1525/MP.2017.34.4.438.

905 22. Honing, H. On the biological basis of musicality. *Ann. N. Y. Acad. Sci.* (2018) doi:10.1111/nyas.13638.

906 23. Tarr, B., Slater, M. & Cohen, E. Synchrony and social connection in immersive Virtual Reality. *Sci. Rep.* (2018)
907   doi:10.1038/s41598-018-21765-4.

908 24. Lense, M. D. & Camarata, S. PRESS-Play: Musical Engagement as a Motivating Platform for Social Interaction and Social
909   Play in Young Children with ASD. *Music Sci.* (2020) doi:10.1177/2059204320933080.

910 25. Fitch, W. T. Empirical approaches to the study of language evolution. *Psychon. Bull. Rev.* **24**, 3–33 (2017).

911 26. Savage, P. E. *et al.* Music as a coevolved system for social bonding. *Behav. Brain Sci.* 1–42 (2020)
912   doi:10.1017/S0140525X20000333.

913 27. Woodruff Carr, K., White-Schwoch, T., Tierney, A. T., Strait, D. L. & Kraus, N. Beat synchronization predicts neural
914   speech encoding and reading readiness in preschoolers. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 14559–14564 (2014).

915 28. Swaminathan, S. & Schellenberg, E. G. Musical Ability, Music Training, and Language Ability in Childhood. *J. Exp.*
916   *Psychol. Learn. Mem. Cogn.* (2019) doi:10.1037/xlm0000798.

917 29. Keller, P. E., Novembre, G. & Hove, M. J. Rhythm in joint action: Psychological and neurophysiological mechanisms for
918   real-time interpersonal coordination. *Philosophical Transactions of the Royal Society B: Biological Sciences* (2014)
919   doi:10.1098/rstb.2013.0394.

920 30. Ladányi, E., Persici, V., Fiveash, A., Tillmann, B. & Gordon, R. L. Is atypical rhythm a risk factor for developmental
921   speech and language disorders? *WIREs Cogn. Sci.* **11**, (2020).

922 31. Moumdjian, L., Sarkamo, T., Leone, C., Leman, M. & Feys, P. Effectiveness of music-based interventions on motricity or
923   cognitive functioning in neurological populations: A systematic review. *European Journal of Physical and Rehabilitation*
924   *Medicine* (2017) doi:10.23736/S1973-9087.16.04429-4.

925 32. Merchant, H., Grahn, J., Trainor, L., Rohrmeier, M. & Fitch, W. T. Finding the beat: a neural perspective across humans
926   and non-human primates. *Philos Trans R Soc L. B Biol Sci* **370**, 20140093 (2015).

927 33. Gordon, C. L., Cobb, P. R. & Balasubramaniam, R. Recruitment of the motor system during music listening: An ALE
928   meta-analysis of fMRI data. *PLoS One* (2018) doi:10.1371/journal.pone.0207213.

929 34. Cannon, J. J. & Patel, A. D. How Beat Perception Co-opts Motor Neurophysiology. *Trends Cogn. Sci.* **25**, 137–150 (2021).

930 35. Dalla Bella, S. *et al.* BAASTA: Battery for the Assessment of Auditory Sensorimotor and Timing Abilities. *Behav. Res.*
931   *Methods* (2017) doi:10.3758/s13428-016-0773-6.

932 36. Pulli, K. *et al.* Genome-wide linkage scan for loci of musical aptitude in Finnish families: evidence for a major locus at
933   4q22. *J Med Genet* **45**, 451–456 (2008).

934 37. Oikkonen, J. *et al.* A genome-wide linkage and association study of musical aptitude identifies loci containing genes
935   related to inner ear development and neurocognitive functions. *Mol. Psychiatry* **20**, 275 (2014).

936 38. Ullén, F., Mosing, M. A., Holm, L., Eriksson, H. & Madison, G. Psychometric properties and heritability of a new online
937   test for musicality, the Swedish Musical Discrimination Test. *Pers. Individ. Dif.* **63**, 87–93 (2014).

938 39. Mosing, M. A., Verweij, K. J. H., Madison, G. & Ullén, F. The genetic architecture of correlations between perceptual
939   timing, motor timing, and intelligence. *Intelligence* **57**, 33–40 (2016).

940 40. Seesjärvi, E. *et al.* The Nature and Nurture of Melody: A Twin Study of Musical Pitch and Rhythm Perception. *Behav.*
941   *Genet.* (2016) doi:10.1007/s10519-015-9774-y.

942 41. Gingras, B., Honing, H., Peretz, I., Trainor, L. J. & Fisher, S. E. Defining the biological bases of individual differences in
943   musicality. *Philos. Trans. R. Soc. B Biol. Sci.* **370**, 20140092 (2015).

944 42. Wray, N. R., Goddard, M. E. & Visscher, P. M. Prediction of individual genetic risk of complex disease. *Curr. Opin. Genet.*
945   *Dev.* **18**, 257–263 (2008).

946 43. Müllensiefen, D., Gingras, B., Musil, J. & Stewart, L. The musicality of non-musicians: an index for assessing musical
947   sophistication in the general population. *PLoS One* **9**, e89642 (2014).

948 44. Musil, J. J., Iversen, J. R. & Müllensiefen. Measuring individual differences in musical beat alignment perception. *Pers.*
949   *Individ. Dif.* **60**, S35 (2014).

950 45. Law, L. N. C. & Zentner, M. Assessing musical abilities objectively: Construction and validation of the Profile of Music
951   Perception Skills. *PLoS One* **7**, e52508 (2012).

952 46. Grahn, J. A. & Brett, M. Rhythm and beat perception in motor areas of the brain. *J. Cogn. Neurosci.* **19**, 893–906 (2007).

953 47. Anglada-Tort, M., Harrison, P. & Jacoby, N. REPP: A robust cross-platform solution for online sensorimotor
954   synchronization experiments. *bioRxiv* (2021) doi:10.1101/2021.01.15.426897.

955 48. Li, M. & Yue, W. VRK2, a Candidate Gene for Psychiatric and Neurological Disorders. *Mol. Neuropsychiatry* **4**, 119–133
956   (2018).

957 49. Dashti, H. S. *et al.* Genome-wide association study identifies genetic loci for self-reported habitual sleep duration
958   supported by accelerometer-derived estimates. *Nat. Commun.* **10**, (2019).

959 50. Chang, D. *et al.* A meta-analysis of genome-wide association studies identifies 17 new Parkinson's disease risk loci. *Nat.*
960   *Genet.* (2017) doi:10.1038/ng.3955.

961 51. D'Angelo, D. *et al.* Defining the effect of the 16p11.2 duplication on cognition, behavior, and medical comorbidities.

JAMA Psychiatry (2016) doi:10.1001/jamapsychiatry.2015.2123.

52.   Hippolyte, L. *et al.* The Number of Genomic Copies at the 16p11.2 Locus Modulates Language, Verbal Memory, and
Inhibition. *Biol. Psychiatry* (2016) doi:10.1016/j.biopsych.2015.10.021.

53.   Bulik-Sullivan, B. K. *et al.* LD Score regression distinguishes confounding from polygenicity in genome-wide association
studies. *Nat Genet* **47**, 291–295 (2015).

54.   Oikkonen, J., Onkamo, P., Järvelä, I. & Kanduri, C. Convergent evidence for the molecular basis of musical traits. *Sci.*
Rep. **6**, 39707 (2016).

55.   Park, H. *et al.* Comprehensive genomic analyses associate UGT8 variants with musical ability in a Mongolian
population. *J Med Genet* **49**, 747–752 (2012).

56.   Leeuw, C. A. de, Stringer, S., Dekkers, I. A., Heskes, T. & Posthuma, D. Conditional and interaction gene-set analysis
reveals novel functional pathways for blood pressure. *Nat. Commun. 2018 91* **9**, 1–13 (2018).

57.   de Leeuw, C. A., Mooij, J. M., Heskes, T. & Posthuma, D. MAGMA: Generalized Gene-Set Analysis of GWAS Data. *PLoS*
Comput. Biol. (2015) doi:10.1371/journal.pcbi.1004219.

58.   Watanabe, K., Taskesen, E., van Bochoven, A. & Posthuma, D. Functional mapping and annotation of genetic
associations with FUMA. *Nat. Commun.* **8**, 1826 (2017).

59.   Consortium, Gte. The Genotype-Tissue Expression (GTEx) pilot analysis: multitissue gene regulation in humans. *Science*
(80-.). **348**, 648–660 (2015).

60.   Finucane, H. K. *et al.* Partitioning heritability by functional annotation using genome-wide association summary
statistics. *Nat. Genet.* **47**, 1228 (2015).

61.   Lindblad-Toh, K. *et al.* A high-resolution map of human evolutionary constraint using 29 mammals. *Nature* (2011)
doi:10.1038/nature10530.

62.   Hujoel, M. L. A., Gazal, S., Hormozdiari, F., van de Geijn, B. & Price, A. L. Disease Heritability Enrichment of Regulatory
Elements Is Concentrated in Elements with Ancient Sequence Age and Conserved Function across Species. *Am J Hum*
Genet **104**, 611–624 (2019).

63.   Finucane, H. K. *et al.* Heritability enrichment of specifically expressed genes identifies disease-relevant tissues and cell
types. *Nat. Genet.* (2018) doi:10.1038/s41588-018-0081-4.

64.   Mithen, S. J. *The Singing Neanderthals: The origins of music, language, mind, and body.* (Harvard University Press,
2005).

65.   Capra, J. A., Erwin, G. D., McKinsey, G., Rubenstein, J. L. & Pollard, K. S. Many human accelerated regions are
developmental enhancers. *Philos Trans R Soc L. B Biol Sci* **368**, 20130025 (2013).

66.   Hubisz, M. J. & Pollard, K. S. Exploring the genesis and functions of Human Accelerated Regions sheds light on their role
in human evolution. *Curr Opin Genet Dev* **29**, 15–21 (2014).

67.   Doan, R. N. *et al.* Mutations in Human Accelerated Regions Disrupt Cognition and Social Behavior. *Cell* **167**, 341-354
e12 (2016).

68.   Todd, E. J. *et al.* Next generation sequencing in a large cohort of patients presenting with neuromuscular disease
before or at birth. *Orphanet J. Rare Dis.* (2015) doi:10.1186/s13023-015-0364-0.

69.   Kichaev, G. *et al.* Leveraging Polygenic Functional Enrichment to Improve GWAS Power. *Am J Hum Genet* **104**, 65–75
(2019).

70.   Davies, G. *et al.* Study of 300,486 individuals identifies 148 independent genetic loci influencing general cognitive
function. *Nat. Commun.* **9**, 2098 (2018).

71.   Niarchou, M., Lin, G. T., Lense, M. D., Gordon, R. L. & Davis, L. K. Medical phenome of musicians: an investigation of
health records collected on 9803 musically active individuals. *Ann. N. Y. Acad. Sci.* nyas.14671 (2021)
doi:10.1111/NYAS.14671.

72.   Bulik-Sullivan, B. *et al.* An atlas of genetic correlations across human diseases and traits. *Nat. Genet.* **47**, 1236–1241
(2015).

73.   Grotzinger, A. D. *et al.* Genomic SEM Provides Insights into the Multivariate Genetic Architecture of Complex Traits
HHS Public Access. *Nat Hum Behav* (2019) doi:10.1038/s41562-019-0566-x.

74.   Shrine, N. *et al.* New genetic signals for lung function highlight pathways and chronic obstructive pulmonary disease
associations across multiple ancestries. *Nat. Genet.* (2019) doi:10.1038/s41588-018-0321-7.

75.   Willems, S. M. *et al.* Large-scale GWAS identifies multiple loci for hand grip strength providing biological insights into
muscular fitness. *Nat. Commun.* (2017) doi:10.1038/ncomms16015.

76.   Finkel, D., Ernsth-Bravell, M. & Pedersen, N. L. Temporal Dynamics of Motor Functioning and Cognitive Aging. *Journals*
Gerontol. - Ser. A Biol. Sci. Med. Sci. (2015) doi:10.1093/gerona/glv110.

77.   Bégel, V., Verga, L., Benoit, C. E., Kotz, S. A. & Dalla Bella, S. Test-retest reliability of the Battery for the Assessment of
Auditory Sensorimotor and Timing Abilities (BAASTA). *Ann. Phys. Rehabil. Med.* (2018)
doi:10.1016/j.rehab.2018.04.001.

78.   Bonacina, S., Krizman, J., White-Schwoch, T., Nicol, T. & Kraus, N. How Rhythmic Skills Relate and Develop in School-
Age Children. *Glob. Pediatr. Heal.* (2019) doi:10.1177/2333794x19852045.

79. Tranchant, P., Vuvan, D. T. & Peretz, I. Keeping the beat: A large sample study of bouncing and clapping to music. *PLoS*
One (2016) doi:10.1371/journal.pone.0160178.
80. Tranchant, P. & Peretz, I. Basic timekeeping deficit in the Beat-based Form of Congenital Amusia. *Sci. Rep.* (2020)
doi:10.1038/s41598-020-65034-9.
81. Coleman, J. R. I. The Validity of Brief Phenotyping in Population Biobanks for Psychiatric Genome-Wide Association
Studies on the Biobank Scale. *Complex Psychiatry* (2021) doi:10.1159/000516837.
82. Abdellaoui, A. & Verweij, K. J. H. Dissecting polygenic signals from genome-wide association studies on human
behaviour. *Nat. Hum. Behav. 2021 56* **5**, 686–694 (2021).
83. Watanabe, K. *et al.* A global overview of pleiotropy and genetic architecture in complex traits. *Nat. Genet.* **51**, 1339–
1348 (2019).
84. Nagel, M. *et al.* Meta-analysis of genome-wide association studies for neuroticism in 449,484 individuals identifies
novel genetic loci and pathways. *Nat Genet* **50**, 920–927 (2018).
85. Pardinas, A. F. *et al.* Common schizophrenia alleles are enriched in mutation-intolerant genes and in regions under
strong background selection. *Nat Genet* **50**, 381–389 (2018).
86. Lévy, J. *et al.* Molecular and clinical delineation of 2p15p16. 1 microdeletion syndrome. *Am. J. Med. Genet. Part A* **173**,
2081–2087 (2017).
87. Jones, S. E. *et al.* Genome-wide association analyses of chronotype in 697,828 individuals provides insights into
circadian rhythms. *Nat. Commun.* (2019) doi:10.1038/s41467-018-08259-7.
88. Grahn, J. A. & Rowe, J. B. Feeling the Beat: Premotor and Striatal Interactions in Musicians and Nonmusicians during
Beat Perception. *J. Neurosci.* **29**, 7540–7548 (2009).
89. Grahn, J. A. & Rowe, J. B. Finding and feeling the musical beat: striatal dissociations between detection and prediction
of regularity. *Cereb. Cortex* **23**, 913–921 (2013).
90. Kung, S.-J., Chen, J. L., Zatorre, R. J. & Penhune, V. B. Interacting cortical and basal ganglia networks underlying finding
and tapping to the musical beat. *J. Cogn. Neurosci.* **25**, 401–420 (2013).
91. Bengtsson, S. L. *et al.* Listening to rhythms activates motor and premotor cortices. *Cortex.* **45**, 62–71 (2009).
92. Teki, S., Grube, M., Kumar, S. & Griffiths, T. D. Distinct Neural Substrates of Duration-Based and Beat-Based Auditory
Timing. *J. Neurosci.* **31**, 3805–3812 (2011).
93. McAuley, J. D., Henry, M. J. & Tkach, J. Tempo mediates the involvement of motor areas in beat perception. *Ann. N. Y.*
Acad. Sci. **1252**, 77–84 (2012).
94. Dissanayake, E. If music is the food of love, what about survival and reproductive success? *Music. Sci.* **12**, 169–195
(2008).
95. Mas-Herrero, E., Marco-Pallares, J., Lorenzo-Seva, U., Zatorre, R. J. & Rodriguez-Fornells, A. Individual differences in
music reward experiences. *Music Percept. An Interdiscip. J.* **31**, 118–138 (2013).
96. Mosing, M. A., Pedersen, N. L., Madison, G. & Ullén, F. Genetic pleiotropy explains associations between musical
auditory discrimination and intelligence. *PLoS One* **9**, e113874 (2014).
97. Tung, J. Y. *et al.* Efficient replication of over 180 genetic associations with self-reported medical data. *PLoS One* **6**,
e23473 (2011).
98. Haegens, S. & Golumbic, E. Z. Rhythmic facilitation of sensory processing: a critical review. *Neurosci. Biobehav. Rev.* **86**,
150–165 (2018).
99. Sowiński, J. & Dalla Bella, S. Poor synchronization to the beat may result from deficient auditory-motor mapping.
Neuropsychologia **51**, 1952–1963 (2013).
100. Jacoby, N. *et al.* Cross-Cultural Work in Music Cognition. *Music Percept. An Interdiscip. J.* (2020)
doi:10.1525/mp.2020.37.3.185.
101. Devaney, J. Eugenics and Musical Talent: Exploring Carl Seashore's Work on Talent Testing and Performance. *Am.*
Music Rev. **48**, (2019).
102. Adam, D. The promise and peril of the new science of social genomics. *Nature* (2019) doi:10.1038/d41586-019-03171-
6.
103. Wray, N. R. *et al.* Research Review: Polygenic methods and their application to psychiatric traits. *Journal of Child*
Psychology and Psychiatry and Allied Disciplines (2014) doi:10.1111/jcpp.12295.
104. Border, R. *et al.* No support for historical candidate gene or candidate gene-by-interaction hypotheses for major
depression across multiple large samples. *Am. J. Psychiatry* **176**, 376–387 (2019).
105. Mosing, M. A., Madison, G., Pedersen, N. L. & Ullén, F. Investigating cognitive transfer within the framework of music
practice: Genetic pleiotropy rather than causality. *Dev. Sci.* **19**, 504–512 (2016).
106. Marees, A. T. *et al.* Genetic correlates of socio-economic status influence the pattern of shared heritability across
mental health traits. *Nat. Hum. Behav. 2021* 1–9 (2021) doi:10.1038/s41562-021-01053-4.
107. Emery, C. F., Finkel, D. & Pedersen, N. L. Pulmonary Function as a Cause of Cognitive Aging. *Psychol. Sci.* (2012)
doi:10.1177/0956797612439422.
108. Finkel, D., Ernsth Bravell, M. & Pedersen, N. L. Role of motor function and lung function in pathways to ageing and

decline. *Aging Clin. Exp. Res.* (2020) doi:10.1007/s40520-020-01494-3.

109.   Duggan, E. C. *et al.* A Multi-study Coordinated Meta-analysis of Pulmonary Function and Cognition in Aging. *Journals*
1080   *Gerontol. - Ser. A Biol. Sci. Med. Sci.* (2019) doi:10.1093/gerona/glz057.

1081   110.   Clouston, S. A. P. *et al.* The dynamic relationship between physical function and cognition in longitudinal aging cohorts.
1082   *Epidemiol. Rev.* (2013) doi:10.1093/epirev/mxs004.

1083   111.   Larsson, M., Richter, J. & Ravignani, A. Bipedal Steps in the Development of Rhythmic Behavior in Humans. *Music Sci.*
1084   (2019) doi:10.1177/2059204319892617.

1085   112.   Provasi, J., Anderson, D. I. & Barbu-Roth, M. Rhythm perception, production, and synchronization during the perinatal
1086   period. *Frontiers in Psychology* (2014) doi:10.3389/fpsyg.2014.01048.

1087   113.   Bernard, J. A., Millman, Z. B. & Mittal, V. A. Beat and metaphoric gestures are differentially associated with regional
1088   cerebellar and cortical volumes. *Hum. Brain Mapp.* (2015) doi:10.1002/hbm.22894.

1089   114.   Gjermunds, N., Brechan, I., Johnsen, S. Å. K. & Watten, R. G. Musicians: Larks, Owls or Hummingbirds? *J. Circadian*
1090   *Rhythms* **17**, 4 (2019).

1091   115.   Martin, J., Taylor, M. J. & Lichtenstein, P. Assessing the evidence for shared genetic risks across psychiatric disorders
1092   and traits. *Psychological Medicine* (2018) doi:10.1017/S0033291717003440.

1093   116.   KS, K., C, C. & SA, B. The impact on estimations of genetic correlations by the use of super-normal, unscreened, and
1094   family-history screened controls in genome wide case-control studies. *Genet. Epidemiol.* **44**, 283–289 (2020).

1095   117.   Mansens, D., Deeg, D. J. H. & Comijs, H. C. The association between singing and/or playing a musical instrument and
1096   cognitive functions in older adults. *Aging Ment. Heal.* (2018) doi:10.1080/13607863.2017.1328481.

1097   118.   Matthews, T. E., Witek, M. A. G., Lund, T., Vuust, P. & Penhune, V. B. The sensation of groove engages motor and
1098   reward networks. *Neuroimage* **214**, 116768 (2020).

1099   119.   Povel, D.-J. & Essens, P. Perception of temporal patterns. *Music Percept. An Interdiscip. J.* **2**, 411–440 (1985).

1100   120.   Grahn, J. A. & McAuley, J. D. Neural bases of individual differences in beat perception. *Neuroimage* **47**, 1894–1903
1101   (2009).

1102   121.   Gordon, R. L., Jacobs, M. S., Schuele, C. M. & Mcauley, J. D. *Perspectives on the rhythm-grammar link and its*
1103   *implications for typical and atypical language development*. *Annals of the New York Academy of Sciences* vol. 1337
1104   (2015).

1105   122.   Wieland, E. A., McAuley, J. D., Dilley, L. C. & Chang, S.-E. Evidence for a rhythm perception deficit in children who
1106   stutter. *Brain Lang.* **144**, 26–34 (2015).

1107   123.   Woods, K. J. P., Siegel, M. H., Traer, J. & McDermott, J. H. Headphone screening to facilitate web-based auditory
1108   experiments. *Attention, Perception, Psychophys.* **79**, 2064–2072 (2017).

1109   124.   Macmillan, N. A. & Creelman, C. D. *Detection theory: a user's guide*. (Cambridge University Press, 1991).

1110   125.   Gordon, R. L. *et al.* Musical rhythm discrimination explains individual differences in grammar skills in children. *Dev. Sci.*
1111   (2015) doi:10.1111/desc.12230.

1112   126.   Berinsky, A. J., Margolis, M. F. & Sances, M. W. Separating the shirkers from the workers? Making sure respondents
1113   pay attention on self-administered surveys. *Am. J. Pol. Sci.* (2014) doi:10.1111/ajps.12081.

1114   127.   Anwyl-Irvine, A. L., Dalmaijer, E. S., Hodges, N., & Evershed, J. Online Timing Accuracy and Precision: A comparison of
1115   platforms, browsers, and participant's devices. *PsyArXiv*.

1116   128.   Bridges, D., Pitiot, A., MacAskill, M. R. & Peirce, J. W. The timing mega-study: Comparing a range of experiment
1117   generators, both lab-based and online. *PeerJ* (2020) doi:10.7717/peerj.9414.

1118   129.   McKinney, M. F., Moelants, D., Davies, M. E. P. & Klapuri, A. Evaluation of audio beat tracking and music tempo
1119   extraction algorithms. *J. New Music Res.* (2007) doi:10.1080/09298210701653252.

1120   130.   Repp, B. H. Rate limits of on-beat and off-beat tapping with simple auditory rhythms: 1. Qualitative observations.
1121   *Music Percept.* (2005) doi:10.1525/mp.2005.22.3.479.

1122   131.   Repp, B. H. & Su, Y. H. Sensorimotor synchronization: A review of recent research (2006-2012). *Psychon. Bull. Rev.*
1123   (2013) doi:10.3758/s13423-012-0371-2.

1124   132.   London, J. *Hearing in Time: Psychological Aspects of Musical Meter*. *Hearing in Time: Psychological Aspects of Musical*
1125   *Meter* (2012). doi:10.1093/acprof:oso/9780199744374.001.0001.

1126   133.   3.5.1., R. D. C. T. A Language and Environment for Statistical Computing. *R Found. Stat. Comput.* (2018).

1127   134.   Ashburner, M. *et al.* Gene ontology: tool for the unification of biology. *Nat. Genet.* **25**, 25 (2000).

1128   135.   The Gene Ontology, C. The Gene Ontology Resource: 20 years and still GOing strong. *Nucleic Acids Res* **47**, D330–D338
1129   (2019).

1130   136.   Vernot, B. *et al.* Excavating Neandertal and Denisovan DNA from the genomes of Melanesian individuals. *Science (80-.*
1131   *).* **352**, 235–239 (2016).

1132   137.   Quinlan, A. R. & Hall, I. M. BEDTools: A flexible suite of utilities for comparing genomic features. *Bioinformatics* (2010)
1133   doi:10.1093/bioinformatics/btq033.

1134   138.   Ge, T., Chen, C.-Y., Ni, Y., Feng, Y.-C. A. & Smoller, J. W. Polygenic prediction via Bayesian regression and continuous
1135   shrinkage priors. *Nat. Commun.* **10**, 1776 (2019).

1136

Acknowledgements

We are grateful to 23andMe participants and employees for their contribution to the study, Nancy Cox and Simon Fisher for suggestions and insight throughout the process, Navya Thakkar for assistance with graphics, and Miriam Lense, Matthew D. Morrison, Aniruddh Patel, and Wim Pouw for thoughtful discussions. We would also like to thank Michaela Novakovic, Yune Lee, and Duane Watson for input during previous stages of the project, and Vanderbilt Trans-Institutional Programs for sparking the initial collaboration. RLG would like to thank Brandy Clark and P!nk for ongoing musical inspiration. Members of the 23andMe Research Team are: Michelle Agee, Stella Aslibekyan, Adam Auton, Robert K. Bell, Katarzyna Bryc, Sarah K. Clark, Sarah L. Elson, Kipper Fletez-Brant, Pierre Fontanillas, Nicholas A. Furlotte, Pooja M. Gandhi, Karl Heilbron, Barry Hicks, Karen E. Huber, Ethan M. Jewett, Yunxuan Jiang, Aaron Kleinman, Keng-Han Lin, Nadia K. Litterman, Jennifer C. McCreight, Matthew H. McIntyre, Kimberly F. McManus, Joanna L. Mountain, Sahar V. Mozaffari, Priyanka Nandakumar, Elizabeth S. Noblin, Carrie A.M. Northover, Jared O'Connell, Steven J. Pitts, G. David Poznik, Anjali J. Shastri, Janie F. Shelton, Suyash Shringarpure, Chao Tian, Joyce Y. Tung, Robert J. Tunney, Vladimir Vacic, and Xin Wang.

Research reported in this publication was supported by the National Institutes of Health Common Fund through the Office of the NIH Director and the National Institute on Deafness And Other Communication Disorders under awards DP2HD098859, K18DC017383, and R01DC016977. JAC was supported by the National Institutes of Health (R35GM127087). The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH. The dataset used for the analyses described were obtained from Vanderbilt University Medical Center's BioVU which is supported by numerous sources: institutional funding, private agencies, and federal grants. These include the NIH funded Shared Instrumentation Grant S10RR025141; and CTSA grants UL1TR002243, UL1TR000445, and UL1RR024975. Genomic data are also supported by investigator-led projects that include U01HG004798, R01NS032830, RC2GM092618, P50GM115305, U01HG006378, U19HL065962, R01HD074711; and additional funding sources listed at https://victr.vumc.org/biovu-funding/. Also, The Genotype-Tissue Expression (GTEx) Project was supported by the Common Fund of the NIH, and by NCI, NHGRI, NHLBI, NIDA, NIMH, and NINDS. The funders had no role in study design, data collection and analysis, decision to publish or preparation of the manuscript.

Author contributions

Conceptualization of study: RLG, LKD*; GWAS data acquisition and GWAS analyses: JFS,* 23andMe Research Team, DH, PS, MN*; GWAS and post-GWAS study design: LKD, RLG, JFS, MN, AJC, DH; Data visualization:* PS, MN, MAT, NJ, RLG, EE, DEG*; Post-GWAS analyses and interpretation: MN, DEG, PS, RLG, EE, EM, JAC, JFS, MM, FU, NC*; Phenotype studies design: RLG, JDM, NJ, DEG, MAT; Phenotype studies *data collection and analysis: MAT, NJ, DEG, EB, PS, MN, RLG, MM; Interpretation of results, writing, editing, and reviewing drafts:* All authors*; Project Supervision:* RLG, LKD, NJ, DH.

Competing interests

JFS, DH, and members of the 23andMe Research Team are employees of 23andMe, Inc., and hold stock or stock options in 23andMe. All other authors declare no competing interests.

Tables

Table 1. Sample demographics for each of the three study samples (GWAS and Phenotype Experiments 1 and 2).

GWAS Sample by Phenotype group (response to Clap-to-beat question)

	Cases (Yes)	Controls (No)
Total	555660	51165
Males	226188	23998
Females	329472	27167
18 to 30 years old	57898	5186
30 to 45 years old	135168	12909
45 to 60 years old	150939	13312
60 years old and over	211655	19758

Phenotype Validation Experiment 1 (rhythm perception)

Full sample who provided demographics	N	Mean Age in years (for N=722 who reported demographics)	SD Age
Total	722	36.08	10.90
Males	386	34.95	10.60
Females	332	37.49	11.07

Phenotype Validation Experiment 2 (beat synchronization and cross-trait phenotypic replication)

Full sample (questionnaires)	N	Mean Age in years	SD Age
Total	1412	36.34	11.93
Males	678	35.53	11.12
Females	728	37.15	12.61

Subset with valid tapping data	n	Mean Age in years	SD Age
Total	542	35.24	11.39
Males	241	35.02	10.62
Females	300	35.43	12.00

Table 2. Genomic loci and sentinel SNPs significantly associated with beat synchronization in the primary GWAS. Further details (e.g., chromosomal location) are provided in Supplementary Table 2.

Genomic Locus	Sentinel SNP	chr	A1	MAF	p-value	gene symbol
11	rs848293	2	G	0.42228	9.23E-18	VRK2
26	rs62340585	4	G	0.20695	1.81E-14	GPM6A
13	rs10168817	2	G	0.49299	1.94E-14	NA
20	rs10779987	3	T	0.38175	2.21E-14	GBE1
28	rs28392605	5	G	0.33904	8.93E-14	NA
45	rs1832909	9	T	0.40687	1.78E-13	NA
2	rs34762587	1	T	0.31379	2.25E-13	FOXO6
60	rs7542	16	G	0.46184	2.41E-13	MAPK3
5	rs10875125	1	C	0.15305	2.61E-13	DPYD
35	rs9400241	6	C	0.28851	4.49E-13	FOXO3
64	rs4792891	17	T	0.34013	7.07E-13	MAPT

39	rs1468701	7	G	0.29172	3.62E-12	*SND1*
50	rs10848650	12	G	0.42192	6.04E-12	*SLC6A13*
29	rs2635634	5	T	0.45317	9.54E-12	*CDH12*
67	rs9626920	22	G	0.41282	1.04E-11	*MIRLET7BHG*
16	rs764299	2	G	0.26719	1.47E-11	*PLEKHM3*
43	rs10984506	9	T	0.36558	1.66E-11	*ANP32B*
53	rs1426371	12	G	0.25919	1.67E-11	*WSCD2*
58	rs12913592	15	T	0.3596	6.13E-11	*NA*
6	rs72700870	1	G	0.14377	1.42E-10	*MCL1*
34	rs9388171	6	G	0.47595	2.16E-10	*NA*
55	rs6572878	14	T	0.39477	3.48E-10	*HAUS4*
4	rs11210206	1	T	0.31286	3.93E-10	*NA*
28	rs72633496	5	T	0.43224	6.21E-10	*NA*
10	rs7586405	2	G	0.30559	7.19E-10	*PPP1CB*
63	rs3024293	17	T	0.23528	8.26E-10	*C1QL1*
1	rs2061843	1	G	0.4001	1.19E-09	*CSMD2*
19	rs1349028	3	T	0.25977	1.54E-09	*EIF4E3*
25	rs4443239	4	T	0.2463	1.68E-09	*C4orf27*
33	rs1901739	5	T	0.47772	2.14E-09	*NA*
7	rs55678522	1	G	0.21629	2.25E-09	*LRRN2*
61	rs8079923	17	T	0.25309	2.88E-09	*AKAP10*
62	rs7501911	17	T	0.18191	3.34E-09	*NLK*
66	rs6087848	20	G	0.44304	3.40E-09	*POFUT1*
54	rs10744255	12	G	0.23229	4.24E-09	*NA*
31	rs13163173	5	C	0.16597	4.51E-09	*MEF2C*
3	rs2819333	1	T	0.37068	4.54E-09	*PTPRF*
51	rs2453873	12	G	0.22254	5.17E-09	*NA*
27	rs67264739	5	G	0.27395	5.54E-09	*ADCY2*
69	rs4898322	X	T	0.14076	5.90E-09	*NA*
56	rs2284901	14	G	0.37485	6.48E-09	*AKAP6*
32	rs1596431	5	T	0.19182	7.42E-09	*NA*
44	rs10978661	9	T	0.12006	7.74E-09	*ZNF462*
23	rs4263335	4	G	0.49483	8.74E-09	*JAKMIP1*
48	rs7939759	11	T	0.23981	1.23E-08	*CTSF*
65	rs9710427	19	G	0.41536	1.32E-08	*TECR*
21	rs12638746	3	G	0.33546	1.37E-08	*EPHA3*
59	rs12909047	15	G	0.48251	1.49E-08	*UBL7*
46	rs2505344	10	G	0.17674	1.51E-08	*EPC1*
24	rs67816799	4	C	0.38188	1.56E-08	*CCSER1*
15	rs10932201	2	G	0.46351	1.59E-08	*CREB1*
49	rs526904	11	T	0.34865	1.60E-08	*PICALM*
68	rs764935655	X	T	0.23454	1.83E-08	*NA*
9	rs6548147	2	T	0.4402	2.05E-08	*TSSC1*

52	rs10877461	12	G	0.29968	2.44E-08	NA
41	rs11996434	8	G	0.27037	2.61E-08	NA
40	rs1996148	8	G	0.31961	2.69E-08	PEBP4
47	rs10885458	10	G	0.28314	2.69E-08	NA
17	rs191373913	2	T	0.43899	2.74E-08	NGEF
38	rs12056186	7	C	0.42875	2.93E-08	ORC5
42	rs7856850	9	C	0.22184	3.07E-08	PTPRD
36	rs13197257	6	T	0.27444	3.23E-08	PTPRK
14	rs10497355	2	T	0.46078	3.43E-08	UBR3
12	rs11692449	2	T	0.37522	3.45E-08	XPO1
30	rs4704043	5	T	0.2827	3.65E-08	TNPO1
18	rs43182	3	T	0.13443	3.80E-08	PTPRG
57	rs62014217	15	G	0.20132	3.91E-08	HERC1
8	rs476141	1	T	0.49868	4.49E-08	NA
37	rs2849543	6	G	0.41591	4.60E-08	PARK2
22	rs571760466	3	C	0.27511	4.81E-08	LSAMP

Abbreviations: *SNP=Single Nucleotide Polymorphism, chr=Chromosome, A1=effect allele, MAF=Minor Allele Frequency, OR=Odds Ratio, Notes: Gene symbol is based on HUGO (HGNC). These are all genes annotated to SNPs in r2>0.1 with the lead SNP; Sentinel SNP in a given locus refers to independent SNP from FUMA. The SNPs were mapped to genes based on ANNOVAR annotation and on being physically located inside a Protein coding gene using 10kb window. NA=when the SNP is not within the 10kb window of a gene. For presentation reasons we only included one gene per SNP. For the full list of genes see Supplementary Table 2.*