**Idiosyncratic liver alterations of five frog species to land use changes in the**

**Brazilian Cerrado**

Lilian Franco-Belussi[1,5*]; Diogo B. Provete[2]; Rinneu Elias Borges[1,3]; Classius de

Oliveira[1]; Lia Raquel de Souza Santos[4]

[1]Department of Biology, São Paulo State University (UNESP), 15054-000, São José do

Rio Preto, São Paulo, Brazil.

[2] Setor de Ecologia, Instituto de Biociências, Universidade Federal de Mato Grosso do

Sul, Campo Grande, 79002-970, Mato Grosso do Sul, Brazil. E-mail:

diogo.provete@ufms.br.

[3] Departamento de Biologia, Universidade de Rio Verde (UniRV), Fazenda Fontes do

Saber, Rio Verde, Goiás, Brazil. E-mail:

[4] Instituto Federal de Educação Ciência e Tecnologia Goiano, Rio Verde, Goiás, Brazil;

* Corresponding author. E-mail: lilian.belussi@gmail.com

[5] Present address: Instituto de Biociências, Universidade Federal de Mato Grosso do Sul, Campo Grande, 79002-970, Mato Grosso do Sul, Brazil.

**\*Highlights (for review : 3 to 5 bullet points (maximum 85 characters including spaces per bullet point)**

- Changes in land use promote amphibian declines worldwide
- Our study focused on the biosphere, anthroposphere, and hydrosphere
- We used method to test for trait-environment relationship
- Species responded differently to land use change
- Land use change can affect internal morphology and physiology of organisms

Abstract

Changes in land use trigger environmental changes that can lead to decreased

biodiversity and species loss. The liver is an essential detoxification organ that reflects

systemic physiological responses to environmental changes. Here, we tested whether

land use changes influence the amount of substances from the hepatic cellular

catabolism and melanomacrophages of five anuran species in the Brazilian Cerrado. We

used routine histological and histochemical techniques. We then use recently developed

ecological methods to relate functional traits to environmental variables. There was an

increase in the amount of melanin in environments with high proportion of agriculture,

as well as variation in the amount of lipofuscin and hemosiderin. Therefore, the area of

melanomacrophages in the liver and the metabolic products in their cytoplasm can be

used as biomarkers of environmental contamination in regions with intense agricultural

activities. Our results add a new perspective to the influence of land use changes on

environmental health by highlighting the effect of environmental changes on internal

morphological aspects of animals.

**Keywords:** Melanomacrophages. Bioindicators. Environmental Stressors. Internal

pigmentation. Melanin. Liver metabolism.

1. **Introduction**

Human-driven land use changes are causing biodiversity loss around the world (Powers and Jetz, 2019). Brazil is one of the countries with the highest proportion of net loss of tree cover in South America, with a loss of 8% from 1982 to 2016 (Song et al., 2018). At the same time, there was a 12% increase in short vegetation cover (Song et al., 2018), which includes shrubs and herbaceous vegetation. This trend was especially significant in the Brazilian Cerrado, of which less of 2% are protected (Françoso et al., 2015; Damasco et al., 2018). Accordingly, more than half of the original 2 million km$^2$ of the Cerrado were transformed into planted pastures and annual cultures by 2005 (Klink and Machado, 2005). Central Brazil is a thriving region for industrial agricultural activities as one of the largest exporters of soybean and cattle meat in the world (Contini and Martha 2010, Reynolds et al., 2016). One of the consequences of land use change for export-oriented agricultural activities is the intensive use of agrochemicals (Schiesari et al., 2013; Aranha and Rocha, 2019). Therefore, land use changes, along with agrochemicals, are currently the main threats for biodiversity conservation and the sustainable use of natural resources in this biome (Reynolds et al., 2016).

Water quality in the Cerrado has been drastically affected by the intense use of fertilizers, with a significant increase in nitrogen and pesticides (Hunke et al., 2014). As a result, freshwater habitats in the Cerrado receive a great load of contaminants, which impact several aspects of aquatic biodiversity (De Marco et al., 2013; Bichsel et al., 2016). For example, there is evidence that the replacement of natural habitats by urban and agricultural areas decrease not only amphibian populations, but also their genetic diversity (Eterovick et al., 2016). Additionally, land use changes can promote rapid transformation in biological communities beyond species composition. Specifically, it can alter phenotypic aspects of several animal groups (e.g., Borges et al., 2019a), which

impact how species interact and adapt to a changing environment. These phenotypic changes include DNA damages (Borges et al., 2019b) and developmental abnormalities (Borges et al., 2019a). However, little is known about the effects of land use changes in internal phenotypic aspects of organisms inhabiting Neotropical agroecosystems.

Amphibians are good bioindicators of environmental quality because they show rapid responses to environmental stress (Brodeur and Candioti, 2017). In addition, they have permeable skin and eggs, making them vulnerable to aquatic contamination and infections. As such, these animals are useful for monitoring changes in both the aquatic and terrestrial environments because they depend on the two environments to complete their life cycle (Brodeur and Candioti, 2017). Amphibians are rapidly declining worldwide due to human-induced changes in the environment (Catenazzi, 2015; Alton and Franklin, 2017). Several factors are involved in this decline, including climate change, increased incidence of ultraviolet (UV) radiation due to ozone depletion, invasive species, environmental contamination, diseases, and habitat change or loss (Collins and Crump, 2009; Alton and Franklin, 2017).

Previous studies have analyzed the effects of agrochemicals on tadpole developmental abnormalities (Borges et al., 2019a) or genotoxic effects in adult anurans. However, environmental alterations that promote morphological and physiological changes at the tissue level are poorly understood. The liver plays a key role in the detoxification of xenobiotics (Pérez-Iglesias, et al., 2018; Fanali et al., 2018), as well as other functions related to the metabolism (Hinton et al., 2001; Fenoglio et al., 2005). The detoxification is performed by hepatocytes and melanomacrophages and may either enzymatic or not. Melanomacrophage centers and their pigments (melanin, lipofuscin, and hemosiderin) are involved in the hepatic response to various toxic

compounds. Thus, the liver is an important organ to evaluate the response of organisms to environmental pollutants (Fenoglio et al., 2005).

Melanomacrophages (MMs) are cells that occur in the hepatic tissue and produce and store melanin in their cytoplasm. The area and occurrence of melanomacrophages in the liver increase in response to environmental stressors, such as temperature (Santos et al., 2014), UV radiation (Franco-Belussi et al., 2016), and xenobiotics (Pérez-Iglesias et al., 2016). In addition to melanin, MMs contain catabolic substances originated from the degradation of red blood cells: hemosiderin and lipofuscin, originated from the degradation of polyunsaturated membrane lipids (Oliveira and Franco-Belussi 2012). Therefore, the density of melanomacrophages can be a useful morphological biomarker for the effect of environmental stressors (Oliveira et al., 2017). Morphofunctional changes in the hepatic parenchyma happen as a result of contaminants, suggesting the action of detoxification by melanomacrophages due to the processing action of some enzymes, besides the protective action of melanin (Fenoglio et al., 2005). Additionally, the functions of melanomacrophages are also related to the maintenance of homeostasis, regulating fibrosis, controlling basophiles and participating in the recycling of red blood cells (Gutierre et al. 2017).

Changes in the amount of catabolic substances within melanomacrophages may be associated with changes in phagocytic activity (Bucke et al., 1992; Fenoglio et al., 2005). For example, lipofuscin is produced as a result of the degradation of cellular components. Thus, its increase may hinder cell renewal and promote an accumulation of damaged cellular components in the tissue. The accumulation of lipofuscin can increase cellular sensitiveness to oxidative stress, since this molecule binds to metals, such as iron and copper (Terman and Brunk, 2004). Hemosiderin is an intermediate metabolite generated during the recycling of components in blood production. Thus, the

accumulation of hemosiderin indicates a disorder in blood cell recycling (Pérez-Iglesias

et al., 2016). In addition, environmental factors that alter the concentration of these

substances in the liver may contribute strongly to the decrease in animal health.

Consequently, this can affect individual fitness and promote population decline.

Therefore, analyzing liver cell physiology may be useful for detecting the effects of

environmental changes by human actions. However, while previous studies (Franco-

Belussi et al. 2013, Santos et al. 2014, Gregorio et al. 2016, Pérez-Iglesias et al. 2016)

have already analyzed the variation of melanin, lipofuscin, and hemosiderin in response

to climatic factors and xenobiotics in laboratory experiments using model species, no

study has investigated how these three substances vary at the same time in response to

changes in land use patterns in wild amphibian populations.

Here, we tested the influence of land use changes and aquatic contaminants on

liver cell morphology and physiology of five frog species. These can be hidden effects

of changes associated with agricultural intensification that are often neglected in

biodiversity assessments that only consider species abundance and occurrence. Aquatic

contaminants may alter organismal health that cannot be assessed by only recording

species presence in a given environment, since the internal morphology of individuals

can be damaged, which can compromise population viability and species persistence in

the environment in the long term.

**2. Material and methods**

*2.1. Study area and specimen sampling*

Field work was carried out in two regions: three ponds in the surroundings of

Rio Verde (17º48'11.28" S; 50º56'24.95" W), and two ponds within the Emas National

Park (18º15'32.11" S; 52º53'14.13" W; Fig. 1), Goiás, central Brazil. Samplings were

conducted during the breeding season between November and March for three consecutive years: 2013, 2014, and 2015. The region around the city of Rio Verde is a thriving agricultural area, mainly covered with planted pastures, and monocultures of soybean, corn, and sorghum. The Emas National Park is an enclave of Cerrado Protected Area with several typical vegetation types of this biome, varying from open formation, such as *campo limpo* and *campo rupestre*, to closed-canopy formations, such as *Cerradão* and Seasonal Deciduous Forest (Valente 2006).

We used a paired design in which we collected five adult males of five frog species in both regions: *Boana albopunctata, Dendropsophus minutus, Scinax fuscomarginatus, Leptodactylus fuscus,* and *Physalaemus cuvieri*. These species were used because they occurred in both regions. Additionally, all of them are widely distributed throughout South America (AmphibiaWeb, 2019) and seem to be generalists, occurring in a wide range of habitats, from open, natural formations, to bush Savannas, to peri-urban areas (Haddad et al. 2015). *Boana albopunctata*, *Scinax fuscomarginatus,* and *Dendropsophus minutus* are hylid tree-frogs that call perched on the vegetation, while *Leptodactylus fuscus* and *Physalaemus cuvieri* are medium-sized leptodactylids that are found calling and usually foraging on the ground near temporary ponds (Haddad et al. 2015). Both *B. albopunctata* and *D. minutus* deposit egg masses directly on water, while *L. fuscus* and *P. cuvieri* build foam nests in which they deposit eggs on the margin (in the case of *P. cuvieri*) or on subterranean chambers (in the case of *L. fuscus*; Haddad et al. 2015) of water bodies. Thus, adults, eggs, and larvae of these species have different degrees of contact with contaminated water. Consequently, these species can potentially be useful as environmental indicators, since they are highly abundant within their geographic range and seem to respond quickly to environmental disturbance.

*2.2. Water quality analysis*

To test if the ponds used by amphibians were contaminated by pesticides, we collected 1-L samples of water in one pond from each region. We quantified organochlorines and organophosphates in water samples from both sites. Quantification was made using standard methodology by A3Q Laboratory of Environmental Quality (Cascavel, Paraná, Brazil). Samplings from the Park did not contain any substance above the references, while Rio Verde had atrazine well above the level allowed by the Brazilian environmental agency (Table S1).

*2.3. Routine histological processing*

Specimens were anesthetized with 5 g/L benzocaine. This procedure was approved by the ethics committee on animal use of our university (protocol #0316, CEUA/UniRV). Liver fragments were extracted and fixed in metacarn solution for 3 h. Subsequently, they were dehydrated in an alcoholic series and included in historesin (Leica®). We obtained 2 μm sections in rotating microtome (RM 2265, Leica, Switzerland), which were stained with hematoxylin and eosin.

Histochemical analyzes were performed for the detection of lipofuscin as follows: sections were incubated for 15 min in Schmorl's solution, composed of 75 mL of 1% ferric chloride, 10 mL of potassium ferricyanide, and 15 mL of distilled water. Then, sections were immersed in neutral red aqueous solution at 1% followed by 1% aqueous solution of eosin. For the detection of hemosiderin, sections were incubated for 15 min in acid solution of ferrocyanide obtained by the dissolution of 2 g of potassium ferrocyanide in 100 mL of 0.75 mol/L hydrochloric acid solution. Subsequently they

were immersed in aqueous solution of 1% neutral red followed by aqueous solution of

1% eosin.

Images were captured in a digital camera coupled with a microscope (Laborana

Lab50AB-S) using an image capture system (Lab3001-C) and analyzed in Image Pro-

Plus v. 6.0 (Media-Cybernetics Inc.). Image analysis was conducted using 25

randomized histological fields for each animal. Specifically, we quantified the area

occupied by each substance by differences in staining intensity in 25 pictures per animal

following Santos et al. (2014).

*2.4. Quantification of land use pattern*

To quantify land use, we used a shape file with land cover and use classes

(IBGE, 2018 URL: http://www.sieg.go.gov.br/produtosIMB.asp?cod=4725). This file

classifies land use and cover into 14 classes based on satellite images from 2014 (see

Appendix II in IBGE 2018). To calculate the area of each land use, we established a

buffer of 500 m radius around each water body sampled and calculated the area

occupied by each land use class in ArcGIS 9.0 software (ESRI 2011). This buffer size

has been commonly used in studies of landscape ecology involving anurans (e.g.,

Almeida-Gomes et al. 2016), since it is usually the dispersal distance of individuals

moving among ponds in agricultural landscapes. The areas of each land use class were

then converted into proportions and used as predictor variables in further analyzes. The

land use types we found around our sampling sites were: natural grassland and shrub

vegetation, artificial area, forest mosaic, grassland mosaic, planted pastures, and natural

pasture, and agricultural area. Since the area of some land use types in our data set was

small, we combined artificial area with grassland mosaic into a class called anthropized

area, and also combined forest mosaic, natural pastures, and agricultural area into a

class called farming to improve data analysis and interpretation of results.

*2.5. Data analysis*

There are currently several methods to test the influence of environmental

variables on species traits (see review in Kleyer et al., 2012), the so-called fourth-corner

problem. However, there is still no consensus as to which method is best, and it seems

that this decision is dependent on the context and trait data type (continuous, categorical

or mixed), although studies show that correlating Community-weighted Mean with

environmental variables always seems to produce larger Type I Errors (Peres-Neto et

al., 2017; ter Braak et al., 2017). Here, we used a double-constrained Correspondence

Analysis (or dc-CA, for short) to test whether the mean area of melanin, hemosiderin,

and lipofuscin (response variables) of the five species are correlated with different

classes of land use (predictor variables). This is a long-proposed method (Lebreton et

al., 1988), but a new algorithm has only recently been proposed (ter Braak et al., 2018).

The advantages of dc-CA are that it takes into account the correlation among

environmental variables and among traits, when relating traits to environment (see ter

Braak et al., 2018), differently from RLQ and CWM-RDA (see Kleyer et al. 2012).

dc-CA uses three tables: a species-by-site matrix **Y**, which can contain either

abundances or incidence; a site-by-environment matrix **E**, and a species-by-trait **T**

matrix. The method starts by conducting a Correspondence Analysis constraining both

the columns (species) by species traits and row scores of the **Y** matrix by environmental

variables (ter Braak et al. 2018). We then produced a triplot showing the relationship

among traits, environmental variables, and species incidence in a single ordination

diagram. Analysis was conducted using R code available in the supplementary material

of ter Braak et al. (2018) and Canoco 5.12 (ter Braak and Šmilauer 2018).

**3. RESULTS**

We found both great intra- and interspecific variation in the amount of each

substance (Fig. 2). Most species had higher amounts of lipofuscin and melanin than

hemosiderin, except *Boana albopunctata* that had lower amounts of lipofuscin than the

other species. Interestingly, *Leptodactylus fuscus* and *Dendropsophus minutus* showed

little intraspecific variation in the amount of all three substances, while *Physalaemus*

*cuvieri* and *B. albopunctata* had high intraspecific variation. The means for the three

substances was similar for *L. fuscus* and *D. minutus*, whereas the other three species had

very distinct means.

The total inertia of the dc-CA model was 0.955. The first axis of dc-CA

accounted for 72% of the variation in the trait-environment relationship, while the

second accounted for 26%. The maximized fourth-corner correlation along the first and

second axes are 0.83 and 0.50, respectively. Melanin was positively correlated with

natural and planted pastures, but negatively correlated with man-made buildings and

agriculture (Table 1). Hemosiderin was negatively correlated natural and planted

pastures, and man-made buildings, but positively correlated with agriculture. The

correlation pattern of lipofuscin was almost identical to hemosiderin with small changes

in the strength of correlation with some land use types (Table 1).

The occurrence of *P. cuvieri* was positively correlated with the percentage of

planted pasture in the landscape (Fig. 3). This was also the species with the highest

amount of melanin in the liver (Fig 3), which was positively correlated (fourth-corner

correlation = 0.473, Table 1) with the percentage of planted pasture (Fig. 3). In contrast,

*L. fuscus* had the lowest amount of melanin (Fig. 3) whose incidence was negatively

correlated with the percentage of planted pasture in the landscape (Fig. 3).

Conversely, the incidence of *D. minutus* and *S. fuscomarginatus* were positively

correlated with area of agriculture and negatively with anthropogenic area, while the

incidence of *B. albopunctata* was positively correlated with anthropogenic area and

negatively with area of agriculture (Fig. 3). *Dendropsophus minutus* and *S.*

*fuscomarginatus* had higher amounts of hemosiderin and lipofuscin, suggesting higher

hepatic metabolism, which was positively correlated with agriculture and negatively

with percentage of anthropogenic area (Fig 3). Conversely, *B. albopunctata* had fewer

hepatic catabolite substances (Fig. 3).

**4. Discussion**

We found that *P. cuvieri* occurred in sites with planted pastures and man-made

buildings, had higher amounts of melanin, while the occurrence of *L. fuscus* was

positively correlated with man-made buildings and negatively correlated with planted

pastures. Additionally, *D. minutus* and *S. fuscomarginatus* had high amount, while *B.*

*albopunctata* had low amount of lipofuscin and hemosiderin and was positively

correlated with man-made buildings. Taken together, these results demonstrate the

effects environmental changes on melanomacrophages and hepatic metabolism of these

frog species.

The amount of melanin was highest in *P. cuvieri* and *B. albopunctata*, but low in

*L. fuscus*, *D. minutus,* and *S. fuscomarginatus*. There was also a high positive

correlation between melanin and planted pastures (0.473), and a less strong correlation

with man-made buildings (-0.033), and natural pastures (0.139). Melanin is a complex

polymer that absorbs and neutralizes free radicals, besides participating in the innate

immune response and protection of tissues in ectotherms (Cesarini 1996). Changes in

the amount of melanin was found in frogs experimentally exposed to several

xenobiotics. In these experiments, the variation in melanin was due to several

compounds (reviewed in Oliveira et al. 2017), such as polycyclic aromatic

hydrocarbons (PAHs; Fanali et al. 2018), herbicides (e.g., atrazine and glyphosate;

Pérez-Iglesias et al. 2019; Bach et al. 2019), drugs, and endocrine disrupters (Gregorio

et al. 2016). Here, we found large amounts of atrazine in the area under agriculture

influence (more than 5,000 time the limit accepted by Brazilian legislation, i.e.

5349.940 μg/L). Atrazine is an endocrine disruptor, which has immunotoxic and

immunosuppressive effects even at concentrations usually found in the environment

(i.e. < 500 μg/L; Rohr ad McCoy 2010). This substance can change swimming ability,

cause malformations, and promote nuclear alterations at higher concentrations in

tadpoles (Pérez-Iglesias et al. 2019). Atrazine causes oxidative stress in several tissues,

which can evolve to function loss.

Although our statistical model explained much variation in the three substances,

other climatic factors that have been changing as a consequence of human activities,

such as temperature and Ultra-violet radiation can also change the amount of melanin in

internal tissues of frogs (Franco-Belussi et al. 2017). Changes in these environmental

variables may promote changes in hepatic metabolism, such as increasing glycogen and

lipid levels (Mizell, 1965; Barni & Bernocchi, 1991; Fenoglio et al. 1992, Barni et al.

2002). Adaptation to natural conditions (i.e., hibernation) may promote an increase in

the amount of melanin in the liver by increasing the number of cells (i.e. hyperplasia) or

the increase in cell size (i.e. hypertrophy), besides an increase in the production of

melanin in cell interior (Barni et al. 2002). These changes occur as a mechanism of

metabolic defense against free radicals. For example, with increased storage of lipids

and glycogen in hepatic tissue due to environmental changes that occur naturally in

hibernating animals (Mizell, 1965; Barni & Bernocchi, 1991; Fenoglio et al. 1992,

Barni et al. 2002). Thus, the liver of anurans is a plastic organ, besides being sensitive

to the alterations of the natural biological rhythms (i.e. reproduction and hibernation),

coordinating these mechanisms to maintain the homeostasis of the organism during

adaptation to environmental changes (Barni et al. 2002). Therefore, cells that produce

and store melanin seem to be involved in the adaptation to environmental stressors at

the tissue level.

*Dendropsophus minutus* and *S. fuscomarginatus* had high amounts, whereas *B.*

*albopunctata* had low amounts of hemosiderin and lipofuscin. Lipofuscin and

hemosiderin are products of cellular catabolism and can be used to measure hepatic

metabolism, since both substances may be altered as a result of environmental stress

following habitat alteration (Santos et al. 2014, Pérez-Iglesias et al. 2016). Low

amounts of these substances indicate decreased phagocytic activity of cells (Bucke et al.

1992; Fenoglio et al. 2005), while accumulation of hemosiderin is related to increased

turnover of blood cells (Fenoglio et al. 2005). Therefore, the increase in recycling of

blood cells in *D. minutus* and *S. fuscomarginatus* indicates that biotic or abiotic factors

resulting from anthropic changes may be causing changes in the liver of these two

species, but so in *B. albopunctata*.

The idiosyncratic responses of species may be related to differences in life

history traits. For example, *Physalaemus* and *Leptodactylus* are terrestrial and can have

more contact with contaminants; while *Boana*, *Scinax,* and *Dendropsophus* are arboreal

and putatively less exposed to aquatic contaminants (Silva et al. 2013). However, it is

noteworthy that any drastic change in land use appears to promote metabolic alterations

related to liver physiological processes in anurans. Therefore, MMCs seem to be an

efficient biomarker that indicated alterations in the liver in response to land use change.

Actions for mitigating the negative effects of industrial agricultural activities must be

taken if we want to reach the goal of making environmentally responsible agricultural

products (ONU 2050 goals), especially in savanna biomes (Parr et al. 2014, Overback et

al. 2015).

Acknowledgements

Funding: This study was supported by Conselho Nacional de Desenvolvimento

Científico e Tecnológico (CNPq) (grant #477044/2013-1) to L.R.S.S and Fundação de

Amparo a Pesquisa do Estado de São Paulo (FAPESP) (grant #2013/02067-5). LFB was

supported by a FAPESP post-doctoral fellowship (#2014/00946-4) during the initial

phase of this study. B. Valverde, W. R. Rezende, R. A. Assis, and G. Leite helped with

histological procedures. R. F. Oliveira in the elaboration of the map with the animal

collection points. M. Almeida-Gomes helped with GIS data processing and extracting

area of each land use type. C. ter Braak clarified some statistical aspects of analysis. CO

has been continuously supported by CNPq (#305081/2015-2). The Instituto Chico

Mendes de Conservação da Biodiversidade provided colleting permit (Sisbio #34485-1)

and Emas National Park provided housing and authorization for field work.

**References**

Agius C, Agbede SA. 1984. An electron microscopical study on the genesis of

lipofuscin, melanin and haemosiderin in the haemopoietic tissues of fish. *Journal*

*Fish Biology*. 24,471-488.

Agius C. 1980. Phylogenetic development of melano-macrophage centers in fish.

*Journal Zoology*. 191,11-31.

Almeida- Gomes, M., Rocha, C. F. and Vieira, M. V. (2016), Local and Landscape

Factors Driving the Structure of Tropical Anuran Communities: Do Ephemeral

Ponds have a Nested Pattern? Biotropica, 48: 365-372. doi:10.1111/btp.12285

Aranha A, Rocha L. "Coquetel" com 27 agrotóxicos foi achado na água de 1 em cada 4

municípios. Reporter Brasil, São Paulo, 2019.

Barni S, Vaccarone R, Bertone V, Fraschini A, Bernini F, Fenoglio C. 2002.

Mechanisms of changes to the liver pigmentary component during the annual cycle

(activity and hibernation) of *Rana esculenta*. *Journal of Anatomy*. 200:185-194.

Barni, S., & Bernocchi, G. (1991). Internalization of erythrocytes into liver

parenchymal cells in naturally hibernating frogs (Rana esculenta L.). *Journal of*

*Experimental Zoology*, *258*(2), 143-150

Beuchle R, Grecchi RC, Shimabukuro YE, Seliger R, Eva HD, Sano E, et al. Land

cover changes in the Brazilian Cerrado and Caatinga biomes from 1990 to 2010

based on a systematic remote sensing sampling approach. Applied Geography 2015;

58: 116-127.

Bichsel D, De Marco P, Bispo AA, Ilg C, Dias-Silva K, Vieira TB, et al. Water quality

of rural ponds in the extensive agricultural landscape of the Cerrado (Brazil).

Limnology 2016; 17: 239-246.

Blaustein AR, Wake DB. Declining amphibian populations: A global phenomenon?

Trends in Ecology and Evolution 1990; 5: 203-204.

Borges, R. E., de Souza Santos, L. R., Assis, R. A., Benvindo-Souza, M., Franco-

Belussi, L., & de Oliveira, C. (2019a). Monitoring the morphological integrity of

neotropical anurans. *Environmental Science and Pollution Research*, *26*(3), 2623-2.

Borges, R. E.; Santos, L. R. S.; Benvindo-Souza, M.; Assis, R. A.; Modesto, R. S.;

Oliveira, C. (2019b) Genotoxic evaluation in tadpoles associated with agricultural in

the central cerrado, Brazil. Archives of Environmental Contamination And

Toxicology, v. 4, p. 1-7, 634.

Buikema AL, Niederlehner BR, Cairns J. 1982. Biological monitoring. Part IV –

Toxicity testing. *Water Research*. 16: 239–262.

Cesarini, J. P. (1996). Melanins and their possible roles through biological

evolution. *Advances in Space Research*, *18*(12), 35-40

Carneiro FF, Pignati W, Rigotto RM, Augusto LGS, Rizollo A, Muller NM, Alexandre

VP, Friedrich K, Mello MSC. 2012. Dossiê ABRASCO - Um alerta sobre os

impactos dos agrotóxicos na saúde. *ABRASCO*, Rio de Janeiro, abril de 2012. 1ª

Parte. 98p.

Carr J, Gentles A, Smith E, Goleman W, Urquidi L, Thuett K, Kendall RJ, Giesy JP,

Gross TS, Solomon KR, Van Der Kraak G. 2003. Response of larval *Xenopus laevis*

to atrazine: assessment of growth, metamorphosis, and gonadal and laryngeal

morphology. *Environmental Toxicology Chemistry*. 22: 396 – 405.

Chutter FM. 1998. Research on the rapid biological assessment of water quality impacts

in streams and rivers. *Water Research Commission Report*. 422.

Contini, E.; Martha, Jr., G.B. 2010. Brazilian agriculture, its productivity and change.

Bertebos Conference on "Food security and the futures of farms: 2020 and toward

2050". Falkenberg: Royal Swedish Academy of Agriculture and Forestry.

De Marco P, Nogueira DS, Correa CC, Vieira TB, Silva KD, Pinto NS, et al. Patterns in

the organization of Cerrado pond biodiversity in Brazilian pasture landscapes.

Hydrobiologia 2013; 723: 87-101.

De Oliveira, C., Franco-Belussi, L., Fanali, L. Z., & Santos, L. R. (2017). Use of

melanin-pigmented cells as a new tool to evaluate effects of agrochemicals and other

emerging contaminants in Brazilian anurans. *Ecotoxicology and Genotoxicology:*

*Non-traditional Terrestrial Models*, *32*, 125.

Dias LC, Pimenta FM, Santos AB, Costa MH, Ladle RJ. Patterns of land use,

extensification, and intensification of Brazilian agriculture. Glob Chang Biol 2016;

22: 2887-903.

Environmental Systems Research Institute (ESRI) (2011). ArcGIS Desktop: Release 9.

Redlands, CA: Environmental Systems Research Institute. Available at:

http://resources.arcgis.com/en/help/main/10.2/index.html.

Fanali, L. Z., Franco-Belussi, L., Bonini-Domingos, C. R., & de Oliveira, C. (2018).

Effects of benzo [a] pyrene on the blood and liver of Physalaemus cuvieri and

Leptodactylus fuscus (Anura: Leptodactylidae). *Environmental pollution*, *237*, 93-

102.

Fenoglio C, Boncompagni E, Fasola M, Gandini C, Comizzoli S, Milanesi G, Barni S.

2005. Effects of environmental pollution on the liver parenchymal cells and Kupffer-

melanomacrophagic cells of the frog *Rana esculenta*. *Ecotoxicology Environmental*.

Saf. 60, 259–268.

Fenoglio, C., Bernocchi, G., & Barni, S. (1992). Frog hepatocyte modifications induced

by seasonal variations: a morphological and cytochemical study. Tissue and Cell,

24(1), 17-29.

Franco-Belussi L, Oliveira C, 2011. Lipopolysaccharides induce changes in the visceral

pigmentation of *Eupemphix nattereri* (Anura: Leiuperidae). *Zoology.* 114:298-305.

Franco-Belussi L, Santos LRS, Zieri R, Oliveira C. 2011. Visceral pigmentation in

*Dendropsophus* (Anura: Hylidae): Occurrence and comparison. *Zoological*

*Anzeinger* 250, 102-110.

Franco-Belussi L, Santos LRS, Zieri R, Vicentini CA, Taboga SR, Oliveira C. 2012.

Liver Anatomy, Histochemistry, and Ultrastructure of *Eupemphix nattereri* (Anura:

Leiuperidae) During the Breeding Season. *Zoological Science*. 29: 844-848

Franco-Belussi, L., de Lauro Castrucci, A. M., & de Oliveira, C. (2013). Responses of

melanocytes and melanomacrophages of Eupemphix nattereri (Anura: Leiuperidae)

to Nle4, D-Phe7-α-melanocyte stimulating hormone and

lipopolysaccharides. *Zoology*, *116*(5), 316-324.

Franco- Belussi, L., Provete, D. B., & De Oliveira, C. (2017). Environmental correlates

of internal coloration in frogs vary throughout space and lineages. *Ecology and*

*evolution*, *7*(22), 9222-9233.

Gregorio, L. S., Franco-Belussi, L., Gomes, F. R., & de Oliveira, C. (2016). Flutamide

effects on morphology of reproductive organs and liver of Neotropical Anura,

*Rhinella schneideri*. *Aquatic Toxicology*, *176*, 181-189.

Griffiths R, Beebee T. 1992. Decline and fall of the amphibians. *New Scientist*. 134

(1827): 25–29.

Guillette LJ Jr, Gross TS, Gross D, Rooney AA, Percival HF. 1995. Gonadal

steroidogenesis in vitro from juvenile alligators obtained from contaminated and

control lakes. *Environmental Health Perspective.* 103(4):31-36

Gutierre, R. C., Jared, C., Antoniazzi, M. M., Coppi, A. A., & Egami, M. I. (2018).

Melanomacrophage functions in the liver of the caecilian *Siphonops annulatus*.

Journal of anatomy, 232(3), 497-508.

Halliday T. 2000. Do frogs make good canaries? *BioScience.*47: 143–146.

Hayes T, Haston K, Tsui M, Hoang A, Haeffele C, Vonk A. 2003. Atrazine-Induced

hermaphroditism at 0.1 ppb in american leopard frogs (*Rana pipiens*): laboratory and

field evidence. *Environmental Health Perspective.* 111:568-575.

Hayes TB, Case P, Chui S, Chung D, Haeffele C, Haston K, Lee M, Mai VP, Marjuoa

Y, Parker J, Tsui M. 2006. Pesticide mixtures, endocrine disruption, and amphibian

declines: are we underestimating the impact? *Environmental Health Perspective.*

114(suppl 1):40-50.

Hayes TB, Collins A, Lee M, Mendoza M, Noriega N, Stuart AA, Vonk A. 2002.

Hermaphroditic, demasculinized frogs after exposure to the herbicide atrazine at low

ecologically relevant doses. *Proceedings of the National Academy of Sciences of the*

*United State of America.* 99(8):5476-5480.

IBGE, 2018. Mudanças na cobertura e uso da terra do Brasil 2014 – 2016. Instituto

Brasileiro de Geografia e Estatística, Rio de Janeiro, Brasil. Disponível em:

https://www.ibge.gov.br/geociencias/informacoes-ambientais/cobertura-e-uso-da-

terra/15831-cobertura-e-uso-da-terra-do-brasil.html?=&t=downloads

Kleyer M, Dray S, Bello Fd, Leps J, Pakeman RJ, Strauss B, et al. Assessing species

and community functional responses to environmental gradients: which multivariate

methods? Journal of Vegetation Science 2012; 23: 805–821.

Klink CA, Machado RB. 2005. Conservation of the Brazilian Cerrado. *Conservation*

*Biology* 1(1): 148-152.

Kopp K, Signorelli L, Bastos RP. 2010. Distribuição temporal e diversidade de modos

reprodutivos de anfíbios anuros no Parque Nacional das Emas e entorno, estado de

Goiás, Brasil. *Iheringia*, Série Zoologica, Porto Alegre, 100(3):192-200.

LARSSON, B. S. (1993). Interaction between chemicals and melanin. Pigment Cell

Research, 6(3), 127-133.

Lebreton J, Chessel D, Richardot-Coulet M, Yoccoz N. L'analyse des relations espèces-

milieu par l'analyse canonique des correspondances. II. Variables de milieu

qualitatives. Acta Oecologia Generalis 1988; 9: 137–151.

McCoy KA, Bortnick LJ, Campbell CM, Hamlin HJ, Guillette Jr LJ, St Mary CM.

2008. Agriculture alters gonadal form and function in the toad *Bufo marinus*.

*Environmental Health Perspective.* 116(11):1526-1532.

Mizell, S. (1965). Seasonal changes in energy reserves in the common frog, Rana

pipiens. *Journal of Cellular and Comparative Physiology*, *66*(2), 251-258.

Muniz KPR, Giaretta AA, Silva WR, Facure KG. 2008. Auto-ecologia de *Hypsiboas*

*albopunctatus* (Anura, Hylidae) em área de Cerrado no sudeste do Brasil. *Iheringia*,

Sér. Zool. 98(2):254-259.

Newbold T, Hudson LN, Hill SL, Contu S, Lysenko I, Senior RA, et al. Global effects

of land use on local terrestrial biodiversity. Nature 2015; 520: 45-50.

Oliveira C, Franco-Belussi L. 2012. Melanic pigmentation in ectothermic vertebrates:

occurrence and function. In: Melanin: Biosynthesis, Functions and Health Effects,

editor Nova Publishers.pp.213-226.

Overbeck GE, Vélez-Martin E, Scarano FR, Lewinsohn TM, Fonseca CR, Meyer ST, et

al. Conservation in Brazil needs to include non-forest ecosystems. Diversity and

Distributions 2015; 21: 1455-1460.

Parr CL, Lehmann CE, Bond WJ, Hoffmann WA, Andersen AN. Tropical grassy

biomes: misunderstood, neglected, and under threat. Trends Ecol Evol 2014; 29:

205-13.

Passantino L, Santamaria N, Zupa R, Pousis C, Garofalo R, Cianciotta A, Jirillo E,

Acone F, Corriero A. 2013. Liver melanomacrophage centres as indicators of

Atlantic bluefin tuna, *Thunnus thynnus* L. well-being *Journal of Fish Diseases.* 2013

Peres-Neto PR, Dray S, ter Braak CJF. Linking trait variation to the environment:

critical issues with community-weighted mean correlation resolved by the fourth-

corner approach. Ecography 2017; 40: 806-816.

Pérez-Iglesias, J. M., Franco-Belussi, L., Moreno, L., Tripole, S., de Oliveira, C., &

Natale, G. S. (2016). Effects of glyphosate on hepatic tissue evaluating

melanomacrophages and erythrocytes responses in neotropical anuran Leptodactylus

latinasus. *Environmental Science and Pollution Research*, *23*(10), 9852-9861.

Pérez-Iglesias, J. M., Franco-Belussi, L., Natale, G. S., & de Oliveira, C. (2019).

Biomarkers at different levels of organisation after atrazine formulation (SIPTRAN

500SC®) exposure in Rhinella schineideri (Anura: Bufonidae) Neotropical

tadpoles. *Environmental pollution*, *244*, 733-746.

Phillips K. 1990. Where have all the frogs and toads gone? *BioScience.* 40: 422–424.

Powers, R. P., & Jetz, W. (2019). Global habitat loss and extinction risk of terrestrial

vertebrates under future land-use-change scenarios. *Nature Climate Change*, 1

Prado VH, Rossa-Feres DD. Multiple Determinants of Anuran Richness and

Occurrence in an Agricultural Region in South-Eastern Brazil. Environ Manage

2014.

R Core Team. 2010. R: a language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. Available at: http://www.R-

project.org (accessed on 20 august 2014).

Rohr, J. R., & McCoy, K. A. (2009). A qualitative meta-analysis reveals consistent

effects of atrazine on freshwater fish and amphibians. *Environmental health

perspectives*, *118*(1), 20-32.

Rohr J, Palmer B. 2005. Aquatic herbicide exposure increases salamander desiccation

risk eight months later in a terrestrial environment. *Environmental Toxicology

Chemistry.* 24: 1253 -1258.

Rosenberg DM, Resh VH. 1993. Freshwater Biomonitoring and benthic

macroinvertebrates, Chapman & Hall, NewYork.

Santos, L. R. S., Franco- Belussi, L., Zieri, R., Borges, R. E., & De Oliveira, C. (2014).

Effects of thermal stress on hepatic melanomacrophages of *Eupemphix nattereri*

(Anura). *The Anatomical Record*, *297*(5), 864-875.

Schiesari L, Waichman A, Brock T, Adams C, Grillitsch B. Pesticide use and

biodiversity conservation in the Amazonian agricultural frontier. Philos Trans R Soc

Lond B Biol Sci 2013; 368: 20120378.

Sichel G, Scalia M, Mondio F, Corsaro C. 1997. The amphibian Kupffer cells build and

demolish melanosomes: an ultrastructural point of view. *Pigment Cell Research*. 10,

271-287.

Silva ASVP, Loiola C, Pereira SRF, Santos RL, Andrade GV, Nunes GS. 2013.

Toxicidade aguda e genotoxicidade do agrotóxico comercial folisuper 600br a

girinos de *Physalaemus cuvieri* (Anura: Leiuperidae). Pesticidas: *Revista de*

*Ecotoxicologia e Meio Ambiente*. v. 23, p. 1-10.

Song XP, Hansen MC, Stehman SV, Potapov PV, Tyukavina A, Vermote EF, et al.

Global land change from 1982 to 2016. Nature 2018; 560: 639-643.

Strassburg BBN, Brooks T, Feltran-Barbieri R, Iribarrem A, Crouzeilles R, Loyola R, et

al. Moment of truth for the Cerrado hotspot. Nat Ecol Evol 2017; 1: 99.

Tavera-Mendoza L, Ruby S, Brousseau P, Fournier M, Cyr D, Marcogliese D. 2002.

Response of the amphibian tadpole (*Xenopus laevis*) to atrazine during sexual

differentiation of the testis. *Environmental Toxicology Chemistry*. 21: 527 - 531.

Van der Heever JA, Grobbelaar JU. 1995 The use of algae in bioassays to detect the

presence of toxic compounds in natural waters. *Water Research Commission Report*.

393.

Ter Braak CJ, Peres-Neto P, Dray S. A critical issue in model-based inference for

studying trait-based community assembly and a solution. PeerJ 2017; 5: e2885.

ter Braak CJF, Šmilauer P, Dray S. Algorithms and biplots for double constrained

correspondence analysis. Environmental and Ecological Statistics 2018; 25: 171-197.

ter Braak, C. J., & Šmilauer, P. (2018). Canoco reference manual and user's guide:

software for ordination (version 5.10). Biometris, Wageningen University &

Research.

Van Vuren JHJ, Du Preez HH, Deacon AR. 1994. Effect of pollutants on the physiology

of fish in the Olifants River (Eastern Transvaal). *Water Research Commission*

*Report.* 350.

Zuasti A, Jiménez-Cervantes C, García-Borrón JC, Ferrer C. 1998. The melanogenic

system of *Xenopus laevis*. *Archives of Histology and Cytology.* 61:305-316.

Figure legends

Fig. 1. Map showing the location of sampling sites in Central Brazil.

Fig. 2. Ternary plot showing the relative proportion of the area of each substance in the

five frog species. Small points indicate individual measurements, while the large dot

represents the mean of each substance for each species.

Fig. 3. Ordination diagram of double-constrained Correspondence Analysis showing the

relationships between species incidence, cell metabolism substances, and land use

classes. Arrow length indicates the importance of variables for the construction of

ordination axes.

Table legend

Table 1. Fourth-corner correlation calculated with between species traits (area of each

substance in liver) and environmental variables (percentage of land use type in 500 m

buffer).

|                        | Melanin | Hemosiderin | Lipofuscin |
|------------------------|---------|-------------|------------|
| Natural pasture        | 0.139   | -0.276      | -0.252     |
| Man-made building area  | -0.033  | -0.188      | -0.228     |
| Planted pastures       | 0.473   | -0.135      | -0.267     |
| Agriculture            | -0.386  | 0.721       | 0.78       |