Reduced Auditory Steady State Responses in Autism Spectrum Disorder

Seymour, R.A[1,2,3]., Rippon, G[1]., Gooding-Williams, G[1],. Sowman, P.F.[2] & Kessler, K[1].

[1] Aston Neuroscience Institute, School of Life and Health Sciences, Aston University, Birmingham, B4 7ET.

[2] Department of Cognitive Science, Macquarie University, Sydney, Australia, 2109.

[3] Wellcome Centre for Human Neuroimaging, Queen Square Institute of Neurology, University College London, 12 Queen Square, London WC1N 3AR, UK

Correspondence: k.kessler@aston.ac.uk
rob.seymour@ucl.ac.uk

Keywords: Auditory Steady State; Autism Spectrum Disorder; MEG; Gamma; Sensory.

Abstract

Background: Auditory steady state responses (ASSRs) are elicited by clicktrains or amplitude-modulated tones, which entrain auditory cortex at their specific modulation rate. Previous research has reported reductions in ASSRs at 40Hz for autism spectrum disorder (ASD) participants and first-degree relatives of people diagnosed with ASD [1,2].

Methods: Using a 1.5s-long auditory clicktrain stimulus, designed to elicit an ASSR at 40Hz, this study attempted to replicate and extend these findings. Magnetencephalography (MEG) data were collected from 18 adolescent ASD participants and 18 typically developing controls.

Results: The ASSR localised to bilateral primary auditory regions. Regions of interest were thus defined in left and right primary auditory cortex (A1). While the transient gamma-band response (tGBR) from 0-0.1s following presentation of the clicktrain stimulus was not different between groups, for either left or right A1, the ASD group had reduced oscillatory power at 40Hz from 0.5 to 1.5s post-stimulus onset, for both left and right A1. Additionally, the ASD group had reduced inter-trial coherence (phase consistency over trials) at 40Hz from 0.64-0.82s for right A1 and 1.04-1.22s for left A1.

Limitations: In this study, we did not conduct a clinical autism assessment (e.g. the ADOS), and therefore it remains unclear whether ASSR power and/or ITC are associated with the clinical symptoms of ASD.

Conclusion: Overall, our results support a specific reduction in ASSR oscillatory power and inter-trial coherence in ASD, rather than a generalised deficit in gamma-band responses. We argue that this could reflect a developmentally relevant reduction in non-linear neural processing.

Background

Autism Spectrum Disorder (ASD) is a neurodevelopmental condition characterised by impairments in social interaction, disrupted communication and repetitive behaviours [3]. Although these features remain the primary diagnostic markers of ASD, the presence of sensory symptoms has recently been given a more central diagnostic role. This change in symptom emphasis reflects the observation that over 90% of ASD individuals experience hyper- and/or hypo-sensitive sensory perception [4,5]. It has been suggested that differences in low-level sensory processing contribute to the atypical developmental trajectories of higher-level cognitive functions in autism [6]. An understanding of the neural circuits involved will therefore prove fruitful for ASD research, and could even facilitate the identification of earlier, brain-based diagnostic markers [7,8].

Dysregulated neural oscillations are a promising neural correlate of atypical sensory

processing in ASD. In particular, atypicalities in high frequency gamma-band oscillations

(30-80Hz) have been reported in ASD across visual, auditory and somatosensory domains

[7,9–13]. Gamma oscillations are generated through excitatory-inhibitory (E-I) neuronal

coupling [14], which facilitates periods of pre- and post-synaptic excitability alignment,

thereby promoting efficient neural communication [15]. Findings of atypical gamma

oscillations in ASD may therefore reflect disrupted E-I interactions within cortical micro-

circuits [16], and concomitant effects on local and global brain connectivity [17].

Within the context of auditory processing, dysregulated gamma-band oscillations in ASD

have been previously reported [7]. One prevalent approach to study auditory gamma-band

activity non-invasively is through amplitude modulated tones called "clicktrains". Such stimuli

produce two distinct gamma-band responses. First, a transient gamma-band response

(tGBR) is generated within 1s of stimulus onset [18]. This tGBR spans frequencies from 30-

60Hz and is generated in primary and secondary auditory cortices. Second, clicktrain stimuli

produce an auditory steady-state response (ASSR), in which neural populations in primary

auditory regions are entrained to the modulation frequency for the duration of the clicktrain

[19]. In adults, the entrainment in primary auditory cortex is greatest for clicktrains modulated

at 40Hz [20]. Measures of inter-trial coherence (ITC) can also be used to measure the

ASSR, by quantifying the degree of phase consistency across trials [21]. One advantage of

ASSRs is their high test re-test reliability which approaches an intraclass correlation of 0.96,

even with a relatively small number of trials [22,23]. Furthermore, ASSRs are modulated by

neural development, increasing in power by approximately 0.01 ITC value per year, until

early adulthood [24,25]. This increase has been linked with the maturation of superficial

cortical layers [26,27]. This makes the ASSR an ideal tool for studying auditory function in

developmental conditions such as ASD.

Two studies have measured ASSRs in an ASD context, that is, in ASD participants and in

the first-degree relatives of people diagnosed with ASD. Wilson and colleagues reported a

reduction in left-hemisphere auditory ASSR power in a group of 10 autistic adolescents,

using an early 37-channel MEG system [2]. The second study reported reduced ITC in first-

degree relatives of people diagnosed with ASD, with maximal reductions at 40Hz across

both hemispheres [1]. Reductions in the ASSR could therefore be an ASD-relevant

endophenotype. Additionally, the finding of reduced ITC suggests that dysregulated phase

dynamics in bilateral primary auditory cortex could underlie reductions in the ASSR in ASD.

However, measures of ITC have not been applied to study the ASSR directly in a group of

autistic participants. Additionally, it remains unclear whether reductions in ASSRs are

bilateral [1] or unilateral [2] in nature.

As discussed above, auditory stimuli also elicit a more broadband, transient gamma-band

response (tGBR) within 0.1s post-stimulus onset [18]. Previously, using auditory clicktrains,

Rojas and colleagues [1] reported equivalent tGBRs between the first-degree relatives of

people diagnosed with ASD, and controls. However, using sinusoidal auditory tones, several

studies have found reduced tGBRs in ASD [8,28,29]. It therefore remains unclear whether

both early evoked *and* later sustained gamma-band activity are dysregulated in ASD (also

see Kessler, Seymour & Rippon [7], for a review). We therefore opted to analyse tGBRs

alongside ASSRs using clicktrain stimuli in a group of autistic participants. However, the

primary focus for this study was on the sustained 40Hz response, given the ASD-related

differences, previously reported using clicktrain stimuli [1,2].

This study attempted to replicate and extend previous findings showing differences in

ASSRs and tGBRs in autism [1,2], within an adolescent population (aged 14-20). We

focussed on this age range because adolescence is a crucial period for brain maturation

[30,31] and ASSR power increases with age [24,25]. Therefore, we reasoned that ASD-

related differences in the ASSR would be more pronounced for an adolescent versus adult

population. We also opted to recruit adolescents rather than children for this study, given

fixed (adult) size of most MEG helmets and higher levels of compliance in adolescent

populations. Data were collected from a group of 18 ASD participants and 18 typically

developing controls using a 306-channel MEG system (Elekta Neuromag). An auditory

clicktrain stimulus was presented binaurally to participants, to elicit bilateral ASSRs at 40Hz.

To investigate prolonged neural entrainment, clicktrain stimuli were presented for a total of

1.5, rather than 0.5s as in previous studies [1,2]. ASSRs were analysed over time, in order

to investigate transient changes in 40Hz power and inter-trial coherence. It was

hypothesised that, compared with the control group, the ASD group would show reduced

ASSR power and ITC at 40Hz for the duration of clicktrain presentation [1,2]. In contrast, it

was hypothesised that tGBRs would be equivalent between groups, as previously reported

by Rojas & colleagues [1].

122 Methods

123 *Participants*

124 Data were collected from 18 participants diagnosed with ASD and 18 age-matched typically

125 developing controls, see Table 1. ASD participants had a confirmed clinical diagnosis of

ASD or Asperger's syndrome from a paediatric psychiatrist. Participants were excluded from participating if they were taking psychiatric medication or reported epileptic symptoms. Control participants were excluded from participating if a sibling or parent was diagnosed with ASD. MEG data from a further 9 participants was collected but excluded, due to: intolerance to MEG resulting in experimental attrition (2 ASD); movement over 0.5cm (2 ASD, 2 control, see MEG Acquisition); metal artefacts (1 ASD, 1 control); AQ score over 30 (1 control). The movement and AQ thresholds were defined before data collection began.

	N	**Age**	**Male/Female**	**Autism Quotient (Adult) /50**	**Raven Matrices Score /60**	**Glasgow Sensory Score /168**	**Mind in the Eyes Score /36**
ASD	18	Mean = 16.67; SD = 3.2; Range = 14-20	14 M; 4 F	Mean = 32.60*; SD = 6.64; Range = 21-46	Mean = 43.84; SD = 7.93; Range = 27-56	Mean = 65.33*; SD = 27.69; Range = 27-126	Mean = 21.88; SD = 4.87; Range = 12-30
Control	18	Mean = 16.89; SD = 2.8; Range = 14-20	15 M; 3 F	Mean = 10.91; SD = 5.43; Range = 6-21	Mean = 48.71; SD = 5.78; Range = 37-56	Mean = 38.70; SD = 6.88; Range = 29-50	Mean = 25.44; SD = 4.03; Range = 17-33

Table 1: Participant demographic and behavioural data. SD = Standard Deviation. * = behavioural scores significantly greater in ASD>control group, t-test, p<.05.

Behavioural Assessments

The severity of autistic traits was assessed using the Autism Quotient (AQ) [32] and sensory traits using the Glasgow Sensory Questionnaire (GSQ) [33]. Using an independent samples t-test, it was shown that both AQ scores, $t(34) = 9.869$, p<.001 and GSQ scores, $t(34) = 3.533$, p=.001, were significantly higher in the ASD group, compared with the control group (see Table 1).

Breaking down the GSQ scores further, we observed that our ASD participants showed a heterogeneous pattern of sensory symptoms, with mixtures of hypo- and hyper-sensitivities, across sensory domains (see Supporting Figure 4). Interestingly, auditory scores were the highest amongst any sensory modality, with a mean of 13.9/24. This means that the ASD participants, on average, answered between "Sometimes" and "Often", when reporting atypical auditory processing on the GSQ.

151 General non-verbal intelligence was assessed using the Raven's Matrices Task [34]. Using
152 an independent samples t-test, it was shown that there were no significant group differences
153 in the Raven Matrices Score, $t(34) = -1.372$, $p=.179$.
154
155 Participants also completed the Mind in the Eyes test [35], however there were no group
156 differences for this test, $t(34) = -1.615$, $p=.116$. The Mind in the Eyes test has been recently
157 criticised for measuring emotion recognition rather than an autism-specific deficit in mental
158 state attribution [36], and therefore these scores were not used to investigate correlations
159 between brain patterns and questionnaire measures.
160
Paradigm
Whilst undergoing MEG, participants performed an engaging sensory task. Each trial started
with a randomised fixation period (1.5, 2.5 or 3.5s), followed by the presentation of a visual
grating or auditory binaural click train stimulus. The visual and auditory stimuli were
presented randomly, rather than in separate experimental blocks. Only the auditory clicktrain
data will be described in this article (please see Seymour et al., 2019 [13] for analysis of the
visual grating data). The auditory clicktrain was created from auditory square wave clicks,
each of 2ms duration delivered every 25ms for a total of 1.5s. Clicktrains were presented at
80dB (verified using a decibel meter after pneumatic transduction and transmission)
binaurally through Etymotic MEG-compatible ear tubes. To keep participants engaged with
the task, cartoon pictures of aliens or astronauts were presented after the auditory clicktrain,
for a maximum of 0.5s. Participants were instructed to press a response-pad as soon as
they were presented with a picture of an alien, but not if they were presented with a picture
of an astronaut (maximum response duration allowed was 1.0s). Correct versus incorrect
responses were conveyed through 0.5s-long audio-visual feedback (correct: green box, high
auditory tone; incorrect responses: red box, low auditory tone). Prior to MEG acquisition, the
nature of the task was fully explained to participants and several practice trials were
performed. MEG recordings lasted 12-13 minutes and included 64 trials with auditory
clicktrain stimuli. Accuracy of picture classification was above 95% for all participants.
180

Figure 1. Experimental procedure. *(A) Participants performed an audiovisual task,*

consisting of 1.5–3.5 s baseline period followed by presentation of an auditory clicktrain

stimulus for a duration of 1.5s. After this, participants were presented with a cartoon alien or

astronaut picture and instructed to only respond when an alien was presented (response

time up to 1.5 s), followed by a green or a red framed box for a correct or an incorrect

response, respectively. The alien/astronaut stimuli were to maintain attention and do not

form part of the analysed data.

MEG Acquisition

MEG data were acquired using a 306-channel Neuromag MEG scanner (Vectorview, Elekta,

Finland) made up of 102 triplets of two orthogonal first-order planar gradiometers and one

magnetometer. Acquisition was conducted in a MaxShield™ magnetically shielded room

using MaxShield's™ patented "four layers in one shell" construction. All data were recorded

at a sampling rate of 1000 Hz, using default on-line filters between 0.1 - 330 Hz. Internal

Active Shielding (Max Shield) was turned off for the recording due to numerous artefacts

generated by this technology. Five head position indicator (HPI) coils were applied for

continuous head position tracking, and visualised post-acquisition using an in-house Matlab

script. Any participant who moved more than a conservative threshold of 5mm in any one

direction (x, y or z) were excluded from subsequent analysis. For MEG-MRI coregistration

purposes, the locations of three anatomical landmarks (nasion, left and right pre-auricular

points), the locations of the HPI coils and 300-500 points from the head surface were

acquired using a Polhemus Fastrak digitizer.

Structural MRI

A structural T1 brain scan was acquired for source reconstruction using a Siemens

MAGNETOM Trio 3T scanner with a 32-channel head coil (TE=2.18ms, TR=2300ms,

TI=1100ms, flip angle=9°, 192 or 208 slices depending on head size, voxel-size =

0.8x0.8x0.8cm).

MEG-MRI Coregistration and Cortical Mesh Construction

MEG data were co-registered with participants' structural MRIs by matching the digitised head-shape data with surface data from the structural scan [37]. Two control participants did not complete a T1 structural MRI and therefore the digitised head-shape data was matched with a database of 95 structural MRIs from the Human Connectome Database [38], using an Iterative Closest Points (ICP) algorithm. The headshape-MRI pair with the lowest ICP error was then used as a 'pseudo-MRI' for subsequent steps. This procedure has been shown to improve source localisation performance, in situations where a subject-specific anatomic MRI is not available [39,40]. The aligned MRI-MEG images were used to create a forward model based on a single-shell description of the inner surface of the skull [41] (3000 vertices), using the segmentation function in SPM8 [42]. The cortical mantle was then extracted to create a cortical mesh, using Freesurfer v5.3 [43], and registered to a standard *fs_LR* mesh, based on the Conte69 brain [44], using an interpolation algorithm from the Human Connectome Project [45] (also see: https://goo.gl/3HYA3L). Finally, the mesh was downsampled to 4002 vertices per hemisphere.

MEG Pre-Processing

Following data inspection, four MEG channels containing large amounts of non-physiological noise (low-frequency drift) were removed from the data, and not included in any of the subsequent pre-processing steps. No channel interpolation was performed. MEG data were pre-processed using Maxfilter (temporal signal space separation, default settings with correlation limit raised to 0.9), which attenuates external sources of noise from outside the head [46]. Further pre-processing steps were performed in Matlab 2014b using the Fieldtrip toolbox v20161024 [47]. Firstly, for each participant the entire recording was band-pass filtered between 0.5-250Hz (Butterworth filter, low-pass order 4, high-pass order 3) and band-stop filtered (49.5-50.5Hz; 99.5-100.5Hz) to remove residual 50Hz power-line contamination and its harmonic. Data were epoched into segments of 4s (1.5s pre, 1.5s post stimulus onset, with 0.5s of padding either side) and each trial was demeaned and detrended. Trials were inspected if the trial-by-channel (magnetometer) variance exceeded 8×10^{-23} (threshold determined from pilot testing), and those containing artefacts (SQUID jumps, eye-blinks, head movement, muscle) were removed. This resulted in the rejection, on average, of 3.4 trials per participant (mean number of trials across participants = 60.6, minimum = 54, maximum = 64). The remaining mean number of trials across participants used for analysis was therefore 60.6 (minimum = 54, maximum = 64). Finally, data were resampled to 200Hz to aid computation time.

Source-Level Spectral Power

Source analysis was conducted using a linearly constrained minimum variance beamformer

[48], which applies a spatial filter to the MEG data at each vertex of the cortical mesh. Only

data from orthogonal first-order planar gradiometers were used for source analysis due to

current controversy over combining both magnetometers and gradiometers (which have

different levels of noise). A covariance matrix for each participant was constructed from the

non-averaged, filtered (see below) data, rather than trial-averaged data (as sensor-level data

will be made more 'correlated' by averaging over trials). Beamformer weights were

calculated by combining the gradiometer covariance matrix with leadfield information, with

data pooled across baseline and clicktrain periods (-1.5s to 1.5s). Based on

recommendations for optimisation of MEG beamforming [49], a regularisation parameter of

lambda 5% was applied, due to the rank-deficiency of the data following the Maxfilter

procedure.

Figure 2: Procedure for Source Analysis. *For ASSR beamforming, a common spatial filter*

was computed using data pooled across ASSR and baseline data. This common filter was

then used to localise ASSR/baseline data separately. This process was repeated for tGBR

data, but the common spatial filter was computed using a different time and frequency band

of interest. ASSR = Auditory Steady State Response. tGBR = Transient Gamma-Band

Response.

Whilst the tGBR and ASSR originate from primary auditory cortex, both responses have

different frequency ranges and underlying neural generators [25]. Therefore we opted to use

separate spatial filters, rather than single spatial filter based on the M100 as used in

previous studies [2,50,51]. This decision was based on recent work suggesting that

beamformer weights should be optimised for specific data of interest [52].

To localise the ASSR, data were band-pass filtered (Butterworth filter, fifth order) between

35-45Hz. A period of 0.0-1.5s following stimulus onset was compared with a 1.5s baseline

period (1.5 to 0.0s before clicktrain onset, also see Figure 2). To localise the tGBR, data

were band-pass filtered between 30-60Hz, and a period of 0.0-0.1s following clicktrain onset

was compared with a 0.1s baseline period (see Figure 2).

ROI definition

Regions of interest (ROI) were selected in bilateral primary auditory (A1) cortices to

investigate ASSRs and tGBRs in greater detail. ROIs were defined using a multi-modal

parcellation from the Human Connectome Project (Supporting Figure 1, [53]). To obtain a

single spatial filter for each ROI (right A1 and left A1 separately), we performed principal

components analysis on the concatenated filters of each ROI, multiplied by the sensor-level

covariance matrix, and extracted the first component, see [54]. Broadband (0.5-250Hz)

sensor-level data were multiplied by this spatial filter to obtain "virtual electrodes".

A1 Spectral Power

A1 gamma power (ASSR, tGBR) was analysed using the multi-taper method, as

implemented in the Fieldtrip toolbox [47], using discrete prolate spheroidal sequences

(Slepian functions). This has been shown to offer an optimal trade-off between time and

frequency resolution, and is preferred to Morlet wavelets for high-frequency gamma-band

activity [55,56]. Oscillatory power was calculated from 35-45Hz using a 0.5s sliding window

(step size 0.02s) with ±5Hz frequency smoothing. Due to the narrow frequency range under

investigation, power values were averaged between 35-45Hz. Finally, the percentage

change in ASSR power was calculated, using the baseline time-period, i.e. 1.5s before

clicktrain presentation. For tGBR power, power values were averaged between 30-60Hz and

across time (0-0.1s versus a baseline window 0.1s before clicktrain presentation). From this,

the percentage change in tGBR power was calculated.

A1 Inter-trial Coherence

To assess band-limited phase consistency across trials, we calculated inter-trial coherence

(ITC). An ITC value of 0, indicates complete absence of phase consistency, whereas a value

of 1 indicates perfect phase consistency across trials [21]. ITC values were converted to Z-
values, as recommended by Maris & colleagues [57], to ensure a normal distribution for the
statistical analysis (see below).
310
Statistical Analysis
For MEG data, statistical analysis was performed using cluster-based permutation tests as
implemented in the Fieldtrip toolbox, which have been shown to adequately control the type-
I error rate for electrophysiological data [58]. Cluster permutation tests consist of two parts:
first an uncorrected independent t-test is performed (two-tailed), and all values exceeding a
5% significance threshold are grouped into clusters. The maximum t-value within each
cluster is carried forward. Second, a null distribution is obtained by randomising the data
labels (e.g. ASD/control and time) 10,000 times and calculating the largest cluster-level t-
value for each permutation. The maximum t-value within each original cluster is then
compared against this null distribution, with values exceeding a 5% significance threshold
(corrected across both tails, i.e. p<.025 for each tail) deemed significant. Given that only two
other MEG studies have used auditory clicktrains to study ASD-related differences [1,2], we
adopted a conservative statistical approach and used two-tailed tests in all instances.
324
For both left and right A1, the following within-group planned statistical contrasts were
performed: ASSR power (0.0s to 1.5s) versus baseline (-1.5s to 0.0s); ASSR ITC (0.0s to
1.5s) versus baseline (-1.5s to 0.0s). In addition, the following between-group planned
statistical contrasts were performed: control versus ASD ASSR power (0-1.5s post clicktrain
onset); control versus ASD ITC (0-1.5s post clicktrain onset); control versus ASD tGBR
power (0.0s to 0.1s).
331

Results

ASSR – Power

Whilst ASSRs are known to originate from bilateral primary auditory cortex [19,59], in order
to confirm successful source localisation with our pipeline, ASSR power (35-45Hz) was
localised on a cortical mesh, using an LCMV beamformer, see Methods. We then calculated
the percentage change in 35-45Hz power between 0.0-1.5s post-clicktrain onset versus a
1.5s baseline period (-1.5 to 0.0). As expected, the control group showed maximal increases
in power for regions overlapping bilateral primary auditory cortex (Figure 3A) [20,23]. For the
ASD group, there were increases in ASSR power for right, but not left, auditory regions,
albeit with lower average values than controls (Figure 3B). For an alternative visualisation of

results featuring un-thresholded whole-brain statistical maps, see Supporting Information,

Figure 3.

The use of beamforming for bilateral auditory responses has been questioned, due to the

potential for mis-localisations resulting from correlated neural sources [48]. However, as

noted by Van Veen & colleagues [48] and later by Sekihara & colleagues [60], complete

suppression of brain activity, only occurs when the cross-correlation of sources exceeds 0.9.

When realistic sources of noise are added to simulated MEG data, complete suppression

does not occur [61]. Instead Qurana & Cheyne [61], have shown that for correlated sources

at realistic signal-to-noise ratios, beamformers produce a single localisation directly in-

between the two sources. Given the clear separation between bilateral auditory sources in

our data, as shown in Figure 3, we argue that systematic mis-localisation is unlikely to have

occurred. Furthermore, following source analysis we used the online Neurosynth and

Neurovault tools to compute the spatial correlation between unthresholded group-level,

whole-brain images (see Figure 3; Supporting Figure 3) and several "concept-based meta-

analysis maps", generated from over 10,000 neuroimaging studies [62]. Results, reported in

Supporting Table 1, showed the highest correlation with the term "auditory" for both the

control (r = .635) and ASD group (r = .471).

ROIs were defined in bilateral auditory cortex (see Supporting Figure 1), to investigate time-

frequency responses in greater detail. Oscillatory power was calculated in steps of 0.02s

using the multitaper method, and post-stimulus periods (0 to 1.5s) were statistically

compared to baseline periods (-1.5 to 0s). Control participants showed increased 35-45z

power from 0.24-1.5s for left A1 and 0.21-1.5s for right A1 (Figure 3A bottom panel, times

passing a p<.05, two-tailed, threshold are indicated with a dotted line). In contrast, the ASD

group showed increased 35-45Hz power for much shorter time windows: in right A1 between

0.31-0.72s and 0.97-1.35s; and for left A1 between 0.41s-0.63s (Figure 3B top panel, times

passing a p<.05, two-tailed, threshold are indicated with a dotted line). Next, we statistically

compared ASSR 35-45Hz power between groups, for both ROIs. It was found that the

control group had greater 35-45Hz power in both right A1, 0.50-1.5s (Figure 4A) and left A1,

0.59 to 1.5s (Figure 4B), compared with the ASD group.

Figure 3. ASSR Power Analysis. *Top panels (A, B): ASSR beamformer localization. The percentage change in ASSR power (35-45Hz) is presented on a 3D cortical mesh, thresholded at values greater than 10% (white dotted line on colour scale) for illustrative purposes (for unthresholded images, see Supporting Figure 3), separately for control (A) and ASD (B) groups. Bottom panels (A,B): ASSR in Regions of Interest (ROIs). ROIs were defined in left and right A1 (see Supporting Figure 1) and ASSR oscillatory power was calculated between 35-45Hz. The time-period 0-1.5s post-clicktrain onset was statistically compared with a 1.5s baseline period. Data are plotted separately for: (A) the control group and (B) the ASD group. Dotted lines under the graph indicate times passing a p<.05 threshold (two-tailed) compared to baseline, with different colours corresponding to right A1 (red) and left A1 (green). ASSR = Auditory Steady State Response.*

Figure 4. ASSR Power Between Groups. *ASSR 35-45Hz power was statistically compared between groups for (A) right A1 and (B) left A1. It was found that the control group had greater ASSR power than the ASD group from 0.52-1.5s in right A1 and 0.60-1.5s for left A1. The black dotted line under the graph indicates times passing a p<.05 threshold (two-tailed) for the Control>ASD contrast.*

ASSR – Intertrial Coherence (ITC)

Next, inter-trial coherence (ITC) was calculated for the A1 ROIs, using the same time-frequency approach as for power. ITC values were Z-scored for statistical analysis [57]. First, we statistically compared the post-clicktrain time-period (0 to 1.5s) with the baseline time-period (-1.5 to 0s). The control group showed statistically significant, p<.05, increases in ITC from 0.1-1.48s for left A1, and 0.14-1.39s for right A1 (Figure 5A). The ASD group showed statistically significant, p<.05, increases in ITC from 0.18-1.50s for left A1, and between 0.12-0.60s, 0.80-0.90, and 1.08-1.5s for right A1 (Figure 5B).

Statistical comparison of ITC between groups showed that the control group had higher ITC in both right A1 (Figure 5C, p<.05) and left A1 (Figure 5D, p<.05), but only within short time-windows from 0.64-0.82s (right A1, Figure 5C) and 1.04-1.22s (left A1, Figure 5D).

Figure 5. ASSR ITC Analysis. *(A-B) 35-45HZ ASSR inter-trial coherence (ITC) results. The time-period 0-1.5s post-clicktrain onset was statistically compared with a 1.5s baseline period. Data are plotted separately for: (A) the control group and (B) the ASD group. Dotted lines under the graph indicate times passing a p<.05 threshold (two-tailed) compared to baseline, with different colours corresponding to right A1 (red) and left A1 (green). (C-D) Between Group 35-45Hz ASSR ITC results. It was found that the control group had greater ASSR ITC than the ASD group from 0.64-0.82s in right A1 (C) and 1.04-1.22s for left A1 (D). The black dotted line under the graph indicates times passing a p<.05 threshold (two-tailed) for the Control (blue) > ASD (red) contrast.*

ASSR – Behavioural Data

Next, we investigated whether ASSR responses in the ASD group were correlated with behavioural questionnaire data collected from participants. ASSR power and ITC values were averaged, separately, over those times showing a significant difference (p<.05) between the control and ASD group, as reported in the previous sections (also, see Figure 4 and 5C-D, black dotted lines). Furthermore, given the similar time-course of group differences across right and left A1, ASSR power values were averaged across the ROIs. This was not the case for ITC Z-values, however, which were not averaged across right and

left A1. These data were correlated with Autism Quotient (AQ) and Glasgow Sensory

Questionnaire (GSQ) scores, for the ASD group, only. There were no significant correlations

for ASSR power (Figure 6A, top (AQ) r = .01, p = .96; bottom (GSQ) r = -.17, p = .50), or ITC

Z-values (Figure 6B, top (AQ, left A1) r = -.36, p = .14; top (AQ, right A1) r = .11, p = .65;

bottom (GSQ, left A1) r = -.35, p = .14; bottom (GSQ, right A1) r = .02, p = .92). The

correlation analysis, was repeated for Glasgow Sensory Questionnaire (GSQ) Scores,

summed across the six auditory questions only, however no significant correlations were

found (p>.05, see Supporting Figure 5).

Figure 6. ASSR – Behaviour relationships. *Scatter plots to show the relationship between*

ASSR power, averaged across left/right A1 (A), or ITC Z-values (B), with Autism Quotient

(AQ) and Glasgow Sensory Questionnaire (GSQ) Scores. There were no significant (p>.05)

correlations for any brain-behaviour relationship. The shaded region indicates 95%

confidence intervals. ITC = Intertrial Coherence; ASSR = Auditory Steady State Response.

tGBR – Source-Level

Transient gamma-band responses to the auditory clicktrain were localised using a

beamforming approach (see Methods). As for the ASSR analysis, we first confirmed that the

cortical generator(s) of the ASSR originated in bilateral auditory cortex. We calculated the

percentage change in 30-60Hz power from 0.0-0.1s post-clicktrain onset compared with a

0.1 baseline period [58]. As expected, both groups group showed maximal increases in

tGBR power for regions overlapping with bilateral primary auditory cortex (Figure 7A)

[20,23].

Paralleling the ASSR analysis, ROIs were defined in left and right A1. For each ROI and

participant, we calculated the percentage change in tGBR power from 30-60Hz, between

0.0-0.1s post-clicktrain onset and a 0.1s baseline period. Statistically comparing groups, it

was found that there were no significant differences, p>.05, in tGBR power (see Figure 7B).

Figure 7. tGBR Analysis. *(A) The percentage change in transient gamma-band response,*

tGBR, power (30-60Hz) is presented on a 3D cortical mesh, thresholded at t > 3.6% (white

dotted line) for illustrative purposes, separately for control (top) and ASD (bottom) groups.

(B) ROIs were defined in left and right A1 (see Supporting Figure 1). The percentage change

in tGBR was plotted separately across ROIs (ASD: blue bar; controls: orange bar). Solid

black lines indicate 95% confidence intervals. There were no significant differences in tGBR

between groups (p>.05).

Discussion

This study examined the oscillatory basis of auditory steady state responses (ASSRs) and

transient gamma-band responses (tGBR) in a group of 18 autistic adolescents and 18

typically developing controls. We utilised robust source-localisation methods and analysed

auditory responses across time. Compared to the ASSR in the control group, we found

reduced ~40Hz power for the ASD group, for regions of interest defined in the left and right

primary auditory cortices. Furthermore, there was reduced inter-trial coherence for the

autistic group at 40Hz, suggesting that phase dynamics in A1 were less consistent over

time. Our results corroborate the notion that auditory brain responses in autism are locally

dysregulated [7], especially during sustained gamma-band entrainment (<0.5s post-stimulus

onset).

Auditory Steady State Responses (ASSR) in Autism

Our results are largely consistent with two previous studies which show reduced ASSRs in

autistic adolescents [2] and first-degree relatives of people diagnosed with autism [1]. Whilst

our study shows reductions in 40Hz power across *both* hemispheres (Figure 3-4), Wilson &

colleagues, observed a selective *left*-hemisphere reduction in power [2]. This might be due

to the monaural stimulation approach, used by Wilson & colleagues, producing larger

hemispheric asymmetries as compared to binaural auditory stimulation [63]. Future work is

clearly needed to clarify hemispheric asymmetries in ASSR power for ASD populations [63].

Our results build on the previous literature in several ways. Firstly, by examining sustained

ASSRs from 0-1.5s we found that group differences emerged beyond 0.5s post-stimulus

onset (Figures 3-4), suggesting that, when driven at gamma frequencies, A1 becomes

increasingly dysregulated in ASD compared to controls in a time-dependent manner. This

raises the intriguing possibility that sustained, rather than transient, oscillatory activity at

gamma-frequencies is affected in autism, perhaps reflecting synaptic dysfunction and an

imbalance between excitatory and inhibitory populations of neurons [16]. To investigate this

further, future work could parametrically modulate clicktrain duration, intensity, and variability

(e.g. perfect 40 Hz vs 38-42Hz, etc.). Secondly, we also found group differences in inter-trial

coherence (ITC), with reductions in the autistic group for two short time periods between 0.5

to 1.2s post-stimulus onset (Figure 6). Importantly, measures of ITC are normalised by

amplitude and have been shown to be more robust for data with lower signal-to-noise ratios

[23]. The reduction in ITC for the autistic group may reflect reduced phase consistency

across trials and more idiosyncratic neural responses in autism [64,65], as previously

reported for evoked data [66]. However, the reductions in ITC could have also emerged

through differences in ASSR power between groups [67]. That said, the time-course of

group-differences does diverge between ITC and power (see Supporting Figure 6), with

maximum ITC group differences not coinciding with maximum ASSR power differences,

which would be expected, if the latter would fully drive the former. In any case, our findings

strengthen the claim of reduced ASSRs in autism.

Transient Gamma-Band Responses in Autism

Unlike ASSRs, there were no group differences in the transient gamma-band (30-60Hz)

responses to the clicktrain stimulus (Figure 7). Whilst one previous study using sinusoidal

tones reported decreased tGBRs for the first-degree relatives of autistic people, a later study

using auditory clicktrains, found no group differences in either power or ITC [1]. More

generally, findings of transient/evoked gamma-band power across sensory domains are very

mixed, with both increases and decreases reported (reviewed in [7]). The divergence

between steady-state and transient gamma in this study has implications for potential

oscillopathies in ASD, as differences in gamma power may depend on the time-period under

investigation as well as the underlying neural circuits generating gamma oscillations [25].

ASSRs as Markers of Dysregulated Local Activity

There has been recent interest in characterising atypical patterns of gamma-band

oscillations in autism, due to their link with local cortical function and connectivity [7]. The

precise E-I mechanisms underlying gamma generation are well characterised, for a review

see [14]. Of particular importance is the functional inhibition of pyramidal neurons by fast-

spiking interneurons via binding of the neurotransmitter gamma-aminobutyric acid (GABA)

[14,68]. Relatedly, there is emerging evidence showing GABA dysfunction in autism [68].

Reduced gamma-band steady-state responses in autism may therefore reflect dysregulated

neuronal inhibition, resulting in E-I imbalance [16]. As argued by Kessler, Seymour & Rippon

[7], this local dysregulation could result in both hyper and hypo-sensitivities in ASD,

depending on the particular sensory input, and the degree of top-down modulatory

processes employed by individuals [13]. To quantify the precise mechanisms underlying

reduced gamma-band ASSRs, future studies could utilise dynamic causal modelling of A1

neuronal circuits [69], combined with parametric modulations of ASSRs (e.g. duration,

frequency) and participant attention [70]. It would also be interesting to use more naturalistic

auditory stimuli, for example speech stimuli [71,72], to investigate whether neural

entrainment is affected more generally in ASD.

It should also be noted that ASSRs are not simply generated via the linear accumulation of

transient evoked responses [20,73,74]. Instead, the ASSR may reflect a sustained non-

linear neural response at the input stimulation frequency and its harmonics, peaking at the

system's preferred modulation rate [20]. In support of this, Edgar & colleagues (2016) report

that in children, ASSRs are difficult to detect, despite measurable auditory evoked

responses [25]. Similarly, our data show intact auditory evoked fields (see Supporting Figure

2) and transient gamma-band responses in autism (Figure 7), in the presence of a reduced

ASSR (Figure 4). Rather than a generalised gamma-band dysfunction in autism, our data

suggest a more nuanced reduction in the non-linear dynamics underlying steady-state

auditory gamma [1]. Interestingly, an MEG study examining somatosensory processing in

ASD showed reduced frequency harmonics at 50Hz [12], while Vilidaite and colleagues

reported a reduction in harmonic EEG responses during visual steady-state stimulation in

autistic adults [75]. Furthermore, two MEG studies revealed reduced alpha-gamma phase-

amplitude coupling in the visual system in ASD [14, 18]. Overall, this suggests that non-

linear aspects of local cortical processing could be dysregulated across sensory domains in

ASD [8].

ASSRs are developmentally relevant, increasing by approximately 0.01 ITC value per year

[24,25,50]. This trajectory may reflect the continuing development of superficial layers of

cortex where gamma-band oscillations predominantly originate [27]. We hypothesise that the

ASD-related reduction in ASSRs reported in this study results from an atypical trajectory of

gamma-band maturation, in line with developmental disconnection theories of autism [76].

Given that the 40Hz ASSRs continue to mature throughout late adolescence and adulthood

[50], it remains to be established, whether the development of ASSRs in ASD is simply

delayed, or whether reductions persist throughout life. To investigate this further, future

studies should use high-powered longitudinal ASD samples and age-appropriate MEG

systems [77], to characterise ASSR development throughout childhood, adolescence and

into adulthood [78]. If confirmed, divergent ASSR trajectories could act as important autism-

relevant markers of intervention efficacy [79].

Limitations

In this study, formal clinical ASD assessment of our participants, e.g. the ADOS [80], was

not performed. We therefore implemented strict participant exclusion criteria, only including

autistic participants with a confirmed clinical diagnosis of ASD or Asperger's syndrome.

Between groups, there were significant differences in autistic and sensory traits, measured

using two self-report questionnaires (Table 1). However, upon closer inspection of

behavioural data (see Supporting Figure 5), the ASD group showed a mixture of hyper- and

hypo-sensitive traits between different sensory modalities making precise brain-behavioural

correlations problematic. This may explain the lack of relationship between ASSR power/ITC

and AQ/GSQ scores in ASD (Figure 6). Brain-behaviour relationships might be better

quantified using MEG in combination with psychophysical tests of auditory perception and

formal clinical assessments.

Declarations

Ethics approval and consent to participate

All experimental procedures complied with the Declaration of Helsinki and were approved by the Aston University, Department of Life & Health Sciences ethics committee. Participants and a parent/guardian gave written informed consent before participating in the study.

Consent for publication

N/A

Availability of data and materials

The data that support the findings of this study are available on reasonable request from corresponding author, R.S., in a preprocessed and de-anonymized form. The raw data are not publicly available due to ethical restrictions.

Competing Interests

The authors wish to declare the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Funding

The Wellcome Trust, Dr Hadwen Trust and Tommy's Fund supported research-scanning costs. Robert Seymour was supported by a cotutelle PhD studentship from Aston University and Macquarie University. Paul F. Sowman was supported by the Australian Research Council [DP170103148].

Author Contributions

RS: Study design; data collection; data analysis; data interpretation; manuscript writing.

KK: Study design; data interpretation; manuscript writing.

GR: Study design; data interpretation; manuscript writing.

PFS: Data interpretation; manuscript writing.

GGW: Data collection; data interpretation.

Acknowledgments

We wish to thank: the volunteers who gave their time to participate in this study; Dr Shu Yau and Dr. Michael Hall for help with MRI data acquisition; and Dr Jon Brock for intellectual contributions to experimental design.

References

1. Rojas DC, Teale PD, Maharajh K, Kronberg E, Youngpeter K, Wilson LB, et al. Transient and steady-state auditory gamma-band responses in first-degree relatives of people with autism spectrum disorder. Mol Autism. 2011;2:11.

2. Wilson TW, Rojas DC, Reite ML, Teale PD, Rogers SJ. Children and adolescents with autism exhibit reduced MEG steady-state gamma responses. Biol Psychiatry. 2007;62:192–197.

3. American Psychological Association. Diagnostic and statistical manual of mental disorders . Arlington: American Psychiatric Association; 2013.

4. Hazen EP, Stornelli JL, O'Rourke JA, Koesterer K, McDougle CJ. Sensory symptoms in autism spectrum disorders. Harv Rev Psychiatry. 2014;22:112–124.

5. Leekam SR, Nieto C, Libby SJ, Wing L, Gould J. Describing the sensory abnormalities of children and adults with autism. J Autism Dev Disord. 2007;37:894–910.

6. Robertson CE, Baron-Cohen S. Sensory perception in autism. Nat Rev Neurosci. 2017;18:671.

7. Kessler K, Seymour RA, Rippon G. Brain oscillations and connectivity in autism spectrum disorders (ASD): new approaches to methodology, measurement and modelling. Neurosci Biobehav Rev. 2016;71:601–20.

8. Roberts TP, Khan SY, Rey M, Monroe JF, Cannon K, Blaskey L, et al. MEG detection of delayed auditory evoked responses in autism spectrum disorders: towards an imaging biomarker for autism. Autism Res. 2010;3:8–18.

9. Simon DM, Wallace MT. Dysfunction of sensory oscillations in Autism Spectrum Disorder. Neurosci Biobehav Rev. 2016;68:848–61.

10. Brown C, Gruber T, Boucher J, Rippon G, Brock J. Gamma abnormalities during perception of illusory figures in autism. Cortex. 2005;41:364–376.

11. Wright B, Alderson-Day B, Prendergast G, Bennett S, Jordan J, Whitton C, et al. Gamma activation in young people with autism spectrum disorders and typically-developing controls when viewing emotions on faces. PLoS One. 2012;7:e41326.

12. Khan, Michmizos K, Tommerdahl M, Ganesan S, Kitzbichler MG, Zetino M, et al. Somatosensory cortex functional connectivity abnormalities in autism show opposite trends, depending on direction and spatial scale. Brain. 2015;138:1394–409.

13. Seymour RA, Rippon G, Gooding-Williams G, Schoffelen JM, Kessler K. Dysregulated oscillatory connectivity in the visual system in autism spectrum disorder. Brain. 2019; 142:3294–3305.

14. Buzsáki G, Wang X-J. Mechanisms of gamma oscillations. Annu Rev Neurosci. 2012;35:203.

15. Fries P. Rhythms for cognition: communication through coherence. Neuron. 2015;88:220–235.

16. Rubenstein JLR, Merzenich MM. Model of autism: increased ratio of excitation/inhibition
in key neural systems. Genes Brain Behav. 2003;2:255–267.

17. Khan S, Gramfort A, Shetty NR, Kitzbichler MG, Ganesan S, Moran JM, et al. Local and
long-range functional connectivity is reduced in concert in autism spectrum disorders. Proc
Natl Acad Sci. 2013;110:3107–12.

18. Pantev C. Evoked and induced gamma-band activity of the human cortex. Brain Topogr.
1995;7:321–330.

19. Hari R, Hämäläinen M, Joutsiniemi S-L. Neuromagnetic steady-state responses to
auditory stimuli. J Acoust Soc Am. 1989;86:1033–1039.

20. Pantev C, Roberts LE, Elbert T, Roß B, Wienbruch C. Tonotopic organization of the
sources of human auditory steady-state responses. Hear Res. 1996;101:62–74.

21. Busch NA, Dubois J, VanRullen R. The phase of ongoing EEG oscillations predicts
visual perception. J Neurosci. 2009;29:7869–7876.

22. McFadden KL, Steinmetz SE, Carroll AM, Simon ST, Wallace A, Rojas DC. Test-Retest
Reliability of the 40 Hz EEG Auditory Steady-State Response. PLOS ONE. 2014;9:e85748.

23. Tan H-R, Gross J, Uhlhaas PJ. MEG—measured auditory steady-state oscillations show
high test–retest reliability: A sensor and source-space analysis. Neuroimage. 2015;122:417–
426.

24. Cho RY, Walker CP, Polizzotto NR, Wozny TA, Fissell C, Chen C-MA, et al.
Development of sensory gamma oscillations and cross-frequency coupling from childhood to
early adulthood. Cereb Cortex. 2015;25:1509–1518.

25. Edgar JC, Fisk CL, Liu S, Pandey J, Herrington JD, Schultz RT, et al. Translating Adult
Electrophysiology Findings to Younger Patient Populations: Difficulty Measuring 40-Hz
Auditory Steady-State Responses in Typically Developing Children and Children with Autism
Spectrum Disorder. Dev Neurosci. 2016; 38:1-14.

26. Moore JK, Guan Y-L. Cytoarchitectural and axonal maturation in human auditory cortex.
J Assoc Res Otolaryngol. 2001;2:297–311.

27. Moore JK, Linthicum Jr FH. The human auditory system: a timeline of development. Int J
Audiol. 2007;46:460–478.

28. Rojas DC, Maharajh K, Teale P, Rogers SJ. Reduced neural synchronization of gamma-
band MEG oscillations in first-degree relatives of children with autism. BMC Psychiatry.
2008;8:66.

29. Rojas DC, Wilson LB. γ-band abnormalities as markers of autism spectrum disorders.
Biomark Med. 2014;8:353–368.

30. Blakemore S-J. Development of the social brain in adolescence. J R Soc Med.
2012;105:111–116.

31. Casey BJ, Getz S, Galvan A. The adolescent brain. Dev Rev. 2008;28:62–77.

32. Baron-Cohen S, Wheelwright S, Skinner R, Martin J, Clubley E. The autism-spectrum
quotient (AQ): Evidence from asperger syndrome/high-functioning autism, males and
females, scientists and mathematicians. J Autism Dev Disord. 2001;31:5–17.

33. Robertson AE, Simmons DR. The relationship between sensory sensitivity and autistic
traits in the general population. J Autism Dev Disord. 2013;43:775–784.

34. Raven JC, Court JH. Raven's progressive matrices and vocabulary scales . Oxford:
Oxford Psychologists Press; 1998.

35. Baron-Cohen S, Wheelwright S, Hill J, Raste Y, Plumb I. The "Reading the Mind in the
Eyes" test revised version: A study with normal adults, and adults with Asperger syndrome
or high-functioning autism. J Child Psychol Psychiatry. 2001;42:241–251.

36. Oakley BF, Brewer R, Bird G, Catmur C. Theory of mind is not theory of emotion: A
cautionary note on the Reading the Mind in the Eyes Test. J Abnorm Psychol.
2016;125:818.

37. Jenkinson M, Smith S. A global optimisation method for robust affine registration of brain
images. Med Image Anal. 2001;5:143–56.

38. Larson-Prior LJ, Oostenveld R, Della Penna S, Michalareas G, Prior F, Babajani-Feremi
A, et al. Adding dynamics to the Human Connectome Project with MEG. Neuroimage.
2013;80:190–201.

39. Gohel B, Lim S, Kim M-Y, Kwon H, Kim K. Approximate Subject Specific Pseudo MRI
from an Available MRI Dataset for MEG Source Imaging. Front Neuroinformatics.
2017;11:50.

40. Seymour R. Macquarie-MEG-Research/MEMES: For Zenodo [Internet]. 2018. Available
from: https://doi.org/10.5281/zenodo.1451031

41. Nolte G. The magnetic lead field theorem in the quasi-static approximation and its use
for magnetoencephalography forward calculation in realistic volume conductors. Phys Med
Biol. 2003;48:3637–52.

42. Litvak V, Mattout J, Kiebel S, Phillips C, Henson R, Kilner J, et al. EEG and MEG data
analysis in SPM8. Comput Intell Neurosci. 2011.

43. Fischl B. FreeSurfer. Neuroimage. 2012;62:774–781.

44. Essen V, C D, Glasser MF, Dierker DL, Harwell J, Coalson T. Parcellations and
Hemispheric Asymmetries of Human Cerebral Cortex Analyzed on Surface-Based Atlases.
Cereb Cortex. 2012;22:2241–62.

45. Van Essen DC, Ugurbil K, Auerbach E, Barch D, Behrens TEJ, Bucholz R, et al. The
Human Connectome Project: a data acquisition perspective. Neuroimage. 2012;62:2222–
2231.

46. Taulu S, Simola J. Spatiotemporal signal space separation method for rejecting nearby
interference in MEG measurements. Phys Med Biol. 2006;51:1759.

47. Oostenveld R, Fries P, Maris E, Schoffelen J-M. FieldTrip: open source software for
advanced analysis of MEG, EEG, and invasive electrophysiological data. Comput Intell
Neurosci. 2010.

48. Van Veen BD, van Drongelen W, Yuchtman M, Suzuki A. Localization of brain electrical
activity via linearly constrained minimum variance spatial filtering. IEEE Trans Biomed Eng.
1997;44:867–80.

49. Brookes MJ, Vrba J, Robinson SE, Stevenson CM, Peters AM, Barnes GR, et al.
Optimising experimental design for MEG beamformer imaging. Neuroimage. 2008;39:1788–
1802.

50. Rojas DC, Maharajh K, Teale PD, Kleman MR, Benkers TL, Carlson JP, et al.
Development of the 40Hz steady state auditory evoked magnetic field from ages 5 to 52.
Clin Neurophysiol. 2006;117:110–117.

51. Rojas DC, Teale PD, Maharajh K, Kronberg E, Youngpeter K, Wilson LB, et al. Transient
and steady-state auditory gamma-band responses in first-degree relatives of people with
autism spectrum disorder. Mol Autism. 2011;2:1.

52. Barratt EL, Francis ST, Morris PG, Brookes MJ. Mapping the topological organisation of
beta oscillations in motor cortex using MEG. NeuroImage. 2018;

53. Glasser MF, Coalson TS, Robinson EC, Hacker CD, Harwell J, Yacoub E, et al. A multi-
modal parcellation of human cerebral cortex. Nature. 2016;536:171–8.

54. Schoffelen J-M, Hultén A, Lam N, Marquand AF, Uddén J, Hagoort P. Frequency-
specific directed interactions in the human brain network for language. Proc Natl Acad Sci.
2017;114:8083–8088.

55. Hoogenboom N, Schoffelen J-M, Oostenveld R, Parkes LM, Fries P. Localizing human
visual gamma-band activity in frequency, time and space. Neuroimage. 2006;29:764–773.

56. Muthukumaraswamy SD. High-frequency brain activity and muscle artifacts in
MEG/EEG: a review and recommendations. Front Hum Neurosci. 2013.

57. Maris E, Schoffelen J-M, Fries P. Nonparametric statistical testing of coherence
differences. J Neurosci Methods. 2007;163:161–75.

58. Maris E, Oostenveld R. Nonparametric statistical testing of EEG- and MEG-data. J
Neurosci Methods. 2007;164:177–90.

59. Popov T, Oostenveld R, Schoffelen JM. FieldTrip Made Easy: An Analysis Protocol for
Group Analysis of the Auditory Steady State Brain Response in Time, Frequency, and
Space. Front Neurosci. 2018;12.

60. Sekihara K, Nagarajan SS, Poeppel D, Marantz A. Performance of an MEG adaptive-
beamformer technique in the presence of correlated neural activities: effects on signal
intensity and time-course estimates. IEEE Trans Biomed Eng. 2002;49:1534–46.

61. Quraan MA, Cheyne D. Reconstruction of correlated brain activity with adaptive spatial
filters in MEG. NeuroImage. 2010;49:2387–400.

62. Gorgolewski, K. J., Varoquaux, G., Rivera, G., Schwarz, Y., Ghosh, S. S., Maumet, C.,
... & Yarkoni, T. NeuroVault.org: a web-based repository for collecting and sharing
unthresholded statistical maps of the human brain . Frontiers in Neuroinformatics. 2015; 9:8.

63. Ross B, Herdman AT, Pantev C. Right hemispheric laterality of human 40 Hz auditory
steady-state responses. Cereb Cortex. 2005;15:2029–2039.

64. Hahamy A, Behrmann M, Malach R. The idiosyncratic brain: distortion of spontaneous
connectivity patterns in autism spectrum disorder. Nat Neurosci. 2015;18:302–309.

65. Nunes AS, Peatfield N, Vakorin V, Doesburg SM. Idiosyncratic organization of cortical
networks in autism spectrum disorder. Neuroimage. 2018;

66. Dinstein I, Pierce K, Eyler L, Solso S, Malach R, Behrmann M, et al. Disrupted neural
synchronization in toddlers with autism. Neuron. 2011;70:1218–1225.

67. van Diepen RM, Mazaheri A. The Caveats of observing Inter-Trial Phase-Coherence in
Cognitive Neuroscience. Sci Rep. Nature Publishing Group; 2018;8:1–9.

68. Coghlan S, Horder J, Inkster B, Mendez MA, Murphy DG, Nutt DJ. GABA system
dysfunction in autism and related disorders: from synapse to symptoms. Neurosci Biobehav
Rev. 2012;36:2044–2055.

69. Moran RJ, Stephan KE, Seidenbecher T, Pape H-C, Dolan RJ, Friston KJ. Dynamic
causal models of steady-state responses. NeuroImage. 2009;44:796–811.

70. Ross B, Picton TW, Herdman AT, Pantev C. The effect of attention on the auditory
steady-state response. Neurol Clin Neurophysiol NCN. 2004;2004:22–22.

71. Destoky F, Philippe M, Bertels J, Verhasselt M, Coquelet N, Vander Ghinst M, et al.
Comparing the potential of MEG and EEG to uncover brain tracking of speech temporal
envelope. NeuroImage. 2019;184:201–13.

72. Meng Q, Hegner YL, Giblin I, McMahon C, Johnson BW. Lateralised cerebral processing
of abstract linguistic structure in clear and degraded speech. bioRxiv. Cold Spring Harbor
Laboratory; 2020;2020.02.05.934604.

73. Azzena GB, Conti G, Santarelli R, Ottaviani F, Paludetti G, Maurizi M. Generation of
human auditory steadystate responses (SSRs). I: Stimulus rate effects. Hear Res.
1995;83:1–8.

74. Santarelli R, Maurizi M, Conti G, Ottaviani F, Paludetti G, Pettorossi VE. Generation of
human auditory steady-state responses (SSRs). II: Addition of responses to individual
stimuli. Hear Res. 1995;83:9–18.

75. Vilidaite G, Norcia AM, West RJ, Elliott CJ, Pei F, Wade AR, et al. Autism sensory
dysfunction in an evolutionarily conserved system. Proc R Soc B. 2018;285:20182255.

76. Geschwind DH, Levitt P. Autism spectrum disorders: developmental disconnection
syndromes. Curr Opin Neurobiol. 2007;17:103–111.

77. Rapaport H, Seymour RA, Sowman PF, Benikos N, Stylianou E, Johnson BW, et al.
Studying brain function in children using magnetoencephalography. JoVE J Vis Exp.
2019;e58909.

78. Loth E, Charman T, Mason L, Tillmann J, Jones EJH, Wooldridge C, et al. The EU-AIMS
Longitudinal European Autism Project (LEAP): design and methodologies to identify and
validate stratification biomarkers for autism spectrum disorders. Mol Autism. 2017;8:24.

79. Mamashli F, Khan S, Bharadwaj H, Losh A, Pawlyszyn SM, Hämäläinen MS, et al.
Maturational trajectories of local and long-range functional connectivity in autism during face
processing. Hum Brain Mapp. 2018;39:4094–4104.

80. Lord C, Risi S, Lambrecht L, Cook EH, Leventhal BL, DiLavore PC, et al. The Autism
Diagnostic Observation Schedule—Generic: A standard measure of social and

communication deficits associated with the spectrum of autism. J Autism Dev Disord.
2000;30:205–223.

Supporting Information:

Reduced Auditory Steady State Responses in Autism Spectrum Disorder

Seymour, R.A., Rippon, G., Gooding-Williams, G,. Sowman, P.F. & Kessler, K.

Regions of Interest

Left A1   *Right A1*

Supporting Figure 1: Regions of interest were defined in left and right Primary Auditory Cortex, according to HCP-MMP 1.0 Atlas.

Supporting Figure 2: Sensor-level Analysis. **(A)** Group average topo-plot for the auditory M100 event-related field, magnetometers shown. **(B)** Group average topo-plot for auditory steady state responses (ASSR) at 40Hz **(C)** Group average topo-plot of the transient gamma-band response (tGBR), 30-60Hz, 0.0-0.1s. Scales represent MEG field strength, baseline-corrected, with units of Tesla/cm.

Supporting Figure 3: Whole-brain maps showing changes in ASSR power, corresponding to Figure 2 in the main text.

Supporting Figure 4: Responses to the Glasgow Sensory Questionnaire were grouped by sensory domain (maximum score = 20) and hypo- / hyper-sensitivity (green and blue bars respectively). Our data show a heterogeneous pattern of sensory symptoms, with mixture of hypo- and hyper-sensitivities. Auditory symptoms scored 13.9/20 corresponding to questionnaire answers closest to "Sometimes".

Supporting Figure 5: Scatter plots to show the relationship between ASSR power, averaged across left/right A1, and Glasgow Sensory Questionnaire (GSQ) Scores, summed across the six auditory questions only. There were no significant (p>.05) correlations for ASSR power, r = -.07, p=.77, or for ITC Z-Value (left A1), r = -.29, p=.24, (right A1), r = -.07, p = .76. The shaded region indicates 95% confidence intervals. ITC = Intertrial Coherence; ASSR = Auditory Steady State Response.

Supporting Figure 6: For ASSR power (top) and ITC power (bottom), the group average for the ASD group was subtracted from the group average for controls (at each time bin) and plotted for each ROI (right A1: red; left A1: green). The black dotted lines link the times maximum ITC group differences for left and right A1 with the corresponding data for ASSR power.

Rank	Control		ASD	
	Search Term	Correlation	Search Term	Correlation
1	auditory	0.635	auditory	0.471
2	superior temporal	0.595	auditory cortex	0.427
3	auditory cortex	0.57	superior temporal	0.424
4	sounds	0.567	sounds	0.408
5	listening	0.557	sound	0.404
6	sound	0.552	listening	0.398
7	temporal	0.55	heschl gyrus	0.392
8	planum temporale	0.539	heschl	0.39
9	temporale	0.539	primary auditory	0.389
10	planum	0.531	planum temporale	0.388

Supporting Table 1: Results of the meta-analytical decoding (top 10 terms shown). The correlation term corresponds to the r-value between the whole-brain unthresholded map and the Neurosynth concept-based meta-analysis maps, generated from over 10,000 neuroimaging studies.