PharmOmics: A Species- and Tissue-specific Drug Signature Database and Online Tool

for Toxicity Prediction and Drug Repurposing

Yen-Wei Chen[1,2,3], Douglas Arneson[2,4], Graciel Diamente[2], Jennifer Garcia[2], Nima Zaghari[2],

Paul Patel[2], Patrick Allard[1,3], and Xia Yang[1,2,4,5,6]

[1]Interdepartmental Program of Molecular Toxicology, University of California, Los Angeles,

Los Angeles, CA

[2]Department of Integrative Biology and Physiology, University of California, Los Angeles,

CA

[3]Institute for Society and Genetics, University of California, Los Angeles, Los Angeles, CA

[4] Interdepartmental Program of Bioinformatics, University of California, Los Angeles, Los

Angeles, CA

[5]Institute for Quantitative and Computational Biosciences, University of California, Los

Angeles, CA

[6]Molecular Biology Institute, University of California, Los Angeles, CA

Correspondence:

Xia Yang, Ph.D.

Dept. Integrative Biology and Physiology, UCLA

Email: xyang123@ucla.edu

Abstract

Drug development has been hampered by a high failure rate in clinical trials due to suboptimal efficacy or safety issues that are not predicted by preclinical studies in model systems. A key contributor to this hurdle is our incomplete understanding of how drug molecules function across organ systems and species. Therefore, elucidating species- and tissue-specific molecular actions of drugs can provide systems level insights into therapeutic efficacy, toxicity, adverse drug reactions (ADRs), and between-species differences that are necessary for more effective translational medicine. Here, we present a comprehensive drug knowledgebase and analytical tool, PharmOmics, comprised of genomic footprints of drugs in individual tissues from multiple species (human, mouse, and rat) based on meta-analysis of global transcriptome data from public repositories (GEO, ArrayExpress, TG-GATEs, drugMatrix). Using multi-species multi-tissue gene expression signatures as molecular indicators of drug functions, we demonstrate the potential of PharmOmics to predict drugs that can be used for new disease indications or that may induce ADRs. PharmOmics was able to accurately predict known drugs for multiple diseases and identify potential drugs targeting non-alcoholic fatty liver disease for which no approved drugs are currently available. In addition, we found that drugs causing similar ADRs tend to have gene signatures with close proximity in gene regulatory networks. Comparison of our PharmOmics platform with other commonly used drug signature databases, most of which are based on data derived from *in vitro* cell lines, demonstrates that our tissue- and species-specific drug signatures serve as a complementary resource to support drug characterization.

50   **Background**

Drug development has been challenging and costly over the past decades due to the

continuously high failure rate in clinical trials (1). Most drugs with excellent efficacy and

safety profiles in preclinical studies often encounter suboptimal efficacy or safety concerns in

humans (2). This translational gap is likely attributable to our incomplete understanding of

the systems level activities of drugs in individual tissues and organ systems (3) as well as the

differences between humans and model systems (4).

Drug activities can be captured by gene expression patterns, commonly referred to as gene

signatures. By measuring how a pharmacological agent affects the gene signature of a cell or

a tissue type in a particular species, we can infer the cell- or tissue-specific biological

pathways involved in therapeutic processes or toxicological responses. This concept has

prompted drug repositioning studies and provided important predictions for repurposing

approved drugs for new disease indications (5–10). Similarly, gene signatures can reveal

mechanisms underlying adverse drug reactions (ADRs) and can therefore be leveraged to

predict these ADRs as previously shown for liver and kidney toxicity (11–13).

A drug may affect different molecular processes between tissues, causing disease treatment

effects in the desired target tissue(s) but toxicity or ADRs through off-target effects in other

tissues. Therefore, tissue-specific drug treatment signatures will offer more systematic

understanding of drug actions *in vivo*. Additionally, rodent models have been commonly used

in toxicology and preclinical studies, yet species-specific effects of drugs has been well

described (14) and underlie the lack of efficacy or unexpected ADRs or toxicity responses of

certain drugs when used in humans (15). Therefore, understanding the species-specific

molecular effects of drugs is also of high biological importance. A detailed species- and

tissue-specific drug genomic signature database will significantly improve our understanding

of the molecular networks affected by drugs and facilitate network-based drug discovery and

ADR prediction for translational medicine.

The tremendous potential of using gene signatures to facilitate target and toxicity

identification has led to several major efforts in characterizing genomic signatures related to

drug treatment (8,16–18). However, none of the existing platforms offer comprehensive

cross-tissue and cross-species *in vivo* assessments of drug activities to allow predictions of

the effects of a drug on individual tissues and to help assess the translational potential of a

drug based on consistencies or discrepancies between species. For instance, the comparative

toxicogenomics database (CTD), a literature-based resource curating chemical-to-

gene/protein associations as well as chemical-to-disease and gene/protein-to-disease

connections (16), lacks the cellular and tissue context of the curated interactions. More

systematic, data-driven databases like CMAP (8) and LINC1000 (17) focus on characterizing

and cataloging the genomic footprints of more than ten thousand chemicals using *in vitro* cell

lines (primarily cancer cell lines) to offer global views of molecular responses to drugs in

individual cellular systems. However, these *in vitro* cell-line based gene signatures may not

capture *in vivo* tissue-specificity of drug activities. To move into the *in vivo* systems, large

toxicogenomics databases like TG-GATEs (19) and DrugMatrix from the National

Toxicology Programs of the National Institute of Environmental Health Sciences

(https://ntp.niehs.nih.gov/drugmatrix/index.html) have become available to provide unbiased

transcriptome assessment for heart, muscle, liver, and kidney systems. However, information

about other organ systems is limited. To broaden the tissue coverage, efforts to meta-analyze

publicly deposited transcriptomic datasets in GEO (20) and ArrayExpress (21) have also been

described (18), but the species- and tissue-specific effects of drugs have not been addressed.

Here, we establish a computational pipeline to meta-analyze 1,314 drug-related

transcriptomic datasets from mouse, human, and rat studies curated in GEO, ArrayExpress,

TG-GATEs and DrugMatrix, covering >20 tissue/organ systems, and summarize the results

into tissue- and species-specific drug treatment signatures. In addition, we demonstrate the

utility and performance of these signatures for gene network-based drug repositioning and

ADR assessment. The tissue- and species-specific drug signatures are hosted on an

interactive web server PharmOmics, to enable public access, download, and integrative

analyses for drug repositioning (http://mergeomics.research.idre.ucla.edu/PharmOmics/).

Construction and content

Curation of tissue- and species-specific drug transcriptomic datasets

As illustrated in **Figure 1**, a list of clinically relevant target drugs were first compiled from

Kyoto Encyclopedia of Genes and Genomes (KEGG) (16), followed by transcriptomic

dataset search in GEO (20), ArrayExpress (21), TG-GATEs (19) and DrugMatrix

(https://ntp.niehs.nih.gov/drugmatrix/index.html) based on matching generic names in either

the title or abstract fields of the dataset records. GEO and ArrayExpress datasets were

identified as of January 2018. Duplicated datasets between repositories were removed. We

developed a semi-automated pipeline combining automated search with manual checking to

identify relevant datasets for drug treatment. The automated process first extracts datasets

containing drug generic names or abbreviations and then inspects the potential datasets for

availability of both drug treatment and control labels in the constituent samples. We also

manually checked the recorded labels identified by the automated process to validate the

labels to decrease potential false detections. Only datasets with n>=3/group in both drug

treatment and control groups were included in our downstream analyses. We combined drug

treatment samples regardless of dose and regimen for the same drug in order to obtain the

consensus responses across conditions. The decision on not further stratifying the studies

based on dose and regimen was to increase the generalizability of the drug signatures and to

mitigate the reduced statistical power due to uneven coverage of dose and regimen

information across data repositories and drugs.

Obtaining drug treatment signatures stratified by species and tissues

Based on the metadata of each dataset, we retrieved the species and tissue labels. Tissue

names were standardized based on the Brenda Tissue Ontology (22). We implemented a

search procedure to climb the ontology tree structure to determine the suitable tissue

annotations. We first defined a tissue/organ priority list in the Brenda Tissue Ontology tree

system based on the widely used organ systems in toxicological research. Tissue/organ

priority order was set to "kidney", "liver", "pancreas", "breast", "ovary", "adipose tissue",

"cardiovascular system", "nervous system", "respiratory system", "urogenital system",

"immune system", "hematopoietic system", "skeletal system", "integument" (endothelial and

skin tissue), "connective tissue", "muscular system", "gland", "gastrointestinal system", and

"viscus" (other non-classified tissue). Tissue terms relevant to each of these tissues or organs

were curated from the ontology tree into a tissue/organ ontology table. Next, in our

tissue/organ ontology table, we looked up the terms in the Cell/Organism/Tissue column of

the metadata in each transcriptomic dataset. If not found, we searched the title and summary

columns of the metadata as well to retrieve additional information. When searching returned

multiple tissue terms, we used the one with the higher priority order as described above.

Manual checking was conducted to confirm the tissue annotation for each dataset.

For each gene expression dataset, normalized data were retrieved, followed by assessment of

quantile distribution of values and, when data was not normally distributed, log2-

transformation using GEO2R (20). Due to the heterogeneity of the datasets in terms of study

design, platforms, sample size, and normalization methods, we chose to use the characteristic

direction method in the GeoDE package (23) to identify genes affected by drug treatment.

GeoDE was designed to accommodate datasets from different settings and outputs

differences between treatment and control groups using a "characteristic direction" measure

analogous but superior to fold change in identifying biologically relevant genes and pathways.

The normalized characteristic directions for all genes were then transformed into a non-

parametric rank representation. Subsequently, gene ranks of a particular drug from the same

tissue/organ system and the same organism were aggregated across datasets using the Robust

Rank Aggregation method (24) to identify differentially expressed genes (DEGs) of the drug

within each tissue for each species. Robust Rank Aggregation provides a non-parametric

meta-analysis across different ranked lists to obtain commonly shared genes across datasets,

which avoids statistical issues associated with heterogeneous datasets. It computes a null

distribution based on randomized gene ranks and then compares the null distribution with the

empirical gene ranks to obtain a p-value for each gene. The robust rank aggregation process

was done for the upregulated and downregulated genes separately to obtain DEGs for both

directions under Bonferroni-adjusted p-value <0.01, a cutoff implemented in the Robust Rank

Aggregation algorithm. To obtain species-level signatures for each drug, we further

aggregated DEGs across different organs tested for each drug within each species. Pathway

analysis of individual drug signatures was conducted using enrichr (25) by intersecting each

signature with pathways or gene sets from KEGG (26), gene ontology biological pathways

(GOBP) (27), Reactome pathways (28), Biocarta pathways (29) and Wikipathways (30).

Pathways at false discovery rate (FDR) <0.05 were considered significant. For cross-tissue

and species comparison described in this article we used all five pathway databases to make it

more comprehensive.

As summarized in **Figure 2,** in total we identified 1,314 transcriptomic datasets across 551

drugs from GEO, ArrayExpress, DrugMatrix, and TG-GATEs. These datasets cover >20

tissue or organ systems in rats, human, and mouse and 289 KEGG drug classes. The

toxicogenomics databases TG-GATEs and DrugMatrix mainly deposited rat liver and kidney

datasets, while public data repositories GEO and ArrayExpress contained datasets with

broader tissue and species coverage (**Figure 2A**). More liver and kidney datasets are from

rats whereas more human and mouse datasets are available for the other tissues (**Figure 2B**).

There is also a species bias between the data repositories, with GEO covering more mouse

and human datasets whereas DrugMatrix and TG-GATEs curate more rat datasets (**Figure 2C**)

(http://mergeomics.research.idre.ucla.edu/PharmOmics/DrugSignatures/).

Comparison of PharmOmics with existing drug signature platforms revealed both shared

and unique properties between platforms

We compared PharmOmics with the existing CREEDS (18) and L1000FWD (31) databases

to assess the degree of agreement in drug signatures across different platforms

(**Supplementary methods**). As shown in **Supplementary Figure 1**, we found partial

agreement among the three databases. Despite the fact that overlaps in gene signatures

between databases were statistically significant for a majority of the drugs, the Jaccard scores

for signature overlaps were low (mostly between 0.02 and 0.04). We were not able to

systematically compare PharmOmics with CMAP for all drugs as done for CREEDS and

L1000FWD, because CMAP is an online query system (32) only feasible to query select

drugs. We searched atorvastatin in CMAP as an example and found that our PharmOmics

signature of atorvastatin from primary hepatocytes matched well with CMAP signatures of

the same drug from several human cancer cell lines including MCF7 (breast), HT29 (colon)

and VCAP (prostate), but surprisingly, not with HepG2 (hepatocellular carcinoma) which is

supposed to better match hepatocytes (**Supplementary Figure 2A**). The *in vivo* and *in vitro*

atorvastatin rat liver signatures from PharmOmics also showed different matching patterns

with CMAP signatures from the human cancer cell lines (**supplementary Figure 2B**). Taken

together, our results highlight important differences between PharmOmics and major existing

platforms. The differences between *in vivo* and *in vitro* drug signatures support the value of

obtaining more physiologically relevant, species-specific and tissue-specific drug signatures.

Tissue-specificity of drug signatures

To examine tissue-specificity of drug signatures, we analyzed the overall overlaps in gene

signatures for each drug across tissues and. Results were visualized with UpSetR package

(33). shown in **Figure 3A**, the overlap rate in the DEGs of the same drug between tissues and

organs is usually less than 5%, indicating a high variability in DEGs between tissues.

However, shared genes across tissues at DEG and pathway levels also exist. As an example,

we examined the HMGCR (β-Hydroxy β-methylglutaryl-CoA receptor) inhibitor atorvastatin

since its mechanism is well known and atorvastatin has been broadly tested in different

tissues under the human species label. We found that two DEGs (*TSC22D3, THBS1*) were

shared across tissues (**Figure 3B**). These genes are involved in extracellular matrix and

inflammation, suggesting these processes as common targets of atorvastatin across tissues.

Among the pathways shared across tissues, immune related pathways were shared between

blood cells and liver cells but not seen in prostate cells from the urogenital system (**Figure

3C, Supplementary Table 2**). Despite these overlaps, we found that the majority of the

DEGs were tissue-specific. Pathway analysis indicated that steroid synthesis and drug

metabolism pathways were altered mostly in liver, as expected, since the known target of

statin drugs is HMGCR, the rate limiting enzyme in cholesterol biosynthesis in liver. Blood

monocyte DEGs indicate changes in inflammation related pathways, while GPCR ligand

binding proteins were altered in prostate cancer cells. The tissue-specificity of drug

signatures revealed through our analysis supports tissue-specific therapeutic response or side

effects and the need for comprehensive maps of drug signatures across tissue systems as

implemented in the PharmOmics framework.

Species-specificity of drug signatures

We also compared gene signatures for each drug between species and found evidence for

species-specificity. Results were visualized with UpSetR package (33). As shown in **Figure**

3D, the pair-wise overlaps in DEGs between species for the same drug is generally lower

than 5%. Here we chose PPAR gamma receptor agonist rosiglitazone as an example because

this drug has datasets across human, rat, and mouse in PharmOmics and its mode of actions is

well-studied. As shown in **Figure 3E** and **3F**, nine genes (*CPT1C, AKR1B1, VNN1, ACSM3,*

CD36, CPT1A, PDK4, ZNF669, ADH1C) and several pathways including PPAR signaling

and fatty acid, triacylglycerol, and ketone body metabolism are consistently identified from

liver DEGs across species (**Supplementary Table 3**), supporting these as the major shared

pharmacological effectors of rosiglitazone across species. Among the species-specific

findings were rat-specific changes in bile acid and retinol metabolism, and adipocytokine

pathways in human datasets. The differences revealed between species highlight the

importance of understanding the physiological differences among model systems to facilitate

drug design with better translational potential. Our cross-species comparative studies also

highlight the urgent need to investigate drugs in multiple species, as only 21% of the drug-

tissue pairs (237 out of 1140) have data from two or more species.

Implementation of a web server to enable public access to PharmOmics

Previously we have developed an Apache web server Mergeomics (34), which allows

integrative analysis of multi-omics datasets to derive disease-associated pathways, networks,

and network regulators (http://mergeomics.research.idre.ucla.edu/). We implemented the

interactive PharmOmics web server with Rshiny (35) and deployed it on the same server

which we use to host Mergeomics, by enabling an instance of Apache to act as a front-end

proxy to the PharmOmics Shiny Server

(http://mergeomics.research.idre.ucla.edu/PharmOmics/). The PharmOmics web server hosts

the following main functions: i) drug signature query, visualization and download

(http://mergeomics.research.idre.ucla.edu/PharmOmics/DrugSignatures/), ii) gene network-

based drug repositioning using our species- and tissue-stratified signatures

(http://mergeomics.research.idre.ucla.edu/PharmOmics/Network_drug_repositioning/). iii)

Jaccard score-based drug repositioning

(http://mergeomics.research.idre.ucla.edu/PharmOmics/Jaccard_repositioning/) iv). Potential

ADR related signature based on meta-analyzing drugs sharing same ADR

(http://mergeomics.research.idre.ucla.edu/PharmOmics/ADRsignatures/). Due to higher

computational cost in the network-based drug repositioning application, a downloadable

shiny version is available at

http://mergeomics.research.idre.ucla.edu/Download/Network_drug_repositioning/.

Applications and Discussions

The value of PharmOmics to extract drug targets and target pathways

To explore the potential applications of our PharmOmics drug signatures, we evaluated

whether the species- and tissue-specific DEGs have the potential to retrieve drug targets and

target pathways. To this end, we first retrieved known drug targets for the drugs included in

PharmOmics from the DrugBanks database (36). Three different methods were used to

evaluate the potential of DEGs for drug target identification. The first method was based on

277 direct overlaps between DEGs and known drug targets, that is, if a known drug target was

278 directly retrieved from the DEG list it would be considered a positive hit. The second method

279 was based on overlaps in pathways, that is, whether the known pathway containing the drug

280 target was retrieved by the pathway analysis of the drug DEGs. The last method was based on

281 whether a known drug target is within the close neighborhood of drug DEGs in molecular

282 networks, including the STRING network (37) and tissue-specific Bayesian networks (BNs)

283 **(Supplementary methods)**.

285 As shown in **Table 1**, we found that all three drug target retrieval methods (gene overlap,

286 pathway overlap, network overlap) support that drug DEGs are more likely to detect true

287 drug targets than random genes. Compared to the gene overlap method which can retrieve

288 known drug targets at a 22.1% rate, the pathway- and network-based methods reached higher

289 target detection rates (62% for pathways, 40% for protein interaction network, 72% for liver

290 BN with matching liver DEGs, and 40% for kidney BN with mismatching liver DEGs),

291 although false positives also increased in the pathway and network-based approaches.

292 Notably, matching the tissue between DEGs and networks improved the target detection rate.

293 These results suggest that while known drug targets may not be directly detected among the

294 DEGs, pathway relatedness and network proximity in tissue specific network can be

295 informative of drug targets and can enhance the chance of target identification from drug

296 DEGs.

298 ***Drug repositioning analyses support that PharmOmics drug signatures can retrieve known***

299 ***therapeutic drugs for various diseases***

We hypothesized that if a drug is useful for treating a disease, the drug signatures and the

disease signatures will likely target similar pathways and therefore will be highly overlapping

or tightly connected in gene networks. To this end, we evaluated the ability of PharmOmics

to identify drugs for diseases based on overlaps in gene signatures matched by tissue using

the gene overlap-based measurement (Jaccard score) as well as a network proximity

measurement between drug and diseases genes which was previously applied to protein

interaction networks (5) **(Supplementary methods)**. Here, we used tissue-specific BN

networks by testing the mean shortest distance between network hubs of the drug DEGs and

disease genes. We further applied Mergeomics pipeline (38) which identifies hub genes on

network (key drivers) in this method. Using key drivers of drug gene signatures can reduce

noise and improve computing speed (**Supplementary Figure 3**) while performing similarly

to drug DEGs (**Figure 4A** and **4B**).

We selected hyperlipidemia as a disease test case, as multiple known drugs are available as

positive controls to evaluate the performance of drug repositioning. We have previously

developed a method Mergeomics (38), which can accurately extract disease related genes,

pathways, and networks (39,40). Because hyperlipidemia is most relevant to LDL and the

liver tissue, we used Mergeomics to retrieve LDL causal genes and pathways in the liver

tissue based on LDL GWAS and liver genetic regulation of gene expression (**Supplementary**

methods).

The Jaccard score method revealed that our PharmOmics drug signatures were able to

differentiate anti-hyperlipidemia drugs from other drugs with 79% AUC when using *in vivo*

rat liver signatures (Wilcoxon signed rank test p value=0.001), 56% AUC when using *in vitro*

liver signatures (non-significant), and 74% AUC when using the aggregated rat liver

signatures from both *in vivo* and *in vitro* settings (referred as combined, p=0.005) (**Figure**

4C). Among the top 5 predicted drugs are known LDL-lowering drug fluvastatin and other

drugs that may affect LDL including loratadine, sertraline, gefitinib, and paroxetine.

The network-based approach performed similarly as the direct gene overlap method above

(**Figure 4D**). The top 5 drugs predicted for hyperlipidemia based on the network approach

included fluvastatin, loratadine, mestranol-lynestrenol mixture, sertraline and estriol.

Between the network and Jaccard Score methods, 3 of the top 5 drugs from the two methods

overlapped, which are fluvastatin, loratadine and sertraline. Of these, fluvastatin is a known

HMG-CoA inhibitor for antihyperlipidemia therapy while oral contraceptives and sertraline

were linked with hyperlipidemia ADR (41–43). Moreover, among the 53 significant

predictions (out of 263 drugs) at $z < -2.52$ ($p < 0.01$) using our PharmOmics drug signatures

(**Supplementary Table 4**), we retrieved 6 out of the 10 approved anti-hyperlipidemia drugs

in the PharmOmics knowledgebase.

Using alternative hyperlipidemia signature from the Comparative Toxicogenomics database

(CTD) (16) to match with our PharmOmics drug signatures also showed that anti-

hyperlipidemia drugs had better Jaccard scores and better network repositioning z-scores

compared to other non-hyperlipidemia drugs. In addition, *in vivo* drug signatures again

showed better repositioning performance than the *in vitro* and combined drug signatures in

the CTD-based hyperlipidemia analysis (**Figure 4E**, **4F**).

We further tested the performance of PharmOmics in retrieving known drugs for other

diseases. Using CTD disease signatures for diabetes and hyperuricemia, we found

PharmOmics signatures of known diabetes drugs had better repositioning scores than those of

non-diabetes drugs (**Supplementary Figure 4**), and retrieved Benzbromarone as the top

ranked drug for hyperuricemia (**Supplementary Table 5**). These results support the general

utility of PharmOmics for drug repositioning.

Better or comparable drug repositioning performance of PharmOmics compared to

existing platforms

We compared the performance of our PharmOmics drug signatures in drug repositioning with

that of the well-established L1000 signatures, which were mainly based on *in vitro* cell

models. We found that PharmOmics signatures achieved better repositioning results than

L1000 signatures aggregated across dose and regimen for both Mergeomics and CTD

hyperlipidemia signatures using both the Jaccard score-based and network-based

repositioning methods (**Figure 4C-E**). Similarly, PharmOmics signatures also performed

better than the raw L1000 signatures when using the best performing dose and regimen

(Supplementary Figure 5). When compared with CMAP, PharmOmics also showed better

performance regardless of whether the aggregate signature across CMAP cell lines

(**Supplementary Figure 6A, Supplementary Table 4**) or the HEPG2-based signature was

used (**Supplementary Figure 6B, Supplementary Table 4**). Notably, PharmOmics was able

to retrieve 6 out of 10 (60%) of the known anti-hyperlipidemia drugs curated in the

PharmOmics database, while CMAP retrieved only 4 out of the 18 anti-hyperlipidemia drugs

examined in CMAP (22%). These results indicate the benefits of using *in vivo* signatures in

PharmOmics for drug repositioning to complement the existing cell-line based drug

repositioning tools to broaden drug repositioning capacity.

We further compared our PharmOmics gene signatures with the CREEDS database (18) for

drug repositioning. Using only the best scoring signature in CREEDS to represent each drug,

we found comparable drug-repositioning ability between the two databases (**Supplementary**

Figure 7). PharmOmics was able to reach faster repositioning since consensus signatures

across studies stratified by species and tissues were used instead of signatures from individual

datasets, and contained signatures for more drugs.

Use of PharmOmics to predict drugs for NAFLD

After establishing the performance of PharmOmics in drug repositioning using

hyperlipidemia, diabetes and hyperuricemia as test cases, we also applied PharmOmics to

predict potential drugs for NAFLD, for which there is currently no approved drugs. Using

NAFLD gene signatures from published studies (40,44) and the CTD database (16), we

predicted PPAR alpha agonists (fenofibrate and clofibrate) and PPAR gamma agonist

rosiglitazone to be strong matches to the NAFLD signatures (**Supplementary Table 6**).

PPAR agonists have been supported as potential drugs for NAFLD (45). The beneficial

effects of several PPAR alpha agonists on NAFLD have been demonstrated in *in vivo* models

(46,47), but clinical evidence is inconclusive at this point (48–50). On the other hand,

rosiglitazone and other PPAR gamma agonists showed beneficial effects in both rodent

models (50–52) and clinical trials (53–56). These lines of evidence substantiate the drug

repositioning potential of our PharmOmics tool. Additional drugs predicted include aspirin

which was also reported beneficial in mitigating NAFLD (57) and numerous other drugs

(**Supplementary Table 6**) which warrant experimental testing in future studies.

PharmOmics drug signatures can predict ADRs in liver

We hypothesize that drugs with the same ADR likely share common gene signatures driving

the specific ADR events. In this case, gene signature overlaps are expected between an ADR

term and the drugs with the ADR. We further hypothesized that network-based modeling can

better characterize hidden gene signatures to improve ADR prediction, which can be tested

based on network distance from drug signature genes to ADR genes. We collected ADR

information and ADR gene signatures were first aggregated across drugs with the same ADR

regardless of their pharmacological classes (**Supplementary methods**). We then tested

whether the ADR signatures can better predict drugs with the ADR than those with permuted

ADR labels, either based on direct gene overlaps or based on network distances (design

illustrated in **Supplementary Figure 8A**).

We focused on rat liver drug signatures as more than 200 drugs were available for this tissue.

The other tissues had more limited coverage of drugs, preventing a comprehensive

investigation. A total of 617 ADR terms each with more than six drugs were tested. For each

ADR, aggregate gene signatures were obtained from all constituent drugs with the ADR. The

gene overlap matrix between ADR signature and individual drug signatures, or the distance

matrix between ADR signatures and the key drivers of drug signatures in liver Bayesian

network were obtained. A random forest model was built with 10-fold-5-repeat cross

validation combined with smote algorithm (58) to accommodate unbalanced ADR samples

(i.e., fewer drugs with an ADR than those without the ADR, **Supplementary Methods**). For

the 617 ADR terms investigated, the distribution of the raw permuted p-values from random

forest models based on the gene overlapping matrix was similar to that based on the network

distance matrix (**Supplementary Figure 8B**, **8C**), both shifting towards the low p value end

(Kolomogorov-Smirnov test p-value <2.2e-16 against uniform distribution)**.** However, the

network-based method achieved better ADR predictions with lower FDR and higher accuracy,

with 21 ADR terms passing FDR < 0.05 (compared to 0 ADR term based on the gene overlap

matrix at the same threshold) and cross-validation accuracy > 0.99 (**Table 2**). Therefore,

PharmOmics signatures can also be used to derive ADR signatures which can predict drugs

with ADRs using a network-based approach. We were also able to retrieve top predictive

genes for many ADR terms based on the random forest models and predicted additional

drugs that may have these ADRs (**Table 2**).

Discussion

Here we present a publicly available tissue- and species-stratified drug signature database

along with an online web server, PharmOmics. PharmOmics fully utilizes published drug-

related transcriptomic datasets and provides unique tissue- and species-stratified gene

signatures that are more reflective of *in vivo* activities of drugs. We also developed a

framework for fast drug repositioning based on tissue-specific gene network models. We

demonstrate that PharmOmics can be used for molecular characterization of drugs,

comparisons of molecular activities between tissues and between species, and various

applications including target and target pathway identification, drug repurposing, and ADR

prediction.

Compared to the well-established CMAP and LINC1000 platforms, PharmOmics focuses

more on *in vivo* settings and likely captures more physiologically relevant drug signatures to

improve drug repositioning performance. Compared to a previous crowdsourcing effort,

CREEDS, which also utilizes publicly available drug datasets (18), our PharmOmics platform

included more curated drugs (343 in PharmOmics vs. 271 drugs in CREEDS) and involved

systematic tissue and species stratification and signature aggregation across different dosing

and treatment regimens to facilitate drug repositioning.

The use of tissue annotation with Brenda Tissue Ontology in PharmOmics helps normalize

organ labels and makes datasets more comparable. Although the tree structure in Brenda

Tissue Ontology may assign a cell or tissue to multiple systems (for example, breast cancer

cell line may be categorized as both epithelial and breast organ), we prioritized the tissue

terms based on the relevance to toxicology to make tissue assignments unique for each

dataset to reduce ambiguity. The unique tissue- and species-specific analyses implemented in

PharmOmics allows for comprehensive molecular insight into the actions of drug molecules

in individual tissues and species. Our results support that different species indeed have

unique drug responses in addition to shared features, thereby implicating that drug responses

and toxicity measures obtained in animal models may require caution before directly

translating the insights into humans. This notion agrees with the long-observed high failure

rate of drug development that has primarily relied on preclinical animal models and argues

for improved appreciation and understanding of inter-species differences in drug actions.

We demonstrate that PharmOmics can be used to retrieve drug targets and target pathways. In

general, gene signatures of drugs reflect cascades of downstream events after drug

administration. The initial drug target(s) may or may not be captured by drug DEGs due to

the lack of dynamic information in the DEGs. For example, changes in the direct drug targets

may have occurred transiently early upon treatment, but the change may have weakened or

disappeared at the time of transcriptome profiling. Nevertheless, our analyses support that

drug signatures are more likely to contain information about drug targets as well as target

pathways and networks. These findings support the value of using drug signatures to reflect,

at least partially, the molecular mechanisms of drugs.

Previous drug repositioning studies support that protein network-based approach is powerful

for drug repositioning. Here we show that combining the drug transcriptomic signatures in

PharmOmics, particularly the *in vivo* signatures, with tissue-specific gene regulatory

Bayesian networks and gene signatures of diseases can retrieve known therapeutic drugs and

predict potential new therapeutic avenues. Compared to other platforms, the use of tissue-

and species drug signatures along with network biology is a unique strength of PharmOmics,

which enables better performance and faster computation in repositioning.

We also demonstrate the potential of PharmOmics in ADR prediction. Although this effort is

limited by the small numbers of existing ADR datasets at each tissue level, we were able to

show the value of incorporating drug signatures with gene network models to perform ADR

predictions. The ADR signatures derived from our study and the predictive models from the

machine learning-based methods can be informative of mechanisms involved in ADRs,

although we acknowledge that further investigations are necessary to validate this potential.

There are several limitations in this study. First, our existing pipeline may not be able to

identify all of the drug datasets from GEO and ArrayExpress database. Along the same line,

the tissue coverage across the species is unbalanced, preventing a thorough cross-tissue and

cross-species comparison of drug signatures. We will keep refining the pipeline and update

our PharmOmics database periodically to include more datasets as they become available to

increase the coverage of datasets and drug signatures across tissues and species. In addition,

aggregating across different dosage and treatment durations may lose dose or time-specific

drug responses. However, we believe our approach can capture the most consistent drug

signatures regardless of treatment conditions, which are more representative of the general

properties of the drugs. The comparable performance of our platform with that of CREEDS,

which details dose and treatment regimens, in drug repurposing supports this notion. Lastly,

more comprehensive applications and experimental evaluation of PharmOmics in drug

repurposing for different disease areas and in ADR prediction are necessary in future studies.

Conclusion

We have established a new drug treatment signature database, PharmOmics, across different

species and tissues, which comprehensively captures the systems level *in vivo* activities of

drug molecules. In addition, we demonstrate the possible means to integrate these signatures

to address drug repositioning needs for disease treatment and to predict and characterize

ADRs. PharmOmics can complement other available drug signature databases to accelerate

drug development and toxicology research. The pipeline and our PharmOmics database will

be updated periodically to include newly available datasets to increase the coverage of these

datasets and drug signatures across tissues and species.

List of abbreviations

ADR   adverse drug reactions

CTD   comparative toxicogenomics database

KEGG   Kyoto Encyclopedia of Genes and Genomes

DEG   differential expressed genes

FDR   false discovery rate

wKDA   weighted key driver analysis

NAFLD   non-alcoholic fatty liver disease

LDL   low-density lipoprotein cholesterol

GWAS   genome-wide association study

BN   Bayesian gene regulatory network

ROC   Receiver operating characteristic

HMGCR β-Hydroxy β-methylglutaryl-CoA receptor

PPAR   Peroxisome proliferator-activated receptor

GPCR   G-protein coupled receptor

Declaration

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

Indexed dataset catalog, pre-computed gene signatures and pre-computed pathway

enrichments for individual drugs are deposited to and accessible through the PharmOmics

web server (http://mergeomics.research.idre.ucla.edu/PharmOmics/DrugSignatures/). We

also implemented functions for same-tissue between-species comparison and same-species

between-tissue comparison. Direct downloading the whole database or interested parts of the

database is also enabled. In addition, network-based drug repositioning analysis using human

and rat liver signatures is available at

http://mergeomics.research.idre.ucla.edu/PharmOmics/Network_drug_repositioning/. Jaccard

score-based drug repositioning analysis for all the tissues is available at

http://mergeomics.research.idre.ucla.edu/PharmOmics/Jaccard_repositioning/. Potential ADR

signatures aggregated from drugs having ADR is available at

http://mergeomics.research.idre.ucla.edu/PharmOmics/ADRsignatures/

Competing interests

The authors declare that they have no competing interests.

Funding

YC is supported by UCLA Eureka fellowship and Burroughs Wellcome Fund Inter-school

Training Program in Chronic Diseases. DA is supported by NIH-NCI National Cancer

Institute (T32CA201160), UCLA dissertation year fellowship and UCLA Hyde fellowship.

PA is supported by R01 NIH/NIEHS ES02748701, R21 NIH/NIAAA AA024889, and the

Burroughs Wellcome Foundation. XY is funded by NIH-DK104363, DK117850 and NIH-

NS103088.

Authors' contributions

YC curated and analyzed data, constructed database, designed and conducted application

studies and wrote the manuscript. DA provided supports in curation and analysis. GA, JG,

NZ, PP assisted with data curation. PA contributed to manuscript writing. XY designed and

supervised the research. All authors contributed to manuscript editing.

Acknowledgements

We thank Avi Ma'ayan (Icahn School of Medicine at Mount Sinai) for providing L1000FWD

and CREEDS data for comparison. We also thank Luoting Zhang for her contribution in data

curation for our research.

References:

1.   Morgan S, Grootendorst P, Lexchin J, Cunningham C, Greyson D. The cost of drug

development: a systematic review. Health Policy. 2011 Apr;100(1):4–17.

2.   Yu HWH. Bridging the translational gap: collaborative drug development and

dispelling the stigma of commercialization. Drug Discov Today [Internet]. 2016;21(2):299–

305. Available from: http://www.sciencedirect.com/science/article/pii/S1359644615003906

3.   Lin Z, Will Y. Evaluation of Drugs With Specific Organ Toxicities in Organ-Specific

Cell Lines. Toxicol Sci [Internet]. 2011 Dec 13;126(1):114–27. Available from:

https://doi.org/10.1093/toxsci/kfr339

4.   Denayer T, Stöhr T, Van Roy M. Animal models in translational medicine: Validation

and prediction. New Horizons Transl Med [Internet]. 2014;2(1):5–11. Available from:

http://www.sciencedirect.com/science/article/pii/S2307502314000022

5.   Cheng F, Desai RJ, Handy DE, Wang R, Schneeweiss S, Barabási A-L, et al.

Network-based approach to prediction and population-based validation of in silico drug

repurposing. Nat Commun [Internet]. 2018;9:2691. Available from:

http://www.ncbi.nlm.nih.gov/pmc/articles/PMC6043492/

6.   Guney E, Menche J, Vidal M, Barábasi A-L. Network-based in silico drug efficacy

screening. Nat Commun [Internet]. 2016;7:10331. Available from:

http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4740350/

7.   Barabasi AL, Gulbahce N, Loscalzo J. Network medicine: a network-based approach

to human disease. Nat Rev Genet. 2010/12/18. 2011;12(1):56–68.

8.   Subramanian A, Narayan R, Corsello SM, Peck DD, Natoli TE, Lu X, et al. A Next

Generation Connectivity Map: L1000 Platform and the First 1,000,000 Profiles. Cell.

2017/12/02. 2017;171(6):1437-1452.e17.

9.   Hall CJ, Wicker SM, Chien A-T, Tromp A, Lawrence LM, Sun X, et al.

Repositioning drugs for inflammatory disease – fishing for new anti-inflammatory agents.

Dis Model & Mech [Internet]. 2014;7(9):1069. Available from:

http://dmm.biologists.org/content/7/9/1069.abstract

10.   Corbett A, Pickett J, Burns A, Corcoran J, Dunnett SB, Edison P, et al. Drug

repositioning for Alzheimer's disease. Nat Rev Drug Discov [Internet]. 2012;11:833.

Available from: http://dx.doi.org/10.1038/nrd3869

11.   Godoy P, Bolt HM. Toxicogenomic-based approaches predicting liver toxicity in

vitro. Arch Toxicol [Internet]. 2012;86(8):1163–4. Available from:

https://doi.org/10.1007/s00204-012-0892-5

12.   Low Y, Uehara T, Minowa Y, Yamada H, Ohno Y, Urushidani T, et al. Predicting

Drug-Induced Hepatotoxicity Using QSAR and Toxicogenomics Approaches. Chem Res

Toxicol [Internet]. 2011;24(8):1251–62. Available from: https://doi.org/10.1021/tx200148a

13.   AbdulHameed MD, Ippolito DL, Stallings JD, Wallqvist A. Mining kidney

toxicogenomic data by using gene co-expression modules. BMC Genomics. 2016/10/12.

2016;17(1):790.

14.   Toutain P-L, Ferran A, Bousquet-Melou A. Species differences in pharmacokinetics

and pharmacodynamics. Handb Exp Pharmacol. 2010;(199):19–48.

15.   Wong CH, Siah KW, Lo AW. Estimation of clinical trial success rates and related

parameters. 2018 Jan 31; Available from: https://dx.doi.org/10.1093/biostatistics/kxx069

16.   Davis AP, Grondin CJ, Johnson RJ, Sciaky D, King BL, McMorran R, et al. The

Comparative Toxicogenomics Database: update 2017. Nucleic Acids Res [Internet].

2017;45(Database issue):D972–8. Available from:

http://www.ncbi.nlm.nih.gov/pmc/articles/PMC5210612/

17.   Wang Z, Clark NR, Ma'ayan A. Drug-induced adverse events prediction with the

LINCS L1000 data. Bioinformatics [Internet]. 2016;32(15):2338–45. Available from:

http://dx.doi.org/10.1093/bioinformatics/btw168

18.   Wang Z, Monteiro CD, Jagodnik KM, Fernandez NF, Gundersen GW, Rouillard AD,

et al. Extraction and analysis of signatures from the Gene Expression Omnibus by the crowd.

Nat Commun [Internet]. 2016;7:12846. Available from:

http://dx.doi.org/10.1038/ncomms12846

19.   Igarashi Y, Nakatsu N, Yamashita T, Ono A, Ohno Y, Urushidani T, et al. Open TG-

GATEs: a large-scale toxicogenomics database. Nucleic Acids Res. 2014/10/15.

2015;43(Database issue):D921-7.

20.   Barrett T, Wilhite SE, Ledoux P, Evangelista C, Kim IF, Tomashevsky M, et al.

NCBI GEO: archive for functional genomics data sets—update. Nucleic Acids Res [Internet].

2013;41(D1):D991–5. Available from: http://dx.doi.org/10.1093/nar/gks1193

21.   Kolesnikov N, Hastings E, Keays M, Melnichuk O, Tang YA, Williams E, et al.

ArrayExpress update—simplifying data submissions. Nucleic Acids Res [Internet].

2015;43(D1):D1113–6. Available from: http://dx.doi.org/10.1093/nar/gku1057

22.   Gremse M, Chang A, Schomburg I, Grote A, Scheer M, Ebeling C, et al. The

BRENDA Tissue Ontology (BTO): the first all-integrating ontology of all organisms for

enzyme sources. Nucleic Acids Res [Internet]. 2011;39(Database issue):D507–13. Available

from: http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3013802/

23.   Clark NR, Hu KS, Feldmann AS, Kou Y, Chen EY, Duan Q, et al. The characteristic

direction: a geometrical approach to identify differentially expressed genes. BMC

Bioinformatics [Internet]. 2014;15:79. Available from:

https://www.ncbi.nlm.nih.gov/pubmed/24650281

24.   Kolde R, Laur S, Adler P, Vilo J. Robust rank aggregation for gene list integration

and meta-analysis. Bioinformatics [Internet]. 2012;28(4):573–80. Available from:

https://www.ncbi.nlm.nih.gov/pubmed/22247279

25.   Kuleshov M V, Jones MR, Rouillard AD, Fernandez NF, Duan Q, Wang Z, et al.

Enrichr: a comprehensive gene set enrichment analysis web server 2016 update. Nucleic

Acids Res [Internet]. 2016;44(Web Server issue):W90–7. Available from:

http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4987924/

26.   Kanehisa M, Furumichi M, Tanabe M, Sato Y, Morishima K. KEGG: new

perspectives on genomes, pathways, diseases and drugs. Nucleic Acids Res [Internet].

2017;45(Database issue):D353–61. Available from:

http://www.ncbi.nlm.nih.gov/pmc/articles/PMC5210567/

27.   Expansion of the Gene Ontology knowledgebase and resources. Nucleic Acids Res.

2016/12/03. 2017;45(D1):D331-d338.

28.   Fabregat A, Jupe S, Matthews L, Sidiropoulos K, Gillespie M, Garapati P, et al. The

Reactome Pathway Knowledgebase. Nucleic Acids Res. 2017/11/18. 2018;46(D1):D649-

d655.

29.   Nishimura D. BioCarta. Biotech Softw Internet Rep [Internet]. 2001;2(3):117–20.

Available from: https://doi.org/10.1089/152791601750294344

30.   Slenter DN, Kutmon M, Hanspers K, Riutta A, Windsor J, Nunes N, et al.

WikiPathways: a multifaceted pathway database bridging metabolomics to other omics

research. Nucleic Acids Res [Internet]. 2018;46(Database issue):D661–7. Available from:

http://www.ncbi.nlm.nih.gov/pmc/articles/PMC5753270/

31.　Wang Z, Lachmann A, Keenan AB, Ma'ayan A. L1000FWD: fireworks visualization

of drug-induced transcriptomic signatures. Bioinformatics [Internet]. 2018 Jun

15;34(12):2150–2. Available from: http://dx.doi.org/10.1093/bioinformatics/bty060

32.　Lamb J, Crawford ED, Peck D, Modell JW, Blat IC, Wrobel MJ, et al. The

Connectivity Map: using gene-expression signatures to connect small molecules, genes, and

disease. Science (80-). 2006/09/30. 2006;313(5795):1929–35.

33.　Conway JR, Lex A, Gehlenborg N. UpSetR: an R package for the visualization of

intersecting sets and their properties. Bioinformatics [Internet]. 2017;33(18):2938–40.

Available from: http://dx.doi.org/10.1093/bioinformatics/btx364

34.　Arneson D, Bhattacharya A, Shu L, Mäkinen V-P, Yang X. Mergeomics: a web

server for identifying pathological pathways, networks, and key regulators via

multidimensional data integration. BMC Genomics [Internet]. 2016;17(1):722. Available

from: https://doi.org/10.1186/s12864-016-3057-8

35.　Chang W, Cheng J, Allaire JJ, Xie Y, McPherson J. shiny: Web Application

Framework for R [Internet]. 2017. Available from: https://cran.r-project.org/package=shiny

36.　Wishart DS, Feunang YD, Guo AC, Lo EJ, Marcu A, Grant JR, et al. DrugBank 5.0: a

major update to the DrugBank database for 2018. Nucleic Acids Res. 2017/11/11.

2018;46(D1):D1074-d1082.

37.　Szklarczyk D, Morris JH, Cook H, Kuhn M, Wyder S, Simonovic M, et al. The

STRING database in 2017: quality-controlled protein–protein association networks, made

broadly accessible. Nucleic Acids Res [Internet]. 2017;45(Database issue):D362–8. Available

from: http://www.ncbi.nlm.nih.gov/pmc/articles/PMC5210637/

38.   Shu L, Zhao Y, Kurt Z, Byars SG, Tukiainen T, Kettunen J, et al. Mergeomics:

multidimensional data integration to identify pathogenic perturbations to biological systems.

BMC Genomics [Internet]. 2016;17(1):874. Available from: https://doi.org/10.1186/s12864-

016-3198-9

39.   Shu L, Chan KHK, Zhang G, Huan T, Kurt Z, Zhao Y, et al. Shared genetic

regulatory networks for cardiovascular disease and type 2 diabetes in multiple populations of

diverse ethnicities in the United States. PLoS Genet. 2017/09/29. 2017;13(9):e1007040.

40.   Chella Krishnan K, Kurt Z, Barrere-Cain R, Sabir S, Das A, Floyd R, et al.

Integration of Multi-omics Data from Mouse Diversity Panel Highlights Mitochondrial

Dysfunction in Non-alcoholic Fatty Liver Disease. Cell Syst. 2018/01/24. 2018;6(1):103-

115.e7.

41.   Kesim M, Tiryaki A, Kadioglu M, Muci E, Kalyoncu NI, Yaris E. The effects of

sertraline on blood lipids, glucose, insulin and HBA1C levels: A prospective clinical trial on

depressive patients. J Res Med Sci [Internet]. 2011 Dec;16(12):1525–31. Available from:

https://www.ncbi.nlm.nih.gov/pubmed/22973359

42.   Naz F, Jyoti S, Akhtar N, Afzal M, Siddique YH. Lipid profile of women using oral

contraceptive pills. Pakistan J Biol Sci PJBS. 2012 Oct;15(19):947–50.

43.   Connelly PW, Stachenko S, MacLean DR, Petrasovits A, Little JA. The prevalence of

hyperlipidemia in women and its association with use of oral contraceptives, sex hormone

replacement therapy and nonlipid coronary artery disease risk factors. Canadian Heart Health

Surveys Research Group. Can J Cardiol. 1999 Apr;15(4):419–27.

44.   Kurt Z, Barrere-Cain R, LaGuardia J, Mehrabian M, Pan C, Hui ST, et al. Tissue-

specific pathways and networks underlying sexual dimorphism in non-alcoholic fatty liver

disease. Biol Sex Differ. 2018 Oct;9(1):46.

45. Liss KHH, Finck BN. PPARs and nonalcoholic fatty liver disease. Biochimie. 2017

May;136:65–74.

46. Abd El-Haleim EA, Bahgat AK, Saleh S. Resveratrol and fenofibrate ameliorate

fructose-induced nonalcoholic steatohepatitis by modulation of genes expression. World J

Gastroenterol. 2016 Mar;22(10):2931–48.

47. Zhang N, Lu Y, Shen X, Bao Y, Cheng J, Chen L, et al. Fenofibrate treatment

attenuated chronic endoplasmic reticulum stress in the liver of nonalcoholic fatty liver

disease mice. Pharmacology. 2015;95(3–4):173–80.

48. Laurin J, Lindor KD, Crippin JS, Gossard A, Gores GJ, Ludwig J, et al.

Ursodeoxycholic acid or clofibrate in the treatment of non-alcohol-induced steatohepatitis: a

pilot study. Hepatology. 1996 Jun;23(6):1464–7.

49. Fernandez-Miranda C, Perez-Carreras M, Colina F, Lopez-Alonso G, Vargas C,

Solis-Herruzo JA. A pilot trial of fenofibrate for the treatment of non-alcoholic fatty liver

disease. Dig Liver Dis. 2008 Mar;40(3):200–5.

50. Hertz R, Bar-Tana J. Peroxisome proliferator-activated receptor (PPAR) alpha

activation and its consequences in humans. Toxicol Lett. 1998 Dec;102–103:85–90.

51. Kawaguchi K, Sakaida I, Tsuchiya M, Omori K, Takami T, Okita K. Pioglitazone

prevents hepatic steatosis, fibrosis, and enzyme-altered lesions in rat liver cirrhosis induced

by a choline-deficient L-amino acid-defined diet. Biochem Biophys Res Commun. 2004

Feb;315(1):187–95.

52. Nan Y-M, Fu N, Wu W-J, Liang B-L, Wang R-Q, Zhao S-X, et al. Rosiglitazone

prevents nutritional fibrosis and steatohepatitis in mice. Scand J Gastroenterol.

2009;44(3):358–65.

53.   Neuschwander-Tetri BA, Brunt EM, Wehmeier KR, Oliver D, Bacon BR. Improved

nonalcoholic steatohepatitis after 48 weeks of treatment with the PPAR-gamma ligand

rosiglitazone. Hepatology. 2003 Oct;38(4):1008–17.

54.   Torres DM, Jones FJ, Shaw JC, Williams CD, Ward JA, Harrison SA. Rosiglitazone

versus rosiglitazone and metformin versus rosiglitazone and losartan in the treatment of

nonalcoholic steatohepatitis in humans: a 12-month randomized, prospective, open- label trial.

Hepatology. 2011 Nov;54(5):1631–9.

55.   Ratziu V, Charlotte F, Bernhardt C, Giral P, Halbron M, Lenaour G, et al. Long-term

efficacy of rosiglitazone in nonalcoholic steatohepatitis: results of the fatty liver improvement

by rosiglitazone therapy (FLIRT 2) extension trial. Hepatology. 2010 Feb;51(2):445–53.

56.   Ratziu V, Giral P, Jacqueminet S, Charlotte F, Hartemann-Heurtier A, Serfaty L, et al.

Rosiglitazone for nonalcoholic steatohepatitis: one-year results of the randomized placebo-

controlled Fatty Liver Improvement with Rosiglitazone Therapy (FLIRT) Trial.

Gastroenterology. 2008 Jul;135(1):100–10.

57.   Simon TG, Henson J, Osganian S, Masia R, Chan AT, Chung RT, et al. Daily Aspirin

Use Associated With Reduced Risk For Fibrosis Progression In Patients With Nonalcoholic

Fatty Liver Disease. Clin Gastroenterol Hepatol. 2019 May;

58.   Chawla N V., Bowyer KW, Hall LO, Kegelmeyer WP. SMOTE: Synthetic Minority

Over-sampling Technique. J Artif Intell Res. 2016;16:321–57.

Tables

Table 1 The capacity of PharmOmics drug signatures to retrieve known drug targets based on

various methods.

Method	Median gene number count	Target detection rate	Target detection rate from randomized gene symbols
Jaccard gene overlap	427	22.1%	10.5% (p < 0.001)
Pathway (GOBP and KEGG)	1402	62.4%	26.2% (p < 0.001)
Network (one-edge neighbors of DEGs in STRING network)	1143	41.8%	17.4% (p < 0.001)
Network (one-edge neighbors of liver DEGs in liver BN)	4258	71.8%	44.7% (p < 0.001)
Network (one-edge neighbors of liver DEGs in mismatched kidney BN)	861	40.5%	19.2% (p < 0.001)

771   **Table 2** Predictive ADR signatures in rat liver, top predictive genes and new drugs predicted

772   to have the corresponding ADRs.

ADR	FDR	Top predictive genes	New drugs predicted to have ADR
Abdominal cramps	1.8e-2	*Atg10, Fgf1, Plekhb1, Gne, Fabp5*	
Acute generalised exanthematous pustulosis	1.8e-2	*Ptpn4, Arhgdib, Slc26a8, Acly, Itgal*	Cyproterone
Bursitis	1.8e-2	*Esm1, Lgals2, Zfp395, Fnip2, Gpx3*	Chloroquine, Estriol
Calculus of kidney	1.5e-2	*Cenpf, Mvd, Esco2, Top2a, Clec4a1*	Anastrozole, Ethylestrenol
Cardiovascular disorder	4.9e-2	*Chd1l, Gfra3, Rcan1, Esm1, Dnase2b*	
Folliculitis	1.8e-2	*Ahr, Comt, Cds1, Nucb2, Ccnb2*	
Generalised oedema	1.8e-2	*Slc15a1, Fads2, Fosl1, Clec4a3, Cxcl1*	Cyclophosphamide, Tolbutamide
Hostility	4.6e-2	*Otc, Col4a2, Gbp2, Zfp395, Sema3e*	Chloroquine, Estriol
Hyperreflexia	1e-2	*S100a6, Il9r, Ddr1, Mlc1, Dcn*	
Hyperuricaemia	1.8e-2	*Psat1, Fnip2, Syngr1, Gpnmb, Slc22a8*	Benzethonium chloride, Tolbutamide
Lymphocytosis	1.8e-2	*Sema3c, Larp4, S100a6, Tmem132a, Pdlim2*	
Meningitis	1.8e-2	*Tradd, Chka, Cldn1, Usp2, Nampt*	Pramoxine hydrochloride
Nail disorder	8e-3	*Bach2, Lcp1, Coro1a, Rhpn2, Hao2*	Dexamethasone, Diazepam
Nephritis	2e-3	*Dusp6, Tgfb3, Cdkn1a, Jup, Nr1d1*	Cimetidine, Flutamide
Nephrotoxicity	4.6e-3	*Cidea, Prlr, Msl3l2, Dmbt1, Slc25a30*	Ranitidine, Tacrine
Pelvic pain	1.6e-2	*Paqr9, Top2a, Kif20a, Gpx2, Gpx3*	Chloroquine, Estriol
Pruritus	1.8e-2	*Ncapd2, Cited2, Dusp9, Mpzl1, Klf15*	
Psoriasis	1.8e-2	*Hao2, B3galt1, Gdf15, Tekt1, Acsm5*	Nefazodone, Tolbutamide
Slurred speech	1.8e-2	*Ppp1r3b, Adora2b, Paqr9, Pttg1, Slc6a13*	
Tenosynovitis	2e-3	*Dpys, Bub1b, Gcat, Ldhd, Aldoc*	
Thrombosis	1.8e-2	*Fnip2, Rhbdd2, Atf3, Fgf1, Pdlim2*	Acarbose, Tolbutamide

Figure legends

Figure 1 PharmOmics data processing pipeline. A list of clinically relevant drugs was

selected from KEGG database, followed by searching across GEO, ArrayExpress, TG-

GATEs and drugMatrix data repositories for datasets where drug treatment and control

samples were available. Individual transcriptome datasets were analyzed with the GeoDE

method to obtain ranked gene list for each treatment experiment. Datasets were annotated

with tissue and species information and meta-analyzed within tissue, species, and drug to

obtain genes that are altered consistently across datasets for each drug as "drug signatures".

These signatures were used to conduct drug repositioning analysis and ADR prediction

analysis based on direct gene overlaps or a gene network-based approach.

Figure 2. Summary of available datasets based on data sources, tissues, and species. Y-

axis indicates unique dataset counts and X-axis indicates (A) tissue and data resources, (B)

tissue and species, and (C) data resources and species.

Figure 3. Cross-tissue and cross-species comparisons of drug signatures in PharmOmics. (A)

Distribution of drug signature overlap percentages between tissue pairs in matching species

from PharmOmics. Arrow points to the pairs of tissues for drugs with high overlap in gene

signatures. (B) Upset plot of cross-tissue comparison for atorvastatin signatures genes. Y-axis

indicates number of genes. (C) Upset plot of cross-tissue comparison for pathways enriched

in atorvastatin signatures. Y-axis indicates number of pathways. (D) Distribution of drug

signature overlap percentages between pairs of species for matching tissues from

PharmOmics. Arrow points to the species pair with high gene signature overlap for a

matching drug. (E) Upset plot of cross-species comparison for rosiglitazone liver gene

signatures. (F) Upset plot of cross-species comparison for pathways enriched in rosiglitazone

liver signatures. Pairs of tissues with shared drug signature genes or pathways are connected

with black vertical lines in the bottom portion of the Upset plots.

Figure 4. AUC plot for drug repositioning for hyperlipidemia, comparing anti-

hyperlipidemia drugs against other drugs under different testing conditions. Hyperlipidemia

signatures were derived from either the Mergeomics tool or the CTD database. We first

compared key drivers (KD) of DEGs vs DEGs on network-based drug repositioning using

disease signatures from (A) MergeOmics and (B) CTD. The drug repositioning performance

of PharmOmics was then compared with L1000 (C-F). Only matching drugs between L1000

and PharmOmics were used to make results comparable. (C) AUC plot based on the Jaccard

score method and the hyperlipidemia signature from MergeOmics. (D) AUC plot based on

the gene network method and the hyperlipidemia signature from MergeOmics. (E) AUC plot

based on the Jaccard score method and the hyperlipidemia signature from CTD. (F) AUC plot

based on the gene network repositioning method and the hyperlipidemia signature from CTD.

Wilcoxon signed rank test was used to calculate significance between Jaccard/Network z-

scores between groups. *, **, *** inducates $p < 0.05$, $p < 0.01$ and $p < 0.001$ repectively.

822   **Figures**

823   **Figure 1.**

829 **Figure 2.**

A

B

C

    Figure 3.

Figure 4.