Transposon expression in the *Drosophila* brain is driven by neighbouring genes and diversifies the neural transcriptome

Christoph D. Treiber* & Scott Waddell*

Centre for Neural Circuits and Behaviour, University of Oxford, Tinsley Building, Mansfield Road, Oxford OX1 3SR, UK

*Corresponding authors.

Email: christoph.treiber@cncb.ox.ac.uk, scott.waddell@cncb.ox.ac.uk

Abstract

Somatic transposition in neural tissue could contribute to neuropathology and individuality, but its prevalence is debated. We used single-cell mRNA sequencing to map transposon expression in the *Drosophila* midbrain. We found that neural transposon expression is driven by cellular genes. Every expressed transposon is resident in at least one cellular gene with a matching expression pattern. A new long-read RNA sequencing approach revealed that coexpression is a physical link in the form of abundant chimeric transposon-gene mRNAs. We identified 148 genes where transposons introduce cryptic splice sites into the nascent transcript and thereby produce many additional mRNAs. Some genes exclusively produce chimeric mRNAs with transposon sequence and on average transposon-gene chimeras account for 20% of the mRNAs produced from a given gene. Transposons therefore significantly expand the neural transcriptome. We propose that chimeric mRNAs produced by splicing into polymorphic transposons may contribute to functional differences between individual cells and animals.

Introduction

Transposons comprise almost half of every eukaryote genome (Britten and Kohne, 1968; International Human Genome Sequencing Consortium et al., 2001; Ketchum et al., 2000) and their mobilization in the germline contributes to chromosome evolution. Non-heritable *de novo* transposon activity in neural tissue has been proposed to contribute to functional heterogeneity in the brain and to neurological disease (Baillie et al., 2011; Coufal et al., 2009; Evrony et al., 2012; Kazazian, 2011; Muotri et al., 2005). However, it is difficult to faithfully map rare *de novo* transposon insertions using whole-genome DNA sequencing (Baillie et al., 2011; Evrony et al., 2012, 2016; Perrat et al., 2013; Treiber and Waddell, 2017; Upton et al., 2015). A growing number of studies have therefore correlated the development of neurodegeneration in animal models with changes in transposon expression (Guo et al., 2018; Krug et al., 2017; Li et al., 2013; Li et al., 2012; Sun et al., 2018). Using expression as a proxy for mobility could be misleading because high-level transposon expression does not appear to result in elevated *de-novo* somatic transposition in the brain (Evrony et al., 2012, 2016; Treiber and Waddell, 2017). It is therefore important to understand what controls the expression of transposon-derived sequences in the brain and whether their elevated expression relates to neural function.

An early study of the human LINE-1 (L1) promoter demonstrated that its activity was heavily influenced by flanking cellular sequences, and concluded that expression of a given L1 depended on its location in the genome (Lavie et al., 2004). Such a locus specific model could more generally explain the apparent cell-type restricted nature of transposon expression and mobilization in the brain. So far, studies of somatic transposon expression have either focused on single transposon families or have been based on bulk sequencing of tissues or cultured cells (Chung et al., 2019; Faulkner et al., 2009; Li et al., 2013; Philippe et al., 2016; Rangwala et al., 2009). However, answering this question on a brain- and genome-wide scale requires a means to relate the cellular expression of each transposon in the genome to that of their neighbouring genes. Recent technical developments in whole-genome DNA sequencing and high-throughput single-cell transcriptomics of complex tissues now make this possible (Macosko et al., 2015).

Here we used single-cell transcriptomics to map transposon expression to individual cells in the *Drosophila* midbrain. We found that many transposons are expressed with cell-specificity that is often highly correlated to that of a neighbouring gene. A more detailed analysis revealed that >90% of transposon expression can be linked to co-expression with their host gene, indicating that these genes are the main driver of somatic transposon expression. Long-read sequencing showed that the transposon and neighbouring gene are alternatively

spliced together becoming part of the same chimeric mRNAs. Sometimes all mRNAs

produced from a particular gene in which a transposon resides include transposon

sequence. Therefore, transposons produce genome-wide diversification of cellular

transcripts. Analysis of sequencing data produced from other fly strains demonstrated large

differences in their chimeric transcriptomes. Inter-strain and individual differences in

transposon complement therefore constrain the cellular specificity and likelihood of

transposon-directed pathology.

Results

Single-cell transcriptomics reveals cell-type restricted transposon expression

The *Drosophila* genome contains 112 families of transposons and the number of an individual types varies from a few to hundreds of copies (Kaminker et al., 2002). Conventional single-cell RNAseq (scRNAseq) analysis pipelines typically discard sequencing reads that align to multiple genomic loci and so they overlook transposon expression. We therefore devised an analysis pipeline to map the expression of all transposons within scRNAseq data. We masked all repetitive sequences in the reference genome and then added a single copy of the consensus sequence for every known transposon to the masked genome. In essence this produces a *Drosophila* reference genome with one copy of each type of transposon. We first used this modified reference genome to map transposon expression onto single cells of the midbrain in a fly strain expressing mCherry in $\alpha\beta$ Kenyon cells (KCs) of the mushroom body (MB); from here called $\alpha\beta$Cherry flies. We found evidence for expression of both the sense and the antisense strand of most transposons, which comprised 76.2 and 23.8% (+/- 1.7% SD) of all transposon expression, respectively (Figure 1 – figure supplement 1). We first performed principal component decomposition of cellular genes and clustered cells from the midbrain by constructing a k-Nearest-Neighbour graph on the Euclidean distances in the PCA space, optimizing the modularity using the Louvain algorithm (Butler et al., 2018). This analysis grouped cells into many discrete clusters. We next assigned many of these clusters to cell types in the midbrain using the expression patterns of known marker genes (Croset et al., 2018) (Figure 1a). Displaying the expression of individual types of transposons on the cluster plot revealed that some transposons are up-regulated in specific cell types. For example, the long-terminal repeat (LTR) retrotransposons COPIA and NOMAD showed elevated expression in the $\alpha\beta$, $\alpha'\beta'$ and γ Kenyon Cells (KCs) classes (Figure 1b, first graph) and $\alpha'\beta'$ KCs (Figure 1c, first graph), respectively. Other LTR retrotransposons such as MICROPIA were upregulated in the ellipsoid body (Figure 1d, first graph) whereas BLOOD and 412 were higher in glia (Figure 1e,f, first graphs).

Transposon expression correlates with that of cellular genes they are inserted within.

Transposons could be elevated in specific cell types because they are inserted in genes that are highly expressed in the same cells. To test this hypothesis, we re-used our previously published high-coverage gDNA sequence of $\alpha\beta$Cherry flies. We mapped all the germline transposon insertions in these flies using TEchim, a new custom-built transposon analysis program. TEchim first generates long nucleotide contigs from either gDNA or cDNA sequencing reads, then creates in-silico paired-end reads and screens them for cases where one in-silico end maps to a cellular gene and the mate read maps to a specific transposon.

Since these paired-end reads are derived from contiguous sequences, TEchim permits one to subsequently refer back to the original long reads to determine the precise nucleotide sequence of the transposon-gene breakpoints. Using TEchim, we found copies of COPIA, NOMAD, MICROPIA, BLOOD and 412 inside the genes *twin-of-eyeless* (*toy*), *mushroom-body expressed* (*mub*), *Rhodopsin-7* (*Rh7*), *CG32521* and *tequila* (*teq*), respectively. The expression of each of these genes mirrored the expression pattern of the transposon they harboured (Figure 1b-f, second graphs). The expression of these transposons in the brain of $\alpha\beta$Cherry flies therefore appears to be driven by their relevant host genes.

We next tested whether all transposons exhibit patterned expression throughout the midbrain. We re-clustered the single-cell data of the fly using only transposon expression. This generated 78 cell clusters that mostly contained cells from all 8 biological replicates in the data (Figure 1g, Figure 1 – figure supplement 2). This result suggests that transposon expression is stereotyped across samples derived from different flies collected from the same strain. We then analysed the expression of cellular genes across the transposon clusters and found many clusters also preferentially expressed certain genes. For example, the cluster of cells expressing the LTR of STALKER2 was enriched for cells that also expressed the *Hsp70Bc* gene (Figure 1h), and cells in the DOC-positive cluster showed increased expression of *Furin1* (*Fur1*). By referring back to the gDNA, we found that $\alpha\beta$Cherry flies harbour a copy of STALKER2 within *Hsp70Bc* and a copy of the LINE-like DOC element inside *Fur1*. Again, these data suggest that expression of STALKER2 and DOC is driven by a neighbouring gene.

Quantitative analysis reveals high fidelity transposon-gene co-expression

Our gDNA analysis also revealed many transposon insertions inside genes that were more broadly expressed across the brain. In total, we identified 1952 germline transposon insertions within genes. Of these, 881 cases were inserted in the sense direction to the open reading frame of the gene and 1071 were in the antisense orientation (Supplementary Table 1). To quantify the correlated expression of transposons and cellular genes we devised a method based on the established Hardy-Weinberg principle for quantifying linkage equilibrium of two alleles in population genetics (Lewontin and Kojima, 1960) (Figure 2a). We first binarized our scRNAseq data to generate the equivalent of bi-allelic traits in a population. We then calculated the proportion of cells expressing a specific transposon, multiplied it by the proportion of cells expressing a certain gene, and then subtracted this value from the proportion of cells that expressed both the transposon and the gene. We termed this value the Coexpression Disequilibrium, CD. We also normalized these CD values to account for the variable abundance of each transposon and gene in every

transposon-gene pair and repeated the analysis for all transposon-transposon and gene-

gene pairs. These normalized values were then ranked within each of the 8 biological

replicates. P-values were corrected for multiple comparisons and describe the probability

that a transposon-gene pair would have such a highly ranked CD value across multiple

replicates if they were expressed independently.

We combined the list of all germline transposon insertions in $\alpha\beta$Cherry flies with the scRNA

seq data generated from the same population of flies and calculated the CD values between

every transposon and the gene in which it was inserted. We tested every transposon that

contributed at least 0.1% of the overall transposon expression in our midbrain $\alpha\beta$Cherry fly

samples. This cut-off left 59 different transposons, 34 of which were expressed in both the

sense- and the antisense direction, 22 only in sense, and 3 only in antisense (Figure 2b,

Supplementary Table 2). For 56 of these transposons we found at least one copy inside a

gene that exhibited a correlated expression pattern (Benjamini-Hochberg corrected p-val

>0.05). For those cases where the transposon was inserted in the same orientation as the

transcription unit of the gene the expression of the sense strand of the transposon correlated

to that of the gene. In contrast, the antisense strand of reverse orientation transposons was

correlated with the host gene. Importantly, the average number of correlated genes for each

transposon was significantly less if equivalent values were calculated using random

assignment of transposons to host genes (Figure 2c). These analyses therefore

demonstrate that the genomic locus strongly influences the expression patterns of almost all

transposons in the fly brain. We did not identify a neighbouring gene with a correlated

expression pattern for the transposons TART-A, P-element and HMSBEAGLE. This is

expected for TART-A, which is a telomeric retrotransposon, and for P-element, which is a

remnant of transgenic intervention in $\alpha\beta$Cherry flies. It is conceivable that the HMSBEAGLE

retrotransposon is the only element that is expressed independently of a cellular gene.

However, despite our high coverage sequence of $\alpha\beta$Cherry flies we may have missed a

germline HMSBEAGLE insertion that sits near a gene that is driving its expression.

Transposons are exonised into cellular mRNAs

Recent work has shown that mRNAs from the *Arc* gene contain transposon-like sequence in

the coding sequence and 3′ UTR (Pastuzyn et al., 2018; Zhang et al., 2015). We therefore

tested whether chimeric mRNAs might occur more broadly and extend to all transposons.

We extracted mRNA from $\alpha\beta$Cherry fly heads and generated 250 basepair long reads which

were screened using TEchim for chimeric reads. We also incorporated a function in TEchim

that maintains strand-specificity of the input reads which enabled us to unambiguously

assign chimera to cellular genes. This analysis revealed that a large number of transposons

inside introns lead to the formation of chimeric mRNAs. In total, we found chimeric mRNA from 887 transposon insertions (Figure 3a, Supplementary Table 3). Chimera included sequences from LTR, LINE-like and DNA transposons attached to mRNAs from genes involved in a broad range of biological processes. For example, we found sequence from the LTR-retrotransposon GYPSY in transcripts of the ubiquitin gene *Ubi-P5E* and of the neuron specific ubiquitin ligase *highwire* (*hiw*), the non-LTR element DOC in *Fur1*, encoding a synaptic membrane bound protease, and the TIR element HOBO attached to transcripts from *Shaker*, which encodes a voltage-gated potassium channel (Izquierdo, 1994; Kaplan and Trout, 1969; Roebroek et al., 1991; Wan et al., 2000).

Previous studies of chimeric sequencing reads have established that *in vitro* amplification of genetic material often leads to chimeric amplification artefacts (Evrony et al., 2016; Treiber and Waddell, 2017). It was thus important to account for similar errors in our data. We therefore calculated the rate of these artefacts in our mRNA data by selecting 10 sets of 167 exons (with each set providing a number of sequences corresponding to the 112 different transposon types and 55 LTRs) with matching expression levels in the brain. These exons lack the ability to relocate in gDNA so we refer to them as immobile genetic elements (IGEs). Since IGEs should only occur as single copies in the gDNA from $\alpha\beta$Cherry, chimeric reads between IGEs and other genes most likely represent amplification artefacts. We found the rate of generating IGE chimeras was directly correlated to the expression level of the IGE and the gene that it formed a chimeric molecule with. Critically, the IGE chimera rate was substantially lower than that of chimera formed between genes and transposons. We therefore used the rate of IGE chimera to filter the transposon chimera detected using TEchim, defining a false discovery rate (FDR) of 0.1% (Figure 3b, Supplementary Table 3). All examples of chimeric transcripts that are presented in detail in this study have supporting evidence that exceeds this FDR.

LTR retrotransposon expression is predominantly non-autonomously

Given that transposon expression was highly correlated with at least one neighbouring gene, we hypothesized that all transposon expression in the brain might occur as co-expression with cellular genes. To test this, we focused on LTR retrotransposons. We quantified the number of reads that spanned an LTR-gene breakpoint, and compared it to the number of reads that crossed the LTR-transposon breakpoint within the transposon (Figure 4a). Autonomously expressed, full-length transposons only generate the latter type of read, whilst non-autonomous expression should generate both kinds of reads, at varying proportions. We found that around 95% of the highly expressed LTR transposons produced a roughly equivalent number of gene-transposon and LTR-transposon reads, suggesting that LTR

217 transposons are expressed as chimeras with cellular genes, rather than being autonomously

218 expressed (Figure 4b, Supplementary Table 4).

Many transposons introduce cryptic alternative splice-sites into cellular genes

Given the abundance of reads that span a gene-transposon breakpoint, we next investigated

the structure of these transcripts in more detail. Our transposon mapping identified 887 loci

with a germline transposon insertion in the intron of a gene. For each of these we found

mRNA molecules where one section mapped to the beginning or end of the transposon and

the other section corresponded to the flanking intronic sequence. These reads could

represent nascent unspliced chimeric pre-mRNAs from which transposon-derived sequence

could be removed by splicing to yield intact host mRNAs, and full-length transposon

sequences. However, we also found 148 examples where breakpoint-spanning reads

indicate that specific sections of transposon sequence are spliced into host-gene transcripts

(Supplementary Table 5).

Analysis of the breakpoints inside transposons at these 148 sites revealed that chimera are

formed at conserved locations in each type of transposon. For example, in cases where an

antisense ROO resided within an intron, we found transcripts where the 3´-end of an

upstream exon had formed a new phosphate bond to a section of ROO at positions 5460

and 2094 at 19 and 3 different genomic loci, respectively, and also at several additional

breakpoints with lower frequency (Figure 5a,b). In addition, we identified transcripts where

sections of ROO were bound to the 5´-end of a downstream exon. We found breakpoints at

position 5191 from 24 genes, two at 2783, and several others at unique positions (note that

the numbering runs backward because it relates to the forward orientation of ROO).

Whereas intronic antisense ROO provides gene-transposon breakpoints for 28 exons, and

transposon-gene breakpoints for 33, intronic sense ROO only introduced 4 and 1 (Figure

5c). Similarly, the LTR BLOOD also introduced more breakpoints when it was inserted in the

antisense orientation relative to the host gene (14 vs. 6, Figure 5d,e).

We screened the transposon sections around breakpoints for consensus splice- acceptor

(SA) and donor (SD) sequence motifs and found many cases where the gene-to-transposon

chimera had been formed at SA consensus motifs, and transposon-to-gene chimera at SD

motifs (Stephens and Schneider, 1992) (Figure 4 – figure supplement 1, Supplementary

Table 6). For example, all breakpoints in antisense BLOOD that were formed with more than

one exon were precisely located at the predicted SA and SD splice site (Figure 5d).

Interestingly, we did not find a consensus SD motif at the transposon-gene breakpoint at

position 5191 of antisense ROO, although it frequently provided 5´- sequence to transposon-

gene chimeric RNAs. However, the sequence around position 5191 matched the consensus

motif, with the exception of a GT-to-GC conversion (see Figure 4 – figure supplement 1).

Taken together, our analysis revealed that transposons introduce many alternative splice

sites, which are recognised by the host cell spliceosome to combine cellular exonic

sequences with sections of transposon sequence.

We also identified cases of alternative splicing to different sites within the same transposon

insertion. Again using ROO as an example, αβCherry flies harbour a reverse orientation

ROO in the intron between exons 10 and 11 of the pan-neurally expressed *mustard* (*mtd*)

gene, which to date has only been implicated in fly innate immunity (Wang et al., 2012)

(Figure 5f). RNAseq revealed a complex collection of *mtd* splice variants that incorporated

different fragments of ROO (Figure 5g). SD sites upstream of this ROO came from the end

of either *mtd* exon 11. or 13. and these spliced in to the corresponding SA at position 5462

within ROO (Figure 5g). We also found three different SD sites (at positions 641, 2784 and

5191) within ROO, which spliced out to the closest downstream SA (exon 6) of *mtd*.

Therefore, this ROO element substantially increases the *mtd* mRNA isoform repertoire.

Without ROO the *mtd* locus can express 23 isoforms whereas with ROO it can generate 68

differentially spliced mRNAs.

We identified 147 other genes whose transcript diversity was similarly increased by

transposon insertions. These alternative transcripts incorporate 43 different transposon

families which each introduce cryptic SA and/or SD sites into host genes (see

Supplementary Table 5). For example, we found a sense insertion of BLOOD inside the

Dscam2 gene which encodes the transmembrane Down Syndrome cell adhesion molecule

2. Chimeric reads indicate that transcription of *Dscam2* is frequently initiated in BLOOD,

which is then spliced into exon 33 (the second exon) of the gene. This splicing event

combines ORF2 of BLOOD with the remaining exons of *Dscam2* and correctly aligns the

reading frames of the two transcripts, generating a novel N-terminus (Figure 5 – figure

supplement 1). We also observed cases where transposon chimera resulted in exon

skipping (e.g. the above mentioned ROO in *mtd* and 412 inside *tequila*, Figure 5 – figure

supplement 2). Most transposon chimera resulted from intronic insertions, but we also

detected one case where a HOBO insertion resided in the exon of the *CG31705* gene. This

HOBO introduced a cryptic SA site which was spliced to the upstream SD from the first

exon, creating a truncated *CG31705* transcript (Figure 5 – figure supplement 3). Together

these data show that a broad range of *Drosophila* transposons are alternatively spliced into

mRNAs producing many more isoforms of a large number of neurally expressed genes.

Alternative splicing into and out of transposons can be highly penetrant

Chimeric transcripts could be inconsequential to a cell if they only constitute a small percentage of the overall transcript repertoire of a given gene. We therefore quantified the percentage of mRNAs produced from a transposon harbouring gene that include transposon sequence. To do this we analysed sites where transposons were spliced into an exon-intron junction of a gene. For each gene we counted the number of reads spanning transposon-exon boundaries, and the number of reads that spanned the exon immediately up- and downstream of the transposon insertion (Figure 6a). This analysis showed that for some genes, all derived mRNAs contained transposon sequences. For example, all spliced copies of the isoform B of the Allatostatin A receptor 1 (*AstA-R1*) contained a section of ROO, and every transcript of *Piezo-like* (*Pzl*), a gene encoding a predicted mechanosensitive ion channel, ended in the I-element instead of exon 1 (Hu et al., 2019). All mRNAs retrieved from *Mec2* contained TRANSPAC as the most 5′ sequence, suggesting that transcripts might initiate within the TRANSPAC transposon and spliced to the SD of *Mec2* exon 2. On average, transposons contributed to 14% of transcripts derived from a specific gene (Figure 6b, Supplementary Table 7). Insertions in genes on the X chromosome, resulted in a higher average percentage of chimeric transcripts (27% vs. 13% for the rest of the genome) and a larger number of cases where 75% or more of transcripts were chimeric. Insertions on the X chromosome are hemizygous in male flies and our samples were generated by pooling an equal number of male and female flies. The increased chimera rate on the X might therefore reflect the smaller number of different X chromosomes when compared to the rest of the genome. For example, we found that the X-linked *cacophony* (*cac*) gene, which encodes a voltage-gated calcium channel, harboured a sense-orientation BLASTOPIA (Smith et al., 1996). This transposon insertion resulted in 54.7% of *cac* transcripts being truncated in αβCherry fly samples, potentially missing the last 8-11 coding exons, suggesting that many flies in this strain are likely mutant for the *cac* gene (Figure 5 – figure supplement 4). Another interesting example on the X chromosome of αβCherry flies is *Beadex* (*Bx*) which encodes a long-term memory relevant LIM-type transcription factor (Hirano et al., 2016). A sense NOMAD insertion gives rise to at least two new *Bx* transcript isoforms (Figure 5 – figure supplement 5), which constitute 9.8% of all *Bx* transcripts.

Splicing into transposons is common and varies between strains

The transposon complement is highly variable between fly strains. We therefore tested whether other fly strains express chimeric gene:transposon mRNAs by analysing three previously published mRNA sequencing data sets (Croset et al., 2018; Daines et al., 2011; MacKay et al., 2012). Although these prior studies generated shorter paired-end RNAseq reads, than those collected here, we were still able to find chimeric mRNAs in all three data

sets (Supplementary Table 8). Some transposon:gene chimera were conserved across all

strains, whilst others appeared to be strain-specific. 318 of the 887 chimera identified in our

$\alpha\beta$Cherry flies were present in at least one of the three other data sets, whereas 120 of

those occurred in at least three of the four strains. Chimera that were not detected in data

from other strains could either reflect genomic heterogeneity between the tested fly strains,

or the absence of evidence could result from lower sequencing coverage. Nevertheless,

these results demonstrate the prevalence of cellular mRNAs containing transposon

sequence.

Discussion

Somatic transposition in neurons has been proposed to contribute to age-dependent neuronal decline in wildtype and disease models of *Drosophila* and (Li, e al., 2013; Guo et al., 2018; Sun et al., 2018). Although the frequency of neural transposition is debated, expression is a prerequisite for movement. Therefore, transposons can only mobilize in cells that express full-length elements, or transposon mRNAs that encode enzymes permitting other elements to move in *trans*. It is therefore important to understand how transposon expression is controlled in the brain. Here, we combined single-cell expression data from the *Drosophila* midbrain with high-coverage gDNA sequence data of the same fly strain and find that the majority, if not all, expressed transposon sequences are parts of chimeric mRNAs with cellular genes.

Transposons residing in introns have previously been shown to contribute novel exons to several genes (Nekrutenko and Li, 2001). In addition long non-coding (lnc) RNAs frequently contain transposon sequences (Kapusta et al., 2013). We found transposon exonisation to be highly prevalent in the *Drosophila* brain. Transposons within the coding regions of genes and those in introns dramatically increase the transcript repertoire by introducing new splice variants. At this stage it is difficult to definitively determine the whole-genome functional consequences of splicing into transposons because we most often only retrieve the sequence across the splice junctions. Furthermore, although each transposon has a known consensus sequence individual copies are highly polymorphic. Nevertheless, our sequencing shows that transposon exonisation often truncates and/or changes the amino acid sequence of the encoded gene products, potentially changing protein structure and function. We did also identify several examples where the inclusion of transposon sequence conserved the reading frame of the host gene, likely generating a novel chimeric protein. Amongst the 148 transposon harbouring genes identified in this study, there are several that we have described in detail for which the observed locus disruption and altered expression would be expected to have significant consequences for neural function. Flies harbouring HOBO in *Sh* and BLASTOPIA in *cac* might exhibit altered voltage-gated currents, whereas those with ROO in *AstA-R1* will respond differently to the modulatory Allatostatin A neuropeptide. We also described insertions of 412 in *teq* and NOMAD in *Bx*, two genes which have been implicated in long-term memory formation.

Some transposons provide more splice sites than others. Our analysis of the abundant ROO retrotransposon (found in sense orientation inside 48 genes, and in antisense orientation inside 59 genes in our strain) provides a good example, and the insertion in *mtd* exemplifies the extended variance transposons can introduce to cellular transcription units. Interestingly,

antisense insertions of ROO provide more potential SA and SD sites than sense orientation

ROO. Insertions of ROO that are sense to the expression of the host gene were found to be

mainly part of pre-mRNA, from which functional reverse transcriptase could potentially be

translated. In contrast, antisense ROO introduces over 10 times more cryptic splice sites

(compare Figure 5b and c).

mRNA molecules that contain transposon sequences might be susceptible to short-

nucleotide mediated gene-silencing (Malone and Hannon, 2009), as recently reported for

LINE-2 containing mRNAs and LINE-2 derived microRNAs in humans (Petri et al., 2019).

This would provide a challenge for the cells expressing transposon-targeting piRNAs in

differentiating between chimeric transcripts and full-length transposon sequences. It is worth

noting that transposon-directed piRNAs have been identified in the fly brain. It is therefore

conceivable that transposon sequence permits cellular mRNAs to be selectively regulated.

In addition, transposon sequence might confer the capacity to be specifically trafficked within

the cell, and even between cells (Ashley et al., 2018; Pastuzyn et al., 2018).

The process of transposable elements acquiring new cellular functions that benefit the host

cell is called transposon exaptation (Gould and Vrba, 2013). Several examples of exaptation

events during the evolution of eukaryotes have been reported. How these functional

transitions occur is, however, not fully understood (Joly-Lopez and Bureau, 2018). Stress-

induced transposon mobilisation has been observed in many species, and it has been

hypothesized that these mobilisation events trigger the formation of new transposon-gene

chimera. Our results reveal a new mechanism by which transposons participate in the

generation of new gene variants. We show that transposons do not need to mobilise in order

to form chimeric mRNA molecules. Instead, the cellular gene spicing machinery frequently

uses intronic transposon insertions to increase transcriptional diversity.

Transposon sequences in the gDNA can deliver enhancer elements to cellular genes and

thereby influence their expression. However, our results do not support the idea that

transposons provide enhancer sequences contributing to neural expression of neighbouring

genes. If this was the case, we would expect to have found that cellular genes that harbour

the same type of transposon are more likely to be expressed in the same cells. Our data

instead show that transposon expression is dictated by the genes they are inserted within.

This is further supported by the fact that many genes that share transposon insertions with

neurally expressed genes were not expressed in the brain.

Our studies also introduce important practical concerns for the analysis of transposon

expression in somatic tissue and disease models. Several previous studies have used

Quantitative PCR (qPCR) to measure levels of transposon expression in mutant flies and

disease models. Since qPCR probes are usually not designed to cross transposon-gene

breakpoints, they cannot distinguish autonomous transposon expression from chimeric

mRNA. Similarly, standard RNA sequencing protocols rely on the alignment of cDNA

fragments to the reference genome and would not identify chimeric reads as non-

autonomous transposon expression. Baseline and changing cell-specific expression of host

genes that form chimeric transcripts with transposons can therefore be misinterpreted as

cell-restricted autonomous transposon expression and mobilization.

Our data also constrain somatic transposition. If all transposon expression depends on the

neighbouring host gene, only cells expressing that host gene can be susceptible to

transposition of that element. This would mean that GYPSY could only be active in glia, if

the fly strains being studied harbour a copy of GYPSY in a glial-expressed gene.

Interestingly, although we found sense GYPSY sequences in mRNAs for 14 different genes,

we did not detect glial GYPSY expression in our $\alpha\beta$Cherry flies.

Analysis of RNA-seq data from other fly strains suggests that more than half of the chimeric

transposon transcripts we identified in $\alpha\beta$Cherry flies are unique to this strain. This finding

alone demonstrates the incredible heterogeneity of the transposon complement between

strains. In addition, our prior genome sequencing revealed large differences between

individual $\alpha\beta$Cherry flies. Given the broad range of target genes that we have identified to

form chimeric mRNA with transposable elements, it will be important to understand the

functional consequences of fixed and variable transposons inside genes for the host

organism. Nevertheless, it seems highly likely that polymorphism of transposon load and the

distribution of transposons across the host genome could contribute towards heterogeneity

of neural function, and neurological pathology, between individual animals.

Methods

Fly strains

All experiments were performed on $\alpha\beta$Cherry flies, which were generated by crossing MB008b females (Aso et al., 2014) with w-; +; UAS-mCherry males. Flies were raised on standard molasses at 25°C, 40-50% humidity and 12 hr:12 hr dark-light cycles.

Long-read mRNA sequencing

For RNA extraction, groups of ~50 flies were frozen in liquid nitrogen and vortexed for 6 x 30 sec to separate heads from abdomens. Heads were isolated using a sieve. To avoid gDNA contamination, mRNA was purified with a combination of protocols. Samples were first processed with a column-based kit (RNeasy Mini kit, Qiagen, UK), including the optional on-column DNAseI digestion. Next, mRNA was extracted from total RNA using oligo-dT magnetic beads (NEB, Ipswich, MA) and the mRNA was purified again using the RNA columns. Finally, sequencing libraries were generated using oligo-dT magnetic beads from a strand-specific mRNA library preparation kit (TruSeq, Illumina, San Diego, CA), with 17 cycles of PCR amplification. Fragmentation was optimized to obtain ~350nt long fragments. Whole-genome sequencing was performed on a HiSeq 2500, with 250nt paired-end reads. We mapped the long reads using MRTemp, as previously described(Treiber and Waddell, 2017).

Single-cell read alignments

The Drosophila melanogaster reference genome release 6.25 was used for all sequence alignments. Transposon reference sequences were taken from Repbase (Jurka, 2000; Kaminker et al., 2002). Repetitive sequences in the *Drosophila* reference genome were masked using Repeatmasker (Smit et al.). Single-cell sequencing data was processed with a custom-built data processing pipeline, which is available on GitHub. The masked reference genome, as well as a gene reference file (refFlat) with all genes and each unique reference transposon sequence is provided as supplementary data.

Single-cell data analysis

Digital Gene Expression (DGE) matrices were generated as previously described (Butler et al., 2018). Rscripts can be downloaded as a supplementary file. In summary, DGE's were filtered (≥ 800 UMIs, ≥ 400 features) and 8 replicates were merged. Gene and transposon expressions levels were normalised separately. Marker genes were taken from Croset et al. (2018).

Co-expression analysis

Co-expression was quantified by calculating the Co-expression Disequilibrium (CD, see main text). The R-code snippet is available through Github. For the statistical analysis, a non-parametric test was performed. CD values between every gene- and transposon combination were ranked within each biological replicate, p-values were calculated using the student t-test and corrected for multiple comparisons using the Benjamini-Hochberg correction. In addition, CD values were calculated for every tested transposon with a set of 10 randomly assigned genes (or transposons).

Mapping germline transposon insertions in the fly

Germline transposon insertions were mapped with single-nucleotide resolution using previously published gDNA data from the $\alpha\beta$Cherry fly strain and MRTemp (Treiber and Waddell, 2017). A new, purpose-built, multi-functional sequence analysis pipeline called TEchim was developed. TEchim has 5 key functions: 1. generation of support files, including a masked reference genome and endogenous intron-exon junctions. (input files: reference genome, list of genes, list of transposon sequences). 2. alignment of un-stranded genomic DNA sequence data of multiple sequencing lanes and multiple biological replicates, detection of chimeric sequence fragments with single-nucleotide resolution, and the generation of summary output tables. 3. alignment of stranded cDNA data, detection of chimeric fragments, quantification of reads 4. generation of matching immobile genetic elements (IGE, see main text), analysis of these IGEs. These data are then used to determine sample-specific detection thresholds. 5. Quantification of LTR-gene and LTR-transposon reads (see Figure 4). Detailed descriptions and manuals are available on Github.

Splice acceptor (SA) and donor (SD) motif analysis

SA and SD motifs in the *Drosophila melanogaster* reference genome were generated by randomly selecting 500 known SA and SD sites from exons and screening for motifs in these sequences using the MEME suite (Bailey and Elkan, 1994) (Figure 4 – figure supplement 1). These motifs were then searched in sequence sections across transposon-gene breakpoints using FIMO (Grant et al., 2011).

References

Ashley, J., Cordy, B., Lucia, D., Fradkin, L.G., Budnik, V., and Thomson, T. (2018).
Retrovirus-like Gag Protein Arc1 Binds RNA and Traffics across Synaptic Boutons. Cell *172*,
262-274.e11.

Aso, Y., Hattori, D., Yu, Y., Johnston, R.M., Iyer, N.A., Ngo, T.T.B., Dionne, H., Abbott, L.F.,
Axel, R., Tanimoto, H., et al. (2014). The neuronal architecture of the mushroom body
provides a logic for associative learning. Elife *3*, e04577.

Bailey, T.L., and Elkan, C. (1994). Fitting a mixture model by expectation maximization to
discover motifs in biopolymers. Proceedings. Int. Conf. Intell. Syst. Mol. Biol. *2*, 28–36.

Baillie, J.K., Barnett, M.W., Upton, K.R., Gerhardt, D.J., Richmond, T.A., De Sapio, F.,
Brennan, P., Rizzu, P., Smith, S., Fell, M., et al. (2011). Somatic retrotransposition alters the
genetic landscape of the human brain. Nature *479*, 534–537.

Britten, R.J., and Kohne, D.E. (1968). Repeated sequences in DNA. Hundreds of thousands
of copies of DNA sequences have been incorporated into the genomes of higher organisms.
Science *161*, 529–540.

Butler, A., Hoffman, P., Smibert, P., Papalexi, E., and Satija, R. (2018). Integrating single-
cell transcriptomic data across different conditions, technologies, and species. Nat.
Biotechnol. *36*, 411–420.

Chung, N., Jonaid, G.M., Quinton, S., Ross, A., Sexton, C.E., Alberto, A., Clymer, C.,
Churchill, D., Navarro Leija, O., and Han, M. V. (2019). Transcriptome analyses of tumor-
adjacent somatic tissues reveal genes co-expressed with transposable elements. Mob. DNA
10, 1–22.

Coufal, N.G., Garcia-Perez, J.L., Peng, G.E., Yeo, G.W., Mu, Y., Lovci, M.T., Morell, M.,
O'Shea, K.S., Moran, J. V., and Gage, F.H. (2009). L1 retrotransposition in human neural
progenitor cells. Nature *460*, 1127–1131.

Croset, V., Treiber, C.D., and Waddell, S. (2018). Cellular diversity in the Drosophila
midbrain revealed by single-cell transcriptomics. Elife *7*.

Daines, B., Wang, H., Wang, L., Li, Y., Han, Y., Emmert, D., Gelbart, W., Wang, X., Li, W.,
Gibbs, R., et al. (2011). The Drosophila melanogaster transcriptome by paired-end RNA
sequencing. Genome Res. *21*, 315–324.

Evrony, G.D., Cai, X., Lee, E., Hills, L.B., Elhosary, P.C., Lehmann, H.S., Parker, J.J.,
Atabay, K.D., Gilmore, E.C., Poduri, A., et al. (2012). Single-neuron sequencing analysis of
l1 retrotransposition and somatic mutation in the human brain. Cell *151*, 483–496.

Evrony, G.D., Lee, E., Park, P.J., and Walsh, C.A. (2016). Resolving rates of mutation in the
brain using single-neuron genomics. Elife *5*, 1–32.

Faulkner, G.J., Kimura, Y., Daub, C.O., Wani, S., Plessy, C., Irvine, K.M., Schroder, K.,

Cloonan, N., Steptoe, A.L., Lassmann, T., et al. (2009). The regulated retrotransposon

transcriptome of mammalian cells. Nat. Genet. *41*, 563–571.

Gould, S.J., and Vrba, E.S. (2013). Exaptation-A Missing Term in the Science of Form

Exaptation-a missing term in the science of form. Paleobiology *8*, 4–15.

Grant, C.E., Bailey, T.L., and Noble, W.S. (2011). FIMO: Scanning for occurrences of a

given motif. Bioinformatics *27*, 1017–1018.

Guo, C., Jeong, H.H., Hsieh, Y.C., Klein, H.U., Bennett, D.A., De Jager, P.L., Liu, Z., and

Shulman, J.M. (2018). Tau Activates Transposable Elements in Alzheimer's Disease. Cell

Rep. *23*, 2874–2880.

Hirano, Y., Ihara, K., Masuda, T., Yamamoto, T., Iwata, I., Takahashi, A., Awata, H.,

Nakamura, N., Takakura, M., Suzuki, Y., et al. (2016). Shifting transcriptional machinery is

required for long-term memory maintenance and modification in Drosophila mushroom

bodies. Nat. Commun. *7*, 1–14.

Hu, Y., Wang, Z., Liu, T., and Zhang, W. (2019). Piezo-like Gene Regulates Locomotion in

Drosophila Larvae. Cell Rep. *26*, 1369-1377.e4.

International Human Genome Sequencing Consortium, Eric S. Lander, Lauren M. Linton,

Bruce Birren, Chad Nusbaum, Michael C. Zody, Jennifer Baldwin, Keri Devon, Ken Dewar,

and Michael Doyle (2001). Initial sequencing and analysis of the human genome. Nature

409, 860–921.

Izquierdo, M. (1994). Ubiquitin genes and ubiquitin protein location in polytene

chromosomes of Drosophila. Chromosoma *103*, 193–197.

Joly-Lopez, Z., and Bureau, T.E. (2018). Exaptation of transposable element coding

sequences. Curr. Opin. Genet. Dev. *49*, 34–42.

Jurka, J. (2000). Repbase Update: A database and an electronic journal of repetitive

elements. Trends Genet. *16*, 418–420.

Kaminker, J.S., Bergman, C.M., Kronmiller, B., Carlson, J., Svirskas, R., Patel, S., Frise, E.,

Wheeler, D.A., Lewis, S.E., Rubin, G.M., et al. (2002). The transposable elements of the

Drosophila melanogaster euchromatin: a genomics perspective. Genome Biol *3*.

Kaplan, W.D., and Trout, W.E. (1969). The behavior of four neurological mutants of

Drosophila. Genetics *61*, 399–409.

Kapusta, A., Kronenberg, Z., Lynch, V.J., Zhuo, X., Ramsay, L.A., Bourque, G., Yandell, M.,

and Feschotte, C. (2013). Transposable Elements Are Major Contributors to the Origin,

Diversification, and Regulation of Vertebrate Long Noncoding RNAs. PLoS Genet. *9*.

Kazazian, H.H. (2011). Mobile DNA transposition in somatic cells. BMC Biol. *9*, 2–5.

Ketchum, K., Ketchum, K., Hoskins, R., Hoskins, R., Wang, X., Wang, X., Smith, T., Smith,

T., Gocayne, J., Gocayne, J., et al. (2000). The genome sequence of Drosophila

melanogaster. Science (80-.). *287*, 2185–2195.

Krug, L., Chatterjee, N., Borges-Monroy, R., Hearn, S., Liao, W.W., Morrill, K., Prazak, L., Rozhkov, N., Theodorou, D., Hammell, M., et al. (2017). Retrotransposon activation contributes to neurodegeneration in a Drosophila TDP-43 model of ALS.

Lavie, L., Maldener, E., Brouha, B., Meese, E.U., and Mayer, J. (2004). The human L1 promoter: Variable transcription initiation sites and a major impact of upstream flanking sequence on promoter activity. Genome Res. *14*, 2253–2260.

Lewontin, R.C., and Kojima, K. (1960). The Evolutionary Dynamics of Complex Polymorphisms. Evolution (N. Y). *14*, 458–472.

Li, W., Prazak, L., Chatterjee N., Grüninger, S., Krug, L., Theodorou, D., Dubnau, J. (2013). Activation of transposable elements during aging and neuronal decline in Drosophila. Neurosci. Nat. *16*, 529–531.

Li, W., Jin, Y., Prazak, L., Hammell, M., and Dubnau, J. (2012). Transposable Elements in TDP-43-Mediated Neurodegenerative Disorders. PLoS One *7*, 1–10.

MacKay, T.F.C., Richards, S., Stone, E.A., Barbadilla, A., Ayroles, J.F., Zhu, D., Casillas, S., Han, Y., Magwire, M.M., Cridland, J.M., et al. (2012). The Drosophila melanogaster Genetic Reference Panel. Nature *482*, 173–178.

Macosko, E.Z., Basu, A., Satija, R., Nemesh, J., Shekhar, K., Goldman, M., Tirosh, I., Bialas, A.R., Kamitaki, N., Martersteck, E.M., et al. (2015). Highly parallel genome-wide expression profiling of individual cells using nanoliter droplets. Cell *161*, 1202–1214.

Malone, C.D., and Hannon, G.J. (2009). Small RNAs as Guardians of the Genome. Cell *136*, 656–668.

Muotri, A.R., Chu, V.T., Marchetto, M.C.N., Deng, W., Moran, J. V., and Gage, F.H. (2005). Somatic mosaicism in neuronal precursor cells mediated by L1 retrotransposition. Nature *435*, 903–910.

Nekrutenko, A., and Li, W.H. (2001). Transposable elements are found in a large number of human protein-coding genes. Trends Genet. *17*, 619–621.

Pastuzyn, E.D., Day, C.E., Kearns, R.B., Kyrke-Smith, M., Taibi, A. V., McCormick, J., Yoder, N., Belnap, D.M., Erlendsson, S., Morado, D.R., et al. (2018). The Neuronal Gene Arc Encodes a Repurposed Retrotransposon Gag Protein that Mediates Intercellular RNA Transfer. Cell *172*, 275-288.e18.

Perrat, P.N., Dasgupta, S., Wang, J., Theurkauf, W., Weng, Z., Rosbash, M., and Waddell, S. (2013). Heterogeneity in the Drosophila Brain. *340*, 91–96.

Petri, R., Brattås, P.L., Sharma, Y., Jönsson, M.E., Pircs, K., Bengzon, J., and Jakobsson, J. (2019). LINE-2 transposable elements are a source of functional human microRNAs and target sites. PLOS Genet. *15*, e1008036.

Philippe, C., Vargas-Landin, D.B., Doucet, A.J., Van Essen, D., Vera-Otarola, J., Kuciak, M., Corbin, A., Nigumann, P., and Cristofari, G. (2016). Activation of individual L1

retrotransposon instances is restricted to cell-type dependent permissive loci. Elife *5*, 1–30.

Rangwala, S.H., Zhang, L., and Kazazian, H.H. (2009). Many LINE1 elements contribute to the transcriptome of human somatic cells. Genome Biol. *10*, 1–18.

Roebroek, A.J.M., Pauli, I.G.L., Zhang, Y., and van de Ven, W.J.M. (1991). cDNA sequence of a Drosophila melanogaster gene, Dfur1, encoding a protein structurally related to the subtilisin-like proprotein processing enzyme furin. FEBS Lett. *289*, 133–137.

Smit, A.F.A., Hubley, R., and Green, P. RepeatMasker.

Smith, L.A., Wang, X.J., Peixoto, A.A., Neumann, E.K., Hall, L.M., and Hall, J.C. (1996). A Drosophila calcium channel α1 subunit gene maps to a genetic locus associated with behavioral and visual defects. J. Neurosci. *16*, 7868–7879.

Stephens, R.M., and Schneider, T.D. (1992). Features of spliceosome evolution and function inferred from an analysis of the information at human splice sites. J. Mol. Biol. *228*, 1124–1136.

Sun, W., Samimi, H., Gamez, M., Zare, H., and Frost, B. (2018). Pathogenic tau-induced piRNA depletion promotes neuronal death through transposable element dysregulation in neurodegenerative tauopathies. Nat. Neurosci. *21*, 1038–1048.

Treiber, C.D., and Waddell, S. (2017). Resolving the prevalence of somatic transposition in drosophila. Elife *6*.

Upton, K.R., Gerhardt, D.J., Jesuadian, J.S., Richardson, S.R., Sánchez-Luque, F.J., Bodea, G.O., Ewing, A.D., Salvador-Palomeque, C., Van Der Knaap, M.S., Brennan, P.M., et al. (2015). Ubiquitous L1 mosaicism in hippocampal neurons. Cell *161*, 228–239.

Wan, H.I., DiAntonio, A., Fetter, R.D., Bergstrom, K., Strauss, R., and Goodman, C.S. (2000). Highwire regulates synaptic growth in Drosophila. Neuron *26*, 313–329.

Wang, Z., Berkey, C.D., and Watnick, P.I. (2012). The Drosophila Protein Mustard Tailors the Innate Immune Response Activated by the Immune Deficiency Pathway. J. Immunol. *188*, 3993–4000.

Zhang, W., Wu, J., Ward, M.D., Yang, S., Chuang, Y.A., Xiao, M., Li, R., Leahy, D.J., and Worley, P.F. (2015). Structural basis of arc binding to synaptic proteins: Implications for cognitive disease. Neuron *86*, 490–500.

Author Contributions

C.D.T. and S.W. conceived the project and wrote the manuscript. C.D.T. performed and analysed all experiments.

Statement of competing interests

Both authors declare no financial and non-financial competing interests.

Figure Legends

Figure 1. Single-cell transcriptomics reveals patterned transposon expression in the *Drosophila* midbrain.

A Two-dimensional reduction (tSNE) of 14,804 *Drosophila* midbrain cells, based on gene expression levels. Colours represent cell clusters (at SNN resolution of 3.5). **B-F** Mean expression of transposons and neighbouring cellular genes in the relevant cell groups in 8 biological replicates and tSNE representation of cell-type restricted expression. **B** COPIA and *twin-of-eyeless* (*toy*) in all Kenyon Cell (KC) classes. **C** NOMAD and *mushroom-body expressed* (*mub*) in α´β´ KCs. **D** MICROPIA and *Rhodopsin 7* (*Rh7*) in the ellipsoid body **E and F** BLOOD and *CG32521*, and 412 and *tequila* (*teq*) in glia. Values represent the mean normalised number of unique molecular identifiers (UMI's) in an average cell from each cell type, and from the rest of the midbrain. Error bar indicates standard error of mean (SEM). Note that transposon- and gene levels were normalised separately. Blue schematic shows location of cell cluster (pink) in tSNE plot. **G** Two-dimensional reduction of 14,804 *Drosophila* midbrain cells, based exclusively on transposon expression levels. Colours represent cell clusters (at SNN resolution of 3.5). **H and I** Mean expression of STALKER2 and *Hsp70Bc* and DOC and *Furin 1* (*Fur1*) in their relevant transposon clusters and the position of the cluster in the overall transposon-based tSNE (indicated in pink).

Figure 1 – figure supplement 1. Sense strand transposon transcripts are twice as abundant as antisense strand transcripts in the *Drosophila* midbrain.

Graph showing mean expression levels across the entire midbrain of all sense and antisense transposon sequences. Each data point (cross) represents one biological replicate.

Figure 1 – figure supplement 2. Transposon expression patterns are stereotyped across biological replicates.

tSNE based on transposon expression levels showing all 8 biological replicates. Each replicate contributes cells to each cluster.

Figure 2. Transposons are co-expressed with neighbouring genes.

A Schematic and formulae describing the calculation of Co-expression Disequilibrium values. **B** Examples of transposon-gene pairs that are neighbouring in the genome and co-expressed across the midbrain. Pink bar represents the total transposon expression. Examples are selected from the most highly expressing transposons. See Supplementary Table 2 for entire list of correlated transposon-gene pairs. **C** Graph illustrating that more

neighbouring genes are correlated with their resident transposons than if transposons are
randomly assigned to genes.

Figure 3. Chimeric transposon-gene mRNA is abundant in the midbrain.
A Schematic of the structure of chimeric mRNA molecules, and illustrating the data
representation in **B**. **B** Examples of transposon-gene pairs that form chimeric mRNA. The
blue box contains the gene name, the section of the gene that forms a chimera and the
breakpoint, which is always the endogenous exon-intron junction. The red box contains the
transposon name, and the breakpoint(s) on the transposon. Note that for consistency,
transposon breakpoints are always taken from the sense orientation, starting from the core
transposon sequence, unless specifically indicated as [LTR]. Examples shown here are the
same transposon-gene pairs as in Figure 2b. For the entire list of chimera, see
Supplementary Table 5. **C** Graph showing the number of chimeric reads, and the combined
expression levels of each transposon-gene pair (pink), as well as for all 10 sets of IGE-gene
pairs (grey). Combined expression levels are the square root of the product of reads in both
transcripts of a transposon/IGE-gene pair. IGEs were used to calculate a threshold for a
False Discovery Rate that is less than 0.1%.

Figure 4. LTR retrotransposon expression is predominantly non-autonomously
A Illustration showing method of calculating the percentage of chimeric transcripts vs.
autonomously expressed transposon transcripts. Non-autonomous expression should result
in an approximate value of 0.5 for reads spanning the LTR section of transposons and the
intron of a neighbouring gene over the number of reads spanning the LTR and the core
section of the transposon. In contrast, autonomous expression would not result in LTR-gene
spanning reads. **B** List of all LTR transposons analysed in our mRNA data. We identified
LTR-gene spanning reads for every LTR transposon that is expressed in the midbrain. Error
bars represent standard deviation. Values have been capped at 0.5 in this graph. However,
some transposons produced a much higher number of LTR-gene reads (see Supplementary
Table 4).

Figure 4 – figure supplement 1. Splice acceptor and donor motifs.
A Splice acceptor motif, taken from 500 randomly chosen exon-intron junctions of
Drosophila genes. **B** splice donor motif.

Figure 5. Transposons introduce splice sites at conserved locations.
A Illustration of the labelling scheme in panels **B-E**. The pink bar represents the transposon;
light pink ends indicate the LTRs and the dark pink the core sequence. The positions of the

dots above the bar represent the site on the transposon where an upstream exon splice

donor site has merged. Every dot represents a different gene. Black lines in the top half of

the pink bar represent splice acceptor (SA) motifs in the transposon. Dots below the pink bar

indicate the location of breakpoints on the transposon that are spliced to upstream exonic

SA sites of different genes. Bars in the lower half indicate splice donor (SD) motifs. **B-E**

Representations of sense- and antisense ROO and BLOOD (to scale), with all breakpoints

to SA and SD sites of neighbouring genes. Note that the frequently used site on antisense

ROO at position 5191 is a non-consensus SD site, which lacks the expected GT motif at the

immediate breakpoint. The sequence around 5191 resembles the consensus SD motif,

although the GT is a GC. Compare TTTGGCAAGTT to motif in Figure 4 – figure supplement

1a. **F** Illustration of antisense ROO insertion in the *mustard* (*mtd*) gene. Only one isoform of

mtd is shown. Yellow box represents the 5´UTR, blue boxes are exons, orange box the

3´UTR, pink represents ROO transposon with white arrows indicating the LTRs. Breakpoint-

spanning gDNA reads reveal Target Site Duplication (TSD, inset). **G** Schematic of original

mtd transcript, and of three new splice isoforms.

Figure 5 – figure supplement 1. Sense BLOOD insertion in *Dscam2*.

Schematics of DScam2 mRNAs produced from locus containing BLOOD. Top shows the

nascent transcript spliced around the intronic full-length sense BLOOD insertion. Bottom

illustrates a new mRNA splice isoform, which reads through in frame from the ORF2

sequence of BLOOD into exon 2 of *DScam2*. The breakpoint in BLOOD is a consensus SD

motif. 23.2% of all *Dscam2* transcripts in fly heads are chimeric with BLOOD.

Figure 5 – figure supplement 2. Sense 412 insertion in *tequila*.

Schematics of *teq* mRNAs produced from locus containing full-length sense orientation 412

insertion. Top, original transcripts of *teq* splicing around 412. Bottom, and new *teq* splice

isoforms that include 412 sequence. 83.9% of *teq* transcripts contain 412 transposon

sequence. In addition, 22.4 % of 412 containing mRNAs skip exon 5 of *teq*. In 0.5% of cases

exons 2-5 are skipped.

Figure 5 – figure supplement 3. Antisense HOBO insertion in *CG31705*.

Schematic of *CG31705* mRNAs produced from locus containing exonic antisense HOBO

insertion. Transcripts containing unspliced HOBO and two additional new splice isoforms

that are generated by alternative splicing into HOBO are shown.

Figure 5 – figure supplement 4. Sense BLASTOPIA insertion in *cacophony*.

Schematic *cac* transcripts produced from locus containing a full-length intronic sense
BLASTOPIA insertion. A regular cac transcript is produced by splicing around the
BLASTOPIA insertion and a new truncated *cac* isoform results from splicing into
BLAsTOPIA. The *cac* gene is on the X chromosome, and 54.7% of *cac* transcripts in the fly
head are truncated by splicing into BLASTOPIA.

Figure 5 – figure supplement 5. Sense NOMAD insertion in *Bx*.
Schematic showing mRNAs produced from locus containing sense intronic NOMAD
insertion. Original transcript of *Bx* is generated by splicing around NOMAD. New splice
isoforms contain fragments of NOMAD. 9.8% of transcripts that start with the *Bx* 5´UTR are
spliced into NOMAD.

Figure 6. High penetrance of transposon-containing splice isoforms.
A Schematic showing method used to calculate the frequency of transposon chimera
produced from a given gene. The number of reads that span the exon-transposon junction
were divided by the total number of reads that map beyond the exon-intron junction (exon-
transposon reads plus reads that partially map to the exon immediately downstream of the
transposon insertion. **B** Examples of chimeric transposon-gene transcripts detected in fly
heads. Many genes are almost exclusively expressed as chimeric transcripts with
transposon sequence. See Supplementary Table 7 for complete list.

Figure 1

mixed: 2,9,13,15,18,19,28,69

Figure 1 - Figure supplement 1

Figure 1 - figure supplement 2

Figure 2

A

● Gene A (q_A)
● TE X (q_x)
● X and A (q_{AX})

(i) $CD_{AX} = q_{AX} - (q_A \times q_X)$

(ii) $r = \dfrac{CD_{AX}}{\sqrt{q_A \times (1 - q_A) \times q_X \times (1 - q_X)}}$

C

Number of genes with correlated expression pattern (y-axis, 0 to 60)

x-axis: neigbouring genes, random genes

B

Examples of transposon insertions in genes that exhibit correlated expression pattern

Total midbrain transposon expression

COPIA (9.6%) in *toy (& ...)*
ROO (6.6%) in *Glut1* (& ...)
TABOR (6.0%) in *CG17698* (& ...)
DOC (4.7%) in *Fur1* (& ...)
NOMAD (3.5%) in *mub* (& ...)
NEG-HOPPER (2.4%) in *sxc* (& ...)
HOBO (2.0%) in *Sh* (& ...)
MDG3 (1.7%) in *SelR* (& ...)
I-element (1.6%) in *Pde1c* (& ...)
NEG-ROO (1.1%) in *mtd* (& ...)
JOCKEY (1.1%) in *Desat1* (& ...)
NEG-DOC (1.1%) in *Ac3* (& ...)
STALKER2 (0.7%) in *Hsp70Bc*
GYPSY (0.2%) in *hiw* (& ...)
MICROPIA in *Rh7*
For full list, see Supplementary Table 2

Figure 3

A

B

C

Figure 4

A

B

Proportion of reads that span $n_\blacksquare$ / $n_\blacksquare$
LTR-intron vs. LTR-transposon

*no transposon expression detected

Figure 4 - figure supplement 1

A

B

Figure 5

Figure 5 - figure supplement 1

original *Dscam2* transcript:

new *Dscam2* variant:

5721

23.2% new variant

*in frame with BLOOD ORF2

original *teq* transcripts:

new *teq* variants:

83.9% contain transposon sequence

22.4% skip exon 5

0.5% skip exon 2-5

Figure 5 - figure supplement 3

original *CG31705* transcript:

new *CG31705* variants:

Figure 5 - figure supplement 4

Figure 5 - figure supplement 5

Figure 6

A

B

Gene	Transposon	$\dfrac{n_{TE}}{n_{TE}+n_{gene}}$	
AstA-R1	NEG-ROO	100.0%	
Pzl	I	100.0%	
nAChRalpha1	NEG-DOC6	100.0%	
Mec2	TRANSPAC	100.0%	
Thd1	MAX	100.0%	
CR45197	NEG-GYPSY	100.0%	
CG45263	NEG-POGO	100.0%	
Mocs1	P-ELEMENT	100.0%	
AstC-R1	NEG-FW	94.1%	
teq	412	83.9%	
Cyp4p2	COPIA	80.4%	

(for full list, see Supplementary Table 7)