Visualization of the HIV-1 Env Glycan Shield Across Scales

Zachary T. Berndsen[1,2,#], Srirupa Chakraborty[5,6,#], Xiaoning Wang[2,3] , Christopher A. Cottrell[1,2], Jonathan L. Torres[1], Jolene K. Diedrich[3], Cesar A. López[5], John R. Yates III[3] , Marit J. van Gils[7], James C. Paulson[2,3,4], S Gnanakaran[5] *, Andrew B. Ward[1,2] *.

[1]Department of Integrative Structural and Computational Biology, The [2]IAVI Neutralizing Antibody Center and Collaboration of AIDS Vaccine Discovery, Center for HIV/AIDS Vaccine Immunology and Immunogen Discovery (CHAVI-ID), [3]Department of Molecular Medicine, and [4]Department of Immunology and Microbiology, The Scripps Research Institute, La Jolla, CA 92037, USA. [5]Theoretical Biology and Biophysics Group and [6]Center for Nonlinear Studies, Los Alamos National Laboratory, Los Alamos, New Mexico 87545, United States. [7]Department of Medical Microbiology, Amsterdam UMC, location AMC, University of Amsterdam, 1105 AZ Amsterdam, Netherlands

[#]These authors contributed equally

*Correspondence: andrew@scripps.edu, gnana@lanl.gov

Abstract

The dense array of N-linked glycans on the HIV-1 Envelope Glycoprotein (Env), known as the "glycan shield", is a key determinant of immunogenicity, yet intrinsic heterogeneity confounds typical structure-function analysis. Here we present an integrated approach of single-particle electron cryomicroscopy (cryo-EM), computational modeling, and site-specific mass-spectrometry (MS) to probe glycan shield structure and behavior at multiple levels. We found that dynamics lead to an extensive network of inter-glycan interactions that drive the formation of higher-order structure within the glycan shield. This structure defines diffuse boundaries between buried and exposed protein surface and creates a mapping of potentially immunogenic sites on Env. Analysis of Env expressed in different cell lines revealed how cryo-EM can detect subtle changes in glycan occupancy, composition, and dynamics that impact glycan shield structure and epitope accessibility. Importantly, this identified unforeseen changes in the glycan shield of Env obtained from expression in the same CHO cell line used for GMP production. Finally, by capturing the enzymatic deglycosylation of Env in a time-resolved manner we found that highly connected glycan clusters are resistant to digestion and help stabilize the pre-fusion trimer, suggesting the glycan shield may function beyond immune evasion.

Significance Statement

The HIV-1 Env "glycan shield" masks the surface of the protein from immune recognition, yet intrinsic heterogeneity defies a typical structure-function description. Using a complementary approach of cryo-EM, computational modeling, and mass-spectrometry we show how heterogeneity and dynamics affect glycan shield structure across scales. Our combined approach facilitated the development of new cryo-EM data analysis methods and allowed for validation of

models against experiment. Comparison of Env across a range of glycosylation states revealed how subtle differences in composition impact glycan shield structure and affect the accessibility of epitopes on the surface. Finally, time-resolved cryo-EM experiments uncovered how highly connected glycan clusters help stabilize the pre-fusion trimer, suggesting the glycan shield may function beyond immune evasion.

Introduction

The Human Immunodeficiency Virus Type 1 (HIV-1) Envelope Glycoprotein (Env) is the sole antigen on the surface of the virion and has evolved several tactics for evading the adaptive immune system, chief among which is extensive surface glycosylation [1-3]. Env has one of the highest densities of N-linked glycosylation sites known, with glycans accounting for ~1/2 the mass of the molecule [4-8]. This sugar coat, or "glycan shield," is common among viral fusion proteins and is believed to be a primary hurdle in the development of neutralizing antibodies against Env during infection and vaccination [9, 10]. Therefore, arriving at a consistent and general description of its structure and dynamics may prove necessary in designing an effective Env-based immunogen and be of broader importance towards understanding the structure and function of other densely glycosylated proteins.

N-linked glycans can be highly dynamic, however, and can vary substantially in length, connectivity, and chemical composition [11, 12]. On the Env ectodomain surface there are ~95 glycosylation sites on average [6], and mass-spectrometry analysis (MS) has revealed that compositional heterogeneity exists between sites as well as at the same site between different copies of Env [13]. While heterogeneity and dynamics are critical for immune evasion, they are incompatible with a typical structure-function description. When the first structures of the fully glycosylated Env were solved with single-particle electron cryomicroscopy (cryo-EM) [14, 15] most glycans were not resolved beyond the first or second sugar ring except when stabilized by antibodies. This suggested that the glycan shield is highly dynamic, however, due to the missing high-resolution details, structural analysis remained focused on Env-antibody interactions, while experimental investigations into the glycan shield shifted primarily towards chemical analysis via MS [11]. One crystallography study specifically addressing glycan shield structure was recently published and the data shows evidence for highly stabilized glycans engaging in a wide range of inter-glycan contacts [6], which is at odds with the cryo-EM results. Such a disagreement between the two techniques highlights the difficulty in studying the glycan shield and the need for more focused experiments.

Computational modeling, in particular molecular dynamics (MD) simulations, has proven useful as a complementary approach for characterizing glycoprotein structure and dynamics [16-19], including Env [6, 20-22]. Three sets of atomistic dynamics simulations of fully glycosylated Env trimers were recently reported and these predicted that dynamics would lead to extensive shielding of antigenic surface and the formation of complicated networks among interacting glycans [6, 21, 22]. In one study, glycans were observed clustering into distinct micro-domains over long timescales, and it was found that neutralizing antibodies preferentially target the interfaces

between these domains, suggesting the glycan shield may possess biologically relevant large-scale structure. Although

MD is a widely accepted method for modeling glycoproteins, atomistic simulations for systems as large as Env require

substantial computational resources to access biologically relevant spatial and temporal scales and can suffer from

initial model bias. Therefore, more computationally efficient methods for modeling fully glycosylated Env are needed.

Furthermore, there are still no established methods for experimentally validating the results of these simulations.

Here, we describe an integrated approach combining cryo-EM, computational modeling, and site-specific MS

aimed at illuminating glycan shield structure and behavior at multiple levels. First, we introduce a model system for

our cryo-EM experiments and describe the complete structure of the native Env glycan shield for the first time using

scale-space and 3-D variability analysis. In parallel, we developed a high-throughput modeling pipeline for rapidly

generating diverse ensembles of fully glycosylated Env at atomistic resolution, enabling quantification of glycan-

specific geometric properties and concerted behavior within the glycan shield as a whole. We then used these

ensembles as ground-truth for the creation of simulated cryo-EM data, which facilitated the development of new

analysis tools and enabled the validation of theoretical models against cryo-EM experiments.

With our integrated approach, we show that the glycan shield is highly dynamic but exhibits variation

in dynamics between glycans due in part to crowding and other geometric and energetic constraints. We found that

dynamics give rise to a network of inter-glycan interactions that drive the formation of higher-order structure within

the glycan shield. This blurry low-resolution structure creates diffuse boundaries between buried and exposed protein

surface that define potential sites of vulnerability. Using Env expressed in three common cell lines, we show how

differences in glycan composition and occupancy can be detected by cryo-EM and result in changes to glycan shield

structure and dynamics that affect the accessibility of epitopes on the surface. To better inform our results we present

site-specific mass-spectrometry data for the same samples and demonstrate that cryo-EM can complement such

studies. Finally, by exposing Env to endoglycosidase digestion and capturing reaction intermediates with cryo-EM in a

time-resolved manner, we found that highly connected glycan clusters are resistant to digestion and act to stabilize

the pre-fusion trimer structure, implying the glycan shield may function beyond immune evasion.

Results

Env in complex with a base-specific antibody as a model system for cryo-EM analysis of the native glycan shield

We choose BG505 SOSIP.664 as a model system to demonstrate our approach because it is the first and most widely

studied of the soluble pre-fusion stabilized Envs (23, 24). Native Env is a transmembrane protein that forms a trimer

of dimers composed of two gene products, gp120 and gp41, which are responsible for receptor binding and

membrane fusion respectively [add citation]. The acronym *SOSIP* is in reference to the engineered disulfide bonds

between gp120 and gp41 and the isoleucine to proline mutation in the HR1 helix, which act to stabilize the pre-fusion

conformation, while 664 is the final residue in truncated ectodomain (23). BG505 SOSIP.664 is also currently in the

first ever human clinical trials of a trimeric subunit-based HIV vaccine

(https://clinicaltrials.gov/ct2/show/NCT03699241). In Fig. 1 we present the cryo-EM reconstruction and refined

atomic model of BG505 SOSIP.664 purified from HEK293F cells in complex with three copies of the fragment antigen

binding domain (Fab) of the non-neutralizing antibody, RM20A3. This complex will be referred to as BG505_293F from

here on. The RM20A3 Fab was included because it improves orientation bias, as illustrated by the complete and near-

uniform angular distribution (Fig. 1*C*), and it binds to a non-native epitope at the base of the SOSIP trimer that is

devoid of glycans (Fig. 1*D*). This leaves the native glycan shield undisturbed and means that RM2OA3 Fab binding is

independent of glycosylation, making the SOSIP/base-Fab complex ideal for this study.

The global resolution of this C3-symmetric reconstruction determined by Fourier shell correlation (FSC) is

~3.1Å (*SI Appendix*, Fig. S1*A*), and the bulk of the protein is at or near this resolution (Fig. 1*E*). ~80% of Env residues

could be confidently built into this map, while most glycans are ill-defined beyond the core N-acetylglucosamine

(NAG), accounting for only ~15% of the total predicted glycan mass (Fig. 1). Glycans located on the poorly resolved

flexible loops (N185e, N185h, N398, N406, N411, N462) could not be identified at all, however, site-specific MS data

confirmed they are indeed glycosylated (25) (*SI Appendix*, Fig. S2*A*). N-linked glycans on Env can be roughly classified

into two types, high-mannose and complex, based on the extent of intracellular processing, and they differ in

composition and structure (one possible configuration of each type is depicted in Fig. S 1F). Even if the glycans were

better resolved, averaging of individual projections during 3-D reconstruction means we cannot uniquely identify the

specific type of glycan at most sites because they possess a mixture of several different types (26) (*SI Appendix*, Fig.

S2*A*). Of those we could identify, a few were found to be near-uniformly processed (designated as >75% complex), for

example the glycans at N611 and N618 on gp41, and it should be possible to confirm this by the presence of a core

fucose residue. However, we were unable to identify clear density for fucose at any of these sites (Fig. 1*F*). Together,

these observations show that the glycan shield is highly dynamic with respect to the protein core but exhibits some

differences in ordering between glycans, and that it is not possible to distinguish between the different types of

glycans from a single cryo-EM map when the glycans are not stabilized by antibodies.

Scale-space and 3-D variance analysis reveal interconnectivity and higher-order structure within the glycan shield

Although we lack the resolution to build precise atomic models, there is still a great deal of information about glycan

shield structure and dynamics that can be gleaned from the cryo-EM data. At low isosurface thresholds noise appears

surrounding Env's well resolved protein core where the missing glycan mass should be (Fig. 1*E*). To look for meaningful

structure within this noisy signal we analyzed properties of the map across a range of scales by progressively

smoothing it with a Gaussian filter of increasing standard deviation (SD). Resolution in the glycan shield falls off with

distance from the protein surface, therefore as the map is smoothed more and more glycan signal should become

visible until the map encompasses the majority of the space occupied by the ensemble of configurations. This process

could be captured by measuring the volume of the map across the scale-space. To calculate the volume, however,

one must first choose an appropriate threshold. For our analysis we implemented an auto-thresholding method based

a simple measure of topological connectivity described in the *SI Appendix* (*SI Appendix*, Fig. S3*A-F*). We refer to the

resulting intensity as the *noise threshold* since it represents the lowest threshold before the appearance of noise. Fig.

2*A* shows the noise threshold and volume of the map at the noise threshold as a function of Gaussian filter width.

Both curves have a sigmoidal shape; once the map is sufficiently smoothed it undergoes a period of rapid expansion

in volume and reduction in noise threshold that is presumably capturing the appearance of signal from the poorly

resolved regions of the map. The curves then plateau around ~1.5-2 SD, suggesting there is negligible gain in signal

with further smoothing.

Examination of these filtered maps confirms the progressive emergence of more structure in the glycan

shield (Fig. 2*B*). By taking 2-D slices through six representative maps and displaying them on top of one another we

see how glycosylated surfaces undergo substantially more dilation (red arrows) than the non-glycosylated surfaces.

Once the isosurface has expanded to include signal from glycan antennae, neighboring glycans begin to merge,

creating a canopy-like effect that occludes underlying protein surfaces (black arrows). This implies that neighboring

glycans sample overlapping volumes and have the potential to interact. Also apparent is the negligible gain in signal

moving from 1.5 to 2 SD (blue and pink). We therefore concluded that the 1.5 SD Gaussian filtered map is the ideal

scale for interpreting glycan shield structure in this reconstruction. Visualization of this filtered map across a range of

thresholds reveals a hierarchy of structural features (Fig. 2*C*). At high thresholds, the glycan shield is composed

primarily of isolated and well-defined glycans stems, while the majority of the protein surface is exposed (gray map).

Moving to lower thresholds, it progresses through a series of higher-order structural states defined by isolated clusters

of multiple overlapping glycans, until finally becoming completely interconnected near the noise threshold. This low

threshold structure occludes large regions of protein surface and represents the full extent of the glycan shield.

As a complementary approach for visualizing the dynamic glycan shield, we performed 3-D variability analysis

in the SPARX software package (27, 28) (Fig. 2*D*). 3-D variability, which is closely related to 3-D variance, should be

high anywhere there is significant heterogeneity in the map. Indeed, we see high variability around known flexible

regions such as the constant domains of the Fabs (clipped from view) as well as variable loops and the exterior of the

protein surface at the sites of N-linked glycans. Variability in the glycans is highest at the distal ends of the glycan

stems and expands outward as the threshold is reduced, revealing extensive interconnectivity (Fig. 2*D*). 2-D slices

through the variability map highlight the canopy effect between neighboring glycans, which acts to occlude underlying

protein surfaces (red arrows - Fig. 2*D*). Overall the shape and topology are similar to the Gaussian filtered map. A

more complete description of the 3-D variability can be found in the *SI Appendix*.

A high-throughput atomistic modeling (HT-AM) pipeline for generating large ensembles of glycosylated Env

Cryo-EM does not capture atomic details of individual molecules, so to better understand the results presented above,

we performed atomistic simulations of fully glycosylated BG505 SOSIP.664. To overcome the sampling limitations of

MD simulations, we employed a high throughput atomistic modeling (HT-AM) pipeline based around the ALLOSMOD

package of the MODELLER software suite (29), which has been used previously to generate small ensembles of

glycosylated Env (30). This methodology accounts for both energetic and spatial constraints on glycan sampling by a

combination of empirical energy minimization-based structural relaxation and simulated annealing (described in detail

by Guttman et al. (29), along with initial randomization of glycan orientations. However, it does not provide temporal

information about dynamics. Here, we have built on this by incorporating 10 different protein scaffolds and additional

variability in the V2 and V4 loops (Fig. 3*A*) and utilized it to generate much larger ensembles of 1000 fully glycosylated

Env structures with uniform mannose-9 (Man9) glycans. Ten such models are shown in Fig. 3*C*, one from each protein

scaffold, while the full set of models at a single glycosylation site is shown in Fig. 3*D*. We also repeated the simulation

with uniform mannose-5 (Man5) glycosylation for comparison.

The HT-AM pipeline capture spatial and energetic constraints on glycan flexibility

To assess how spatial and energetic constraints affect the flexibility of individual glycans in our simulations we

calculated the root mean-squared fluctuation (rmsf) for each glycan across all 1000 models after aligning the protein

scaffold. Our method captures variability between glycans (Fig. 3*E*), and the additional flexibility imparted by the 10

starting protein scaffolds can be appreciated by comparison to the glycan only rmsf (Fig. 3*F*). For example, glycans

located on the flexible loops have a much higher rmsf than all other glycans (185e, 185h, 406 and 411) whereas

glycans at the N262, N301, N332, N448, and N611 sites have lower rmsf. This leads to a large difference in sampled

volumes between the most and least dynamic glycans (*SI Appendix*, Fig. S4*A*). We also see an increase in average rmsf

of the individual glycan residues starting from the core NAG and moving outward to the tips of each antenna (*SI*

Appendix, Fig. S5*A-C*), which is in line with the cryo-EM results showing reduced resolution beyond the core NAG (Fig.

1*D*).

We observed a similar trend for the Man5 ensemble (*SI Appendix*, Fig. S5*G-H*), however, by comparing the

average rmsf values of the 5th mannose residue alone, we found a slight increase in rmsf and sampled volume at most

sites compared to the Man9 ensemble (Fig. 3*G* and *SI Appendix*, Fig. S5*I-J*). We attribute this effect to increased

crowding in the glycan canopy from the more massive Man9 glycans. In support of this, there is a significant positive

correlation between glycan flexibility glycan crowding (see *SI Appendix*), however, only when considering relatively

large neighborhoods (*SI Appendix*, Fig. S6*A*). A similar trend was observed by Stewart-Jones et al. which they attributed

to different "shells" of influencing glycans (6). Such higher-order dynamic effects could be possible in light of our

structural observations.

In addition to crowding effects, the local protein structure will also influence glycan dynamics. In our

modeling pipeline, the protein backbone was kept harmonically restrained close to the template to allow for extensive

sampling of glycan conformations using simulated annealing. Thus, we see that the Asn sidechains of residues 88, 160,

197, 234 and 262 all have very low rmsf (*SI Appendix*, Fig. S5*F*), possibly stemming from limited torsional space

available during modeling. The glycosylated Asn residues in gp41 have relatively low rmsf as well (N611, N618, N625,

and N637), being situated on stable helical bundles (*SI Appendix*, Fig. S4*B*). This ultimately results in a relative reduction

of the glycan dynamics at some of these sites (Fig. 3*C*). Correcting for the contribution to fluctuations coming from

the underlying protein, we observed that the rmsf between the different glycans are comparable, ranging from 3 Å to

5 Å, with similar scale of SD (Fig. 3*F*).

210 Simulated cryo-EM maps reproduce defining features of the experimental data

Given the single-molecule nature of cryo-EM datasets, the ensembles generated by the HT-AM pipeline can be seen

as representing the individual particles that go into a 3-D reconstruction. With that in mind, we established a protocol

for transforming the ensembles into simulated cryo-EM datasets (Fig. 4*A*). For comparison, we replicated the process

for the Man5 ensemble as well as a protein-only ensemble (BG505_Man9, BG505_Man5 and BG505_PO respectively).

The simulated maps reproduced some of the defining features of the experimental data. For instance, refinement was

dominated by the stable protein core and only the first few sugar residues at each site are defined when the map is

filtered to the global FSC resolution (Fig. 4*A*). This leads to a similar scale-space structure (Fig. 4*B*), with a plateau again

appearing around 1.5-2 SD when the maps are filtered to the same initial resolution, suggesting that the HT-AM

pipeline is capturing physiologically relevant sampling in the glycan shield, at least globally. Importantly, we found that

the volume of the 1.5 SD Gaussian filtered map at the noise threshold closely approximates the total volume sampled

by the ensemble (Fig. 4*B* - dashed lines). Comparing the curves for all three simulated reconstructions (Fig. 4*B*) we see

how differences in glycosylation manifest as differences in volume at low resolutions, suggesting cryo-EM can be used

to capture global changes in glycan shield composition between two reconstructions of the same Env. Visualization of

the 1.5 SD Gaussian filtered BG505_Man9 map at high and low thresholds reveals a similar evolution towards a more

connected topology as the threshold is reduced and extensive shielding of protein surface (Fig. 4*C*). We also calculated

the 3-D variability map for the BG505_Man9 reconstruction and observed a similar result to the experimental data

(Fig. 4*D*). In addition, we compared the SPARX variability map to the true 3-D variance (*SI Appendix*, Fig. S7*B*) and

found negligible differences between the two (See the *SI Appendix* for details).

230 Measuring glycan dynamics in cryo-EM maps

Measuring intensity around individual glycans in the cryo-EM maps should allow us to assess their relative dynamics.

To confirm this, we used the simulated BG505_Man9 map since we could make direct comparison to the rmsf values.

However, only the first 1-2 glycan residues are defined at most sites in the high-resolution map (Figure5A) and their

dynamics deviate significantly from the rest of the glycan (*SI Appendix*, Fig. S8*A*). The third glycan residue, β-mannose

(BMA), more closely approximates the average rmsf (mean deviation ~2Å) and the relative differences between sites

(*SI Appendix*, Fig. S8*A-B*). In addition, the glycan stems are well defined in the 1.5 SD Gaussian filtered map, which

allowed us to identify their locations in the map with reasonable accuracy. So, we built and relaxed glycan stems into

the 1.5 SD Gaussian filtered BG505_Man9 simulated map (Fig. 5*A*) at every glycosylation site we could confidently

identify (23/28 per protomer). We could not identify the V2 and V4 loop glycans at N185e, N185h, N406, and N411,

which had the highest rmsf values in the simulation, nor the glycan at N339, which has a lower rmsf but projects

directly towards the heterogeneous V4 loop. The positions of the BMA residues were then used to analyze local map

statistics within a spherical probe around each glycan (Fig. 5*A*). We observe a strong positive correlation between the
inverse rmsf and normalized mean intensity (Fig. 5*B-C*), with the peak correlation coefficient of ~0.88 (p=2.1e-8)
occurring when using the unfiltered map and a probe radius of 2.3Å. Thus, local intensity around BMA residues
accurately captures ground-truth differences in relative dynamics between glycans and can be used to make direct
comparisons between the simulated and experimental cryo-EM maps.

The HT-AM pipeline reproduces physiologically relevant trends in glycan dynamics measured by cryo-EM
With a method in place for measuring glycan dynamics from cryo-EM maps, we then made direct comparisons with
the experimental data. We built and relaxed glycan stems into the 1.5 SD Gaussian filtered BG505_293F map as
described above and could identify clear density at 21/28 glycosylation sites per monomer, two less than from the
simulated map (N398 and N426). The other five missing glycans were common to both maps, meaning the HT-AM
pipeline captured physiologically relevant dynamics at these sites, at least up to the detection limits of this method.
Overall, we found that the HT-AM pipeline captured a similar trend in glycan dynamics with a correlation coefficient
between the two of ~0.46 (*p* = 0.03) (Fig. 5*D*).

In addition to the V4 and V5 loop glycans at N398 and N462, we also saw a large deviation at the N137 glycan
on the V1 loop, as well as the N262 and N301 glycans. In the BG505_293F map, the N262 glycan was the most ordered
due to stabilizing contacts with the gp120 core, and these interactions may not have been accurately captured by the
simulation given the restricted protein dynamics. In gp41, a large deviation also occurred at the N611 glycan, which
we attribute to restrictive sampling of the protein backbone as previously discussed. To complicate the comparison,
the N618 and N625 sites are under-occupied as revealed by MS (*SI Appendix*, Fig. S2*A*). Although sub-occupancy will
cause reduced signal intensity due to averaging, the severity of this effect is diminished when dynamics at that site
are large relative to other glycans, which is the case for the glycan at N625.

Detecting site-specific changes in glycan dynamics, occupancy and chemical composition from cryo-EM maps.
To test whether these methods could be used to detect site-specific changes in dynamics and occupancy between
cryo-EM maps of differentially glycosylated Env, we performed a comparative analysis between the BG505_Man9 and
BG505_Man5 simulated maps. Given that stems of a Man9 and Man5 glycan are identical, the only changes in
intensity around the BMA residue should arise from differences in dynamics alone. On average, we see a ~17%
reduction in intensity indicative of increased dynamics, which is in line the rmsf data (Fig. 6*A*). We also accurately
detect the largest increase and decrease in dynamics at the N262 and N234 sites respectively. To verify we could
detect changes in occupancy, we removed the glycan at the N625 site from half of the models and re-refined the data
(referred to as BG505_Man9HO for "half occupancy"). Not surprisingly, we see a ~50% reduction in mean intensity
from the fully occupied reconstruction around this site (Fig. 6*A*).

Another technique that should be sensitive to subtle changes between two cryo-EM maps is difference
mapping. Indeed, the change in occupancy at the N625 site is apparent in the BG505_Man9 - BG505_Man9HO

difference map (Fig. 6*B*). At high threshold, the signal is localized around the glycan stem and extends to the protein
surface. In the BG505_Man9 - BG505_Man5 difference map (Fig. 6*C*), however, the difference signal is strongest
where the distal tips of the Man9 glycans would be and does not extend to the protein surface. These results establish
cryo-EM as a tool for measuring glycan dynamics as well as changes in chemical composition and occupancy between
differentially glycosylated Envs.
282
Insights gained from analysis of simulated data allows improved characterization of cell-type specific differences
in glycan shield
To test our methods experimentally, we collected cryo-EM data on complexes of RM20A3 Fab and BG505 SOSIP.664
expressed in two additional common cell lines that produce major and minor changes in glycosylation; HEK293S cells,
and a stable CHO cell line (referred to as BG505_293S and BG505_CHO respectively). The stable CHO cell line
expressed Env sample was provided to us by IAVI as part of the demo run conducted prior to the currently on-going
human clinical trials, and should be identical to the final vaccine product (31). Both new datasets refined to ~3Å-
resolution (*SI Appendix*, Fig. S1*A*) and the three atomic structures are nearly identical (Cα RMSDs ~ 1Å - *SI Appendix*,
Fig. S1*B*). Scale-space (*SI Appendix*, Fig. S3*G-I*) and 3-D variability analysis (*SI Appendix*, Fig. S9*A-C*) also revealed similar
low-resolution structural features to BG505_293F, however, comparison of local map intensity around BMA residues
uncovered some significant difference in the gp41 glycans, particularly at the N611 and N625 sites, potentially from
changes in occupancy (Fig. 7*A*). This is further supported by site-specific MS data collected for the two additional
samples (*SI Appendix*, Fig. S2*A*).
The 293S sample contains only high-mannose type glycans and should therefore allow detection of complex
glycans via difference mapping, and to a lesser extent, different distributions of high-mannose glycans. Indeed, we
see strong difference signal around the primarily complex gp41 glycans in both the difference maps (Fig. 7*B-C*). We
also observe signal around the primarily complex glycan at N88 (Fig. 7*F*). In addition, clear difference signal appeared
around the V2, V4, and V5 loops, specifically near the glycans at N185e and h, N398, and N462, all of which are shown
to be complex by MS. The MS data also shows a subtle difference in both occupancy and percentage of complex
glycans at the N398 site between the 293F and CHO sample, which could explain why there is strong difference signal
in one map and not the other. According to the MS data, the only significant differences in occupancy between the
293F and 293S samples occurred at the N137, N133, and N611 sites, however the difference maps do not contain any
signal around the N611 glycan, suggesting there may be discrepancies between the two methods.
The MS analysis also detected differences in occupancy between the CHO sample compared to the 293F
sample at multiple sites (*SI Appendix*, Fig. S2*A*). Indeed, upon closer examination we found that the difference signal
around the gp41 glycans at N611, N618, and N625 extends all the way to the protein surface (Fig. 7*G*), indicative of
changes in occupancy. In this difference map, there is also clear signal around the N137 site, and to a lesser extent at
the tip of the N332 glycan stem (Fig. 7*H*). Given the proximity of N137 and N332, it is plausible that sub-occupancy at
one is driving changes in dynamics and/or glycan distribution at the other. Similar higher-order effects have also been

observed via MS when comparing glycoform distributions on Env before and after knocking out specific glycans (26,

32, 33).

A probabilistic glycan-glycan interaction network reveals highly connected glycan clusters

Both the experimental and simulated cryo-EM maps showed extensive interconnectivity among glycans, so we sought

to quantify this more precisely. With each glycan in our models sampling a particular region of space, neighboring

glycans can explore overlapping volumes, and the fraction of this overlap gives a measure of their interaction

probability. Fig. 8*A* shows a heat map of the normalized glycan-glycan volume overlap matrix. By interpreting this as

an adjacency matrix we can model the glycan shield as a network. Here, each glycan was defined as a node, and two

nodes were connected by an edge if there was at least 5% overlap between them (Fig. 8*B*). The only glycans from the

neighboring protomers that were considered are those with an inter-protomer edge. Fig. 8*C* shows the network

obtained for the Man9 ensemble unfolded in 2-D. It can be seen that overall there are three main regions of overlap,

– the V1/V2 apex, the gp41 base, and the densely occupied gp120 outer and inner domains that includes the high-

mannose patch (HMP). The inter-protomer overlaps are contributed mainly by the V1/V2 glycans. The nomenclature

was inspired by the established nomenclature for gp120 domains (34), however, our definitions were adapted to

better capture glycan shield structure (*SI Appendix*, Fig. S10*A*). The analysis was repeated for the Man5 ensemble and

we observed a reduced overall connectivity as measured by the mean node degree (from ~7 to ~5) and the maximum

network diameter (from 5 to 8 hops), consistent with their smaller size (*SI Appendix*, Fig. S*11*).

With a network in place we could then analyze the relative influence of each glycan on the whole system and

examine its long-range structure. To do this we calculated the relative eigenvector centrality of the nodes, which is a

measure of importance in the network, and the results are projected on the network as a colormap in Fig. 8*C*. We see

that the gp120 outer domain glycans at N332, N339, N363, N386, and N392, which are densely connected within the

network, have strong eigencentrality measures. In the apex region, the V1/V2 loop glycans at N133, N160 and

N185e/h also have relatively high eigencentrality due to their increased flexibility, while the glycans at N88, N234,

N276, and those in gp41 have low eigencentrality, reflecting low interaction probabilities. Incorporated intrinsically

into the network is a set of stable sub-graphs that represent highly connected glycan clusters. To illustrate this

structural hierarchy, we progressively stripped the network using tighter overlap cutoffs (*SI Appendix*, Fig. S10*B*). As

the network is degraded, we see the formation of two large sub-graphs; one composed of the V1/V2 apex and the

gp120 outer domains; and the second composed of the gp120 inner domain along with gp41 base glycans. With an

even stricter cutoff, the sparsely connected glycans with low eigencentrality separate out and the two sub-graphs split

again to form four sub-graphs; a V1/V2 apex domain; a gp120 outer domain; a gp120 inner domain; and lastly, the

group of unconnected gp41 base glycans.

Highly connected glycan clusters are resistant to enzymatic digestion and help stabilize of the pre-fusion trimer

To determine if the glycan shield affects Env structure and dynamics we exposed BG505_293S (already in complex

with RM20A3 Fab) to digestion by endoglycosidase H (Endo H), which cleaves high-mannose glycans between the first

and second sugar rings (Fig. 9*A*), and collected cryo-EM data on the deglycosylated sample. In addition, we

hypothesized that highly connected glycans would be protected from enzymatic digestion, and vice versa. Therefore,

if we could measure the order in which glycans were cleaved by Endo H it could provide indirect validation of our

network models. To test this, we collected cryo-EM data on a partially deglycosylated sample as well. The reaction

times and Endo H concentrations for achieving partial (2hrs) and complete (16hrs) digestion were determined prior

to performing cryo-EM. The datasets, referred to as BG505_EndoH2 and BG505_EndoH16, reconstructed to ~3.2Å

and ~3.5Å-resolution respectively, with similar overall quality, resulting in nearly identical atomic models to the other

three datasets (*SI Appendix*, Fig. S1*A-B*). The progressive digestion of the glycan shield was confirmed by SDS-PAGE

and SEC chromatography (*SI Appendix*, Fig. S12), while scale-space analysis of the cryo-EM maps revealed a clear

progressive reduction of glycan signal (*SI Appendix*, Fig. S3*G-I*).

358   Indeed, analysis of local map intensity around each glycan revealed that digestion occurred non-uniformly

(Fig. 9*B-C*). This was also confirmed by changes in the 3-D variability maps (*SI Appendix*, Fig. S9*D-E*). By assuming a

linear relationship between intensity and occupancy and using the MS data to set the initial occupancy, we calculated

the occupancy at each site (Fig. 9*C*). After 2 hours the gp41 glycans (N611-637) were completely digested, while some

glycans, particularly those in the densely packed gp120 outer domain, remained mostly intact. Although we could not

identify unique signal for the individual V2, V4, and V5 loop glycans, there was clear signal around these sites in the

0-2hr difference map only (Fig. 9*B*), indicating the dynamic V-loop glycans were completely digested within 2hrs. We

also found partial signal reduction at a few sites, indicative of non-uniform digestion. For example, the apex cluster

composed of the N156 and N160 glycans as well as the glycans at N133, N197, and N234. After 16 hours the glycan

shield was almost completely digested, however, we still detected some residual glycan signal at the previously

discussed cluster composed of the N363, N386, and N197 glycans, as well as a cluster composed of the N295, N332,

and N448 glycans. In addition, the highly protected glycan at N262 remained completely intact. By quantifying the

degree of protection from Endo H (see Methods) and comparing it to the predicted network eigencentralities for the

Man9 ensemble (Fig. 9*E*), we obtained a correlation coefficient of ~0.8 (p=1.14e-05), suggesting highly connected

glycans are resistant to enzymatic digestion. Also evident is the similarity between the persistent glycan clusters and

the sub-graphs generated by progressively stripping the glycan overlap network (*SI Appendix*, Fig. S10*B*), which could

be seen as mimicking the gradual digestion by Endo H.

375   Interestingly, 3-D classification of the Endo H treated datasets revealed an increasing degree of protein

unfolding and subunit dissociation initiating at the V1-3 loops in the trimer apex. Classification consistently converged

to four 4 classes that appeared to capture found heterogeneous intermediates in the unfolding process (Fig. 10*A* and

SI Appendix, Fig. S13-17). We interpreted these as representing (1) stable trimers, (2) trimers with unfolded or partially

unfolded V1-V3 loops, (3) trimers with fully unfolded/dissociated gp120s, and (4) monomers/dimers. As the reaction

progressed, the percentages of unfolded trimers increased, while the percentage of stably folded trimers decreased

(Fig. 10*B*). Although it cannot be easily confirmed, the unfolded trimers in each dataset were likely more completely

deglycosylated than the particles that make up the stable trimeric classes. We also observed a small percentage of

dissociated gp120s in the CHO sample, perhaps as a result of the higher degree of sub-occupancy. When viewed in

context of the results presented above, we concluded that the highly connected glycans and glycan clusters that were

resistant to digestion are also critical to maintaining structural stability of the pre-fusion Env trimer, suggesting the

glycan shield may have additional functions beyond its role in immune evasion.

Discussion

Prior to this study, there were conflicting reports of glycan shield structure from cryo-EM and X-ray

crystallography measurements (6, 14, 35). Our results clearly show that the native glycan shield is highly dynamic

with respect to the protein core. We did not observe any stabilized glycan-glycan interactions like those reported in

crystal structures (6), suggesting methodological differences may be responsible for the apparently contradictory

results. For example, the crystal structures included two bound bnAbs per monomer to facilitate crystal packing, both

of which engage multiple glycans and disrupt the native dynamics and higher-order structure. Desolvation of mobile

waters embedded within the glycan shield during crystallization could also potentially induce the stabilized glycan-

glycan contacts observed. A favorable interpretation that also fits with the conclusions of this paper is that the crystal

structures captured artificially stabilized yet physiologically relevant conformations and are thus suggestive of the

types transient, of non-covalent interactions possible between glycans.

Our work represents the first case of using cryo-EM to validate glycoprotein ensembles, and in turn, their

use in guiding cryo-EM analysis. We found that the HT-AM pipeline captured physiologically relevant glycan

sampling at global and local scales, however, deviated from the experimental observations when the flexibility of the

underlying protein was greater than could be captured by the current pipeline. Thus, exploring methods to enhance

sampling of the protein backbone, particularly at Env variable loops, represents a clear route for improving the

accuracy of the pipeline (36). Another logical next step is to use the experimental cryo-EM maps to steer the

ensemble modeling process, which has been successful for non-glycosylated proteins (37-39), however, extending

such techniques to capture the extreme levels of heterogeneity present in the glycan shield will be challenging.

Experimentally, our ability to capture differences in glycan dynamics from cryo-EM maps is limited when using local

map intensity around BMA residues alone, and such limitations could also be contributing to the observed

discrepancies. Again, using the cryo-EM maps to steer the modeling process could provide more accurate and

comprehensive assessments of variations in relative dynamics between glycan in addition to improving modeling

accuracy. Finally, utilizing sequence engineering and/or different expression systems to achieve a more uniformly

glycosylated Env sample, or conversely, incorporating experimentally relevant levels of heterogeneity into the

modeling pipeline will allow for a more accurate comparison between theory and experiment.

Using graph theory, we mapped out the complex network of potential interactions within the glycan shield.

A similar network-based approach was introduced previously to capture concerted behavior within the glycan shield

during all-atom MD simulations **(22)**. Although the HT-AM pipeline does not capture temporal dynamics, we found

that the probabilistic networks reported here closely resembled the ones derived from the MD simulations, and the

two lead to similar predictions. For instance, the clustering of glycans into larger microdomains (*SI Appendix*, Fig. S10*A*-

B). Moreover, the overlap probability between glycans gives a measure of the structural shielding rather than the

energetic interactions between glycans. At a qualitative level, the hierarchy of structural features we observed in the

Gaussian filtered and 3-D variability maps (Fig. 2*C-D*) could be seen as the ensemble-average manifestation of this

clustering behavior. Given the preference of bnAbs for targeting the interfaces between higher-order glycan clusters

(22), experimentally mapping such long-range structure could be important. At a more quantitative level, we found

that the average degree of protection from Endo H digestion correlated strongly with network eigencentrality for

those glycans we could identify in the cryo-EM maps, and that the order of digestion closely matched the order in

which glycans were removed from the network when applying stricter overlap thresholds. It should be noted,

however, that because of the somewhat simplified approach we have taken to modeling the most dynamic segments

of the variable loops, in particular in the V4 loop segment containing the N398, N406, and N411 glycans, the

eigencentrality of these glycans may be artificially high, leading to minor discrepancies between digestion

susceptibility and network centrality. Nonetheless, these correlations could be seen as providing indirect validation of

the network models, where the high centrality glycans are either located in densely packed regions where Endo H

would have a difficult time accessing (*SI Appendix*, Fig. S6*D-E*) or are protected by strong interaction probability with

their neighbors.

We demonstrated that cryo-EM is capable of detecting and quantifying changes in glycan dynamics, as well

as occupancy and chemical composition between differentially glycosylated Envs, the latter of which were exclusively

provided by MS prior to this study. Thus, cryo-EM provides validation for these measurements while contributing

insight into the structural impact of changes in glycosylation. For instance, semi-quantitative methods to assess the

distribution of unoccupied sites have only recently become available **(13)**, and here we showed that cryo-EM is highly

sensitive to occupancy. Importantly, we found evidence for more extensive changes in glycosylation between the

293F and CHO expressed samples than detected by MS, which has clear relevance to the currently on-going human

clinical trials based on the CHO expressed Env. In addition to these results, we found that the extent of processing

measured by the percentage of high-mannose type glycans at each site was strongly correlated with ordering in the

cryo-EM map (*SI Appendix*, Fig. S2*C-D*), suggesting it may be possible to predict the degree of processing at each site

from cryo-EM maps alone. The relationship between these variables likely reflects their mutual dependence upon

glycan crowding (*SI Appendix*, Fig. S6*B-C*), as well as the properties surrounding protein structure, both of which can

reduce dynamics and restrict the access of Endo H and glycan processing enzymes in the cell.

The observation that enzymatic deglycosylation leads to progressive destabilization of the Env trimer was

somewhat surprising because recent investigations into the effect of glycan knockouts and deglycosylation on Env

stability and viral infectivity have come to different conclusions (40-44). However, there is ample biophysical

evidence that glycans influence protein stability, dynamics, and folding, even in the case of Env (17, 45-56),

although it is not obvious which mechanisms are important here. One possible mechanism is stabilizing interactions

between the core NAG and neighboring side chains, which are observed throughout Env and are common in other

glycoproteins. However, Endo H leaves the core NAG attached, so the stabilizing effect must arise by other

mechanisms. It has been proposed that glycans stabilize proteins primarily by destabilizing the unfolded state and by

increasing the height of the unfolding energy barrier (47). The same study also found that glycosylation reduced the

entropy of the folded state, possibly by imposing pressure on the protein and dampening dynamics, similar to the

effects of molecular crowding or confinement, although the effect was canceled out by an equivalent change in

enthalpy. Along these lines, the slightly lower resolution of the BG505_EndoH16 map compared the fully glycosylated

maps could be indicative of increased dynamics. In addition, our observation that the Man9 glycans were less dynamic

than the smaller Man5 glycans in our simulations could be attributed to more crowding in the canopy, which in turn

could further stabilize the underlying protein. Also, the stabilizing effects were found to depend strongly on the

number of glycans as well as their positions on the protein (47), which is in line with the results of our Endo H

digestion experiment. Therefore, given the high density of glycans on the surface of Env and the importance of their

tertiary arrangement into high density clusters, it is plausible these stabilizing effects could be amplified. Furthermore,

glycan-glycan interactions such as those observed in the crystal structures solved by Stewart-Jones et. al. (6) could

also play a role in stabilizing folded state. Although our results clearly show that such interactions must be more

dynamic than the crystal structures suggest, the combined effect of many such short-lived interactions could become

significant.

The time-resolved Endo H digestion experiment provide an explicit mapping of the importance of each glycan

toward maintaining the integrity of the glycan shield and the overall stability of Env. In addition to the glycan at N262,

the two highly connected clusters in the gp120 outer domain centered around the N386 and N295 glycans were the

most important for trimer stability, and to a lesser extent the apex cluster composed primarily of the N156 and N160

glycans. In support of this, previous studies have shown that knocking out the glycans at N386 or N262 individually

have dramatic effects on trimer stability (33), whereas deletion of up to 5 glycans at once peripheral to the CD4bs,

which are not part of these cluster, are well tolerated (10, 57, 58). In addition, the glycans at N386 and N295 in

particular were found to strongly control the processing of the other glycans near them, reinforcing their importance

in maintaining the integrity of large-scale glycan shield structure (33). These results have implications for rational

vaccine design where removing certain glycans could have undesired effects on protein stability and suggest that

there are limits to the number glycan deletions that can be tolerated, with particular emphasis on maintaining the

integrity of the aforementioned clusters. In addition, we found that instability originated at the trimer apex, around

the V1-3 loops. This region of the structure is known to be metastable and to change conformation upon CD4 and co-

receptor binding(34, 59). Therefore, it is plausible the glycans and glycan clusters in this region are contributing to

maintaining Env in a metastable state poised for receptor binding. Given the high conservation of the glycan clusters

identified here, our results are likely universal to all Env sequences. Lastly, it is possible that glycan induced

stabilization of Env may enable hyper-mutation of the underlying protein surface, thereby endowing Env with the

ability to escape immune pressure with otherwise deleterious mutations.

It is known that immune responses to Env preferentially target glycan-depleted surface area (9, 10), and

the results presented here provide the first experimentally determined mapping of this surface. Under reasonable

physical assumptions, the ensemble-averaged structure captured by cryo-EM should be identical to the time-averaged

structure of a single thermally fluctuating Env molecule at the temperature prior to vitrification. The dynamics of the

glycans create a cloud/shield over the protein, thus an approaching antibody with relatively torpid dynamics will

effectively "see" this blurry ensemble-average structure, which is consistent with the relatively slow on-rates of most

HIV bnAbs. We quantified the shielding effect with a simple "rolling sphere" method for BG505_CHO and visualized

the results as a colormap projected onto the surface of Env (Fig. 11). White surfaces are strongly masked by nearby

glycans and red surfaces are weakly masked, highlighting sites of potential vulnerability to neutralizing antibodies (see

the *SI Appendix* for details). Although the cryo-EM maps presented here do not perfectly capture the in-vivo structure

of a single fluctuating Env because they are ensemble averages of chemically heterogeneous molecules, if one

considers the net serum response to a vaccine that itself contains the exact same heterogeneity as the sample we

analyzed here (60, 61), they still provide an accurate representation of the average surface exposure of the entire

ensemble.

Looking beyond the results of this paper, our integrated approach can be easily extended to other Env clones

and to heavily glycosylated spike proteins from other viruses such as Influenza, Ebola, Lassa, and Coronaviruses, and

represents a potentially powerful approach for studying the structure and dynamics of glycoproteins in general.

Materials and Methods

BG505 SOSIP.664v3 was expressed and purified from HEK293F and HEK293S suspension cell culture in house while

the CHO cell-line derived BG505 SOSIP.664 was provided to us by IAVI and expressed and purified as described in

Dey, et al., 2017 (31). The monoclonal antibody RM20A3 was isolated from a BG505 SOSIP.664 immunized rhesus

macaque(62) and the Fab was expressed and purified from HEK293F cells as described previously(63). The BG505

SOSIP.664 samples expressed in HEK293F and CHO cell lines were prepared for MS analysis as described

previously(25), and the HEK293S sample was prepared with slight modifications on that protocol. All samples were

prepared for cryo-EM analysis via the same protocol after being complexed with RM20A3 Fab. Imaging was

performed on either an FEI Titan Krios or Talos Arctica (Thermo-Fisher) microscope. All non-custom cryo-EM data

processing steps were performed using a combination RELION-2/3(64, 65) and CryoSparc v1(66). Model building

and refinement was carried out with UCSF Chimera (67), Coot (68, 69), and Rosetta(70). All custom analysis of

cryo-EM maps were performed in MatLab (2018b) {MATLAB and Image Processing Toolbox Release 2018b, The

MathWorks, Inc., Natick, Massachusetts, United States.} 3-D variability analysis was performed in SPARX(27). The

HT-AM was carried out by implementing the ALLOSMOD(29, 71) package of MODELLER(72, 73) in a

streamlined pipeline. All graph theory and network-based analysis were performed using Python {Python Software

Foundation, https://www.python.org/} and Matlab_R2018a packages. Simulated cryo-EM data generation and

analysis was performed in RELION. All cryo-EM map and model visualization was performed with UCSF Chimera. A

comprehensive description of all the methods can be found in the *SI Appendix.*

Acknowledgements

Z.T.B would like to acknowledge Bill Anderson and Hannah Turner for assistance with microscopy, Charles Bowman

and JC Ducom for computational support, Pawel Penczek, Francisco Asturias, Jack Johnson, and Gabe Lander for

helpful discussions on 3-D variance analysis and cryo-EM data processing, Jesper Pallesen and JH Lee for early

support with data processing, Jeffrey Copps for help with protein expression and purification, Lauren Holden for

critical reading of the manuscript, and finally IAVI, Antu Dey, and John P. Moore for supplying demo run material and

MS analysis. S.C and S.G. acknowledge Timothy Travers for proposing ALLOSMOD for this work, Nicolas Hengartner

for helpful discussions on graph theory and Bette Korber and Kshitij Wagh for insights on glycosylations in HIV. S.C.

and S.G. acknowledge the LANL High Performance Computing Division for providing computational facilities. We

thank Robin Sung Kyu Park and Titus Jung for help with analysis of MS data for site specific glycosylation.

Additional Information

Competing Interests

The authors declare no competing interests.

Funding

This work was supported by the National Institute of Allergy and Infectious Diseases grants: UM1 AI100663 (to A.B.W.,

J.C.P.), UM1 AI144462 (to A.B.W., J.C.P.), AI113867 (to J.C.P.), and P41GM103533 (to J.R.Y.). This work was supported

by the Bill and Melinda Gates Foundation through the Collaboration for AIDS Vaccine Discovery (CAVD) grants

OPP1100639 and OPP1115782 (A.B.W.). This work was supported by the IAVI Neutralizing Antibody Center through

the Collaboration for AIDS Vaccine Discovery grant OPP1196345 funded by the Bill and Melinda Gates Foundation

(A.B.W.). This work was funded by grants from the NIH (Center for HIV/AIDS Vaccine Immunology and immunogen

Discovery, UM1 AI100645; Consortia for HIV/AIDS Vaccine Development, UM1 AI144371). S.C. was also partially

supported by the Center for Nonlinear Studies (CNLS) at Los Alamos National Laboratory (LANL).

Author contributions

Conceptualization, Z.T.B, S.C., C.A.L, S.G. and A.B.W.; Methodology, Z.T.B, S.C., C.A.L, X.W., J.C.P., S.G. and A.B.W.;

Protein expression and purification, Z.T.B., C.A.C, and J.L.T; Antibody isolation, M.J.vG; Mass-spectrometry

experiments and data analysis, X.W, J.K.D, J.R.Y., and J.C.P.; Cryo-EM experiments, Z.T.B.; Modeling and Simulations,

S.C.; Formal Analysis, Z.T.B. and S.C.; Visualization, Z.T.B. and S.C.; Manuscript Preparation, Z.T.B, S.C., C.A.L, J.C.P.,

S.G. and A.B.W.; Supervision, S.G. and A.B.W.

Data Availability

All data needed to evaluate the conclusions in the paper are present in the paper and/or the Supplementary

Information. Additional data related to this paper may be requested from the authors. The electron potential maps

for BG505_293F, BG505_CHO, BG505_293S, BG505_EndoH2, and BG505_EndoH16 have been deposited in the

Electron Microscopy Data Bank under accession codes 22108, 22109, 22110, 22111, and 22112 respectively, along

with all corresponding half-maps, masks, and SPARX 3-D variability maps. Atomic models have been deposited in the

Protein Data Bank under accession codes 6X9R, 6X9S, 6X9T, 6X9U, 6X9V.

References

1. Fenouillet E, Gluckman JC, Jones IM (1994) Functions of HIV envelope glycans. *Trends in Biochemical*

Sciences 19(2):65–70.

2. Grundner C, et al. (2004) Factors limiting the immunogenicity of HIV-1 gp120 envelope glycoproteins.

Virology 330(1):233–248.

3. Dacheux L, et al. (2004) Evolutionary dynamics of the glycan shield of the human immunodeficiency virus

envelope during natural infection and implications for exposure of the 2G12 epitope. *J Virol* 78(22):12625–

12637.

4. Lasky LA, et al. (1986) Neutralization of the AIDS retrovirus by antibodies to a recombinant envelope

glycoprotein. *Science* 233(4760):209–212.

5. Doores KJ, et al. (2010) Envelope glycans of immunodeficiency virions are almost entirely oligomannose

antigens. *Proc Natl Acad Sci USA* 107(31):13800–13805.

6. Stewart-Jones GBE, et al. (2016) Trimeric HIV-1-Env Structures Define Glycan Shields from Clades A, B, and

G. *Cell* 165(4):813–826.

7. Korber B, et al. (2001) Evolutionary and immunological implications of contemporary HIV-1 variation. *Br*

Med Bull 58(1):19–42.

8. Leonard CK, et al. (1990) Assignment of intrachain disulfide bonds and characterization of potential

glycosylation sites of the type 1 recombinant human immunodeficiency virus envelope glycoprotein

(gp120) expressed in Chinese hamster ovary cells. *Journal of Biological Chemistry* 265(18):10373–10382.

9. Wei X, et al. (2003) Antibody neutralization and escape by HIV-1. *Nature* 422(6929):307–312.

10. Zhou T, et al. (2017) Quantification of the Impact of the HIV-1-Glycan Shield on Antibody Elicitation. *Cell*
Rep 19(4):719–732.

11. Ward AB, Wilson IA (2017) The HIV-1 envelope glycoprotein structure: nailing down a moving target.
Immunol Rev 275(1):21–32.

12. Schwarz F, Aebi M (2011) Mechanisms and principles of N-linked protein glycosylation. *Curr Opin Struct*
Biol 21(5):576–582.

13. Cao L, et al. (2017) Global site-specific N-glycosylation analysis of HIV envelope glycoprotein. *Nat Commun*
8:14954.

14. Lyumkis D, et al. (2013) Cryo-EM structure of a fully glycosylated soluble cleaved HIV-1 envelope trimer.
Science 342(6165):1484–1490.

15. Lee JH, Ozorowski G, Ward AB (2016) Cryo-EM structure of a native, fully glycosylated, cleaved HIV-1
envelope trimer. *Science* 351(6277):1043–1048.

16. Kumar S, Cieplak P (2018) Role of N-glycosylation in activation of proMMP-9. A molecular dynamics
simulations study. *PLoS ONE* 13(1):e0191157.

17. Lee HS, Qi Y, Im W (2015) Effects of N-glycosylation on protein conformation and dynamics: Protein Data
Bank analysis and molecular dynamics simulation study. *Sci Rep* 5(1):8926.

18. Dong C, et al. (2018) Long-ranged Protein-glycan Interactions Stabilize von Willebrand Factor A2 Domain
from Mechanical Unfolding. *Sci Rep* 8(1):16017–11.

19. Hang I, et al. (2015) Analysis of site-specific N-glycan remodeling in the endoplasmic reticulum and the
Golgi. *Glycobiology* 25(12):1335–1349.

20. Ferreira R-C, et al. (2018) Structural Rearrangements Maintain the Glycan Shield of an HIV-1 Envelope
Trimer After the Loss of a Glycan. *Sci Rep* 8(1):15031–15.

21. Yang M, Huang J, Simon R, Wang L-X, MacKerell AD (2017) Conformational Heterogeneity of the HIV
Envelope Glycan Shield. *Sci Rep* 7(1):4435–15.

22. Lemmin T, Soto C, Stuckey J, Kwong PD (2017) Microsecond Dynamics and Network Analysis of the HIV-1
SOSIP Env Trimer Reveal Collective Behavior and Conserved Microdomains of the Glycan Shield. *Structure*
25(10):1631–1639.e2.

23. Sanders RW, et al. (2013) A next-generation cleaved, soluble HIV-1 Env trimer, BG505 SOSIP.664 gp140,
expresses multiple epitopes for broadly neutralizing but not non-neutralizing antibodies. *PLoS Pathog*
9(9):e1003618.

24. Sanders RW, Moore JP (2017) Native-like Env trimers as a platform for HIV-1 vaccine design. *Immunol Rev*
275(1):161–182.

25. Cao L, et al. (2018) Differential processing of HIV envelope glycans on the virus and soluble recombinant
trimer. *Nat Commun* 9(1):3693.

26. Behrens A-J, et al. (2016) Composition and Antigenic Effects of Individual Glycan Sites of a Trimeric HIV-1
Envelope Glycoprotein. *Cell Rep* 14(11):2695–2706.

27. Hohn M, et al. (2007) SPARX, a new environment for Cryo-EM image processing. *J Struct Biol* 157(1):47–
55.

28. Cheng Y, Grigorieff N, Penczek PA, Walz T (2015) A primer to single-particle cryo-electron microscopy. *Cell*
161(3):438–449.

29. Guttman M, Weinkam P, Sali A, Lee KK (2013) All-atom ensemble modeling to analyze small-angle x-ray
scattering of glycosylated proteins. *Structure* 21(3):321–331.

30. Liang Y, et al. (2016) Changes in Structure and Antigenicity of HIV-1 Env Trimers Resulting from Removal of
a Conserved CD4 Binding Site-Proximal Glycan. *J Virol* 90(20):9224–9236.

31. Dey AK, et al. (2017) cGMP production and analysis of BG505 SOSIP.664, an extensively glycosylated,
trimeric HIV-1 envelope glycoprotein vaccine candidate. *Biotechnol Bioeng*. doi:10.1002/bit.26498.

32. Behrens A-J, et al. (2018) Integrity of Glycosylation Processing of a Glycan-Depleted Trimeric HIV-1
Immunogen Targeting Key B-Cell Lineages. *J Proteome Res*:acs.jproteome.7b00639.

33. Pritchard LK, et al. (2015) Glycan clustering stabilizes the mannose patch of HIV-1 and preserves
vulnerability to broadly neutralizing antibodies. *Nat Commun* 6:7479.

34. Kwong PD, et al. (1998) Structure of an HIV gp120 envelope glycoprotein in complex with the CD4 receptor
and a neutralizing human antibody. *Nature* 393(6686):648–659.

35. Lee JH, de Val N, Lyumkis D, Ward AB (2015) Model Building and Refinement of a Natively Glycosylated
HIV-1 Env Protein by High-Resolution Cryoelectron Microscopy. *Structure* 23(10):1943–1951.

36. Chakraborty S, et al. (2020) Quantification of the Resilience and Vulnerability of HIV-1 Native Glycan Shield
at Atomistic Detail. *bioRxiv*:846071.

37. Bonomi M, Pellarin R, Vendruscolo M (2018) Simultaneous Determination of Protein Structure and
Dynamics Using Cryo-Electron Microscopy. *Biophysical Journal* 114(7):1604–1613.

38. Bonomi M, et al. (2019) Bayesian Weighing of Electron Cryo-Microscopy Data for Integrative Structural
Modeling. *Structure* 27(1):175–188.e6.

39. Bonomi M, Vendruscolo M (2019) Determination of protein structural ensembles using cryo-electron
microscopy. *Curr Opin Struct Biol* 56:37–45.

40. Rathore U, et al. (2017) Glycosylation of the core of the HIV-1 envelope subunit protein gp120 is not
required for native trimer formation or viral infectivity. *J Biol Chem* 292(24):10197–10219.

41. Depetris RS, et al. (2012) Partial enzymatic deglycosylation preserves the structure of cleaved recombinant
HIV-1 envelope glycoprotein trimers. *J Biol Chem* 287(29):24239–24254.

42. Wang W, et al. (2013) A systematic study of the N-glycosylation sites of HIV-1 envelope protein on
infectivity and antibody-mediated neutralization. *Retrovirology* 10(1):14.

43. Lee J-Y, et al. (2016) Crystal structure of the human sterol transporter ABCG5/ABCG8. *Nature*
533(7604):561–564.

44. Ohgimoto S, et al. (1998) Location-specific, unequal contribution of the N glycans in simian
immunodeficiency virus gp120 to viral infectivity and removal of multiple glycans without disturbing
infectivity. *J Virol* 72(10):8365–8370.

45. Helenius A, Aebi M (2001) Intracellular functions of N-linked glycans. *Science* 291(5512):2364–2369.

46. Solá RJ, Rodríguez-Martínez JA, Griebenow K (2007) Modulation of protein biophysical properties by
chemical glycosylation: biochemical insights and biomedical implications. *Cell Mol Life Sci* 64(16):2133–
2152.

47. Shental-Bechor D, Levy Y (2008) Effect of glycosylation on protein folding: a close look at thermodynamic
stabilization. *Proc Natl Acad Sci USA* 105(24):8256–8261.

48. Imperiali B, O'Connor SE (1999) Effect of N-linked glycosylation on glycopeptide and glycoprotein structure.
Curr Opin Chem Biol 3(6):643–649.

49. Tian J, et al. (2016) Effect of Glycosylation on an Immunodominant Region in the V1V2 Variable Domain of
the HIV-1 Envelope gp120 Protein. *PLoS Comput Biol* 12(10):e1005094.

50. Wood NT, et al. (2013) The influence of N-linked glycans on the molecular dynamics of the HIV-1 gp120 V3
loop. *PLoS ONE* 8(11):e80301.

51. Gerken TA, Butenhof KJ, Shogren R (1989) Effects of glycosylation on the conformation and dynamics of O-
linked glycoproteins: carbon-13 NMR studies of ovine submaxillary mucin. *Biochemistry* 28(13):5536–
5543.

52. Mer G, Hietter H, Lefèvre JF (1996) Stabilization of proteins by glycosylation examined by NMR analysis of
a fucosylated proteinase inhibitor. *Nat Struct Biol* 3(1):45–53.

53. Joao HC, Dwek RA (1993) Effects of glycosylation on protein structure and dynamics in ribonuclease B and
some of its individual glycoforms. *Eur J Biochem* 218(1):239–244.

54. Lu D, Yang C, Liu Z (2012) How hydrophobicity and the glycosylation site of glycans affect protein folding
and stability: a molecular dynamics simulation. *J Phys Chem B* 116(1):390–400.

55. Shental-Bechor D, Levy Y (2009) Folding of glycoproteins: toward understanding the biophysics of the
glycosylation code. *Curr Opin Struct Biol* 19(5):524–533.

56. Hanson SR, et al. (2009) The core trisaccharide of an N-linked glycoprotein intrinsically accelerates folding
and enhances stability. *Proc Natl Acad Sci USA* 106(9):3131–3136.

57. Dubrovskaya V, et al. (2019) Vaccination with Glycan-Modified HIV NFL Envelope Trimer-Liposomes Elicits
Broadly Neutralizing Antibodies to Multiple Sites of Vulnerability. *Immunity* 51(5):915–929.e7.

58. Dubrovskaya V, et al. (2017) Targeted N-glycan deletion at the receptor-binding site retains HIV Env NFL
trimer integrity and accelerates the elicited antibody response. *PLoS Pathog* 13(9):e1006614.

59. Ozorowski G, et al. (2017) Open and closed structures reveal allostery and pliability in the HIV-1 envelope
spike. *Nature* 15(7663):690–363.

60. Bianchi M, et al. (2018) Electron-Microscopy-Based Epitope Mapping Defines Specificities of Polyclonal
Antibodies Elicited during HIV-1 BG505 Envelope Trimer Immunization. *Immunity* 49(2):288–300.e8.

61. Nogal B, et al. (2020) Mapping Polyclonal Antibody Responses in Non-human Primates Vaccinated with HIV
Env Trimer Subunit Vaccines. *Cell Rep* 30(11):3755–3765.e7.

62. Cottrell CA, et al. (2020) Mapping the immunogenic landscape of near-native HIV-1 envelope trimers in
non-human primates. *bioRxiv*:2020.02.05.936096.

63. Steichen JM, et al. (2019) A generalized HIV vaccine design strategy for priming of broadly neutralizing
antibody responses. *Science* 366(6470):eaax4380.

64. Kimanius D, Forsberg BO, Scheres SH, Lindahl E (2016) Accelerated cryo-EM structure determination with
parallelisation using GPUs in RELION-2. *Elife* 5:19.

65. Zivanov J, et al. (2018) New tools for automated high-resolution cryo-EM structure determination in
RELION-3. *Elife* 7:163.

66. Punjani A, Rubinstein JL, Fleet DJ, Brubaker MA (2017) cryoSPARC: algorithms for rapid unsupervised cryo-
EM structure determination. *Nat Methods* 14(3):290–296.

67. Pettersen EF, et al. (2004) UCSF Chimera--a visualization system for exploratory research and analysis. *J
Comput Chem* 25(13):1605–1612.

68. Emsley P, Crispin M (2018) Structural analysis of glycoproteins: building N-linked glycans with Coot. *Acta
Crystallogr D Struct Biol* 74(Pt 4):256–263.

69. Emsley P, Cowtan K (2004) Coot: model-building tools for molecular graphics. *Acta Crystallogr D Biol
Crystallogr* 60(Pt 12 Pt 1):2126–2132.

70. Wang RY-R, et al. (2016) Automated structure refinement of macromolecular assemblies from cryo-EM
maps using Rosetta. *Elife* 5:352.

71. Weinkam P, Pons J, Sali A (2012) Structure-based model of allostery predicts coupling between distant
sites. *Proc Natl Acad Sci USA* 109(13):4875–4880.

72. Eswar N, et al. (2006) Comparative protein structure modeling using Modeller. *Curr Protoc Bioinformatics*
Chapter 5(1):Unit–5.6.

73. Sali A (1995) Comparative protein modeling by satisfaction of spatial restraints. *Mol Med Today* 1(6):270–
277.

74. Varki A, et al. (2015) Symbol Nomenclature for Graphical Representations of Glycans. *Glycobiology*
25(12):1323–1324.

Figures

Fig. 1 | Soluble SOSIP in complex with a base-specific Fab as a model system for cryo-EM analysis of the native

HIV-1 Env glycan shield. (A) Segmented cryo-EM map of BG505 SOSIP.664 in complex with three copies of the

RM20A3 Fab. The green circle is meant to represents the glycan shield. **(B)** Refined atomic model with stars

highlighting several N-linked glycans. **(C)** 3-D angular distribution histogram. **(D)** Surface representation of the refined

atomic model viewed from the side and from the bottom with the Fab chains removed. **(E)** Sharpened 3.1Å-resolution

cryo-EM map at high and low threshold colored by local resolution. **(F)** SNFG depiction (74) of a representative high-

mannose and complex type glycan (L). Cryo-EM map density for 6 representative glycans from the refined atomic

model (numbers corresponding to the stars in panel B).

730
Fig. 2 | Scale-space and 3-D variability analysis reveal low resolution structure in the glycan shield. (A) Noise

threshold and volume at the noise threshold as a function of Gaussian filter width measured in standard deviation

(SD) showing the emergence of glycan signal at low resolutions and plateau around 1.5 SD. **(B)** BG505_293F cryo-EM

map at 6 representative scales along with 2-D slices through the maps overlaid on top of one another showing the

emergence of glycan signal and new glycan-glycan contacts visible at low resolutions and thresholds (red arrows).

Black arrows are pointing to protein surfaces occluded by glycans. **(C)** Gaussian filtered (1.5 SD) BG505_293F map

(blue) visualized at three intensity thresholds along with a protein-only map (gray). **(D)** SPARX 3-D variability map

(light blue) visualized at three intensity thresholds along with the BG505_293F map (gray) and 2-D slices through the

top and side of the map with red arrows highlighting glycan-glycan contacts.

Fig. 3 | ALLOSMOD-based HT-AM pipeline for fast and robust sampling of fully glycosylated Env. (A) 10

homology models used as protein scaffolds. **(B)** One of the ten models with a single relaxed Man9 glycan at each

site. **(C)** One fully glycosylated and relaxed model for each of the ten protein scaffolds. **(D)** Close up of the N88

glycan showing a single glycan pose (L) and all 1000 poses (R). **(E)** Full glycan average rmsf for all 1000 poses. **(F)**

Glycan only rmsf for all 1000 poses. **(E)** Full glycan average rmsf for all 1000 poses at the 5th mannose residue for

both the Man9 and Man5 ensembles.

Fig. 4| Simulated cryo-EM maps from the HT-AM ensembles reproduce defining features of the experimental

data. (A) Schematic of the simulated data creation pipeline. Each model is projected at 100 uniformly distributed

angles (representative 2-D projections with and without noise are shown), then the combined dataset of 100K

projections is refined using RELION (the resulting 3-D reconstruction and angular distribution histogram are shown).

(B) Noise threshold and volume at the noise threshold as a function of Gaussian filter width (SD) for the

BG505_Man9, BG505_Man5, and BG505_PO simulated cryo-EM maps with dashed lines indicating the volume

sampled by each ensemble. **(C)** 1.5 SD Gaussian filtered BG505_Man9 (teal) and BG505_PO (gray) maps along with

the SPARX 3-D variability map (light blue) viewed from the side and top at high and low ntensity thresholds.

Fig. 5 | Measuring glycan dynamics from cryo-EM maps reveals close agreement between simulation and

experiment. (A) 1.5 SD Gaussian filtered BG505_Man9 simulated cryo-EM map (transparent) and atomic model

used for local intensity analysis. Also shown is a single glycan stem with a spherical probe around the BMA residue.

(**B**) Pearson correlation coefficient and p-value between mean intensity and inverse rmsf as a function of probe

radius for 5 different Gaussian filtered maps. The dashed line designates the confidence threshold (p-value = 0.05).

(**C**) Normalized mean intensity around BMA residues for BG505_Man9 (probe radius = 2.3Å) and the inverse

normalized full glycan average rmsf. Pearson correlation coefficient ~0.88 (p-value = 2.1e-8). **(D)** Normalized mean

intensity around BMA residues for BG505_293F and BG505_Man9 along with the absolute difference at each site.

Pearson correlation coefficient ~0.46 (p = 0.03).

Fig. 6 | Detecting changes in glycan dynamics, occupancy, and chemical composition from simulated cryo-EM maps. (A) Mean intensity around each glycan BMA residue for the BG505_Man9, BG505_Man5, and BG505_Man9HO maps along with the percent change from BG505_Man9. **(B)** BG505_Man9 - BG505_Man9HO and **(C)** BG505_Man9 - BG505_Man5 difference maps (Gaussian filtered) at two intensity thresholds (right - green) along with high-resolution sharpened maps (left).

Fig. 7 | Detection of cell-type specific changes in glycan shield composition and dynamics with cryo-EM. (**A**) Mean intensity at each glycan BMA residue for the BG505_293F, BG505_293S, and BG505_CHO cryo-EM maps along with percent change from BG505_293F. (**B**) BG505_293F − BG505_293S, (**C**) BG505_CHO − BG505_293S, and (**C**) BG505_293F - BG505_CHO difference maps at two intensity thresholds. All difference maps were multiplied by a soft mask around the RM20A3 Fabs then smoothed with a 2 SD Gaussian filter. (**E-H**) Fly-outs of the regions outlined by dashed red boxes.

Fig. 8 | Probabilistic glycan-glycan interaction network for the Man9 ensemble. (A) Glycan-glycan volume overlap

matrix. Inter-protomer overlap is given by suffix "_2" and the variable loop glycans are indicated by tan color bars.

The overlap fraction is normalized with at least 50% overlap being designated as 1. **(B)** Cartoon model of the glycan-

glycan interaction network generated by interpreting the overlap matrix as an adjacency matrix. The edge length is

drawn inversely proportional to the overlap value. Edges represent overlap between two glycans represented by

nodes. **(C)** Glycan-glycan interaction network calculated from the matrix in (A) with nodes colored by normalized

eigenvector centrality.

Fig. 9 | Time-resolved cryo-EM reveals that highly connected glycan clusters are resistant to enzymatic

digestion. (A) Cartoon schematic of the Endo H digestion of high-mannose glycans and an illustration of the three

reaction times captured by time-resolved cryo-EM. **(B)** Mean intensity around glycan BMA residues for the

BG505_293S, BG505_EndoH2, and BG505_EndoH16 cryo-EM maps along with the percent occupancy at each site.

Occupancy was calculated assuming a linear relationship between occupancy and intensity with the initial

occupancy determined by MS. **(C)** BG505_293S − BG505_EndoH2 and BG505_EndoH2 − BG505_EndoH16 difference

maps at two intensity thresholds. Difference maps were first multiplied by a soft mask around the RM20A3 Fabs and

smoothed with a 2 SD Gaussian filter. Also shown is the residual glycan signal remaining after 16hrs of digestion

isolated by masking out the protein density. **(D)** Overlay of the three maps from panel B colored according to

susceptibility to Endo H digestion. **(E)** Normalized Endo H protection score measured as the cumulative occupancy

at each site in the two digestion intermediates and the normalized eigencentrality from the Man9 volume overlap

network. Pearson correlation coefficient of ~0.8 (p=1.14e-05).

A *stable trimer* *partially unfolded V1-V3 loops* *dissociated/ unfolded gp120* *monomer/dimer*

B *293F* *CHO* *293S* *EndoH2* *EndoH16*

100% 84% / 16% 100% 58% / 28% / 14% 32% / 39% / 12% / 17%

■ *stable trimer* ■ *unfolding V1-3 loops* ■ *dissociated/unfolded gp120* ■ *monomer/dimer*

Fig. 10 | Digestion of the glycan shield leads to progressive destabilization of the Env timer. **(A)** Representative cryo-EM maps of the unfolding intermediates captured by 3-D classification. **(B)** Pie charts showing the relative percentages of each class present in the five cryo-EM datasets presented here.

Fig. 11 | Quantifying the glycan shielding effect from cryo-EM maps. The normalized glycan shielding effect for BG505_CHO defined as the total glycan signal within a ~7Å radius spherical probe centered at each voxel. White surfaces are strongly shield by glycans.