Size, not temperature, drives cyclopoid copepod predation of invasive

mosquito larvae

Marie C. Russell[1][†], Alima Qureshi[1], Christopher G. Wilson[1], and Lauren J. Cator[1]

[1]Department of Life Sciences, Imperial College London, Silwood Park Campus, Ascot SL5

7PY, UK

[†]Corresponding author. E-mail: m.russell17@imperial.ac.uk

Abstract

During range expansion, invasive species can experience new thermal regimes. Differences

between the thermal performance of local and invasive species can alter species interactions,

including predator-prey interactions. The Asian tiger mosquito, *Aedes albopictus*, is a known

vector of several viral diseases of public health importance. It has successfully invaded many

regions across the globe and currently threatens to invade regions of the UK where conditions

would support seasonal activity. We assessed the functional response and predation

efficiency (percentage of prey consumed) of the cyclopoid copepods *Macrocyclops albidus*

and *Megacyclops viridis* from South East England, UK against newly-hatched French *Ae.*

albopictus larvae across a relevant temperature range (15, 20, and 25°C). Predator-absent

controls were included in all experiments to account for background prey mortality. We

found that both *M. albidus* and *M. viridis* display type II functional response curves, and that

both would therefore be suitable biocontrol agents in the event of an *Ae. albopictus* invasion

in the UK. No significant effect of temperature on the predation interaction was detected by

either type of analysis. However, the predation efficiency analysis did show differences due

to predator species. The results suggest that *M. viridis* would be a superior predator against

invasive *Ae. albopictus* larvae due to the larger size of this copepod species, relative to *M.*

albidus. Our work highlights the importance of size relationships in predicting interactions

between invading prey and local predators.

Keywords

biocontrol; cyclopoid copepods; functional response; invasion; predation efficiency; public

health; range expansion; vector-borne.

Introduction

An invasion of the Asian tiger mosquito, *Aedes albopictus*, into the UK is a major public

health concern. This species is not only an aggressive nuisance biter, but also a known vector

of arboviruses such as dengue, chikungunya, yellow fever, and Zika (Centers for Disease and

Prevention, 2017). The ability of *Ae. albopictus* mosquitoes to lay desiccation-resistant eggs

has enabled their introduction into cities across the globe, often via used tire shipments (Eritja

et al., 2005, Juliano and Lounibos, 2005, Lounibos, 2002, Medlock et al., 2006, Benedict et

al., 2007). According to the European Centre for Disease Prevention and Control, *Ae.*

albopictus populations have already established throughout the vast majority of Italy,

southern France, and as far north as the French department of Aisne (European Centre for

Disease and Control, 2019). Although this species has not yet established in the UK, it has

been introduced in Kent, a coastal county in the southeast of the UK, where 37 *Ae. albopictus*

eggs were found in September of 2016 (European Centre for Disease and Control, 2019,

Medlock et al., 2017).

Approximately one decade prior to the first recorded introduction of *Ae. albopictus* into the

UK, a model based on abiotic factors including photoperiod, rainfall, and temperature

suggested that upon introduction, *Ae. albopictus* adults would be active from May to

September in areas near London and the southern coastal ports (Medlock et al., 2006). A

more recent study that also considered diurnal temperature range and human population

density found that the most suitable areas for *Ae. albopictus* in the UK currently are centered

around London (Metelmann et al., 2019). Owing to the warming climate, the UK is expected

to become increasingly suitable for *Ae. albopictus* establishment; projections of climate and

human population conditions into future decades suggest that if introduced, the vector species

could establish throughout most of England and southern Wales during the 2060s

(Metelmann et al., 2019). Another recent model predicts that the UK will report *Ae.*

albopictus presence by either 2050 or 2080, depending on the patterns of future greenhouse

gas emissions (Kraemer et al., 2019).

Predation by copepods has been identified as a biological method for controlling *Ae.*

albopictus in Europe following the success of past field trials (Baldacchino et al., 2015). The

cyclopoid copepod *Macrocyclops albidus* has previously been used to eliminate *Ae.*

albopictus populations from tire piles in New Orleans, Louisiana, USA (Marten, 1990a). In

1994, *Mesocyclops aspericornis* were distributed into wells in Charters Towers, Queensland,

Australia to eliminate *Ae. aegypti* populations (Russell et al., 1996). In addition, *Mesocyclops*

copepods were used to effectively control populations of *Ae. aegypti* and *Ae. albopictus* in six

communes in northern Vietnam (Kay et al., 2002). A semi-field study conducted in Bologna,

Italy in 2007 found that *M. albidus* reduced the density of *Ae. albopictus* in experimental

drums by greater than 99% (Veronesi et al., 2015). In the UK, laboratory experiments using

cyclopoid copepods from Northern Ireland against *Culex pipiens* mosquito larvae from

Surrey, UK and *Ae. albopictus* larvae from Montpellier, France have supported the use of

copepods as biocontrol agents (Cuthbert et al., 2019b, Cuthbert et al., 2018b). Previous work

indicates that the attack rates of *M. albidus* and *Megacyclops viridis* predators against *Cx.*

pipiens prey tend to increase with temperature (Cuthbert et al., 2018b). However, the use of

UK copepods as predators of invasive *Ae. albopictus* larvae has not been thoroughly

examined over the range of temperatures that the invasive larvae are predicted to experience.

Due to the potentially negative impacts of exporting copepods to non-native regions for

biocontrol purposes (Coelho and Henry, 2017), it is important to investigate the performance

of copepods local to the predicted sites of *Ae. albopictus* establishment: London and South

East England.

Populations of *M. albidus* and *M. viridis* cyclopoid copepods from the benthos of the

Cumbrian lakes in the UK have previously been studied at temperatures from 5 to 20°C

(Laybourn-Parry et al., 1988). Based on the dry body masses of adult specimens, males were

consistently smaller than females, and *M. albidus* copepods were consistently smaller than *M.*

viridis (Laybourn-Parry et al., 1988). The general adult body length ranges for *M. albidus* and

M. viridis are 1.3 – 2.5 mm (Einsle, 1993) and 1.2 – 3 mm (Dussart, 1969, Einsle, 1988,

Kiefer, 1960), respectively. These are small enough to enable the distribution of copepod

cultures to mosquito larval habitats using "a simple backpack sprayer with a 5 mm hole in the

nozzle," as has been previously recommended (Marten, 1990b). Copepods reproduce

sexually, and females that can produce new egg sacs every 3-6 days tend to predominate in

mature populations (Marten and Reid, 2007). Cyclopoid copepods are considered sit-and-wait

ambush predators because of their attack behaviors, which have been described in six steps:

encounter, aiming, stalking, attack, capture, and ingestion (Awasthi et al., 2012). When

preying on mosquito larvae, copepods often avoid ingesting the head and thorax (Awasthi et

al., 2012).

Functional response curves were originally developed to relate the number of prey attacked

by an invertebrate predator to the prey density (Holling, 1959, Holling, 1966). Previous work

has suggested that the best predators to use in an "inundative release" biocontrol program are

those that have type II functional responses to prey density, so that the predators' attack rates

are high even at low prey densities (Daane et al., 1996). Predators that display a type III

functional response against an invasive prey species —such as signal crayfish (*Pacifastacus*

leniusculus) against New Zealand mud snails (*Potamopyrgus antipodarum*)— may be able to

limit the spread of the invader, but cannot prevent it from establishing (Twardochleb et al.,

2012). Cyclopoid copepods from the UK have previously exhibited type II functional

response curves when provided with *Ae. albopictus* prey at a single temperature setting

(Cuthbert et al., 2019b).

A meta-analysis of fifty functional response curves of cyclopoid copepod predators showed a

"monotonically increasing effect of temperature" on the attack rate parameter (Kalinoski and

DeLong, 2016). In addition, gut clearance rate, another measure of foraging rate, has been

demonstrated to increase linearly with temperature in planktonic copepods (Dam and

Peterson, 1988, Irigoien, 1998). However, five data points that were excluded from the linear

regression analysis and described as "distinctly separate" would have suggested a decrease in

gut clearance rate at higher temperatures (Irigoien, 1998). Furthermore, recent literature

shows that the temperature dependence of attack rate is unimodal, or "hump-shaped,"

especially when the temperature range exceeds thermal optima, and suggests that the impact

of climate change on predation will depend on the thermal optima of consumers (Englund et

al., 2011, Uiterwaal and Delong, 2020).

If UK cyclopoid copepods are used as biocontrol agents against *Ae. albopictus*, they are

likely to experience temperatures higher than those of their natural habitats not only because

of climate change, but also because the predation interaction would occur mostly in artificial

containers, where low volumes of water can warm more quickly, relative to the warming

expected in lakes and ponds. The usual environmental temperatures experienced by *M.*

albidus and *M. viridis*, 5 to 20°C (Laybourn-Parry et al., 1988), are expected to influence

their thermal optima (Dell et al., 2011). An analysis of 2,083 functional responses

demonstrated that the thermal optima of attack rate and handling time parameters often fall

within the range of 15-25°C (Uiterwaal and Delong, 2020). Among 69 unimodal thermal

responses of freshwater organisms, such as cyclopoid copepods, a mean optimal temperature

of 21°C was observed (Dell et al., 2011). However, the thermal optimum of the prey species

can also influence the temperature dependence of predation interactions. The optimal

temperature for French *Ae. albopictus* is likely to be higher than that of UK copepods and has

been recorded at 29.7°C for immature *Ae. albopictus* originating from La Réunion, an island

in the southwest region of the Indian Ocean (Delatte et al., 2009). Such asymmetries in the

thermal response of consumer and prey species may modify the effect of temperature on the

predation interaction (Dell et al., 2014, Grigaltchik et al., 2012).

In this study, we determined the type of functional response of *M. albidus* and *M. viridis*

copepods from Surrey, UK against *Ae. albopictus* larvae from Montpellier, France at three

different temperatures (15, 20, and 25°C) likely to be experienced by container-dwelling

mosquito larvae in London and South East England. We also calculated predation efficiency,

the percentage of larvae consumed, of both copepod species at a constant mosquito prey

density under the same three temperatures.

Materials and methods

Collection of field temperature data:

Previous work has suggested that the microclimates experienced by *Ae. albopictus* are not

represented well by climate data reported from local weather stations (Murdock et al., 2017).

Since used tire shipments are a common source of *Ae. albopictus* introductions (Eritja et al.,

2005, Juliano and Lounibos, 2005, Lounibos, 2002, Medlock et al., 2006, Benedict et al.,

2007), we measured the temperature of rainwater collected in used car tires that were divided

between an urban London site and a suburban site. These temperature data provided rough

guidelines for our choice of the three temperatures to be tested in our functional response and

predation efficiency experiments. The minimum water temperature was 9°C, the 25th

percentile was 16.1°C, the median was 18.1°C, the 75th percentile was 20°C, and the

maximum was 25°C. Thus, our choice of 15, 20, and 25°C as the three main temperatures of

interest is representative of more than 75% of the tire water temperatures recorded in the field

from May through September of 2018 (Supporting Information).

Local copepod cultures:

Adult gravid female copepods were collected in August of 2018 from the edge of Longside

Lake in Egham, Surrey, UK (N 51° 24.298', W 0° 32.599') using a 53 µm sieve (Reefphyto

Ltd, UK). Separate cultures were started from each gravid female. The copepods were kept in

3 L containers of spring water (Highland Spring, UK) at a 12:12 light/dark cycle, and 20 ±

1°C, a temperature previously found to increase the portion of their life cycle spent in the

reproductive phase (Laybourn-Parry et al., 1988). *Chilomonas paramecium* and *Paramecium*

caudatum (Sciento, UK) were provided *ad libitum* as food for the copepods (Suarez et al.,

1992). The ciliates were cultured in 2 L flasks containing boiled wheat seeds; boiled wheat

seeds were also added to the copepod containers (Suarez et al., 1992). Adult copepods were

identified as *Macrocyclops albidus* (Jurine, 1820) and *Megacyclops viridis* (Jurine, 1820) by

Dr. Maria Hołyńska from the Museum and Institute of Zoology in Warsaw, Poland.

Temperate *Ae. albopictus* colony care:

A colony of *Ae. albopictus* mosquitoes (original collection Montpellier, France 2016 obtained

through Infravec2) was maintained at $27 \pm 1°C$, 70% relative humidity, and a 12:12 light/dark

cycle. The colony maintenance temperature of 27°C was chosen because it falls at the upper

bound of the 95% confidence interval that was fitted around mean daily Montpellier

temperatures from 1996-2005 in the summer months, and these temperatures are expected to

increase in the coming decades (Zidon et al., 2016). Larvae were fed fish food (Cichlid Gold

Hikari®, Japan), and adults were given 10% sucrose solution and horse blood (First Link Ltd,

UK) administered through a membrane feeding system (Hemotek®, Blackburn, UK).

Design of functional response experiments:

Adult non-gravid female copepods, identified by larger relative size, were removed from their

culture and each was placed in a Petri dish (diameter: 50 mm, height: 20.3 mm) holding 20

mL of spring water. At approximately 11am, the copepods were placed in three different

controlled environments set to 15 ± 1, 20 ± 1, and $25 \pm 1°C$, all at a 12:12 light/dark cycle, to

begin a 24 h starvation and temperature-acclimation period for the predators (Fig. 1). Each

combination of copepod species (*M. albidus* or *M. viridis*) and temperature had 28 copepods,

each one held in its own Petri dish; there were 168 copepods in total (Fig. S3a). Ten

additional gravid females were randomly selected from each species of copepod and

preserved in 80% ethanol for size measurements.

At noon, groups of 87 newly-hatched *Ae. albopictus* larvae (Supporting Information) were

each pipetted into a small plastic tub containing 150 mL of spring water to strongly dilute any

residual food from the hatching media. Each tub of 87 larvae was then split into seven Petri

dishes each containing 20 mL of spring water and the following larval densities: 1, 2, 4, 8, 16,

24, and 32. All larvae were counted into dishes at room temperature and then divided into the

three different controlled environments, allowing a 12 h temperature acclimation period

before predator introduction (Fig. 1). For each combination of copepod species and

temperature, there were 35 Petri dishes containing a total of 435 newly-hatched *Ae.*

albopictus larvae, with a total of 2,610 larvae used across all predator treatments and controls

(Fig. S3a). This sample size allowed for four replicates of each larval density and one

predator-absent control at each density, for each combination of copepod species and

temperature (Fig. S3a).

Fig. 1. General schedule of both functional response and predation efficiency experiments

At approximately 11am the next day, the copepod predators held at three different

temperatures were introduced to larval Petri dishes of matching controlled temperatures (Fig.

1). The copepods were removed after a 6 h period of predation (Fig. 1), which follows the

schedule of similar experiments (Cuthbert et al., 2018a, Cuthbert et al., 2019a). Each copepod

was preserved in 80% ethanol so that its body size could later be measured and matched to

the larval count data. Immediately following the removal of the copepods, the number of

surviving larvae in each Petri dish was counted and recorded. In addition, the body lengths of

the 10 gravid females selected from each species of copepod, as well as the body lengths of

all copepods included as predators in the functional response experiments, were measured

from the front of the cephalosome to the end of the last urosomite (Dussart and Defaye,

2001).

Design of predation efficiency experiment:

In addition to functional response, predation efficiency has previously been measured to

assess copepod predators of mosquito larvae (Baldacchino et al., 2017). We measured

predation efficiency using a constant density of 24 larvae in 20 mL of spring water. This

density was chosen from the higher end of the range of densities used in the functional

response experiments, where the curves generally start to plateau (Juliano, 2001) (Fig. 2a).

Ae. albopictus larvae were hatched (Supporting Information) and any residual food was

diluted in spring water following the same procedure used for the functional response

experiments. Adult non-gravid female copepods were each placed in a Petri dish at

approximately 11am for a 24 h period of starvation and acclimation to three different

temperature settings: 15 ± 1, 20 ± 1, and $25 \pm 1°C$, all at a 12:12 light/dark cycle (Fig. 1). The

largest non-gravid copepods of each species were selected to minimize the risk of selecting

males or immature stages. The Petri dishes containing larvae were split into the three

different temperature settings 12 h prior to the introduction of copepod predators, and there

was a 6 h period of predation between 11am and 5pm (Fig. 1). At the end of the 6 h period,

the copepods were removed, and each was stored in 80% ethanol and labelled according to its

larval Petri dish. The number of surviving larvae in each Petri dish was recorded immediately

after removing the copepods.

At each of the three temperature settings, there was a total of 24 Petri dishes, each containing

24 larvae (1,728 larvae across all temperatures); the 24 dishes were divided into three groups

(n = 8): one with *M. albidus* predation, one with *M. viridis* predation, and one as a control

(Fig. S3b). Every predator treatment was matched to a control that had been held at the same

temperature (Fig. S3b). Each temperature setting had a total of 192 control larvae. Larval

background mortality was 5% at 15°C, 11% at 20°C, and 3% at 25°C. A total of 24 *M.*

albidus and 24 *M. viridis* copepods were used in this experiment. The body lengths of these

copepods were measured after each was preserved in 80% ethanol.

Functional response curve analysis:

The general shapes of the functional response curves were determined for each combination

of copepod species and temperature using predator-present data. When polynomial logistic

functions are fitted to describe the relationship between the proportion of larvae killed and the

initial larval density, a negative first-order term indicates a type II response, and a positive

first-order term indicates a type III response (Juliano, 2001, Pritchard et al., 2017b). The

"frair_test" function ("frair" package, R version 3.4.2) was used to determine the shapes of

the functional response curves according to the sign and significance of first-order and

second-order terms in logistic regressions (Pritchard et al., 2017a, Pritchard et al., 2017b).

Three observations were excluded from the *M. albidus* at 25°C group — two due to copepod

deaths during the starvation period, and one because two copepod predators were introduced

into the same Petri dish. Three observations were excluded from the *M. viridis* at 25°C group,

and one was excluded from the *M. viridis* at 15°C group; all four *M. viridis* exclusions were

due to insufficient body length of the copepod indicating that these predators could have been

either males or immatures, rather than adult non-gravid females.

Mortality of *Ae. albopictus* larvae was observed in controls for five out of the six

combinations of copepod species and temperature. *M. viridis* predators at 25°C was the only

group without larval mortality in any of the controls. For *M. albidus*, across all control larvae

(n=87) at 15°C, three deaths were observed (3% mortality); four deaths were observed at

20°C (5% mortality); and eleven deaths were observed at 25°C (13% mortality). For *M.*

viridis, two deaths were observed among control larvae at 15°C (2% mortality), and three

deaths were observed at 20°C (3% mortality). Mortality rates of aedine larvae in predator-

absent controls have previously been observed to be as high as 20% (Baldacchino et al.,

2017). To account for both background mortality and prey depletion caused by predation, the

functional response parameters (attack coefficient and handling time) were estimated by

fitting the following differential equation:

$$\frac{dN}{dt} = -\frac{bN^{1+q}}{1 + bhN^{1+q}} P - mN \qquad \text{eqn 1}$$

where *dN/dt* is the change in prey density over time, *b* is the attack coefficient, *q* is an

exponent that can fall between a type II response (q = 0) and a type III response (q = 1), *h* is

the handling time, *P* is the predator density, and *m* is the mortality rate (Rosenbaum and Rall,

2018).

Since background prey mortality should not alter the general shape of the functional response

curve (Rosenbaum and Rall, 2018), the q parameter was fixed to either 0 or 1 based on the

results of the "frair_test" function, which only considers data from predator-present

treatments (Pritchard et al., 2017a, Pritchard et al., 2017b). The attack coefficient (b) was

given a starting value of one, the handling time (h) had a starting value equivalent to the

inverse of the maximum feeding rate, and the mortality rate (m), if applicable, was given a

starting value of 0.01 (Rosenbaum and Rall, 2018). The attack coefficient and handling time

parameters were estimated based on our empirical data using an iterative maximum

likelihood method: "mle2" function, "bbmle" package, R version 3.4.2 (Bolker et al., 2020,

Bolker, 2008, Rosenbaum and Rall, 2018).

The "nll.ode.general.mort" function (Rosenbaum and Rall, 2018) was used to fit data that

included some background prey mortality to the statistical model (eqn1). In cases where the

mortality rate was shown to be insignificant, or where no larval mortality was originally

observed, the "nll.bolker" function (Rosenbaum and Rall, 2018), which assumes a mortality

rate of zero, was used. Both functions are negative log-likelihood functions that are

minimized by applying maximum likelihood estimation methods (Rosenbaum and Rall,

2018). Confidence intervals for the model parameters of interest, attack coefficient and

handling time, were computed by the "confint" function, base package, R version 3.4.2. To

detect differences in functional response parameter estimates due to either temperature

category or copepod species, 95% confidence intervals were compared; this method for

comparing parameter estimates across temperatures has previously been used in a similar

analysis of copepod predation (Novich et al., 2014).

Predation efficiency analysis:

For each pair of predator treatment and control (Fig. S3b), the copepod's predation efficiency

was calculated according to Abbott's formula (Abbott, 1987, Baldacchino et al., 2017):

Predation efficiency $= \frac{\text{Number alive in control} - \text{Number alive in treatment}}{\text{Number alive in control}}$ (100) eqn 2

Copepod body lengths, measured in mm, were converted to estimates of body mass (mg)

using an equation from previous studies (Alcaraz and Strickler, 1988, Klekowski and

Shushkina, 1966, Novich et al., 2014):

Mass $= 0.055$ x length$^{2.73}$ eqn 3

Two linear regression models were fitted to explain predation efficiency. The first tested

temperature and copepod species as predictors (eqn S1), and the second tested copepod body

mass in place of species (eqn S2). Both models were also fitted without temperature

predictors (Supporting Information).

Body mass difference between copepod species by experimental design:

Among the predators used in the functional response experiments, each *M. albidus* copepod

was randomly paired with an *M. viridis* copepod (80 pairs). Predation efficiency predators

were also randomly paired, yielding 23 pairs of the two species. For each pair, the *M. albidus*

body mass was subtracted from that of *M. viridis*, generating 103 differences in body mass

due to species. A Wilcoxon rank sum test was used to determine if the body mass differences

between copepod species varied by experimental design.

Analyses were completed in R version 3.4.2, and all data will be made accessible from the

Dryad Digital Repository.

Results

Functional response curves:

A type II functional response, demonstrated through logistic regression results (Table S1),

was observed in every combination of copepod species and temperature (Fig. 2a). There was

a significant effect of background mortality on the change in prey density over time for *M.*

albidus at 20 and 25°C (Table S2). Within each copepod species, there was no significant

difference in either attack coefficient or handling time due to temperature (Fig. 2b & 2c,

Table S4). Within each temperature setting, there was no significant difference in either type

of parameter estimate due to copepod species (Fig. 2b & 2c, Table S4). The 95% confidence

interval around the attack coefficient estimate for *M. albidus* at 25°C was extremely wide,

representing high uncertainty (Fig. 2b). This imprecise estimate was likely due to the highly

significant effect of mortality rate (p-value = 0.0040) on the change in prey density over time

(Table S2). In addition, the mortality rate's standard error was higher for *M. albidus* at 25°C

than it was for any other data subset that included background larval mortality (Table S2),

and the model deviance (-2 log L = 111.25) was higher than that of any other model used to

generate parameter estimates (Tables S2 & S3).

Fig. 2a.) Type II functional response curves for each combination of copepod species and temperature (Points are shown in the "jitter" position.) **b.)** Attack coefficient estimates and 95% confidence intervals by copepod species and temperature **c.)** Handling time estimates and 95% confidence intervals by copepod species and temperature

Predation efficiency:

Predation efficiency values ranged from 5.0% to 58.3%, with a median of 21.7%, and a slightly right-skewed distribution (skewness = 0.61, kurtosis = 2.9).

Table 1. Linear regression of predation efficiency by copepod species and temperature (n = 47)

Parameter	Estimate	Standard Error	p-value	Adjusted R^2	AIC
Intercept	18.19	3.30	1.92×10^{-6}	0.068	373.1
Species (*M. viridis*, ref = *M. albidus*)	7.54	3.33	0.0288		
Temperature (20°C, ref = 15°C)	0.80	4.04	0.8432		
Temperature (25°C, ref = 15°C)	4.56	4.10	0.2731		

A linear model testing copepod species and temperature category as predictors of predation efficiency explained 6.8% of the variance in predation efficiency (Table 1). The predation efficiency of *M. viridis* copepods was 25.7%, significantly higher (p-value = 0.0288) than that of *M. albidus* copepods, which had a predation efficiency of 18.2%, when controlling for differences in temperature (Table 1, Fig. 3a). Interactions between species and temperature were not significant (Species x Temperature$_{20}$, p-value = 0.150; Species x Temperature$_{25}$, p-value = 0.466). Removing the temperature predictors from the model increased the adjusted R^2 value and decreased the AIC (Tables 3 & S5).

412 **Table 2.** Linear regression of predation efficiency by copepod body mass and temperature (n

413 = 47)

Parameter	Estimate	Standard Error	p-value	Adjusted R^2	AIC
Intercept	9.72	4.88	0.0528	0.140	369.3
Body Mass (mg)	34.44	11.37	0.0042		
Temperature (20°C, ref = 15°C)	0.83	3.88	0.8305		
Temperature (25°C, ref = 15°C)	4.04	3.94	0.3106		

415 **Table 3.** Model selection

Parameters	Degrees of Freedom	Adjusted R^2	AIC[a]
Species, Temperature$_{20}$, Temperature$_{25}$	43	0.068	373.1
Species, Temperature$_{20}$, Temperature$_{25}$, Species x Temperature$_{20}$, Species x Temperature$_{25}$	41	0.071	390.7
Species	45	0.081	362.6
Mass, Temperature$_{20}$, Temperature$_{25}$	43	0.140	369.3
Mass, Temperature$_{20}$, Temperature$_{25}$, Mass x Temperature$_{20}$, Mass x Temperature$_{25}$	41	0.107	388.8
Mass	45	0.156	358.5

416   a.) A difference in AIC value greater than or equal to 2 is the suggested minimum

417   difference for model selection purposes (Burnham and Anderson, 2002).

419 A linear model testing copepod body mass and temperature category as predictors of

420 predation efficiency explained 14.0% of the variance in predation efficiency (Table 2). For

each 0.1 mg increase in copepod body mass, the predation efficiency increased by

approximately 3.4 percentage points (p-value = 0.0042), when controlling for differences in

temperature (Table 2, Fig. 3b). Interactions between body mass and temperature were not

significant (Mass x Temperature$_{20}$, p-value = 0.611; Mass x Temperature$_{25}$, p-value = 0.565).

Removing the temperature predictors from the model improved model fit (Tables 3 & S6).

Fig. 3a.) Bar chart of predation efficiency by copepod species and temperature (Error bars

represent ± the standard error.) **b.)** Predation efficiency by copepod body mass (Points are

shown in the "jitter" position.)

Body mass differences between copepod species by experimental design:

M. viridis copepods were larger than *M. albidus* copepods in both the functional response and

the predation efficiency experiments, and both copepod species were larger in the predation

efficiency experiments than in the functional response experiments (Supporting Information,

Fig. S4).

The body mass difference between copepod species was greater in the predation efficiency

experiments than in the functional response experiments (p-value = 0.0136). In the predation

efficiency experiments, the median increase from *M. albidus* to *M. viridis* was 0.24 mg, while

in the functional response experiments, the median increase was 0.12 mg.

Discussion

Both *M. albidus* and *M. viridis* copepods collected from Surrey, UK would be appropriate for

use in an inundative release biocontrol program because both predator species display type II

functional response curves when presented with invasive *Ae. albopictus* larvae (Table S1).

Type II functional response curves are preferred because they demonstrate the predator's

ability to eliminate invasive populations at low prey densities (Daane et al., 1996). Other

studies using *M. albidus* and *M. viridis* from Northern Ireland against mosquito larvae prey

have also found evidence of type II functional response curves (Cuthbert et al., 2019b,

Cuthbert et al., 2018b).

Our functional response analyses did not detect any differences in response type, attack

coefficient, or handling time due to copepod species or temperature, tested at 15, 20, and

25°C (Fig. 2, Table S1). In addition, we found that temperature categorical variables were not

significant predictors of copepod predation efficiency against *Ae. albopictus* larvae (Tables 1-

3). A meta-analysis of 648 functional responses from 86 different studies found that although

attack rate increased with temperature, the increase was less steep than that which had been

expected based on metabolic theory (Rall et al., 2012). In addition, a previous study of

predator-prey interactions involving sit-and-wait copepod predators found no significant

difference in the attack parameter of the functional response across three different

temperatures: 18, 22, and 26°C (Novich et al., 2014). Since the effect of temperature on sit-

and-wait predation is driven mainly by the velocity of the prey (Dell et al., 2014), our results

suggest that the velocity of newly-hatched *Ae. albopictus* larvae does not change dramatically

over a temperature range of 15-25°C. This is consistent with previous observations of high

cold tolerance in aedine larvae. A recent study showed that *Ae. aegypti* larvae from a colony

maintained at $22 \pm 1°C$ continued to move in response to probing as the temperature was

gradually lowered; the larvae did not stop responding with movement until the temperature

reached 6°C, 16°C lower than the temperature at which their colony was maintained (Jass et

al., 2019). From a public health perspective, the absence of significantly lower predation

efficiency at lower temperatures is useful because *Ae. albopictus* larvae that develop at lower

temperatures have been shown to be more susceptible to chikungunya infection as adults

(Westbrook et al., 2010).

The term "handling time" has been defined differently by different studies. For example, one

study defined it as "the time lapsed between an attack of the prey by the copepod and its

resumption of movement after ingestion," and found that *M. thermocyclopoides* copepods

displayed an average handling time of 83 s for first instar *Anopheles stephensi* larvae and 185

s for first instar *Cx. quinquefasciatus* (Kumar and Rao, 2003). These times are consistent with

directly-observed ingestion times reported by another study, which states that a mosquito

larva is usually consumed by a cyclopoid copepod within a few minutes (Marten and Reid,

2007). However, the original methods designed for fitting functional response curves assume

that the predator only has "two time-consuming behaviours – searching and handling of prey"

(Holling, 1959). In this context, the "handling time" has been defined as "the time lost from

searching per resource consumed," not as the ingestion time per resource consumed

(Uiterwaal and Delong, 2020). Our handling time estimates (Fig. 2c, Table S4) are based on

the original framework of functional response analysis, and therefore represent average

predator search times per larva consumed. One study found that when directly-observed

handling times were adjusted to include both the time that the predator takes to move to the

prey, and the ingestion time, there was no significant difference between directly-observed

handling times and handling times estimated by fitting functional response curves (Poole et

al., 2007). The prey-searching techniques of *M. albidus* and *M. viridis* have previously been

described as "[relying] on bumping into … prey during the course of … meanderings" (Fryer,

1957a). This description of rather inefficient prey searching is consistent with our handling

time estimates of approximately 30 min to 1 h (Fig. 2c, Table S4).

We found that *M. viridis* was a significantly more efficient predator of *Ae. albopictus* than *M.*

albidus (Tables 1 & S5, Fig. 3). An analysis of the gut contents of English Lake District

copepods previously found that 15.7% of *M. viridis* had consumed dipterous larvae,

compared to only 7.4% of *M. albidus* (Fryer, 1957b). Thus, it is possible that English *M.*

viridis may be better adapted to feeding on mosquito larvae than English *M. albidus*. A study

using *M. albidus* and *M. viridis* from Northern Ireland against *Culex* mosquitoes from Surrey,

UK suggested that *M. albidus* is more effective than *M. viridis* as a biocontrol (Cuthbert et

al., 2018b). It appears this is not the case for invasive *Ae. albopictus* prey. A previous study

showed that an aedine species (*Ae. aegypti*) was more vulnerable to predation by *M. viridis*

than *Cx. pipiens*, despite the larger head capsule width of *Ae. aegypti* (Blaustein and Margalit,

1994). In addition, *M. viridis* was among three cyclopoid copepod species from Northern

Ireland that displayed a preference for *Ae. albopictus* larvae over *Cx. pipiens* prey (Cuthbert

et al., 2019b). Higher predation efficiency against *Aedes* than against *Culex* may be due to

differences in larval anatomy and activity patterns. Previous work has attributed higher

survival of *Culex* larvae to lower levels of motility, and to their long bristles, or spines, that

either prevent copepod attacks from being successful, or deter attacks entirely by creating the

illusion of larger size (Dieng et al., 2003, Marten and Reid, 2007, Soumare and Cilek, 2011).

Another study of *M. albidus* and *M. viridis* from Northern Ireland as predators of both

Paramecium caudatum and *Cx. pipiens* reported that *M. viridis* consumed more prey

organisms overall than *M. albidus* (Cuthbert et al., 2019a).

The larger size of *M. viridis* contributes to the higher predation efficiency observed among

that species against *Ae. albopictus* (Fig. 3b). Our analysis shows that copepod body mass was

a better predictor of predation efficiency than species identity (Tables 1-3). The linear

regression model with the highest adjusted R-squared value, 16%, was the model with

copepod body mass as the sole predictor of predation efficiency (Tables 3 & S6). This

adjusted R-squared value is still low, indicating substantial variation within the predation

interaction. The positive relationship we observed between copepod body mass and predation

efficiency (Tables 2 & S6, Fig. 3b) is consistent with the findings of a previous meta-analysis

on crustacean predation of immature fish, which showed that the predation rate was

negatively related to the prey/predator size ratio (Paradis et al., 1996). Our data are also in

agreement with the positive linear relationship between body mass and ingestion rate that has

been demonstrated in other taxa, including herbivorous and carnivorous endotherms, as well

as carnivorous poikilothermic tetrapods (Peters, 1983). Although a recent study using *Ae.*

japonicus larvae as prey found that *M. viridis* predation rates began to decrease when

predators exceeded 1.8 mm in length (Früh et al., 2019), our data represent copepods of up to

2.5 mm in length, and there is no evidence of lower predation efficiency among the largest

predators (Fig. 3b). According to the theory that, at equilibrium, "the rate of food intake is

equal to the rate at which food is leaving the gut" (Charnov and Orians, 1973), our results

suggest that *M. viridis* may have a higher gut clearance rate than *M. albidus*, possibly due to

its larger size.

Despite the higher predation efficiency observed among *M. viridis* (Tables 1 & S5, Fig. 3),

we did not detect any differences between the two copepod species in their functional

response parameter estimates, when controlling for temperature (Fig. 2b & 2c, Table S4); this

was perhaps because the predation efficiency experiment had a larger sample size of

observations for each combination of predictors, and therefore more power to detect

differences. In addition, while the size difference between *M. albidus* and *M. viridis* was

significant in both experimental designs (Supporting Information, Fig. S4), there was a

greater difference in size due to species among the copepods used in the predation efficiency

experiment than among the copepods used in the functional response experiment.

Although we did not detect any difference in functional response parameter estimates or

predation efficiency due to temperature, the temperature at which copepods are cultured

impacts their size (Laybourn-Parry et al., 1988), and our results show that copepod size is a

determinant of their predation efficiency. A previous study, in which UK *M. albidus* and *M.*

viridis were cultured at six different temperatures between 5 and 20°C, showed a consistent

negative relationship between adult body mass of both species and culturing temperature

(Laybourn-Parry et al., 1988). Evidence of this relationship between size and rearing

temperature had already been recorded for freshwater copepods collected in Paris (Coker,

1933). Although cultures tend to have higher rates of reproduction at higher temperatures

(Laybourn-Parry et al., 1988), we recommend that the importance of individual body mass be

taken into consideration when determining the optimal temperature for mass copepod

culturing.

Fecundity data from older studies (Laybourn-Parry et al., 1988, Maier, 1994) have recently

been incorporated into metrics used to compare the efficacy of different copepod species as

biocontrol agents (Cuthbert et al., 2019b, Cuthbert et al., 2018b). Caution is recommended

when interpreting these metrics if the fecundity variable is either based on an inadequate

number of observations or derived from multiple populations of copepods. For example, a

previous calculation used to determine the "reproductive effort" of *M. viridis* incorporated a

clutch weight value based on the dimensions of a single egg sac from a female collected at an

unrecorded temperature in South Germany, female body weight data from a population in

South Germany, and embryonic development time data from English Lake District copepods

held at 20°C (Abdullahi and Laybourn-Parry, 1985, Maier, 1994). This reproductive effort

value was then applied to calculate a metric for *M. viridis* from Northern Ireland that were

cultured at 25°C (Cuthbert et al., 2019b). While it is time-consuming to collect robust and

relevant fecundity data experimentally, biocontrol metrics can only be meaningful if they are

based on appropriate original data.

Life history characteristics other than fecundity, particularly lifespan and resistance to

starvation, also impact the efficiency of biocontrol agents, and these characteristics have been

shown to positively correlate with cyclopoid copepod body size (Dieng et al., 2003, Maier,

1994). The relationship between lifespan and body size is recognized as a general rule in

ecology and has been exemplified by comparing the lifespan of an elephant to that of a mouse

(Brown et al., 2004). *M. viridis* females from English Lake District cultures have previously

been observed to spend approximately 120 days (at 15°C) and 80 days (at 20°C) in the adult

life stage (Laybourn-Parry et al., 1988). The fecundity and long-term stability of field

copepod populations can be difficult to predict. A year-round field study conducted in

Florida, USA of cyclopoid copepod populations in tire piles found that the number of

copepods observed was influenced by the age and amount of leaf litter in each tire (Schreiber

et al., 1996). In addition, cyclopoid copepod populations survive longer in tires near shade or

vegetation because the water in the tire is less likely to evaporate (Marten et al., 1994). In the

UK, where *Ae. albopictus* adults are only expected to be active from May to September

(Medlock et al., 2006), the stability of year-round copepod populations in tires may not be a

priority. The use of copepods as biocontrol agents in UK field sites might benefit from bi-

monthly re-applications of well-maintained laboratory cultures, and perhaps more frequent

reinforcements in case of long droughts followed by substantial precipitation. A control

strategy that is based on precipitation patterns and the estimated survival time of adult

copepods in the field would be less risky than a strategy that relies on the viability of

subsequent copepod generations.

As the invasion of *Ae. albopictus* into the UK progresses, it may become more relevant to

conduct predation experiments on mosquito larvae that are hatched from colonies acclimated

to lower temperatures. In our experiments, the *Ae. albopictus* larvae were from a colony that

was reared under conditions experienced on warm summer days in Montpellier, France. This

approach is currently valid because *Ae. albopictus* has not yet established in the UK and eggs

that could be introduced via used tire shipments would be naïve to UK weather conditions.

However, in future decades, it may be more appropriate to use *Ae. albopictus* larvae from a

colony reared under conditions reflective of the diurnal temperature range experienced in the

southern UK.

Author Contributions

LJC and MCR conceived the ideas and designed the methodology of the predation

experiments; AQ, LJC, and MCR collected the data; CGW provided critical advice on

cultivating the copepods; AQ improved the *Ae. albopictus* hatching procedure; MCR

analyzed the data and led the writing of the manuscript. All authors contributed critically to

the drafts and gave final approval for publication.

Acknowledgements

Many thanks to Flora Dickie, Dr. Maria Hołyńska, and Dr. Ross Cuthbert for their advice and

assistance. In addition, we express our gratitude to Dr. Samraat Pawar for his guidance on the

functional response literature. This project has received resources funded by the European

Union's Horizon 2020 research and innovation program under grant agreement No 731060

(Infravec2). The work was primarily funded by a President's PhD Scholarship from Imperial

College London awarded to Marie C. Russell.

Literature Cited

ABBOTT, W. S. 1987. A method of computing the effectiveness of an insecticide. 1925. *J Am Mosq*
Control Assoc, 3**,** 302-3.
ABDULLAHI, B. A. & LAYBOURN-PARRY, J. 1985. The effect of temperature on size and development
in three species of benthic copepod. *Oecologia,* 67**,** 295-297.
ALCARAZ, M. & STRICKLER, J. R. 1988. LOCOMOTION IN COPEPODS - PATTERN OF MOVEMENTS AND
ENERGETICS OF CYCLOPS. *Hydrobiologia,* 167**,** 409-414.
AWASTHI, A. K., WU, C. H., TSAI, K. H., KING, C. C. & HWANG, J. S. 2012. How Does the Ambush
Predatory Copepod Megacyclops formosanus (Harada, 1931) Capture Mosquito Larvae of
Aedes aegypti? *Zoological Studies,* 51**,** 927-936.
BALDACCHINO, F., BRUNO, M. C., VISENTIN, P., BLONDEL, K., ARNOLDI, D., HAUFFE, H. C. & RIZZOLI,
A. 2017. Predation efficiency of copepods against the new invasive mosquito species Aedes
koreicus (Diptera: Culicidae) in Italy. *European Zoological Journal,* 84**,** 43-48.
BALDACCHINO, F., CAPUTO, B., CHANDRE, F., DRAGO, A., DELLA TORRE, A., MONTARSI, F. & RIZZOLI,
A. 2015. Control methods against invasive Aedes mosquitoes in Europe: a review. *Pest*
management science, 71**,** 1471-1485.
BENEDICT, M. Q., LEVINE, R. S., HAWLEY, W. A. & LOUNIBOS, L. P. 2007. Spread of the tiger: Global
risk of invasion by the mosquito Aedes albopictus. *Vector-Borne and Zoonotic Diseases,* 7**,**
76-85.
BLAUSTEIN, L. & MARGALIT, J. 1994. DIFFERENTIAL VULNERABILITY AMONG MOSQUITO SPECIES TO
PREDATION BY THE CYCLOPOID COPEPOD, ACANTHOCYCLOPS-VIRIDIS. *Israel Journal of*
Zoology, 40**,** 55-60.

BOLKER, B., TEAM', R. D. C. & GINE-VAZQUEZ, I. 2020. *bbmle: Tools for general maximum liklihood*
estimation. R package version 1.0.23.1 [Online]. Available: https://CRAN.R-
project.org/package=bbmle [Accessed].
BOLKER, B. M. 2008. *Ecological models and data in R,* Princeton, NJ, Princeton University Press.
BROWN, J. H., GILLOOLY, J. F., ALLEN, A. P., SAVAGE, V. M. & WEST, G. B. 2004. Toward a metabolic
theory of ecology. *Ecology,* 85**,** 1771-1789.
BURNHAM, K. P. & ANDERSON, D. R. 2002. Formal Inference From More Than One Model:
Multimodel Inference (MMI). *Model Selection and Multimodel Inference: A Practical*
Information-Theoretic Approach. 2nd ed. New York: Springer.
CENTERS FOR DISEASE, C. & PREVENTION 2017. Surveillance and Control of Aedes aegypti and Aedes
albopictus in the United States.
CHARNOV, E. & ORIANS, G. 1973. *Optimal Foraging: Some Theoretical Explorations*.
COELHO, P. & HENRY, R. 2017. *Copepods Against Aedes Mosquitoes: A Very Risky Strategy*.
COKER, R. E. 1933. Influence of temperature on size of freshwater copepods (Cyclops). *International*
Review of Hydrobiology, 29**,** 406-436.
CUTHBERT, R. N., CALLAGHAN, A. & DICK, J. T. A. 2018a. Dye another day: the predatory impact of
cyclopoid copepods on larval mosquito Culex pipiens is unaffected by dyed environments.
Journal of Vector Ecology, 43**,** 334-336.
CUTHBERT, R. N., CALLAGHAN, A. & DICK, J. T. A. 2019a. The Effect of the Alternative Prey,
Paramecium caudatum (Peniculida: Parameciidae), on the Predation of Culex pipiens
(Diptera: Culicidae) by the Copepods Macrocyclops albidus and Megacyclops viridis
(Cyclopoida: Cyclopidae). *Journal of Medical Entomology,* 56**,** 276-279.
CUTHBERT, R. N., CALLAGHAN, A. & DICK, J. T. A. 2019b. A novel metric reveals biotic resistance
potential and informs predictions of invasion success. *Scientific Reports,* 9.
CUTHBERT, R. N., DICK, J. T. A., CALLAGHAN, A. & DICKEY, J. W. E. 2018b. Biological control agent
selection under environmental change using functional responses, abundances and
fecundities; the Relative Control Potential (RCP) metric. *Biological Control,* 121**,** 50-57.
DAANE, K. M., YOKOTA, G. Y., ZHENG, Y. & HAGEN, K. S. 1996. Inundative release of common green
lacewings (Neuroptera: Chrysopidae) to suppress Erythroneura variabilis and E-elegantula
(Homoptera: Cicadellidae) in vineyards. *Environmental Entomology,* 25**,** 1224-1234.
DAM, H. G. & PETERSON, W. T. 1988. THE EFFECT OF TEMPERATURE ON THE GUT CLEARANCE RATE-
CONSTANT OF PLANKTONIC COPEPODS. *Journal of Experimental Marine Biology and*
Ecology, 123**,** 1-14.
DELATTE, H., GIMONNEAU, G., TRIBOIRE, A. & FONTENILLE, D. 2009. Influence of Temperature on
Immature Development, Survival, Longevity, Fecundity, and Gonotrophic Cycles of Aedes
albopictus, Vector of Chikungunya and Dengue in the Indian Ocean. *Journal of medical*
entomology, 46**,** 33-41.
DELL, A. I., PAWAR, S. & SAVAGE, V. 2014. Temperature dependence of trophic interactions are
driven by asymmetry of species responses and foraging strategy. *Journal of Animal Ecology,*
83**,** 70-84.
DELL, A. I., PAWAR, S. & SAVAGE, V. M. 2011. Systematic variation in the temperature dependence
of physiological and ecological traits. *Proceedings of the National Academy of Sciences of the*
United States of America, 108**,** 10591-10596.
DIENG, H., BOOTS, M., TUNO, N., TSUDA, Y. & TAKAGI, M. 2003. Life history effects of prey choice by
copepods: Implications for biocontrol of vector mosquitoes. *Journal of the American*
Mosquito Control Association, 19**,** 67-73.
DUSSART, B. 1969. *Les copépodes des eaux continentales d'Europe Occidentale. 2: Cyclopoïdes et*
Biologie, Paris, Editions N Boubée & Cie.
DUSSART, B. H. & DEFAYE, D. 2001. *Introduction to the Copepoda,* Leiden, Backhuys Publishers.
EINSLE, U. 1993. *Crustacea: Copepoda: Calanoida und Cyclopoida*, Gustav Fischer.

EINSLE, U. K. 1988. TAXONOMY OF THE GENUS MEGACYCLOPS (CRUSTACEA, COPEPODA) -
MORPHOMETRY AND THE USE OF ENZYME ELECTROPHORESIS. *Hydrobiologia,* 167**,** 387-391.
ENGLUND, G., OHLUND, G., HEIN, C. L. & DIEHL, S. 2011. Temperature dependence of the functional
response. *Ecology Letters,* 14**,** 914-921.
ERITJA, R., ESCOSA, R., LUCIENTES, J., MARQUES, E., MOLINA, R., ROIZ, D. & RUIZ, S. 2005.
Worldwide invasion of vector mosquitoes: present European distribution and challenges for
Spain. *Biological Invasions,* 7**,** 87-97.
EUROPEAN CENTRE FOR DISEASE, P. & CONTROL. 2019. *Aedes albopictus - current known*
distribution: January 2019.
FRÜH, L., KAMPEN, H., SCHAUB, G. A. & WERNER, D. 2019. Predation on the invasive mosquito
Aedes japonicus (Diptera: Culicidae) by native copepod species in Germany. *J Vector Ecol,* 44**,**
241-247.
FRYER, G. 1957a. The feeding mechanism of some freshwater cyclopoid copepods. *Proceedings of*
the Zoological Society of London, 129.
FRYER, G. 1957b. The food of some freshwater cyclopoid copepods and its ecological significance.
Journal of Animal Ecology, 26**,** 263-286.
GRIGALTCHIK, V. S., WARD, A. J. W. & SEEBACHER, F. 2012. Thermal acclimation of interactions:
differential responses to temperature change alter predator-prey relationship. *Proceedings*
of the Royal Society B-Biological Sciences, 279**,** 4058-4064.
HOLLING, C. S. 1959. Some Characteristics of Simple Types of Predation and Parasitism. *The*
Canadian Entomologist, 91**,** 385-398.
HOLLING, C. S. 1966. The Functional Response of Invertebrate Predators to Prey Density. *Memoirs of*
the Entomological Society of Canada, 98**,** 5-86.
IRIGOIEN, X. 1998. Gut clearance rate constant, temperature and initial gut contents: a review.
Journal of Plankton Research, 20**,** 997-1003.
JASS, A., YERUSHALMI, G. Y., DAVIS, H. E., DONINI, A. & MACMILLAN, H. A. 2019. An impressive
capacity for cold tolerance plasticity protects against ionoregulatory collapse in the disease
vector Aedes aegypti. *Journal of Experimental Biology,* 222.
JULIANO, S. A. 2001. Non-linear curve fitting: predation and functional response curves. *In:*
SCHEINER, S. M. & GUREVITCH, J. (eds.) *Design and Analysis of Ecological Experiments.*
Oxford: Oxford University Press.
JULIANO, S. A. & LOUNIBOS, L. P. 2005. Ecology of invasive mosquitoes: effects on resident species
and on human health. *Ecology Letters,* 8**,** 558-574.
KALINOSKI, R. M. & DELONG, J. P. 2016. Beyond body mass: how prey traits improve predictions of
functional response parameters. *Oecologia,* 180**,** 543-550.
KAY, B. H., NAM, V. S., VAN TIEN, T., YEN, N. T., PHONG, T. V., DIEP, V. T. B., NINH, T. U., BEKTAS, A. &
AASKOV, J. G. 2002. Control of Aedes vectors of dengue in three provinces of Vietnam by use
of Mesocyclops (Copepoda) and community-based methods validated by entomologic,
clinical, and serological surveillance. *American Journal of Tropical Medicine and Hygiene,* 66**,**
40-48.
KIEFER, F. 1960. *Ruderfußkrebse (Copepoden),* Stuttgart, Kosmos-Verlag Franckh.
KLEKOWSKI, R. Z. & SHUSHKINA, E. A. 1966. *Ernährung, Atmung, Wachstum und Energie-Umformung*
in Macrocyclops albidus Jurine.
KRAEMER, M. U. G., REINER, R. C., JR., BRADY, O. J., MESSINA, J. P., GILBERT, M., PIGOTT, D. M., YI,
D., JOHNSON, K., EARL, L., MARCZAK, L. B., SHIRUDE, S., DAVIS WEAVER, N., BISANZIO, D.,
PERKINS, T. A., LAI, S., LU, X., JONES, P., COELHO, G. E., CARVALHO, R. G., VAN BORTEL, W.,
MARSBOOM, C., HENDRICKX, G., SCHAFFNER, F., MOORE, C. G., NAX, H. H., BENGTSSON, L.,
WETTER, E., TATEM, A. J., BROWNSTEIN, J. S., SMITH, D. L., LAMBRECHTS, L., CAUCHEMEZ,
S., LINARD, C., FARIA, N. R., PYBUS, O. G., SCOTT, T. W., LIU, Q., YU, H., WINT, G. R. W., HAY,
S. I. & GOLDING, N. 2019. Past and future spread of the arbovirus vectors Aedes aegypti and
Aedes albopictus. *Nat Microbiol,* 4**,** 854-863.

KUMAR, R. M. & RAO, T. R. 2003. Predation on mosquito larvae by Mesocyclops thermocyclopoides (Copepoda : Cyclopoida) in the presence of alternate prey. *International Review of Hydrobiology,* 88**,** 570-581.

LAYBOURN-PARRY, J., ABDULLAHI, B. A. & TINSON, S. V. 1988. Temperature-Dependent Energy Partitioning in the Benthic Copepods Acanthocyclops-Viridis and Macrocyclops-Albidus. *Canadian Journal of Zoology-Revue Canadienne De Zoologie,* 66**,** 2709-2714.

LOUNIBOS, L. P. 2002. Invasions by insect vectors of human disease. *Annual Review of Entomology,* 47**,** 233-266.

MAIER, G. 1994. PATTERNS OF LIFE-HISTORY AMONG CYCLOPOID COPEPODS OF CENTRAL-EUROPE. *Freshwater Biology,* 31**,** 77-86.

MARTEN, G. G. 1990a. Elimination of Aedes-Albopictus from Tire Piles by Introducing Macrocyclops-Albidus (Copepoda, Cyclopidae). *Journal of the American Mosquito Control Association,* 6**,** 689-693.

MARTEN, G. G. 1990b. Issues in the development of cyclops for mosquito control. *Arbovirus Research in Australia,* 5**,** 159-164.

MARTEN, G. G., BORDES, E. S. & NGUYEN, M. 1994. Use of Cyclopoid Copepods for Mosquito-Control. *Hydrobiologia,* 293**,** 491-496.

MARTEN, G. G. & REID, J. W. 2007. Cyclopoid copepods. *Journal of the American Mosquito Control Association,* 23**,** 65-92.

MEDLOCK, J. M., AVENELL, D., BARRASS, I. & LEACH, S. 2006. Analysis of the potential for survival and seasonal activity of Aedes albopictus (Diptera : Culicidae) in the United Kingdom. *Journal of Vector Ecology,* 31**,** 292-304.

MEDLOCK, J. M., VAUX, A. G., CULL, B., SCHAFFNER, F., GILLINGHAM, E., PFLUGER, V. & LEACH, S. 2017. Detection of the invasive mosquito species Aedes albopictus in southern England. *Lancet Infect Dis,* 17**,** 140.

METELMANN, S., CAMINADE, C., JONES, A. E., MEDLOCK, J. M., BAYLIS, M. & MORSE, A. P. 2019. The UK's suitability for Aedes albopictus in current and future climates. *Journal of the Royal Society Interface,* 16.

MURDOCK, C. C., EVANS, M. V., MCCLANAHAN, T. D., MIAZGOWICZ, K. L. & TESLA, B. 2017. Fine-scale variation in microclimate across an urban landscape shapes variation in mosquito population dynamics and the potential of Aedes albopictus to transmit arboviral disease. *Plos Neglected Tropical Diseases,* 11**,** e0005640.

NOVICH, R. A., ERICKSON, E. K., KALINOSKI, R. M. & DELONG, J. P. 2014. The temperature independence of interaction strength in a sit-and-wait predator. *Ecosphere,* 5.

PARADIS, A. R., PEPIN, P. & BROWN, J. A. 1996. Vulnerability of fish eggs and larvae to predation: Review of the influence of the relative size of prey and predator. *Canadian Journal of Fisheries and Aquatic Sciences,* 53**,** 1226-1235.

PETERS, R. H. 1983. Ingestion. *In:* BECK, E., BIRKS, H. J. B. & CONNOR, E. F. (eds.) *The ecological implications of body size.* Cambridge: Cambridge University Press.

POOLE, A. E., STILLMAN, R. A., WATSON, H. K. & NORRIS, K. J. 2007. Searching efficiency and the functional response of a pause-travel forager. *Functional Ecology,* 21**,** 784-792.

PRITCHARD, D., BARRIOS-O'NEILL, D., BOVY, H. & PATERSON, R. 2017a. *frair: Tools for functional response analysis, R package version 0.5.100* [Online]. Available: https://CRAN.R-project.org/package=frair [Accessed].

PRITCHARD, D. W., PATERSON, R. A., BOVY, H. C. & BARRIOS-O'NEILL, D. 2017b. FRAIR: an R package for fitting and comparing consumer functional responses. *Methods in Ecology and Evolution,* 8**,** 1528-1534.

RALL, B. C., BROSE, U., HARTVIG, M., KALINKAT, G., SCHWARZMULLER, F., VUCIC-PESTIC, O. & PETCHEY, O. L. 2012. Universal temperature and body-mass scaling of feeding rates. *Philosophical Transactions of the Royal Society B-Biological Sciences,* 367**,** 2923-2934.

ROSENBAUM, B. & RALL, B. C. 2018. Fitting functional responses: Direct parameter estimation by
simulating differential equations. *Methods in Ecology and Evolution,* 9**,** 2076-2090.
RUSSELL, B. M., MUIR, L. E., WEINSTEIN, P. & KAY, B. H. 1996. Surveillance of the mosquito Aedes
aegypti and its biocontrol with the copepod Mesocyclops aspericornis in Australian wells and
gold mines. *Medical and Veterinary Entomology,* 10**,** 155-160.
SCHREIBER, E. T., HALLMON, C. F., ESKRIDGE, K. M. & MARTEN, G. G. 1996. Effects of Mesocyclops
longisetus (Copepoda: Cyclopidae) on mosquitoes that inhabit tires: Influence of litter type,
quality, and quantity. *Journal of the American Mosquito Control Association,* 12**,** 688-694.
SOUMARE, M. K. F. & CILEK, J. E. 2011. The Effectiveness of Mesocyclops Longisetus (Copepoda) for
the Control of Container-Inhabiting Mosquitoes in Residential Environments. *Journal of the*
American Mosquito Control Association, 27**,** 376-383.
SUAREZ, M. F., MARTEN, G. G. & CLARK, G. G. 1992. A Simple Method for Cultivating Fresh-Water
Copepods used in Biological-Control of Aedes-Aegypti. *Journal of the American Mosquito*
Control Association, 8**,** 409-412.
TWARDOCHLEB, L. A., NOVAK, M. & MOORE, J. W. 2012. Using the functional response of a
consumer to predict biotic resistance to invasive prey. *Ecological Applications,* 22**,** 1162-
1171.
UITERWAAL, S. F. & DELONG, J. P. 2020. Functional responses are maximized at intermediate
temperatures. *Ecology*.
VERONESI, R., CARRIERI, M., MACCAGNANI, B., MAINI, S. & BELLINI, R. 2015. Macrocyclops Albidus
(Copepoda: Cyclopidae) for the Biocontrol of Aedes Albopictus and Culex Pipiens in Italy.
Journal of the American Mosquito Control Association, 31**,** 32-43.
WESTBROOK, C. J., REISKIND, M. H., PESKO, K. N., GREENE, K. E. & LOUNIBOS, L. P. 2010. Larval
environmental temperature and the susceptibility of Aedes albopictus Skuse (Diptera:
Culicidae) to Chikungunya virus. *Vector Borne Zoonotic Dis,* 10**,** 241-7.
ZIDON, R., TSUEDA, H., MORIN, E. & MORIN, S. 2016. Projecting pest population dynamics under
global warming: the combined effect of inter- and intra-annual variations. *Ecol Appl,* 26**,**
1198-210.

<u>Supporting Information</u>

Materials and methods

Collection of field temperature data:

In late February and early March of 2018, six empty car tires, each leaning on a wooden

crate, were placed in six different locations (Fig. S1), spread between two main sites: South

Kensington (London, UK) and Silwood Park (Berkshire, UK). Although most tires were at

ground-level near vegetation or shade, the "First Floor Balcony" tire (Fig. S1) had more

direct sun exposure, and thus, may be more representative of the conditions present in large

tire piles. Each tire was surveyed once a week from the 16^{th} of April until the 30^{th} of October

between 11am and 5pm, which matches the time period of predation for both our functional

response and predation efficiency experiments. If water was present in a tire, the water

temperature was taken using a hand-held Hanna Instruments thermometer (model name:

"Checktemp 1"). Temperature data from Silwood Park and South Kensington were recorded

from May through September of 2018 (Fig. S2), the months of predicted *Ae. albopictus* adult

activity (Medlock et al., 2006). There were 105 observations in total; 16 temperature

recordings were missing because the tire was dry, and two were excluded because they were

recorded outside the time period of 11am to 5pm.

865 Silwood Park sites   South Kensington sites

872 **Fig. S1.** Tire locations

881 **Fig. S2.** May through September of 2018 weekly tire water temperatures (Dotted lines

882 represent the three temperatures tested in the functional response and predation efficiency

883 experiments.)

a.)

• 1 mosquito larva prey

1 copepod predator

predator treatments

controls

b.)

M. albidus predators   *M. viridis* predators   predator-absent controls

Fig. S3. Experimental design: (a) functional response repeated six times, once for each combination of predator species and temperature; (b) predation efficiency repeated three

915 times, once for each temperature (Source of copepod silhouette accessed August 2019:

916 http://phylopic.org/name/f79c5b86-7b73-468a-9ad4-995646398f99)

918 *Ae. albopictus* hatching procedure for functional response and predation efficiency

919 experiments:

921 *Ae. albopictus* eggs were collected from the colony on filter papers and stored in plastic bags

922 containing damp paper towels to maintain humidity. Stored egg papers were submerged in 3

923 mg/L nutrient broth solution (Sigma-Aldrich © 70122 Nutrient Broth No 1) and oxygen was

924 displaced by vacuum suction for 30 min. Immediately following oxygen displacement,

925 ground fish food (Cichlid Gold Hikari®, Japan) was added *ad libitum*. The eggs were left in

926 this solution at $27 \pm 1°C$ for 12 h (Hanson and Craig, 1994) before the newly-hatched larvae

927 were counted (Fig. 1). The hatching temperature was kept high, relative to the experimental

928 temperatures (15-25°C), in order to maximize the hatch rate over a short, semi-synchronous

929 time period, especially since *Ae. albopictus* often hatch in multiple installments (Hanson and

930 Craig, 1994). The larval densities used in the functional response experiments were chosen

931 based on previous studies (Cuthbert et al., 2018, Cuthbert et al., 2019).

933 Predation efficiency linear regression models:

935 The following linear regression model was fitted to test temperature setting and copepod

936 species as predictors of predation efficiency:

938 $Y = \beta_0 + \beta_1(\text{Species}) + \beta_2(\text{Temperature}_{20}) + \beta_3(\text{Temperature}_{25}) + \varepsilon$   eqn S1

where Y represents the predation efficiency; β_0 is the Y-intercept; Species is a binary variable

denoted as 0 for *M. albidus* and 1 for *M. viridis*; Temperature$_{20}$ is a binary variable denoted

as 0 for 15°C and 1 for 20°C; Temperature$_{25}$ is a binary variable denoted as 0 for 15°C and 1

for 25°C; and ε is a random error term, assumed $\sim N(0, \sigma^2)$. Forty-seven observations were

included in the model; one was excluded because the copepod predator (*M. viridis* at 25°C)

was found to have died at some point during the 6 h predation period. The statistical

significance of each independent parameter estimate was analyzed at $\alpha = 0.05$.

An additional linear regression model was fitted to test temperature setting and copepod body

mass as predictors of predation efficiency:

$Y = \beta_0 + \beta_1(\text{Mass}) + \beta_2(\text{Temperature}_{20}) + \beta_3(\text{Temperature}_{25}) + \varepsilon$ eqn S2

where Mass is a continuous variable referring to each copepod's body mass in mg; and all

other notation is identical to that used in the previous model.

Both linear regression models above were also fitted without any temperature predictors.

Akaike information criterion (AIC) values were calculated for model selection, and the

Shapiro-Wilk test was used to confirm that the residuals of each model were normally

distributed.

965 **Results**

967 **Table S1.** Results of logistic regression of proportional consumption data as a function of

968 initial prey density using the "frair_test" function

Species	Temperature (°C)	n[a]	First Order Term[b]	p - value
M. albidus	15	28	-0.06	<0.0001
M. albidus	20	28	-0.08	<0.0001
M. albidus	25	25	-0.04	0.0007
M. viridis	15	27	-0.09	<0.0001
M. viridis	20	28	-0.08	<0.0001
M. viridis	25	25	-0.09	<0.0001

969   1.) Controls are not included for determining the type of functional response.

970   2.) The response is classified as "type II" when there is a significant negative

971   first-order term.

973 **Table S2.** Results of fitting the "nll.ode.general.mort" function for functional response curves

974 in which background mortality was observed

Species	Temp. (°C)	n[a]	Parameter	Estimate	Standard Error	p-value	-2 log L
M. albidus	15	35	attack coefficient	0.2379	0.0921	0.0098	102.73
			handling time	1.1583	0.2782	<0.0001	
			mortality rate	0.0063	0.0036	0.0816	
M. albidus	20	35	attack coefficient	0.4561	0.1156	<0.0001	90.98
			handling time	0.6036	0.0838	<0.0001	
			mortality rate[2]	0.0076	0.0038	0.0426	
M. albidus	25	32	attack coefficient	0.4909	0.3523	0.1634	111.25
			handling time	0.9163	0.3601	0.0109	
			mortality rate[b]	0.0289	0.0100	0.0040	
M. viridis	15	34	attack coefficient	0.4911	0.1214	<0.0001	100.14
			handling time	0.5163	0.0705	<0.0001	
			mortality rate	0.0038	0.0027	0.1569	
M. viridis	20	35	attack coefficient	0.4511	0.1063	<0.0001	93.30
			handling time	0.4848	0.0667	<0.0001	
			mortality rate	0.0057	0.0033	0.0823	

975   a.) Number of empirical observations includes both predator and control treatments.

976   b.) Background prey mortality had a significant impact on fitting the model for *M.*

977   *albidus* at 20 and 25°C.

980   **Table S3.** Results of fitting the "nll.bolker" function for functional response curves in which

981   background mortality was either not observed or insignificant

Species	Temp. (°C)	n	Parameter	Estimate	Standard Error	p-value	-2 log L
M. albidus	15	28	attack coefficient	0.2166	0.0716	0.0025	89.71
			handling time	0.9189	0.1696	<0.0001	
M. viridis	15	27	attack coefficient	0.4803	0.1137	<0.0001	92.14
			handling time	0.4853	0.0619	<0.0001	
M. viridis	20	28	attack coefficient	0.4430	0.0989	<0.0001	83.57
			handling time	0.4447	0.0554	<0.0001	
M. viridis	25	25	attack coefficient	0.5629	0.1438	<0.0001	85.53
			handling time	0.6527	0.0725	<0.0001	

983   **Table S4.** Functional response parameter estimates and 95% confidence intervals

Species	Temp. (°C)	Attack Coefficient (attacks per hour)	Handling Time (hours)
M. albidus	15	0.217, (0.116 – 0.425)	0.919, (0.596 – 1.28)
M. albidus	20	0.456, (0.277 – 0.747)	0.604, (0.452 – 0.798)
M. albidus	25	0.491, (0.178 – 13.9)	0.916, (0.451 – 1.99)
M. viridis	15	0.480, (0.305 – 0.777)	0.485, (0.368 – 0.615)
M. viridis	20	0.443, (0.287 – 0.691)	0.445, (0.337 – 0.559)
M. viridis	25	0.563, (0.338 – 0.921)	0.653, (0.519 – 0.808)

Table S5. Linear regression of predation efficiency by copepod species (n = 47)

Parameter	Estimate	Standard Error	p-value	Adjusted R²	AIC
Intercept	19.97	2.314	4.25 x 10⁻¹¹	0.081	362.6
Species (*M. viridis*, ref = *M. albidus*)	7.42	3.308	0.0299		

Table S6. Linear regression of predation efficiency by copepod body mass (n = 47)

Parameter	Estimate	Standard Error	p-value	Adjusted R²	AIC
Intercept	11.19	4.33	0.0130	0.156	358.5
Body Mass (mg)	34.75	11.26	0.0035		

Copepod body sizes by experimental design and species:

The ten gravid female copepods of each species that were measured as a validation set for the size of adult females included in the functional response experiments ranged from 1.2 to 1.7 mm for *M. albidus*, and from 1.4 to 2.3 mm for *M. viridis*. The non-gravid copepods used as predators in these experiments ranged from 1.2 to 1.9 mm (n = 81) for *M. albidus*, and from 1.4 to 2.5 mm (n = 80) for *M. viridis*. The eighty-one *M. albidus* copepods included in the functional response curve experiments ranged from 0.09 to 0.32 mg in body mass (median = 0.20 mg), and the eighty *M. viridis* copepods included in the functional response curve experiments ranged from 0.14 to 0.67 mg in body mass (median = 0.32 mg). Results of the Shapiro-Wilk test showed that the distribution of body masses included in the functional response experiments was significantly different from a normal distribution for both *M. albidus* (p-value = 0.0019) and *M. viridis* (p-value = 0.0001). The results of a Wilcoxon rank

1003   sum test showed that *M. viridis* used in the functional response experiments were generally

1004   larger than *M. albidus* used in the same experiments (p-value < 0.0001, Fig. S4).

1005

Twenty-four *M. albidus* copepods were used as predators in the predation efficiency

experiments, ranging in body mass from 0.14 to 0.36 mg (mean = 0.24 mg, standard

deviation = 0.06 mg), and 23 *M. viridis* were used, ranging in body mass from 0.23 to 0.67

mg (mean = 0.48 mg, standard deviation = 0.10 mg). Results of the Shapiro-Wilk test showed

that the distribution of body masses included in the predation efficiency experiments was not

significantly different from a normal distribution for both *M. albidus* (p-value = 0.0665) and

M. viridis (p-value = 0.2128). The results of a Welch two sample t-test for samples of

unequal variance showed that *M. viridis* used in the predation efficiency experiments were

significantly larger than *M. albidus* used in the same experiments (p-value < 0.0001, Fig. S4).

The Wilcoxon rank sum test was used to identify differences in copepod body mass by

experimental design, controlling for species. Copepods used in the predation efficiency

experiment were generally larger than those used in the functional response experiment for

both *M. albidus* (p-value = 0.0008) and *M. viridis* (p-value = 0.0003).

1019

1020

Fig. S4. Boxplot of copepod body mass by experimental design. Sample sizes: Functional

response, *M. albidus* = 81; Functional response, *M. viridis* = 80; Predation efficiency, *M.*

albidus = 24; Predation efficiency, *M. viridis* = 23.

Literature Cited

CUTHBERT, R. N., CALLAGHAN, A. & DICK, J. T. A. 2018. Dye another day: the

predatory impact of cyclopoid copepods on larval mosquito Culex pipiens is

unaffected by dyed environments. *Journal of Vector Ecology,* 43**,** 334-336.

CUTHBERT, R. N., CALLAGHAN, A. & DICK, J. T. A. 2019. The Effect of the Alternative

Prey, Paramecium caudatum (Peniculida: Parameciidae), on the Predation of Culex

pipiens (Diptera: Culicidae) by the Copepods Macrocyclops albidus and Megacyclops

viridis (Cyclopoida: Cyclopidae). *Journal of Medical Entomology,* 56**,** 276-279.

HANSON, S. M. & CRAIG, G. B., JR. 1994. Cold acclimation, diapause, and geographic

origin affect cold hardiness in eggs of Aedes albopictus (Diptera: Culicidae). *J Med*

Entomol, 31**,** 192-201.

1036   MEDLOCK, J. M., AVENELL, D., BARRASS, I. & LEACH, S. 2006. Analysis of the

1037   potential for survival and seasonal activity of Aedes albopictus (Diptera : Culicidae)

1038   in the United Kingdom. *Journal of Vector Ecology,* 31**,** 292-304.

1039

1040