**Environmental variables determining the distribution of an avian parasite: the**

**case of the *Philornis torquans* complex (Diptera: Muscidae) in South America**

Pablo F. Cuervo[1], Alejandro Percara[1], Lucas Monje[1], Pablo M. Beldomenico[1], Martín

A. Quiroga[1,2,3]

[1]Laboratorio de Ecología de Enfermedades, Instituto de Ciencias Veterinarias del

Litoral (ICIVET - Litoral), Universidad Nacional del Litoral (UNL) / Consejo Nacional

de Investigaciones Científicas y Técnicas (CONICET), Esperanza, Santa Fe, Argentina

[2]The Peregrine Fund, Boise, Idaho, USA

[3]Facultad de Ciencia y Tecnología (FCyT), Universidad Autónoma de Entre Ríos

(UADER), Oro Verde, Entre Ríos, Argentina

Running title: *P. torquans* complex in South America

Corresponding author: Martín A. Quiroga. Laboratorio de Ecología de Enfermedades,

Instituto de Ciencias Veterinarias del Litoral (ICIVET - Litoral), Universidad Nacional

del Litoral (UNL) / Consejo Nacional de Investigaciones Científicas y Técnicas

(CONICET), Esperanza, Santa Fe, Argentina. The Peregrine Fund, Boise, Idaho,

USA.[3]Facultad de Ciencia y Tecnología (FCyT), Universidad Autónoma de Entre Ríos

(UADER), Oro Verde, Entre Ríos, Argentina. mquiroga06@hotmail.com

**Abstract**

*Philornis* flies are the major cause of myasis in altricial nestlings of neotropical birds.

Its impact ranges from subtle to lethal, being of major concern in endangered bird

species with geographically-restricted, fragmented and small-sized populations. In spite

of its relevance for bird conservation, there is little information about the environmental

dimensions determining their geographical range. We identified for the first time the

macro-environmental variables constraining the abiotic niche of the *P. torquans*

complex in South America, and provided a model map of its potential distribution based

on environmental suitability. We used maximum entropy to model the ecological niche

of the *P. torquans* complex, and to determine the contributing scenopoetic variables.

We applied most of the recommendations to produce robust ecological niche models:

accounting for sampling bias and spatial autocorrelation; using an accessible area for

model calibration; testing multiple combinations of model parameters; using multiple

statistical criteria; using a number of model replicates to determine consistency and

uncertainty in geographical predictions. We identified the minimum temperature of the

coldest month as the most relevant variable, associated with the largest decrease in

habitat suitability in Brazil and northern South America. Furthermore, the mean

temperature of the warmest quarter limited suitability mostly along the Andean range.

In addition, humidity and moisture are influential factors in most of Argentina, northern

Chile and coastal Peru. The geographical projection of the niche model suggests that

environments in most of central-eastern Argentina, and in a broad area in central Chile,

are suitable for the presence of the *P. torquans* complex. Besides of contributing to the

knowledge of the ecology of the genus, this study is of relevance as a tool for bird

conservation and represents a good reference for future work on the distribution of this

parasite genus.

Keywords: *Philornis torquans* complex; distribution model; southern South America;

Maxent

**Introduction**

In the Neotropics, *Philornis* Meinert 1890 [Diptera: Muscidae] is the major cause of

myasis in birds (Guimarães & Papavero, 1999; Dudaniec & Kleindorfer, 2006).

*Philornis* adults are free living, while larvae are largely associated with altricial nestling

birds, establishing two types of parasitic associations: semi-haematophagous or

subcutaneous (Teixeira, 1999). The impact of *Philornis* spp. on hosts ranges from subtle

(Dudaniec & Kleindorfer, 2006) to strong and lethal (Dudaniec *et al.*, 2007; Quiroga &

Reboreda, 2012), being of particular concern in vulnerable and endangered bird species

(Bulgarella *et al.*, 2019). While in healthy populations effects of nest parasitism could

sometimes be negligible, in geographically-restricted, fragmented and small-sized

populations, the effects of *Philornis* parasitism could be exerting additional pressure

(Bulgarella *et al.*, 2019); thus, pushing these populations to the edge of extinction.

Evidence that these parasitic flies can be a serious threat to bird conservation is the

severe impact caused by *Philornis downsi* [Dodge & Aitkens 1968] on Darwin's finches

(i.e. Kleindorfer & Dudaniec, 2016) and by *P. pici* [Macquart, 1854] on Ridgway's

hawks (*Buteo ridgwayi* - Hayes *et al.*, 2018). Indeed, it may cause up to 75% mortality

of nestlings (Fessl *et al.*, 2006; Hayes *et al.*, 2018).

Despite this genus comprises about 50 fly species (Teixeira, 1999) and parasitize more

than 250 bird species in the Neotropics (Couri, 1999; Dudaniec & Kleindorfer, 2006;

Antoniazzi *et al.*, 2011), there are enormous gaps of information about aspects of its

biology and ecology (Dudaniec & Kleindorfer, 2006; Patitucci *et al.*, 2017). One aspect

of particular interest and poorly known is the environmental dimensions determining the

geographical range of *Philornis* spp. A better understanding of the distribution of this

genus may be of relevance to identify locations with high presence of these parasitic

flies that might threaten populations of endangered bird species. Furthermore, it may be

also appropriate to locate habitats with low risk of *Philornis* occurrence, useful for the

re-introduction of bird species, the reinforcement of their populations, or to identify

*Philornis*-free areas that are prone to be colonized.

The present study is aimed to identify what macro-environmental variables drive the

geographical distribution of *Philornis* species in South America by assessing their

relative contribution to the environmental niche. We focus on *P. torquans* and *P. seguyi*,

a relatively well-known complex of species within *Philornis*, to start understanding the

role of scenopoetic dimensions on the geographical distribution of *Philornis*. As

temperature and rainfall are presumed to be the most relevant abiotic parameters

affecting *Philornis* spp. life cycle (Dudaniec *et al.*, 2007; Antoniazzi *et al.*, 2011;

Manzoli *et al.*, 2013), we hypothesise that low temperatures and reduced precipitation

will constrain the occurrence of these species. Ultimately, our contribution to the

knowledge on the geographical distribution of these parasitic flies might prove a

powerful tool relevant for bird conservation all over the neotropics.

**Materials and Methods**

*Study species*

*Philornis torquans* [Nielsen, 1913] and *P. seguyi* [Garcia, 1952] are the two

subcutaneous *Philornis* species most commonly occurring in southern South America

(Couri *et al.*, 2009; Silvestri *et al.*, 2011), where eight *Philornis* species have been

described but only half of them are recognized as valid by Couri *et al.* (2009) (namely,

*P. torquans* , *P. seguyi* , *P. blanchardi* [Garcia, 1952] and *P. downsi*). Specimens

identified morphologically as either *P. torquans* or *P. seguyi* are indistinguishable at the

molecular level (Monje *et al.*, 2013). This is the reason why it was suggested that the

subcutaneous *Philornis* morphs resembling *P. torquans* or *P. seguyi* in southern South

America should be treated as *Philornis torquans* complex (Quiroga *et al.*, 2016). This

complex of parasitic flies was selected because: i) its biology and ecoepidemiology have

been studied for more than a decade (Antoniazzi *et al.*, 2011; Quiroga & Reboreda,

2012; Manzoli *et al.*, 2013, 2018; Saravia-Pietropaolo *et al.*, 2018); ii) this is the group

of flies with the highest number of confirmed morphology-based records; and iii) it is

known to affect at least 34 species of birds (see Table S1), and suspected to parasitize

endangered species, such as the Saffron-Cowled Blackbird (*Xanthopsar flavus* -

BirdLife International, 2019) and the Yellow Cardinal (*Gubernatrix cristata* - BirdLife

International, 2018). Both bird species were reported as hosts of unidentified *Philornis*

species (Domínguez *et al.*, 2015; Pucheta *et al.*, 2019) in the vicinity of *P. torquans*

complex records.

*Species occurrence data*

Occurrence data of *Philornis* species was obtained from two different sources: i) field

surveys conducted in Argentina; and ii) a thorough review of existing literature.

Regarding field surveys, *Philornis* larvae and pupae were recovered from hand-

inspected nestlings (during the breeding season) or disassembled nest remains (year-

round) and reared to adulthood in captivity by AP. Adults were identified to species

level by MAQ using the available taxonomic key (Couri, 1999; Couri *et al.*, 2009). Our

literature survey included 265 peer-reviewed publications ranging from 1853 until 2019,

which we then systematically reviewed for localities were parasitism occurred. Due to

the lack of occurrence data of free-living adults, information obtained from immature

*Philornis* stages represents a valid surrogate to assess the environmental suitability for

the species.

The specimens identified based on morphology as *P. torquans* or *P. seguyi* were

considered as members of the *Philornis torquans* complex (see Quiroga *et al.*, 2016),

which accounted for 80 occurrence records (most of them concerning different host

species in the same site). After removing duplicate localities, the dataset was reduced to

34 occurrence sites (Table S1). Since spatial autocorrelation in environmental

information and consequent non-independence of occurrence data could bias predictions

(Boria *et al.*, 2014), we spatially filtered sites preserved for calibration with the package

"spThin" in R Statistical Software 3.5.0 (Aiello-Lammens *et al.*, 2015; R Core Team,

2018). Spatial thinning of occurrence records provides an easy-to-implement and

relatively straightforward method to alleviate the effects of sampling bias (Boria *et al.*,

2014; Aiello-Lammens *et al.*, 2015). To attain a uniform distribution of occurrences and

to avoid excessive clustering within any particular region, we explored different

filtering distances in sequential steps, visually inspecting results. Once a satisfactory

spatial distribution of occurrence data was obtained (i.e., when spatial clustering of

records was no longer evident), we ended up with 18 sites for the *P. torquans* complex

(Fig. 1). We split the *P. torquans* complex locations randomly in equal quarts: one set

of locations for calibration (75%) and another for evaluation during model calibration

(25%).

**Figure 1.** Geographical coordinates representing locations of *Philornis torquans* complex and accessible area (M) used in the construction of models.

*Environmental data and selection of variables*

To characterize climatic conditions meaningful to biological species, we used the so-called "bioclimatic" variables based on WorldClim from the CHELSA database (www.chelsa-climate.org, Karger *et al.*, 2017). In addition, we included the climatic variables from the ENVIREM database, derived from and complementing the bioclimatic variables from WorldClim (www.envirem.github.io, Title & Bemmels, 2018). We excluded nine data layers (detailed in supplementary Table S2), as these variables include artefacts that create abrupt differences between neighbouring pixels. We coarsed environmental data to 10 geographical minutes (~17 x ~17 km) spatial

resolution, which approximately matches the uncertainty inherent in the occurrence

data.

As a first step and in order to test our hypothesis, we selected a set of five variables

based on current knowledge on the biology of *Philornis* spp.: i) mean temperature of

warmest quarter (Bio10); ii) minimum temperature of coldest month (Bio06); iii)

precipitation of driest month (Bio14); iv) precipitation of wettest quarter (Bio16) (the

four of them from CHELSA); v) CMI, a climatic moisture index from ENVIREM. The

"mean temperature of warmest quarter" (Bio10) was selected because this variable is

related to the sum of degree days available for development, a usual method used to

predict the life stages of other dipteran species (e.g., Psychodidae, Stratiomyidae,

Calliphoridae) (Kasap & Alten, 2005; Harnden & Tomberlin, 2016; Yanmanee *et al.*,

2016). The "minimum temperature of the coldest month" (Bio06) was selected due to

its negative effect on developmental times and survival of pupae and adults of a wide

variety of insects (Sinclair, 2015), therefore arguably limiting *Philornis* distribution.

Furthermore, Manzoli *et al.* (2013) found that preceding increases in rainfall were

highly correlated with the abundance of *Philornis torquans* parasitic stages. This aspect

was considered by including the "precipitation of the driest month" (Bio14) and the

"precipitation of the wettest quarter" (Bio16). The "climatic moisture index" (CMI) was

included as an integrative marker of environmental conditions, as it is a metric of

potential water availability imposed solely by climate (ranges from –1 to +1, with wet

climates showing positive values, and dry climates negative (Vörösmarty *et al.*, 2005)).

Additionally, we created six other sets of variables to be considered in the evaluation

and calibration of models. The variables from the original set of data were grouped in

three different sets: i) a dataset including only the 15 bioclimatic variables from

CHELSA (referred to hereafter as the *bioclim* dataset); ii) a dataset using the *bioclim*

dataset plus the 11 climatic ENVIREM variables (hereafter referred to as the *climatic*

dataset); and iii) a dataset using the *climatic* dataset plus the two topographic variables

from ENVIREM (hereafter referred to as the *environmental* dataset). Then, we reduced

model complexity and collinearity by selecting two simpler sets of variables from each

of these three datasets, based on the calibration of preliminary Maxent models (details

below). For the selection of these simpler sets of variables, we applied two procedures

available in the R package "SDMtune" (Vignali *et al.*, 2019): removal of highly

correlated variables ($R > 0.7$), and removal of variables with low importance for the

model performance (percent contribution $< 5\%$). Finally, for the calibration of models,

we retained six sets of environmental variables: i) a set based on current knowledge on

the biology of *Philornis* spp. (Set 1); ii) three sets of uncorrelated variables from the

*bioclim*, *climatic* and *environmental* datasets (Sets 2a, 3a and 4); and iii) two sets of

relevant variables selected from the *bioclim* and *climatic* datasets (Sets 2b and 3b) (the

selection from the *climatic* and *environmental* datasets resulted in two equivalent sets of

relevant variables) (Datasets detailed in Table S2).

*Niche modelling and assessment of variables importance*

We used Maxent 3.4.1 for developing models (Phillips *et al.*, 2006). Maxent is a

machine-learning method that estimates the species potential geographical distribution

by finding the probability distribution of maximum entropy (closest to uniform), subject

to the constraint of the expected values of the environmental predictors. Maxent was

developed to use presence-only data by contrasting presences against background

locations (Phillips *et al.*, 2006; Merow *et al.*, 2013), and has shown to outperform other

algorithms, even when applied to small data sets like ours (Hernandez *et al.*, 2006; Wisz

*et al.*, 2008; van Proosdij *et al.*, 2016).

To minimize the impact of assumptions about absences from areas that are not

accessible to the species (Barve *et al.*, 2011), we restricted model calibration to the area

to which the species likely had access via dispersal. Considering there is no available

information on the dispersal capacity of the *P. torquans* complex, we defined the

adequate accessible area (M) as a maximum area within 1000 km of any of the

occurrences (Fig. 1), based on: i) the average geographical distance between a centroid

point and all the occurrences (~730 km); ii) the high dispersal capacity (Dudaniec *et al.*,

2008) and assisted dispersion of *Philornis downsi* to the Galapagos Islands (Kleindorfer

& Dudaniec, 2016); and iii) the dispersal capacity of other Muscidae species (Sabrosky,

1961; Hogsette & Ruff, 1985).

The complexity of models built with Maxent can be adjusted with the inclusion of

additional feature classes (i.e. transformations of the original predictor variables) that

allow for increasingly complex models, as well as with a regularization multiplier that

contributes to select those features and reduces overfitting (Warren & Seifert, 2011;

Merow *et al.*, 2013). The preliminary models for each of the three data sets (the *bioclim*,

*climatic* and *environmental* datasets), later used for variable selection, were calibrated

using the "kuenm" R package (Cobos *et al.*, 2019). To determine the optimal model

complexity, we explored all combinations of [a] 17 values of the regularization

parameter (0.1–1.0 at intervals of 0.1, 2–6 at intervals of 1, and 8 and 10), and [b] nine

sets of potential combinations of four feature classes: linear "l", quadratic "q", product

"p", and hinge "h" ("l", "lq", "lp", "lqp", "h", "lh", "lqh", "lph", "lqph")".

After the selection of variables, we conducted a detailed model selection exercise,

exploring all combinations of [a] the 17 values of the regularization parameter, [b] the

nine sets of potential combinations of four feature classes, and [c] the six sets of

bioclimatic variables (see above). As such, a total of 918 models were calibrated, and

each model was evaluated for statistical significance (partial ROC tests), performance

(omission rate), and the Akaike Information Criterion corrected for small sample sizes

(AICc). Final models were selected under the three criteria (first, statistical significance,

then omission rate, followed by AICc) (Cobos *et al.*, 2019).

We used the best parameter settings chosen to create final models. For each final model,

we ran 30 bootstrap replicates, with no clamping or extrapolation, and retaining a

random partition of 25% of the points from each run. These final models, though only

calibrated across M, were transferred to the extent of South America, particularly to

southern South America, which is the focus of this contribution. We used the cloglog

output format to assess median values across replicates as an estimate of the spatial

distribution of suitable and unsuitable conditions for the *P. torquans* complex across

South America. The uncertainty in model predictions was estimated using the range in

the suitability values (i.e., maximum minus minimum) across the model replicates. To

provide an additional assessment on the reliability of our model transfers, we calculated

the mobility-oriented parity (MOP) metric (Owens *et al.*, 2013) to offer a view of the

novelty of environmental values and combinations of values across the transfer area,

with special attention paid to areas of strict extrapolation (i.e., transfer areas with values

outside the range of climates in the calibration area).

We assessed the importance of each variable to determine the environmental suitability

for the *P. torquans* complex with median estimates across replicates of percent

contribution, permutation importance and Jackknife analysis. In order to determine the

most limiting variables, we used the Maxent limiting factor mapping tool described in

Elith *et al*. (2010), and implemented via the package "rmaxent" (Baumgartner *et al.*,

2017). The limiting factors are an interesting insight into the drivers of predictions, and

provide a powerful basis for the statement of hypotheses concerning physiological

tolerance and aptitude (Elith *et al.*, 2010).

**Results**

We assessed 918 models for the final model calibration. Seven hundred and twenty-

seven of them were statistically significant as compared with a null model of random

prediction. Of such significant models, 137 (~19%) met the omission criterion of < 5%.

Of the significant, low-omission models, only three met the AICc criterion of ΔAICc <

2. These three models were calibrated with the set of variables selected based on current

knowledge (Set 1), and were variants of the regularization multiplier (Table 1). Models

based on current knowledge selection of variables outperformed the ones calibrated

with automated selected datasets, as the best of models calibrated with Set 1 presented a

ΔAICc = 21.8 when compared with the best model calibrated with other set of variables.

**Table 1**. Performance metrics for parameter settings regarding regularization multiplier

(RMs), feature classes (FCs), and sets of environmental predictors (Pred. sets), used for

creating final models for the *P. torquans* complex. FCs are as follows: linear = l,

quadratic = q, product = p, and hinge = h.

| RMs | FCs | Pred. Sets | Partial ROC | Omission rate 5% | AICc | ΔAICc | Weight AICc | # of parameters |
|-----|-----|------------|-------------|------------------|------|-------|-------------|-----------------|
| 0.8 | Lq | Set 1 | 0.00 | 0.00 | 362.95 | 0 | 0.21 | 4 |
| 0.9 | Lq | Set 1 | 0.00 | 0.00 | 363.18 | 0.22 | 0.19 | 4 |
| 1 | Lq | Set 1 | 0.00 | 0.00 | 363.41 | 0.46 | 0.17 | 4 |

The "minimum temperature of the coldest month" (Bio06) was the most influential

variable, as evidenced by the percent contribution, permutation importance and

Jackknife analysis (Table 2 and Fig. S1). Its response curve had a bell shape, with a

central range optimal for prediction, while in both extremes its influence appeared to be

detrimental (Fig. S2). The second most important variable was the "mean temperature

of warmest quarter" (Bio10), followed by the "climatic moisture index" (CMI). The

CMI response curve was similar to the one from Bio06, while the response curve from

Bio10 had a sigmoidal shape with positive relation (Fig. S2). The "precipitation of the

driest month" (Bio14) and the "precipitation of the wettest quarter" (Bio16) had nearly

no influence on the distribution modelling of the *P. torquans* complex.

**Table 2**. Variable importance: percent contribution and permutation importance

(median and sd). References: Bio06 = minimum temperature of coldest month; Bio10 =

mean temperature of warmest quarter; Bio14 = precipitation of driest month; Bio16 =

precipitation of wettest quarter; CMI = climatic moisture index.

| Variable | Percent Contribution | Permutation Importance |
|---|---|---|
| **Bio06** | 54.75 (5.67) | 70.59 (8.09) |
| **Bio10** | 25.10 (5.77) | 18.71 (5.93) |
| **Bio14** | 1.38 (3.29) | 0.64 (4.27) |
| **Bio16** | 0.19 (1.42) | 0.88 (4.43) |
| **CMI** | 12.89 (9.31) | 6.44 (7.07) |

287    The minimum temperature of coldest month (Bio06) was the predictor associated with

288    the largest decrease in habitat suitability in Brazil and northern South America (Fig. 2).

289    For the majority of Argentina, the climatic moisture index (CMI) was the most limiting

290    factor, while the mean temperature of warmest quarter (Bio10) limited suitability

291    mostly along the Andean range (Fig. 2).

**Figure 2**. Limiting factors (for any point, the limiting factor is the variable whose value

at that point most influences the model prediction). References: Bio06 = minimum

temperature of coldest month; Bio10 = mean temperature of warmest quarter; Bio14 =

precipitation of driest month; Bio16 = precipitation of wettest quarter; CMI = climatic

moisture index. [Colours after the Golden-Collared Tanager *Iridosornis jelskii*]. Patterns

of colours for figures were generated with the package "tanagR" (Cadena & Zapata,

2019), inspired by the plumage of passerine birds in the tanager family (Thraupidae)

from Central and South America.

As more than one best model was selected, we used the median and range of all

replicates across parameters to consolidate results for the *P. torquans* complex. The

median of the selected models (Fig. 3A) identified areas with different levels of habitat

suitability for the *P. torquans* complex across South America. Highly suitable areas

were concentrated on the central-east of Argentina and Uruguay, comprising the

Pampean and Chaco (eastern Chaco district) biogeographical provinces (Fig. S3, see in

Morrone, 2017). Suitability declined towards western Argentina, southern Brazil and

Paraguay. A broad area in central Chile was identified as environmentally suitable,

although it may not be accessible to the *P. torquans* complex by natural dispersion. The

model showed moderate uncertainty areas mainly throughout central-western Argentina

(the Monte biogeographical province), central Chile and the Andean range (Fig. 3B and

Fig. S3). The MOP analysis revealed high environmental similarity in most of South

America, and thus almost no risk of strict model extrapolation (Fig. 3C).

**Figure 3**. Climatologically suitable regions for the *Philornis torquans* complex

distribution in South America. A) Median prediction. B) Uncertainty (range) associated

with the median prediction of suitable regions for the *Philornis torquans* complex

distribution in South America. C) Mobility-oriented parity analysis revealing similarity

of environmental values in South America for the detection of areas of strict

extrapolation. [Colours after Scarlet-Breasted Dacnis *Dacnis berlepschi* and Moss-

backed Tanager *Bangsia edwardsi*]

**Discussion**

In this study, we identified for the first time the macro-environmental (scenopoetic)

variables constraining the abiotic niche of the *P. torquans* complex in South America,

and provided a model map of its potential distribution based on environmental

suitability. Knowledge on the potential distribution of this group of parasitic flies might

allow to identify areas where threatened or endemic population of birds may be affected

by these parasites; thus being of avian conservation concern.

This study took into consideration most of the recent methodological recommendations

to produce robust ecological niche models, as outlined by Fen*g et al.* (2019). As in any

modelling approach, model quality is clearly influenced by the number of records used

in model building. Thus, it could be argued that the reduced number of occurrences

used for modelling in this study is not adequate to produce a reliable niche model.

However, Maxent is among the algorithms with best predictive power across a wide

range of sample sizes (as low as ten records) (see Hernandez *et al.*, 2006; Wisz *et al.*,

2008), having moderate sample size sensitivity combined with excellent predictive

ability (Wisz *et al.*, 2008). Maxent's strong and consistent performance across sample

sizes may be explained by the way it uses regularization to avoid over-fitting (Wisz *et*

*al.*, 2008). The amount of regularization varies flexibly with sample size to ensure

consistent performance (Warren & Seifert, 2011; Shcheglovitova & Anderson, 2013),

which we have considered when evaluating a number of models across a range of

regularization values and features classes (as suggested in Morales *et al.*, 2017). All the

same, this modelling approach and its results could benefit from future additions of

confirmed occurrence records to the database.

We hypothesised that low temperatures and reduced precipitation will restrain the

occurrence of the *Philornis torquans* complex, as these abiotic parameters play a

predominant role in establishing the distribution limits of organisms and have already

been related with the intensity of parasitism by *Philornis* larvae (Dudaniec *et al.*, 2007;

Antoniazzi *et al.*, 2011; Manzoli *et al.*, 2013). Since insects are ectothermic, their rate of

development and survival can vary greatly in response to many biotic and abiotic

factors (Adams, 1979; Couret *et al.*, 2014). Winters are a critical period for insects at

higher latitudes since overwintering conditions strongly affect their fitness (Irwin &

Lee, 2000). Indeed, our model identified the minimum temperature of coldest month

(Bio06) as the most relevant variable (Table 2 and Fig. S1), supporting our hypothesis.

Morphs suggested to belong to the *P. torquans* complex (*P. torquans and P. seguyi*)

(Monje *et al.*, 2013) are native to southern South America (with origins somewhere near

the border between Brazil and Uruguay - Löwenberg-Neto & De Carvalho, 2009).

Representing the southernmost species of the genus, and unlike the majority of

*Philornis* species, the *P. torquans* complex must be adapted to endure colder, longer

and drier winters, typical of temperate areas.

Furthermore, our model revealed an unexpected detrimental effect in both sides of a

suitable range of winter temperatures (signalled by Bio06, Fig. S2), suggesting that the

*P. torquans* complex might present a reaction norm of its thermal aptitude along

latitudes. Besides the expected negative effect of low temperatures during winter, our

model suggests that warm winters may be limiting the northern distribution of the *P.*

*torquans* complex (Fig. 2). Such scenario seems plausible, as insects adapted to cold

winters may not perform well in mild winters (Irwin & Lee, 2000), since warmer

temperatures may increase energy consumption during this period (Irwin & Lee, 2000;

Sinclair, 2015). Therefore, these individuals may not be able to reach the next breeding

season or may leave them without enough energy reserves for early reproduction

(Sinclair, 2015).

As aforementioned, the developmental rate of insects is related to environmental

conditions, particularly as temperature impacts critical rate processes throughout

development and adulthood (Kingsolver & Huey, 2008). The temperature dependence

of ectotherms' developmental processes is commonly represented as the cumulative

sum of degree-time products required for the development of key life stages, such as

egg maturation or emergence from pupae (e.g., Kasap & Alten, 2005; Harnden &

Tomberlin, 2016; Yanmanee *et al.*, 2016). In these regards, the mean temperature of the

warmest quarter (Bio10) proved to be the second most important variable (Table 2 and

Fig. S1), showing a positive relation with sigmoidal shape between predicted

probability of presence and Bio10 (Fig. S2). This sigmoidal shape implies a threshold

temperature (apparently around 10°C), above which the predicted probability of

presence increases exponentially. The former makes sense when considering that the

increase of Bio10 might favour the development of temperature-dependent non-parasitic

stages, such as pupae maturation, which ultimately relates with the abundance of adults.

Otherwise, temperatures below the apparent threshold seems to limit the distribution of

the *P. torquans* complex, as evinced along the Andean range (Fig. 2).

The third most relevant variable, the climatic moisture index (CMI), was the most

influential factor in most of Argentina, northern Chile and coastal Peru (Fig. 2). The

CMI values observed through the actual distribution of the *P. torquans* complex

indicate that the species must endure mostly a dry environment when compared with

most of South America. As reported for species laying eggs into a relatively dry

environment (Grzywacz *et al.*, 2012), the eggs of the *P. torquans* morphs present a

small portion of the egg surface that functions as a plastron (Patitucci *et al.*, 2017), a

respiratory structure that facilitates oxygen income minimizing water loss (Hinton,

1960). The reduced size of the plastron of the *P. torquans* morph (Patitucci *et al.*, 2017)

may have evolved as an adaptation allowing the radiation to the drier conditions typical

of central Argentina (see Löwenberg-Neto & De Carvalho, 2009).

Opposed to our prediction, the ecological niche model suggests that precipitation has no

direct influence on limiting the environmental suitability of the *P. torquans* complex.

The precipitation of the driest month (Bio14) and the precipitation of the wettest quarter

(Bio16) had a poor contribution to the model performance (Table 2), and showed a

weak explanatory power in the jackknife analysis (Fig. S1). Though the humidity and

moisture are clearly relevant factors determining the *P. torquans* complex distribution,

as reflected by CMI and supported by this model, the proximal influence of

precipitation seems to be mainly related with the intensity of parasitism by *Philornis*

species, as previously reported (Arendt, 1985, 2000; Dudaniec *et al.*, 2007; Antoniazzi

*et al.*, 2011; Manzoli *et al.*, 2013).

The outcome of the geographical projection of the environmental niche model was a

map of the potential distribution of the *P. torquans* complex in South America.

According to our results, the *P. torquans* complex is potentially distributed across

central-eastern Argentina, mostly restricted to the Pampean and Chaco biogeographical

provinces, and declining with some uncertainty in the surrounding Monte

biogeographical province (for more details see in Morrone, 2017) (Fig. 3 and Fig. S3).

Despite this niche model is based on scenopoetic variables constrained by accessibility

(see Soberón, 2010), but ignoring biotic interactions, the projected potential distribution

of the *P. torquans* complex does not seem to be limited by the availability of host

species. In fact, the *P. torquans* complex was reported to parasitize at least 34 bird

species (Table S1), some of them fairly abundant and widely distributed in South

America (e.g., *Pitangus sulphuratus*, *Troglodytes aedon*, *Zonotrichia capensis*).

The area predicted as suitable for the *P. torquans* complex in central Chile should be

considered with care. First, the estimated suitability is based solely on environmental

(scenopoetic) conditions, and thus biotic interactions should prove adequate for the

occurrence of the complex in that area. Second, it is required to verify the present

occurrence of a *Philornis* species in the predicted area, which could represent either the

finding of a new population of the *P. torquans* complex, or the discovery of a sister

species. Third, the Andean range might represent a pronounced topographical barrier to

be sorted out by the *P. torquans* complex upon its arrival to the estimated suitable area,

either by natural dispersion or by human intervention.

In conclusion, this represents the first study to identify macro-environmental variables

driving the geographical distribution of a *Philornis* species by evaluating their

contribution to its' environmental niche. Besides of contributing to the knowledge of the

ecology of the genus and integrating current methodological recommendations to

produce a robust ecological niche prediction, our model is of relevance as a tool for bird

conservation and represents a good reference for future work on the distribution of this

parasite genus.

**Acknowledgements**

We greatly acknowledge the contribution made by Salvador S.A., who kindly provided

the geographical coordinates from the sites mentioned in Salvador & Bodrati (2013).

We also thank Dra. Bulgarella for the critical reading of this manuscript.

This work was funded by Consejo Nacional de Investigaciones Científicas y Técnicas

[PIP11220100100261; PIP 11220130100561CO]; and Universidad Autónoma de Entre

Ríos [PIDAC S013805/15]. The funders had no role in study design, data collection and

analysis, decision to publish, or preparation of the manuscript.

**Contribution of authors**

PFC and MAQ participated in the design and coordination of the study, and drafted the

manuscript; PFC performed the data analysis and modelling; AP carried out field

surveys and revised the existing literature; LM and PMB provided financial support and

critically reviewed the manuscript. All authors revised the manuscript and gave final

approval for publication.

**Competing interests**

We have no competing interests.

**References**

Adams, T.S. (1979) The reproductive physiology of the Screwworm, *Cochliomyia*

*hominivorax* (Diptera: Calliphoridae) II. Effect of constant temperatures on oogenesis.

*Journal of Medical Entomology*, **15**, 484–487.

Aiello-Lammens, M.E., Boria, R.A., Radosavljevic, A., Vilela, B. & Anderson, R.P.

(2015) spThin: An R package for spatial thinning of species occurrence records for use

in ecological niche models. *Ecography*, **38**, 541–545.

Antoniazzi, L.R., Manzoli, D.E., Rohrmann, D., Saravia, M.J., Silvestri, L. &

Beldomenico, P.M. (2011) Climate variability affects the impact of parasitic flies on

Argentinean forest birds. *Journal of Zoology*, **283**, 126–134.

Arendt, W.J. (1985) Philornis Ectoparasitism of Pearly-eyed Thrashers. I. Impact on

Growth and Development of Nestlings. *The Auk*, **102**, 270–280.

Arendt, W.J. (2000) Impact of nest predators, competitors, and ectoparasites on Pearly-

eyed Thrashers, with comments on the potential implications for Puerto Rican Parrot

recovery. *Ornitologia Neotropical*, 13–63.

Barve, N., Barve, V., Jiménez-Valverde, A., Lira-Noriega, A., Maher, S.P., Peterson,

A.T., *et al.* (2011) The crucial role of the accessible area in ecological niche modeling

and species distribution modeling. *Ecological Modelling*, **222**, 1810–1819.

Baumgartner, J., Wilson, P. & Esperón-Rodríguez, M. (2017) rmaxent: Tools for

working with Maxent in R. R package version 0.8.3.9000.

BirdLife International. (2018) Gubernatrix cristata [WWW Document]. *The IUCN Red

List of Threatened Species*. URL http://dx.doi.org/10.2305/IUCN.UK.2018-

2.RLTS.T22721578A131888081.en [accessed on 2018].

BirdLife International. (2019) Xanthopsar flavus [WWW Document]. *The IUCN Red

List of Threatened Species*. URL https://dx.doi.org/10.2305/IUCN.UK.2019-

3.RLTS.T22724673A153660526.en [accessed on 2019].

Boria, R.A., Olson, L.E., Goodman, S.M. & Anderson, R.P. (2014) Spatial filtering to

reduce sampling bias can improve the performance of ecological niche models.

*Ecological Modelling*, **275**, 73–77.

Bulgarella, M., Quiroga, M.A. & Heimpel, G.E. (2019) Additive negative effects of

Philornis nest parasitism on small and declining Neotropical bird populations. *Bird

Conservation International*, **29**, 339–360.

Cadena, D. & Zapata, F. (2019) tanagR: Tanager-Inspired Color Palettes. R package

version 0.1.0.

Cobos, M.E., Peterson, A.T., Barve, N. & Osorio-Olvera, L. (2019) kuenm: an R

package for detailed development of ecological niche models using Maxent. *PeerJ*, **7**,

e6281.

Couret, J., Dotson, E. & Benedict, M.Q. (2014) Temperature, larval diet, and density

effects on development rate and survival of *Aedes aegypti* (Diptera: Culicidae). *PLOS*

*ONE*, **9**, e87468.

Couri, M. (1999) Myiasis caused by obligatory parasites. Ia. *Philornis* Meinert

(Muscidae). In *Myiasis in man and animals in the neotropical region* (ed. by

Guimarães, J.H. & Papavero, N.). Editora Plêiade, pp. 51–70.

Couri, M.S., Antoniazzi, L.R., Beldomenico, P. & Quiroga, M. (2009) Argentine

*Philornis* Meinert species (Diptera: Muscidae) with synonymic notes. *Zootaxa*, **62**, 52–

62.

Domínguez, M., Reboreda, J.C. & Mahler, B. (2015) Impact of Shiny Cowbird and

botfly parasitism on the reproductive success of the globally endangered Yellow

Cardinal *Gubernatrix cristata*. *Bird Conservation International*, **25**, 294–305.

Dudaniec, R.Y., Fessl, B. & Kleindorfer, S. (2007) Interannual and interspecific

variation in intensity of the parasitic fly, *Philornis downsi*, in Darwin's finches.

*Biological Conservation*, **139**, 325–332.

Dudaniec, R.Y., Gardner, M.G., Donnellan, S. & Kleindorfer, S. (2008) Genetic

variation in the invasive avian parasite, *Philornis downsi* (Diptera, Muscidae) on the

Galápagos archipelago. *BMC Ecology*, **8**, 13.

Dudaniec, R.Y. & Kleindorfer, S. (2006) Effects of the parasitic flies of the genus

*Philornis* (Diptera: Muscidae) on birds. *Emu*, **106**, 13–20.

Elith, J., Kearney, M. & Phillips, S. (2010) The art of modelling range-shifting species.

*Methods in Ecology and Evolution*, **1**, 330–342.

Feng, X., Park, D.S., Walker, C., Peterson, A.T., Merow, C. & Papeş, M. (2019) A

checklist for maximizing reproducibility of ecological niche models. *Nature Ecology &*

*Evolution*, **3**, 1382–1395.

Fessl, B., Sinclair, B.J. & Kleindorfer, S. (2006) The life-cycle of *Philornis downsi*

(Diptera: Muscidae) parasitizing Darwin's finches and its impacts on nestling survival.

*Parasitology*, **133**, 739–747.

Grzywacz, A., Szpila, K. & Pape, T. (2012) Egg morphology of nine species of Pollenia

Robineau-Desvoidy, 1830 (Diptera: Calliphoridae). *Microscopy Research and*

*Technique*, **75**, 955–967.

Guimarães, J.H. & Papavero, N. (1999) *Myiasis in man and animals in the Neotropical*

*Region*. Editora Plêiade, São Paulo.

Harnden, L.M. & Tomberlin, J.K. (2016) Effects of temperature and diet on black

soldier fly, *Hermetia illucens* (L.) (Diptera: Stratiomyidae), development. *Forensic*

*Science International*, **266**, 109–116.

Hayes, C.D., Hayes, T.I., McClure, C.J.W., Quiroga, M., Thorstrom, R.K., Anderson,

D.L., *et al.* (2018) Native parasitic nest fly impacts reproductive success of an island-

endemic host. *Animal Conservation*, **22**, 157–164.

Hernandez, P.A., Graham, C.H., Master, L.L., Albert, D.L., Deborah, L. & Albert, D.L.

(2006) The effect of sample size and species characteristics on performance of different

species distribution modeling methods. *Ecography*, **29**, 773–785.

Hinton, H.E. (1960) The chorionic plastron and its role in the eggs of the Muscinae

(Diptera). *Journal of Cell Science*, **s3-101**, 313–332.

Hogsette, J.A. & Ruff, J.P. (1985) Stable Fly (Diptera: Muscidae) Migration in

Northwest Florida. *Environmental Entomology*, **14**, 170–175.

Irwin, J.T. & Lee, R.E. (2000) Mild winter temperatures reduce survival and potential

fecundity of the goldenrod gall fly, *Eurosta solidaginis* (Diptera: Tephritidae). *Journal*

*of Insect Physiology*, **46**, 655–661.

Karger, D.N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R.W., *et al.*

(2017) Climatologies at high resolution for the earth's land surface areas. *Scientific*

*Data*, **4**, 170122.

Kasap, O.E. & Alten, B. (2005) Laboratory estimation of degree-day developmental

requirements of *Phlebotomus papatasi* (Diptera: Psychodidae). *Journal of Vector*

*Ecology*, **30**, 328–333.

Kingsolver, J.G. & Huey, R.B. (2008) Size, temperature, and fitness: three rules.

*Evolutionary Ecology Research*, **10**, 251–268.

Kleindorfer, S. & Dudaniec, R.Y. (2016) Host-parasite ecology, behavior and genetics:

A review of the introduced fly parasite *Philornis downsi* and its Darwin's finch hosts.

*BMC Zoology*.

Löwenberg-Neto, P. & Carvalho, C.J.B. De. (2009) Areas of endemism and spatial

diversification of the Muscidae (Insecta: Diptera) in the Andean and Neotropical

regions. *Journal of Biogeography*, **36**, 1750–1759.

Manzoli, D.E., Antoniazzi, L.R., Saravia, M.J., Silvestri, L., Rorhmann, D. &

Beldomenico, P.M. (2013) Multi-level determinants of parasitic fly infection in forest

passerines. *PLoS ONE*, **8**, e67104.

Manzoli, D.E., Saravia-Pietropaolo, M.J., Antoniazzi, L.R., Barengo, E., Arce, S.I.,

Quiroga, M.A., *et al.* (2018) Contrasting consequences of different defence strategies in

a natural multihost- parasite system. *International Journal of Parasitology*.

Merow, C., Smith, M.J. & Silander, J.A. (2013) A practical guide to MaxEnt for

modeling species' distributions: What it does, and why inputs and settings matter.

*Ecography*, **36**, 1058–1069.

Monje, L.D., Quiroga, M., Manzoli, D.E., Couri, M.S., Silvestri, L., Venzal, J.M., *et al.*

(2013) Sequence analysis of the internal transcribed spacer 2 (ITS2) from *Philornis*

*seguyi* (García, 1952) and *Philornis torquans* (Nielsen, 1913) (Diptera: Muscidae).

*Systematic Parasitology*, **86**, 43–51.

Morales, N.S., Fernández, I.C. & Baca-González, V. (2017) MaxEnt's parameter

configuration and small samples: are we paying attention to recommendations? A

systematic review. *PeerJ*, **5**, e3093.

Morrone, J.J. (2017) *Neotropical biogeography. Regionalization and evolution*. CRC

Press, Inc.

Owens, H.L., Campbell, L.P., Dornak, L.L., Saupe, E.E., Barve, N., Soberón, J., *et al.*

(2013) Constraints on interpretation of ecological niche models by limited

environmental ranges on calibration areas. *Ecological Modelling*, **263**, 10–18.

Patitucci, L.D., Quiroga, M., Couri, M.S. & Saravia-Pietropaolo, M.J. (2017)

Oviposition in the bird parasitic fly *Philornis torquans* (Nielsen, 1913) (Diptera:

Muscidae) and eggs' adaptations to dry environments. *Zoologischer Anzeiger*, **267**, 15–

20.

Phillips, S.J., Anderson, R.P. & Schapire, R.E. (2006) Maximum entropy modeling of

species geographic distributions. *Ecological Modelling*, **190**, 231–259.

Proosdij, A.S.J. van, Sosef, M.S.M., Wieringa, J.J. & Raes, N. (2016) Minimum

required number of specimen records to develop accurate species distribution models.

*Ecography*, **39**, 542–552.

Pucheta, M.F., Patitucci, L.D., Bulgarella, M., Pereda, M.I., Giacomo, A.S. Di &

Kopuchian, C. (2019) Primer registro de parasitismo de *Philornis* en pichones de Tordo

Amarillo (*Xanthopsar flavus*). In *XVIII Reunión Argentina de Ornitología (RAO)*.

Tandil, Argentina, p. 103.

Quiroga, M.A., Monje, L.D., Arrabal, J.P. & Beldomenico, P.M. (2016) Datos

moleculares nuevos sobre *Philornis* (Diptera: Muscidae) subcutáneas del sur de

Sudamérica sugieren la existencia de un complejo de especies. *Revista Mexicana de*

*Biodiversidad*, **87**, 1383–1386.

Quiroga, M.A. & Reboreda, J.C. (2012) Lethal and sublethal effects of Botfly

(*Philornis seguyi*) parasitism on House Wren nestlings. *The Condor*, **114**, 197–202.

R Core Team. (2018) R: A language and environment for statistical computing.

Sabrosky, C.W. (1961) Our First Decade with the Face Fly, *Musca autumnalis. Journal*

*of Economic Entomology*, **54**, 761–763.

Saravia-Pietropaolo, M.J., Arce, S.I., Manzoli, D.E., Quiroga, M. & Beldomenico, P.M.

(2018) Aspects of the life cycle of the avian parasite *Philornis torquans* (Diptera:

Muscidae) under laboratory rearing conditions. *The Canadian Entomologist*, **150**, 317–

325.

Shcheglovitova, M. & Anderson, R.P. (2013) Estimating optimal complexity for

ecological niche models: A jackknife approach for species with small sample sizes.

*Ecological Modelling*, **269**, 9–17.

Silvestri, L., Antoniazzi, L.R., Couri, M.S., Monje, L.D. & Beldomenico, P.M. (2011)

First record of the avian ectoparasite *Philornis downsi* Dodge & Aitken, 1968 (Diptera:

Muscidae) in Argentina. *Systematic Parasitology*, **80**, 137–140.

Sinclair, B.J. (2015) Linking energetics and overwintering in temperate insects. *Journal*

*of Thermal Biology*, **54**, 5–11.

Soberón, J.M. (2010) Niche and area of distribution modeling: a population ecology

perspective. *Ecography*, **33**, 159–167.

Teixeira, D. (1999) Myiasis caused by obligatory parasites. Ib. General observations on

the biology of species of the genus Philornis Meinert, 1890 (Diptera, Muscidae). In

*Myiasis in man and animals in the Neotropical Region* (ed. by Guimarães, J.H. &

Papavero, N.). Pleaide/FAPESP, São Paulo, pp. 71–96.

Title, P.O. & Bemmels, J.B. (2018) ENVIREM: an expanded set of bioclimatic and

topographic variables increases flexibility and improves performance of ecological

niche modeling. *Ecography*, **41**, 291–307.

Vignali, S., Barras, A. & Braunisch, V. (2019) SDMtune: Species Distribution Model

Selection.

Vörösmarty, C.J., Douglas, E.M., Green, P.A. & Revenga, C. (2005) Geospatial

indicators of emerging water stress: an application to Africa. *AMBIO: A Journal of the*

*Human Environment*, **34**, 230–236.

Warren, D.L. & Seifert, S.N. (2011) Ecological niche modeling in Maxent: the

importance of model complexity and the performance of model selection criteria.

*Ecological Applications*, **21**, 335–342.

Wisz, M.S., Hijmans, R.J., Li, J., Peterson, A.T., Graham, C.H. & Guisan, A. (2008)

Effects of sample size on the performance of species distribution models. *Diversity and*

*Distributions*, **14**, 763–773.

Yanmanee, S., Husemann, M., Benbow, M.E. & Suwannapong, G. (2016) Larval

development rates of *Chrysomya rufifacies* Macquart, 1842 (Diptera: Calliphoridae)

within its native range in South-East Asia. *Forensic Science International*, **266**, 63–67.