Environmental variables determining the presence of an avian parasite: the case of the *Philornis torquans* complex (Diptera: Muscidae) in South America

Pablo F. Cuervo[1], Alejandro Percara[1], Lucas Monje[1], Pablo M. Beldomenico[1], Martín A. Quiroga[1,2,3]

[1]Laboratorio de Ecología de Enfermedades, Instituto de Ciencias Veterinarias del Litoral (ICIVET - Litoral), Universidad Nacional del Litoral (UNL) / Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Esperanza, Santa Fe, Argentina

[2]The Peregrine Fund, Boise, Idaho, USA

[3]Facultad de Ciencia y Tecnología (FCyT), Universidad Autónoma de Entre Ríos (UADER), Oro Verde, Entre Ríos, Argentina

Running title: *P. torquans* complex in South America

Corresponding author: Martín A. Quiroga. Laboratorio de Ecología de Enfermedades, Instituto de Ciencias Veterinarias del Litoral (ICIVET - Litoral), Universidad Nacional del Litoral (UNL) / Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Esperanza, Santa Fe, Argentina. The Peregrine Fund, Boise, Idaho, USA.[3]Facultad de Ciencia y Tecnología (FCyT), Universidad Autónoma de Entre Ríos (UADER), Oro Verde, Entre Ríos, Argentina. mquiroga06@hotmail.com

Abstract

1. *Philornis* flies are exclusive of birds and their nests, being some of them of

conservation concern. Despite many studies addressed different aspects of this

problem, none have focused on the geographical distribution of these species.

2. In this study, besides providing a model map of environmental suitability of the *P.*

torquans complex in South America, we also contributed with data on the

constraints of its abiotic niche.

3. We provided baseline data (supplementary information) on sites where *P. torquans*

complex is present in southern South America, valuable for future *Philornis*

distribution modelling efforts.

4. According to our results, the *P. torquans* complex is potentially distributed across

central-eastern Argentina, mostly restricted to the Pampean and Chaco (eastern

Chaco district) biogeographical provinces.

5. The potential distribution of the *P. torquans* complex mainly relates to the "minimum

temperature of the coldest month" and "mean temperature of warmest quarter". In

addition, moisture/aridity seems to be exerting additional constraints.

6. Our work is the first one to identify macro-environmental variables driving the

geographical distribution of a *Philornis* species by evaluating their contribution to

its' environmental niche.

7. Besides the relevance of our model as a tool for bird's conservation, it integrates the

most up-to-date methodological recommendations to predict a robust ecological

niche; thus representing a methodological reference for future work on the

distribution of this group of parasites.

Keywords: *Philornis torquans* complex; distribution model; southern South America;

Maxent

Introduction

The infestation of live vertebrates with dipteran larvae which feed on the host's dead or

living tissue and body fluids is known as myiasis (Zumpt, 1965). In birds, myiasis is

mainly caused by species of the genera *Protocalliphora* [Hough, 1899 -Diptera:

Calliphoridae], *Passeromyia* [Rodhain & Villeneuve, 1915 - Diptera: Muscidae] and

Philornis [Meinert, 1890 - Diptera: Muscidae] (Little, 2008), *Carnus* [Nitzsch, 1818 -

Diptera: Carnidae] (Petrescu & Adam, 2000), *Neottiophilum* [von Frauenfeld, 1868 -

Diptera: Neottiophilidae] (Krivosheina *et al.*, 2017). Of these genera, the only one

occurring in the Neotropics is *Philornis*, comprising about 50 described species (Couri,

1999; Dudaniec & Kleindorfer, 2006). *Philornis* adults are free living, while larvae are

strictly associated with bird nestlings, establishing different types of associations:

coprophagous, semi-haematophagous or subcutaneous (Dudaniec & Kleindorfer, 2006).

Despite more than 250 bird species have been reported as *Philornis* hosts in South

America (Couri, 1999; Dudaniec & Kleindorfer, 2006; Löwenberg-Neto, 2008;

Antoniazzi *et al.*, 2011), there are enormous gaps of information on *Philornis* parasitism

and its effect on host birds; especially concerning vulnerable and endangered bird

species. Impact of *Philornis* spp. on hosts ranges from subtle (Kinsella & Winegarner,

1974; Oniki, 1983; Young, 1993; Fessl *et al.*, 2006) to strong and lethal (Dudaniec *et*

al., 2006, 2007; Quiroga & Reboreda, 2012), an depends on the outcome of the

interaction between traits of both parasite and host (Bronstein, 1994; Manzoli *et al.*,

2018).

While in healthy populations effects of nest parasitism could sometimes be negligible;

in geographically-restricted, fragmented and small-sized populations, the effects of

Philornis parasitism could be exerting additional pressure (Bulgarella *et al.*, 2019); thus,

pushing these populations to the edge of extinction. Evidence that this parasite can be a

serious threat to bird conservation is the severe impact caused by *Philornis downsi*

[Dodge & Aitkens 1968] on Darwin's finches (i.e. Koop *et al.*, 2011) and by *P. pici*

[Macquart, 1854] on Ridgway's hawks (*Buteo ridgwayi* - Hayes *et al.*, 2018). Indeed, it

may cause up to 75% of nestlings mortality (Fessl *et al.*, 2006; Hayes *et al.*, 2018).

Therefore, more research should be conducted on the occurrence and effects of these

parasites. One aspect of particular interest and poorly known is the environmental

dimensions determining the geographical distribution of *Philornis* spp. Such knowledge

is essential for better understanding the ecology of the genus, and may be useful to

identify *Philornis*-free areas that are prone to be colonized, to identify locations with

high presence of these flies that might threaten populations of endangered bird species,

or to locate habitats with low risk of *Philornis* occurrence for the re-introduction of bird

species or reinforcement of their populations.

The present study is aimed to identify what macro-environmental variables drive the

geographical distribution of *Philornis* species in South America by assessing their

relative contribution to the environmental niche. We focus on *P. torquans* and *P. seguyi*,

a relatively well-known species complex within *Philornis*, to start understanding the

role of environmental dimensions on the geographic distribution of *Philornis*. As

temperature and rainfall are presumed to be the most relevant abiotic parameters

affecting *Philornis* spp. life cycle (Dudaniec *et al.*, 2007; Antoniazzi *et al.*, 2011;

Manzoli *et al.*, 2013), we hypothesise that low temperatures and reduced precipitation

will restrain the occurrence of these species.

Materials and Methods

Study species

Philornis torquans [Nielsen, 1913] and *P. seguyi* [Garcia, 1952] are the two

subcutaneous *Philornis* species most commonly occurring in southern South America

(Couri *et al.*, 2009; Silvestri *et al.*, 2011), where eight *Philornis* species have been

described but only half of them are recognized as valid by Couri *et al.* (2009) (namely,

P. torquans , *P. seguyi* , *P. blanchardi* [Garcia, 1952] and *P. downsi*). Specimens

identified morphologically as either *P. torquans* or *P. seguyi* are undistinguishable at

the molecular level (Monje *et al.*, 2013). This is the reason why it was suggested that

the subcutaneous *Philornis* morphs resembling *P. torquans* or *P. seguyi* in southern

South America should be treated as *Philornis torquans* complex (Quiroga *et al.*, 2016).

This complex of parasitic flies was selected because: i) its biology and ecoepidemiology

have been studied for more than a decade (Antoniazzi *et al.*, 2011; Quiroga &

Reboreda, 2012; Manzoli *et al.*, 2013, 2018; Saravia-Pietropaolo *et al.*, 2018); ii) this is

the group of flies with the highest number of confirmed morphology-based records; and

iii) it could be potentially affecting the endangered Yellow Cardinal (*Gubernatrix*

cristata - BirdLife International, 2018), as this bird species hosts an unidentified

Philornis species (Domínguez *et al.*, 2015; Beier & Fontana, 2019) in the vicinity of *P.*

torquans complex records.

Species occurrence data

Occurrence data of *Philornis* species was obtained from field surveys conducted in

Argentina and from existing literature. In every case, *Philornis* adults were obtained

from larvae or pupae recovered from nestlings or nest remains, and were identified to

species level following Couri (1999). Due to the absence of occurrence data of free-

living adults, information obtained from immature *Philornis* stages represents a valid

surrogate to assess the environmental suitability for the species. The specimens

identified based on morphology as *P. torquans* or *P. seguyi* were considered as

members of the *Philornis torquans* complex (see Quiroga *et al.*, 2016), which accounted

for 80 occurrence records. After removing duplicate localities, the data was reduced to

34 occurrence sites (Table S1). Since spatial autocorrelation in environmental

information and consequent non-independence of occurrence data could bias predictions

(Kramer-Schadt *et al.*, 2013), we spatially filtered sites preserved for calibration with

the package "spThin" in R Statistical Software 3.5.0 (Aiello-Lammens *et al.*, 2015; R

Core Team, 2018). Spatial thinning of occurrence records provides an easy-to-

implement and relatively straightforward method to alleviate the effects of sampling

bias (Boria *et al.*, 2014; Aiello-Lammens *et al.*, 2015). To attain a uniform distribution

of occurrences and to avoid excessive clustering within any particular region, we

explored different filtering distances in sequential steps, visually inspecting results.

Once a satisfactory distribution of occurrence data was obtained (i.e., when spatial

clustering of records was no longer evident), we ended with 18 sites for the *P. torquans*

complex (Fig. 1). We split the *P. torquans* complex locations randomly in equal quarts:

one set of locations for calibration (75%) and another for evaluation during model

calibration (25%).

Figure 1. Geographic coordinates representing locations of *Philornis torquans* complex

and accessible area (M) used in the construction of distribution models.

Environmental data and selection of variables

To characterize climatic conditions meaningful to biological species, we used the so-

called "bioclimatic" variables based on WorldClim from the CHELSA database

(www.chelsa-climate.org, Karger *et al.*, 2017). In addition, we included the climatic

variables from the ENVIREM database, derived from and complementing the

bioclimatic variables from WorldClim (www.envirem.github.io, Title & Bemmels,

2018). We excluded nine data layers (mean temperature of wettest quarter, mean

temperature of driest quarter, precipitation of warmest quarter, precipitation of driest

quarter, mean PET of coldest quarter, mean PET of warmest quarter, mean PET of

driest quarter, mean PET of wettest quarter, and count of the number of months with

mean temperature greater than 10°C), as these variables include artefacts that create

abrupt differences between neighbouring pixels (Escobar *et al.*, 2014; Raghavan *et al.*,

2019). We coarsed environmental data to 10' (~17 x ~17 km) spatial resolution, which

approximately matches the uncertainty inherent in the occurrence data.

As a first step and in order to test our hypothesis, we selected a set of five variables

based on current knowledge on the biology of *Philornis* spp.: i) mean temperature of

warmest quarter (Bio10); ii) minimum temperature of coldest month (Bio06); iii)

precipitation of driest month (Bio14); iv) precipitation of wettest quarter (Bio16) (the

four of them from CHELSA); v) a climatic moisture index (CMI, a metric of relative

wetness and aridity, from ENVIREM). The "mean temperature of warmest quarter"

(Bio10) was selected because this variable is related to the sum of degree days available

for development, a usual method used to predict the life stages of other dipteran species

(e.g., Kasap & Alten, 2005; Nieto *et al.*, 2006; Gutierrez & Ponti, 2014; Harnden &

Tomberlin, 2016; Yanmanee *et al.*, 2016). The "minimum temperature of the coldest

month" (Bio06) was selected due to its negative effect on developmental times and

survival of pupae and adults (Sinclair, 2015), therefore arguably limiting *Philornis*

distribution. Furthermore, Manzoli *et al.* (2013) found that preceding increases in

rainfall were highly correlated with the abundance of *Philornis torquans* parasitic

stages. This aspect was considered by including the "precipitation of the driest month"

(Bio14) and the "precipitation of the wettest quarter" (Bio16). A "climatic moisture

index" (CMI) was included as an integrative marker of environmental conditions,

strongly related to temperature and precipitation.

Additionally, we created six other sets of variables to be considered in the evaluation

and calibration of models. The variables from the original set of data were grouped in

three different sets: i) a dataset including only the 15 bioclimatic variables from

CHELSA (referred to hereafter as the *bioclim* dataset); ii) a dataset using the *bioclim*

dataset plus the 11 climatic ENVIREM variables (hereafter referred to as the *climatic*

dataset); and iii) a dataset using the *climatic* dataset plus the two topographic variables

from ENVIREM (hereafter referred to as the *environmental* dataset). Then, we reduced

model complexity and collinearity by selecting two simpler sets of variables from each

of these three datasets, based on the calibration of preliminary Maxent models (details

below). For the selection of these simpler sets of variables, we applied two procedures

available in the R package "SDMtune" (Vignali *et al.*, 2019): removal of highly

correlated variables ($R > 0.7$), and removal of variables with low importance for the

model performance (percent contribution $< 5\%$). Finally, for the calibration of models,

we retained six sets of environmental variables: i) a set based on current knowledge on

the biology of *Philornis* spp. (Set 1); ii) three sets of uncorrelated variables from the

bioclim, *climatic* and *environmental* datasets (Sets 2a, 3a and 4); and iii) two sets of

relevant variables selected from the *bioclim* and *climatic* datasets (Sets 2b and 3b) (the

selection from the *climatic* and *environmental* datasets resulted in two equivalent sets of

relevant variables) (Datasets in Table S2).

Niche modelling and assessment of variables importance

We used Maxent 3.4.1 for developing models (Phillips *et al.*, 2006). Maxent is a

machine-learning method that estimates the species potential geographic distribution by

finding the probability distribution of maximum entropy (closest to uniform), subject to

the constraint of the expected values of the environmental predictors. Maxent was

developed to use presence-only data by contrasting presences against background

locations (Phillips *et al.*, 2006; Merow *et al.*, 2013), and has shown to outperform other

algorithms, even when applied to small data sets like ours (Elith *et al.*, 2006; Hernandez

et al., 2006; van Proosdij *et al.*, 2016).

To minimize the impact of assumptions about absences from areas that are not

accessible to the species (Barve *et al.*, 2011), we restricted model calibration to the area

to which the species likely had access via dispersal. We defined the adequate accessible

area (M) as a maximum area within 1000 km of any of the occurrences, based on the

average geographic distance between a centroid point and all the occurrences (~730

km), the assisted dispersion of *Philornis downsi* to the Galapagos Islands (Kleindorfer

& Dudaniec, 2016) and the dispersal capacity of other Muscidae species (Labud *et al.*,

2003; Showler & Osbrink, 2015).

The complexity of models built with Maxent can be adjusted with the inclusion of

additional feature classes (i.e. transformations of the original predictor variables) that

allow for increasingly complex models, as well as with a regularization multiplier that

contributes to select those features and reduces overfitting (Merow *et al.*, 2013; Warren

et al., 2016). The preliminary models for each of the three data sets (the *bioclim*,

climatic and *environmental* datasets), later used for variable selection, were calibrated

using the "kuenm" R package (Cobos *et al.*, 2019). To determine the optimal model

complexity, we explored all combinations of [a] 17 values of the regularization

parameter (0.1–1.0 at intervals of 0.1, 2–6 at intervals of 1, and 8 and 10), and [b] nine

sets of potential combinations of four feature classes: linear "l", quadratic "q", product

"p", and hinge "h" ("l", "lq", "lp", "lqp", "h", "lh", "lqh", "lph", "lqph")".

After the selection of variables, we conducted a detailed model selection exercise,

exploring all combinations of [a] the 17 values of the regularization parameter, [b] the

nine sets of potential combinations of four feature classes, and [c] the six sets of

bioclimatic variables (see above). As such, a total of 918 models were calibrated, and

each model was evaluated for statistical significance (partial ROC tests), performance

(omission rate), and the Akaike Information Criterion corrected for small sample sizes

(AICc). Final models were selected under the three criteria (first, statistical significance,

then omission rate, followed by AICc) (Cobos *et al.*, 2019).

We used the best parameter settings chosen to create final models. For each final model,

we ran 30 bootstrap replicates, with no clamping or extrapolation, and retaining a

random partition of 25% of the points from each run. These final models, though only

calibrated across M, were transferred to the extent of South America, particularly to

southern South America, which is the focus of this contribution. We used the cloglog

output format to assess median values across replicates as an estimate of the spatial

distribution of suitable and unsuitable conditions for the *P. torquans* complex across

South America. The uncertainty in model predictions was estimated using the range in

the suitability values (i.e., maximum minus minimum) across the model replicates. To

provide an additional assessment on the reliability of our model transfers, we calculated

the mobility-oriented parity (MOP) metric (Owens *et al.*, 2013) to offer a view of the

novelty of environmental values and combinations of values across the transfer area,

with special attention paid to areas of strict extrapolation (i.e., transfer areas with values

outside the range of climates in the calibration area).

We assessed the importance of each variable to determine the environmental suitability

for the *P. torquans* complex with median estimates across replicates of percent

contribution, permutation importance and Jackknife analysis. In order to determine the

most limiting variables, we used the Maxent limiting factor mapping tool described in

Elith *et al*. (2010), and implemented via the package "rmaxent" (Baumgartner *et al.*,

2017). The limiting factors are an interesting insight into the drivers of predictions, and

provide a powerful basis for comparisons with physiological knowledge (Elith *et al.*,

2010).

Patterns of colours for figures were generated with the package "tanagR" (Cadena &

Zapata, 2019), inspired by the plumage of passerine birds in the tanager family

(Thraupidae). The package currently includes palettes based on 16 tanager species from

Central and South America.

Results

We assessed 918 models for the final model calibration. Seven hundred and twenty-

seven of them were statistically significant as compared with a null model of random

prediction. Of such significant models, 137 (~19%) met the omission criterion of < 5%.

Of the significant, low-omission models, only three met the AICc criterion of ΔAICc <

2. These three models were calibrated with the set of variables selected based on current

knowledge (Set 1), and were variants of the regularization multiplier (Table 1). Models

based on current knowledge selection of variables outperformed the ones calibrated

with automated selected datasets, as the best of models calibrated with Set 1 presented a

ΔAICc = 21.8 when compared with the best model calibrated with other set of variables.

Table 1. Performance metrics for parameter settings regarding regularization multiplier

(RMs), feature classes (FCs), and sets of environmental predictors (Pred. sets), used for

creating final models for the *P. torquans* complex. FCs are as follows: linear = l,

quadratic = q, product = p, and hinge = h.

RMs	FCs	Pred. Sets	Partial ROC	Omission rate 5%	AICc	ΔAICc	Weight AICc	# of parameters
0.8	Lq	Set 1	0.00	0.00	362.95	0	0.21	4
0.9	Lq	Set 1	0.00	0.00	363.18	0.22	0.19	4
1	Lq	Set 1	0.00	0.00	363.41	0.46	0.17	4

The "minimum temperature of the coldest month" (Bio06) was the most influential

variable, as evidenced by the percent contribution, permutation importance and

Jackknife analysis (Table 2 and Fig. 2). Its response curve evidence a bell shape, with a

central range optimal for prediction, while in both extremes its influence appeared to be

detrimental (Fig. S1). The second most important variable was the "mean temperature

of warmest quarter" (Bio10), followed by the "climatic moisture index" (CMI). The

CMI response curve was similar to the one from Bio06, while the response curve from

Bio10 evinced a sigmoidal shape with positive relation (Fig. S1). The "precipitation of

the driest month" (Bio14) and the "precipitation of the wettest quarter" (Bio16) had

nearly no influence on the distribution modelling of the *P. torquans* complex.

Table 2. Variable importance: percent contribution and permutation importance

(median and sd). References: Bio10 = mean temperature of warmest quarter; Bio06 =

minimum temperature of coldest month; Bio14 = precipitation of driest month; Bio16 =

precipitation of wettest quarter; CMI = climatic moisture index.

Variable	Percent Contribution	Permutation Importance
Bio06	54.75 (5.67)	70.59 (8.09)
Bio10	25.10 (5.77)	18.71 (5.93)
Bio14	1.38 (3.29)	0.64 (4.27)
Bio16	0.19 (1.42)	0.88 (4.43)
CMI	12.89 (9.31)	6.44 (7.07)

Figure 2. Results of the jackknife test of variable importance, when a variable is used in isolation or it is omitted from modelling. References: Bio10 = mean temperature of warmest quarter; Bio06 = minimum temperature of coldest month; Bio14 = precipitation of driest month; Bio16 = precipitation of wettest quarter; CMI = climatic moisture index. [Colours after the Green-Grass Tanager *Chlorornis riefferii*]

The minimum temperature of coldest month (Bio06) was the predictor associated with the largest decrease in habitat suitability in Brazil and northern South America (Fig. S2). For the majority of Argentina, the climatic moisture index (CMI) was the most limiting factor, while the mean temperature of warmest quarter (Bio10) limited suitability mostly along the Andean range (Fig. S2).

Figure 3. Climatologically suitable regions for the *Philornis torquans* complex

distribution in South America. a) Median prediction. b) Uncertainty (range) associated

with the median prediction of suitable regions for the *Philornis torquans* complex

distribution in South America. c) Mobility-oriented parity analysis revealing areas in S.

America for which the model predictions are based strictly on extrapolation. [Colours

after Scarlet-Breasted Dacnis *Dacnis berlepschi* and Moss-backed Tanager *Bangsia*

edwardsi]

As more than one best model was selected, we used the median and range of all

replicates across parameters to consolidate results for the *P. torquans* complex. The

median of the selected models (Fig 2A) identified areas with different levels of habitat

suitability for the *P. torquans* complex across South America. Highly suitable areas

were concentrated on the central-east of Argentina and Uruguay, comprising the

Pampean and Chaco (eastern Chaco district) biogeographical provinces (Fig. S3, see in

Morrone, 2017). Suitability declined towards western Argentina, southern Brazil and

Paraguay. A broad area in central Chile was identified as suitable, although it may not

be accessible to the *P. torquans* complex by natural dispersion. The model showed

moderate uncertainty areas mainly throughout central-western Argentina (the Monte

biogeographical province), central Chile and the Andean range (Fig 2B and Fig. S3).

The uncertainty observed was mainly due to the variability among replicates (~95%)

(Fig. S3). The MOP analysis revealed that disparate areas in southern Chile and

northern South America would present model extrapolation (Fig 2C).

Discussion

In this study, we provided a model map of environmental suitability for the *P. torquans*

complex in South America and investigated what constrains its abiotic niche. Ecological

niche modelling is a powerful tool to identify macro-environmental variables defining

the environmental niche, and thus driving the geographical distribution of species. This

study took into consideration many of the recent methodological recommendations to

produce robust ecological niche models, as outlined by Fen*g et al.* (2019): filtering and

thinning occurrences (Boria *et al.*, 2014; Aiello-Lammens *et al.*, 2015); using an

accessible area for model calibration (Barve *et al.*, 2011); testing a number of candidate

sets of environmental variables and multiple combinations of model parameters

(features, regularization multipliers, and sets of variables); using multiple statistical

criteria (partial ROC, omission rates, and AICc) for model selection (Cobos *et al.*,

2019); replicating the model selected a number of times to determine that results are

consistent (Morales *et al.*, 2017); and accounting for model uncertainty in geographical

predictions (Owens *et al.*, 2013). As in any modelling approach, having a large sample

size is a central issue in ecological niche modelling. As suggested by van Proosdij *et al.*

(2016), the final sample size in our study after spatial thinning seems appropriate to

produce useful results. Moreover, selecting the best combination of parameters (features

and regularization multiplier) further improves our ability to select the most appropriate

model when dealing with a reduced number of occurrences (Morales *et al.*, 2017).

Among a number of causes driving uncertainty in niche modelling, the nature of the

used occurrence dataset might represent a concern to our modelling approach. As far as

we know, there are no published records of free-living adults of any *Philornis* species

trapped in the wild, with the sole exception of the introduced *P. downsi* in the

Galapagos Islands (Causton *et al.*, 2019). Therefore, we modelled the ecological niche

of the *P. torquans* complex from reported records of immature (parasitic) stages, as a

proxy for the presence of adults. To date, no *Philornis* species is known to migrate, thus

we assume that the adults are present year-round in the same area, as observed in *P.*

downsi (Causton *et al.*, 2019). Despite modelling the niche of the adult *P. torquans* flies

from its parasitic stage, we have no reason to suspect that the true geographical

distribution might differ among life stages.

We hypothesised that low temperatures and reduced precipitation will restrain the

occurrence of the *Philornis torquans* complex, as these abiotic parameters play a

predominant role in establishing the distribution limits of organisms and have already

been related with the intensity of parasitism by *Philornis* larvae (Dudaniec *et al.*, 2007;

Antoniazzi *et al.*, 2011; Manzoli *et al.*, 2013). Since insects are ectothermic, their rate of

development and survival can vary greatly in response to many biotic and abiotic

factors (Adams, 1979; Couret *et al.*, 2014). Winters are critical period for insects at

higher latitudes since overwintering conditions strongly affect their fitness (Irwin &

Lee, 2000). Indeed, our model identified the minimum temperature of coldest month

(Bio06) as the most relevant variable (Table 2 and Fig. 2), revealing a detrimental effect

in both sides of a suitable range of temperatures (Fig. S1).

Morphs suggested to belong to the *P. torquans* complex (Monje *et al.*, 2013) are

considered to be originated in southern South America (somewhere near the border

between Brazil and Uruguay - Löwenberg-Neto & De Carvalho, 2009). Representing

the southernmost species of the genus, and unlike the majority of *Philornis* species, the

P. torquans complex is adapted to colder, longer and drier winters; typical of temperate

areas. The observed data across presence sites of the *P. torquans* complex indicates that

minimum temperatures of the coldest month ranges between 3°C and 18°C. Therefore,

it is highly likely that the *P. torquans* complex evolved to overwinter as adults or pupae,

as known for other Diptera species (Protocalliphoridae, Culicidae, Muscidae) (Bellamy

& Reeves, 1963; Stiner, 1969; Skidmore, 1985; Sabrosky *et al.*, 1989; Nasci *et al.*,

2001). Despite being unknown, we infer that the *P. torquans* complex may overwinter

as adults, a well-known strategy for *Protocalliphora* blowflies (Sabrosky *et al.*, 1989)

and recently reported for *P. downsi* in Galapagos (Causton *et al.*, 2019). The former is

sustained in the fact that *Philornis* adults are known to survive over three months in

captivity (Lincango & Causton, 2008; Saravia-Pietropaolo *et al.*, 2018) and had been

trapped at the same territories all year round (Causton *et al.*, 2019).

On the other hand, insects adapted to cold winters may not perform well in mild winters

(Irwin *et al.*, 2001). Our model suggests that warm winters (signalled by Bio06) may be

limiting the northern distribution of the *P. torquans* complex (Fig. S2). Such scenario

seems plausible, as warmer winter temperatures may increase energy consumption,

(Tauber *et al.*, 1986; Harrington *et al.*, 2001; Irwin & Lee Jr, 2003; Sinclair, 2015).

Therefore, these individuals may not be able to reach the next breeding season or may

leave them without enough energy reserves for early reproduction (Harrington *et al.*,

2001).

As aforementioned, the developmental rate of insects is related to environmental

conditions, particularly as temperature impacts critical rate processes throughout

development and adulthood (Kingsolver & Huey, 2008). The temperature dependence

of ectotherms' developmental processes is commonly represented as the cumulative

sum of degree-time products required for the development of key life stages, such as

egg maturation or emergence from pupae (e.g., Kasap & Alten, 2005; Nieto *et al.*, 2006;

Gutierrez & Ponti, 2014; Harnden & Tomberlin, 2016; Yanmanee *et al.*, 2016). In these

regards, the mean temperature of the warmest quarter (Bio10) proved to be the second

most important variable (Table 2 and Fig. 2), showing a positive relation with sigmoidal

shape between predicted probability of presence and Bio10 (Fig. S1). This sigmoidal

shape implies a threshold temperature (apparently around 10°C), above which the

predicted probability of presence increases exponentially. The former makes sense

when considering that the increase of Bio10 might favour the development of

temperature-dependent non-parasitic stages, such as pupae maturation, which lately

relates with the abundance of adults. Otherwise, temperatures below the apparent

threshold seems to limit the distribution of the *P. torquans* complex, as evinced along

the Andean range (Fig. S2).

The third most relevant variable, the climatic moisture index (CMI), resulted to be the

most influential factor in most of Argentina, northern Chile and coastal Peru (Fig. S2).

The CMI is an aggregate measure of potential water availability imposed solely by

climate, ranging from –1 to +1, with wet climates showing positive values, dry climates

negative (Vörösmarty *et al.*, 2005). The CMI values observed through the actual

distribution of the *P. torquans* complex (mean -0.25, sd 0.19) indicate that the species

must endure mostly a dry environment when compared with most of South America. As

reported for species laying eggs into a relatively dry environment (Grzywacz *et al.*,

2012), the eggs of the *P. torquans* morphs present a small portion of the egg surface that

functions as a plastron (Patitucci *et al.*, 2017), a respiratory structure that facilitates

oxygen income minimizing water loss (Hinton, 1960). The reduced size of the plastron

of the *P. torquans* morph (Patitucci *et al.*, 2017) may have evolved as an adaptation

allowing the radiation to the drier conditions typical of central Argentina (see

Löwenberg-Neto & De Carvalho, 2009).

Opposed to our prediction, the ecological niche model suggests that precipitation has no

direct influence on limiting the environmental suitability of the *P. torquans* complex.

The precipitation of the driest month (Bio14) and the precipitation of the wettest quarter

(Bio16) had a poor contribution to the model performance (Table 2), and showed a

weak explanatory power in the jackknife analysis (Fig. 2). Despite it is clear that

humidity and/or moisture are relevant environmental factors determining the *P.*

torquans complex distribution, as reflected by CMI and supported by this model, the

proximal influence of precipitation seems to be mainly related with the intensity of

parasitism by *Philornis* species, as previously reported (Arendt, 1985, 2000; Dudaniec

et al., 2007; Antoniazzi *et al.*, 2011; Manzoli *et al.*, 2013).

The outcome of the geographical projection of the environmental niche model was a

map of the potential distribution of the *P. torquans* complex in South America.

According to our results, the *P. torquans* complex is potentially distributed across

central-eastern Argentina, mostly restricted to the Pampean and Chaco biogeographical

provinces, and declining with some uncertainty in the surrounding Monte

biogeographical province (Fig. 3 and Fig. S3). The later makes sense due to the

historical relationship among these three biogeographical provinces of the Chacoan

dominion (Morrone, 2017). Indeed, the Chacoan and the Monte provinces are closely

related (revised in Morrone, 2017), and a sequence of biotic impoverishment from the

Chaco province to the Pampean province, with the Monte province being intermediate

between both, has been suggested (Cabrera, 1976; Morrone, 2017). Regarding the broad

area predicted as suitable in central Chile, and despite its relative high uncertainty, the

Andean range might represent a pronounced topographic barrier impeding the natural

dispersion of the *P. torquans* complex to this area.

In conclusion, we present the first study to identify macro-environmental variables

driving the geographical distribution of a *Philornis* species by evaluating their

contribution to its' environmental niche. Our model is of relevance as a tool for bird

conservation, and it integrates current methodological recommendations to produce a

robust ecological niche prediction; thus representing a good reference for future work

on the distribution of this parasite genus.

Acknowledgements

We greatly acknowledge the contribution made by Salvador S.A., who kindly provided

the geographic coordinates from the sites mentioned in Salvador & Bodrati (2013). We

also thank Dra. Bulgarella for the critical reading of this manuscript.

This work was funded by Consejo Nacional de Investigaciones Científicas y Técnicas

[PIP11220100100261; PIP 11220130100561CO]; and Universidad Autónoma de Entre

Ríos [PIDAC S013805/15]. The funders had no role in study design, data collection and

analysis, decision to publish, or preparation of the manuscript.

Contribution of authors

PFC and MAQ participated in the design and coordination of the study, and drafted the

manuscript; PFC performed the data analysis and modelling; AP carried out field

surveys and revised the existing literature; LM and PMB provided financial support and

critically reviewed the manuscript. All authors revised the manuscript and gave final

approval for publication.

Competing interests

We have no competing interests.

References

Adams, T.S. (1979) The reproductive physiology of the Screwworm, *Cochliomyia*

hominivorax (Diptera: Calliphoridae) II. Effect of constant temperatures on oogenesis.

Journal of Medical Entomology, **15**, 484–487.

Aiello-Lammens, M.E., Boria, R.A., Radosavljevic, A., Vilela, B. & Anderson, R.P.

(2015) spThin: An R package for spatial thinning of species occurrence records for use

in ecological niche models. *Ecography*, **38**, 541–545.

Antoniazzi, L.R., Manzoli, D.E., Rohrmann, D., Saravia, M.J., Silvestri, L. &

Beldomenico, P.M. (2011) Climate variability affects the impact of parasitic flies on

Argentinean forest birds. *Journal of Zoology*, **283**, 126–134.

Arendt, W.J. (1985) Philornis Ectoparasitism of Pearly-eyed Thrashers. I. Impact on

Growth and Development of Nestlings. *The Auk*, **102**, 270–280.

Arendt, W.J. (2000) Impact of nest predators, competitors, and ectoparasites on Pearly-

eyed Thrashers, with comments on the potential implications for Puerto Rican Parrot

recovery. *Ornitologia Neotropical*, 13–63.

Barve, N., Barve, V., Jiménez-Valverde, A., Lira-Noriega, A., Maher, S.P., Peterson,

A.T., *et al.* (2011) The crucial role of the accessible area in ecological niche modeling

and species distribution modeling. *Ecological Modelling*, **222**, 1810–1819.

Baumgartner, J., Wilson, P. & Esperón-Rodríguez, M. (2017) rmaxent: Tools for

working with Maxent in R. R package version 0.8.3.9000.

Beier, C. & Fontana, C. (2019) Breeding biology of the endangered Yellow Cardinal

Gubernatrix cristata in Brazil. *Revista Brasileira de Ornitologia - Brazilian Journal of*

Ornithology, **27**.

Bellamy, R.E. & Reeves, W.C. (1963) The winter biology of *Culex tarsalis* (Diptera:

Culicidae) in Kern County, California. *Annals of the Entomological Society of America*,

56, 314–323.

BirdLife International. (2018) *Gubernatrix cristata* [WWW Document]. *The IUCN Red*

List of Threatened Species. URL http://dx.doi.org/10.2305/IUCN.UK.2018-

2.RLTS.T22721578A131888081.en [accessed on 2018].

Boria, R.A., Olson, L.E., Goodman, S.M. & Anderson, R.P. (2014) Spatial filtering to

reduce sampling bias can improve the performance of ecological niche models.

Ecological Modelling, **275**, 73–77.

Bronstein, J.L. (1994) Conditional outcomes in mutualistic interactions. *Trends in*

Ecology and Evolution, **9**, 214–217.

Bulgarella, M., Quiroga, M.A. & Heimpel, G.E. (2019) Additive negative effects of

Philornis nest parasitism on small and declining Neotropical bird populations. *Bird*

Conservation International, **29**, 339–360.

Cabrera, A.L. (1976) Regiones fitogeográficas argentinas. In *Enciclopedia Argentina de*

Agricultura y Jardinería (ed. by Kugler, W.F.). ACME, Buenos Aires, pp. 1–85.

Cadena, D. & Zapata, F. (2019) tanagR: Tanager-Inspired Color Palettes. R package

version 0.1.0.

Causton, C.E., Moon, R.D., Cimadom, A., Boulton, R.A., Cedeño, D., Lincango, M.P.,

et al. (2019) Population dynamics of an invasive bird parasite, *Philornis downsi*

(Diptera: Muscidae), in the Galapagos Islands. *PLOS ONE*, **14**, 1–21.

Cobos, M.E., Peterson, A.T., Barve, N. & Osorio-Olvera, L. (2019) kuenm: an R

package for detailed development of ecological niche models using Maxent. *PeerJ*, **7**,

e6281.

Couret, J., Dotson, E. & Benedict, M.Q. (2014) Temperature, larval diet, and density

effects on development rate and survival of *Aedes aegypti* (Diptera: Culicidae). *PLOS*

ONE, **9**, e87468.

Couri, M. (1999) Myiasis caused by obligatory parasites. Ia. *Philornis* Meinert

(Muscidae). In *Myiasis in man and animals in the neotropical region* (ed. by

Guimarães, J.H. & Papavero, N.). Editora Plêiade, pp. 51–70.

Couri, M.S., Antoniazzi, L.R., Beldomenico, P. & Quiroga, M. (2009) Argentine

Philornis Meinert species (Diptera: Muscidae) with synonymic notes. *Zootaxa*, **62**, 52–

62.

Domínguez, M., Reboreda, J.C. & Mahler, B. (2015) Impact of Shiny Cowbird and

botfly parasitism on the reproductive success of the globally endangered Yellow

Cardinal *Gubernatrix cristata*. *Bird Conservation International*, **25**, 294–305.

Dudaniec, R.Y., Fessl, B. & Kleindorfer, S. (2007) Interannual and interspecific

variation in intensity of the parasitic fly, *Philornis downsi*, in Darwin's finches.

Biological Conservation, **139**, 325–332.

Dudaniec, R.Y. & Kleindorfer, S. (2006) Effects of the parasitic flies of the genus

Philornis (Diptera: Muscidae) on birds. *Emu*, **106**, 13–20.

Dudaniec, R.Y., Kleindorfer, S. & Fessl, B. (2006) Effects of the introduced

ectoparasite *Philornis downsi* on haemoglobin level and nestling survival in Darwin's

Small Ground Finch (*Geospiza fuliginosa*). *Austral Ecology*, **31**, 88–94.

Elith, J., Graham, C.H., Anderson, R.P., Dudík, M., Ferrier, S., Guisan, A., *et al.* (2006)

Novel methods improve prediction of species' distributions from occurrence data.

Ecography, **29**, 129–151.

Elith, J., Kearney, M. & Phillips, S. (2010) The art of modelling range-shifting species.

Methods in Ecology and Evolution, **1**, 330–342.

Escobar, L.E., Lira-Noriega, A., Medina-Vogel, G. & Townsend Peterson, A. (2014)

Potential for spread of the white-nose fungus (*Pseudogymnoascus destructans*) in the

Americas: Use of Maxent and NicheA to assure strict model transference. *Geospatial*

Health, **9**, 221–229.

Feng, X., Park, D.S., Walker, C., Peterson, A.T., Merow, C. & Papeş, M. (2019) A

checklist for maximizing reproducibility of ecological niche models. *Nature Ecology &*

Evolution, **3**, 1382–1395.

Fessl, B., Sinclair, B.J. & Kleindorfer, S. (2006) The life-cycle of *Philornis downsi*

(Diptera: Muscidae) parasitizing Darwin's finches and its impacts on nestling survival.

Parasitology, **133**, 739–747.

Grzywacz, A., Szpila, K. & Pape, T. (2012) Egg morphology of nine species of Pollenia

Robineau-Desvoidy, 1830 (Diptera: Calliphoridae). *Microscopy Research and*

Technique, **75**, 955–967.

Gutierrez, A.P. & Ponti, L. (2014) The new world screwworm: Prospective distribution

and role of weather in eradication. *Agricultural and Forest Entomology*, **16**, 158–173.

Harnden, L.M. & Tomberlin, J.K. (2016) Effects of temperature and diet on black

soldier fly, *Hermetia illucens* (L.) (Diptera: Stratiomyidae), development. *Forensic*

Science International, **266**, 109–116.

Harrington, R., Fleming, R.A. & Woiwod, I.P. (2001) Climate change impacts on insect

management and conservation in temperate regions: can they be predicted? *Agricultural*

and Forest Entomology, **3**, 233–240.

Hayes, C.D., Hayes, T.I., McClure, C.J.W., Quiroga, M., Thorstrom, R.K., Anderson,

D.L., *et al.* (2018) Native parasitic nest fly impacts reproductive success of an island-

endemic host. *Animal Conservation*, **22**, 157–164.

Hernandez, P.A., Graham, C.H., Master, L.L., Albert, D.L., Deborah, L. & Albert, D.L.

(2006) The effect of sample size and species characteristics on performance of different

species distribution modeling methods. *Ecography*, **29**, 773–785.

Hinton, H.E. (1960) The chorionic plastron and its role in the eggs of the Muscinae

(Diptera). *Journal of Cell Science*, **s3-101**, 313–332.

Irwin, J.T., Bennett, V.A. & Lee Jr, R.E. (2001) Diapause development in frozen larvae

of the Goldenrod Gall fly, *Eurosta solidaginis* fitch (Diptera: Tephritidae). *Journal of*

Comparative Physiology B, **171**, 181–188.

Irwin, J.T. & Lee Jr, R.E. (2003) Cold winter microenvironments conserve energy and

improve overwintering survival and potential fecundity of the Goldenrod Gall fly,

Eurosta solidaginis. *Oikos*, **100**, 71–78.

Irwin, J.T. & Lee, R.E. (2000) Mild winter temperatures reduce survival and potential

fecundity of the goldenrod gall fly, *Eurosta solidaginis* (Diptera: Tephritidae). *Journal*

of Insect Physiology, **46**, 655–661.

Karger, D.N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R.W., *et al.*

(2017) Climatologies at high resolution for the earth's land surface areas. *Scientific*

Data, **4**, 170122.

Kasap, O.E. & Alten, B. (2005) Laboratory estimation of degree-day developmental

requirements of *Phlebotomus papatasi* (Diptera: Psychodidae). *Journal of Vector*

Ecology, **30**, 328–333.

Kingsolver, J.G. & Huey, R.B. (2008) Size, temperature, and fitness: three rules.

Evolutionary Ecology Research, **10**, 251–268.

Kinsella, J. & Winegarner, C. (1974) Notes on the life history of *Neomusca porteri*

(Dodge), parasitic on nestlings of the great crested flycatcher in Florida. *Journal of*

Medical Entomology, **11**, 633.

Kleindorfer, S. & Dudaniec, R.Y. (2016) Host-parasite ecology, behavior and genetics:

A review of the introduced fly parasite *Philornis downsi* and its Darwin's finch hosts.

BMC Zoology.

Koop, J.A.H., Huber, S.K., Laverty, S.M. & Clayton, D.H. (2011) Experimental

demonstration of the fitness consequences of an introduced parasite of Darwin's

finches. *PLoS ONE*, **6**, e19706.

Kramer-Schadt, S., Niedballa, J., Pilgrim, J.D., Schröder, B., Lindenborn, J., Reinfelder,

V., *et al.* (2013) The importance of correcting for sampling bias in MaxEnt species

distribution models. *Diversity and Distributions*, **19**, 1366–1379.

Krivosheina, N.P., Morozov, N.S. & Khudyakov, V. V. (2017) On the biology of the

bird parasite *Neottiophilum praeustum* (Meigen, 1826) (Diptera, Neottiophilidae).

Entomological Review, **97**, 1048–1052.

Labud, V.A., Semenas, L.G. & Laos, F. (2003) Diptera of sanitary importance

associated with composting of biosolids in Argentina. *Revista de Saude Publica*, **37**,

722–728.

Lincango, P. & Causton, C. (2008) *Crianza en cautiverio de* Philornis downsi*, en las*

Islas Galápagos.

Little, S.E. (2008) Myasis in wild birds. In *Parasitic diseases of wild birds* (ed. by

Atkinson, C.T., Thomas, N.J. & Hunter, D.B.). Wiley Blackwell, pp. 546–556.

Löwenberg-Neto, P. (2008) The structure of the parasite-host interactions between

Philornis (Diptera: Muscidae) and neotropical birds. *Journal of Tropical Ecology*, **24**,

575–580.

Löwenberg-Neto, P. & Carvalho, C.J.B. De. (2009) Areas of endemism and spatial

diversification of the Muscidae (Insecta: Diptera) in the Andean and Neotropical

regions. *Journal of Biogeography*, **36**, 1750–1759.

Manzoli, D.E., Antoniazzi, L.R., Saravia, M.J., Silvestri, L., Rorhmann, D. &

Beldomenico, P.M. (2013) Multi-level determinants of parasitic fly infection in forest

passerines. *PLoS ONE*, **8**, e67104.

Manzoli, D.E., Saravia-Pietropaolo, M.J., Antoniazzi, L.R., Barengo, E., Arce, S.I.,

Quiroga, M.A., *et al.* (2018) Contrasting consequences of different defence strategies in

a natural multihost- parasite system. *International Journal of Parasitology*.

Merow, C., Smith, M.J. & Silander, J.A. (2013) A practical guide to MaxEnt for

modeling species' distributions: What it does, and why inputs and settings matter.

Ecography, **36**, 1058–1069.

Monje, L.D., Quiroga, M., Manzoli, D.E., Couri, M.S., Silvestri, L., Venzal, J.M., *et al.*

(2013) Sequence analysis of the internal transcribed spacer 2 (ITS2) from *Philornis*

seguyi (García, 1952) and *Philornis torquans* (Nielsen, 1913) (Diptera: Muscidae).

Systematic Parasitology, **86**, 43–51.

Morales, N.S., Fernández, I.C. & Baca-González, V. (2017) MaxEnt's parameter

configuration and small samples: are we paying attention to recommendations? A

systematic review. *PeerJ*, **5**, e3093.

Morrone, J.J. (2017) *Neotropical biogeography. Regionalization and evolution*. CRC

Press, Inc.

Nasci, R.S., Savage, H.M., White, D.J., Miller, J.R., Cropp, B.C., Godsey, M.S., *et al.*

(2001) West Nile virus in overwintering Culex mosquitoes, New York City, 2000.

Emerging Infectious Diseases, **7**, 742–744.

Nieto, P., Malone, J.B. & Bavia, M.E. (2006) Ecological niche modeling for visceral

leishmaniasis in the state of Bahia, Brazil, using genetic algorithm for rule-set

prediction and growing degree day-water budget analysis. *Geospatial health*, **1**, 115–

126.

Oniki, Y. (1983) Notes on fly (Muscidae) parasitism of nestlings of South American

birds. *Le Gerfaut*, **73**, 281–286.

Owens, H.L., Campbell, L.P., Dornak, L.L., Saupe, E.E., Barve, N., Soberón, J., *et al.*

(2013) Constraints on interpretation of ecological niche models by limited

environmental ranges on calibration areas. *Ecological Modelling*, **263**, 10–18.

Patitucci, L.D., Quiroga, M., Couri, M.S. & Saravia-Pietropaolo, M.J. (2017)

Oviposition in the bird parasitic fly *Philornis torquans* (Nielsen, 1913) (Diptera:

Muscidae) and eggs' adaptations to dry environments. *Zoologischer Anzeiger*, **267**, 15–

20.

Petrescu, A. & Adam, C. (2000) *Carnus hemapterus* Nitzsch, 1818 (Diptera: Carnidae)

parasite on *Merops apiaster* L.(Aves: Meropidae) in southern Romania. *Travaux du*

Muséum National d'Histoire Naturelle "Grigore Antipa," **42**, 221–224.

Phillips, S.J., Anderson, R.P. & Schapire, R.E. (2006) Maximum entropy modeling of

species geographic distributions. *Ecological Modelling*, **190**, 231–259.

Proosdij, A.S.J. van, Sosef, M.S.M., Wieringa, J.J. & Raes, N. (2016) Minimum

required number of specimen records to develop accurate species distribution models.

Ecography, **39**, 542–552.

Quiroga, M.A., Monje, L.D., Arrabal, J.P. & Beldomenico, P.M. (2016) Datos

moleculares nuevos sobre *Philornis* (Diptera: Muscidae) subcutáneas del sur de

Sudamérica sugieren la existencia de un complejo de especies. *Revista Mexicana de*

Biodiversidad, **87**, 1383–1386.

Quiroga, M.A. & Reboreda, J.C. (2012) Lethal and sublethal effects of Botfly

(*Philornis seguyi*) parasitism on House Wren nestlings. *The Condor*, **114**, 197–202.

R Core Team. (2018) R: A language and environment for statistical computing.

Raghavan, R.K., Peterson, A.T., Cobos, M.E., Ganta, R. & Foley, D. (2019) Current

and future distribution of the Lone Star Tick, *Amblyomma americanum* (L .) (Acari:

Ixodidae) in North America. *PLoS ONE*, **14**, e0209082.

Sabrosky, C.W., Bennett, G.F., Whitworth, T.L. & others. (1989) *Bird blow flies*

(Protocalliphora) in North America (Diptera: Calliphoridae) with notes on Palearctic

species. Smithsonian Institution Press, Washington, D.C.

Saravia-Pietropaolo, M.J., Arce, S.I., Manzoli, D.E., Quiroga, M. & Beldomenico, P.M.

(2018) Aspects of the life cycle of the avian parasite *Philornis torquans* (Diptera:

Muscidae) under laboratory rearing conditions. *The Canadian Entomologist*, **150**, 317–

325.

Showler, A.T. & Osbrink, W.L.A. (2015) Stable Fly, *Stomoxys calcitrans* (L.), dispersal

and governing factors. *International Journal of Insect Science*, **7**, IJIS.S21647.

Silvestri, L., Antoniazzi, L.R., Couri, M.S., Monje, L.D. & Beldomenico, P.M. (2011)

First record of the avian ectoparasite *Philornis downsi* Dodge & Aitken, 1968 (Diptera:

Muscidae) in Argentina. *Systematic Parasitology*, **80**, 137–140.

Sinclair, B.J. (2015) Linking energetics and overwintering in temperate insects. *Journal*

of Thermal Biology, **54**, 5–11.

Skidmore, P. (1985) *The biology of the Muscidae of the World*. Dr W. Junk Publishers,

Dordrecht, The Netherlands.

Stiner, F. (1969) Overwintering by *Protocalliphora metallica* (Diptera: Calliphoridae).

Annals of the Entomological Society of America, **62**, 1205–1206.

Tauber, M.J., Tauber, C.A. & Masaki, S. (1986) *Seasonal adaptations of insects*.

Oxford University Press.

Title, P.O. & Bemmels, J.B. (2018) ENVIREM: an expanded set of bioclimatic and

topographic variables increases flexibility and improves performance of ecological

niche modeling. *Ecography*, **41**, 291–307.

Vignali, S., Barras, A. & Braunisch, V. (2019) SDMtune: Species Distribution Model

Selection.

Vörösmarty, C.J., Douglas, E.M., Green, P.A. & Revenga, C. (2005) Geospatial

indicators of emerging water stress: an application to Africa. *AMBIO: A Journal of the*

Human Environment, **34**, 230–236.

Warren, D.L., Seifert, S.N., Warren, D.L. & Seifert, S.N. (2016) Ecological niche

modeling in Maxent: the importance of model complexity and the performance of

model selection criteria. *Ecological Applications*, **21**, 335–342.

Yanmanee, S., Husemann, M., Benbow, M.E. & Suwannapong, G. (2016) Larval

development rates of *Chrysomya rufifacies* Macquart, 1842 (Diptera: Calliphoridae)

within its native range in South-East Asia. *Forensic Science International*, **266**, 63–67.

Young, B.E. (1993) Effects of the parasitic botfly Philornis carinatus on nestling house

wrens, *Troglodytes aedon*, in Costa Rica. *Oecologia*, **93**, 256–262.

Zumpt, F. (1965) *Myiasis in man and animals in the Old World. A textbook for*

physicians, veterinarians and zoologists. London, Butterworth. Butterworth & Co Ltd.,

London, United Kingdom.