Environmental variables determining the presence of an avian parasite: the case of the *Philornis torquans* complex (Diptera: Muscidae) in South America

Pablo F. Cuervo[1], Alejandro Percara[1], Lucas Monje[1], Pablo M. Beldomenico[1], Martín A. Quiroga[1,2,3]

[1]Laboratorio de Ecología de Enfermedades, Instituto de Ciencias Veterinarias del Litoral (ICIVET - Litoral), Universidad Nacional del Litoral (UNL) / Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Esperanza, Santa Fe, Argentina

[2]The Peregrine Fund, Boise, Idaho, USA

[3]Facultad de Ciencia y Tecnología (FCyT), Universidad Autónoma de Entre Ríos (UADER), Oro Verde, Entre Ríos, Argentina

Running title: *P. torquans* complex in South America

Corresponding author: Martín A. Quiroga. Laboratorio de Ecología de Enfermedades, Instituto de Ciencias Veterinarias del Litoral (ICIVET - Litoral), Universidad Nacional del Litoral (UNL) / Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Esperanza, Santa Fe, Argentina. The Peregrine Fund, Boise, Idaho, USA.[3]Facultad de Ciencia y Tecnología (FCyT), Universidad Autónoma de Entre Ríos (UADER), Oro Verde, Entre Ríos, Argentina. mquiroga06@hotmail.com

Abstract

1. *Philornis* flies are exclusive parasites of birds and their nests, being some of them of conservation concern. Despite many studies addressed different aspects of this problem, none have focused on the geographical distribution of these species; relevant information of conservation concern.

2. In this study, besides providing a model map of environmental suitability of the *P. torquans* complex in South America, we also contributed with data related to the constraints of its abiotic niche.

3. We provided baseline data (supplementary information) on sites where *P. torquans* complex is present in southern South America, valuable for future *Philornis* distribution modelling efforts.

4. According to our results, the *P. torquans* complex is potentially distributed across central-eastern Argentina, mostly restricted to the Pampean and Chaco (eastern Chaco district) biogeographical provinces.

5. The potential distribution of the *P. torquans* complex is mainly related to the "minimum temperature of the coldest month" and "mean temperature of warmest quarter". In addition, moisture/aridity seems to be exerting additional constraints.

6. Our work is the first one to identify macro-environmental variables driving the geographical distribution of a *Philornis* species by evaluating their contribution to its' environmental niche.

7. Besides the relevance of our model as a tool for bird's conservation, our model integrates the most up-to-date methodological recommendations to produce a robust ecological niche; thus representing a methodological reference for future work on the distribution of this group of parasites.

50   Keywords: *Philornis torquans* complex; distribution model; southern South America;

51   MaxEnt

Introduction

The infestation of live vertebrates with dipteran larvae which feed on the host's dead or living tissue and body fluids is known as myiasis (Zumpt, 1965). In birds, myiasis is mainly caused by species of the genera *Protocalliphora* [Hough, 1899 -Diptera: Calliphoridae], *Passeromyia* [Rodhain & Villeneuve, 1915 - Diptera: Muscidae] and *Philornis* [Meinert, 1890 - Diptera: Muscidae] (Little, 2008). The only of these genera present in the Neotropical regions is *Philornis*, comprising about 50 described species (Couri, 1999; Dudaniec & Kleindorfer, 2006). *Philornis* adults are free living, while larvae are strictly associated with bird nestlings, establishing different types of associations: coprophagous, semi-haematophagous or subcutaneous (Dudaniec & Kleindorfer, 2006).

Despite more than 250 bird species have been reported as *Philornis* hosts in South America (Couri, 1999; Dudaniec & Kleindorfer, 2006; Löwenberg-Neto, 2008; Antoniazzi *et al.*, 2011), there are still huge gaps of information on the parasitism by *Philornis* and its effect on bird hosts; especially concerning vulnerable and endangered species. Impact of *Philornis* spp. on hosts ranges from subtle (Kinsella & Winegarner, 1974; Oniki, 1983; Young, 1993; Fessl *et al.*, 2006) to strong and lethal (Dudaniec *et al.*, 2006, 2007; Quiroga & Reboreda, 2012), and depends on the interaction between parasite and host (Bronstein, 1994; Manzoli *et al.*, 2018).

While in healthy populations effects of nest parasitism could sometimes be negligible; in geographically restricted, fragmented and small sized populations, the effects of parasitism by *Philornis* could be devastating (Bulgarella *et al.*, 2018). Indeed, evidence that this parasite can be a serious threat to conservation is the severe impact caused by *Philornis downsi* [Dodge & Aitkens 1968] on Darwin's finches (i.e.: Koop *et al.*, 2011) and by *P. pici* [Macquart, 1854] on Ridgway's Hawks (*Buteo ridgwayi* - Hayes *et al.*,

2019). On already small or declining populations, infestation with *Philornis* spp. seems

to be exerting additional pressure (Bulgarella *et al.*, 2018), as it may cause up to 75% of

nestlings mortality (Fessl *et al.*, 2006; Hayes *et al.*, 2018); thus, pushing these

populations to the edge of extinction.

For all the above, there is need to pursue research on the occurrence of these parasites.

One aspect of particular interest and, so far little investigated, is the environmental

dimensions determining the geographical distribution of *Philornis* spp. Such knowledge

is essential for a better comprehension of the ecology of the genus, and may be useful to

identify *Philornis*-free areas that are prone to be colonized, or to identify locations with

high presence of these flies that might threaten populations of endangered bird species,

or to locate habitats with low risk of *Philornis* occurrence for the re-introduction of bird

species or reinforcement of their populations.

The present study is aimed to identify macro-environmental variables driving the

geographical distribution of *Philornis* species in South America by assessing their

contribution to its' environmental niche. For an initial insight on the role of

environmental dimensions on the geographic distribution of *Philornis*, the focus was

placed on *P. torquans* and *P. seguyi*, a fairly well-known complex of species of the

genus. As temperature and rainfall are presumed to be the most relevant abiotic

parameters affecting *Philornis* sp. life cycle (Dudaniec *et al.*, 2007; Antoniazzi *et al.*,

2011; Manzoli *et al.*, 2013), we hypothesise that low temperatures and reduced

precipitation will restrain the occurrence of these *Philornis* species.

Materials and Methods

Study species

Philornis torquans [Nielsen, 1913] and *Philornis seguyi* [Garcia, 1952] are the

subcutaneous species most commonly reported in southern South America (Couri *et al.*,

2009; Silvestri *et al.*, 2011), where eight *Philornis* species have been described but only

half of them were recognized as valid (namely, *P. torquans* , *P. seguyi* , *P. blanchardi*

[Garcia, 1952] and *P. downsi*). However, specimens morphologically identified either

as *P. torquans* or *P. seguyi* proved to be undistinguishable at a molecular level (Monje

et al., 2013), reason why it was suggested that the subcutaneous *Philornis* morphotypes

resembling *P. torquans* or *P. seguyi* in southern South America should be treated as

Philornis torquans complex (Quiroga *et al.*, 2016). This complex of parasitic flies was

selected because: i) its biology and ecoepidemiology have been studied for more than a

decade (Antoniazzi *et al.*, 2011; Quiroga & Reboreda, 2012; Manzoli *et al.*, 2013, 2018;

Saravia-Pietropaolo *et al.*, 2018); ii) this is the group of flies with the highest number of

confirmed morphology-based detections; and iii) could potentially affect the endangered

Yellow Cardinal (*Gubernatrix cristata* - BirdLife International, 2018), reported as a

host of an unidentified *Philornis* species (Domínguez *et al.*, 2015; Beier & Fontana,

2019) in the vicinity of *P. torquans* complex records.

Species occurrence data

Occurrence data of *Philornis* species was obtained from field surveys conducted in

Argentina and from existing literature. In every case, *Philornis* adults were obtained

from larvae or pupae recovered from nestlings or nests remains, and were identified to

species level mostly following Couri (1999). Due to the absence of occurrence data of

free-living adults, information obtained from immature *Philornis* stages represents a

valid surrogate to assess the environmental suitability for the species. The specimens

identified based on morphology as *P. torquans* or *P. seguyi* were considered as

members of the *Philornis torquans* complex (see Quiroga *et al.*, 2016), which accounted

for 80 occurrence records. After removing duplicate localities, the data was reduced to

34 occurrence sites (Table S1). Since spatial autocorrelation in environmental

information and consequent non-independence of occurrence data could bias predictions

(Kramer-Schadt *et al.*, 2013), we spatially filtered occurrence sites preserved for

calibration with the package "spThin" in R Statistical Software 3.5.0 (Aiello-Lammens

et al., 2015; R Core Team, 2018). Spatial thinning of occurrence records provides an

easy-to-implement and relatively straightforward method to alleviate the effects of

sampling bias (Boria *et al.*, 2014; Aiello-Lammens *et al.*, 2015). To attain a uniform

distribution of occurrences and to avoid excessive clustering within any particular

region, we explored different filtering distances in sequential steps, visually inspecting

results. Once a satisfactory distribution of occurrence data was obtained (i.e., when

spatial clustering of records was no longer evident), we ended with 18 occurrence sites

for the *P. torquans* complex (Fig. 1). We split the *P. torquans* complex locations

randomly in equal quarts: one set of locations for calibration (75%) and another for

evaluation during model calibration (25%).

Figure 1. Geographic coordinates representing locations of *Philornis torquans* complex and accessible area (M) used in the construction of distribution models.

Environmental data and selection of variables

To characterize climatic conditions meaningful to biological species, we used the so-called "bioclimatic" variables based on WorldClim from the CHELSA database (www.chelsa-climate.org, Karger *et al.*, 2017). In addition, we included the climatic variables from the ENVIREM database, derived from and complementing the bioclimatic variables from WorldClim (www.envirem.github.io, Title & Bemmels, 2018). We excluded nine data layers (mean temperature of wettest quarter, mean temperature of driest quarter, precipitation of warmest quarter, precipitation of driest quarter, mean PET of coldest quarter, mean PET of warmest quarter, mean PET of driest quarter, mean PET of wettest quarter, and count of the number of months with mean temperature greater than 10°C), as these variables include artefacts that create abrupt differences between neighbouring pixels (Escobar *et al.*, 2014; Raghavan *et al.*, 2019). We coarsed environmental data to 10' (~17 x ~17 km) spatial resolution, which approximately matches the uncertainty inherent in the occurrence data.

As a first step and in order to test our hypothesis, we selected a set of five variables based on current knowledge on the biology of *Philornis* spp.: Bio10 = mean temperature of warmest quarter, Bio06 = minimum temperature of coldest month, Bio14 = precipitation of driest month, Bio16 = precipitation of wettest quarter (the four of them from CHELSA), and CMI = climatic moisture index (a metric of relative wetness and aridity, from ENVIREM). The "mean temperature of warmest quarter" (Bio10) was selected because this variable is related to the sum of degree days available for development, a usual method used to predict the life stages of other Dipteran species (e.g., Kasap & Alten, 2005; Nieto *et al.*, 2006; Gutierrez & Ponti, 2014; Harnden &

Tomberlin, 2016; Yanmanee *et al.*, 2016). The "minimum temperature of the coldest

month" (Bio06) was selected due to its negative effect on developmental times and

survival of pupae and adults (Sinclair, 2015), therefore arguably limiting *Philornis*

distribution. Furthermore, Manzoli *et al.* (2013) found that preceding increases in

rainfall were highly correlated with the abundance of *Philornis torquans* parasitic

stages. This aspect was taken into account by including the "precipitation of driest

month" (Bio14) and the "precipitation of the wettest quarter" (Bio16). A "climatic

moisture index" (CMI) was included as an integrative marker of environmental

conditions, strongly related to temperature and precipitation.

Additionally, we created six other sets of variables to be considered in the evaluation

and calibration of models. The variables from the original set of data were grouped in

three different sets: i) a dataset including only the 15 bioclimatic variables from

CHELSA (referred to hereafter as the *bioclim* dataset); ii) a dataset using the *bioclim*

dataset plus the 11 climatic ENVIREM variables (hereafter referred to as the *climatic*

dataset); and iii) a dataset using the *climatic* dataset plus the two topographic variables

from ENVIREM (hereafter referred to as the *environmental* dataset). Then, we reduced

model complexity and collinearity by selecting two simpler sets of variables from each

of these three datasets, based on the calibration of preliminary Maxent models (details

below). For the selection of these simpler sets of variables, we applied two procedures

available in the R package "SDMtune" (Vignali *et al.*, 2019): removal of highly

correlated variables ($R > 0.7$), and removal of variables with low importance for the

model performance (percent contribution $< 5\%$). Finally, for the calibration of models,

we retained six sets of environmental variables: i) a set based on current knowledge on

the biology of *Philornis* spp. (Set 1); ii) three sets of uncorrelated variables from the

bioclim, *climatic* and *environmental* datasets (Sets 2a, 3a and 4); and iii) two sets of

relevant variables selected from the *bioclim* and *climatic* datasets (Sets 2b and 3b) (the

selection from the *climatic* and *environmental* datasets resulted in two equivalent sets of

relevant variables) (Datasets in Table S2).

Niche modelling and assessment of variables importance

We used MaxEnt 3.4.1 for developing models (Phillips *et al.*, 2006). MaxEnt is a

machine-learning method that estimates the species potential geographic distribution by

finding the probability distribution of maximum entropy (closest to uniform), subject to

the constraint of the expected values of the environmental predictors. MaxEnt was

developed to use presence-only data by contrasting presences against background

locations (Phillips *et al.*, 2006; Merow *et al.*, 2013), and has shown to outperform other

algorithms, even when applied to small data sets like ours (Elith *et al.*, 2006; Hernandez

et al., 2006; van Proosdij *et al.*, 2016).

To minimize the impact of assumptions about absences from areas that are not

accessible to the species (Barve *et al.*, 2011), we restricted model calibration to the area

to which the species likely had access via dispersal. We defined the adequate accessible

area (M) as a maximum area within 1000 km of any of the occurrences, based on the

average geographic distance between a centroid point and all the occurrences (~730

km), the assisted dispersion of *Philornis downsi* to the Galapagos Islands (Kleindorfer

& Dudaniec, 2016) and the dispersal capacity of other Muscidae species (Labud *et al.*,

2003; Showler & Osbrink, 2015).

The complexity of models built with MaxEnt can be adjusted with the inclusion of

additional feature classes (i.e. transformations of the original predictor variables) that

allow for increasingly complex models, as well as with a regularization multiplier that

contributes to select those features and reduces overfitting (Merow *et al.*, 2013; Warren

et al., 2016). The preliminary models for each of the three data sets (the *bioclim*,

climatic and *environmental* datasets), later used for variable selection, were calibrated

using the "kuenm" R package (Cobos *et al.*, 2019). To determine the optimal model

complexity, we explored all combinations of [a] 17 values of the regularization

parameter (0.1–1.0 at intervals of 0.1, 2–6 at intervals of 1, and 8 and 10), and [b] nine

sets of potential combinations of four feature classes: linear "l", quadratic "q", product

"p", and hinge "h" ("l", "lq", "lp", "lqp", "h", "lh", "lqh", "lph", "lqph")".

After the selection of variables, we conducted a detailed model selection exercise,

exploring all combinations of [a] the 17 values of the regularization parameter, [b] the

nine sets of potential combinations of four feature classes, and [c] the six sets of

bioclimatic variables (see above). As such, a total of 918 models were calibrated, and

each model was evaluated for statistical significance (partial ROC tests), performance

(omission rate), and the Akaike Information Criterion corrected for small sample sizes

(AICc). Final models were selected under the three criteria (first, statistical significance,

then omission rate, followed by AICc) (Cobos *et al.*, 2019).

We used the best parameter settings chosen to create final models. For each final model,

we ran 30 bootstrap replicates, with no clamping or extrapolation, and retaining a

random partition of 25% of the points from each run. These final models, though only

calibrated across M, were transferred to the extent of South America, particularly to

southern South America, which is the focus of this contribution. We used the cloglog

output format to assess median values across replicates as an estimate of the spatial

distribution of suitable and unsuitable conditions for the *P. torquans* complex across

South America. The uncertainty in model predictions was estimated using the range in

the suitability values (i.e., maximum minus minimum) across the model replicates. To

provide an additional assessment on the reliability of our model transfers, we calculated

the mobility-oriented parity (MOP) metric (Owens *et al.*, 2013) to offer a view of the

novelty of environmental values and combinations of values across the transfer area,

with special attention paid to areas of strict extrapolation (i.e., transfer areas with values

outside the range of climates in the calibration area).

We assessed the importance of each variable to determine the environmental suitability

for the *P. torquans* complex with median estimates across replicates of percent

contribution, permutation importance and Jackknife analysis. In order to determine the

most limiting variables, we used the Maxent limiting factor mapping tool described in

Elith *et al*. (2010), and implemented via the package "rmaxent" (Baumgartner *et al.*,

2017). The limiting factors are an interesting insight into the drivers of predictions, and

provide a powerful basis for comparisons with physiological knowledge (Elith *et al.*,

2010).

Patterns of colours for figures were generated with the package "tanagR" (Cadena &

Zapata, 2019), inspired by the plumage of passerine birds in the tanager family

(Thraupidae). The package currently includes palettes based on 16 tanager species from

Central and South America.

Results

We assessed 918 models for the final model calibration. Seven hundred and twenty-

seven of them were statistically significant as compared with a null model of random

prediction. Of such significant models, 137 (~19%) met the omission criterion of 5%.

Of the significant, low-omission models, only three met the AICc criterion of ΔAICc <

2. These three models were calibrated with the set of variables selected based on current

knowledge (Set 1), and were variants of the regularization multiplier (Table 1). Models

based on current knowledge selection of variables outperformed the ones calibrated

with automated selected datasets, as the best of models calibrated with Set 1 presented a

ΔAICc = 21.8 when compared with the best model calibrated with other set of variables.

Table 1. Performance metrics for parameter settings regarding regularization multiplier

(RMs), feature classes (FCs), and sets of environmental predictors (Pred. sets), used for

creating final models for the *P. torquans* complex. FCs are as follows: linear = l,

quadratic = q, product = p, and hinge = h.

RMs	FCs	Pred. Sets	Partial ROC	Omission rate 5%	AICc	ΔAICc	Weight AICc	N° of parameters
0.8	lq	Set 1	0.00	0.00	362.95	0	0.21	4
0.9	lq	Set 1	0.00	0.00	363.18	0.22	0.19	4
1	lq	Set 1	0.00	0.00	363.41	0.46	0.17	4

The "minimum temperature of the coldest month" (Bio06) was the most influential

variable by far, as evidenced by the percent contribution, permutation importance and

Jackknife analysis (Table 2 and Fig. 2). Its response curve evidence a bell shape, with a

central range optimal for prediction, while in both extremes its influence appeared to be

detrimental (Fig. S1). The second most important variable was the "mean temperature

of warmest quarter" (Bio10), followed by the "climatic moisture index" (CMI). The

CMI response curve was similar to the one from Bio06, while the response curve from

Bio10 evinced a sigmoidal shape with positive relation (Fig. S1). The "precipitation of

driest month" (Bio14) and the "precipitation of the wettest quarter" (Bio16) had nearly

no influence on the distribution modelling of the *P. torquans* complex.

Table 2. Variable importance: percent contribution and permutation importance

(median and sd). References: Bio10 = mean temperature of warmest quarter; Bio06 =

minimum temperature of coldest month; Bio14 = precipitation of driest month; Bio16 =

precipitation of wettest quarter; CMI = climatic moisture index.

Variable	Percent Contribution	Permutation Importance
Bio06	54.75 (5.67)	70.59 (8.09)
Bio10	25.10 (5.77)	18.71 (5.93)
Bio14	1.38 (3.29)	0.64 (4.27)
Bio16	0.19 (1.42)	0.88 (4.43)
CMI	12.89 (9.31)	6.44 (7.07)

Figure 2. Results of the jackknife test of variable importance, when variable is used in isolation or is omitted from modelling. References: Bio10 = mean temperature of warmest quarter; Bio06 = minimum temperature of coldest month; Bio14 = precipitation of driest month; Bio16 = precipitation of wettest quarter; CMI = climatic moisture index. [Colours after the Green-Grass Tanager *Chlorornis riefferii*]

Bio06 was the predictor associated with the largest decrease in suitability in Brasil and northern South America (Fig. S2). In most of the territory of Argentina, Bio10 was the

most limiting factor, while Bio14 limited suitability mostly along the Andean range

(Fig. S2).

As more than one best model was selected, we used the median and range of all

replicates across parameters to consolidate results for the *P. torquans* complex. The

median of the selected models (Fig 2A) identified areas with different levels of

suitability for the *P. torquans* complex across South America. Highly suitable areas

were concentrated on the central-east of Argentina and Uruguay, comprising the

Pampean and Chaco (eastern Chaco district) biogeographical provinces (Fig. S3, see in

Morrone, 2017). Suitability declined towards western Argentina, southern Brazil and

Paraguay. A broad area in central Chile was identified as suitable, although it may not

be accessible to the *P. torquans* complex by natural dispersion. Disparate areas along

the Andean range appeared to be suitable as well. The model showed moderate

uncertainty areas mainly throughout central-western Argentina (the Monte

biogeographical province), central Chile and the Andean range (Fig 2B and Fig. S3).

The uncertainty observed was mainly due to the variability among replicates (~95%)

(Fig. S3). The MOP analysis revealed that disparate areas in southern Chile and

northern South America would present model extrapolation (Fig 2C).

Figure 3. Climatologically suitable regions for the *Philornis torquans* complex

distribution in South America. a) Median prediction. b) Uncertainty (range) associated

with the median prediction of suitable regions for the *Philornis torquans* complex

distribution in South America. c) Mobility-oriented parity analysis revealing areas in S.

America for which the model predictions are based strictly on extrapolation. [Colours

after Scarlet-Breasted Dacnis *Dacnis berlepschi* and Moss-backed Tanager *Bangsia*

edwardsi]

Discussion

In this study, besides providing a model map of environmental suitability of the *P.*

torquans complex in South America, we also contributed with data related to the

constraints of its abiotic niche. Ecological niche modelling is a powerful tool to identify

macro-environmental variables defining the environmental niche, and thus driving the

geographical distribution of species. This study took into consideration many of the

recent methodological recommendations to produce robust ecological niche models, as

outlined by Feng *et al.* (2019): filtering and thinning occurrences (Boria *et al.*, 2014;

Aiello-Lammens *et al.*, 2015); using an accessible area for model calibration (Barve *et*

al., 2011); testing a number of candidate sets of environmental variables and multiple

combinations of model parameters (features, regularization multipliers, and sets of

variables); using multiple statistical criteria (partial ROC, omission rates, and AICc) for

model selection (Cobos *et al.*, 2019); replicating the model selected a number of times

(e.g., 30) to determine that results are consistent (Morales *et al.*, 2017); and accounting

for model uncertainty in geographical predictions (Owens *et al.*, 2013). As in any

modelling approach, counting with a sufficient sample size is a central issue in

ecological niche modelling. As suggested (van Proosdij *et al.*, 2016), the final sample

size in our study after spatial thinning seems appropriate to produce useful results.

Moreover, selecting the best combination of parameters (features and regularization

multiplier) further improves our ability to select the most appropriate model when

dealing with a reduced number of occurrences (Morales *et al.*, 2017).

Among a number of causes driving uncertainty in niche modelling, the nature of the

used occurrence dataset might represent a concern to our modelling approach. As far as

we know, there are no published records of free-living adults of any *Philornis* species,

with the sole exception of the introduced *P. downsi* in the Galapagos Islands (Causton *et*

al., 2014). Therefore, we modelled the ecological niche of the *P. torquans* complex

from reported records of immature (parasitic) stages, as an undeniable proxy of the

presence of adults. To date, no *Philornis* species is known to migrate, thus we assume

that there are adults present year-round in the same area, as also observed in *P. downsi*

(Causton *et al.*, 2014). Despite we are indeed modelling the niche of the *P. torquans*

complex from its parasitic state, we have no reason to suspect that the true geographical

distribution might differ among states.

We hypothesised that low temperatures and reduced precipitation will restrain the

occurrence of the *Philornis torquans* complex, as these abiotic parameters play a

predominant role in establishing the distribution limits of organisms and have already

been related with the intensity of parasitism by *Philornis* larvae (Dudaniec *et al.*, 2007;

Antoniazzi *et al.*, 2011; Manzoli *et al.*, 2013). Since insects are poikilothermic entities,

their rate of development and survival can vary greatly in response to many biotic and

abiotic factors (Adams, 1979; Couret *et al.*, 2014). Winters are the most critical period

for insects at higher latitudes since overwintering conditions strongly affect their fitness

(Irwin & Lee, 2000). Indeed, our model identified the minimum temperature of coldest

month (Bio06) as the most relevant variable (Table 2 and Fig. 2), revealing a

detrimental effect in both sides of a suitable range of temperatures (Fig. S1).

Morphotypes suggested to belong to the *P. torquans* complex (Monje *et al.*, 2013) are

considered to be originated in southern South America (somewhere near the border

between Brazil and Uruguay - Löwenberg-Neto & De Carvalho, 2009). Representing

the southernmost species of the genus, and unlike the majority of *Philornis* species, the

P. torquans complex must have adapted to colder, longer and drier winters; typical of

temperate areas. The observed data across presence sites of the *P. torquans* complex

indicates that minimum temperatures of coldest month ranges between 3°C and 18°C.

Therefore, it is highly likely that the *P. torquans* complex evolved to overwinter as

adults or pupae, as known for other Diptera species (Protocalliphoridae, Culicidae,

Muscidae) (Bellamy & Reeves, 1963; Stiner, 1969; Skidmore, 1985; Sabrosky *et al.*,

1989; Nasci *et al.*, 2001). Despite being unknown, we infer that the *P. torquans*

complex may overwinter as adults, a well-known strategy for Protocalliphora blowflies

(Sabrosky *et al.*, 1989). The former is sustained in the fact that *Philornis* adults are

known to survive over three months in captivity (Lincango & Causton, 2008; Saravia-

Pietropaolo *et al.*, 2018) and had been trapped at the same territories all year round

(Causton *et al.*, 2014).

On the other hand, insects adapted to cold winters may be handicapped by mild

overwintering temperatures (Irwin *et al.*, 2001). Our model suggests that warm winters

(signalled by Bio06) may be limiting the northern distribution of the *P. torquans*

complex (Fig. S2). Such scenario seems plausible, as warmer winter temperatures may

prevent adults to enter dormancy, thus increasing energy consumption and depleting

vital reserves in a period of potential food shortage (Tauber *et al.*, 1986; Harrington *et*

al., 2001; Irwin & Lee Jr, 2003; Sinclair, 2015). Therefore, these individuals may not be

able to reach the next breeding season or may leave them without enough energy

reserves for early reproduction (Harrington *et al.*, 2001).

As aforementioned, the developmental rate of insects is related to environmental

conditions, particularly as temperature impacts critical rate processes throughout

development and adulthood (Kingsolver & Huey, 2008). The temperature dependence

of poikilotherms' developmental processes is commonly represented as the cumulative

sum of degree-time products required for the development of key life stages, such as

egg maturation or emergence from pupae (e.g., Kasap & Alten, 2005; Nieto *et al.*, 2006;

Gutierrez & Ponti, 2014; Harnden & Tomberlin, 2016; Yanmanee *et al.*, 2016). In these

regards, the mean temperature of warmest quarter (Bio10) proved to be the second most

important variable (Table 2 and Fig. 2), showing a positive relation with sigmoidal

shape between predicted probability of presence and Bio10 (Fig. S1). This sigmoidal

shape implies a threshold temperature (apparently around 10°C), above which the

predicted probability of presence increases exponentially. The former makes sense

when considering that the increase of Bio10 might favour the development of

temperature-dependent non-parasitic stages, such as pupae maturation, which lately

relates with the abundance of adults. Otherwise, temperatures below the apparent

threshold seems to limit the distribution of the *P. torquans* complex, as evinced along

the Andean range (Fig. S2).

The third most relevant variable, the climatic moisture index (CMI), resulted to be the

most influential factor in most of Argentina, northern Chile and coastal Peru (Fig. S2).

The CMI is an aggregate measure of potential water availability imposed solely by

climate, ranging from –1 to +1, with wet climates showing positive values, dry climates

negative (Vörösmarty *et al.*, 2005). The CMI values observed through the actual

distribution of the *P. torquans* complex (mean -0.25, sd 0.19) indicate the species must

endure mostly a dry environment when compared with most of South America. As

reported for species laying eggs into a relatively dry environment (Grzywacz *et al.*,

2012), the eggs of the *P. torquans* morphotype present a smaller portion of the egg

surface that functions as a plastron (Patitucci *et al.*, 2017), a respiratory structure that

facilitates oxygen income minimizing water loss (Hinton, 1960). This reduced size of

the plastron of the *P. torquans* morphotype may have evolved as an adaptation allowing

the radiation to drier conditions typical of central Argentina from its centre of origin

(see Löwenberg-Neto & De Carvalho, 2009).

Opposed to our prediction, the ecological niche model suggests that precipitation has no

direct influence on limiting the environmental suitability of the *P. torquans* complex.

The "precipitation of driest month" (Bio14) and the "precipitation of the wettest

quarter" (Bio16) had a poor contribution to the model performance (Table 2), and

evinced a weak explanatory power in the jackknife analysis (Fig. 2). Despite it is clear

that humidity and/or moisture are relevant environmental factors determining the *P.*

torquans complex distribution, as reflected by CMI and supported by this model, the

proximal influence of precipitation seems to be mainly related with the intensity of

parasitism by *Philornis* species, as previously reported (Arendt, 1985, 2000; Dudaniec

et al., 2007; Antoniazzi *et al.*, 2011; Manzoli *et al.*, 2013).

The outcome of the geographical projection of the environmental niche model was a

map of the potential distribution of the *P. torquans* complex in South America.

According to our results, the *P. torquans* complex is potentially distributed across

central-eastern Argentina, mostly restricted to the Pampean and Chaco (eastern Chaco

district) biogeographical provinces, and declining with some uncertainty in the

surrounding Monte biogeographical province (Fig. 3 and Fig. S3). The later makes

sense due to the historical relationship among these three biogeographical provinces of

the Chacoan dominion (Morrone, 2017). Cabrera (1976) hypothesized a sequence of

biotic impoverishment from the Chaco province to the Pampean province, with the

Monte province being intermediate between both. Furthermore, the Chacoan and the

Monte provinces are closely related, as evidenced by the Chacoan affinities of the

entomofauna of the Monte and a cladistic biogeographic analysis based on beetle and

plant taxa (revised in Morrone, 2017). Regarding the broad area predicted as suitable in

central Chile, and despite its relative high uncertainty, the Andean range might represent

a pronounced topographic barrier impeding the natural dispersion of the *P. torquans*

complex to the area.

In conclusion, our work is the first one to identify macro-environmental variables

driving the geographical distribution of a *Philornis* species by evaluating their

contribution to its' environmental niche. Besides the relevance of our model as a tool

for bird's conservation, our model integrates the most up-to-date methodological

recommendations to produce a robust ecological niche; thus representing a

methodological reference for future work on the distribution of this group of parasites.

Acknowledgements

We greatly acknowledge the contribution made by Salvador S.A., who kindly provided

the geographic coordinates from the sites mentioned in Salvador & Bodrati (2013).

This work was funded by Consejo Nacional de Investigaciones Científicas y Técnicas

[PIP11220100100261; PIP 11220130100561CO]; and Universidad Autónoma de Entre

Ríos [PIDAC S013805/15]. The funders had no role in study design, data collection and

analysis, decision to publish, or preparation of the manuscript.

Contribution of authors

PFC and MAQ participated in the design and coordination of the study, and drafted the

manuscript; PFC performed the data analysis and modelling; AP carried out field

surveys and revised the existing literature; LM and PMB provided financial support and

critically reviewed the manuscript. All authors revised the manuscript and gave final

approval for publication.

Competing interests

We have no competing interests.

References

Adams, T.S. (1979) The reproductive physiology of the Screwworm, *Cochliomyia*

hominivorax (Diptera: Calliphoridae) II. Effect of constant temperatures on oogenesis.

Journal of Medical Entomology, **15**, 484–487.

Aiello-Lammens, M.E., Boria, R.A., Radosavljevic, A., Vilela, B. & Anderson, R.P.

(2015) spThin: An R package for spatial thinning of species occurrence records for use

in ecological niche models. *Ecography*, **38**, 541–545.

Antoniazzi, L.R., Manzoli, D.E., Rohrmann, D., Saravia, M.J., Silvestri, L. &

Beldomenico, P.M. (2011) Climate variability affects the impact of parasitic flies on

Argentinean forest birds. *Journal of Zoology*, **283**, 126–134.

Arendt, W.J. (1985) Philornis Ectoparasitism of Pearly-eyed Thrashers. I. Impact on

Growth and Development of Nestlings. *The Auk*, **102**, 270–280.

Arendt, W.J. (2000) Impact of nest predators, competitors, and ectoparasites on Pearly-

eyed Thrashers, with comments on the potential implications for Puerto Rican Parrot

recovery. *Ornitologia Neotropical*, 13–63.

Barve, N., Barve, V., Jiménez-Valverde, A., Lira-Noriega, A., Maher, S.P., Peterson,

A.T., *et al.* (2011) The crucial role of the accessible area in ecological niche modeling

and species distribution modeling. *Ecological Modelling*, **222**, 1810–1819.

Baumgartner, J., Wilson, P. & Esperón-Rodríguez, M. (2017) rmaxent: Tools for

working with Maxent in R. R package version 0.8.3.9000.

Beier, C. & Fontana, C. (2019) Breeding biology of the endangered Yellow Cardinal

Gubernatrix cristata in Brazil. *Revista Brasileira de Ornitologia - Brazilian Journal of*

Ornithology, **27**.

Bellamy, R.E. & Reeves, W.C. (1963) The winter biology of *Culex tarsalis* (Diptera:

Culicidae) in Kern County, California. *Annals of the Entomological Society of America*,

56, 314–323.

BirdLife International. (2018) *Gubernatrix cristata* [WWW Document]. *The IUCN Red*

List of Threatened Species. URL http://dx.doi.org/10.2305/IUCN.UK.2018-

2.RLTS.T22721578A131888081.en [accessed on 2018].

Boria, R.A., Olson, L.E., Goodman, S.M. & Anderson, R.P. (2014) Spatial filtering to

reduce sampling bias can improve the performance of ecological niche models.

Ecological Modelling, **275**, 73–77.

Bronstein, J.L. (1994) Conditional outcomes in mutualistic interactions. *Trends in*

Ecology and Evolution, **9**, 214–217.

Bulgarella, M., Quiroga, M.A. & Heimpel, G.E. (2018) Additive negative effects of

Philornis nest parasitism on small and declining Neotropical bird populations. *Bird*

Conservation International.

Cabrera, A.L. (1976) Regiones fitogeográficas argentinas. In *Enciclopedia Argentina de*

Agricultura y Jardinería (ed. by Kugler, W.F.). ACME, Buenos Aires, pp. 1–85.

Cadena, D. & Zapata, F. (2019) tanagR: Tanager-Inspired Color Palettes. R package

version 0.1.0.

Causton, C., Lincango, P., Lahuatte, P., Cedeño, D. & Castenada, J. (2014) *Philornis*

research update. Puerto Ayora, Ecuador.

Cobos, M.E., Peterson, A.T., Barve, N. & Osorio-Olvera, L. (2019) kuenm: an R

package for detailed development of ecological niche models using Maxent. *PeerJ*, **7**,

e6281.

Couret, J., Dotson, E. & Benedict, M.Q. (2014) Temperature, larval diet, and density

effects on development rate and survival of *Aedes aegypti* (Diptera: Culicidae). *PLOS*

ONE, **9**, e87468.

Couri, M. (1999) Myiasis caused by obligatory parasites. Ia. *Philornis* Meinert

(Muscidae). In *Myiasis in man and animals in the neotropical region* (ed. by

Guimarães, J.H. & Papavero, N.). Editora Plêiade, pp. 51–70.

Couri, M.S., Antoniazzi, L.R., Beldomenico, P. & Quiroga, M. (2009) Argentine

Philornis Meinert species (Diptera: Muscidae) with synonymic notes. *Zootaxa*, **62**, 52–

62.

Domínguez, M., Reboreda, J.C. & Mahler, B. (2015) Impact of Shiny Cowbird and

botfly parasitism on the reproductive success of the globally endangered Yellow

Cardinal *Gubernatrix cristata*. *Bird Conservation International*, **25**, 294–305.

Dudaniec, R.Y., Fessl, B. & Kleindorfer, S. (2007) Interannual and interspecific

variation in intensity of the parasitic fly, *Philornis downsi*, in Darwin's finches.

Biological Conservation, **139**, 325–332.

Dudaniec, R.Y. & Kleindorfer, S. (2006) Effects of the parasitic flies of the genus

Philornis (Diptera: Muscidae) on birds. *Emu*, **106**, 13–20.

Dudaniec, R.Y., Kleindorfer, S. & Fessl, B. (2006) Effects of the introduced

ectoparasite *Philornis downsi* on haemoglobin level and nestling survival in Darwin's

Small Ground Finch (*Geospiza fuliginosa*). *Austral Ecology*, **31**, 88–94.

Elith, J., Graham, C.H., Anderson, R.P., Dudík, M., Ferrier, S., Guisan, A., *et al.* (2006)

Novel methods improve prediction of species' distributions from occurrence data.

Ecography, **29**, 129–151.

Elith, J., Kearney, M. & Phillips, S. (2010) The art of modelling range-shifting species.

Methods in Ecology and Evolution, **1**, 330–342.

Escobar, L.E., Lira-Noriega, A., Medina-Vogel, G. & Townsend Peterson, A. (2014)

Potential for spread of the white-nose fungus (*Pseudogymnoascus destructans*) in the

Americas: Use of Maxent and NicheA to assure strict model transference. *Geospatial*

Health, **9**, 221–229.

Feng, X., Park, D.S., Walker, C., Peterson, A.T., Merow, C. & Papeş, M. (2019) A

checklist for maximizing reproducibility of ecological niche models. *Nature Ecology &*

Evolution, **3**, 1382–1395.

Fessl, B., Sinclair, B.J. & Kleindorfer, S. (2006) The life-cycle of *Philornis downsi*

(Diptera: Muscidae) parasitizing Darwin's finches and its impacts on nestling survival.

Parasitology, **133**, 739–747.

Grzywacz, A., Szpila, K. & Pape, T. (2012) Egg morphology of nine species of Pollenia

Robineau-Desvoidy, 1830 (Diptera: Calliphoridae). *Microscopy Research and*

Technique, **75**, 955–967.

Gutierrez, A.P. & Ponti, L. (2014) The new world screwworm: Prospective distribution

and role of weather in eradication. *Agricultural and Forest Entomology*, **16**, 158–173.

Harnden, L.M. & Tomberlin, J.K. (2016) Effects of temperature and diet on black

soldier fly, *Hermetia illucens* (L.) (Diptera: Stratiomyidae), development. *Forensic*

Science International, **266**, 109–116.

Harrington, R., Fleming, R.A. & Woiwod, I.P. (2001) Climate change impacts on insect

management and conservation in temperate regions: can they be predicted? *Agricultural*

and Forest Entomology, **3**, 233–240.

Hayes, C.D., Hayes, T.I., McClure, C.J.W., Quiroga, M., Thorstrom, R.K., Anderson,

D.L., *et al.* (2018) Native parasitic nest fly impacts reproductive success of an island-

endemic host. *Animal Conservation*, **22**, 157–164.

Hernandez, P.A., Graham, C.H., Master, L.L., Albert, D.L., Deborah, L. & Albert, D.L.

(2006) The effect of sample size and species characteristics on performance of different

species distribution modeling methods. *Ecography*, **29**, 773–785.

Hinton, H.E. (1960) The chorionic plastron and its role in the eggs of the Muscinae

(Diptera). *Journal of Cell Science*, **s3-101**, 313–332.

Irwin, J.T., Bennett, V.A. & Lee Jr, R.E. (2001) Diapause development in frozen larvae of the Goldenrod Gall fly, *Eurosta solidaginis* fitch (Diptera: Tephritidae). *Journal of Comparative Physiology B*, **171**, 181–188.

Irwin, J.T. & Lee Jr, R.E. (2003) Cold winter microenvironments conserve energy and improve overwintering survival and potential fecundity of the Goldenrod Gall fly, *Eurosta solidaginis*. *Oikos*, **100**, 71–78.

Irwin, J.T. & Lee, R.E. (2000) Mild winter temperatures reduce survival and potential fecundity of the goldenrod gall fly, *Eurosta solidaginis* (Diptera: Tephritidae). *Journal of Insect Physiology*, **46**, 655–661.

Karger, D.N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R.W., *et al.* (2017) Climatologies at high resolution for the earth's land surface areas. *Scientific Data*, **4**, 170122.

Kasap, O.E. & Alten, B. (2005) Laboratory estimation of degree-day developmental requirements of *Phlebotomus papatasi* (Diptera: Psychodidae). *Journal of Vector Ecology*, **30**, 328–333.

Kingsolver, J.G. & Huey, R.B. (2008) Size, temperature, and fitness: three rules. *Evolutionary Ecology Research*, **10**, 251–268.

Kinsella, J. & Winegarner, C. (1974) Notes on the life history of *Neomusca porteri* (Dodge), parasitic on nestlings of the great crested flycatcher in Florida. *Journal of Medical Entomology*, **11**, 633.

Kleindorfer, S. & Dudaniec, R.Y. (2016) Host-parasite ecology, behavior and genetics: A review of the introduced fly parasite *Philornis downsi* and its Darwin's finch hosts. *BMC Zoology*.

Koop, J.A.H., Huber, S.K., Laverty, S.M. & Clayton, D.H. (2011) Experimental demonstration of the fitness consequences of an introduced parasite of Darwin's

finches. *PLoS ONE*, **6**, e19706.

Kramer-Schadt, S., Niedballa, J., Pilgrim, J.D., Schröder, B., Lindenborn, J., Reinfelder, V., *et al.* (2013) The importance of correcting for sampling bias in MaxEnt species distribution models. *Diversity and Distributions*, **19**, 1366–1379.

Labud, V.A., Semenas, L.G. & Laos, F. (2003) Diptera of sanitary importance associated with composting of biosolids in Argentina. *Revista de Saude Publica*, **37**, 722–728.

Lincango, P. & Causton, C. (2008) *Crianza en cautiverio de* Philornis downsi*, en las Islas Galápagos*.

Little, S.E. (2008) Myasis in wild birds. In *Parasitic diseases of wild birds* (ed. by Atkinson, C.T., Thomas, N.J. & Hunter, D.B.). Wiley Blackwell, pp. 546–556.

Löwenberg-Neto, P. (2008) The structure of the parasite-host interactions between *Philornis* (Diptera: Muscidae) and neotropical birds. *Journal of Tropical Ecology*, **24**, 575–580.

Löwenberg-Neto, P. & Carvalho, C.J.B. De. (2009) Areas of endemism and spatial diversification of the Muscidae (Insecta: Diptera) in the Andean and Neotropical regions. *Journal of Biogeography*, **36**, 1750–1759.

Manzoli, D.E., Antoniazzi, L.R., Saravia, M.J., Silvestri, L., Rorhmann, D. & Beldomenico, P.M. (2013) Multi-level determinants of parasitic fly infection in forest passerines. *PLoS ONE*, **8**, e67104.

Manzoli, D.E., Saravia-Pietropaolo, M.J., Antoniazzi, L.R., Barengo, E., Arce, S.I., Quiroga, M.A., *et al.* (2018) Contrasting consequences of different defence strategies in a natural multihost- parasite system. *International Journal of Parasitology*.

Merow, C., Smith, M.J. & Silander, J.A. (2013) A practical guide to MaxEnt for modeling species' distributions: What it does, and why inputs and settings matter.

Ecography, **36**, 1058–1069.

Monje, L.D., Quiroga, M., Manzoli, D.E., Couri, M.S., Silvestri, L., Venzal, J.M., *et al.*
(2013) Sequence analysis of the internal transcribed spacer 2 (ITS2) from *Philornis*
seguyi (García, 1952) and *Philornis torquans* (Nielsen, 1913) (Diptera: Muscidae).
Systematic Parasitology, **86**, 43–51.

Morales, N.S., Fernández, I.C. & Baca-González, V. (2017) MaxEnt's parameter
configuration and small samples: are we paying attention to recommendations? A
systematic review. *PeerJ*, **5**, e3093.

Morrone, J.J. (2017) *Neotropical biogeography. Regionalization and evolution*. CRC
Press, Inc.

Nasci, R.S., Savage, H.M., White, D.J., Miller, J.R., Cropp, B.C., Godsey, M.S., *et al.*
(2001) West Nile virus in overwintering Culex mosquitoes, New York City, 2000.
Emerging Infectious Diseases, **7**, 742–744.

Nieto, P., Malone, J.B. & Bavia, M.E. (2006) Ecological niche modeling for visceral
leishmaniasis in the state of Bahia, Brazil, using genetic algorithm for rule-set
prediction and growing degree day-water budget analysis. *Geospatial health*, **1**, 115–
126.

Oniki, Y. (1983) Notes on fly (Muscidae) parasitism of nestlings of South American
birds. *Le Gerfaut*, **73**, 281–286.

Owens, H.L., Campbell, L.P., Dornak, L.L., Saupe, E.E., Barve, N., Soberón, J., *et al.*
(2013) Constraints on interpretation of ecological niche models by limited
environmental ranges on calibration areas. *Ecological Modelling*, **263**, 10–18.

Patitucci, L.D., Quiroga, M., Couri, M.S. & Saravia-Pietropaolo, M.J. (2017)
Oviposition in the bird parasitic fly *Philornis torquans* (Nielsen, 1913) (Diptera:
Muscidae) and eggs' adaptations to dry environments. *Zoologischer Anzeiger*, **267**, 15–

20.

Phillips, S.J., Anderson, R.P. & Schapire, R.E. (2006) Maximum entropy modeling of

species geographic distributions. *Ecological Modelling*, **190**, 231–259.

Proosdij, A.S.J. van, Sosef, M.S.M., Wieringa, J.J. & Raes, N. (2016) Minimum

required number of specimen records to develop accurate species distribution models.

Ecography, **39**, 542–552.

Quiroga, M.A., Monje, L.D., Arrabal, J.P. & Beldomenico, P.M. (2016) Datos

moleculares nuevos sobre *Philornis* (Diptera: Muscidae) subcutáneas del sur de

Sudamérica sugieren la existencia de un complejo de especies. *Revista Mexicana de*

Biodiversidad, **87**, 1383–1386.

Quiroga, M.A. & Reboreda, J.C. (2012) Lethal and sublethal effects of Botfly

(*Philornis seguyi*) parasitism on House Wren nestlings. *The Condor*, **114**, 197–202.

R Core Team. (2018) R: A language and environment for statistical computing.

Raghavan, R.K., Peterson, A.T., Cobos, M.E., Ganta, R. & Foley, D. (2019) Current

and future distribution of the Lone Star Tick, *Amblyomma americanum* (L .) (Acari:

Ixodidae) in North America. *PLoS ONE*, **14**, e0209082.

Sabrosky, C.W., Bennett, G.F., Whitworth, T.L. & others. (1989) *Bird blow flies*

(Protocalliphora) in North America (Diptera: Calliphoridae) with notes on Palearctic

species. Smithsonian Institution Press, Washington, D.C.

Saravia-Pietropaolo, M.J., Arce, S.I., Manzoli, D.E., Quiroga, M. & Beldomenico, P.M.

(2018) Aspects of the life cycle of the avian parasite *Philornis torquans* (Diptera:

Muscidae) under laboratory rearing conditions. *The Canadian Entomologist*, **150**, 317–

325.

Showler, A.T. & Osbrink, W.L.A. (2015) Stable Fly, *Stomoxys calcitrans* (L.), dispersal

and governing factors. *International Journal of Insect Science*, **7**, IJIS.S21647.

Silvestri, L., Antoniazzi, L.R., Couri, M.S., Monje, L.D. & Beldomenico, P.M. (2011) First record of the avian ectoparasite *Philornis downsi* Dodge & Aitken, 1968 (Diptera: Muscidae) in Argentina. *Systematic Parasitology*, **80**, 137–140.

Sinclair, B.J. (2015) Linking energetics and overwintering in temperate insects. *Journal of Thermal Biology*, **54**, 5–11.

Skidmore, P. (1985) *The biology of the Muscidae of the World*. Dr W. Junk Publishers, Dordrecht, The Netherlands.

Stiner, F. (1969) Overwintering by *Protocalliphora metallica* (Diptera: Calliphoridae). *Annals of the Entomological Society of America*, **62**, 1205–1206.

Tauber, M.J., Tauber, C.A. & Masaki, S. (1986) *Seasonal adaptations of insects*. Oxford University Press.

Title, P.O. & Bemmels, J.B. (2018) ENVIREM: an expanded set of bioclimatic and topographic variables increases flexibility and improves performance of ecological niche modeling. *Ecography*, **41**, 291–307.

Vignali, S., Barras, A. & Braunisch, V. (2019) SDMtune: Species Distribution Model Selection.

Vörösmarty, C.J., Douglas, E.M., Green, P.A. & Revenga, C. (2005) Geospatial indicators of emerging water stress: an application to Africa. *AMBIO: A Journal of the Human Environment*, **34**, 230–236.

Warren, D.L., Seifert, S.N., Warren, D.L. & Seifert, S.N. (2016) Ecological niche modeling in Maxent: the importance of model complexity and the performance of model selection criteria. *Ecological Applications*, **21**, 335–342.

Yanmanee, S., Husemann, M., Benbow, M.E. & Suwannapong, G. (2016) Larval development rates of *Chrysomya rufifacies* Macquart, 1842 (Diptera: Calliphoridae) within its native range in South-East Asia. *Forensic Science International*, **266**, 63–67.

695   Young, B.E. (1993) Effects of the parasitic botfly Philornis carinatus on nestling house

696   wrens, *Troglodytes aedon*, in Costa Rica. *Oecologia*, **93**, 256–262.

697   Zumpt, F. (1965) *Myiasis in man and animals in the Old World. A textbook for*

698   *physicians, veterinarians and zoologists. London, Butterworth.* Butterworth & Co Ltd.,

699   London, United Kingdom.