Phylogenetic and metabolic diversity have contrasting effects on the ecological functioning of bacterial communities

Constantinos Xenophontos[1,2], Martin Taubert[1], W. Stanley Harpole[2,3,4], Kirsten Küsel[1,2]

1. Institute of Biodiversity, Friedrich Schiller University Jena, Jena 07743, Germany
2. German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, 04103 Leipzig, Germany
3. Department of Physiological Diversity, Helmholtz-Center for Environmental Research (UFZ), 04318 Leipzig, Germany
4. Institute of Biology, Martin Luther University of Halle-Wittenberg, 06120 Halle, Germany

Correspondence:

Kirsten Küsel kirsten.kuesel@uni-jena.de

Abstract

Quantifying the relative contributions of microbial species to ecosystem functioning is

challenging, because of the distinct mechanisms associated with microbial phylogenetic and

metabolic diversity. We constructed bacterial communities with different diversity traits and

employed exoenzyme activities (EEAs) and total available carbon (TAC) from substrates as

proxies of bacterial functioning to test the independent effects of these two aspects of

biodiversity. We expected that metabolic diversity, but not phylogenetic diversity would be

associated with greater ecological function. Phylogenetically relatedness should intensify

species interactions and coexistence, therefore amplifying the influence of metabolic

diversity. We examined the effects of each diversity treatment using linear models, while

controlling for the other, and found that phylogenetic diversity strongly influenced

community functioning, positively and negatively. Metabolic diversity, however, exhibited

negative or non-significant relationships with community functioning. When controlling for

different substrates, EEAs increased along with phylogenetic diversity but decreased with

metabolic diversity. The strength of diversity effects was related to substrate chemistry and

the molecular mechanisms associated with each substrate's degradation. EEAs of

phylogenetically similar groups were strongly affected by within-genus interactions. These

results highlight the unique flexibility of microbial metabolic functions that must be

considered in further ecological theory development.

40 Keywords

41 ecosystem functions; phylogenetic diversity; metabolic diversity; biodiversity; bacteria;

42 synthetic communities; exoenzymes, community interactions

Introduction

Bacterial activities are driving ecosystem functioning (Finlay, Maberly and Cooper

1997) and recent research has highlighted the contribution of microbial diversity to the

biological processes occurring in ecosystems (Petchey and Gaston 2006; Cardinale *et al.*

2012; Galand *et al.* 2018; Le Bagousse-Pinguet *et al.* 2019; Johnson and Pomati 2020). As

such, a direct assessment of metabolic diversity is a great predictor of community ecosystem

functions (Mokany, Ash and Roxburgh 2008; Salles *et al.* 2009; Krause *et al.* 2014).

Microbial metabolic traits can be determined based on, e.g., functional gene-targeting

microarrays and metabolic activity assays (Litchman and Klausmeier 2008; Naeem *et al.*

2009; Paula *et al.* 2014; Wang *et al.* 2017). Such assays have shown that metabolic diversity

is positively correlated with ecologically important community functions, such as organic

matter decomposition and CO_2 production (Rivett *et al.* 2016; Evans *et al.* 2017). Microbial

functions like organic matter degradation are often conveyed by exoenzymes. The production

and release of exoenzymes is an important mechanism that bacteria use to access exogenous,

complex carbon and energy sources. Inevitably, exoenzymes are shared among different

species (Drescher *et al.* 2014; Smith and Schuster 2019), and thus their production is expected

to be affected by community species interactions (Folse and Allison 2012). Exoenzyme

activities (EEAs) can also link multiple bacterial metabolic processes to ecosystem functions

like carbon cycling (Malik *et al.* 2019). Moreover, metabolic traits affect fitness, as they relate

to energy acquisition and expenditure, placing them under the influence of strong

evolutionary selection processes. If we aim to improve our understanding of how metabolic

diversity influences microbial community ecosystem functioning, the evolutionary history and

relationship of microbes must be considered.

Phylogenetic relationships are usually assessed using neutral genetic markers like the

bacterial 16S rRNA gene. But as a neutral marker it is not subject to selective pressures. So

ecologists have been debating the usefulness of phylogenetic diversity in the study of

ecosystem functioning, as studies have yielded mixed results (Cadotte *et al.* 2009; Flynn *et al.*

2011; Alexandrou *et al.* 2015; Gerhold *et al.* 2015; Venail *et al.* 2015). Nevertheless, closely

related bacterial species are often more likely to have similar metabolic traits (Martiny,

Treseder and Pusch 2013; Goberna and Verdú 2016). Phylogenetic diversity is a good

predictor for metabolic functions with a strong phylogenetic clustering: For example,

ammonia oxidation is restricted to a group of chemolithoautotrophic bacteria (Kowalchuk and

Stephen 2001), traced in deep phylogenetic branches. Yet, especially for broad, simple

functions such as carbon assimilation from amino acids and simple sugars (Martiny, Treseder

and Pusch 2013), it is a poor predictor of ecological functioning (Srivastava *et al.* 2012;

Martiny, Treseder and Pusch 2013; Goberna and Verdú 2016). Nevertheless, under certain

circumstances, phylogenetic relationships can predict growth rates (Morrissey *et al.* 2016),

species interactions via ecological networks (Srivastava *et al.* 2012) and species coexistence

(Germain, Weir and Gilbert 2016). Hence, we can link phylogenetic relatedness to properties

that are under natural selection and therefore contribute to ecological functioning (Morrissey

et al. 2016).

The precise connection behind the phylogenetic and metabolic diversities is still

unclear, because the underlying mechanisms are either unknown or very complex (Losos

2008). Since phylogenetic diversity is connected with species coexistence and interactions,

which should interact with metabolic diversity (Pascual-García, Bonhoeffer and Bell 2020)

and possibly modify or intensify the effects metabolic diversity has on community

functioning (Nealson and Popa 2005). For the further development of micro-ecological theory

(Krause *et al.* 2014; Shade *et al.* 2018) and biodiversity research (Flynn *et al.* 2011; Roger *et*

al. 2016), potential interactions between phylogenetic and metabolic diversity have to be

identified to understand their influence on community ecosystem functions (Thompson,

Davies and Gonzalez 2015).

Here, we aim to disentangle this interaction by using synthetic microbial communities

in a multifactorial experimental design, where we control for each diversity aspect while

studying the other's effect on community ecological functions. We used groundwater

bacterial isolates to form simplified four-member communities with different phylogenetic

and metabolic diversity treatments, allowing us to test how these two aspects of biodiversity

interact as they shape community ecological functions. EEA assays were used for the

metabolic classification of bacterial isolates and to assess the functioning of synthetic

communities. We hypothesised that 1) when controlling for phylogenetic diversity, metabolic

diversity has a positive effect on community ecological functions. In contrast, 2) when

controlling for metabolic diversity, phylogenetic diversity, based on a neutral marker, should

have no effect on community ecological functions. Finally, 3) the effect of metabolic diversity

is proportionally stronger in communities that are phylogenetically similar, as

phylogenetically dissimilar communities hold species with less niche overlap and therefore

negative interactions like competition are weaker.

Material and Methods

Selection of bacterial isolates and culture setups

The twenty-seven bacterial strains used in this study (Appendix S1: Table S1) were

isolated from pristine groundwater (Geesink *et al.* 2018), obtained from the Hainich Critical

Zone Exploratory located in Thuringia, Germany (Küsel *et al.* 2016). The isolates were

selected from a pool of more than 340 groundwater isolates as representatives of the most

abundant taxonomic families within the total groundwater bacterial population, based on 16S

rRNA gene MiSeq amplicon sequencing data (Kumar *et al.* 2017; Schwab *et al.* 2017),

throughout the taxonomic classes of *Actinobacteria*, *Bacilli*, *Flavobacteriia*, *Sphingobacteria*,

Alpha- and *Gammaproteobacteria*.

The isolates were maintained on S2P medium agar (Geesink *et al.* 2018). Before each

of the experimental assays, isolates were cultured aerobically in 20 ml of liquid S2P medium,

in cap-perforated 50 ml Falcon tubes, horizontally shaken (125 rpm) at room temperature, for

48 h to early stationary phase. Growth in the liquid cultures was being determined via optical

density measurements at a wavelength of 600 nm (Synergy™ H4 Hybrid Multi-Mode

Microplate Reader, BioTek, Germany).

Molecular analysis and taxonomic classification of bacterial isolates

DNA from bacterial isolates was extracted with GenElute™ Bacterial Genomic DNA

Kit (NA2110-1KT, Sigma-Aldrich, Germany). Sterile inoculation loops were used to gather 1

µl-worth of biomass from the agar medium cultures, for DNA extraction. For the 16S rRNA

gene amplification, the primers 8F (5′-AGA GTT TGA TCC TGG CTC AG-3′) (Turner *et al.*

1999) and 1492R (5'-GGT TAC CTT GTT ACG ACT T-3') (Turner *et al.* 1999) were used.

PCR was performed with a PEQSTAR thermocycler (PEQLAB, Germany) using the

following cycling conditions: initial denaturation at 95 °C for 10 min, 30 cycles of 95 °C

(denaturation), 55 °C (primer annealing) and 72 °C (elongation) for 30 sec, 45 sec and 60 sec,

respectively, followed by 10 min at 72 °C (final elongation). DNA purification and

sequencing of the 1484bp-length PCR products was carried out by Macrogen (Amsterdam,

Netherlands) with the 27F (5'-AGA GTT TGA TCM TGG CTC AG-3') primer.

The sequences were assigned to their closest reference relative using BLASTn (Zhang

et al. 2000) (v2.8.0+) and cross-referenced with SILVA Incremental Aligner (SINA)

(Pruesse, Peplies and Glöckner 2012) (v1.2.11). A rooted maximum-likelihood phylogenetic

tree for the 16S rRNA gene taxonomic classification of the organisms was constructed with

MEGA X (Kumar *et al.* 2018) (v10.1.5).

Exoenzyme activity (EEA) assays

EEA assays for each bacterial isolate were performed using eleven substrates

(Appendix S1: Table S2) (Sigma-Aldrich, Germany), each carrying a fluorescent moiety

allowing spectrophotometric measurements of activity using a microplate reader (Synergy™

H4 Hybrid Multi-Mode Microplate Reader). The sensor was set to detect fluorescence at

wavelengths 360/9 nm for excitation and 450/9 nm for emission.

Prior to the EEA assays, the optical density of the bacterial cultures was determined

spectrophotometrically (600 nm). For the EEA assays, a modified version of the protocols

from Frossard *et al.* (Frossard *et al.* 2012) and Rivett *et al.* (Rivett *et al.* 2016) was used here,

as follows: a black 96-well plate setup, with 100 µl HEPES buffer at 100 mM stock

concentration in each well, was inoculated with 100 µl of bacterial culture and to each test

well 50 µl of substrate solution at 40 µM final concentration was added. Negative controls

consisted of wells containing 100 µl of sterile S2P liquid medium and wells with just 100 µl

sterile Milli-Q water (Merck Millipore, Germany). Fluorescence standards for the MUB

substrates were measured using 4-Methylumbelliferone with final concentrations of 10 to

0.3125 µM by halving dilutions. Fluorescence standards for the AMC substrates were

measured using 7-Amino-4-methylcoumarin with final concentrations of 2 to 0.0625 µM by

halving dilutions. The standards were used to transform relative fluorescence units (RFU) to

concentration of degraded substrate. Each assay plate setup held technical triplicates for the

cultures, controls and standards. Triplicates were averaged and their standard deviation

propagated for the data analyses. Background fluorescence, measured in negative controls

with either substrate solution or Milli-Q water, was subtracted from the respective assay

plates. Measurement of the assay plates begun immediately after the substrate addition and

continued for 90 min, at 2.5 min measurement intervals.

Determining phylogenetic and metabolic diversity of isolates

To examine effects of diversity on the functioning of synthetic bacterial communities,

the 16S rRNA gene taxonomy and EEAs of the twenty-seven bacterial isolates were used as

the basis of the community traits: phylogenetic and metabolic diversity. Using seventeen of

the tested isolates, we formed four trait-groups, each consisting of five isolates, being either

phylogenetically and metabolically similar (Psim-Msim), phylogenetically similar but

metabolically dissimilar (Psim-Mdis), phylogenetically dissimilar and metabolically similar

(Pdis-Msim) as well as phylogenetically and metabolically similar (Pdis-Mdis). In total, the

isolates covered ten taxonomic families of five classes (Appendix S1: Figure S1). In the Psim

groups, isolates were affiliated with the same genus; Psim-Msim group held *Flavobacteria*

isolates and Psim-Mdis held *Pseudomonas* isolates. Pdis groups were formed so that all

members of a group were related to different families. Subsequently, Psim and Pdis group

phylogenetic diversity quantitative measures were obtained using the "*Compare within group*

mean distance" function of MEGA X (Kumar *et al.* 2018) (v10.1.5) and verified to be

statistically insignificant for Psim communities and significantly different for Pdis

communities using paired t-tests (Appendix S1: Table S3). Hierarchical clustering based on

Euclidean distance was used to quantise functioning of the total EEAs for each isolate (Figure

1) and categorise them as metabolically similar or dissimilar. The clustering analysis was

performed in R (v. 3.6.1) (R Core Team 2019) using the core stats functions *dist(x,*

method = "euclidean", p = 2) to calculate the Euclidean dissimilarity distance matrix (Borg

and Groenen 2005), followed by *hclust(d, method = "ward.D")* to hierarchically rank and

cluster the EEA dissimilarities between the bacterial isolates using the Ward hierarchical

clustering method (Ward 1963; Murtagh and Legendre 2014). Msim groups were formed

from isolates of the same cluster, Mdis groups were formed so that each member of the group

belonged to a different metabolic cluster (Figure 1). Quantitative measures of principal

component analysis (PCA) based on the metabolic traits of the isolates confirmed the higher

metabolic diversity between isolates of Psim-Mdis and Pdis-Mdis compared to group Psim-

Msim and Pdis-Msim (Appendix S1: Figure S2).

Bacterial community setups and experimental design

For each of the four trait groups with five isolates (Psim-Msim, Psim-Mdis, Pdis-

Msim, Pdis-Mdis), we designed five distinct synthetic communities of bacterial species by

omitting one isolate each per community. This resulted in twenty communities, five for each

trait group, identified as C1 to C5. We confirmed the phylogenetic diversity between the

communities via the quantitative estimates of isolate phylogenetic distance and their

metabolic diversity based on the Euclidean distance matrix of isolate EEAs as previously

mentioned (Appendix S1: Table S3). The phylogenetic and metabolic diversity values for

each community showed a distinct differentiation of the communities from each of the four

groups, confirming that we successfully separated the bacterial isolates and communities into

the four intended groups (Figure 2).

We grew cultures of each isolate in liquid S2P medium as described above. Optical

density (600 nm) was used to calculate the required final volume for each isolate, so that all

members of each community were to be introduced into the communities at the same amount

of biomass. The communities were established in cultures by mixing a predetermined amount

of liquid isolate culture for a total volume of 2.5 ml. The community cultures were incubated

in deep 96-well plates (2.5 ml volume), at room temperature and sealed with rayon film seals

(391-1262, VWR, Germany) to allow gas exchange. The EEAs of the communities were

assessed after 0 h, 6 h and 24 h of incubation. Additionally, at the 0 h and 24 h marks, a

200 µl culture sample was pelleted by centrifugation ($17.000 \times g$, 5 min) and stored at -80 °C,

for subsequent molecular analysis.

Ecological functioning data analysis

As a measure of multifunctionality, we used the total available carbon (TAC) from

each substrate for a direct comparison of communities. This was done by multiplying the total

number of carbon atoms in each substrate (excluding the fluorescent moiety) with the molar

amount of degraded substrate. TAC was summed as a unified measure of ecological

functioning for each isolate and community. 4-MUB phosphate (PHOS) and 4-MUB sulfate

(SULF) were excluded because they contained no carbon atoms.

In addition, the effect of the diversity treatments on each individual EEA was

determined. The communities constructed in this study were composed out of different

isolates with widely differing metabolic activities. A comparison of EEAs between

communities alone would thus only reflect the differences between the constituent isolates.

Therefore, EEA differences of the communities were normalized to the respective isolates and

expressed as proportional changes of EEA. This was done with the following steps (visualised

in Appendix S1: Figure S3): The measured EEAs of the isolates within each community

group were used to calculate expected EEAs for each of the group's communities. The

expected EEAs were compared to the measured EEAs of the experimental communities and

the result was represented as % proportional changes of EEAs. This resulted in values from -

100% to 385%.

To evaluate the significance of phylogenetic diversity, metabolic diversity was

controlled by subsetting into metabolically similar and dissimilar communities. Likewise, to

examine the significance of metabolic diversity, phylogenetic diversity was controlled for in

the same way. These analyses were performed for the 0 h timepoint data, using linear

regression models in R. For the linear models, the TAC response variable data, were log2-

transformed. To also enable logarithmic scaling of the EEA % proportional change data,

101% was added to all datapoints, to avoid negative values and zeroes, since the smallest

values were -100%; As the TAC values were all > 0, this was not necessary. To avoid

pseudoreplication in the models, the 11 EEA % proportional change datapoints of each

community were averaged in the diversity treatments. For example, group Psim-Fsim

community C1 had eleven proportional change values, one for each substrate, that were

averaged into one. This was repeated for all communities in all group and resulted in a total of

20 datapoints. The group Pdis-Mdis showed no activity in the SULF assay, and so this assay

was excluded from the models. For the TAC data, 20 datapoints were also used, one

calculated value per community. Through subsetting the predictor variable data, to control for

each diversity effect, 10 datapoints were used for each model, for both EEA % proportional

change data as well as the TAC data. Furthermore, predictor variables, phylogenetic and

metabolic diversity metrics, were z-score normalised, for the linear regression models, to

make them comparable.

For the analyses for the influences of substrate biochemistry on the diversity treatment

effects, the EEA% proportional change data were not subset by diversity treatment but by

each substrate. Therefore, the EEA % proportional change response variable for each

substrate model was 20 datapoints or 15 for SULF. Following that, ANOVA of linear

regression models was performed as described above.

Effects of diversity in EEA changes over 24 h

Time is a crucial driver of bacterial interactions and therefore we evaluated how the

EEA proportional changes of communities developed over 24 h, by comparing the 0 h

timepoint to the 6 h and 24 h timepoint data. This was done for all 20 communities of all 4

diversity treatment groups, and for each substrate, the average of each group was determined.

The significance of temporal changes was tested using the Student's t-test (2-tailed) by

comparing the EEA proportional change of all communities in a treatment group at 0 h to all

EEA proportional changes of the same group for timepoints 6 h and 24 h, for each individual

substrate. False discovery rate was controlled with the Benjamini–Hochberg procedure

(McDonald 2014).

Substrate acclimation in EEA assays

To study influences of acclimation of isolates and communities to the added substrates

on the observed effects of diversity treatments on community ecological functions, we added

a pre-incubation period to the EEA assays. This was performed only with the Psim-Mdis

group. After mixing of the communities, substrates were added after 0 h and 6 h to isolates

and communities, and EEA assays were performed as described above (no-acclimation

treatments). In addition, isolates and communities with substrates added 0 h after mixing were

incubated for a total of 4.5 h, before another series of EEA assays were conducted

(acclimation treatments).

Assay duration was 90 min, as in previous assays. To achieve a direct comparability of

the activities measured with and without acclimation, the determined EEAs were normalized

to optical density of isolates and communities, which was measured before each assay.

Following that, the change in EEAs from 0 h to 4.5 h and 6 h was compared between isolates

and communities.

16S rRNA gene amplicon sequencing of synthetic communities

Community DNA was extracted as described above. DNA concentration was

measured spectrophotometrically (NanoDrop 1000, PEQLAB, Germany) and samples were

stored at -20 °C.

For Illumina MiSeq amplicon sequencing of 16S rRNA genes, PCR was performed

using the primer set Bakt_341F (5'-CCT ACG GGN GGC WGC AG-3') (Herlemann *et al.*

2011) and Bakt_805R (5'-GAC TAC HVG GGT ATC TAA TCC-3') (Herlemann *et al.* 2011)

targeting the V3-V4 hypervariable region of the rRNA gene. DNA amplification was carried

out with a PEQSTAR 2X thermocycler (PEQLAB, Germany) in the following conditions:

initial denaturation at 95 °C for 15 min, 30 cycles of 94 °C (denaturation), 55 °C (primer

annealing) and 72 °C (elongation) at 45 sec for each step, followed by 10 min at 72 °C (final

elongation). The PCR DNA products were purified using NucleoSpin Gel and PCR Clean-up

(Ref. 740609, Macherey-Nagel, Germany) following the manufacturer's protocol. The final

step, DNA recovery from the columns, was performed twice for a total volume of 60 µl

purified PCR product samples.

MiSeq-generated sequence data were analysed using mothur (Schloss *et al.* 2009)

(v.1.39.1; mothur.org) along with the mothur MiSeq SOP developed by Kozich *et al.* (Kozich

et al. 2013) (accessed November 19th 2018). Paired-end reads were combined and sequences

below 400 bp or above 429 bp as well as sequences with ambiguous bases or homopolymer

runs above 8 bp were removed. Sequences were aligned to the SILVA reference database

release SSU 132 (Quast *et al.* 2012) and clustered when differed in four or fewer bases.

Chimeric sequences were removed with uchime (Edgar 2016) using the GOLD database

(Mukherjee *et al.* 2019) as a reference. OTUs were binned from sequences with a 3% identity

cut-off and were classified using the same SILVA database. Raw data have been deposited in

the European Nucleotide Archive under the study with accession number PRJEB34832

(samples ERR3588683 - ERR3588722). The amplicon sequencing data were used to examine

shifts in the Pdis community species relative abundances that might explain changes in the

ecological functioning of communities that cannot be attributed to the diversity treatments.

Results

Effects of diversity on community ecological functions

We examined the absolute level of functioning of communities, using TAC turned

over by EEA in each community as a unified measure (Figure 3; Appendix S1: Table S4).

Linear regression analyses revealed that when controlling for metabolic diversity,

phylogenetic diversity had a significant negative effect on TAC in metabolically similar

communities (-0.305 ± 0.102, estimate $\pm$ std. error, Adj. $R^2 = 0.467$, p = 0.018) (Figure 3A).

This shows that communities with phylogenetically diverse isolates were able to acquire less

carbon than communities with phylogenetically similar isolates, when these isolates had a

high metabolic similarity. For metabolically dissimilar communities, phylogenetic diversity

had a positive effect (0.313 ± 0.105, estimate $\pm$ std. error, Adj. $R^2 = 0.467$, p = 0.018). While

controlling for phylogenetic diversity, metabolic diversity had a negative effect in

phylogenetically similar communities (-0.611 ± 0.151, estimate $\pm$ std. error, Adj. $R^2 = 0.632$,

p = 0.004) (Figure 3B), but no effect was observed for phylogenetically dissimilar

communities (0.002 ± 0.115, estimate $\pm$ std. error, Adj. $R^2 = -0.125$, p = 0.986).

In addition to TAC, similar trends were also observed when performing linear

regression analyses on community EEA changes (Appendix S1: Figure S4, Table S5). For

metabolically similar communities, phylogenetic diversity had a significant negative effect on

the EEAs (-0.279 ± 0.121, estimate $\pm$ std. error; Adj. $R^2 = 0.325$; p = 0.050) when controlling

for metabolic diversity (Appendix S1: Figure S4A). In contrast, for metabolically dissimilar

communities, a significant positive effect of phylogenetic diversity was observed

(1.094 ± 0.231, estimate $\pm$ std. error, Adj. $R^2 = 0.704$, p = 0.002). While controlling for

phylogenetic diversity phylogenetically similar communities exhibited a strong negative

relationship, (-1.535 ± 0.366, estimate $\pm$ std. error, Adj. $R^2 = 0.648$, p = 0.003), while

phylogenetically dissimilar communities' relationship was below the significance threshold

(-0.121 ± 0.113, estimate ± std. error, Adj. R^2 = 0.017, p = 0.314) (Appendix S1: Figure S4B).

Hence, biodiversity effects in the carbon acquisition potential of these communities were also

reflected on the community EEAs.

Interestingly, we furthermore observed that EEA changes of the synthetic four-species

communities exhibited strong differences between substrates (Figure 4; Appendix S1: Figure

S5). EEA changes showed sharp decreases, down to -100%, and increases, up to 385%, at 0 h

after mixing. Substrates with glycosidic bonds like 4-MUB-N-acetyl-β-D-glucosaminide

(NAG), 4-MUB-β-D-cellobioside (CB) and 4-MUB-β-D-xylopyranoside (XYL) exhibited the

highest relative increases in EEA (see Appendix S1: Table S2 for complete list of substrate

abbreviations). The EEA changes for each one substrate within a treatment group, however,

were typically consistent (< 50% EEA proportional change from the average) for 33 of the 44

(75%) community-substrate combinations. This indicated that, in most cases, community

behaviour was not strongly influenced by the excluded fifth isolate of each group. The

consistency was more prominent for substrates with ester bonds like 4-MUB acetate (ACE),

4-MUB-butyrate (BUT) and 4-MUB oleate (OLE). A complete overview of cleaved substrate

concentrations for isolates, communities and estimated community EEAs for each treatment

group, can be found in the supplement (Appendix 1: Tables S6, S7, S8, S9). When comparing

EEA changes between the groups (Appendix S1: Figure S5B-E), we observed that Pdis

communities had relatively small to no changes in EEA: 76% of community-substrate

combinations of group Pdis-Msim and Pdis-Mdis remained between +50% and -50% EEA

proportional change. Contrarily, communities of Psim groups differed greatly in activities

compared to their constituent isolates. Group Psim-Msim exhibited most of the largest EEA

increases (up to 330%) with 27% of the group's 55 community-substrate combinations

measuring > +50% EEA proportional change. Group Psim-Mdis communities recorded the

greatest amount of negative EEA proportional changes (53% of the group's 55 community-

substrate combinations having < -50% EEA proportional change) with the maximum EEA

proportional change increase at only 5.4%. These results show that phylogenetically similar

communities exhibited more variable and genus-specific responses than phylogenetically

dissimilar communities.

Chemical structure differences of substrates influence the strength of diversity

effects

364   Investigating the datasets for all substrates individually using linear models, we

observed broad differences in the relationships of EEAs to the metabolic and phylogenetic

diversity between them (Appendix S1: Table S10). When controlling for substrates, the

diversity treatments have a consistent pattern of influence, positive for phylogenetic diversity

and negative for metabolic diversity (Figure 5). For example, CB showed a significant

positive relationship with phylogenetic diversity (1.029 ± 0.444, estimate $\pm$ std. error;

Adj. $R^2 = 0.520$, $p = 0.034$) and for metabolic diversity, a significantly negative relationship (-

1.604 ± 0.441, estimate $\pm$ std. error; Adj. $R^2 = 0.520$, $p = 0.002$). The other two

polysaccharide-analogue substrates, XYL and NAG, behaved similarly to CB (Appendix S1:

Table S10). We observed that the strength of the effect was related with the chemical

structure of the substrates (Figure 5). Esters (ACE, BUT, OLE) exhibited no significant

relationship with diversity treatments, with linear slopes ranging from -0.095 to 0.038 for

phylogenetic diversity and -0.227 to -0.007 for metabolic diversity. Monosaccharide (aG, bG)

linear slopes ranged from 0.211 to 0.368 and -0.760 to -0.336, a relationship that was overall

similar but weaker than for the polysaccharides (CB, XYL, NAG), whose linear slopes were

1.029 to 1.470 and -2.019 to -1.604 for phylogenetic and metabolic diversity, respectively. On

average, by controlling for substrates alone, we observed that phylogenetic diversity had a

positive effect on EEA changes (0.643 ± 0.193, estimate $\pm$ std. error; Adj. $R^2 = 0.645$,

$p = 0.004$) (Table S10). Contrastingly, metabolic diversity had a negative effect (-0.897 ±

0.191, estimate ± std. error; Adj. $R^2 = 0.645$, $p = 0.0003$).

Effects of diversity in EEA changes over 24 h

We further investigated how the community EEA changed with increasing time after

community mixing, by comparing EEA results from time points of 0 h, 6 h and 24 h (Figure

6). Significant positive changes ranged from 10 - 240% and negative changes ranged from -10

- -137%. Both Msim groups as well as Pdis-Mdis exhibited negative or non-significant

changes over the 24 h. However, the Psim-Mdis group showed both significantly negative and

positive changes between EEAs measured over 24 h after mixing. Changes in the Pdis

communities' composition within 24 h did not explain the observed activity changes; 16S

rRNA gene amplicon sequencing results showed that all the introduced species appear to

persist in the community cultures during 24 h post-mixing, with only close to ~1-fold change

of relative abundance for most species and a 2-fold change for two species

(Appendix S1: Figure S6). Thus, the observed differences in changes in EEA over time

between diversity treatments were not the result of a shifting community composition, but of

diversity treatment effects.

Substrate acclimation in EEA assays

Acclimation describes any phenotypic plasticity an organism can undergo to adjust to

changes in its environment. Effects of substrate acclimation were observed directly by

measuring the increase of EEA due to potential upregulation of exoenzyme production,

during the incubation with a substrate. Here we show the results from the assay with aG, as an

example. Comparing the timepoint of 0 h to the acclimated cultures timepoint of 4.5 h, we

observed increases of the EEAs for both. However, these increases were much higher for

communities, with ~5-fold increases on average, than for isolates, with average increases of

~1-fold (Figure 7). This was also reflected in the highest changes observed: For isolate

activities in the aG substrate assay, 2.3 times increase, from 0.110 ± 0.003 to

0.255 ± 0.008 µM min^{-1}, was the largest change observed. For community activities, increases

up to 10 times, from 0.020 ± 0.001 to 0.199 ± 0.012 µM min^{-1} were observed, making

community acclimation capacity almost 5 times higher compared to isolates. This increase in

activity did not match the EEA changes observed for communities after 6 h of mixing without

the addition of substrates. Since both isolate and community activities were normalised to

biomass, the increased activity in communities could not be attributed to differences in cell

density. In addition, the acclimation effect could not have been the result of species

abundance changes since even the most active of the isolates only had ~2 times increase of

EEA due to acclimation (0.225 ± 0.006 to 0.454 ± 0.003 µM min^{-1}).

Discussion

Using synthetic microbial communities, we showed that both phylogenetic and

metabolic diversity influence community ecological functioning. Because both diversity

aspects are so intricately linked, we used a multifactorial treatment experiment that allowed us

to examine what the effects of each diversity were when controlling for the other. Since

metabolic diversity has a direct impact on ecological functions, we expected that higher

metabolic diversity would lead to higher functioning of bacterial communities. What we see is

that the interaction between diversity treatments causes contrasting effects, which did not

meet our expectations.

Metabolic diversity has either no effect, for Pdis communities, or a negative

relationship, for Psim communities for both community TAC and community EEA %

proportional changes (Figure 3B, Appendix S1: Figure S4B). In Pdis communities,

differences in nutrient acquisition and interaction strategies of different species could be

playing a role for the lack of effect of metabolic diversity. Such a mixture of species can

engage in varying competitive or cooperative interactions (Gross 2008; Violle *et al.* 2011).

This complementary set of community species interactions might be masking any observable

influence of metabolic diversity (Maynard, Crowther and Bradford 2017). On the other hand,

in Psim communities, related species are likely to be reacting similarly to each other and

resource competition. However, increased conspecific competition can result in different

outcomes. With lower metabolic diversity, upregulation of exoenzyme activity as a response

to competition for resource exploitation, might be translating to increased community

functioning. The mechanism behind this might be how the cost:benefit ratio of exoenzyme

production manifests in low diversity communities if they are all producers and strong

competitors (Allison *et al.* 2011). Consequently, higher metabolic diversity results in lower

community functioning as species can diversify their resource preferences and therefore do

not contribute to the same substrate degradation in the assays. Moreover, species of different

genera might have distinct ways of interacting with species of their own genus, further

influencing competitive outcomes (Mayfield and Levine 2010; Maynard, Crowther and

Bradford 2017).

We also predicted that as phylogenetic diversity is based on a neutral genetic marker,

it would have no effect on community ecological functioning. We found that for both TAC

and EEA % changes, phylogenetic diversity has a positive relationship with the ecological

functioning of Mdis communities and a negative relationship for Msim communities (Figure

3A; Appendix S1: Figure S4A). In the Mdis communities, organisms can engage in a variety

of metabolic functions. This increases the capacity for different kinds of community species

interactions (Weber and Strauss 2016), facilitated by phylogenetic diversity and coexistence.

More diverse species interactions might translate into more cooperative communities,

metabolic complementarity and increased overall functioning. Contrastingly, Msim

communities rely on a limited number of metabolic functions. The limited metabolic

functions do not allow for elaborate interaction strategies, such as cooperation, to be enacted.

As such, any influence of phylogenetic diversity is severely decreased. In this setup, resource

competition and antagonistic interactions with multiple species cause a negative relationship

to emerge.

Lastly, we hypothesised that because Psim community species would have less diverse

strategies in responding to situations of competition or cooperation, metabolic diversity would

have a much stronger effect in their ecological functioning, compared to their Pdis community

member counterparts. Indeed, this is true as we show that Psim communities had a strong,

albeit negative relationship with EEA changes and TAC, while Pdis communities had none.

Moreover, the consistency of results between the single and multifunctional measures of

EEAs and TAC increases the confidence of the diversity relationships we have observed. This

should not be surprising, as we already established a deep connection between phylogenetic

and metabolic diversity, which would explain why neither of them is the sole driver of

community ecological functions.

The contrasting diversity effects were also revealed when controlling for the substrates

in the treatments. We identified as a general trend that phylogenetic diversity has a positive

effect on EEA changes and metabolic diversity has a negative influence (Figure 5). In

addition to that, we saw that the diversity effects were operating on a strength gradient, with

groups of biochemically similar substrates appearing in distinct positions along that gradient

(Figure 5). Such differences between substrates and groups of substrates can be related to

inherent variations in exoenzyme production, maintenance and regulation and how different

classes of enzymes behave to break down their substrates. For example, polysaccharides are

targeted by hydrolytic exoenzymes that are produced on-demand (Warren 1996; Lynd *et al.*

2002). In contrast, enzymes involved in the degradation of fatty acids belonging to the ester

group of substrates are part of bacterial core metabolism, and thus can be continuously

produced. Our results show that stronger diversity effects are more important to ecosystem

functions that rely on environmental factors, such as the temporal or spatial presence of plant-

derived polysaccharides. Further adding to the complexity of experimental outcomes, time

also influences different substrates and diversity treatments in different ways (Figure 6).

Hence, not only the diversity of ecosystem functions tested (Huston 1997, 2000; Tilman

1997; Loreau 2000) but also their underlying biochemical mechanisms have a great effect on

the outcomes of biodiversity research and the choice, effect and timing of different functions

in experimental designs needs to considered.

To understand the very strong contrasting changes in EEAs between the Psim groups,

their genus-specific capacity for co-existence with its related species is crucial. Psim-Msim

communities contained *Flavobacteria* species which are usually non-motile organisms, prefer

aquatic environments and have strong differences between species. The genus includes

generalist species that can use a wide range of carbohydrates, to more specialist

asaccharolytic species that prefer nitrogenous substrates (i.e. amino acids) (Bernardet and

Bowman 2006). These physiological characteristics might allow the *Flavobacteria*

communities to increase their EEA as different species could be utilising different carbon

sources in their growth medium. Communities of these specialist *Flavobacteria* could be

facilitating community species interactions that translate into increases in their community

EEAs. In contrast, the *Pseudomonas* species in Psim-Mdis communities are generally motile-

planktonic bacteria, ubiquitous in almost all oxic environments, including soil and water, due

to their expansive metabolic repertoire, simple nutritional requirements and rapid growth rates

(Moore *et al.* 2006). However, the reduced capacity of many *Pseudomonas* species to

withstand starvation (Moore *et al.* 2006) might lead them to adapt more selfish nutrient

acquisition strategies (Reintjes *et al.* 2017, 2019; Arnosti, Reintjes and Amann 2018). As

generalist species, *Pseudomonas* might be more competitive, resulting in *Pseudomonas*

communities with lower EEAs than their respective isolates. Thus, bacteria with a broad

substrate range might not benefit when living in communities of the same genus but of

dissimilar metabolism. Contrastingly, in the Pdis communities these genus-specific effects in

nutrient acquisition strategies can be balanced due to the increased species diversity.

Nevertheless, even competing generalists like *Pseudomonas* species will profit from species

interactions when incubated for longer time in the presence of certain substrates. Using the

Psim-Mdis communities to run EEA substrate acclimated vs. non-acclimated assays (Figure

7), we identified that the amplifying effect of substrate acclimation on the EEAs is much

stronger for the communities than for the constituent isolates on substrate aG. In addition,

since the increased EEAs are not present in the non-acclimated 6 h timepoint communities,

we can conclude that the effect was indeed substrate acclimatisation of the bacteria and not an

effect of mixing the community cultures. Consequently, the observed acclimation effect was

the result of community species interactions that enhanced species' ability to acclimate to the

substrate. These observations link back to our first hypothesis, where we predict that the

metabolic diversity of the group will enhance the functioning of communities. But

oversimplifying the experimental design and just looking at one substrate or one group, would

have let to an underestimation of the complexity of diversity-ecosystem functioning

relationships, possibly explaining as to why we see studies showing opposite effects of

biodiversity (Balvanera *et al.* 2006).

Altogether, our results concur with previous studies that focus on separating the

different aspects of biodiversity and examining the relationship between biodiversity and

ecosystem functioning (Griffiths *et al.* 2001; Balvanera *et al.* 2006; Zhang and Zhang 2006;

Naeem *et al.* 2009; Jousset *et al.* 2011; Purschke *et al.* 2013; Thompson, Davies and Gonzalez

2015; Gamfeldt and Roger 2017). A meta-analysis demonstrated that biodiversity effects

depend on the organisation level: with positive effects at the community level, positive but

weaker effects at the ecosystem level and negative effects at the population level (Balvanera

et al. 2006). Experiments done by Griffiths *et al.* (Griffiths *et al.* 2001) recognised no

consistent effect from biodiversity on soil functions. Thus, the need for more research

including the effects of the environmental context, like nutrient availability, on the

relationship between biodiversity and ecosystem functioning was recognized (Zhang and

Zhang 2006). Our study fills this gap by focussing not only on the phylogenetic and metabolic

diversity of organisms, but also measuring a broad diversity of activities controlled by

different underlying mechanisms of enzyme regulation. Such a complex approach is required

to tackle the complex dimensionality of biodiversity research. We show that biodiversity had

a significant impact on most EEAs, but the strength of the effect depended on the substrate

context. This might explain why the relationship between biodiversity and community

ecosystem functioning cannot always be predicted to be positive as previously observed

(Aarssen 1997; Tilman 1997; Huston 2000; Carroll, Cardinale and Nisbet 2011). Natural,

larger scale experiments struggle with introducing different levels of complexity in their

experimental design (Cairns *et al.* 2018). Looking forward, we should aim to understand how

diversity effects can mechanistically influence ecosystems, synthetic and natural (Fox and

Harpole 2008; Naeem *et al.* 2009; Turnbull *et al.* 2013), and not take the effects as granted

(Shade 2017).

Acknowledgements

We gratefully acknowledge the support of the German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) (DFG FZT 118, 202548816). This project has been conducted in the framework of the iDiv-Flexpool – the internal funding mechanism of iDiv. Martin Taubert was funded by the DFG under Germany's Excellence Strategy – EXC 2051 – Project-ID 390713860. We would like to express our gratitude to Patricia Geesink for the bacterial isolation work under the framework of the CRC AquaDiva of the Friedrich Schiller University Jena, funded by the DFG (DFG SFB 1076). Additionally, we would like to thank Will A. Overholt and Lijuan Yan for their invaluable statistical analysis input.

Author Contributions

KK and SWH secured funding for the project. CX, MT, and KK designed the experiments. CX performed the experiments and carried out all assays and molecular analyses; the data analysis was performed by CX closely supported by MT. Statistical analysis was performed by CX supported by MT and SWH. All co-authors assisted in interpreting the results; CX initiated the manuscript writing, which was finalized with substantial contributions from MT, KK and SWH.

Competing Interests

The authors declare no conflicts of interest.

References

Aarssen LW. High productivity in grassland ecosystems: effected by species diversity or

productive species? test. *Oikos* 1997;**80**:183–4.

Alexandrou MA, Cardinale BJ, Hall JD *et al.* Evolutionary relatedness does not predict

competition and co-occurrence in natural or experimental communities of green algae.

Proc Biol Sci 2015;**282**:20141745.

Allison SD, Weintraub MN, Gartner TB *et al.* Evolutionary-Economic Principles as

Regulators of Soil Enzyme Production and Ecosystem Function. In: Shukla G, Varma

A (eds.). *Soil Enzymology*. Berlin, Heidelberg: Springer Berlin Heidelberg, 2011, 229–

43.

Arnosti C, Reintjes G, Amann R. A mechanistic microbial underpinning for the size-reactivity

continuum of dissolved organic carbon degradation. *Marine Chemistry* 2018;**206**:93–

9.

Balvanera P, Pfisterer AB, Buchmann N *et al.* Quantifying the evidence for biodiversity

effects on ecosystem functioning and services. *Ecology Letters* 2006;**9**:1146–56.

Bernardet J-F, Bowman JP. The Genus Flavobacterium. *The Prokaryotes*. New York, NY:

Springer New York, 2006, 481–531.

Borg I, Groenen PJF. *Modern Multidimensional Scaling: Theory and Applications*. Springer,

New York, NY, 2005.

Cadotte MW, Cavender-Bares J, Tilman D *et al.* Using phylogenetic, functional and trait

diversity to understand patterns of plant community productivity. *PLoS One*

2009;**4**:e5695.

Cairns J, Jokela R, Hultman J *et al.* Construction and Characterization of Synthetic Bacterial

Community for Experimental Ecology and Evolution. *Frontiers in Genetics* 2018;**9**,

DOI: 10.3389/fgene.2018.00312.

Cardinale BJ, Duffy JE, Gonzalez A *et al.* Biodiversity loss and its impact on humanity.

Nature 2012;**486**:59–67.

Carroll IT, Cardinale BJ, Nisbet RM. Niche and fitness differences relate the maintenance of

diversity to ecosystem function. *Ecology* 2011;**92**:1157–65.

Drescher K, Nadell CD, Stone HA *et al.* Solutions to the public goods dilemma in bacterial

biofilms. *Curr Biol* 2014;**24**:50–5.

Edgar R. UCHIME2: improved chimera prediction for amplicon sequencing. *bioRxiv*

2016:074252.

Evans R, Alessi AM, Bird S *et al.* Defining the functional traits that drive bacterial

decomposer community productivity. *The ISME Journal* 2017;**11**:1680–7.

Finlay BJ, Maberly SC, Cooper JI. Microbial Diversity and Ecosystem Function. *Oikos*

1997;**80**:209.

Flynn DFB, Mirotchnick N, Jain M *et al.* Functional and phylogenetic diversity as predictors

of biodiversity–ecosystem-function relationships. *Ecology* 2011;**92**:1573–81.

Folse HJ 3rd, Allison SD. Cooperation, competition, and coalitions in enzyme-producing

microbes: social evolution and nutrient depolymerization rates. *Front Microbiol*

2012;**3**:338.

Fox JW, Harpole WS. Revealing How Species Loss Affects Ecosystem Function: The Trait-

Based Price Equation Partition. *Ecology* 2008;**89**:269–79.

Frossard A, Gerull L, Mutz M *et al.* Disconnect of microbial structure and function: enzyme

activities and bacterial communities in nascent stream corridors. *The ISME journal*

2012;**6**:680–91.

Galand PE, Pereira O, Hochart C *et al.* A strong link between marine microbial community

composition and function challenges the idea of functional redundancy. *ISME J*

2018;**12**:2470–8.

Gamfeldt L, Roger F. Revisiting the biodiversity–ecosystem multifunctionality relationship.

Nature Ecology & Evolution 2017;**1**:0168.

Geesink P, Tyc O, Küsel K *et al.* Growth promotion and inhibition induced by interactions of

groundwater bacteria. *FEMS Microbiology Ecology* 2018;**94**:1–8.

Gerhold P, Cahill JF, Winter M *et al.* Phylogenetic patterns are not proxies of community

assembly mechanisms (they are far better). Venail P (ed.). *Funct Ecol* 2015;**29**:600–

14.

Germain RM, Weir JT, Gilbert B. Species coexistence: macroevolutionary relationships and

the contingency of historical interactions. *Proc Biol Sci* 2016;**283**:20160047.

Goberna M, Verdú M. Predicting microbial traits with phylogenies. *The ISME Journal*

2016;**10**:959–67.

Griffiths BS, Ritz K, Wheatley R *et al.* An examination of the biodiversity–ecosystem

function relationship in arable soil microbial communities. *Soil Biology and*

Biochemistry 2001;**33**:1713–22.

Gross K. Positive interactions among competitors can produce species-rich communities. *Ecol*

Lett 2008;**11**:929–36.

Herlemann DPR, Labrenz M, Jürgens K *et al.* Transitions in bacterial communities along the

2000 km salinity gradient of the Baltic Sea. *ISME Journal* 2011;**5**:1571–9.

Huston MA. Hidden treatments in ecological experiments: re-evaluating the ecosystem

function of biodiversity. *Oecologia* 1997;**110**:449–60.

Huston MA. No Consistent Effect of Plant Diversity on Productivity. *Science*

2000;**289**:1255a–1255.

Johnson DR, Pomati F. A brief guide for the measurement and interpretation of microbial

functional diversity. *Environ Microbiol* 2020;**22**:3039–48.

Jousset A, Schmid B, Scheu S *et al.* Genotypic richness and dissimilarity opposingly affect

ecosystem functioning. *Ecology Letters* 2011;**14**:537–45.

Kowalchuk GA, Stephen JR. Ammonia-oxidizing bacteria: a model for molecular microbial

ecology. *Annu Rev Microbiol* 2001;**55**:485–529.

Kozich JJ, Westcott SL, Baxter NT *et al.* Development of a Dual-Index Sequencing Strategy

and Curation Pipeline for Analyzing Amplicon Sequence Data on the MiSeq Illumina

Sequencing Platform. *Applied and Environmental Microbiology* 2013;**79**:5112–20.

Krause S, Le Roux X, Niklaus PA *et al.* Trait-based approaches for understanding microbial

biodiversity and ecosystem functioning. *Frontiers in Microbiology* 2014;**5**:1–10.

Kumar S, Herrmann M, Thamdrup B *et al.* Nitrogen Loss from Pristine Carbonate-Rock

Aquifers of the Hainich Critical Zone Exploratory (Germany) Is Primarily Driven by

Chemolithoautotrophic Anammox Processes. *Frontiers in Microbiology* 2017;**8**:1–13.

Kumar S, Stecher G, Li M *et al.* MEGA X: Molecular Evolutionary Genetics Analysis across

Computing Platforms. Battistuzzi FU (ed.). *Molecular Biology and Evolution*

2018;**35**:1547–9.

Küsel K, Totsche KU, Trumbore SE *et al.* How deep can surface signals be traced in the

critical zone? Merging biodiversity with biogeochemistry research in a central German

Muschelkalk landscape. *Frontiers in Earth Science* 2016;**4**:1–18.

Le Bagousse-Pinguet Y, Soliveres S, Gross N *et al.* Phylogenetic, functional, and taxonomic

richness have both positive and negative effects on ecosystem multifunctionality. *Proc*

Natl Acad Sci U S A 2019;**116**:8419–24.

Litchman E, Klausmeier C a. Trait-Based Community Ecology of Phytoplankton. *Annual*

Review of Ecology, Evolution, and Systematics 2008;**39**:615–39.

Loreau M. Biodiversity and ecosystem functioning: recent theoretical advances. *Oikos*

2000;**91**:3–17.

Losos JB. Phylogenetic niche conservatism, phylogenetic signal and the relationship between

phylogenetic relatedness and ecological similarity among species. *Ecol Lett*

2008;**11**:995–1003.

Lynd LR, Weimer PJ, van Zyl WH *et al.* Microbial Cellulose Utilization: Fundamentals and

Biotechnology. *Microbiol Mol Biol Rev* 2002;**66**:506–77.

Malik AA, Martiny JBH, Brodie EL *et al.* Defining trait-based microbial strategies with

consequences for soil carbon cycling under climate change. *ISME J* 2019, DOI:

10.1038/s41396-019-0510-0.

Martiny AC, Treseder K, Pusch G. Phylogenetic conservatism of functional traits in

microorganisms. *The ISME journal* 2013;**7**:830–8.

Mayfield MM, Levine JM. Opposing effects of competitive exclusion on the phylogenetic

structure of communities. *Ecol Lett* 2010;**13**:1085–93.

Maynard DS, Crowther TW, Bradford MA. Competitive network determines the direction of

the diversity–function relationship. *Proceedings of the National Academy of Sciences*

2017:201712211.

McDonald JH. *Handbook of Biological Statistics*. 3rd ed. Baltimore, Maryland: Sparky House

Publishing, 2014.

Mokany K, Ash J, Roxburgh S. Functional identity is more important than diversity in

influencing ecosystem processes in a temperate native grassland. *J Ecol* 2008;**96**:884–

93.

Moore ERB, Tindall BJ, Martins Dos Santos VAP *et al.* Nonmedical: Pseudomonas. *The*

Prokaryotes. New York, NY: Springer New York, 2006, 646–703.

Morrissey EM, Mau RL, Schwartz E *et al.* Phylogenetic organization of bacterial activity. *The*

ISME Journal 2016;**10**:2336–40.

Mukherjee S, Stamatis D, Bertsch J *et al.* Genomes OnLine database (GOLD) v.7: updates

and new features. *Nucleic Acids Research* 2019;**47**:D649–59.

Murtagh F, Legendre P. Ward's Hierarchical Agglomerative Clustering Method: Which

Algorithms Implement Ward's Criterion? *J Classification* 2014;**31**:274–95.

Naeem S, Bunker DE, Hector A *et al. Biodiversity, Ecosystem Functioning, and Human*

Wellbeing. Oxford University Press, 2009.

Nealson KH, Popa R. Metabolic diversity in the microbial world: relevance to exobiology.

Micro-Organisms and Earth Systems. Cambridge University Press, 2005, 151–72.

Pascual-García A, Bonhoeffer S, Bell T. Metabolically cohesive microbial consortia and

ecosystem functioning. *Philos Trans R Soc Lond B Biol Sci* 2020;**375**:20190245.

Paula FS, Rodrigues JLM, Zhou J *et al.* Land use change alters functional gene diversity,

composition and abundance in Amazon forest soil microbial communities. *Molecular*

Ecology 2014;**23**:2988–99.

Petchey OL, Gaston KJ. Functional diversity: back to basics and looking forward. *Ecol Lett*

2006;**9**:741–58.

Pruesse E, Peplies J, Glöckner FO. SINA: Accurate high-throughput multiple sequence

alignment of ribosomal RNA genes. *Bioinformatics* 2012;**28**:1823–9.

Purschke O, Schmid BC, Sykes MT *et al.* Contrasting changes in taxonomic, phylogenetic

and functional diversity during a long-term succession: insights into assembly

processes. Fridley J (ed.). *Journal of Ecology* 2013;**101**:857–66.

Quast C, Pruesse E, Yilmaz P *et al.* The SILVA ribosomal RNA gene database project:

improved data processing and web-based tools. *Nucleic Acids Research*

2012;**41**:D590–6.

R Core Team. R: A Language and Environment for Statistical Computing. 2019.

Reintjes G, Arnosti C, Fuchs B *et al.* Selfish, sharing and scavenging bacteria in the Atlantic

Ocean: a biogeographical study of bacterial substrate utilisation. *The ISME Journal*

2019;**13**:1119–32.

Reintjes G, Arnosti C, Fuchs BM *et al.* An alternative polysaccharide uptake mechanism of

marine bacteria. *The ISME Journal* 2017;**11**:1640–50.

Rivett DW, Scheuerl T, Culbert CT *et al.* Resource-dependent attenuation of species

interactions during bacterial succession. *The ISME journal* 2016;**10**:2259–68.

Roger F, Bertilsson S, Langenheder S *et al.* Effects of multiple dimensions of bacterial

diversity on functioning, stability and multifunctionality. *Ecology* 2016;**97**:2716–28.

Salles JF, Poly F, Schmid B *et al.* Community niche predicts the functioning of denitrifying

bacterial assemblages. *Ecology* 2009;**90**:3324–32.

Schloss PD, Westcott SL, Ryabin T *et al.* Introducing mothur: Open-Source, Platform-

Independent, Community-Supported Software for Describing and Comparing

Microbial Communities. *Applied and Environmental Microbiology* 2009;**75**:7537–41.

Schwab VF, Herrmann M, Roth VN *et al.* Functional diversity of microbial communities in

pristine aquifers inferred by PLFA- and sequencing-based approaches. *Biogeosciences*

2017;**14**:2697–714.

Shade A. Diversity is the question, not the answer. *The ISME Journal* 2017;**11**:1–6.

Shade A, Dunn RR, Blowes SA *et al.* Macroecology to Unite All Life, Large and Small.

Trends in Ecology & Evolution 2018;**33**:731–44.

Smith P, Schuster M. Public goods and cheating in microbes. *Curr Biol* 2019;**29**:R442–7.

Srivastava DS, Cadotte MW, Macdonald AAM *et al.* Phylogenetic diversity and the

functioning of ecosystems. *Ecology Letters* 2012;**15**:637–48.

Thompson PL, Davies TJ, Gonzalez A. Ecosystem Functions across Trophic Levels Are

Linked to Functional and Phylogenetic Diversity. *PLOS ONE* 2015;**10**:e0117595.

Tilman D. Distinguishing between the Effects of Species Diversity and Species Composition.

Oikos 1997;**80**:185.

Turnbull LA, Levine JM, Loreau M *et al.* Coexistence, niches and biodiversity effects on

ecosystem functioning. *Ecology Letters* 2013;**16**:116–27.

Turner S, Pryer KM, Miao VPW *et al.* Investigating Deep Phylogenetic Relationships among

Cyanobacteria and Plastids by Small Subunit rRNA Sequence Analysis. *The Journal*

of Eukaryotic Microbiology 1999;**46**:327–38.

Venail P, Gross K, Oakley TH *et al.* Species richness, but not phylogenetic diversity,

influences community biomass production and temporal stability in a re-examination

of 16 grassland biodiversity studies. Fox C (ed.). *Funct Ecol* 2015;**29**:615–26.

Violle C, Nemergut DR, Pu Z *et al.* Phylogenetic limiting similarity and competitive

exclusion. *Ecol Lett* 2011;**14**:782–7.

Wang Y, Zhang R, He Z *et al.* Functional Gene Diversity and Metabolic Potential of the

Microbial Community in an Estuary-Shelf Environment. *Frontiers in Microbiology*

2017;**8**:1–12.

Ward JH. Hierarchical Grouping to Optimize an Objective Function. *J Am Stat Assoc*

1963;**58**:236–44.

Warren RA. Microbial hydrolysis of polysaccharides. *Annu Rev Microbiol* 1996;**50**:183–212.

Weber MG, Strauss SY. Coexistence in Close Relatives: Beyond Competition and

Reproductive Isolation in Sister Taxa. *Annu Rev Ecol Evol Syst* 2016;**47**:359–81.

Zhang Q-G, Zhang D-Y. Resource availability and biodiversity effects on the productivity,

temporal variability and resistance of experimental algal communities. *Oikos*

2006;**114**:385–96.

Zhang Z, Schwartz S, Wagner L *et al.* A Greedy Algorithm for Aligning DNA Sequences.

Journal of Computational Biology 2000;**7**:203–14.

Figure legends

Figure 1

Metabolic and phylogenetic classification and grouping of groundwater isolates. EEAs

are shown for twenty-seven groundwater isolates incubated with eleven fluorescent substrate

analogues. The heatmap scale shows the concentration of substrate that was broken down in

the EEA assays. Empty (white) boxes denote activity outside the detectable range. The left

dendrogram shows clusters of isolates with similar metabolic traits based on Euclidean

hierarchical diversity distances. The topmost dendrogram shows hierarchical clustering of

substrates that produced similar activity patterns in the assays. The assignment of isolates to

the communities of each treatment group is depicted to the right of the heatmap. Double-

coloured boxes indicate isolates that were assigned to more than one group. In the legend, the

colours indicate the community group and the phylogenetic (P) and metabolic (M) traits of the

communities are indicated as similar (sim) or dissimilar (dis).

Figure 2

Individual communities plotted against the phylogenetic and metabolic diversity within

and between groups. Colours indicate the community groups; red for Psim-Msim group,

green for Psim-Mdis group, blue for Pdis-Msim group and purple for Pdis-Mdis group.

Phylogenetic and metabolic diversity metrics have been z-score normalised. Variation of

diversity scores exists within groups, but the groups are cleanly separated from each other

(see Appendix 1: Table S3 for statistical significance).

Figure 3

Effects of phylogenetic diversity (A) and metabolic diversity (B) on the total available

carbon (TAC) of communities. Black regression lines show the effect of each treatment

when controlling for the dissimilar groups of the second treatment. Grey regression lines

show the effect of each treatment when controlling for the similar groups of the second

treatment. Statistical significance for each linear regression is shown in each plot, with the

respective coloured text (grey or black). Coloured squares indicate community groups: Psim-

Msim in red, Psim-Mdis in green, Pdis-Msim in blue and Pdis-Mdis in purple. The y axes

have been log2 transformed and the x axes have been z-scale transformed.

Figure 4

Heatmap of community EEA % proportional changes. The heatmap scale depicts

proportional changes from -100 to 100%. Communities that exhibited changes over 100%

have their percent change value shown in the heatmap matrix. Red colours depict a higher

activity of the community; blue colours depict a lower activity of the community than

estimated, based on the activities of the individual isolates. White boxes show measurements

outside the detection limits. On the right, the label for each row indicates the community and

on the left the group from where the activities are taken from; red for Psim-Msim group,

green for Psim-Mdis group, blue for Pdis-Msim group and purple for Pdis-Mdis group

communities. Columns indicate the substrates used in the EEA assays. The dendrogram

depicts hierarchical clustering of each substrate's activities, showing groups of substrates with

similar activity responses.

Figure 5

Substrate chemistry drives the strength and direction of phylogenetic and metabolic

diversity treatment effects. (A) The linear regression model slopes for all 11 substrates and

the average for all substrates (Appendix 1: Table S10) are plotted here as points. In (B) the

average for all substrates, community EEA, is positively related with phylogenetic diversity

($p < 0.001$). In (C) the average for all substrates, community EEA, is negatively related with

metabolic diversity ($p < 0.001$). Coloured points in (A) indicate different substrates and the

average (red). Biochemically similar substrates are circled in dashed ellipses: blue for fatty

acids, orange for monosaccharides and green for polysaccharides. Coloured squares in (B)

and (C) indicate community groups: Psim-Msim in red, Psim-Mdis in green, Pdis-Msim in

blue and Pdis-Mdis in purple.

Figure 6

Temporal changes of EEA % proportional changes between communities and isolates

from 0 h to 24 h after community mixing. Rows show community groups and columns

show the substrate where the activity change was noted. The average changes between the

timepoints are indicated by the colour of the arrows and their statistical significance

(Student's *t*-test, 2-tailed) showed on top of the arrows. P values below 0.001 are shown as

<0.001 and values over 0.05 are marked as non-significant (*ns*).

Figure 7

Substrate acclimation effect on isolates and communities. EEA of Psim-Mdis group

isolates and communities on 4-MUB-α-D-glucopyranoside (aG) are shown as an example of

the substrate acclimation effect. In green are shown the non-acclimated EEAs of cultures as a

fold-change of 1, to which all other activities were normalised. In orange, the cultures

received the substrate at the same time as the previous cultures and left to acclimate for 4.5 h.

Lastly, the EEAs of cultures that had been mixed for 6 h but not left to acclimate, as shown in

blue. Bars depict average fold-changes in the EEA of isolates and communities and error bars

Phylogenetic vs. metabolic diversity

show the standard deviation (n = 5). Community EEAs showed a ~5-fold increase when

acclimated with the substrate and isolates only showed a ~1-fold increase.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7