Phylogenetic and metabolic diversity have contrasting effects on the ecological functioning of bacterial communities

Constantinos Xenophontos[1,2], Martin Taubert[1], W. Stanley Harpole[2,3,4], Kirsten Küsel[1,2]

1. Institute of Biodiversity, Friedrich Schiller University Jena, Jena 07743, Germany
2. German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, 04103 Leipzig, Germany
3. Department of Physiological Diversity, Helmholtz-Center for Environmental Research (UFZ), 04318 Leipzig, Germany
4. Institute of Biology, Martin Luther University of Halle-Wittenberg, 06120 Halle, Germany

Correspondence:

Kirsten Küsel kirsten.kuesel@uni-jena.de

Abstract

Biodiversity can affect microbial ecosystem functioning because greater numbers of species,

and their associated genetic and metabolic differences, can contribute at different times and

contexts to overall ecosystem functioning. Quantifying the relative contributions of microbial

species to ecosystem functioning is challenging, because of the distinct mechanisms that are

associated with their phylogenetic diversity vs. their metabolic diversity. We used synthetic

bacterial communities to test the independent effects of two aspects of biodiversity,

phylogenetic and metabolic diversity, on community ecological functioning. We expected that

metabolic diversity, as a direct measure of species functional traits, would be associated with

greater ecological function. Contrastingly, since phylogenetic diversity is inherently a metric

based on neutral genetic markers, it should have no influence. Moreover, we hypothesised that

phylogenetically related species would impact species interactions and coexistence, therefore

amplifying the influence of metabolic diversity. To study these effects, we used bacterial

isolates to construct communities with different diversity traits and employed exoenzyme

activities (EEAs) and total available carbon (TAC) from substrates as proxies of bacterial

functioning. Using linear models to examine the effects of one diversity treatment while

controlling for the other, we found that they did not meet most of our expectations. Phylogenetic

diversity strongly influenced community ecological functioning and metabolic diversity

exhibited significantly negative relationships. When controlling for different substrates, EEAs

increased along with phylogenetic diversity but decreased with metabolic diversity. In addition,

the strength of the diversity effects was depended on the substrates. EEAs on carbohydrates

showed a strong positive relationship with phylogenetic diversity and a strong negative

relationship with metabolic diversity, while EEAs on fatty acids typically showed no

relationship at all. Furthermore, EEAs of phylogenetically similar groups were strongly affected

by within-genus interactions. We concluded that both phylogenetic and metabolic diversity

have complicated and contrasting effects, which furthermore depended on the ecological

function under examination – diversity effects were depended on substrate chemistry and the

molecular mechanisms related to each substrate's degradation. This highlights the importance

of investigating individual aspects of biodiversity but also their interactions and the variations

within the biological mechanisms and processes that underscore them, to improve ecological

theory.

Keywords

49 ecosystem functions; phylogenetic diversity; metabolic diversity; biodiversity; bacteria;

50 synthetic communities; exoenzymes

51 Introduction

More diverse ecosystems are typically more productive (Lehman and Tilman 2000, Byrnes et

al. 2014, Maron et al. 2018), perform more functions (Hector and Bagchi 2007, Maron et al.

2018), are more resistant to disturbances (Ives and Carpenter 2007, Gross et al. 2014) and

increase community temporal stability (Lehman and Tilman 2000). Bacterial activities are

inextricable from ecosystem functioning (Finlay et al. 1997) and recent research has highlighted

that microbial diversity contributes to the functioning of ecosystems, challenging earlier ideas

that microbes might have a high degree of functional redundancy (Yin et al. 2000, Louca et al.

2018, Taubert et al. 2019). For ecological functioning, the metabolic diversity of microbes is

especially relevant (Petchey and Gaston 2006, Cardinale et al. 2012, Johnson and Pomati 2020).

Metabolic diversity directly describes genetic variation of functional genes. Evolutionary

selection acts directly on such traits because they influence organism fitness, making metabolic

diversity a direct result of evolution (Fischbach et al. 2008). To gain deeper insights into the

evolutionary history and relationship of microbes, information about their phylogenetic

diversity is required.

Phylogenetic diversity is usually assessed using neutral genetic markers like the bacterial 16S

rRNA gene. Neutral markers are not under evolutionary pressure by selection. However, they

are affected by genetic drift, leading to stable mutation rates that can be used as molecular

clocks. But as phylogenetic diversity rests on a neutral marker, it should hold no information

regarding functioning. For this reason, ecologists have been debating the usefulness of

phylogenetic diversity in the study of ecosystem functioning. Consequently, studies that have

used phylogenetic diversity have yielded mixed results when trying to explain ecosystem

functioning (Cadotte et al. 2009, Flynn et al. 2011, Alexandrou et al. 2015, Gerhold et al. 2015,

Venail et al. 2015).

Despite that, phylogenetic diversity is linked to metabolic diversity because closely related

bacterial species are often more likely to have similar metabolic traits (Martiny et al. 2013,

Goberna and Verdú 2016).

The connection between metabolic traits and phylogenetic relatedness is called phylogenetic

signal (Losos 2008). Phylogenetic diversity is a good predictor for metabolic functions with a

strong phylogenetic signal, such as ammonia oxidation mediated by a restricted group of

chemolithoautotrophic bacteria (Kowalchuk and Stephen 2001), which can be traced in deep

phylogenetic branches. Yet, especially for broad, simple functions such as carbon assimilation

from amino acids and simple sugars (Martiny et al. 2013), it is a poor predictor of ecological

functioning (Srivastava et al. 2012, Martiny et al. 2013, Goberna and Verdú 2016). However,

phylogenetic distance can be used to infer niche differences and fitness differences, given that

those differences influence coexistence (Godoy et al. 2014, Weber and Strauss 2016). As such,

under certain circumstances, phylogenetic relationships can predict growth rates (Morrissey et

al. 2016), species interactions via ecological networks (Srivastava et al. 2012) and species

coexistence (Germain et al. 2016). Hence, we can link species evolutionary relatedness to

properties that are under natural selection and therefore contribute to ecological functioning

(Morrissey et al. 2016). Therefore, even though phylogenetic diversity is based on neutral

markers, it should correlate with functional differences because it influences species

interactions that lead to the evolution of different functions. The precise connection behind the

phylogenetic and metabolic diversities is still unclear, because the underlying mechanisms are

either unknown or very complex (Losos 2008). For this reason, efforts to study the effects of

phylogenetic diversity and, as an extend, the usefulness of phylogenetic signals have been

mixed and relatively limited (Venail et al. 2015).

A direct assessment of metabolic diversity hence allows a better prediction of community

ecosystem functions (Mokany et al. 2008, Salles et al. 2009, Krause et al. 2014). Metabolic

diversity, however, is more difficult to define and quantify (Mouchet et al. 2010, Cadotte et al.

2011, Steudel et al. 2016) as it requires the measurement of functional trait differences of

organisms (Petchey and Gaston 2002, 2006, Naeem et al. 2009). Microbial functional traits can

be determined based on, e.g., functional gene-targeting microarrays and metabolic activity

assays (Litchman and Klausmeier 2008, Naeem et al. 2009, Paula et al. 2014, Wang et al. 2017).

Such assays can also be used as a measure for community functioning itself along with other

ecologically important community functions, such as nutrient cycling, organic matter

decomposition and CO_2 production, which were shown to be positively correlated with

metabolic diversity (Rivett et al. 2016, Evans et al. 2017).

While phylogenetic and metabolic diversities are associated with distinct evolutionary

mechanisms and processes, they should also be intricately linked, making it difficult to

understand their individual effects on community ecological functions. Since phylogenetic

diversity influences species coexistence and interactions, such processes should interact with

metabolic diversity (Pascual-García et al. 2020) and possibly modify the effects metabolic

diversity has on community functioning. For the further development of micro-ecological

theory (Krause et al. 2014, Shade et al. 2018) and biodiversity research (Flynn et al. 2011,

Roger et al. 2016), potential interactions between phylogenetic and metabolic diversity have to

be identified to understand their influence on community ecosystem functions (Thompson et

al. 2015).

Here, we aim to disentangle this interaction by using synthetic microbial communities in a

multifactorial experimental design, where we control for each diversity aspect while studying

the other's effect on community ecological functions. To study the combined effects of

phylogenetic and metabolic diversity, we used bacterial groundwater isolates to form simplified

four-member communities with different phylogenetic and metabolic diversity treatments,

allowing us to test how these two aspects of biodiversity interact as they shape community

ecological functions. We hypothesised that 1) when controlling for phylogenetic diversity,

metabolic diversity has a positive effect on community ecological functions. In contrast, 2)

when controlling for metabolic diversity, phylogenetic diversity, based on a neutral marker,

should have no effect on community ecological functions. Finally, 3) the effect of metabolic

diversity is proportionally stronger in communities that are phylogenetically similar than in

communities that are phylogenetically dissimilar, as phylogenetically dissimilar communities

have less niche overlap and therefore interact less strongly in terms of competition.

To determine community ecological functioning, we targeted exoenzymes related to nutrient

assimilation. Production and release of exoenzymes is an important mechanism that bacteria

use to access exogenous, complex carbon and energy sources. Moreover, exoenzymes can be

produced constantly or on-demand. As such, exoenzymes are "public goods" that are shared

among different species (Drescher et al. 2014, Smith and Schuster 2019), and thus their

production is expected to be affected by microbial community interactions. Exoenzyme

activities (EEAs) can be measured in an easy, standardised way to link multiple bacterial

metabolic processes to ecosystem functions like carbon cycling (Malik et al. 2019). We used

EEA assays to provide a metabolic classification of bacterial isolates, allowing us to construct

communities with predetermined diversity treatments. Furthermore, EEAs were used to

measure functioning in the synthetic communities. Overall, the measure of EEAs allowed us to

examine how differences in the phylogenetic and metabolic diversity between species can affect

community ecological functions.

Material and Methods

Selection of bacterial isolates and culture setups

The twenty-seven bacterial strains used in this study (Table S1) were isolated from pristine groundwater (Geesink et al. 2018), obtained from the Hainich Critical Zone Exploratory located in Thuringia, Germany (Küsel et al. 2016). The isolates were selected from a pool of more than 340 groundwater isolates as representatives of the most abundant taxonomic families within the total groundwater bacterial population, based on 16S rRNA gene MiSeq amplicon sequencing data (Schwab et al. 2017, Kumar et al. 2017), throughout the taxonomic classes of *Actinobacteria*, *Bacilli*, *Flavobacteriia*, *Sphingobacteria*, *Alpha-* and *Gammaproteobacteria*. The isolates were maintained on S2P medium agar (Geesink et al. 2018). Before each of the experimental assays, isolates were cultured aerobically in 20 ml of liquid S2P medium, in cap-perforated 50 ml Falcon tubes, horizontally shaken (125 rpm) at room temperature, for 48 h to early stationary phase. Growth in the liquid cultures was being determined via optical density measurements at a wavelength of 600 nm (Synergy™ H4 Hybrid Multi-Mode Microplate Reader, BioTek, Germany).

Molecular analysis and taxonomic classification of bacterial isolates

DNA from bacterial isolates was extracted with GenElute™ Bacterial Genomic DNA Kit (NA2110-1KT, Sigma-Aldrich, Germany). Sterile inoculation loops were used to gather 1 µl-worth of biomass from the agar medium cultures, for DNA extraction. For the 16S rRNA gene amplification, the primers 8F (5′-AGA GTT TGA TCC TGG CTC AG-3′) (Turner et al. 1999) and 1492R (5'-GGT TAC CTT GTT ACG ACT T-3') (Turner et al. 1999) were used. PCR was performed with a PEQSTAR thermocycler (PEQLAB, Germany) using the following cycling conditions: initial denaturation at 95 ºC for 10 min, 30 cycles of 95 ºC (denaturation), 55 ºC

(primer annealing) and 72 °C (elongation) for 30 sec, 45 sec and 60 sec, respectively, followed

by 10 min at 72 °C (final elongation). DNA purification and sequencing of the 1484bp-length

PCR products was carried out by Macrogen (Amsterdam, Netherlands) with the 27F (5'-AGA

GTT TGA TCM TGG CTC AG-3') primer.

The sequences were assigned to their closest reference relative using BLASTn (Zhang et al.

2000) (v2.8.0+) and cross-referenced with SILVA Incremental Aligner (SINA) (Pruesse et al.

2012) (v1.2.11). A rooted maximum-likelihood phylogenetic tree for the 16S rRNA gene

taxonomic classification of the organisms was constructed with MEGA X (Kumar et al. 2018)

(v10.1.5).

Exoenzyme activity assays (EEAs)

EEAs for each bacterial isolate were performed using eleven substrates (Table S2) (Sigma-

Aldrich, Germany), each carrying a fluorescent moiety allowing spectrophotometric

measurements of activity using a microplate reader (Synergy™ H4 Hybrid Multi-Mode

Microplate Reader). The sensor was set to detect fluorescence at wavelengths 360/9 nm for

excitation and 450/9 nm for emission.

Prior to the EEA assays, the optical density of the bacterial cultures was determined

spectrophotometrically (600 nm). For the EEA assays, a modified version of the protocols from

Frossard *et al.* (Frossard et al. 2012) and Rivett *et al.* (Rivett et al. 2016) was used here, as

follows: a black 96-well plate setup, with 100 µl HEPES buffer at 100 mM stock concentration

in each well, was inoculated with 100 µl of bacterial culture and to each test well 50 µl of

substrate solution at 40 µM final concentration was added. Negative controls consisted of wells

containing 100 µl of sterile S2P liquid medium and wells with just 100 µl sterile MilliQ water

(Merck Millipore, Germany). Fluorescence standards for the MUB substrates were measured

using 4-Methylumbelliferone with final concentrations of 10 to 0.3125 µM by halving dilutions.

Fluorescence standards for the AMC substrates were measured using 7-Amino-4-

methylcoumarin with final concentrations of 2 to 0.0625 μM by halving dilutions. The

standards were used to transform relative fluorescence units (RFU) to concentration of

degraded substrate. Background fluorescence, measured in negative controls with either

substrate solution or MilliQ water, was subtracted from the respective assay plates.

Measurement of the assay plates begun immediately after the substrate addition and continued

for 90 min, at 2.5 min measurement intervals.

Determining phylogenetic and metabolic diversity of isolates

To examine effects of diversity on the functioning of synthetic bacterial communities, the 16S

rRNA gene taxonomy and EEAs of the twenty-seven bacterial isolates were used as the basis

of the community traits: phylogenetic and metabolic diversity. We formed four trait-groups,

each consisting of five isolates, being either phylogenetically and metabolically similar (Psim-

Msim), phylogenetically similar but metabolically dissimilar (Psim-Mdis), phylogenetically

dissimilar and metabolically similar (Pdis-Msim) as well as phylogenetically and metabolically

similar (Pdis-Mdis). In total, the isolates covered ten taxonomic families of five classes (Figure

S1). In the Psim groups, isolates were affiliated with the same genus; Psim-Msim group held

Flavobacteria isolates and Psim-Mdis held *Pseudomonas* isolates. Pdis groups were formed so

that all members of a group were related to different families. Subsequently, Psim and Pdis

group phylogenetic diversity quantitative measures were obtained using the "*Compare within*

group mean distance" function of MEGA X (Kumar et al. 2018) (v10.1.5) and verified to be

statistically insignificant for Psim communities and significantly different for Pdis communities

using paired t-tests (Table S3). Hierarchical clustering based on Euclidean distance was used

to quantise functioning of the total EEAs for each isolate (Figure 1) and categorise them as

metabolically similar or dissimilar. The clustering analysis was performed in R (v. 3.6.1) (R

Core Team 2019) using the core stats functions *dist(x, method = "euclidean", p = 2)* to

calculate the Euclidean dissimilarity distance matrix (Borg and Groenen 2005), followed by

hclust(d, method = "ward.D") to hierarchically rank and cluster the EEA dissimilarities

between the bacterial isolates using the Ward hierarchical clustering method (Ward 1963,

Murtagh and Legendre 2014). Msim groups were formed from isolates of the same cluster,

Mdis groups were formed so that each member of the group belonged to a different metabolic

cluster (Figure 1). Quantitative measures of principal component analysis (PCA) based on the

metabolic traits of the isolates confirmed the higher metabolic diversity between isolates of

Psim-Mdis and Pdis-Mdis compared to group Psim-Msim and Pdis-Msim (Figure S2).

Bacterial community setups and experimental design

We designed synthetic communities of bacterial species according to their metabolic and

phylogenetic diversity, based on the four trait groups: Psim-Msim, Psim-Mdis, Pdis-Msim,

Pdis-Mdis. From the initial twenty-seven isolates tested, a total of seventeen isolates were used

for designing communities. Within each trait group, five distinct communities were formed by

omitting one isolate out of the five, per community, allowing us to test whether within-group

community EEA variations were due to the activity of a specific community member. This

resulted in twenty communities, five for each trait group, identified as C1 to C5. We confirmed

the phylogenetic diversity between the communities via the quantitative estimates of isolate

phylogenetic distance and their metabolic diversity based on the Euclidean distance matrix of

isolate EEAs as previously mentioned (Table S3). The phylogenetic and metabolic diversity

values for each community showed a distinct differentiation of the communities from each of

the four groups, confirming that we successfully separated the bacterial isolates and

communities into the four intended groups (Figure 2).

We grew cultures of each isolate in liquid S2P medium as described above. Optical density

(600 nm) was used to calculate the required final volume for each isolate, so that all members

of each community were to be introduced into the communities at the same amount of biomass.

The communities were established in cultures by mixing a predetermined amount of liquid

isolate culture for a total volume of 2.5 ml. The community cultures were incubated in deep 96-

well plates (2.5 ml volume), at room temperature and sealed with rayon film seals (391-1262,

VWR, Germany) to allow gas exchange. The EEAs of the microcosms were assessed after 0 h,

6 h and 24 h of incubation. Additionally, at the 0 h and 24 h marks, a 200 µl culture sample was

pelleted by centrifugation (17.000 $\times$ g, 5 min) and stored at -80 °C, for subsequent molecular

analysis.

Ecological functioning data analysis

The communities constructed in this study were composed out of different isolates with widely

differing metabolic activities. A comparison of EEAs between communities alone would thus

only reflect the differences between the constituent isolates. Therefore, EEA differences of the

communities were normalized to the respective isolates and expressed as proportional changes

of EEA. This was done with the following steps (visualised in Figure S3): The measured EEAs

of the isolates within each community group were used to calculate expected EEAs for each of

the group's communities. The expected EEAs were compared to the measured EEAs of the

experimental communities and the result was represented as % proportional changes of EEAs.

This resulted in values from -100% to 385%. In addition, the total available carbon (TAC) from

each substrate were determined. This was done by multiplying the total number of carbon atoms

in each substrate (excluding the fluorescent moiety) with the molar amount of degraded

substrate. TAC was summed as a unified measure of ecological functioning for each isolate and

community. 4-MUB phosphate and 4-MUB sulphate were excluded because they contained no

carbon atoms. TAC was used to directly compare communities as an absolute measure of

functioning.

To evaluate the significance of phylogenetic diversity, metabolic diversity was controlled in the

community EEA proportional change data by subsetting into metabolically similar and

dissimilar communities. Likewise, to examine the significance of metabolic diversity,

phylogenetic diversity was controlled for in the same way. These analyses were performed for

the 0 h timepoint data, using ANOVA of linear regression models in R. For the linear models,

the EEA % proportional change data as well as the TAC data were log2-transformed. To enable

logarithmic scaling of the EEA % proportional change data, 101% was added to all datapoints,

to avoid negative values and zeroes, since the smallest values were -100%; For the TAC

datapoints, this was not necessary. Furthermore, phylogenetic and metabolic diversity metrics

were z-score normalised, for the linear regression models, to make them comparable. The group

Pdis-Mdis showed no activity in the 4-MUB sulphate assay, and so this assay was excluded

from the models, resulting in a total of 215 datapoints of EEA % proportional changes. For the

TAC data, 20 datapoints were used, one per community.

For the analyses for the influences of substrate biochemistry on the diversity treatment effects,

the EEA proportional change data were not subset by diversity treatment but by each substrate.

Following that, ANOVA of linear regression models was performed as described above.

Effects of diversity in EEA changes over 24 h

Time is a crucial driver of bacterial interactions and therefore we evaluated how the EEA

proportional changes of communities developed over 24 h, by comparing the 0 h timepoint to

the 6 h and 24 h timepoint data. This was done for all 20 communities of all 4 diversity

treatment groups, and for each substrate, the average of each group was determined. The

significance of temporal trends was tested using the Student's t-test (2-tailed) by comparing the

EEA proportional change of all communities in a treatment group at 0 h to all EEA proportional

changes of the same group for timepoints 6 h and 24 h, for each individual substrate. False

discovery rate was controlled with the Benjamini–Hochberg procedure (McDonald 2014).

Substrate acclimation in EEA assays

To study influences of acclimation of isolates and communities to the added substrates on the

observed effects of diversity treatments on community ecological functions, we added a pre-

incubation period to the EEA assays. This was performed only with the Psim-Mdis group. After

mixing of the communities, substrates were added after 0 h and 6 h to isolates and communities,

and EEA assays were performed as described above (no-acclimation treatments). In addition,

isolates and communities with substrates added 0 h after mixing were incubated for a total of

4.5 h, before another series of EEA assays were conducted (acclimation treatments).

Assay duration was 90 min, as in previous assays. To achieve a direct comparability of the

activities measured with and without acclimation, the determined EEAs were normalized to

optical density of isolates and communities, which was measured before each assay. Following

that, the change in EEAs from 0 h to 4.5 h and 6 h was compared between isolates and

communities.

Molecular and diversity analysis of synthetic communities

Community DNA was extracted as described above. DNA concentration was measured

spectrophotometrically (NanoDrop 1000, PEQLAB, Germany) and samples were stored

at -20 °C.

For Illumina MiSeq amplicon sequencing of 16S rRNA genes, PCR was performed using the

primer set Bakt_341F (5'-CCT ACG GGN GGC WGC AG-3') (Herlemann et al. 2011) and

Bakt_805R (5'-GAC TAC HVG GGT ATC TAA TCC-3') (Herlemann et al. 2011) targeting

the V3-V4 hypervariable region of the rRNA gene. DNA amplification was carried out with a

PEQSTAR 2X thermocycler (PEQLAB, Germany) in the following conditions: initial

denaturation at 95 °C for 15 min, 30 cycles of 94 °C (denaturation), 55 °C (primer annealing)

and 72 °C (elongation) at 45 sec for each step, followed by 10 min at 72 °C (final elongation).

The PCR DNA products were purified using NucleoSpin Gel and PCR Clean-up (Ref. 740609,

Macherey-Nagel, Germany) following the manufacturer's protocol. The final step, DNA

recovery from the columns, was performed twice for a total volume of 60 μl purified PCR

product samples.

MiSeq-generated sequence data were analysed using mothur (Schloss et al. 2009) (v.1.39.1;

mothur.org) along with the mothur MiSeq SOP developed by Kozich *et al*. (Kozich et al. 2013)

(accessed November 19th 2018). Paired-end reads were combined and sequences below 400 bp

or above 429 bp as well as sequences with ambiguous bases or homopolymer runs above 8 bp

were removed. Sequences were aligned to the SILVA reference database release SSU 132

(Quast et al. 2012) and clustered when differed in four or fewer bases. Chimeric sequences were

removed with uchime (Edgar 2016) using the GOLD database (Mukherjee et al. 2019) as a

reference. OTUs were binned from sequences with a 3% identity cut-off and were classified

using the same SILVA database. Raw data have been deposited in the European Nucleotide

Archive under the study with accession number PRJEB34832 (samples ERR3588683 -

ERR3588722).

Results

Effects of diversity on community ecological functions

EEAs changes of the synthetic four-species communities exhibited strong differences between substrates (Figure 3; Figure S4). In addition, EEA changes showed sharp decreases, down to 100%, and increases, up to 385%, at 0 h after mixing. Substrates with glycosidic bonds like 4-MUB-N-acetyl-β-D-glucosaminide (NAG), 4-MUB-β-D-cellobioside (CB) and 4-MUB-β-D-xylopyranoside (XYL) exhibited the highest relative increases in EEA (see Table S2 for complete list of substrate abbreviations). The EEA changes for each one substrate within a treatment group, however, were typically consistent ($< 50\%$ EEA proportional change from the average) for 33 of the 44 (75%) community-substrate combinations. This indicated that, in most cases, community behaviour was not strongly influenced by the excluded fifth isolate of each group. The consistency was more prominent for substrates with ester bonds like 4-MUB acetate (ACE), 4-MUB-butyrate (BUT) and 4-MUB oleate (OLE). Tables S4, S5, S6 and S7 show a complete overview of cleaved substrate concentrations for isolates, communities and estimated community EEAs for each treatment group. When comparing EEA changes between the groups (Figure S4B-E), we observed that Pdis communities had relatively small to no changes in EEA: 76% of community-substrate combinations of group Pdis-Msim and Pdis-Mdis remained between +50% and -50% EEA proportional change. Contrarily, communities of Psim groups differed greatly in activities compared to their constituent isolates. Group Psim-Msim exhibited most of the largest EEA increases (up to 330%) with 27% of the group's 55 community-substrate combinations measuring $> +50\%$ EEA proportional change. Group Psim-Mdis communities recorded the greatest amount of negative EEA proportional changes (53% of the group's 55 community-substrate combinations having $< -50\%$ EEA proportional change) with the maximum EEA proportional change increase at only 5.4%. These results show that

phylogenetically similar communities exhibited more variable and genus-specific responses

than phylogenetically dissimilar communities.

Using linear regression analyses, we determined the effects of phylogenetic diversity on

community EEA changes while controlling for the metabolic diversity treatments (Figure 4A;

Table S8). For metabolically similar communities, phylogenetic diversity had a significant

negative effect on the EEAs (-0.279 $\pm$ 0.094, estimate $\pm$ std. error; Adj. R^2 = 0.068; p = 0.004).

In contrast, for metabolically dissimilar communities, a significant positive effect of

phylogenetic diversity was observed (1.094 $\pm$ 0.198, estimate $\pm$ std. error, Adj. R^2 = 0.222,

p = 2.35×10^{-7}). The effect of metabolic diversity was similarly shown while controlling for

the phylogenetic diversity treatments. For both phylogenetically similar and dissimilar

communities, metabolic diversity had a negative effect on EEAs (Figure 4B). This effect,

however, was much stronger in the case of phylogenetically similar communities,

(-1.535 $\pm$ 0.233, estimate $\pm$ std. error,   Adj. R^2 = 0.280,   p = 1.64×10^{-9}),   than   for

phylogenetically dissimilar communities, where it was below the significance threshold

(-0.123 $\pm$ 0.090, estimate $\pm$ std. error, Adj. R^2 = 0.008, p = 0.177).

Furthermore, we examined the absolute level of functioning of communities, using TAC turned

over by EEA in each community as a unified measure (Figure S5; Table S9). Linear regression

analyses revealed similar trends than observed for the proportional changes of community

EEAs: When controlling for metabolic diversity, phylogenetic diversity had a significant

negative   effect   on   TAC   in   metabolically   similar   communities

(-0.305 $\pm$ 0.102, estimate $\pm$ std. error,   Adj. R^2 = 0.467,   p = 0.018)   (Figure   S5A).   For

metabolically   dissimilar   communities,   phylogenetic   diversity   had   positive   effect

(0.313 $\pm$ 0.105, estimate $\pm$ std. error,   Adj. R^2 = 0.467,   p = 0.018).   While   controlling   for

phylogenetic diversity, metabolic diversity had a negative effect in phylogenetically similar

communities (-0.611 $\pm$ 0.151, estimate $\pm$ std. error, Adj. R^2 = 0.632, p = 0.004) (Figure S5B),

but   no   effect   was   observed   for   phylogenetically   dissimilar   communities

(0.002 ± 0.115, estimate ± std. error, Adj. R^2 = -0.125, p = 0.986). Hence, biodiversity effects

on the community EEAs were also reflected in the carbon acquisition potential of these

communities.

Chemical structure differences of substrates influence the strength of diversity effects

Investigating the datasets for all substrates individually using linear models, we observed broad

differences in the relationships of EEAs to the metabolic and phylogenetic diversity between

them (Table S10). When controlling for substrates, the diversity treatments have a consistent

pattern of influence, positive for phylogenetic diversity and negative for metabolic diversity

(Figure 7). For example, CB showed a significant positive relationship with phylogenetic

diversity (1.029 ± 0.444, estimate ± std. error; Adj. R^2 = 0.520, p = 0.034) and for metabolic

diversity, a significantly negative relationship (-1.604 ± 0.441, estimate ± std. error;

Adj. R^2 = 0.520, p = 0.002). The other two polysaccharide-analogue substrates, XYL and

NAG, behaved similarly to CB (Table S10). We observed that the strength of the effect was

related with the chemical structure of the substrates (Figure 7A). Esters (ACE, BUT, OLE)

exhibited no relationship with diversity treatments, with linear slopes ranging from -0.095 to

0.038 for phylogenetic diversity and -0.227 to -0.007 for metabolic diversity. Monosaccharide

(aG, bG) linear slopes ranged from 0.211 to 0.368 and -0.760 to -0.336, a relationship that was

overall similar but weaker than for the polysaccharides (CB, XYL, NAG), whose linear slopes

were 1.029 to 1.470 and -2.019 to -1.604 for phylogenetic and metabolic diversity, respectively.

On average, by controlling for substrates alone, we observed that phylogenetic diversity had a

positive effect on EEA changes (0.643 ± 0.126, estimate ± std. error; Adj. R^2 = 0.286,

p = 7.44×10^{-7}) (Figure 7B). Accordingly, metabolic diversity had a negative effect (-0.898 ±

0.125, estimate ± std. error; Adj. R^2 = 0.286, p = 1.33×10^{-11}) (Figure 7C).

Effects of diversity in EEA changes over 24 h

We further investigated how the community EEA changed with increasing time after community mixing, by comparing EEA results from time points of 0 h, 6 h and 24 h (Figure 5). Both Msim groups exhibited only negative trends over the 24 h. The Psim-Mdis group showed both significantly negative and positive trends between EEAs measured over 24 h after mixing. Changes in the Pdis communities' composition within 24 h did not explain the observed activity changes, because all the introduced species appear to persist in the community cultures during 24 h post-mixing, with only close to ~1-fold change of relative abundance for most species and a 2-fold change for two species (Figure S6). Thus, the observed changes in EEA over time should not be the result of a shifting community composition supporting our hypothesis, that EEA changes are due to diversity treatment effects.

Substrate acclimation in EEA assays

Acclimation describes any phenotypic plasticity an organism can undergo to adjust to changes in its environment. Effects of substrate acclimation were observed directly by measuring the increase of EEA due to potential upregulation of exoenzyme production, during the incubation with a substrate. Here we show the results from the assay with aG, as an example. Comparing the timepoint of 0 h to the acclimated cultures timepoint of 4.5 h, we observed increases of the EEAs for both. However, these increases were much higher for communities, with ~5-fold increases on average, than for isolates, with average increases of ~1-fold (Figure 6). This was also reflected in the highest changes observed: For isolate activities in the aG substrate assay, 2.3 times increase, from 0.110 ± 0.003 to 0.255 ± 0.008 µM min^{-1}, was the largest change observed. For community activities, increases up to 10 times, from 0.020 ± 0.001 to 0.199 ± 0.012 µM min^{-1} were observed, making community acclimation capacity almost 5 times higher compared to isolates. This increase in activity did not match the EEA changes

observed for communities after 6 h of mixing without the addition of substrates. Since both

isolate and community activities were normalised to biomass, the increased activity in

communities could not be attributed to differences in cell density. In addition, the acclimation

effect could not have been the result of species abundance changes since even the most active

of the isolates only had ~2 times increase of EEA due to acclimation (0.225 ± 0.006 to

$0.454 \pm 0.003 \ \mu M \ min^{-1}$).

Discussion

Using synthetic microbial communities, we show that both phylogenetic and metabolic diversity influence community ecological functioning. Because both diversity aspects are so intricately linked, we used a multifactorial treatment experiment that allowed us to examine what the effects of each diversity are when controlling for the other. Since metabolic diversity has a direct impact on ecological functions, we expected that higher metabolic diversity would lead to higher functioning of bacterial communities. What we see is that the interaction between diversity treatments causes contrasting effects which did not meet our expectations. Metabolic diversity has either no effect, for Pdis communities, or a negative relationship, for Psim communities for both community EEA proportional changes and community TAC (Figure 4B, Figure S5B). In Pdis communities, differences in nutrient acquisition and interaction strategies of different species could be playing a role for the lack of effect of metabolic diversity. Such a mixture of species can engage in varying competitive or cooperative interactions (Gross 2008, Violle et al. 2011). This complementary set of community interactions might be masking any observable influence of metabolic diversity (Maynard et al. 2017). On the other hand, in Psim communities, related species are likely to be reacting similarly to each other and resource competition. However, increased conspecific competition can result in different outcomes. With lower metabolic diversity, upregulation of exoenzyme activity as a response to competition for resource exploitation, might be translating to increased community functioning. The mechanism behind this might be how the cost:benefit ratio of exoenzyme production manifests in low diversity communities if they are all producers and strong competitors (Allison et al. 2011). Consequently, higher metabolic diversity results in lower community functioning as species can diversify their resource preferences and therefore do not contribute to the same substrate degradation in the assays. Moreover, species of different genera

might have distinct ways of interacting with species of their own genus, further influencing

competitive outcomes (Mayfield and Levine 2010, Maynard et al. 2017).

We also predicted that as phylogenetic diversity is based on a neutral genetic marker, it would

have no effect on community ecological functioning. Instead, we find that for both EEA

changes and TAC, phylogenetic diversity has a positive relationship with the ecological

functioning of Mdis communities and a negative relationship for Msim communities (Figure

4A; Figure S5A). In the Mdis communities, organisms can engage in a variety of metabolic

functions. This increases the capacity for different kinds of community interactions (Weber and

Strauss 2016), facilitated by phylogenetic diversity and coexistence. More diverse interactions

might translate into more cooperative communities, metabolic complementarity and increased

overall functioning. Contrastingly, Msim communities rely on a limited number of metabolic

functions. The limited metabolic functions do not allow for elaborate interaction strategies, such

as cooperation, to be enacted. As such, any influence of phylogenetic diversity is severely

decreased. In this setup, resource competition and antagonistic interactions with multiple

species cause a negative relationship to emerge.

Lastly, we hypothesised that because Psim communities would have less diverse strategies in

dealing with competition or cooperation, metabolic diversity would have a much stronger effect

in their ecological functioning, compared to their Pdis counterparts. Indeed, this is true as we

show that Psim communities had a strong, albeit negative relationship with EEA changes and

TAC, while Pdis communities had none. Moreover, the consistency of results between the

single and multifunctional measures of EEAs and TAC increases the confidence of the diversity

relationships we have observed. This should not be surprising, as we already established a deep

connection between phylogenetic and metabolic diversity, which would explain why neither of

them is the sole driver of community ecological functions.

The contrasting diversity effects were also revealed when controlling for the substrates in the

treatments. We identified as a general trend that phylogenetic diversity has a positive effect

(Figure 7B) on EEA changes and metabolic diversity has a negative influence (Figure 7C). In

addition to that, we saw that the diversity effects were operating on a strength gradient, with

groups of biochemically similar substrates appearing in distinct positions along that gradient

(Figure 7A). Such differences between substrates and groups of substrates can be related to

inherent variations in exoenzyme production, maintenance and regulation and how different

classes of enzymes behave to break down their substrates. For example, polysaccharides are

targeted by hydrolytic exoenzymes that are produced on-demand (Warren 1996, Lynd et al.

2002). In contrast, enzymes involved in the degradation of fatty acids belonging to the ester

group of substrates are part of bacterial core metabolism, and thus can be continuously

produced. Our results show that stronger diversity effects are more important to ecosystem

functions that rely on environmental factors, such as the temporal or spatial presence of plant-

derived polysaccharides. Further complicating experimental outcomes, time also influences

different substrates and diversity treatments in different ways (Figure 5). Hence, not only the

diversity of ecosystem functions tested (Huston 1997, 2000, Tilman 1997, Loreau 2000) but

also their underlying biochemical mechanisms have a great effect on the outcomes of

biodiversity research and the choice, effect and timing of different functions in experimental

designs needs to considered.

To understand the very strong contrasting changes in EEAs between the Psim groups, their

genus-specific capacity for co-existence with its related species is crucial. Psim-Msim

communities contained *Flavobacteria* species which are usually non-motile organisms, prefer

aquatic environments and have strong differences between species. The genus includes

generalist species that can use a wide range of carbohydrates, to more specialist asaccharolytic

species that prefer nitrogenous substrates (i.e. amino acids) (Bernardet and Bowman 2006).

These physiological characteristics might allow the *Flavobacteria* communities to increase

their EEA as different species could be utilising different carbon sources in their growth

medium. Communities of these specialist *Flavobacteria* could be facilitating community

interactions that translate into increases in their community EEAs. In contrast, the

Pseudomonas species in Psim-Mdis communities are generally motile-planktonic bacteria,

ubiquitous in almost all oxic environments, including soil and water, due to their expansive

metabolic repertoire, simple nutritional requirements and rapid growth rates (Moore et al.

2006). However, the reduced capacity of many *Pseudomonas* species to withstand starvation

(Moore et al. 2006) might lead them to adapt more selfish nutrient acquisition strategies

(Reintjes et al. 2017, 2019, Arnosti et al. 2018). As generalist species, *Pseudomonas* might be

more competitive, resulting in *Pseudomonas* communities with lower EEAs than their

respective isolates. Thus, bacteria with a broad substrate range might not benefit when living

in communities of the same genus but of dissimilar metabolism. Contrastingly, in the Pdis

communities these genus-specific effects in nutrient acquisition strategies can be balanced due

to the increased species diversity.

Nevertheless, even competing generalists like *Pseudomonas* species will profit from species

interactions when incubated for longer time in the presence of certain substrates. Using the

Psim-Mdis communities to run EEA substrate acclimated vs. non-acclimated assays (Figure 6),

we identified that the amplifying effect of substrate acclimation on the EEAs is much stronger

for the communities than for the constituent isolates on substrate aG. In addition, since the

increased EEAs are not present in the non-acclimated 6 h timepoint communities, we can

conclude that the effect was indeed substrate acclimatisation of the bacteria and not an effect

of mixing the community cultures. Consequently, the observed acclimation effect was the result

of community interactions that enhanced species' ability to acclimate to the substrate. These

observations link back to our first hypothesis, where we predict that the metabolic diversity of

the group will enhance the functioning of communities. But oversimplifying the experimental

design and just looking at one substrate or one group, would have let to an underestimation of

the complexity of diversity-ecosystem functioning relationships, possibly explaining as to why

we see studies showing opposite effects of biodiversity (Balvanera et al. 2006).

Altogether, our results concur with previous studies that focus on separating the different

aspects of biodiversity and examining the relationship between biodiversity and ecosystem

functioning (Griffiths et al. 2001, Zhang and Zhang 2006, Balvanera et al. 2006, Naeem et al.

2009, Jousset et al. 2011, Purschke et al. 2013, Thompson et al. 2015, Gamfeldt and Roger

2017). A meta-analysis demonstrated that biodiversity effects depend on the organisation level:

with positive effects at the community level, positive but weaker effects at the ecosystem level

and negative effects at the population level (Balvanera et al. 2006). A modelling approach

showed that when considering increasing number of functions, the effect of biodiversity is lost

(Gamfeldt and Roger 2017). Experiments done by Griffiths *et al*. (Griffiths et al. 2001)

recognised no consistent effect from biodiversity on soil functions. Thus, the need for more

research including the effects of the environmental context, like nutrient availability, on the

relationship between biodiversity and ecosystem functioning was recognized (Zhang and Zhang

2006). Our study fills this gap by focussing not only on the phylogenetic and metabolic diversity

of organisms, but also measuring a broad diversity of activities controlled by different

underlying mechanisms of enzyme regulation. Such a complex approach is required to tackle

the complex dimensionality of biodiversity research. We show that biodiversity had a

significant impact on most EEAs, but the strength of the effect depended on the substrate

context. As such, biodiversity's effects on community ecosystem functioning cannot always be

predicted as a positive relationship between species diversity and enhanced ecosystem function

(Aarssen 1997, Tilman 1997, Huston 2000, Carroll et al. 2011). Natural, larger scale

experiments struggle with introducing different levels of complexity in their experimental

design (Cairns et al. 2018). Looking forward, we should aim to understand how diversity effects

can mechanistically influence ecosystems, synthetic and natural (Fox and Harpole 2008, Naeem

et al. 2009, Turnbull et al. 2013), and not take the effects as granted (Shade 2017).

Acknowledgements

We gratefully acknowledge the support of the German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) (DFG FZT 118). This project has been conducted in the framework of the iDiv-Flexpool – the internal funding mechanism of iDiv. We would like to express our gratitude to Patricia Geesink for the bacterial isolation work under the framework of the CRC AquaDiva of the Friedrich Schiller University Jena, funded by the DFG (DFG SFB 1076). Additionally, we would like to thank Will A. Overholt and Lijuan Yan for their invaluable statistical analysis input. Lastly, we want to thank the two reviewers for the very positive and constructive feedback, which has helped to greatly improve this manuscript.

Author Contributions

KK and SWH secured funding for the project. CX, MT, and KK designed the experiments. CX performed the experiments and carried out all assays and molecular analyses; the data analysis was performed by CX closely supported by MT. Statistical analysis was performed by CX supported by MT and SWH. All co-authors assisted in interpreting the results; CX initiated the manuscript writing, which was finalized with substantial contributions from MT, KK and SWH.

Competing Interests

The authors declare no conflicts of interest.

References

Aarssen, L. W. 1997. High productivity in grassland ecosystems: effected by species diversity

or productive species? test. Oikos 80:183–184.

Alexandrou, M. A., B. J. Cardinale, J. D. Hall, C. F. Delwiche, K. Fritschie, A. Narwani, P. A.

Venail, B. Bentlage, M. S. Pankey, and T. H. Oakley. 2015. Evolutionary relatedness

does not predict competition and co-occurrence in natural or experimental communities

of green algae. Proceedings. Biological sciences / The Royal Society 282:20141745.

Allison, S. D., M. N. Weintraub, T. B. Gartner, and M. P. Waldrop. 2011. Evolutionary-

Economic Principles as Regulators of Soil Enzyme Production and Ecosystem

Function. Pages 229–243 *in* G. Shukla and A. Varma, editors. Soil Enzymology.

Springer Berlin Heidelberg, Berlin, Heidelberg.

Arnosti, C., G. Reintjes, and R. Amann. 2018. A mechanistic microbial underpinning for the

size-reactivity continuum of dissolved organic carbon degradation. Marine Chemistry

206:93–99.

Balvanera, P., A. B. Pfisterer, N. Buchmann, J.-S. He, T. Nakashizuka, D. Raffaelli, and B.

Schmid. 2006. Quantifying the evidence for biodiversity effects on ecosystem

functioning and services. Ecology Letters 9:1146–1156.

Bernardet, J.-F., and J. P. Bowman. 2006. The Genus Flavobacterium. Pages 481–531 The

Prokaryotes. Springer New York, New York, NY.

Borg, I., and P. J. F. Groenen. 2005. Modern Multidimensional Scaling: Theory and

Applications. Springer, New York, NY.

Byrnes, J. E. K., L. Gamfeldt, F. Isbell, J. S. Lefcheck, J. N. Griffin, A. Hector, B. J. Cardinale,

D. U. Hooper, L. E. Dee, and J. Emmett Duffy. 2014. Investigating the relationship

between biodiversity and ecosystem multifunctionality: Challenges and solutions.

Methods in Ecology and Evolution 5:111–124.

Cadotte, M. W., K. Carscadden, and N. Mirotchnick. 2011. Beyond species: functional diversity

and the maintenance of ecological processes and services. Journal of Applied Ecology

48:1079–1087.

Cadotte, M. W., J. Cavender-Bares, D. Tilman, and T. H. Oakley. 2009. Using phylogenetic,

functional and trait diversity to understand patterns of plant community productivity.

PloS one 4:e5695.

Cairns, J., R. Jokela, J. Hultman, M. Tamminen, M. Virta, and T. Hiltunen. 2018. Construction

and Characterization of Synthetic Bacterial Community for Experimental Ecology and

Evolution. Frontiers in Genetics 9.

Cardinale, B. J., J. E. Duffy, A. Gonzalez, D. U. Hooper, C. Perrings, P. Venail, A. Narwani,

G. M. MacE, D. Tilman, D. A. Wardle, A. P. Kinzig, G. C. Daily, M. Loreau, J. B.

Grace, A. Larigauderie, D. S. Srivastava, and S. Naeem. 2012. Biodiversity loss and its

impact on humanity. Nature 486:59–67.

Carroll, I. T., B. J. Cardinale, and R. M. Nisbet. 2011. Niche and fitness differences relate the

maintenance of diversity to ecosystem function. Ecology 92:1157–1165.

Drescher, K., C. D. Nadell, H. A. Stone, N. S. Wingreen, and B. L. Bassler. 2014. Solutions to

the public goods dilemma in bacterial biofilms. Current biology: CB 24:50–55.

Edgar, R. 2016. UCHIME2: improved chimera prediction for amplicon sequencing.

bioRxiv:074252.

Evans, R., A. M. Alessi, S. Bird, S. J. McQueen-Mason, N. C. Bruce, and M. A. Brockhurst.

2017. Defining the functional traits that drive bacterial decomposer community

productivity. The ISME Journal 11:1680–1687.

Finlay, B. J., S. C. Maberly, and J. I. Cooper. 1997. Microbial Diversity and Ecosystem

Function. Oikos 80:209.

Fischbach, M. A., C. T. Walsh, and J. Clardy. 2008. The evolution of gene collectives: How

natural selection drives chemical innovation. Proceedings of the National Academy of

Sciences of the United States of America 105:4601–4608.

Flynn, D. F. B., N. Mirotchnick, M. Jain, M. I. Palmer, and S. Naeem. 2011. Functional and

phylogenetic diversity as predictors of biodiversity–ecosystem-function relationships.

Ecology 92:1573–1581.

Fox, J. W., and W. S. Harpole. 2008. Revealing How Species Loss Affects Ecosystem Function:

The Trait-Based Price Equation Partition. Ecology 89:269–279.

Frossard, A., L. Gerull, M. Mutz, and M. O. Gessner. 2012. Disconnect of microbial structure

and function: enzyme activities and bacterial communities in nascent stream corridors.

The ISME journal 6:680–691.

Gamfeldt, L., and F. Roger. 2017. Revisiting the biodiversity–ecosystem multifunctionality

relationship. Nature Ecology & Evolution 1:0168.

Geesink, P., O. Tyc, K. Küsel, M. Taubert, C. van de Velde, S. Kumar, and P. Garbeva. 2018.

Growth promotion and inhibition induced by interactions of groundwater bacteria.

FEMS Microbiology Ecology 94:1–8.

Gerhold, P., J. F. Cahill, M. Winter, I. V. Bartish, and A. Prinzing. 2015. Phylogenetic patterns

are not proxies of community assembly mechanisms (they are far better). Functional

ecology 29:600–614.

Germain, R. M., J. T. Weir, and B. Gilbert. 2016. Species coexistence: macroevolutionary

relationships and the contingency of historical interactions. Proceedings. Biological

sciences / The Royal Society 283:20160047.

Goberna, M., and M. Verdú. 2016. Predicting microbial traits with phylogenies. The ISME

Journal 10:959–967.

Godoy, O., N. J. B. Kraft, and J. M. Levine. 2014. Phylogenetic relatedness and the

determinants of competitive outcomes. Ecology letters 17:836–844.

Griffiths, B. S., K. Ritz, R. Wheatley, H. L. Kuan, B. Boag, S. Christensen, F. Ekelund, S. J.

Sørensen, S. Muller, and J. Bloem. 2001. An examination of the biodiversity–ecosystem

function relationship in arable soil microbial communities. Soil Biology and

Biochemistry 33:1713–1722.

Gross, K. 2008. Positive interactions among competitors can produce species-rich

communities. Ecology letters 11:929–936.

Gross, K., B. J. Cardinale, J. W. Fox, A. Gonzalez, M. Loreau, H. Wayne Polley, P. B. Reich,

and J. van Ruijven. 2014. Species Richness and the Temporal Stability of Biomass

Production: A New Analysis of Recent Biodiversity Experiments. The American

Naturalist 183:1–12.

Hector, A., and R. Bagchi. 2007. Biodiversity and ecosystem multifunctionality. Nature

448:188–190.

Herlemann, D. P. R., M. Labrenz, K. Jürgens, S. Bertilsson, J. J. Waniek, and A. F. Andersson.

2011. Transitions in bacterial communities along the 2000 km salinity gradient of the

Baltic Sea. ISME Journal 5:1571–1579.

Huston, M. A. 1997. Hidden treatments in ecological experiments: re-evaluating the ecosystem

function of biodiversity. Oecologia 110:449–460.

Huston, M. A. 2000. No Consistent Effect of Plant Diversity on Productivity. Science

289:1255a–11255.

Ives, A. R., and S. R. Carpenter. 2007. Stability and Diversity of Ecosystems. Science 317:58–

62.

Johnson, D. R., and F. Pomati. 2020. A brief guide for the measurement and interpretation of

microbial functional diversity. Environmental microbiology.

Jousset, A., B. Schmid, S. Scheu, and N. Eisenhauer. 2011. Genotypic richness and dissimilarity

opposingly affect ecosystem functioning. Ecology Letters 14:537–545.

Kowalchuk, G. A., and J. R. Stephen. 2001. Ammonia-oxidizing bacteria: a model for

678   molecular microbial ecology. Annual review of microbiology 55:485–529.

679   Kozich, J. J., S. L. Westcott, N. T. Baxter, S. K. Highlander, and P. D. Schloss. 2013.

680   Development of a Dual-Index Sequencing Strategy and Curation Pipeline for Analyzing

681   Amplicon Sequence Data on the MiSeq Illumina Sequencing Platform. Applied and

682   Environmental Microbiology 79:5112–5120.

683   Krause, S., X. Le Roux, P. A. Niklaus, P. M. Van Bodegom, J. T. Lennon, S. Bertilsson, H.-P.

684   Grossart, L. Philippot, and P. L. E. Bodelier. 2014. Trait-based approaches for

685   understanding microbial biodiversity and ecosystem functioning. Frontiers in

686   Microbiology 5:1–10.

687   Kumar, S., M. Herrmann, B. Thamdrup, V. F. Schwab, P. Geesink, S. E. Trumbore, K.-U.

688   Totsche, and K. Küsel. 2017. Nitrogen Loss from Pristine Carbonate-Rock Aquifers of

689   the Hainich Critical Zone Exploratory (Germany) Is Primarily Driven by

690   Chemolithoautotrophic Anammox Processes. Frontiers in Microbiology 8:1–13.

691   Kumar, S., G. Stecher, M. Li, C. Knyaz, and K. Tamura. 2018. MEGA X: Molecular

692   Evolutionary Genetics Analysis across Computing Platforms. Molecular Biology and

693   Evolution 35:1547–1549.

694   Küsel, K., K. U. Totsche, S. E. Trumbore, R. Lehmann, and M. Herrmann. 2016. How deep

695   can surface signals be traced in the critical zone? Merging biodiversity with

696   biogeochemistry research in a central German Muschelkalk landscape. Frontiers in

697   Earth Science 4:1–18.

698   Lehman, C. L., and D. Tilman. 2000. Biodiversity, Stability, and Productivity in Competitive

699   Communities. The American Naturalist 156:534–552.

700   Litchman, E., and C. a. Klausmeier. 2008. Trait-Based Community Ecology of Phytoplankton.

701   Annual Review of Ecology, Evolution, and Systematics 39:615–639.

Loreau, M. 2000. Biodiversity and ecosystem functioning: recent theoretical advances. Oikos
91:3–17.

Losos, J. B. 2008. Phylogenetic niche conservatism, phylogenetic signal and the relationship
between phylogenetic relatedness and ecological similarity among species. Ecology
letters 11:995–1003.

Louca, S., M. F. Polz, F. Mazel, M. B. N. Albright, J. A. Huber, M. I. O'Connor, M.
Ackermann, A. S. Hahn, D. S. Srivastava, S. A. Crowe, M. Doebeli, and L. W. Parfrey.
2018. Function and functional redundancy in microbial systems. Nature Ecology &
Evolution 2:936–943.

Lynd, L. R., P. J. Weimer, W. H. van Zyl, and I. S. Pretorius. 2002. Microbial Cellulose
Utilization: Fundamentals and Biotechnology. Microbiology and Molecular Biology
Reviews 66:506–577.

Malik, A. A., J. B. H. Martiny, E. L. Brodie, A. C. Martiny, K. K. Treseder, and S. D. Allison.
2019. Defining trait-based microbial strategies with consequences for soil carbon
cycling under climate change. The ISME journal.

Maron, P.-A., A. Sarr, A. Kaisermann, J. Lévêque, O. Mathieu, J. Guigue, B. Karimi, L.
Bernard, S. Dequiedt, S. Terrat, A. Chabbi, and L. Ranjard. 2018. High Microbial
Diversity Promotes Soil Ecosystem Functioning. Applied and Environmental
Microbiology 84:e02738-17.

Martiny, A. C., K. Treseder, and G. Pusch. 2013. Phylogenetic conservatism of functional traits
in microorganisms. The ISME journal 7:830–838.

Mayfield, M. M., and J. M. Levine. 2010. Opposing effects of competitive exclusion on the
phylogenetic structure of communities. Ecology letters 13:1085–1093.

Maynard, D. S., T. W. Crowther, and M. A. Bradford. 2017. Competitive network determines
the direction of the diversity–function relationship. Proceedings of the National
Academy of Sciences:201712211.

McDonald, J. H. 2014. Handbook of Biological Statistics. 3rd edition. Sparky House

Publishing, Baltimore, Maryland.

Mokany, K., J. Ash, and S. Roxburgh. 2008. Functional identity is more important than

diversity in influencing ecosystem processes in a temperate native grassland. Journal of

Ecology 96:884–893.

Moore, E. R. B., B. J. Tindall, V. A. P. Martins Dos Santos, D. H. Pieper, J.-L. Ramos, and N.

J. Palleroni. 2006. Nonmedical: Pseudomonas. Pages 646–703 The Prokaryotes.

Springer New York, New York, NY.

Morrissey, E. M., R. L. Mau, E. Schwartz, J. G. Caporaso, P. Dijkstra, N. van Gestel, B. J.

Koch, C. M. Liu, M. Hayer, T. A. McHugh, J. C. Marks, L. B. Price, and B. A. Hungate.

2016. Phylogenetic organization of bacterial activity. The ISME Journal 10:2336–2340.

Mouchet, M. A., S. Villéger, N. W. H. Mason, and D. Mouillot. 2010. Functional diversity

measures: an overview of their redundancy and their ability to discriminate community

assembly rules. Functional Ecology 24:867–876.

Mukherjee, S., D. Stamatis, J. Bertsch, G. Ovchinnikova, H. Y. Katta, A. Mojica, I.-M. A.

Chen, N. C. Kyrpides, and T. B. K. Reddy. 2019. Genomes OnLine database (GOLD)

v.7: updates and new features. Nucleic Acids Research 47:D649–D659.

Murtagh, F., and P. Legendre. 2014. Ward's Hierarchical Agglomerative Clustering Method:

Which Algorithms Implement Ward's Criterion? Journal of Classification 31:274–295.

Naeem, S., D. E. Bunker, A. Hector, M. Loreau, and C. Perrings. 2009. Biodiversity, Ecosystem

Functioning, and Human Wellbeing. Oxford University Press.

Pascual-García, A., S. Bonhoeffer, and T. Bell. 2020. Metabolically cohesive microbial

consortia and ecosystem functioning. Philosophical transactions of the Royal Society of

London. Series B, Biological sciences 375:20190245.

Paula, F. S., J. L. M. Rodrigues, J. Zhou, L. Wu, R. C. Mueller, B. S. Mirza, B. J. M. Bohannan,

K. Nüsslein, Y. Deng, J. M. Tiedje, and V. H. Pellizari. 2014. Land use change alters

functional gene diversity, composition and abundance in Amazon forest soil microbial

communities. Molecular Ecology 23:2988–2999.

Petchey, O. L., and K. J. Gaston. 2002. Functional diversity (FD), species richness and

community composition. Ecology Letters 5:402–411.

Petchey, O. L., and K. J. Gaston. 2006. Functional diversity: back to basics and looking

forward. Ecology Letters 9:741–758.

Pruesse, E., J. Peplies, and F. O. Glöckner. 2012. SINA: Accurate high-throughput multiple

sequence alignment of ribosomal RNA genes. Bioinformatics 28:1823–1829.

Purschke, O., B. C. Schmid, M. T. Sykes, P. Poschlod, S. G. Michalski, W. Durka, I. Kühn, M.

Winter, and H. C. Prentice. 2013. Contrasting changes in taxonomic, phylogenetic and

functional diversity during a long-term succession: insights into assembly processes.

Journal of Ecology 101:857–866.

Quast, C., E. Pruesse, P. Yilmaz, J. Gerken, T. Schweer, P. Yarza, J. Peplies, and F. O.

Glöckner. 2012. The SILVA ribosomal RNA gene database project: improved data

processing and web-based tools. Nucleic Acids Research 41:D590–D596.

R Core Team. 2019. R: A Language and Environment for Statistical Computing.

Reintjes, G., C. Arnosti, B. Fuchs, and R. Amann. 2019. Selfish, sharing and scavenging

bacteria in the Atlantic Ocean: a biogeographical study of bacterial substrate utilisation.

The ISME Journal 13:1119–1132.

Reintjes, G., C. Arnosti, B. M. Fuchs, and R. Amann. 2017. An alternative polysaccharide

uptake mechanism of marine bacteria. The ISME Journal 11:1640–1650.

Rivett, D. W., T. Scheuerl, C. T. Culbert, S. B. Mombrikotb, E. Johnstone, T. G. Barraclough,

and T. Bell. 2016. Resource-dependent attenuation of species interactions during

bacterial succession. The ISME journal 10:2259–2268.

Roger, F., S. Bertilsson, S. Langenheder, O. A. Osman, and L. Gamfeldt. 2016. Effects of

multiple dimensions of bacterial diversity on functioning, stability and

multifunctionality. Ecology 97:2716–2728.

Salles, J. F., F. Poly, B. Schmid, and X. Le Roux. 2009. Community niche predicts the

functioning of denitrifying bacterial assemblages. Ecology 90:3324–3332.

Schloss, P. D., S. L. Westcott, T. Ryabin, J. R. Hall, M. Hartmann, E. B. Hollister, R. A.

Lesniewski, B. B. Oakley, D. H. Parks, C. J. Robinson, J. W. Sahl, B. Stres, G. G.

Thallinger, D. J. Van Horn, and C. F. Weber. 2009. Introducing mothur: Open-Source,

Platform-Independent, Community-Supported Software for Describing and Comparing

Microbial Communities. Applied and Environmental Microbiology 75:7537–7541.

Schwab, V. F., M. Herrmann, V. N. Roth, G. Gleixner, R. Lehmann, G. Pohnert, S. Trumbore,

K. Küsel, and K. U. Totsche. 2017. Functional diversity of microbial communities in

pristine aquifers inferred by PLFA- and sequencing-based approaches. Biogeosciences

14:2697–2714.

Shade, A. 2017. Diversity is the question, not the answer. The ISME Journal 11:1–6.

Shade, A., R. R. Dunn, S. A. Blowes, P. Keil, B. J. M. Bohannan, M. Herrmann, K. Küsel, J.

T. Lennon, N. J. Sanders, D. Storch, and J. Chase. 2018. Macroecology to Unite All

Life, Large and Small. Trends in Ecology & Evolution 33:731–744.

Smith, P., and M. Schuster. 2019. Public goods and cheating in microbes. Current biology: CB

29:R442–R447.

Srivastava, D. S., M. W. Cadotte, A. A. M. Macdonald, R. G. Marushia, and N. Mirotchnick.

2012. Phylogenetic diversity and the functioning of ecosystems. Ecology Letters

15:637–648.

Steudel, B., C. Hallmann, M. Lorenz, S. Abrahamczyk, K. Prinz, C. Herrfurth, I. Feussner, J.

W. R. Martini, and M. Kessler. 2016. Contrasting biodiversity-ecosystem functioning

relationships in phylogenetic and functional diversity. New Phytologist 212:409–420.

Taubert, M., J. Stähly, S. Kolb, and K. Küsel. 2019. Divergent microbial communities in
groundwater and overlying soils exhibit functional redundancy for plant-polysaccharide
degradation. PloS one 14:e0212937.

Thompson, P. L., T. J. Davies, and A. Gonzalez. 2015. Ecosystem Functions across Trophic
Levels Are Linked to Functional and Phylogenetic Diversity. PLOS ONE 10:e0117595.

Tilman, D. 1997. Distinguishing between the Effects of Species Diversity and Species
Composition. Oikos 80:185.

Turnbull, L. A., J. M. Levine, M. Loreau, and A. Hector. 2013. Coexistence, niches and
biodiversity effects on ecosystem functioning. Ecology Letters 16:116–127.

Turner, S., K. M. Pryer, V. P. W. Miao, and J. D. Palmer. 1999. Investigating Deep
Phylogenetic Relationships among Cyanobacteria and Plastids by Small Subunit rRNA
Sequence Analysis. The Journal of Eukaryotic Microbiology 46:327–338.

Venail, P., K. Gross, T. H. Oakley, A. Narwani, E. Allan, P. Flombaum, F. Isbell, J. Joshi, P.
B. Reich, D. Tilman, J. Ruijven, and B. J. Cardinale. 2015. Species richness, but not
phylogenetic diversity, influences community biomass production and temporal
stability in a re-examination of 16 grassland biodiversity studies. Functional ecology
29:615–626.

Violle, C., D. R. Nemergut, Z. Pu, and L. Jiang. 2011. Phylogenetic limiting similarity and
competitive exclusion. Ecology letters 14:782–787.

Wang, Y., R. Zhang, Z. He, J. D. Van Nostrand, Q. Zheng, J. Zhou, and N. Jiao. 2017.
Functional Gene Diversity and Metabolic Potential of the Microbial Community in an
Estuary-Shelf Environment. Frontiers in Microbiology 8:1–12.

Ward, J. H. 1963. Hierarchical Grouping to Optimize an Objective Function. Journal of the
American Statistical Association 58:236–244.

Warren, R. A. 1996. Microbial hydrolysis of polysaccharides. Annual review of microbiology
50:183–212.

Weber, M. G., and S. Y. Strauss. 2016. Coexistence in Close Relatives: Beyond Competition

and Reproductive Isolation in Sister Taxa. Annual review of ecology, evolution, and

systematics 47:359–381.

Yin, B., D. Crowley, G. Sparovek, W. J. De Melo, and J. Borneman. 2000. Bacterial functional

redundancy along a soil reclamation gradient. Applied and environmental microbiology

66:4361–4365.

Zhang, Q.-G., and D.-Y. Zhang. 2006. Resource availability and biodiversity effects on the

productivity, temporal variability and resistance of experimental algal communities.

Oikos 114:385–396.

Zhang, Z., S. Schwartz, L. Wagner, and W. Miller. 2000. A Greedy Algorithm for Aligning

DNA Sequences. Journal of Computational Biology 7:203–214.

Figure legends

Figure 1

Metabolic and phylogenetic classification and grouping of groundwater isolates. EEAs are

shown for twenty-seven groundwater isolates incubated with eleven fluorescent substrate

analogues. The heatmap scale shows the concentration of substrate that was broken down in

the EEA assays. Empty (white) boxes denote activity outside the detectable range. The left

dendrogram shows clusters of isolates with similar metabolic traits based on Euclidean

hierarchical diversity distances. The topmost dendrogram shows hierarchical clustering of

substrates that produced similar activity patterns in the assays. The assignment of isolates to

the communities of each treatment group is depicted to the right of the heatmap. Double-

coloured boxes indicate isolates that were assigned to more than one group. In the legend, the

colours indicate the community group and the phylogenetic (P) and metabolic (M) traits of the

communities are indicated as similar (sim) or dissimilar (dis).

Figure 2

Individual communities plotted against the phylogenetic and metabolic diversity within

and between groups. Colours indicate the community groups; red for Psim-Msim group, green

for Psim-Mdis group, blue for Pdis-Msim group and purple for Pdis-Mdis group. Phylogenetic

and metabolic diversity metrics have been z-score normalised. Variation of diversity scores

exists within groups, but the groups are cleanly separated from each other (see Table S3 for

statistical significance).

Phylogenetic vs. metabolic diversity

 Figure 3

 Heatmap of community EEA % proportional changes. The heatmap scale depicts

 proportional changes from -100 to 100%. Communities that exhibited changes over 100% have

 their percent change value shown in the heatmap matrix. Red colours depict a higher activity

 of the community; blue colours depict a lower activity of the community than estimated, based

 on the activities of the individual isolates. White boxes show measurements outside the

 detection limits. On the right, the label for each row indicates the community and on the left

 the group from where the activities are taken from; red for Psim-Msim group, green for Psim-

 Mdis group, blue for Pdis-Msim group and purple for Pdis-Mdis group communities. Columns

 indicate the substrates used in the EEA assays. The dendrogram depicts hierarchical clustering

 of each substrate's activities, showing groups of substrates with similar activity responses.

 Figure 4

 Effects of phylogenetic diversity (A) and metabolic diversity (B) on the EEA proportional

 changes of communities. Black regression lines show the effect of each treatment when

 controlling for the dissimilar groups of the second treatment. Grey regression lines show the

 effect of each treatment when controlling for the similar groups of the second treatment.

 Statistical significance for each linear regression is shown in each plot, with the respective

 coloured text (grey or black). Coloured squares in (B) and (C) indicate community groups:

 Psim-Msim in red, Psim-Mdis in green, Pdis-Msim in blue and Pdis-Mdis in purple. The y axes

 have been log2 transformed and the x axes have been z-scale transformed.

 Figure 5

 Temporal trends of EEA changes between communities and isolates from 0 h to 24 h after

 community mixing. Rows show community groups and columns show the substrate where the

 activity change was noted. The average changes between the timepoints are indicated by the

colour of the arrows and their statistical significance (Student's *t*-test, 2-tailed) showed on top

of the arrows. P values below 0.001 are shown as *<0.001* and values over 0.05 are marked as

non-significant (*ns*).

Figure 6

Substrate acclimation effect on isolates and communities. EEA of Psim-Mdis group isolates

and communities on 4-MUB-α-D-glucopyranoside (aG) are shown as an example of the

substrate acclimation effect. In green are shown the non-acclimated EEAs of cultures as a fold-

change of 1, to which all other activities were normalised. In orange, the cultures received the

substrate at the same time as the previous cultures and left to acclimate for 4.5 h. Lastly, the

EEAs of cultures that had been mixed for 6 h but not left to acclimate, as shown in blue. Bars

depict average fold-changes in the EEA of isolates and communities and error bars show the

standard deviation (n = 5). Community EEAs showed a ~5-fold increase when acclimated with

the substrate and isolates only showed a ~1-fold increase.

Figure 7

Substrates drive the strength and direction of phylogenetic and metabolic diversity

treatment effects. (A) The linear regression model slopes for all 11 substrates and the average

for all substrates (Table S10) are plotted here as points. In (B) the average for all substrates,

community EEA, is positively related with phylogenetic diversity ($p < 0.001$). In (C) the

average for all substrates, community EEA, is negatively related with metabolic diversity

($p < 0.001$). Coloured points in (A) indicate different substrates and the average (red).

Biochemically similar substrates are circled in dashed ellipses; blue for fatty acids, orange for

monosaccharides and green for polysaccharides. Coloured squares in (B) and (C) indicate

community groups: Psim-Msim in red, Psim-Mdis in green, Pdis-Msim in blue and Pdis-Mdis

in purple.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7