Astrocyte-like glia-specific gene *deathstar* is crucial for normal development, adult

locomotion and lifespan of male *Drosophila*

Hadi Najafi[1], Kyle Wong[1], Ammar Salkini[1], Hongyu Miao[2], and Woo Jae Kim [1,2,*]

[1]Department of Cellular and Molecular Medicine, Faculty of Medicine, University of Ottawa,

Ottawa, Ontario K1H 8M5, Canada

[2]The HIT Center for Life Sciences, Harbin Institute of Technology, Harbin, China.

*Correspondence to: wkim@hit.edu.cn

HN: h.najafi2999@gmail.com KW: kwong132@uottawa.ca AS: asalk017@uottawa.ca HM:

20210125l@stu.hit.edu.cn

ABSTRACT

The nervous system of *Drosophila* is composed of diverse cell types with substantial homologies

to various cell types of more complex species, including the human. Among all nervous system

cell types, astrocyte-like glia (ALG) is necessary for the proper physiology and behavior of the

fly and human. In this study, we exploited the gene expression profile of single cells in

Drosophila optic lobe to identify the genes with specific expression pattern in each subtype glia.

Through bioinformatics analysis of the data, a novel ALG-specific gene (here assigned as

deathstar, des) was identified. Immunostaining of *des* in the central nervous system (CNS)

showed its presence in specific regions of *Drosophila* brain and ventral nerve cord, previously

characterized as ALG cells. Consistent with the bioinformatics analysis, *des*-related signals were

overlapped with the signals of the previously-reported ALG marker. At physiological level,

RNAi-mediated suppression of *des* gene impeded the normal development of male flies without

any effects on females. Cell type-specific expression of *des* RNAi showed that *des* affects

locomotion behaviour and lifespan of *D. melanogaster*, in an ALG-specific manner. Taken

together, we showed that bioinformatically identified ALG-specific gene *deathstar* play critical

roles in development and physiology of *Drosophila*.

KEYWORDS: Astrocyte-like glia, *CG11000*, Cell type, Male development, *Drosophila*

melanogaster, Single cell RNA-seq, Specific marker, *deathstar, des*

INTRODUCTION

The brain of *D. melanogaster* is a complex organ with a diverse set of cells serving analogous

functions to those of human [1]. With differential expression of the genes, each cell in the brain

can determine its own fate to be a particular cell type with specific functional characteristics [2].

Previous studies have identified and reported the existence of several cell types in nervous

system of *D. melanogaster* assigned as; perineural glia (PNG), subperineural glia (SPG), cortex

glia (CG), neuropil ensheathing glia (EGN), tract ensheathing glia (EGT), astrocyte-like glia

(ALG) and neurons (Neu) [3].

The function of the central nervous system (CNS) depends on a complex network of several

neuronal and glial cell types [4,5]. Due to the feasibility of genetic manipulation and behavioral

analysis, *Drosophila melanogaster* has emerged as a powerful model organism for studying the

development and function of the nervous system [6]. There is a functional homology between the

cell types in the nervous system of *D. melanogaster* and those of more advanced organisms,

including human. Glial cells constitute an essential part of CNS [7], and until now, several

distinct types of these cells were also identified in the *Drosophila* nervous system [1,3]. Each

cell types have a distinct profile of gene expression [8] and have their specific functions matched

to their physical territories or physiological states [9]. Based on the differences in gene

expression profile, each cell type of *Drosophila* nervous system can be distinguished by a set of

genes, which are known as molecular markers for each cell type [3].

In addition to the existed molecular markers for each cell type, there may be some

uncharacterized genes in *D. melanogaster*, which are capable of discerning a specific cell type in

the nervous system. Moreover, a specific cell type likely differs only in the expression of the

limited number of genes which confer a new function to it [10,11]. In this study, we used a

publicly available expression profile of 55,000 single cells of *Drosophila* optic lobe in order to

find novel molecular markers for specific cell types, following by experimental confirmation of

the predicted results.

RESULTS

Cell type identification of the single cells of *Drosophila* optic lobe based on their specific

markers

To identify the cells corresponding to distinct subtypes of glia and neurons, the expression

profile of the sequenced single cells by Konstantinides et al. [2] was analyzed based on their

previously-reported marker genes (Fig. 1A). According to the expression profile of 55000

sequenced single cells, the cells with the highest expression of each cell type-specific marker

were selected and classified. A total of forty-two cells were identified and classified into seven

cell types. The heatmap of gene expression for these selected cells confirmed the unique

expression pattern of each marker in the corresponding cell types (Fig. 1B). *CG4797* and *gem* are

differentially expressed in PNG, and SPG were distinguished by the three

markers *Mdr65*, *Moody* and *Gli*. Other glial cells were successfully distinguished by single

marker; *Cyp4g15*, *CG9657*, *CG34340* and *Eaat1*. Moreover, these genes could individually

separate CG, EGN, EGT and ALG, respectively. Neuronal cells were determined based on the

highest expression of *nSyb* and *elav* genes (Fig. 1B).

Hierarchical clustering of *Drosophila* neural cell subtypes based on their similarities in

transcriptome

As shown above, different types of *Drosophila* neural cells could be classified based on some

specific marker genes (Fig. 1). In order to know whether the classified cells in each category

have the same expression profile or not, correlation analysis between the selected cells was

carried out based on their transcriptome analysis. Results showed that most of the cells

expressing the same genetic marker(s) tend to be clustered together, suggesting their similar

transcriptome and accuracy of the identification of cell types (Fig. 2A). However, the cells with

the highest expression of SPG markers and some other cells could not be classified in a single

cluster suggesting their transcriptome dissimilarity or partial insufficiency of the available

genetic markers for clustering (e.g. expression heterogeneity of the SPG

markers: *Mdr65*, *Moody* and *Gli*).

With at least 3 out of 6 similar cells in each category, differential expression analysis was

performed in order to find the genes with specific expression in each cell type. While the

identified PNG were positive for the expression of their two known genetic markers

(*CG4797* and *gem*) (Fig. 1B), there were two sub-clusters of PNGs (here assigned as PNG-α and

PNG-β) in our transcriptome analysis which suggests the presence of a novel subtype for PNG

(Fig. 2A). Additional similar analysis was performed on another set of 42 cells of 55,000 cells

and similar clustering results were obtained, which could support the accuracy of cell type

characterization and clustering (Supplemental Fig.S1). The list of differentially expressed genes

(DEGs) for each cell type in addition to the DEGs between the two subtypes of PNG is presented

in Supplemental file S2.

Using the transcriptome data of the studied cell types and their clustering results, we assigned the

degree of differences between them as a dendrogram (Fig. 2B). Accordingly, the Venn diagram

was used to illustrate the number of unique and shared DEGs for each cell type. The extent of

unique DEGs for each cell type is as follows: 98.33% (59 of 60) for PNG, 100% (1 of 1) for

SPG, 90% (36 of 40) for CG, 60% (3 of 5) for EG, 78.95 (75 of 95) for ALG and 96.39 (454 of

471) for neurons (Fig. 2C).

Identification of *des* as a gene with specific enrichment in astrocyte-like glia of *D.*

melanogaster

After cell type identification, we asked whether specific genes can also characterize the

identified cell types. To identify the genes capable of distinguishing different cell types, we

performed differential expression analysis in order to compare the transcriptome of the cluster of

interest with that of all other cell types. The list of the DEGs for each cell type are presented in

Supplemental file S2.

Among all the clustered cells, ALG could be sufficiently discriminated by a single

uncharacterized gene named *CG11000* (here assigned as *deathstar, des)* [4]. Expression level

analysis of this gene using the RNA-seq profile of 84 single cells in *Drosophila* optic lobe [2]

demonstrated that among the 48 cells analyzed, 12 cells (which correspond to ALG) express *des*

at highest levels (Fig. 3A). In the same RNA-seq expression data (accession number:

GSE103771), a significant positive correlation (R^2=0.6637; *p*-value = 0.0001) was observed

between the expression level of the candidate gene (*des*) and that of *Eaat1*, which previously was

reported as an ALG-specific marker [7] (Fig. 3B). In addition, the expression pattern of *des* and

Eaat1 across developmental time of *D. melanogaster*, derived from a different RNA-seq data

(accession number: GSE107049) showed a positive correlation between these two genes (Fig. 3C

and 3D), supporting the expression specificity of *des* gene in ALG of *D. melanogaster*.

To corroborate the assigned specificity of *des*, we tested whether the promoter-*GAL4* lines of this

gene could label ALG in CNS of *D. melanogaster*. Consistently, data showed that *des* (Red

signals) is mostly co-labelled with the ALG marker *Eaat1* (Green signals), in specific bilateral

parts of *Drosophila* optic lobe, pars intercerebrails (PI) region, and small part of subesophageal

ganglion (SOG) in male but not female brain (Fig. 4A). Intriguingly, *des-GAL4* identified 4-5

cells with large cell body towards the upper border of SOG in each hemisphere that were not

labeled by *Eaat1-lexA* driver (bottom panels in Fig. 4A) or previously characterized *ALG-GAL4*

driver [12] (Fig. 4B). We discovered that des-GAL4 further marks ALG cells in the MesoNM of

both male and female VNC (Fig. 4C). As shown in the brain SOG area, des-GAL4 identified 10

to 12 cells with large cell bodies in the ProNM/AMNp and MetaNm/ANn regions. Two of these

male-specific cells are uniquely designated by Eaat1-lexA, showing that male-specific des

expression in these distinct cell lineages (Compare MesoNm panels between male and female in

Fig. 4C). These cells with big cell bodies cannot be labeled using the previously published ALG-

GAL4 reagent [12] (Fig. 4D). All of these results indicate that *des-GAL4* identifies *Eaat1-lexA-*

labeled ALG cells as well as particular brain and VNC cells.

Sex-biased developmental effect of *des* gene in *D. melanogaster*

We crossed GAL4 strains that are expressed in each neural cell types of *D. melanogaster* with

the *des-RNAi* to knockdown the expression of *des* gene in a cell type-specific manner. Our

results showed that most of the males expressing *des-RNAi* in all neural cell types are lethal

during development. However, the females with the same genotype developed normally to adult.

Both males and females in the control group (harboring *des-RNAi* without *GAL4* expression)

showed normal development (Fig. 5A). This result suggests that the *des* gene may have a

developmental function exclusively in male flies. However, such developmental effect was not

cell type-specific. As a control, the Drosophila CG15765 gene, which is less likely to express

with des in our analysis, was evaluated using the same methodology, and a standard male-to-

female ratio (~1) was obtained for all genotypes (Fig. 5B). The exact number of the progenies of

the corresponding crosses are summarized in the Supplemental Table S1.

Knockdown of *des* gene affects locomotion activity of *D. melanogaster*

It has been previously reported that ALG cells that cover specific territories of *Drosophila* CNS

have significant effects on locomotion behaviours [13]. To investigate the role of *des* gene in

locomotion behavior in *D. melanogaster*, we knocked down the *des* gene expression through

expressing a *des*-specific RNAi within different cell types of *Drosophila* CNS. Then we

performed the climbing assay. Results showed that the flies that specifically express *des-RNAi* in

astrocytes exhibited significant defects in climbing ability (Fig. 6A), suggesting the crucial role

of this gene in this cell type. Interestingly, we found that *des* knockdown does not affect the

climbing activity in female flies (Fig. 6B). When we knocked down the *des* in EGN and SPG,

however, female flies showed severe defects in climbing activity (Fig. 6B). These data suggest

that the expression of *des* in ALG cells is crucial only in males, not females regarding the

climbing activity.

To test whether the behavioural locomotive effect of *des* gene is altered during the ageing of *D.*

melanogaster, the climbing assay was performed on the flies with different ages. Results showed

that the older flies (three- and four-week-old flies) expressing *des-RNAi* in all neural cell types

exhibited significant climbing defect (Fig. 6C). When we tested the same genotype of flies in the

one- and two-week stages, the climbing defects appeared in ALG-specific manner (Fig. 6C).

However, climbing defects were not striking for the flies with neuron-specific suppression

of *des*, suggesting the climbing defects related to *des* is glial-specific in old flies (Fig. 6C).

Unlike the male flies, both young and old females did not exhibit any meaningful pattern of

climbing defects (Fig. 6D). All these data suggest that the expression of *des* in male ALG cells is

crucial for climbing ability in the early adult stages

Astrocyte-specific suppression of *des* gene shortens lifespan in *D. melanogaster*

To test the possible function of *des* gene in *D. melanogaster* lifespan, we knocked

down *des* expression in distinct CNS cell types (CG, PNG, ALG, EGN, EGT, SPG, Glia and

Neurons) and measured their life span for 40~50 days. We observed a significant reduction in

lifespan for the flies that had ALG-specific expression of *des* RNAi, supporting the specific

effect of the *des* gene in ALG on the life span of *D. melanogaster* (Fig. 7). The ALG-specific

effect of *des* RNAi on lifespan was observed in both male and female flies (Fig. 7A and 7B),

suggesting that the role of *des* on the lifespan is not sexually dimorphic. However, in the female

flies, such reducing effects of *des* RNAi were also observed in other cell types (EGN, EGT, PNG

and neurons) but at lower levels than that of ALG (Fig. 7B).

DISCUSSION

In this study, we analyzed a newly released scRNA seq dataset including typical glial cell marker

genes (Fig. 1) and uncovered intriguing characteristics of the *Drosophila* glial lineage (Fig. 2).

Among the unique marker genes revealed throughout our investigation, the *deathstar* (*des*) gene,

CG11000, might precisely designate ALG (Fig. 2). *des* expresses significantly in ALG

throughout early adulthood (Fig. 3). The expression of *des-GAL4* demonstrates that *des* and

Eaat1 are significantly coexpressed in the adult head and VNC (Fig, 4). *des* gene knockdown

using subtype-specific glia-*GAL4s* reveals glial roles of *des* throughout development (Fig. 5),

adult locomotion (Fig. 6), and lifetime (Fig. 7).

Recent advances of single-nuclear RNA-sequencing (snRNA-seq) technologies enable fly

community to establish a cell atlas for the entire adult *Drosophila* [14]. Using the publicly

accessible Fly Cell Atlas annotation dataset in SCope website, we examined the expression

patterns of *deathstar* (*des*) in several tissues with distinct annotations. In the entire tissues of both

male and female flies, the expression of *des* strongly overlapped with glial cells (yellow dots in

Fig. S2A-B). *des* expression is also detectable in annotated neurons (magenta dots in Fig. S2C-

D). When consolidated, tiny clusters of glial cells could be distinguished from neurons (black-

dotted circles in Fig. S2E-F).

To determine the ALG expression of *des*, we employed the previously identified ALG markers

astrocytic leucine-rich repeat molecule (*alrm*) and *Eaat1*. The SCope platform does not yet

support well-defined ALG annotations. Intriguingly, the glial cell clusters observed in Fig. S1E-

F were also recognized by *alrm* (Fig. S3A-B) and *Eaat1* (Fig. S3C-D) in all tissues. We believe

that these tiny clusters are ALG cells in adult flies. This ALG cluster starts to be appeared from

the 3 days old adult stage (Fig. S2G-J). The head exhibited more ALG cells than the entire body

(Figure S3E-H), and the annotated glial cells in SCope demonstrate co-expression of *des* with

alrm and *Eaat1* in a unique ALG cluster (Fig. S3I-J).

To investigate if this unique cluster of cell types is ALG, we examined every subtype of glial cell

identified in SCope. The ALG clusters in the head (gray-dotted region in Fig. S4) did not overlap

with neuron (Fig. S4A-B), adult brain cell body glial cell (Fig. S4C-D), adult brain perineurial

glial cell (Fig. S4E-F), adult lamina epithelial/marginal glial cell (Fig. S4G-H), adult optic

chiasma glial cell (Fig. S4O-P). The ALG cluster in the body (gray-dotted region in Fig. S5) did

not overlap with neuron (Fig. S5A-B), CNS surface associated glial cell (Fig. S5C-D), cell body

glial cell (Fig. S5E-F), subperineurial glial cell (Fig. S5G-H), and perineurial glial sheath (Fig.

S5I-J). However, we found that *des* expression pattern in the body overlaps with adult reticular

neuropil associated glial cell (Fig. S5K-L). These data suggest that our bioinformatics analysis of

ALG-associated *des* expression is also confirmed by FCA dataset. We believe the *des*-specific

ALG cluster in the body represents the *des-GAL4*-labeled large cell body cells in the brain and

VNC (Fig. 4).

There are pieces of evidence for ALG-enriched gene that have different patterns of expression

during early and late stages of *D. melanogaster* development. ALG-enriched genes in larval

stages are mainly involved in metabolism, energy production, and protein synthesis, consistent

with the known role of astrocytes in metabolic support of neurons. In comparison with the larval

astrocytes, ALG cells in adult *D. melanogaster* acquire a new function for regulation of

behaviors, additional to their functions in earlier stages [15]. Therefore, it is expected that the

suppression of the *des* gene in larval stages is lethal in males. However, this suppression at

adulthood stages mediates climbing defects in males (Fig. 5 and 6). In accordance of this idea,

the *des* protein has a 'Pleckstrin homology domain' (PH domain). PH domain can bind

phosphatidylinositol within biological membranes and proteins, such as the beta/gamma subunits

of heterotrimeric G proteins and protein kinase C, and play a role in recruiting proteins to

different membranes, thereby targeting them to appropriate cellular compartments or allowing

them to interact with other components of signal transduction pathways [16].

Analysis of lifespan in the flies expressing *des* RNAi in different cell types showed the greatest

effects when *des* is suppressed specifically in ALG cells which was observed in both sexes,

providing another supportive evidence for specificity of the gene to ALG cell type (Fig. 7). It is

known that *Drosophila* astrocytes (ALG) are significant contributors to modulate

neurotransmitter homeostasis within synapses [17] and also regulators of circadian rhythm [18].

Therefore, it is likely that abolishing the functions of ALG or ALG-specific genes could interfere

with normal locomotion properties and lifespan of *D. melanogaster*, which are the direct

outcome of dysregulations in neurotransmission and circadian rhythm, respectively. Moreover, it

was recently reported that mitochondrial dysfunctions in ALG cells contribute to

neurodegeneration [17,19], which could give rise to shortened lifespan of *D. melanogaster* [20].

Taken together, *des* (previously known as *CG11000*) was reported as a protein-coding gene in

fruit fly with 332 amino acids. Its functional role was not known yet in *Drosophila*, and it has no

homologue gene in human. However, it is predicted to be involved in regulation of cell

differentiation process (Flybase). In this study, we firstly annotated *des* as an ALG-specific gene

which could mediate the developmental process of the male flies additional to its functions in

locomotion and lifespan properties of *D. melanogaster*. Further investigations will shed insight

on how this membrane protein with a PH domain may specify the identity of ALG from neurons

and influence *Drosophila* physiology.

MATERIALS AND METHODS

Dataset

A single-cell RNA-sequencing (RNA-seq) data of a recently published paper [2] was selected

from the data repository NCBI Gene Expression Omnibus (GEO) (accession number:

GSE103771). It contained the expression profile of 17272 *Drosophila* genes in a population of

55000 single cells from the optic lobe of *Drosophila* brain. Also, the data of expression profile

for the whole brain of *D. melanogaster* across four different developmental stages (days 5, 10, 20

and 40 after eclosion) were downloaded from NCBI GEO (accession number: GSE107049).

Bioinformatics analyses

For expression analysis of the genes in single cells of *D. melanogaster* brain, the expression

matrix belonging to the study of Konstantinides et al. [2] was downloaded for downstream

analyses. We analyzed the expression matrix of the single cell RNA-seq data of *Drosophila* optic

lobe (accession number: GSE103771) based on the expression level of known markers,

previously reported for each cell type. Based on the expression pattern of the specific markers,

the type of the cells in the matrix was characterized. For each cell type, six single cells were

selected according to the expression level of the markers. Forty-two cells comprising seven cell

types of *Drosophila* nervous system (assigned as perineural glia (PNG), subperineural glia

(SPG), cortex glia (CG), neuropil ensheathing glia (EGN), tract ensheathing glia (EGT),

astrocyte-like glia (ALG) and neurons (Neu)) were selected for downstream analyses. For

confirmation of this cell type classification, hierarchical clustering was performed for the

selected forty-two cells based on their similarities in transcriptome and presented as a heatmap

with color codes for the Pearson correlation coefficient. The clusters with at least three cells (of

all six cells) were validated as unique cell type. For further confirmation, this analysis was

performed similarly on another set of forty-two cells from the 55,000 single cells. To identify the

genes enriched in a specific cell type, we first identified differentially expressed genes (DEGs)

between the examined cell types and all other cells by the R package limma [21]. Then, after

applying adjusted *P*-values (Benjamini–Hochberg correction) of < 0.05 and $\log_2 FC \geq 1.5$, the

specificity of the candidate genes with highest degree of enrichment was determined by analysis

of their expression across all the examined single cells (unpaired t-test; *P*-value < 0.05). Venn

diagram was obtained from the web-based tool on

http://bioinformatics.psb.ugent.be/webtools/Venn/.

Fly stocks and crosses

All the *Drosophila melanogaster* lines in this study were prepared from Bloomington Drosophila

Stock Center (BDSC). After receiving the stocks, they were cultured in standard conditions

(standard cornmeal-agar yeast fly food at 25° C on a 12-h light:12-h dark condition) as

previously described [22]. CO_2 was used as an anesthetic. For functional analysis of the genes,

the *GAL4/UAS-RNAi* system was used to deplete their functions in a cell-type specific manner.

The fly stocks together with their crosses for functional analyses of *des* gene are summarized in

table 1. After crossing the virgin females of *UAS-des*-RNAi line with males of different GAL4

driver lines (*X-GAL4*), ensuring expression of RNAi in cell type-specific manner, the transgenic

progenies (*X-GAL4 > UAS-RNAi*) were collected based on their genotypes and assessed for their

behaviors (Table 1). The specific *GAL4* driver gene for each cell type was as below: *CG4797*

gene for PNG, *Mdr65* gene for SPG, *Cyp4g15* gene for CG, *CG9657* gene for EGN, *CG34340*

gene for EGT, *Eaat1* gene for ALG, *repo* gene for all glial cells (glia) and *nSyb* gene for

neurons. For control of the experiment flies without expression of RNAi were used. These flies

had *RNAi* in their genome without expression of *GAL4*. Therefore, no functional *RNAi* was

produced in the control flies. All of our experiments were performed in Canton-S/wild type

background to rule out all the genetic background effects.

For the crosses of fluorescent labeling experiments, the two binary systems of *GAL4/UAS-*

mCD8GFP and *LexA/LexAoP-mCD8RFP* were used simultaneously in an individual male and

female flies. The ALG cells of *Drosophila* CNS were labelled by the expression of a membrane-

bound GFP (*mCD8GFP*) under the control of *Eaat1* gene promoter, fused to the sequence of

LexA (*Eaat1*-LexA). For tracing the expression pattern of *des* gene in *Drosophila* CNS, a

transgenic fly line carrying *des-GAL4* construct (BDSC#: 23418) was used. Expression of RFP

under the control of *des* gene promoter depicts the expression distribution of *des* gene across

CNS. The crossing scheme for this experiment is summarized in table 2.

Immunostaining and imaging

As described before [21], brains dissected from adults 5 days after eclosion were fixed in 4%

formaldehyde for 30 min at room temperature, washed with 1% PBT three times (30 min each)

and blocked in 5% normal donkey serum for 30 min. The brains were then incubated with

primary anti-bodies in 1% PBT at 4° C overnight followed with fluorophore-conjugated

secondary antibodies for 1 hour at room temperature. Brains were mounted with antifade

mounting solution (Invitrogen, catalog #S2828) on slides for imaging.

Primary antibodies used here are chicken anti-GFP (Aves Labs, 1:1000), rabbit anti-DsRed

express (Clontech, 1:250) and mouse anti-Bruchpilot (nc82) (DSHB, 1:50). Fluorophore-

conjugated secondary antibodies used here are Alexa Fluor 488-conjugated goat anti-chicken

(Invitrogen, 1:100). Alexa Fluor 488-conjugated donkey anti-rabbit (Invitrogen, 1:100), RRX-

conjugated donkey anti-rabbit (Jackson Laboratory, 1:100), and Dylight 405-conjugated donkey

anti-mouse (Jackson Laboratory, 1:100).

Images were collected using a Zeiss LSM 7 MP microscope scanning confocal microscope.

Projection of the confocal stacks, adjusting the brightness and contrast of the images were

performed by NIH ImageJ [26], then presented by Adobe Illustrator CC 2019 for windows.

Developmental lethality assays

To assess the possible function of *des* gene in the development of *D. melanogaster* from pupae to

adulthood stage, the newly eclosed *D. melanogaster* adults of the crosses in Supplemental Table

1, were collected and counted based on their genotypes. After counting, data were presented as

the percentage of male and female flies with *des* RNAi to the total number of eclosed flies.

Lifespan assay

Lifespan was measured at room temperature according to standard protocols. In brief, newly

ecloded flies (0 to 3 days) were collected (50 per genotype) and then placed in vials (25 flies per

vial) and transferred to fresh vials every three days. Survival was recorded for each genotype

(two vials). We scored flies stacked in the food as death events in all the vials analyzed. We

created survival curves with Prism version 6 (GraphPad Software, San Diego, CA, USA) using

the method of Kaplan and Meier. For statistical significance of the experiment, one-way

ANOVA method was used and the death rate of each genotype was compared to the Canton-S

(CS) flies as the control. *P-value < 0.05, ** P-value < 0.01.

Climbing assay

For the climbing assay, flies were separated into groups of 25 adults per each vial and kept in 29°

C. At the time of experiment, the groups of 25 flies were placed in an empty climbing vial and

then tapped down to the bottom. They were allowed to climb from the bottom to top of the vial

(8 cm) for 20 seconds. In order to evaluate the climbing pattern of each genotype, the climbing

height achieved by each genotype was calculated in every 5 seconds after the tapping and

initiating negative geotaxis. The results of climbing assay were obtained by video recording of

climbing flies then processed by ImageJ software. Data was presented as graphs using Prism

version 6 (GraphPad Software, San Diego, CA, USA). Flies with wild-type/*UAS-RNAi* genotype

were used as control group.

Statistical analysis

Statistical analysis and graphic representations were performed using Prism 6.00 for Windows

(GraphPad Software, San Diego, CA, USA). Unpaired t test or one-way ANOVA (using the

Bonferroni's multiple comparison) were applied depending on the measurements analyzed in the

corresponding experiment. In all cases averages with SEM are plotted. *P*-values were calculated

by two-tailed unpaired Student's t-test unless otherwise specified, using GraphPad Prism 6

software. All data are presented as the mean ± SEM. $*P < 0.05$; $**P < 0.01$; and $***P < 0.001$.

Single-nucleus RNA-sequencing Analyses - Data and Code Availability

snRNAseq dataset analyzed in this paper is published in [14] and available at the Nextflow

pipelines (VSN, https:// github.com/vib-singlecell-nf), the availability of raw and processed

datasets for users to explore, and the development of a crowd-annotation platform with voting,

comments, and references through SCope (https://flycellatlas.org/ scope), linked to an online

analysis platform in ASAP (https://asap.epfl.ch/fca).

Gene expression pattern analyses in different tissues

For the gene expression pattern of the 10 genes involved in SMD in each cell type of leg and

other tissues, we used the single-cell RNA-seq data from https://flycellatlas.org [14], and the 10x

Genomics stringent loom files were downloaded. The cell types are split by FCA.

The digital expression matrices were analyzed with the Seurat 4.1.0 R package [23]. The dot

plots of the 10 genes involved in SMD in each cell type of different tissues were then made using

the 'DotPlot' function with broad annotation (broad cell types) and the annotation (detailed cell

types), respectively.

LIST OF ABBREVIATIONS

ALG: Astrocyte-like glia; **BBB**: Blood-brain barrier; **CG**: Cortex glia; **CNS**: Central nervous

system; **CS**: Canton S; **DEG**: Differentially-expressed gene; **dsRNA**: double strand RNA; **EGN**:

Neuropil ensheathing glia; **EGT**: Tract ensheathing glia; **GEO**: Gene expression Omnibus;

NEU: Neurons; **PNG**: Perineural glia; **RNAi**: RNA interference; **SPG**: Subperineural glia; **GFP**:

Green fluorescent protein; **RFP**: Green fluorescent protein.

DECLARATIONS

Ethics approval and consent to participate: Not applicable.

Consent for publication: Not applicable.

Availability of data and materials: Not applicable.

Competing interests:

The authors declare that they have no competing interests.

Funding: This research was supported a University of Ottawa Startup grant to WJK, a

University of Ottawa Brain and Mind Research Institute/Center for Neural Dynamics Open call

project grant to WJK, a University of Ottawa Interdisciplinary Research Group Funding

Opportunity (IRGFO stream 1 and 2) Grant to WJK, a Natural Sciences and Engineering

Research Council of Canada (NSERC) Discovery grant (reference: 211406) to WJK, a Mitacs

Globalink Research Internship Program grant to WJK, and Startup funds from HIT Center for

Life Science to WJK. This research was also supported by the Brain Pool Program of the

National Research Foundation in Korea to WJK, Burroughs Wellcome Fund Collaborative

Research Travel Grants (reference: 1017486) to WJK and a NVIDIA Academic Hardware Grant

Program to WJK.

AUTHOR CONTRIBUTIONS:

Concept and design of experiments: WJK, HN and KW; performing the experiments: HN, KW,

WJK and AS; acquisition of data: HN, KW, WJK and AS; analysis and interpretation of data:

HN, KW and WJK; revision experiments and analysis: HM; drafting or revising the article: HN

and WJK.

ACKNOWLEDGMENTS:

Stocks obtained from the Bloomington Drosophila Stock Center (NIH P40OD018537) were used

in this study. We also thank Khoi-Nguyen Ha Nguyen for his contribution to experiments

performed in this manuscript.

FIGURE LEGENDS

Figure 1. Identification of *Drosophila* neural cell types through analysis of their specific

marker genes. (A) Specific marker genes, previously reported as markers for each cell type of

Drosophila glial cells. (B) Heatmap shows the expression pattern of the markers across 42

selected cells from 55000 single sequenced cells, in which, six different cells were identified as a

unique cell type (denoted by different colors). Cells' numbers correspond to the numbering

method of Konstantinides et al. Red and black colors indicate the marker-positive and marker-

negative cells, respectively. Color intensities in the heatmap denote the level of gene expression

based on RPKM per single cell.

Figure 2. Hierarchical clustering of different neural cell types of *D. melanogaster* based on

their transcriptome. (A) Hierarchical cluster analysis of gene expression for 42 selected cells

from *Drosophila* optic lobe based on the expression level of 17,272 genes for each single cell.

Except SPG, all the studied cell types were clustered together suggesting their similarities in

expression profile. PNG cells were strongly clustered in two different clusters, suggesting

presence of two distinct PNG subtypes in *Drosophila* (PNG-α and PNG-β). (B) Dendrogram

shows the similarity degree of the clustered cells based on whole transcriptome. (C) Number of

unique and shared DEGs for each cell type of *D. melanogaster* nervous system. Total number of

DEGs for each cell type are presented in parenthesis. PNG= perineural glia, SPG= subperineural

glia, CG= cortex glia, ALG= astrocyte-like glia, EG= ensheathing glia, and NEU= neurons.

Figure 3. Expression specificity of *des* gene in ALG cells of *D. melanogaster*. (A) While the

ALG cells exhibited high level of *des* mRNA (Blue points), other glia cells as well as neurons

showed low or no expression of this gene (gray points). (B) Positive correlation between

expression level of known ALG marker (*Eaat1* gene) and that of *des* gene. Blue and gray points

respectively denote expression of the genes in ALG and non-ALG single cells. (C) Similar

expression pattern of *des* (dark blue) and *Eaat1* (bright blue) genes during development of male

D. melanogaster. (D) Similar expression pattern of *des* (dark blue) and *Eaat1* (bright blue) genes

during development of female *D. melanogaster*. Developmental times are presented as the days

after eclosion of the flies. SPG= subperineural glia, PNG= perineural glia, CG= cortex glia,

ALG= astrocyte-like glia, EGN= neuropil ensheathing glia, EGT= tract ensheathing glia, and

NEU= neurons.

Figure 4. *des* gene is sex-specifically expressed in *Drosophila* ALG cells. Immunostaining

result of (A) brain and (C) VNC in male and female flies. In both images, the ALG cells were

stained with anti-GFP (green), and the red signals (anti-RFP) reflect *UAS-mCD8RFP* driven by a

GAL4, tagged by the promoter region of *des* gene (*des-GAL4*). Anti-nc82 was used to label

neuropil (Blue). Co-labeling of astrocytic cells using *Eaat1-lexA/lexAop-mCD8GFP* (green)

together with *des-GAL4/UAS-mCD8RFP* (Red) showed ALG-specific expression of *des* gene

among other types of *Drosophila* glia cells. Scale bars represent 100 µm in top panels and 50 µm

in others. Dotted boxes represent the magnified area in the next panels. Dotted entangles indicate

the regions of *des*-positive and *Eaat1*-positive cells found. Red arrows indicate des-positive but

Eaat1-negative large cell body cells and Empty red-line arrows indicate those cells are empty in

Eaat1-lexA/lexAop-mCD8GFP panels. (B and D) Female (B) brain and (D) VNC of flies

expressing ALG-GAL4/UAS-mCD8GFP. Source image lsm files (~500MB) were downloaded

from FlyLight platform constructed by Janelia Farm Research Center (JFRC) then reconstructed

using ImageJ https://www.janelia.org/project-team/flylight. PI: pars intercerebrails; OL: optic

lobe; SOG; subesophageal ganglion; VNC; ventral nerve cord; ProNm: Prothoracic neuromere;

AMNp: Accessory Mesothoracic neuropil; MesoNm: Mesothoracic neuromere; MetaNm:

Metathoracic neuromere; ANm: bdominal neuromere [24].

Figure 5. Male-specific developmental effect of *des* gene in *D. melanogaster*. (A) While the

females with *des-RNAi* showed normal development, male flies with the same genotypes could

not be matured to adults and showed overall less frequencies in comparison with the expected

ratio of each genotype. (B) Using another RNAi as control (*CG15765* RNAi), equal number of

males and females were observed which were consistent with the expected ratio of each

genotype. $*P < 0.05$; $**P < 0.01$; $***P < 0.001$ and $****P < 0.0001$, n.s.= non-significant ($P \geq$

0.05).

Figure 6. Effects of *des* gene suppression on locomotion behavior of *D. melanogaster*. (A)

ALG-specific reduction of climbing activity in male flies under suppression of *des* gene. (B)

Slight EGN- and SPG-specific reduction of climbing behavior in female *D. melanogaster* under

suppression of *des* gene. (C) Climbing behaviors of male *D. melanogaster* expressing *des* RNAi

during different times of development (weeks 1-4 post-eclosion) showing ALG-specific effect of

the gene. (D) Climbing behaviors of female *D. melanogaster* expressing *des* RNAi during

different times of development (weeks 1-4 post-eclosion) showed no significant effect in ageing

flies.

Figure 7. Lifespan effect of *des* gene suppression in *D. melanogaster*. Suppression of *des* gene

shortened lifespan of male and female flies in an ALG-specific manner. Only the males with

ALG-specific suppression of *des* showed significant reduction in survival. In females, the

greatest level of lifespan reduction was observed for the flies with ALG-specific suppression of

des. However, females carrying *des* RNAi in their EGN, EGT, PNG and neurons showed such an

effect with lower extent. CS refers wild-type/*des-RNAi* control flies.

SUPPLEMENTARY FIGURE LEGENDS

Figure S1. Hierarchical clustering of single cells selected based on specific markers for

neural cell types of *D. melanogaster*. Among all cell types, the ALG and neurons had the best

clustering pattern.

Figure S2. Whole tissue Fly SCope dataset. (A-F) Annotations and gene names are color-coded

using red, green, and blue words. When cells overlap, the color of the dots is either yellow, cyan,

or magenta. Colored arrows indicate cells that have been double- or triple-labeled by annotations

and genes. (G-J) Whole tissue Fly SCope dataset in day1, 2, 3, and 5 with the expression of *des,*

alrm, and *Eaat1*.

Figure S3. Fly SCope dataset. (A-J) Tissues, sexes, annotations and gene names are color-coded

using red, green, and blue words. When cells overlap, the color of the dots is either yellow, cyan,

or magenta. Dotted entangles indicate probabl cell populations of *des*-positive ALG cells.

Figure S4. Head region of Fly SCope dataset. (A-P) Tissues, sexes, annotations and gene names

are color-coded using red, green, and blue words. When cells overlap, the color of the dots is

either yellow, cyan, or magenta. Dotted entangles indicate probabl cell populations of *des*-

positive ALG cells.

Figure S5. Body region of Fly SCope dataset. (A-L) Tissues, sexes, annotations and gene names

are color-coded using red, green, and blue words. When cells overlap, the color of the dots is

either yellow, cyan, or magenta. Dotted entangles indicate probabl cell populations of *des*-

positive ALG cells.

Table 1. Summary of crosses and experimental genotypes for behavioral analyses. In order

to generate the transgenic flies with expression of *des*-RNAi in cell type-specific manner, the

male flies of different *GAL4* driver lines were crossed with the females harboring *des*-dsRNAi in

their genome (BDSC#: 51918). All the females (♀) for these crosses were virgin. Among the

progenies of these crosses, the flies with proper genotypes were selected based on the absence of

balancer chromosomes (CyO, TM3 and TM6B).

Table 2. Crossing scheme for imaging experiments of CNS in *Drosophila*. The line A

contains RFP and GFP reporter genes under regulation of UAS and LexAop sequences,

respectively. Line B expresses lexA protein in an ALG-specific manner. By crossing the lines A

and B (Parents), the fly line C (F_1 generation) was produced and then crossed with *des-GAL4*

line (D) to produce F_2 generation. Among the flies of F_2, the males and females that lack

balancer chromosomes but contain the components of the binary systems *UAS/GAL4* and

LexA/LexAop were selected for central nervous system (CNS) dissection. The females (♀) of all

crosses were virgin.

SUPPLEMENTAL DATA CAPTIONS

Supplemental file 1. List of differentially expressed genes for each cell type of _Drosophila_

central nervous system. The genes with $\log_2$fold change (Log_2FC) ≥ 1.5 and _P_-value < 0.05

were considered as differentially expressed genes (Denoted as highlighted cells).

Supplemental Table 1. Number of newly-eclosed flies under suppression of _des_ gene in

neural cell types of _D. melanogaster_. The number of male progenies was less than the females

when the _des_ expression was suppressed in neural cell types of _D. melanogaster_ (RNAi carriers).

As control of the experiment, a fly line which had no expression of _des_ RNAi (referred as

balancer carrier) was used. Equal number of males and females was observed for this control

(CS) group.

REFERENCES

1. Freeman MR. Drosophila central nervous system glia. Cold Spring Harbor perspectives in
biology. 2015;7: a020552.

2. Konstantinides N, Kapuralin K, Fadil C, Barboza L, Satija R, Desplan C. Phenotypic
convergence: distinct transcription factors regulate common terminal features. Cell. 2018;174:
622-635. e13.

3. Kremer MC, Jung C, Batelli S, Rubin GM, Gaul U. The glia of the adult D rosophila nervous
system. Glia. 2017;65: 606–638.

4. Zeng Z, Miao N, Sun T. Revealing cellular and molecular complexity of the central nervous
system using single cell sequencing. Stem cell research & therapy. 2018;9: 234.

5. Brody T, Odenwald WF. Cellular diversity in the developing nervous system: a temporal view
from Drosophila. Development. 2002;129: 3763–3770.

6. Olesnicky E, Wright E. Drosophila as a model for assessing the function of RNA-binding
proteins during neurogenesis and neurological disease. Journal of developmental biology.
2018;6: 21.

7. Yildirim K, Petri J, Kottmeier R, Klämbt C. Drosophila glia: Few cell types and many
conserved functions. Glia. 2019;67: 5–26. doi:10.1002/glia.23459

8. Shih M-FM, Davis FP, Henry GL, Dubnau J. Nuclear transcriptomes of the seven neuronal
cell types that constitute the Drosophila mushroom bodies. G3: Genes, Genomes, Genetics.
2019;9: 81–94.

9. Croset V, Treiber CD, Waddell S. Cellular diversity in the Drosophila midbrain revealed by
single-cell transcriptomics. Elife. 2018;7: e34550.

10. Henry GL, Davis FP, Picard S, Eddy SR. Cell type–specific genomics of Drosophila neurons.
Nucleic acids research. 2012;40: 9691–9704.

11. Hilu-Dadia R, Hakim-Mishnaevski K, Levy-Adam F, Kurant E. Draper-mediated JNK
signaling is required for glial phagocytosis of apoptotic neurons during Drosophila
metamorphosis. Glia. 2018;66: 1520–1532.

12. Kremer MC, Jung C, Batelli S, Rubin GM, Gaul U. The glia of the adult Drosophila nervous
system. Glia. 2017;65: 606–638. doi:10.1002/glia.23115

13. Peco E, Davla S, Camp D, Stacey SM, Landgraf M, Meyel DJ van. Drosophila astrocytes
cover specific territories of the CNS neuropil and are instructed to differentiate by Prospero, a
key effector of Notch. Development. 2016;143: 1170–1181.

14. Li H, Janssens J, Waegeneer MD, Kolluru SS, Davie K, Gardeux V, et al. Fly Cell Atlas: A
single-nucleus transcriptomic atlas of the adult fruit fly. Science. 2022;375: eabk2432.
doi:10.1126/science.abk2432

15. Liu L, Zhang K, Sandoval H, Yamamoto S, Jaiswal M, Sanz E, et al. Glial lipid droplets and
ROS induced by mitochondrial defects promote neurodegeneration. Cell. 2015;160: 177–190.

16. Wang DS, Shaw R, Hattori M, Arai H, Inoue K, Shaw G. Binding of Pleckstrin Homology
Domains to WD40/β-Transducin Repeat Containing Segments of the Protein Product of the Lis-
1 Gene. Biochem Bioph Res Co. 1995;209: 622–629. doi:10.1006/bbrc.1995.1545

17. Kounatidis I, Chtarbanova S. Role of glial immunity in lifespan determination: a Drosophila
perspective. Frontiers in immunology. 2018;9: 1362.

18. Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. limma powers differential
expression analyses for RNA-sequencing and microarray studies. Nucleic acids research.
2015;43: e47–e47.

19. Guo A, Li L, Xia S, Feng C, Wolf R, Heisenberg M. Conditioned visual flight orientation in
Drosophila: dependence on age, practice, and diet. Learning & Memory. 1996;3: 49–59.

20. Brand AH, Perrimon N. Targeted gene expression as a means of altering cell fates and
generating dominant phenotypes. development. 1993;118: 401–415.

21. Kim WJ, Jan LY, Jan YN. A PDF/NPF Neuropeptide Signaling Circuitry of Male Drosophila
melanogaster Controls Rival-Induced Prolonged Mating. Neuron. 2013;80: 1190–1205.
doi:10.1016/j.neuron.2013.09.034

22. Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis.
Nature methods. 2012;9: 671.

23. Satija R, Farrell JA, Gennert D, Schier AF, Regev A. Spatial reconstruction of single-cell
gene expression data. Nat Biotechnol. 2015;33: 495–502. doi:10.1038/nbt.3192

24. Court R, Namiki S, Armstrong JD, Börner J, Card G, Costa M, et al. A Systematic
Nomenclature for the Drosophila Ventral Nerve Cord. Neuron. 2020;107: 1071-1079.e2.
doi:10.1016/j.neuron.2020.08.005

A

Cell type	Marker gene
PNG	CG4797
	gem
SPG	Mdr65
	moody
	Gli
CG	Cyp4g15
EGN	CG9657
EGT	CG34340
ALG	Eaat1
NEU	nSyb
	elav

— PNG: Perineurial glia
— CG: Cortex glia
— SPG: Subperineurial glia
— EGN: Neuropile ensheathing glia
— EGT: Tract ensheathing glia
— ALG: Astrocyte-like glia
— NEU: Neurons

B

Hadi., Fig.1

Hadi., Fig.2

A

deathstar gene

B

R² = 0.6637
P-value < 0.0001
n = 84

C

Male

D

Female

Hadi., Fig.3

A *des-GAL4, Eaat1-lexA/UAS-mCD8RFP, lexAop-mCD8GFP*

B *ALG-GAL4(R86E01)/UAS-mCD8RFP*

Hadi., Fig.4

C *des-GAL4, Eaat1-lexA/UAS-mCD8RFP, lexAop-mCD8GFP*

D *ALG-GAL4(R86E01)/UAS-mCD8RFP*

Hadi., Fig.4

Hadi., Fig.5

Hadi., Fig.6

Hadi., Fig.7

Hadi., Fig.S1

Hadi., Fig.S2

Hadi., Fig.S3

Hadi., Fig.S4

A Body · *des* · ♂ · glial cell · neuron · •des⁺ neurons · •des⁺ glia

B Body · *des* · ♀ · glial cell · neuron · •des⁺ neurons · •des⁺ glia

C Body · *des* · ♂ · glial cell · CNS surface associated glial cell

D Body · *des* · ♀ · glial cell · CNS surface associated glial cell

E Body · *des* · ♂ · glial cell · cell body glial cell

F Body · *des* · ♀ · glial cell · cell body glial cell

G Body · *des* · ♂ · glial cell · subperineurial glial cell

H Body · *des* · ♀ · glial cell · subperineurial glial cell

I Body · *des* · ♂ · glial cell · perineurial glial sheath

J Body · *des* · ♀ · glial cell · perineurial glial sheath

K Body · *des* · ♂ · glial cell · adult reticular neuropil associated glial cell

L Body · *des* · ♀ · glial cell · adult reticular neuropil associated glial cell

Hadi., Fig.S5

Parents for crosses		Progenies
UAS- RNAi lines (♀)	GAL4 lines (♂)	
w+ ; +/+ ; *deathstar*-RNAi (# 51918)	wild type	wild type
w+ ; +/+ ; *deathstar*-RNAi	+ ; +/+ ; PNG-GAL4/ TM3	Non-TM3
w+ ; +/+ ; *deathstar*-RNAi	+ ; +/+ ; SPG-GAL4/ TM3	Non-TM3
w+ ; +/+ ; *deathstar*-RNAi	+ ; +/+ ; CG-GAL4/ TM3	Non-TM3
w+ ; +/+ ; *deathstar*-RNAi	+ ; +/+ ; EGN-GAL4/ TM3	Non-TM3
w+ ; +/+ ; *deathstar*-RNAi	+ ; +/+ ; EGT-GAL4/ TM3	Non-TM3
w+ ; +/+ ; *deathstar*-RNAi	+ ; +/+ ; ALG-GAL4/ TM6B	Non-TM6B
w+ ; +/+ ; *deathstar*-RNAi	+ ; +/+ ; Neuron-GAL4/ TM6B	Non-TM6B

Table 1. Summary of crosses and experimental genotypes for behavioral analyses. In order to generate the transgenic flies with expression of *deathstar*-RNAi in cell type-specific manner, the male flies of different GAL4 driver lines were crossed with the females harboring UAS-*deathstar*-dsRNAi in their genome (BDSC#: 51918). All the females (♀) for these crosses were virgin. Among the progenies of these crosses, the flies with proper genotypes were selected based on the absence of balancer chromosomes (CyO, TM3 and TM6B).

Lines	Fly stocks (Genotypes and Bloomington ID)	Cross scheme
A	UAS-mCD8RFP, lexAop-mCD8GFP ; Sp/CyO ; TM2/TM6B (#32229)	P: A ♀ × B ♂ F$_1$: C ♂ × D ♀ F$_2$: E (♂ or CNS dissec
B	ALG-LexA/CyO ; TM2/TM6B (#45914)	
C	UAS-RFP, lexAop-GFP ; ALG-LexA/CyO ; TM2/TM6B	
D	*deathstar* -GAL4 (# 23416)	
E	UAS-RFP, lexAop-mCD8GFP ; ALG-LexA/+ ; TM6B/ *deathstar* -GAL4	

Table 2. Crossing scheme for imaging experiments of CNS in *Drosophila*. The line A contains RFP and GFP reporter genes under regulation of UAS and LexAop sequences, respectively. Line B expresses LexA protein in an ALG-specific manner. By crossing the lines A and B (Parents), the fly line C (F$_1$ generation) was produced and then crossed with *deathstar* -GAL4 line (D) to produce F$_2$ generation. Among the flies of F$_2$, the males and females that lack balancer chromosomes but contain the components of the binary systems UAS/GAL4 and LexA/LexAop were selected for central nervous system (CNS) dissection. The females (♀) of all crosses were virgin.

Supplemental Table S1. Number of newly-eclosed flies under suppression of *Deathstar* gene in neural cell types of *D. melanogaster*. The number of male progenies was less than the females when the *Deathstar* expression was suppressed in neural cell types of *D. melanogaster* (RNAi carriers). As control of the experiment, a fly line which had no expression of *Deathstar* RNAi (referred as balancer carrier) was used. Equal number of males and females was observed for this control (CS) group.

Day of collection	Fly genotypes	CS ♂	CS ♀	CG ♂	CG ♀	PNG ♂	PNG ♀	ALG ♂	ALG ♀	EGN ♂	EGN ♀	EGT ♂	EGT ♀	SPG ♂	SPG ♀	Glia ♂	Glia ♀	Neu. ♂	Neu. ♀
1	RNAi carriers	7	15	2	6	4	6	4	12	7	24	1	5	1	12	7	9	12	11
1	Balancer carriers	2	4	13	12	10	12	18	6	22	24	6	11	15	10	5	39	26	24
2	RNAi carriers	8	8	1	7	3	5	3	4	1	6	1	1	0	9	4	5	4	6
2	Balancer carriers	2	6	7	16	4	6	4	3	4	14	5	8	5	9	8	32	8	16
3	RNAi carriers	2	1	2	10	7	5	4	5	0	15	2	5	1	10	2	4	2	4
3	Balancer carriers	4	3	16	10	5	6	4	8	2	13	8	12	13	9	4	14	17	19
4	RNAi carriers	12	12	0	10	1	4	1	4	2	3	0	1	2	4	3	6	6	8
4	Balancer carriers	3	7	7	12	2	3	8	2	4	1	4	2	2	8	7	6	8	10
5	RNAi carriers	18	16	6	17	4	5	10	12	3	9	1	1	6	23	2	3	12	20
5	Balancer carriers	11	9	29	25	11	10	20	22	20	25	7	6	24	23	30	21	45	35
6	RNAi carriers	12	11	3	5	2	7	3	15	3	9	2	4	1	9	6	19	0	6
6	Balancer carriers	5	6	6	6	5	12	15	11	30	25	5	4	10	15	14	16	7	7