RUNNING HEAD: Discordant genomic and phenotypic clines

TITLE: When polymorphism and monomorphism meet: discordant genomic and phenotypic

clines across a lizard contact zone

AUTHORS: Caroline M. Dong[1,2]*, Claire A. McLean[1,2], Adam Elliott[1], Adnan Moussalli[2]†,

Devi Stuart-Fox[1]†

[1]School of BioSciences, The University of Melbourne, Parkville, Vic., Australia

[2]Sciences Department, Museums Victoria, Carlton Gardens, Vic., Australia

*Author for correspondence

†Joint last author

CORRESPONDING AUTHOR

Caroline M. Dong

School of BioSciences, The University of Melbourne, Parkville, Victoria 3010, Australia.

E-mail: caroline.dong@unimelb.edu.au

Abstract

Colour polymorphism can promote rapid evolution and speciation, particularly when

populations differ in the number or composition of morphs. The contact zone between

Ctenophorus modestus (swift dragon) and *C. decresii* (tawny dragon) is a compelling study

system in which to examine evolutionary processes and outcomes when polymorphic and

monomorphic populations meet. *Ctenophorus modestus* is polymorphic for male throat

coloration and lacks ultraviolet (UV) reflectance while *C. decresii* is monomorphic with UV-

blue throats. We characterised genomic and phenotypic clines across the contact zone based

on single nucleotide polymorphisms, the mitochondrial ND4 gene, and male colour traits, and

concurrently assessed the phenotype of captive-bred F1 hybrids. Our results indicate that

genomic introgression is asymmetric, with high frequencies of backcrossing to *C. modestus*

but not *C. decresii*, accompanied by the prevalence of the *C. modestus* mtDNA haplotype in

hybrids. The clines for throat phenotype are abrupt and displaced towards the range of *C.*

decresii, relative to the genetic and dorsolateral phenotype clines. By contrast, both throat and

dorsolateral phenotypes in captive-bred F1 hybrids are intermediate. Our results are consistent

with the hypothesis that throat coloration, a polymorphic sexual signal in *C. modestus*, is the

target of selection during incipient speciation and provide insight into the microevolutionary

processes that may link polymorphism and speciation.

Keywords: coloration, *Ctenophorus decresii*, *Ctenophorus modestus*, ddRAD, hybridisation,

reproductive isolation, speciation

Introduction

Colour polymorphism, the co-existence of multiple colour morphs within a single interbreeding population (Huxley, 1955), is possibly the best studied model of discrete intraspecific phenotypic variation. Recently, there has been great interest in the role of colour polymorphism in promoting rapid phenotypic evolution and speciation, particularly when populations differ in the number, type, and frequency of morphs (Corl, Davis, Kuchta, & Sinervo, 2010; Forsman, Ahnesjö, Caesar, & Karlsson, 2008; Gray & McKinnon, 2007; Hugall & Stuart-Fox, 2012; McLean & Stuart-Fox, 2014; West-Eberhard, 1986). However, we lack detailed studies on the microevolutionary processes linking colour polymorphism and speciation. In particular, the genomic and phenotypic consequences of secondary contact between polymorphic and monomorphic lineages remain unclear, especially where the monomorphic lineage represents a novel morph.

Based on the widespread importance of colour signals in reproductive isolation (Boughman, 2001; Hooper, Griffith, & Price, 2019; Jiggins, Naisbit, Coe, & Mallet, 2001; Mavárez et al., 2006; Sætre et al., 1997; Seehausen et al., 2008) and the predicted genetic architecture of polymorphism (Sinervo & Svensson, 2002), there is expected to be a high probability of incompatibility during secondary contact between populations that differ in morph number, type, or frequency. Gene flow may result in the breakdown of adaptive genetic correlations and generate genetic incompatibilities between morphs (Pryke & Griffith, 2009). More generally, incompatibilities may arise when coadapted alleles are brought together for the first time in hybrids (Dobzhansky-Muller incompatibilities; Dobzhansky, 1937; Muller, 1942) or when coadapted nuclear and organelle (e.g. mitochondria and chloroplast) genomes that evolved in allopatry are disrupted (cytonuclear incompatibilities; Burton and Barreto 2012; Hill 2015). Further, a divergence in sexual phenotypic

characteristics may also be accompanied by behavioural incompatibilities in mating

preferences resulting in restricted gene flow and/or selection against hybrids. These

phenotypic and genetic incompatibilities together may facilitate the process of speciation

between populations which differ in morph frequencies and/or composition.

Here, we investigate the outcome of contemporary contact between the colour

polymorphic *Ctenophorus modestus* (swift dragon; Ahl, 1926) and monomorphic *C. decresii*

(tawny dragon*;* Duméril & Bibron, 1837), previously regarded as divergent lineages of *C.*

decresii sensu lato (Dong et al., in press). The two species differ in male dorsolateral and

throat coloration (Houston 1974; McLean et al. 2014b; Figures 1 and 2a). The polymorphic *C.*

modestus (previously northern lineage *C. decresii*) has four discrete male throat colour

morphs that are present in all populations and reflect little to no ultraviolet (UV): orange,

yellow, yellow with orange centre, and grey (Teasdale et al. 2013; Figure 1a). The black

dorsolateral stripe of *C. modestus* is relatively straight-edged and continuous, sometimes with

one distinct blotch near the neck, and with yellow/orange coloration terminating just behind

the shoulder (Figure 2a; McLean, Moussalli, Sass, & Stuart-Fox, 2013). By contrast,

monomorphic *C. decresii* (previously southern lineage) males have blue throat coloration with

a strong UV reflectance peak, representing a novel morph (Figure 1b; McLean et al. 2014b).

Males within an insular population (i.e. Kangaroo Island; "KI") have yellow reticulations in

addition to the UV-blue coloration (Figure 2a; McLean et al. 2014b). The black dorsolateral

stripe of *C. decresii* is discontinuous with two distinct blotches near the neck and coloration

terminating mid-body (Figure 2a; McLean et al., 2013).

The throat colour polymorphism is likely to have evolved in *C. decresii* sensu lato and

then was subsequently lost in *C. decresii* sensu stricto, thereby facilitating the evolution and

fixation of the novel blue morph (McLean, Stuart-Fox, et al., 2014). Genetic divergence

between the species (3.7% net sequence divergence in mtDNA) is consistent with range

contraction to allopatric refugia during glacial-interglacial Pleistocene cycles (Byrne, 2008;

McLean, Stuart-Fox, et al., 2014), where differing site-specific selective pressures may have

led to the evolution of divergent phenotypes. A small number of genetically admixed

individuals have been identified where the species meet geographically, suggesting that

subsequent post-glacial range expansion has resulted in a secondary contact zone (McLean,

Stuart-Fox, et al., 2014).

This is a compelling study system to explore the evolutionary consequences of

hybridisation between colour polymorphic and monomorphic species which differ in morph

number and type. Furthermore, the biology of both species and the polymorphism of *C.*

modestus are well-characterised. *Ctenophorus modestus* and *C. decresii* are sexually

dichromatic: females have cream coloured throats with a cryptic dorsal phenotype. Throat and

dorsolateral coloration are important intraspecific sexual signals, which are emphasised

during complex movement-based territorial and courtship displays (Gibbons, 1977, 1979;

Osborne, Umbers, Backwell, & Keogh, 2012; Ramos & Peters, 2016; Stuart-Fox & Johnston,

2005). Male throat coloration develops at sexual maturity (ca. 18 months), is fixed for life,

and autosomally inherited. The four morphs of *C. modestus* employ alternative behavioural

strategies (Yewers, Pryke, & Stuart-Fox, 2016) corresponding to differing hormonal profiles

(Yewers, Jessop, & Stuart-Fox, 2017), and are associated with MHC supertypes (Hacking,

Stuart-Fox, Godfrey, & Gardner, 2018). The discrete polymorphism is likely controlled by

two independently segregating loci, though the extent of orange or yellow within morphs

behaves as a quantitative (polygenic) trait (Rankin, McLean, Kemp, & Stuart-Fox, 2016).

We used a method of double digest restriction-site associated DNA sequencing to

identify single nucleotide polymorphisms (SNPs), sequenced the mitochondrial ND4 gene,

and characterised male throat and dorsolateral phenotype across a naturally occurring contact

zone between the polymorphic *C. modestus* and monomorphic *C. decresii*. Using geographic

cline analysis, we compared the frequency of these genetic and phenotypic traits along a

transect through the contact zone from pure *C. modestus* to pure *C. decresii* populations.

Furthermore, we conducted captive-breeding to investigate F1 hybrid phenotype independent

of exogenous selection. Specifically, we examined i) the geographic extent of hybridisation,

ii) whether hybridisation is due to recent or earlier admixture, iii) whether hybridisation is

bidirectional or asymmetric, and iv) whether there is evidence of selection on phenotype. We

predicted that selection on throat coloration to maintain the *C. modestus* polymorphism

should result in steeper and narrower clines compared to SNP and dorsolateral phenotype

clines.

Materials and Methods

Field sampling

Ctenophorus modestus and *C. decresii* are small [snout-vent-length (SVL) ≤ 90 mm]

diurnal rock-dwelling agamids endemic to South Australia. *Ctenophorus modestus* occurs in

the Flinders and Olary Ranges whereas *C. decresii* occurs in the Mount Lofty Ranges,

Fleurieu Peninsula, and on Kangaroo Island (Figure 2b). The contact zone is found in an area

of relatively low-lying grassland between the rocky ranges of the parental species, coinciding

with a change from relatively temperate to more semiarid conditions (Houston 1974; McLean

et al. 2014b). Our sampling design aimed to capture the full geographic range of both species

(Figure 2b, Table S1) to reaffirm previously described genetic divergence based on multi-loci

sequence data and microsatellites (McLean, Stuart-Fox, et al., 2014). To extend this previous

work we intensively sampled 14 sites along a north-south 160 km transect through the contact

zone (Figure 2c, Table S2). As the geographic extent of hybridisation was initially unknown,

the transect was anchored on either end by two genetically pure *C. decresii* (southernmost

sites 1 – 2) and *C. modestus* (northernmost sites 13 – 14) populations, which were

geographically connected along a rock range. Sites were demarcated by geographic distance

when possible; however, areas with high densities of sampling were demarcated more finely

to allow greater accuracy in analyses. Lizards were captured with a lasso (telescopic pole and

fishing line) or by hand and released at the site of capture following data collection. A genetic

sample (blood or tail tip stored in 99% ethanol at -20°C) and phenotypic data (described

below and in Supplementary Information) were taken from each individual. We focused on

collecting data from adult males (SVL $\geq$ 65 mm) and collected genetic samples from females

opportunistically. Our dataset comprised 234 genetic samples in total from across both

species' full ranges (Table S1, Figure 2b). This included 152 genetic samples from the 14

sites along the contact zone transect, from which we also obtained phenotypic data from up to

233 males along the contact zone transect, depending on the trait (Table S2).

SNP discovery

Genomic DNA was isolated and purified from blood or tissue using either an E.Z.N.A.

Tissue DNA Kit (Omega Bio-tek, Norcross, GA, USA) or a GenCatch Blood and Tissue

Genomic Mini-Prep Kit (Epoch Life Sciences, Sugar Land, TX, USA). DNA was sent to

Diversity Arrays Technology (Bruce, ACT, Australia) for SNP genotyping using DArTSeq, a

genome complexity reduction method similar to double digest restriction-site associated DNA

(ddRAD) sequencing protocol. Protocols are described in detail for broad applications in

Kilian et al., 2012 and specifically for the genus *Ctenophorus* in Melville et al., 2017. Briefly,

genomic DNA was digested with two restriction endonucleases (PstI/HpaII), PCR amplified

(1 min at 94°C; 30 cycles of 94°C for 20 s, 58°C for 30 s, 72°C for 45 s; final extension at

72°C for 7 min), and sequenced on an Illumina HiSeq2500. 'High-density' DArTseq SNP

arrays use 2,500,000 (±7%) reads per sample in marker calling. Two technical replicates of

each DNA sample were genotyped to assess reproducibility. DArT analysis pipelines and a

calling algorithm (DArTsoft14) were used to process sequences into a matrix of SNP loci. As

reference genomes, we used the *C. decresii* sensu lato draft genome (McLean, Stuart-Fox, and

Moussalli, unpublished data) and the fully annotated central bearded dragon genome (*Pogona*

vitticeps; Georges et al., 2015). In total, we obtained 172,893 SNPs from 234 individuals with

41.54% missing data. Using the R package *dartR* (v1.1.6, Gruber et al. 2018), we filtered this

dataset conservatively with thresholds of 100% reproducibility, ≤1% missing data per locus,

≤10% missing data per individual, and also removed loci that deviated from Hardy-Weinberg

Equilibrium ($P < 0.05$). This resulted in a dataset of 1,333 SNPs from 230 individuals with

0.26% missing data. Additionally, to evaluate potential bias associated with stringent filtering,

we repeated the filtering steps with a threshold of ≤5% missing data per locus (other

thresholds unchanged), resulting in a dataset of 6,889 SNPs from 225 individuals with 1.46%

missing data. Analyses run on this dataset indicated qualitatively similar results (Figure S1).

Mitochondrial DNA

We sequenced an 850 base-pair region of the mitochondrial genome which included

the NADH dehydrogenase subunit four (ND4), and tRNAhis, tRNAser and tRNAleu regions

using the primers ND4F (Forstner, Davis, & Arèvalo, 1995) and tRNA-Leu (Scott & Keogh,

2000) for a subset of 74 individuals from sites across the contact zone transect (Table S1).

PCR was performed following procedures in McLean, Stuart-Fox, et al., 2014. PCR products

were submitted to the Australian Genome Research Facility (Melbourne, Victoria, AUS) for

purification and forward and reverse sequencing on an Applied Biosystems 3730xl DNA

Analyzer. Sequences were edited and aligned with additional ND4 sequences spanning the

range of *C. modestus* and *C. decresii* (McLean, Stuart-Fox, et al., 2014) in Geneious (v6.1.7;

Kearse et al. 2012). Haplotype networks were constructed using the statistical parsimony

algorithms of TCS (v1.21; Clement et al. 2000).

Quantifying population structure and admixture

We examined genome-wide patterns of divergence using the Bayesian model-based

clustering approach implemented in STRUCTURE (v2.3.4; Pritchard, Stephens, & Donnelly,

2000). After an initial burn-in of 500,000 Markov chain Monte Carlo (MCMC) generations,

we ran 500,000 generations with 10 independent replicates for K values ranging from 1 to 5

under the "admixture" model with correlated allele frequencies and prior population

information, all other parameters were kept at default, and convergence of runs was visually

assessed. We used STRUCTURE HARVESTER (Earl & VonHoldt, 2012) to identify the

optimum value of K for our dataset and summarised results across the selected value of K

using CLUMPP (v1.1.2; Jakobsson & Rosenberg, 2007). We used the Bayesian estimation of

the *q*-value (proportion of an individual's genome inherited from each parental population)

with 95% confidence intervals (CI) as a hybrid index. Individuals were identified as pure

parentals or putative hybrids based on their hybrid index and 95% CI: an individual was

classified as parentals if *q* was less than 0.1 (i.e. *C. modestus*) or greater than 0.9 (i.e. *C.*

decresii), or of mixed ancestry if *q* was between 0.1 and 0.9.

We used previously identified population groups within each parental lineage to

further examine population structure (*C. modestus*: Northern Flinders Ranges [NFR],

Southern Flinders Ranges [SFR], Olary Ranges [OR]; *C. decresii*: Mainland South [MS],

Kangaroo Island [KI]; McLean et al. 2014b) and Contact Zone (CZ). We calculated expected

and observed heterozygosity, the number of alleles and allelic richness, estimated population

pairwise F_{ST}, and the inbreeding coefficient (F) for each population using the R packages

dartR (v1.1.6; Gruber et al. 2018) and *adegenet* (v2.1.1; Jombart & Ahmed, 2011). We then

performed a principal coordinate analysis (PCoA) using *dartR* to visualize pairwise genetic

distances between individuals.

Hybrid classification

We subsampled our SNP dataset to obtain a dataset of 200 loci with the highest

average polymorphic information content values using the R package *dartR* (v1.1.6; Gruber et

al. 2018). The dataset was analysed three times in parallel, each with a total of 100,000

MCMC generations after an initial burn-in of 50,000 generations, using the program

NewHybrids (v1.1 Anderson & Thompson, 2002) and the R package *parallelnewhybrid*

(Wringe, Stanley, Jeffery, Anderson, & Bradbury, 2017). We determined the probability that

individuals within the contact zone belonged to one of the six genotypic classes resulting from

the first two generations of hybridisation: parental forms (P1 and P2), F1 (P1 × P2), F2 (F1 ×

F1), and backcrossed (F1 × P1/P2). Although it is not always possible to unambiguously

assign individuals into discrete classes due to the continuum that may be found in wild

populations, classification may be suggestive of situations where contact is very recent or

where F1 hybrids are infertile based on the absence of F2 hybrids and backcrosses, or where

contact is several generations old based on the absence of pure parentals and F1 hybrids.

Inferring population history

We estimated population demographics with the program δaδi v2.0.6 (Gutenkunst,

Hernandez, Williamson, & Bustamante, 2009, 2010), which uses a diffusion approximation

method to analyse joint site frequency spectra. For this analysis we considered only

geographically connected populations of *C. modestus* and *C. decresii*, and therefore excluded

NFR and KI from the dataset (Figure 2). The full set of SNPs (i.e. 172,893 SNPs from 234

individuals) was filtered with a threshold of ≤5% missing data per locus, ≤10% missing data

per individual and 100% reproducibility, and secondaries were removed (i.e. retained one

SNP per locus) to reduce linkage disequilibrium (filtering performed in *dartR* (v1.1.6, Gruber

et al. 2018). This resulted in a dataset of 4,956 SNPs from 212 individuals with 1.48%

missing data, which we converted to δaδi input format using the R package *radiator* (v1.1.5;

Gosselin, Lamothe, Devloo-Delva, & Grewe, 2020).

We implemented a modified version of the model optimisation workflow

(dadi_pipeline) from Portik et al., 2017. We were specifically interested in inferring the age of

the contact zone and therefore tested two biologically realistic models of population history

consistent with the observed structure of the contact zone: 1) "sec_contact_asym_mig", where

the two species split with no migration (T1), followed by a period of asymmetrical migration

(T2; Figure 4); and 2) "sec_contact_asym_mig_three_epoch", a more complex model

involving a period of isolation after secondary contact (T3; Figure S2). Both models

considered two geographically structured 'populations', on either side of the density trough

between sites 2 and 3 (Figure 2): 1) *C. decresii* (MS; Figure 2) and 2) *C. modestus* pooled

with admixed individuals (OR, SFR and CZ; Figure 2). We used folded site frequency

spectra, a grid size of 200, 220, 240 and projection sizes of 60 and 180 alleles for *C. decresii*

and *C. modestus* respectively. For each model, we performed two independent runs with four

consecutive rounds of optimisation, with 100 replicates per round, increasing maximum

iterations (maxiter = 5, 5, 10, 20), and decreasing fold in parameter generation (fold = 3, 2, 2,

1). Models were compared using the log-likelihood and Akaike information criterion (AIC),

and the two independent runs converged on similar parameters (Table S5). We converted

parameters from the top model as follows: reference effective population size as $N_{ref} = \frac{\theta}{4\mu l}$,

where θ was the scaled population parameter, μ was a mutation rate of 9×10^{-10} (Singhal &

Moritz, 2013) and *l* was the length of analysed sequence (4956 SNPs × 70b = 34.7Mb);

population size as $Nu = N_{ref} \times nu$; time as $T = t2N_{ref} \times g$ where g was a generation time

of 2 years; and migration (individuals per generation) as $M_{ij} = \left(\frac{m_{ij}}{2 \times Nref}\right) \times nu_i$.

Captive-breeding and testosterone-induced expression of colour traits

260   To evaluate F1 hybrid phenotype independent of extrinsic selection pressures, we

conducted captive breeding using 36 pure *C. decresii* individuals (female = 17, male = 19)

collected from cline site 1 (southernmost site, longitude: 139.103°, latitude: -33.443°) and 41

pure *C. modestus* individuals (female = 20, male = 21) from cline site 14 (northernmost site,

longitude: 139.103°, latitude: -33.443°; Table S2, Figure 2c) to produce reciprocal cross

hybrids, in addition to pure offspring for comparison. Offspring were housed individually and

reared under laboratory conditions (total = 138; southern = 11, northern = 54, hybrids = 73).

We genotyped adults and offspring at five microsatellite loci (Am41, Cp10, Ctde45, Ctde05,

Ctde12; Schwartz et al. 2007; McLean et al. 2014b) and confirmed paternity using a

maximum likelihood approach in CERVUS (v3.0.7; Kalinowski, Taper, & Marshall, 2007;

Marshall, Slate, Kruuk, & Pemberton, 1998). For all offspring, we induced throat colour

expression by artificially elevating testosterone levels at nine months of age in both sexes

(female = 64, male = 74; supplemental information). Females can be reliably induced to

express discrete throat morphs found in males using testosterone because throat coloration is

autosomally inherited; testosterone may influence the intensity of coloration but morphs

remain objectively classifiable (Rankin & Stuart-Fox, 2015). Dorsolateral colour patterns

remained unchanged in testosterone-treated females therefore dorsolateral phenotype was

analysed only for male offspring. See Supplemental Information for full details of breeding,

rearing, genotyping, and throat colour expression procedures.

Analysis of phenotypic data

To characterise the phenotype of wild-caught males and captive-bred F1 hybrids, we

took standardised ventral (i.e. throat) and dorsal photographs and spectral reflectance

measurements of throat colours (details in Supplementary Information). The standardised

photos were used to classify throats into existing morphs (orange, yellow, orange-yellow,

grey, or blue), and to quantify the proportion of orange and yellow coloration present on the

head and neck from dorsal photographs. We also manually quantified the number of distinct

breaks in the dorsolateral stripe near the neck by visually inspecting all photographs of the

dorsal region. Spectral reflectance measurements (300 – 700 nm) were used to estimate the

receptor quantum catch (QC), the stimulation of retinal photoreceptors, for the violet to UV

wavelength sensitive (UVS) photoreceptor because we were specifically interested in the

diagnostic UV-blue component of throat coloration.

We performed principal component analyses (PCA) to explore variation in throat

coloration variables (proportion of orange, proportion of yellow, proportion of blue) and

dorsolateral phenotype variables (proportion orange, proportion of yellow, breaks in the

dorsolateral stripe) in the R package *FactoMineR* (v1.4.1; Le, Josse, & Husson, 2008). In

each, the lineages were differentiated along the first dimension (PC1) and this was used to

collapse variables into a single measure of overall phenotype and retained for use in

subsequent cline analyses. The quantum catch of the UV sensitive photoreceptor (QC_{UVS}) was

used singly in cline analyses to assess its frequency across the contact zone independent of

throat colour morph. Due to missing data, 204 individuals were used in throat phenotype

analyses whereas 233 individuals were used in dorsolateral phenotype analyses. Linear

discriminant analyses were performed on pure individuals in the R package *MASS* (v7.3.50;

Ripley, 2002) using all throat phenotype variables (proportion of orange, yellow, and blue,

and QC_{UVS}) and dorsolateral phenotype variables as independent variables.

Geographic cline analysis

We used the R package *HZAR* (v0.2-5, Derryberry, Derryberry, Maley, & Brumfield,

2014) to fit a series of equilibrium geographic cline models (Gay, Crochet, Bell, &

Lenormand, 2008; Szymura & Barton, 1986; Szymura & Barton, 1991) to both the molecular

and phenotypic datasets, and in all cases using 1,000,000 MCMC generations following an

initial burn-in of 100,000 generations. Specifically, we estimated cline models for the hybrid

index (H_{Index}; Bayesian *q*-value based on 1,333 loci), mtDNA haplotype frequencies, quantum

catch of the UV wavelength sensitive receptor (QC_{UVS}), PC1 for throat coloration (T_{PC1}), and

PC1 for dorsolateral phenotype (D_{PC1}).

HZAR uses logistic regression equations to fit sigmoidal clines in a maximum-

likelihood framework and describes cline centre (*c*) as the position where estimated trait

frequency is 0.5 and width (*w*) as 1/slope at the cline centre. For molecular clines, we

compared 15 models that varied in the fitting of trait interval (*p*; minimum and maximum

frequencies at tails; fixed at 0 and 1, observed values, estimated values) and exponential

decay tails (none, right only, left only, mirrored, both estimated separately). These 15 models

were compared to an additional null model. For phenotypic trait data we compared models

that varied in the fitting of exponential decay tails (none, right only, left only, mirrored, both

estimated separately) and estimated mean and variance on the ends and in the centre. All

models estimated cline centre and width as well as tail parameters if applicable (δ, distance

from the centre to the tail; τ, tail slope). QC_{UVS} and T_{PC1} were normalised to values between 0

and 1 for plotting purposes.

To address model selection uncertainty, we conducted model averaging in an Akaike's

information criterion (AIC) framework. We calculated the Akaike weight (AIC*w*), a value

between 0 and 1 which can be interpreted as the probability of being the best approximating

model (Burnham & Anderson, 2002). Models were then ranked by AICw downwards until the

cumulative AICw reached 0.95 to attain a 95% confidence set of best-ranked models (Lukacs,

Burnham, & Anderson, 2010; Symonds & Moussalli, 2011). We recalculated the AICw of

models in the 95% confidence set and used those to calculate the model-averaged estimate of

parameters (e.g. centre and width). We produced model-averaged clines by first multiplying

the estimate for a given distance interval by the AICw. This was done for each model within

the 95% confident set of models. The AIC model averaged estimate was then obtained by

summing up weighted estimates for each distance interval. Similarly, we multiplied the

minimum and maximum estimate bounds by the AICw, followed by summation to represent

the credible region. Finally, the coincidence of cline centres was assessed by comparing

model-averaged maximum-likelihood derived CIs (two log-likelihood unit support limits) of

cline centres, with non-overlapping CIs considered statistically supported differences. Akaike

weights were calculated using the R package *MuMIn* (v1.43.6; Bartoń 2018).

Results

Population structure and admixture

We found distinct population genetic structure within *C. modestus* and *C. decresii*, and

evidence for the presence of hybrids in the contact zone. Based on the STRUCTURE analysis,

only two clusters were identified, corresponding to the major split between *C. modestus* and

C. decresii (Figure 2d, Table S3). The *q*-values ranged from 0.999 (pure southern) to 0.001

(pure northern). Across the contact zone transect, individuals from the southernmost sites 1 –

2 had a high probability of belonging to the *C. decresii* genetic cluster ($q \geq 0.90$; mean ± SD =

0.99 ± 0.02) with the exception of two admixed individuals ($q = 0.81$ and 0.57). All

individuals from the northernmost sites 9 – 14 had a high probability of belonging to the *C.*

modestus genetic cluster ($q \leq 0.10$; mean $\pm$ SD $= 0.04 \pm 0.03$). Hybrids were identified within sites 3 – 8 where all individuals were of mixed ancestry and exhibited a highly skewed continuum of *q*-values ranging from 0.53 to 0.11. These sites will hereinafter be referred to as 'the contact zone', encompassing a distance of approximately 22.5 km. We also identified a density trough of approximately 12 km between sites 2 and 3 (Figures 2c–d) across which the genotype changes abruptly and multi-year surveys recovered no individuals. With the exception of two individuals at site 3, all admixed individuals in sites 3 – 13 had *C. modestus* mtDNA haplotypes. The *C. decresii* mtDNA haplotype was fixed in sites 1 – 2, including the only two admixed individuals identified in the geographic range of *C. decresii* (Figure S3).

The first axis of the PCoA constructed from 1,333 loci clustered all populations into two major groups corresponding to *C. modestus* and *C. decresii* (variance explained = 19.4%; Figure 3). We found evidence for the presence of hybrids along this axis, with hybrids clustered more closely to the Olary Ranges (OR) population of *C. modestus*. Population pairwise measures of F_{ST} between the contact zone and *C. modestus* OR/SFR/NFR populations (F_{ST} range: 0.058 – 0.216; Table S4) were lower relative to measures of F_{ST} between the contact zone and *C. decresii* MS/KI populations (F_{ST} range: 0.203 – 0.293). The contact zone had the lowest inbreeding coefficient (F = 0.095) and the highest level of heterozygosity (H_{exp} = 0.084, H_{obs} = 0.079) relative to parental populations, with very similar allelic number and richness to the OR population (Table 1). The second PCoA axis (variance explained = 3.5%; Figure 3) showed population structure within each species, with the Northern Flinders Ranges (NFR) and Kangaroo Island (KI) populations showing differentiation from other populations of their respective species, and from each other (F_{ST} = 0.571; Table S4). This can likely be attributed to the fact that each represents the northernmost and southernmost populations of their species, and KI is an insular population.

Hybrid classification

We found a strong bias in genetic introgression with admixed individuals being

predominantly backcrossed with *C. modestus*. Of individuals within the contact zone, 35%

had intermediate *q*-values (0.4 < *q* < 0.6) and a large proportion (65%) had *q*-values between

0.39 and 0.10 suggestive of backcrossing. NewHybrids analysis classified all individuals of *C.*

modestus and *C. decresii* (0.1 > *q* > 0.9) as pure parental forms (PP ≥ 99.1%). Further, 59 of

79 individuals in the contact zone were classified to one of the six classes with a posterior

probability (PP) of ≥ 95%. Additionally, 6 individuals were classified as pure *C. modestus*

(P1), 36 as F2 hybrids, and 19 as backcrosses to the northern lineage (B1). We did not recover

any F1 hybrids or backcrosses to *C. decresii*, posterior probabilities for these two categories

were uniformly low (average PP = 0.0%). The two admixed individuals found in the

geographic range of *C. decresii* were ambiguously classified as an F2 (PP = 54.2%) and a

backcross to *C. decresii* (PP = 80.8%). The 22 individuals unable to be confidently classified

(PP ≤ 95%) likely represent the continuum present in wild populations, unsuited for discrete

classification, and difficult to distinguish even with many diagnostic markers (Boecklen &

Howard, 1997; Vähä & Primmer, 2006).

Population history

The δaδi analysis showed that the model of secondary contact with ongoing

asymmetrical migration ("sec_contact_asym_mig") was more likely than the model with

subsequent isolation ("sec_contact_asym_mig_three_epoch"), based on log-likelihood and

AIC (Figure S4, Table S5). To aid interpretation, we converted parameter estimates into

biologically meaningful values (Figure 4; Figure S2). For the top model, the total time since

divergence of *C. decresii* and *C. modestus* was ~1.95 million years (T1+T2), with secondary

contact occurring ~170,000 years ago (T2). The 'population' size of *C. modestus* (Nu2) was

approximately 4 times larger than *C. decresii* (Nu1), and migration was asymmetrical with

0.36 individuals from *C. modestus* to *C. decresii* per generation compared to 1.55 individuals

from *C. decresii* to *C. modestus* per generation (Figure 4).

Hybrid throat and dorsolateral phenotypes

The throat phenotype of wild-caught hybrid males was more similar to that of *C.*

modestus, exhibiting all four throat colour morphs with uniformly low levels of UV

reflectance. Linear discriminant analyses (LDA) on throat variables showed *C. modestus* and

C. decresii strongly differentiated along LD1 which was driven primarily by the proportion of

blue and secondarily by QC_{UVS} and the proportion of orange (variance explained = 99.8%;

Figure 5a, Table S6). The 95% confidence ellipses drawn for the throat phenotypes of *C.*

modestus and hybrid males overlie each other. The large majority of hybrid throats cluster

with *C. modestus* throats. Blue coloration was not detected on the throats of pure *C. modestus*

males. A degree of blue coloration ($\geq$ 5% blue pixels in segmentation analysis), however, was

found on a portion of hybrid throats of each morph: grey (47.4%, N = 19), orange (5.2%, N =

39), orange-yellow (6.7%, N = 15), yellow (11.8%, N = 17; Figure 6a). The throat phenotype

of the two hybrid individuals found in the range of *C. decresii* had larger proportions of blue

coloration on the throat relative to other hybrids.

In contrast to wild-caught hybrid throats, captive-bred F1 hybrid throat phenotype was

more intermediate and exhibited a wide range of UV reflectance levels varying from those

characteristic of *C. modestus* to *C. decresii*. Linear discriminant analysis on throat variables of

captive-bred F1 hybrids showed strong differentiation between lineages along LD1 driven

primarily by QC_{UVS} and secondarily by the proportions of blue and yellow (variance

explained = 88.0%; Figure 5b, Table S6). The 95% confidence ellipse drawn for hybrid

throats overlaps those of both *C. modestus* and *C. decresii* throats to a similar extent. Results

were essentially unchanged when considering hybrids sired by a *C. modestus* or *C. decresii*

male and by offspring sex. Similar to wild hybrids, all four *C. modestus* throat colour morphs

were recovered in captive-bred hybrids, and all but the orange-yellow morph had a proportion

of individuals with a blue component: grey (32.3%, N = 31), orange (18.5%, N = 27),

orange-yellow (0%, N = 2), yellow (8.3%, N = 12; Figure 6b). Although the artificial

elevation of testosterone levels has been shown to induce the expression of *C. modestus* throat

colour morphs in juveniles and females (Rankin et al., 2016; Rankin & Stuart-Fox, 2015), we

found that it did not comparably induce the expression of the blue throat colour morph in all

pure *C. decresii* offspring. Therefore, the presence of blue coloration on F1 hybrid offspring

throats may have been underestimated.

Dorsolateral phenotype in both wild-caught and captive-bred hybrids is intermediate

between the lineages (Figure 5c–d). Within the contact zone, differentiation between lineages

along LD1 is driven primarily by the number of breaks in the dorsolateral stripe and

proportion of orange (Table S6). The 95% confidence ellipses drawn for the dorsolateral

phenotypes of hybrids and *C. modestus* males overlap incompletely, likely reflecting the high

frequencies of backcrossing to *C. modestus*. In captive-bred hybrids, comparable results were

found with differentiation driven by breaks in the lateral stripe (Table S6) and the 95%

confidence ellipse for hybrid dorsolateral phenotype overlaps those of both *C. modestus* and

C. decresii dorsolateral phenotypes.

Geographic clines

Model-averaged clines for mtDNA and throat phenotype (QC_{UVS} and T_{PC1}) exhibited

relatively narrower clines with abrupt transitions suggesting selection on these traits,

compared to clines for the hybrid index (H_{Index}) and dorsolateral phenotype (D_{PC1}; Table 2).

The best-fitting clines for the hybrid index and mtDNA had coincident centres with

overlapping CIs but showed discordance in slope where the mtDNA cline was significantly

narrower (H_{Index}: $c = 28.41$ km, $w = 29.77$ km; mtDNA: $c = 27.78$ km, $w = 0.64$ km; Figure

8a–b). The cline fitted for the dorsolateral phenotype ($c = 23.02$ km, $w = 13.97$) was

coincident with that of the hybrid index (Figure 8c). Clines for throat phenotype (QC_{UVS} and

T_{PC1}) were similar in centre and width to each other (QC_{UVS}: $c = 15.90$ km, $w = 8.48$ km;

T_{PC1}: $c = 15.38$ km, $w = 3.22$ km; Figure 8d), their centres were displaced by 12.51 and 13.03

km to the south from the hybrid index cline centre and do not have overlapping CIs with the

hybrid index cline. See Table S7 for full details of cline models in the AIC 95% confidence

set for each trait.

Discussion

Evaluating the evolutionary outcomes of secondary contact between lineages that

differ in morph composition can provide valuable insights into the role of polymorphism in

speciation. We characterised genomic and phenotypic patterns across the contact zone

between the colour polymorphic *C. modestus* and monomorphic *C. decresii* and concurrently

measured the phenotype of captive-bred F1 hybrids. We found that the contact zone is

geographically narrow relative to the parental ranges (approx. 30 km) with no parental forms

or F1 hybrids present in the genetic centre, suggesting that hybridisation is neither recent nor

ongoing. Genomic introgression was asymmetric, with hybrids backcrossing almost

exclusively to *C. modestus*. Formal comparison of two plausible models of population history

for this system supports a prolonged period of secondary contact with asymmetrical

migration. This suggests that despite potential for admixture throughout the last interglacial

period, hybridisation between the species is not ongoing. Dorsolateral phenotype across the

contact zone was consistent with patterns of genetic admixture, whereas the cline for throat

phenotype was displaced relative to clines for the hybrid index and dorsolateral phenotype.

Throat phenotype of wild-caught hybrids resembled that of *C. modestus*, with all four *C.*

modestus throat morphs present and uniformly low UV reflectance. By contrast, both

dorsolateral and throat phenotype were intermediate in captive-bred F1 hybrids. Taken

together, these data are consistent with selection on throat phenotype across the contact zone.

The pattern of asymmetric genomic introgression, with limited backcrossing to the

southern lineage, could reflect multiple evolutionary and demographic scenarios. First,

extrinsic barriers may hinder dispersal and migration, in turn preventing backcrossing.

Second, the pattern of asymmetric genomic introgression coupled with the prevalence of *C.*

modestus mtDNA could potentially be explained by demographic circumstances. For

instance, a greater abundance of *C. modestus* individuals during initial periods of

hybridisation with continued asymmetric migration into the contact zone, or the sex-biased

dispersal of *C. decresii* males to the north, could both result in asymmetric introgression.

Third, intrinsic genomic incompatibilities could limit backcrossing via reinforcement (i.e.

selection against maladaptive hybridsation). We discuss each explanation below in light of

our data and available evidence.

The restricted gene flow to *C. decresii* is associated with an approximately 12 km

density trough (Figure 3b–c). In this region, it appears that hybrid and/or pure populations

either occur in levels below human detection or do not currently exist despite continuous

rocky habitat with no obvious barriers to dispersal (e.g. major roads, waterways, human

structures; Figure S5). The density trough was surveyed thoroughly in 2016 and 2017 and

briefly in 2013. The presence of suitable habitat was confirmed by surveys on foot, satellite

imagery (Figure S5), and/or flying drone surveys; this area is unlikely to constitute an

ecological barrier. Furthermore, a model with prolonged secondary contact and ongoing

asymmetric migration better fits the data than one with subsequent isolation. Secondary

contact over at least the last interglacial period suggests a sufficient period for dispersal across

the 12 km density trough in the absence of extrinsic barriers. Although dispersal has not been

examined in *C. decresii*, dispersal estimates in other lizard species range from 0.01 km/gen to

0.1 km/gen for Dactyloidae and Lacertidae (Calsbeek, Duryea, Parker, & Cox, 2014;

reviewed in McEntee et al., 2020; Vercken, Sinervo, & Clobert, 2012). With a generation

time of two years, dispersal across this distance is consistent with the estimated age of the

contact zone. In general, the width of the genetic contact zone (approx. 30 km), even taking

into consideration the 12 km density trough, is relatively narrow relative to the parental

ranges (Figure 2b). This distance is in accordance with cline widths estimated from genetic

data in other lizards: Teiidae (3.2 – 93.9 km; Dessauer, Cole, & Townsend, 2000; Gifford,

2008), Dactyloidae (3 – 5 km; Case & Williams, 1984; Johansson, Surget-Groba, & Thorpe,

2008), Lacertidae (2.7 – 29.9 km; Miraldo, Faria, Hewitt, Paulo, & Emerson, 2013),

Phrynosomatidae (0.8 – 30.9 km; Leaché & Cole, 2007; Leaché, Grummer, Harris, &

Breckheimer, 2017; J. C. Marshall & Sites Jr., 2001; Sites, Barton, & Reed, 1995).

Demographic factors alone are similarly unlikely to explain the observed asymmetric

introgression. The greater abundance and migration of *C. modestus* individuals into the

contact zone should result in a greater proportion of *C. modestus* genomic background (*C.*

decresii genomic signature quickly eroded by *C. modestus* contribution). However, we found

close to equal genomic proportions in many hybrids in the genetic centre. Despite opportunity

for ongoing migration, albeit asymmetrical, hybrids backcrossed to *C. decresii* were almost

entirely absent. Male-biased dispersal could underlie population genetic structure consistent

with our results (Jockusch & Wake, 2002; Petit & Excoffier, 2009); however, it is unlikely

that male-biased dispersal would exclusively be observed in *C. decresii* and not in *C.*

modestus given their polygynous mating systems (reviewed in Li & Kokko, 2019).

The absence of extrinsic barriers coupled with the narrower and more abrupt transition

exhibited by the mtDNA cline relative to the hybrid index, and the rarity of admixed

individuals containing the *C. decresii* mtDNA haplotype, point to a role for intrinsic genomic

incompatibilities (e.g. Helbig, Salomon, Bensch, & Seibold, 2001; Miller, Lipshutz, Smith, &

Bermingham, 2014). Captive breeding experiments between *C. modestus* and *C. decresii*

show no significant prezygotic barriers to the formation of F1 hybrids with either parental

mtDNA haplotype (Dong, Rankin, McLean, & Stuart-Fox, unpublished data), although their

reproductive viability is unknown. However, stronger incompatibilities and hybrid breakdown

are more frequently seen in F2 and later generation hybrids during the early stages of

divergence. This has been described extensively for both Dobzhansky-Muller

incompatibilities (reviewed in Coyne & Orr, 2004) and for cytonuclear incompatibilities

(Ellison & Burton, 2008; Fishman & Willis, 2006; Meiklejohn et al., 2013; Niehusis, Judson,

& Gadau, 2008). Overall, our results support the hypothesis that intrinsic factors contribute to

limiting gene flow, and backcrossing to *C. decresii*, regardless of mtDNA haplotype, may

result in offspring that are inviable, infertile, or suffer reduced fitness.

Phenotypic patterns recovered from within the contact zone could theoretically result

from selection on the throat colour signal or neutral processes. We found that both throat

phenotype clines were displaced relative to the genetic cline, in contrast to the dorsolateral

phenotype cline. The displaced throat phenotype clines and prevalence of the *C. modestus*

throat phenotype in the contact zone could theoretically arise from genetic dominance. In this

scenario, captive-bred F1 hybrids would be expected to have largely *C. modestus* throat

phenotypes which contrasts with the observed intermediate phenotypes. Another possibility is

that this phenotypic pattern is due to extensive backcrossing to *C. modestus*. While some wild

hybrids may have northern throat phenotypes because they are heavily backcrossed to *C.*

modestus, the prevalence of *C. modestus* phenotype in hybrids with close to equal proportions

of *C. modestus* and *C. decresii* genomic backgrounds points towards selection on throat

colour.

We did not detect the loss of any *C. modestus* throat morphs across the contact zone or

in F1 hybrids, indicating that co-adapted trait complexes of the *C. modestus* polymorphism

are stable despite genetic disruption. The maintenance of all four morphs in all populations of

C. modestus despite environmental selection on morph frequencies suggests that negative

frequency dependent selection is a strong evolutionary force counteracting loss of

polymorphism due to drift (McLean, Stuart-Fox, & Moussalli, 2015). Although hybrid throats

more closely resembled *C. modestus* throat morphs, we observed a proportion of each morph

which exhibited a degree of intermediate nature where an area of blue coloration was present.

There was a notably higher proportion of grey morphs which had blue coloration present.

This may be due to the similar mechanism by which grey and blue skin are produced in *C.*

modestus and *C. decresii* (McLean, Lutz, Rankin, Stuart-Fox, & Moussalli, 2017).

Nevertheless, we did not detect any hybrid throats that exhibited a true *C. decresii* throat

phenotype with both predominantly blue throat coloration and a high level of UV reflectance.

This, in addition to the variable presence of UV reflectance in captive-bred F1 hybrids, points

toward the uncoupling of blue coloration and UV reflectance in hybrids. Hybridisation can

generate novel phenotypes (e.g. blue without UV); for example, admixture between treefrog

lineages which have red and red/blue legs result in purple legs which are produced from the

directional introgression of colour loci inherited from parental lineages (Akopyan et al.,

2020).

Selection on throat coloration could be driven by mating behaviour which evolved via

reinforcement and/or microhabitat characteristics. Mate preference trials with captive *C.*

modestus and *C. decresii* predict a greater frequency of mating between *C. decresii* females

with *C. modestus* males than the reciprocal combination(McLean, Bartle, Dong, Rankin, &

Stuart-Fox, 2020). This contrasts with patterns observed within the contact zone, indicating

that prezygotic barriers between species have not evolved in allopatry. However, postzygotic

barriers (inferred from low frequency of hybrids backcrossed to the south) should select for

the formation of prezygotic barriers to prevent maladaptive hybridisation (i.e. reinforcement).

Extrinsic factors could also contribute to the *C. modestus* throat phenotype having a

competitive advantage in the contact zone, for example, due to a relatively higher

conspicuousness to conspecifics. The throat coloration of each species is more conspicuous

against the predominant background colours of native lichen in their respective ranges

(McLean, Moussalli, & Stuart-Fox, 2014). Within the contact zone, lichen coloration is

similar to that found in the range of *C. modestus*, predominantly pale green. Therefore, sexual

selection may operate through male-male interactions or female choice to result in a

reproductive advantage of the *C. modestus* throat phenotype as a more conspicuous and

detectable visual signal (reviewed in Servedio & Boughman, 2017). Similarly, the greater

overlap of hybrid phenotype (e.g. ventral coloration) with one parental species was associated

with environmental variation in *Hyperolius* frogs (Bell & Irian, 2019). Selection has been

shown to drive the rapid spread of an advantageous trait/phenotype across contact zones, and

in some cases into the other parental lineage (Baldassarre, White, Karubian, & Webster, 2014;

Pardo-Diaz et al., 2012; Stein & Uy, 2006; While et al., 2015). In contrast, the dorsolateral

phenotype cline indicates that it is not under similarly strong selection and appears to play a

lesser role in reproductive isolation in sympatry. Male dorsolateral coloration may be

constrained as it is under both sexual and natural selection because of its dual function as a

intraspecific signal and in minimising conspicuousness to predators against native lichen

(McLean, Moussalli, et al., 2014; Ramos & Peters, 2016).

Conclusions

In summary, our study suggests that sexual signal divergence and intrinsic genetic

incompatibilities play an important role in restricting gene flow between the polymorphic *C.*

modestus and monomorphic *C. decresii*. The species remain distinct despite continued

migration and opportunity for gene flow. The stability of the *C. modestus* polymorphism

despite genetic disruption is consistent with the theoretical genetic architecture of colour

polymorphism and correlated traits, namely control by pleiotropic or tightly linked genes.

Taken together, these results lend support to the view that polymorphism can promote

speciation.

Acknowledgements

This study was funded by the Australian Research Council (DP150101048 and DP1092908)

to D.S-F., Holsworth Wildlife Endowment and The David Lachlan Hay Memorial Fund to

C.M.D., and Holsworth Wildlife Endowment, Alfred Nicholas Fellowship and Nature

Foundation South Australia to C.A.M. We thank Madeleine Yewers, Steven Mesis, Katrina

Rankin and Anne Auslebrook for assistance; private landowners for kindly allowing land

access; and South Australian Museum for providing tissue samples. Data collection was

conducted with permits from the Department of Environment, Water and Natural Resources,

South Australia (permit nos. E25861-1 and Q26428-3), the Department of Environment,

Land, Water and Planning, Victoria (permit nos. 10007000 and 10007751) and with approval

by The University of Melbourne Animals Ethics Committee (approval nos. 1011760.1 and

1413220.4), the Wildlife Ethics Committee of South Australia (approval nos. 18/2010 and

25/2015). We thank Ben Phillips and three anonymous reviewers for insightful comments,

which helped to improve the manuscript.

618
References

Akopyan, M., Gompert, Z., Klonoski, K., Vega, A., Kaiser, K., Mackelprang, R., …
Robertson, J. M. (2020). Genetic and phenotypic evidence of a contact zone between
divergent colour morphs of the iconic red-eyed treefrog. *Molecular Ecology*, 1–15. doi:
10.1111/mec.15639

Anderson, E. C., & Thompson, E. A. (2002). A model-based method for identifying species
hybrids using multilocus genetic data. *Genetics*, *160*(3), 1217–1229.

Baldassarre, D. T., White, T. A., Karubian, J., & Webster, M. S. (2014). Genomic and
morphological analysis of a semipermeable avian hybrid zone suggests asymmetrical
introgression of a sexual signal. *Evolution*, *68*(9), 2644–2657. doi: 10.1111/evo.12457

Bartoń, K. (2018). *MuMIn: Multi-Model Inference. R package version 1.42.1.*
https://CRAN.R-project.org/package=MuMIn.

Bell, R. C., & Irian, C. G. (2019). Phenotypic and genetic divergence in reed frogs across a
mosaic hybrid zone on São Tomé Island. *Biological Journal of the Linnean Society*,
128(3), 672–680. doi: 10.1093/biolinnean/blz131

Boecklen, W. J., & Howard, D. J. (1997). Genetic analysis of hybrid zones: numbers of
markers and power of resolution. *Ecology*, *78*(8), 2611–2616.

Boughman, J. W. (2001). Divergent sexual selection enhances reproductive isolation in
sticklebacks. *Nature*, *411*(6840), 944–948. doi: 10.1038/35082064

Burnham, K., & Anderson, D. (2002). *Model selection and multimodel inference* (2nd ed.).
New York: Springer.

Burton, R. S., & Barreto, F. S. (2012). A disproportionate role for mtDNA in Dobzhansky-
Muller incompatibilities? *Molecular Ecology*, *21*(20), 4942–4957. doi:
10.1111/mec.12006

Byrne, M. (2008). Evidence for multiple refugia at different time scales during Pleistocene
climatic oscillations in southern Australia inferred from phylogeography. *Quaternary*
Science Reviews, *27*, 2576–2585. doi: 10.1016/j.quascirev.2008.08.032

Calsbeek, R., Duryea, M. C., Parker, E., & Cox, R. M. (2014). Sex-biased juvenile dispersal
is adaptive but does not create genetic structure in island lizards. *Behavioral Ecology*,
25(5), 1157–1163. doi: 10.1093/beheco/aru102

Case, S. M., & Williams, E. E. (1984). Study of a Contact Zone in the Anolis distichus
Complex in the Central Dominican Republic. *Herpetologica*, *40*(2), 118–137.

Clement, M., Posada, D., & Crandall, K. A. (2000). TCS: a computer program to estimate
gene genealogies. *Molecular Ecology*, *9*(10), 1657–1659. doi: 10.1046/j.1365-

294x.2000.01020.x

Corl, A., Davis, A. R., Kuchta, S. R., & Sinervo, B. (2010). Selective loss of polymorphic
   mating types is associated with rapid phenotypic evolution during morphic speciation.
   Proceedings of the National Academy of Sciences, *107*(9), 4254–4259. doi:
   10.1073/pnas.0909480107

Coyne, J. A., & Orr, H. A. (2004). Speciation. In *Sinauer Associates*. Sunderland,
   Massachusetts: Sinauer Associates.

Derryberry, E. P., Derryberry, G. E., Maley, J. M., & Brumfield, R. T. (2014). HZAR: hybrid
   zone analysis using an R software package. *Molecular Ecology Resources*, *14*(3), 652–
   663. doi: 10.1111/1755-0998.12209

Dessauer, H. C., Cole, C. J., & Townsend, C. R. (2000). Hybridization among western
   whiptail lizards (Cnemidophorus tigris) in Southwestern New Mexico: Population
   genetics, morphology, and ecology in three contact zones. *Bulletin of the American
   Museum of Natural History*, (246), 1–127.

Dobzhansky, T. (1937). Genetics the Origin and of Species. In *Genetics and the origin of
   species*. New York: Columbia University Press.

Dong, C. M., Johnston, G. R., Stuart-Fox, D., Moussalli, A., Rankin, K. J., & McLean, C. A.
   (2021). Elevation of divergent color polymorphic and monomorphic lizard lineages
   (Squamata: Agamidae) to species level. *Ichthyology and Herpetology*, *109*(1). doi:
   10.1643/h2020064

Earl, D. A., & VonHoldt, B. M. (2012). STRUCTURE HARVESTER: a website and program
   for visualizing STRUCTURE output and implementing the Evanno mothod.
   Conservation Genetics Resources, *4*(2), 359–360. doi: 10.1007/s12686-011-9548-7

Ellison, C. K., & Burton, R. S. (2008). Interpopulation hybrid breakdown maps to the
   mitochondrial genome. *Evolution*, *62*(3), 631–638. doi: 10.1111/j.1558-
   5646.2007.00305.x

Fishman, L., & Willis, J. H. (2006). A cytonuclear incompatibility causes anther sterility in
   Mimulus hybrids. *Evolution*, *60*(7), 1372–1381. doi: 10.1554/05-708.1

Forsman, A., Ahnesjö, J., Caesar, S., & Karlsson, M. (2008). A model of ecological and
   evolutionary consequences of color polymorphism. *Ecology*, *89*(1), 34–40. doi:
   10.1890/07-0572.1

Forstner, M. R., Davis, S. K., & Arèvalo, E. (1995). Support for the hypothesis of anguimorph
   ancestry for the suborder Serpentes from phylogenetic analysis of mitochondrial DNA
   sequences. *Molecular Phylogenetics and Evolution*, *4*(1), 93–102. doi:
   10.1006/mpev.1995.1010

Gay, L., Crochet, P. A., Bell, D. A., & Lenormand, T. (2008). Comparing clines on molecular
   and phenotypic traits in hybrid zones: A window on tension zone models. *Evolution*,
   62(11), 2789–2806. doi: 10.1111/j.1558-5646.2008.00491.x

Georges, A., Li, Q., Lian, J., O'Meally, D., Deakin, J., Wang, Z., … Zhang, G. (2015). High-
   coverage sequencing and annotated assembly of the genome of the Australian dragon
   lizard *Pogona vitticeps*. *GigaScience*, *4*(1), 45. doi: 10.1186/s13742-015-0085-2

Gibbons, J. R. H. (1977). *Comparative ecology and behaviour of lizards of the* Amphibolurus
decresii *species complex*. The University of Adelaide.

Gibbons, J. R. H. (1979). The hind leg pushup display of the *Amphibolurus decresii* species
complex (Lacertilia: Agamidae). *Copeia*, *1*, 29–40. doi: 10.2307/1443725

Gifford, M. E. (2008). Divergent character clines across a recent secondary contact zone in a
Hispaniolan lizard. *Journal of Zoology*, *274*(3), 292–300. doi: 10.1111/j.1469-
7998.2007.00385.x

Gosselin, T., Lamothe, M., Devloo-Delva, F., & Grewe, P. (2020). *radiator: RADseq Data*
Exploration, Manipulation and Visualization using R. doi: 10.5281/zenodo.3687060

Gray, S. M., & McKinnon, J. S. (2007). Linking color polymorphism maintenance and
speciation. *Trends in Ecology & Evolution*, *22*(2), 71–79. doi:
10.1016/j.tree.2006.10.005

Gruber, B., Unmack, P. J., Berry, O. F., & Georges, A. (2018). dartR: An R package to
facilitate analysis of SNP data generated from reduced representation genome
sequencing. *Molecular Ecology Resources*, *18*, 691–699. doi: 10.1111/1755-0998.12745

Gutenkunst, R., Hernandez, R., Williamson, S., & Bustamante, C. (2009). Inferring the joint
demographic history of multiple populations from multidimensional SNP frequency data.
PLoS Genetics, *5*(10). doi: 10.1371/journal.pgen.1000695

Gutenkunst, R., Hernandez, R., Williamson, S., & Bustamante, C. (2010). Diffusion
approximations for demographic inference: DaDi. *Nature Precedings*. doi:
10.1038/npre.2010.4594.1

Hacking, J. D., Stuart-Fox, D., Godfrey, S. S., & Gardner, M. G. (2018). Specific MHC class
I supertype associated with parasite infection and color morph in a wild lizard
population. *Ecology and Evolution*, *8*(19), 9920–9933. doi: 10.1002/ece3.4479

Helbig, A. J., Salomon, M., Bensch, S., & Seibold, I. (2001). Male-biased gene flow across an
avian hybrid zone: Evidence from mitochondrial and microsatellite DNA. *Journal of*
Evolutionary Biology, *14*(2), 277–287. doi: 10.1046/j.1420-9101.2001.00273.x

Hill, G. E. (2015). Mitonuclear ecology. *Molecular Biology and Evolution*, *32*(8), 1917–1927.
doi: 10.1093/molbev/msv104

Hooper, D. M., Griffith, S. C., & Price, T. D. (2019). Sex chromosome inversions enforce
reproductive isolation across an avian hybrid zone. *Molecular Ecology*, *28*(6), 1246–
1262. doi: 10.1111/mec.14874

Houston, T. F. (1974). Revision of the *Amphibolurus decresii* complex (Lacertilia: Agamidae)
of South Australia. *Transactions of The Royal Society of South Australia*, *98*(2), 49–60.
doi: 10.1126/science.97.2526.482-a

Hugall, A. F., & Stuart-Fox, D. (2012). Accelerated speciation in colour-polymorphic birds.
Nature, *485*, 631–634. doi: 10.1038/nature11050

Huxley, J. (1955). Morphism in birds. *Acta International Congress Ornithology XI*, 309–328.

Jakobsson, M., & Rosenberg, N. A. (2007). CLUMPP: A cluster matching and permutation

program for dealing with label switching and multimodality in analysis of population
structure. *Bioinformatics*, *23*(14), 1801–1806. doi: 10.1093/bioinformatics/btm233

Jiggins, C. D., Naisbit, R. E., Coe, R. L., & Mallet, J. (2001). Reproductive isolation caused
by colour pattern mimicry. *Nature*, *411*(6835), 302–305. doi: 10.1038/35077075

Jockusch, E. L., & Wake, D. B. (2002). Falling apart and merging: Diversification of slender
salamanders (Plethodontidae: Batrachoseps) in the American West. *Biological Journal of*
the Linnean Society, *76*(3), 361–391. doi: 10.1046/j.1095-8312.2002.00071.x

Johansson, H., Surget-Groba, Y., & Thorpe, R. S. (2008). The roles of allopatric divergence
and natural selection in quantitative trait variation across a secondary contact zone in the
lizard Anolis roquet. *Molecular Ecology*, *17*(23), 5146–5156. doi: 10.1111/j.1365-
294X.2008.03979.x

Jombart, T., & Ahmed, I. (2011). adegenet 1.3-1: New tools for the analysis of genome-wide
SNP data. *Bioinformatics*, *27*(21), 3070–3071. doi: 10.1093/bioinformatics/btr521

Kalinowski, S. T., Taper, M. L., & Marshall, T. C. (2007). Revising how the computer
program CERVUS accommodates genotyping error increases success in paternity
assignment. *Molecular Ecology*, *16*(5), 1099–1106. doi: 10.1111/j.1365-
294X.2007.03089.x

Kearse, M., Moir, R., Wilson, A., Stones-havas, S., Sturrock, S., Buxton, S., … Drummond,
A. (2012). Geneious Basic: An integrated and extendable desktop software platform for
the organization and analysis of sequence data. *Bioinformatics*, *28*(12), 1647–1649. doi:
10.1093/bioinformatics/bts199

Kilian, A., Wenzl, P., Huttner, E., Carling, J., Xia, L., Caig, V., … Uszynski, G. (2012).
Diversity Arrays Technology: A Generic Genome Profiling Technology on Open
Platforms. In F. Pompanon & A. Bonin (Eds.), *Data Production and Analysis in*
Population Genomics. Methods in Molecular Biology (Methods and Protocols) (Vol.
888, pp. 67–89). doi: 10.1007/978-1-61779-870-2

Lê, S., Josse, J., & Husson, F. F. (2008). FactoMineR: An R Package for Multivariate
Analysis. *Journal of Statistical Software*, *25*(1), 1–18. doi: 10.1016/j.envint.2008.06.007

Leaché, A. D., & Cole, C. J. (2007). Hybridization between multiple fence lizard lineages in
an ecotone: Locally discordant variation in mitochondrial DNA, chromosomes, and
morphology. *Molecular Ecology*, *16*(5), 1035–1054. doi: 10.1111/j.1365-
294X.2006.03194.x

Leaché, A. D., Grummer, J. A., Harris, R. B., & Breckheimer, I. K. (2017). Evidence for
concerted movement of nuclear and mitochondrial clines in a lizard hybrid zone.
Molecular Ecology, *26*(8), 2306–2316. doi: 10.1111/mec.14033

Li, X. Y., & Kokko, H. (2019). Sex-biased dispersal: a review of the theory. *Biological*
Reviews, *94*(2), 721–736. doi: 10.1111/brv.12475

Lukacs, P. M., Burnham, K. P., & Anderson, D. R. (2010). Model selection bias and
Freedman's paradox. *Annals of the Institute of Statistical Mathematics*, *62*(1), 117–125.
doi: 10.1007/s10463-009-0234-4

Marshall, J. C., & Sites Jr., J. W. (2001). A comparison of nuclear and mitochondrial cline

shapes in a hybrid zone in the Sceloporus grammicus complex (Squamata ;
Phrynosomatidae). *Molecular Ecology*, *10*, 435–449.

Marshall, T. C., Slate, J., Kruuk, L. E. B., & Pemberton, J. M. (1998). Statistical confidence
for likelihood-based paternity inference in natural populations. *Molecular Ecology*, *7*,
639–655.

Mavárez, J., Salazar, C. A., Bermingham, E., Salcedo, C., Jiggins, C. D., & Linares, M.
(2006). Speciation by hybridization in *Heliconius* butterflies. *Nature*, *441*(7095), 868–
871. doi: 10.1038/nature04738

McEntee, J. P., Burleigh, J. G., & Singhal, S. (2020). Dispersal predicts hybrid zone widths
across animal diversity: Implications for species borders under incomplete reproductive
isolation. *American Naturalist*, *196*(1), 9–28. doi: 10.1086/709109

McLean, C. A., Bartle, R. A., Dong, C. M., Rankin, K. J., & Stuart-Fox, D. (2020). Divergent
male and female mate preferences do not explain incipient speciation between lizard
lineages. *Current Zoology*, (March), 1–8. doi: 10.1093/cz/zoaa010

McLean, C. A., Lutz, A., Rankin, K. J., Stuart-Fox, D., & Moussalli, A. (2017). Revealing the
biochemical and genetic basis of color variation in a polymorphic lizard. *Molecular
Biology and Evolution*, *34*(8), 1924–1935. doi: 10.1093/molbev/msx136

McLean, C. A., Moussalli, A., Sass, S., & Stuart-Fox, D. (2013). Taxonomic assessment of
the *Ctenophorus decresii* complex (Reptilia: Agamidae) reveals a new species of dragon
lizard from western New South Wales. *Records of the Australian Museum*, *65*(3), 51–63.
doi: 10.3853/j.2201-4349.65.2013.1600

McLean, C. A., Moussalli, A., & Stuart-Fox, D. (2014). Local adaptation and divergence in
colour signal conspicuousness between monomorphic and polymorphic lineages in a
lizard. *Journal of Evolutionary Biology*, *27*(12), 2654–2664. doi: 10.1111/jeb.12521

McLean, C. A., & Stuart-Fox, D. (2014). Geographic variation in animal colour
polymorphisms and its role in speciation. *Biological Reviews*, *89*(4), 860–873. doi:
10.1111/brv.12083

McLean, C. A., Stuart-Fox, D., & Moussalli, A. (2014). Phylogeographic structure,
demographic history and morph composition in a colour polymorphic lizard. *Journal of
Evolutionary Biology*, *27*(10), 2123–2137. doi: 10.1111/jeb.12464

McLean, C. A., Stuart-Fox, D., & Moussalli, A. (2015). Environment, but not genetic
divergence, influences geographic variation in colour morph frequencies in a lizard.
BMC Evolutionary Biology, *15*(1), 156. doi: 10.1186/s12862-015-0442-x

Meiklejohn, C. D., Holmbeck, M. A., Siddiq, M. A., Abt, D. N., Rand, D. M., & Montooth,
K. L. (2013). An incompatibility between a mitochondrial tRNA and its nuclear-encoded
tRNA synthetase compromises development and fitness in *Drosophila*. *PLoS Genetics*,
9(1), e1003238. doi: 10.1371/journal.pgen.1003238

Melville, J., Haines, M. L., Boysen, K., Hodkinson, L., Kilian, A., Smith Date, K. L., …
Parris, K. M. (2017). Identifying hybridization and admixture using SNPs: application of
the DArTseq platform in phylogeographic research on vertebrates. *Royal Society Open
Science*, *4*, 161061. doi: 10.1098/rsos.161061

Miller, M. J., Lipshutz, S. E., Smith, N. G., & Bermingham, E. (2014). Genetic and
phenotypic characterization of a hybrid zone between polyandrous Northern and Wattled
Jacanas in Western Panama. *BMC Evolutionary Biology*, *14*(1), 1–14. doi:
10.1186/s12862-014-0227-7

Miraldo, A., Faria, C., Hewitt, G. M., Paulo, O. S., & Emerson, B. C. (2013). Genetic analysis
of a contact zone between two lineages of the ocellated lizard (*Lacerta lepida* Daudin
1802) in south-eastern Iberia reveal a steep and narrow hybrid zone. *Journal of*
Zoological Systematics and Evolutionary Research, *51*(1), 45–54. doi:
10.1111/jzs.12005

Muller, H. J. (1942). Isolating mechanisms, evolution, and temperature. *Biol Symp*, *6*, 71–125.

Niehuis, O., Judson, A. K., & Gadau, J. (2008). Cytonuclear genic incompatibilities cause
increased mortality in male F2 hybrids of *Nasonia giraulti* and *N. vitripennis*. *Genetics*,
178(1), 413–426. doi: 10.1534/genetics.107.080523

Osborne, L., Umbers, K. D. L., Backwell, P. R. Y., & Keogh, J. S. (2012). Male tawny
dragons use throat patterns to recognize rivals. *Naturwissenschaften*, *99*(10), 869–872.
doi: 10.1007/s00114-012-0968-3

Pardo-Diaz, C., Salazar, C., Baxter, S. W., Merot, C., Figueiredo-Ready, W., Joron, M., …
Jiggins, C. D. (2012). Adaptive introgression across species boundaries in *Heliconius*
butterflies. *PLoS Genetics*, *8*(6), e1002752. doi: 10.1371/journal.pgen.1002752

Petit, R. J., & Excoffier, L. (2009). Gene flow and species delimitation. *Trends in Ecology*
and Evolution, *24*(7), 386–393. doi: 10.1016/j.tree.2009.02.011

Portik, D. M., Leaché, A. D., Rivera, D., Barej, M. F., Burger, M., Hirschfeld, M., … Fujita,
M. K. (2017). Evaluating mechanisms of diversification in a Guineo-Congolian tropical
forest frog using demographic model selection. *Molecular Ecology*, *26*(19), 5245–5263.
doi: 10.1111/mec.14266

Pritchard, J. K., Stephens, M., & Donnelly, P. (2000). Inference of population structure using
multilocus genotype data. *Genetics*, *155*(2), 945–959. doi: 10.1111/j.1471-
8286.2007.01758.x

Pryke, S. R., & Griffith, S. C. (2009). Postzygotic genetic incompatibility between sympatric
color morphs. *Evolution*, *63*(3), 793–798. doi: 10.1111/j.1558-5646.2008.00584.x

Ramos, J. A., & Peters, R. A. (2016). Dragon wars: Movement-based signalling by Australian
agamid lizards in relation to species ecology. *Austral Ecology*, *41*, 302–315. doi:
10.1111/aec.12312

Rankin, K. J., McLean, C. A., Kemp, D. J., & Stuart-Fox, D. (2016). The genetic basis of
discrete and quantitative colour variation in the polymorphic lizard, *Ctenophorus*
decresii. *BMC Evolutionary Biology*, *16*(1), 1–14. doi: 10.1186/s12862-016-0757-2

Rankin, K. J., & Stuart-Fox, D. (2015). Testosterone-induced expression of male colour
morphs in females of the polymorphic tawny dragon lizard, *Ctenophorus decresii*. *PLOS*
ONE, *10*(10), e0140458. doi: 10.1371/journal.pone.0140458

Sætre, G. P., Moum, T., Bureš, S., Král, M., Adamjan, M., Moreno, J., … Moreno, J. (1997).
A sexually selected character displacement in flycatchers reinforces premating isolation.

Nature, *387*(June), 589–592. doi: 10.1038/42451

Schwartz, T. S., Warner, D. A., Beheregaray, L. B., Olsson, M., & Note, P. (2007). Microsatellite loci for Australian agamid lizards. *Molecular Ecology Notes*, *7*(3), 528–531. doi: 10.1111/j.1471-8286.2006.01644.x

Scott, I. A. W. W., & Keogh, J. S. (2000). Conservation genetics of the endangered grassland earless dragon *Tympanocryptis pinguicolla* (Reptilia: Agamidae) in Southeastern Australia. *Conservation Genetics*, *1*(4), 357–363. doi: 10.1023/A:1011542717349

Seehausen, O., Terai, Y., Magalhaes, I. S., Carleton, K. L., Mrosso, H. D. J., Miyagi, R., … Okada, N. (2008). Speciation through sensory drive in cichlid fish. *Nature*, *455*(7213), 620–627. doi: 10.1038/nature07285

Servedio, M. R., & Boughman, J. W. (2017). The role of sexual selection in local adaptation and speciation. *Annual Review of Ecology, Evolution, and Systematics*, *48*(1), 85–109. doi: 10.1146/annurev-ecolsys-110316-022905

Sinervo, B., & Svensson, E. (2002). Correlational selection and the evolution of genomic architecture. *Heredity*, *89*(5), 329–338. doi: 10.1038/sj.hdy.6800148

Singhal, S., & Moritz, C. (2013). Reproductive isolation between phylogeographic lineages scales with divergence. *Proceedings of the Royal Society B: Biological Sciences*, *280*, 20132246. doi: 10.1098/rspb.2013.2246

Sites, J. W., Barton, N. H., & Reed, K. M. (1995). The genetic structure of a hybrid zone between two chromosome races of the *Sceloporus grammicus* complex (Sauria, Phrynosomatidae) in central Mexico. *Evolution*, *49*(1), 9–36. doi: 10.1111/j.1558-5646.1995.tb05955.x

Stein, A. C., & Uy, J. A. C. (2006). Unidirectional introgression of a sexually selected trait across an avian hybrid zone: a role for female choice? *Evolution*, *60*(7), 1476–1485. doi: 10.1111/j.0014-3820.2006.tb01226.x

Stuart-Fox, D. M., & Johnston, G. R. (2005). Experience overrides colour in lizard contests. *Behaviour*, *142*, 329–350. doi: 10.1163/1568539053778265

Symonds, M. R. E., & Moussalli, A. (2011). A brief guide to model selection, multimodel inference and model averaging in behavioural ecology using Akaike's information criterion. *Behavioral Ecology and Sociobiology*, *65*, 13–21. doi: 10.1007/s00265-010-1037-6

Szymura, J. M., & Barton, N. H. (1986). Genetic analysis of a hybrid zone between the fire-bellied toads, *Bombina bombina* and *B . variegata*, near Cracow in Southern Poland. *Evolution*, *40*(6), 1141–1159.

Szymura, J. M., & Barton, N. H. (1991). The genetic structure of the hybrid zone between the fire-bellied toads *Bombina bombina* and *B. variegata*: comparisons between transects and between loci. *Evolution*, *45*(2), 237–261. doi: 10.1111/j.1558-5646.1991.tb04400.x

Teasdale, L. C., Stevens, M., & Stuart-Fox, D. (2013). Discrete colour polymorphism in the tawny dragon lizard (*Ctenophorus decresii*) and differences in signal conspicuousness among morphs. *Journal of Evolutionary Biology*, *26*(5), 1035–1046. doi: 10.1111/jeb.12115

Vähä, J. P., & Primmer, C. R. (2006). Efficiency of model-based Bayesian methods for
detecting hybrid individuals under different hybridization scenarios and with different
numbers of loci. *Molecular Ecology*, *15*(1), 63–72. doi: 10.1111/j.1365-
294X.2005.02773.x

Venables, W. N., & Ripley, B. D. (2002). *Modern Applied Statistics with S* (Fourth).
Retrieved from http://www.stats.ox.ac.uk/pub/MASS4

Vercken, E., Sinervo, B., & Clobert, J. (2012). The importance of a good neighborhood:
Dispersal decisions in juvenile common lizards are based on social environment.
Behavioral Ecology, *23*(5), 1059–1067. doi: 10.1093/beheco/ars075

West-Eberhard, M. J. (1986). Alternative adaptations, speciation, and phylogeny (A Review).
Proceedings of the National Academy of Sciences, *83*, 1388–1392. doi:
10.1073/pnas.83.5.1388

While, G. M., Michaelides, S., Heathcote, R. J. P., Macgregor, H. E. A., Zajac, N., Beninde,
J., … Uller, T. (2015). Sexual selection drives asymmetric introgression in wall lizards.
Ecology Letters, *18*(12), 1366–1375. doi: 10.1111/ele.12531

Wringe, B. F., Stanley, R. R. E., Jeffery, N. W., Anderson, E. C., & Bradbury, I. R. (2017).
parallelnewhybrid: an R package for the parallelization of hybrid detection using
newhybrids. *Molecular Ecology Resources*, *17*(1), 91–95. doi: 10.1111/1755-
0998.12597

Yewers, M. S. C., Jessop, T. S., & Stuart-Fox, D. (2017). Endocrine differences among colour
morphs in a lizard with alternative behavioural strategies. *Hormones and Behavior*, *93*,
118–127. doi: 10.1016/j.yhbeh.2017.05.001

Yewers, M. S. C., Pryke, S. R., & Stuart-Fox, D. (2016). Behavioural differences across
contexts may indicate morph-specific strategies in the lizard *Ctenophorus decresii*.
Animal Behaviour, *111*, 329–339. doi: 10.1016/j.anbehav.2015.10.029

Data Accessibility

The DArTseq genotype data, mtDNA alignment, R code and morphological data that support

the findings of this study are openly available in Dryad at http://doi.org[doi], reference

number [number]. All mtDNA sequences are deposited in GenBank, accession numbers

MW261595 - MW261739.

Data will be made available on Dryad upon acceptance of the manuscript.

Author Contributions

All authors contributed to study design and field sampling. C.M.D. and C.A.M. performed

laboratory work. C.M.D. conducted data analyses and drafted the manuscript. C.A.M, A.M.

and D.S-F provided input on analyses, interpretation, and the manuscript. All authors read and

approved the final manuscript.

935 **Tables**
936

937 **Table 1.** Populations sampled across the ranges of *C. modestus* (Northern Flinders Ranges
938 [NFR], Southern Flinders Ranges [SFR], Olary Ranges [OR]), *C. decresii* (Mainland South
939 [MS], and Kangaroo Island [KI]), and Contact Zone (CZ), with details of sample size for the
940 SNP dataset (*N*), expected and observed heterozygosity (H_{exp}, H_{obs}), inbreeding coefficient
941 (F), allele number (AN), and allelic richness (AR).
942

Population	*N* (♂ / ♀ / NA)	H_{exp}	H_{obs}	F	AN	AR
NFR	7 (2 / 0 / 5)	0.064	0.055	0.177	1629	0.222
SFR	41 (31 / 8 / 2)	0.076	0.070	0.129	2091	0.569
OR	60 (45 / 7 / 8)	0.072	0.060	0.177	2138	0.604
CZ	82 (69 / 13 / 0)	0.084	0.079	0.095	2134	0.601
MS	33 (26 / 4 / 3)	0.051	0.043	0.175	1714	0.286
KI	7 (6 / 1 / 0)	0.023	0.019	0.165	1429	0.072
Total	230					

†Sex was unknown for a portion of loaned museum tissue samples.

Table 2. Model-averaged estimates for cline centre and width based on the AIC 95%
confidence set models, with N representing the number of models within the AIC confident
set. Cline centre is the distance (km) from the southern-most sampling site and width (w) is
1/slope at the centre. Two log-likelihood unit support limits are shown as 'low' and 'high'.
See Table S7 for full details of cline models.
950

Trait		N	Centre	(low – high)	Width	(low – high)
Hybrid Index (H_{Index})		10	28.41	(22.37 – 33.00)	29.77	(15.05 – 55.84)
mtDNA		7	27.78	(14.59 – 28.17)	0.64	(0.08 – 21.85)
Dorsolateral Phenotype (D_{PC1})		2	23.02	(16.33 – 27.61)	13.97	(0.82 – 29.02)
Throat Phenotype	QC_{UVS}	4	15.90	(14.25 – 20.48)	8.48	(0.74 – 17.27)
	T_{PC1}	2	15.38	(15.06 – 16.61)	3.22	(2.43 – 7.94)

951
952

Figures

Figure 1. Average spectral reflectances of male throat colours found in (a) *C. modestus* measured from Southern Flinders Ranges populations: orange (O), yellow (Y), grey (G), and (b) *C. decresii* ultra-violet blue throat colour measured from Mainland South populations. Standard error shown as shaded region (N = 10 for each colour). Ultraviolet wavelengths are shown with a grey bar on the *x*-axis.

963
964

Figure 2. Geographic variation in throat colour morphs and genetic structure between *C.*
modestus and *C. decresii*. (a) Representative throats (*C. modestus*: yellow, orange-yellow,
grey and orange; *C. decresii*: blue) and dorsolateral phenotypes found in each species and the
contact zone. (b) Map showing sampling localities of individuals used in genetic analyses: *C.*
modestus (orange circles; Northern Flinders Ranges [NFR], Southern Flinders Ranges [SFR],
Olary Ranges [OR]), *C. decresii* (blue circles; Mainland South [MS], and Kangaroo Island
[KI]), and genetically admixed individuals (white circles; $0.1 < q < 0.9$). Pie charts indicate
relative frequencies of male throat colour morphs. Proportions for the Southern Flinders
Ranges and northernmost population in the Olary Ranges are adapted from McLean, Stuart-
Fox, & Moussalli, 2015. Throat colour morph data are not available for the Northern Flinders
Ranges. Elevation in meters is represented by grey shading on the scale bar. (c) An enlarged
map section showing sites across the contact zone transect used for geographic cline analyses;
(d) Results of a Bayesian analysis of ancestry in the program STRUCTURE (K = 2). Each
horizontal bar represents an individual and individuals are ordered by latitude and population.
The proportion of orange and blue represents the proportion of *C. modestus* and *C. decresii*
ancestry in each individual. A density trough between sites 2 and 3, where surveys did not
recover any individuals, is indicated in red in panels B and C.

Figure 3. Two-dimensional principal coordinate plot (PCoA) constructed from the SNP data set (1,333 loci) showing pairwise genetic distances between individuals. 95% confidence ellipses of each group are shown (*C. modestus:* Northern Flinders Ranges [NFR], Southern Flinders Ranges [SFR], Olary Ranges [OR]; Contact Zone (CZ); *C. decresii* (Mainland South [MS], and Kangaroo Island [KI]). Two outlying admixed individuals (from the 95% confidence ellipse) were those found in the geographic range of *C. decresii* (MS).

"sec_contact_asym_mig"

996 **Figure 4.** Schematic representation of the top demographic model from δaδi analysis.
997 Populations 1 and 2 are *C. decresii* and *C. modestus* (pooled with admixed individuals)
998 respectively. Converted values are provided for times (T1, T2, T3; in years), migration rates
999 (m12, m21; in individuals per generation) and population sizes (nu1 and nu2; in number of
1000 individuals).

1001

Figure 5. Biplots of linear discriminant analyses on (a, b) throat and (c, d) dorsolateral
phenotype variables for (a, c) wild-caught individuals along the contact zone transect and (b,
d) captive-bred pure and F1 hybrid offspring. 95% confidence ellipses are shown.

1008
1009
1010 **Figure 6.** Percentage of individuals in each throat morph category which exhibit a degree of
1011 blue coloration on the throat for (a) wild-caught male hybrids (grey N = 19, orange N = 39,
1012 orange-yellow N = 15, yellow N = 17) and (b) captive-bred F1 hybrids (both sexes; grey N =
1013 31, orange N = 27, orange-yellow N = 2, yellow N = 12).
1014

1015
1016

Figure 7. AICc model-averaged geographic clines (black) and credible regions, clines comprising the 95% confidence set of models (grey), and observed frequencies at each site for (a) the hybrid index estimated from 1333 loci (N = 154), (b) mtDNA haplotype frequencies (N = 74), (c) dorsolateral phenotype (D_{PC1}; N = 233), and (d) throat phenotype clines (N = 204 for both): quantum catch of UV receptor (QC_{UVS}; lower cline, frequencies shown with '×' symbols) and throat colouration (T_{PC1}; upper cline). Numbers along the top refer to sampling sites (see Table S2 for sample sizes at each site), bolded sites (3 – 8) comprise the contact zone. Transect distance is the cumulative distance from the southern-most population with increasing distance northwards. Arrows indicate cline centres; throat phenotype clines have essentially the same centres and are indicated with one arrow.

1027