Comparative transcriptomics reveal distinct patterns of gene expression conservation

through vertebrate embryogenesis

Megan E. Chan[1,2], Pranav S. Bhamidipati[1,2], Heather J. Goldsby[3], Arend Hintze[3], Hans A.

Hofmann[1,2,4], and Rebecca L. Young[1,2,*]

Affiliations:

[1]Department of Integrative Biology, The University of Texas at Austin, Austin, TX, USA

[2]Center for Computational Biology and Bioinformatics, The University of Texas at Austin, Austin,

TX, USA

[3]Department of Integrative Biology, Michigan State University, East Lansing, MI, USA

[4]Institute for Cellular and Molecular Biology, Institute for Neuroscience, The University of Texas

at Austin, Austin, TX, USA

*Correspondence to Dr. Rebecca L. Young, youngrl@utexas.edu.

Abstract

Despite life's diversity, studies of variation across animals often remind us of our shared evolutionary past. Abundant genome sequencing over the last ~25 years reveals remarkable conservation of genes and recent analyses of gene regulatory networks illustrate that not only genes but entire pathways are conserved, reused, and elaborated in the evolution of diversity. Predating these discoveries, 19[th]-century embryologists observed that though morphology at birth varies tremendously, certain stages of embryogenesis appear remarkably similar across vertebrates. Specifically, while early and late stages are variable across species, anatomy of mid-stages embryos (the 'phylotypic' stage) is conserved. This model of vertebrate development and diversification has found mixed support in recent analyses comparing gene expression across species possibly owing to differences across studies in species, embryonic stages, and gene sets compared. Here we perform a comparative analysis using 186 microarray and RNA-seq expression data sets covering embryogenesis in six vertebrate species spanning ~420 million years of evolution. We use an unbiased clustering approach to group stages of embryogenesis by transcriptomic similarity and ask whether gene expression similarity of clustered embryonic stages deviates from the null hypothesis of no relationship between timing and diversification. We use a phylogenetic comparative approach to characterize expression conservation pattern (i.e., early conservation, hourglass, inverse hourglass, late conservation, or no relationship) of each gene at each evolutionary node. Across vertebrates, we find an enrichment of genes exhibiting early conservation, hourglass, late conservation patterns and a large depletion of gene exhibiting no distinguishable pattern of conservation in both microarray and RNA-seq data sets. Enrichment of genes showing patterned conservation through embryogenesis indicates diversification of embryogenesis may be temporally constrained. However, the circumstances (e.g., gene groups, evolutionary nodes, species) under which each pattern emerges remain unknown and require both broad evolutionary sampling and systematic examination of embryogenesis across species.

Introduction

During embryogenesis, a single-cell zygote develops into a multicellular, functional embryo. Given the complexity of this process and the astonishing diversity of resultant phenotypes, the similarities in the developmental processes and anatomy of embryogenesis across species are striking and have captivated the imagination of biologists for nearly two centuries (von Baer, 1828). For example, vertebrates establish a highly conserved body plan ('*bauplan*') from which species-specific variation and elaborations develop. Yet, whether there are generalizable rules that direct diversification of embryogenesis across distantly-related species remains controversial (Richardson et al., 1997; Bininda-Emonds et al., 2003). Inspired by von Baer's (1828) pioneering observations, one hypothesis suggests that early and late phases of embryogenesis are variable across species (owing to diversity and species specificity of reproductive modes and post-body plan elaboration, respectively), while anatomy mid-embryogenesis is conserved (Fig. 1). According to this 'developmental hourglass' hypothesis (Elinson, 1987), similarity of the mid-embryogenesis 'phylotypic stage' (at the pharyngula stage: Ballard, 1976, 1981; Sander, 1983; Richardson, 1995) reflects developmental constraints of body plan formation, including global signaling interdependence and interactions (Raff, 1996; Galis and Metz, 2001) and temporal and spatial patterns of Hox expression (Duboule, 1994). Still others have hypothesized that development of later stages is dependent on early stages of embryogenesis and that this 'developmental burden' results in highest conservation early in embryogenesis (Riedl, 1978; discussed in Irie and Kuratani, 2014) (Fig. 1).

Over the past decade, an increasing number of studies have leveraged the ever growing genome-scale data and approaches to test omics-level predictions of the hourglass hypothesis and its underlying mechanistic basis (Fig. 2A; Supplementary Fig. S1; Supplementary Table 1). Nevertheless, support for the hourglass model of development varies across studies and the question remains very much unsettled (Fig 2B). A number of studies comparing gene expression variation through embryogenesis across species have found support for an increase in expression conservation mid-embryogenesis (Domazet-Loso and Tautz, 2010; Irie and Kuratani, 2011, 2014; Yanai et al., 2011; Levin et al., 2012; Wang et al., 2013; Gerstein et al., 2014; Zalts and Yanai, 2017). Surprisingly, one study comparing expression divergence of animals from different phyla reported an inverse hourglass – where expression differences were highest mid-embryogenesis

(Levin et al., 2016)– although this analysis did not account for phylogenetic non-independence

(Dunn et al., 2018). Still others found that diversification in gene expression is not consistent with

the hourglass model (Piasecka et al., 2013; Tian et al., 2013; Wu et al., 2019). von Baer's (1828)

original morphological observations were based on vertebrate embryos separated by ~420 million

years of evolution; however, recent analyses varied in the evolutionary distances among species

investigated (Fig. 2C), sometimes even spanning more than one phylum (de Mendoza et al., 2013;

Levin et al., 2016; Hu et al., 2017) and often testing predictions of the hourglass hypothesis in only

one or two species (e.g., zebrafish, Domazet-Loso and Tautz, 2010; soft-shell turtle and chicken,

Wang et al., 2013; Caenorhabditis elegans, Zalts and Yanai, 2017; Fig. 2D). Taken together, even

though numerous studies have used sophisticated –omics level analyses to examine embryogenesis

across diverse species, we still lack conclusive molecular evidence of a developmental hourglass.

Any test of the developmental hourglass hypothesis faces several fundamental challenges. First,

proper alignment of stages of embryogenesis across species is difficult. Despite several valiant

attempts to overcome this difficulty (Bininda-Emonds et al., 2003; Mungall et al., 2012; Gerstein

et al., 2014; Li et al., 2014), nomenclature and sampling conventions that often vary substantially

across different model systems as well as widespread heterochrony of developmental events have

prevented a satisfactory solution. Irie and Kuratani (2011) circumvented this challenge by directly

comparing only a subset of well-defined stages across species, but it is unclear to what extent any

observed temporal pattern might be depend on the stages selected for comparison. Second, what

constitutes an appropriate gene set to compare across species is very much unclear. Most studies

to date have examined conservation of all expressed gene for which orthologs can be identified for

all the taxa in the analysis. However, others have suggested that abundant expression of

housekeeping genes may bias discovery of gene expression conservation across species (Piasecka

et al., 2013). Third, it is clear that the genomic and developmental processes underlying even

anatomically similar and homologous phenotypes can diverge via developmental drift or selective

processes (de Beer, 1971; Wagner, 1989; True and Haag, 2001; Wilkins, 2002; McGary et al.,

2010; Young and Wagner, 2011). As a result, some authors have argued that studies examining

the evolution of organismal phenotypes should focus on a core set of regulatory genes critical to

the initiation of the specific developmental program of that character (designated as "kernels" by

Davidson and Erwin, 2006, or "Character Identity Networks, ChINs" by Wagner, 2007). However,

how to identify the relevant gene set (i.e., kernel or ChIN) that is fundamental for shared

developmental processes of vertebrate embryogenesis is unclear. Finally, it is often not clear what

the appropriate null expectation should be in comparative studies, as it may depend on the type of

data available and the level of analysis (Dunn et al., 2018; Young and Hofmann, 2019; Church and

Extavour, 2020).

While numerous studies have reported patterned expression divergence (i.e., early conservation or

hourglass patterns) through embryogenesis across species, what pattern is followed and whether

that diversification pattern varies over evolutionary time remains unclear. Here, we perform a

comparative analysis using 186 publicly available microarray and RNA-seq expression data sets

covering embryogenesis in six vertebrate species spanning ~420 million years of evolution. We

use an unbiased clustering approach to group stages of embryogenesis by transcriptomic similarity

and ask whether gene expression similarity of clustered embryonic stages deviates from the null

hypothesis that gene expression levels are invariant of developmental time. Second, using a

phylogenetic comparative analysis, we characterized the expression conservation pattern (i.e.,

early conservation, hourglass, inverse hourglass, late conservation, or no relationship) exhibited

by each gene at each evolutionary node and ask whether the number of genes that fall into a given

pattern deviates from a biologically meaningful null expectation. Finally, we discuss challenges of

comparative analyses that rely on publicly available transcriptome data and suggest several novel

approaches for future tests of the hourglass hypothesis.

Methods

Obtaining and preprocessing genome-wide gene expression data from public repositories

Gene expression profiles through embryogenesis were obtained from publicly available

repositories for six vertebrate species. In total 112 microarray data sets from five species and 74

RNA-seq data sets from four species were included in the analysis. Developmental time points

included for each species and gene expression profiling technology are detailed in Supplementary

Tables 2 and 3 (microarray and RNA-seq, respectively). Data sets include: zebrafish, *Danio rerio*

(microarray: Roux and Robinson-Rechavi, 2008; RNA-seq: Comte et al., 2010; Yang et al., 2013);

chicken, *Gallus gallus* (microarray: Irie and Kuratani, 2011; RNA-seq: Wang et al., 2013),

Chinese soft-shell turtle, *Pelodiscus sinensis* (RNA-seq: Wang et al., 2013), mouse, *Mus musculus*

(microarray: Irie and Kuratani, 2011; Xue et al., 2013); African clawed frog, *Xenopus laevis*

(microarray: Yanai et al., 2011), and Western clawed frogs, *Xenopus tropicalis* (microarray: Yanai

et al., 2011; RNA-seq: Tan et al., 2013). *D. rerio* expression data at the zygote developmental

stage (0.25 hpf) was excluded because of likely abundance of maternal transcripts, and time points

after 4 dpf were excluded due to substantial completion of the developmental program (after

Kimmel et al., 1995; Yang et al., 2013).

Preprocessing microarray and RNA-seq data

Affymetrix and Agilent microarray data were imported directly using the R packages *simpleaffy*

and *limma*, respectively (Wilson and Miller, 2005; Ritchie et al., 2015). For both microarray

platforms, preprocessing consisted of RMA background correction with quantile normalization

(Irizarry et al., 2003). This information was automatically attained by *limma* for Agilent data sets.

For Affymetrix and Agilent data, probe sets that mapped to multiple genes or no genes at all were

excluded from further analysis. All expression values were then transformed to log-scale using the

function $log_2(x)$ ("Log2-transformed"). For RNA-seq data, raw reads were checked for quality

using FastQC (Andrews, 2010). All data sets were of good quality with less than 10% adaptor

contamination, thus, no trimming was required. After quality control raw reads were

pseudoaligned to species-specific reference transcriptomes using Kallisto to produce read counts

(Bray et al., 2016). Read counts were transformed to transcripts per million mapped reads (TPMs).

The package *biomaRt* was used to assign corresponding ENSEMBL gene ID(s) to each Affymetrix

probe set and RNA-seq transcript (Durinck et al., 2009). For microarray data, the signals of all

probe sets mapping to the same gene were averaged. For RNA-seq counts mapped to different

transcripts of the same gene were summed. Expression of each gene was averaged across

biological replicates for developmental time point. These were the expression values used for

downstream analysis.

Clustering of embryonic stages

We used transcriptomic similarities to classify the embryonic stages of each species in an unbiased

manner. First, we determined the number of clusters using an elbow plot method. Specifically, for

each species and gene expression profiling technology we performed k-means clustering using

gene expression for all developmental stages. We varied the number of clusters from $1 \leq k \leq 9$ for

microarray and $1 \leq k \leq 9$ for RNA-seq and computed the sums of squares error (SSE, or variance

within cluster) for all k. To determine an appropriate number of clusters, we used a) the "elbow"

effect (or determined the k at which additional cluster no longer results in a large reduction in SSE)

and b) determined the k where clustering of groups maintained temporal order of embryogenesis

(i.e., no late stages cluster with early rather than other late stages). Second, to generate clusters of

embryonic stages for each species, we hierarchically clustered stages of embryogenesis by

similarity in gene expression measured as Spearman's rank correlations. The resulting

dendrograms were partitioned into five groups to determine stage clusters. Description of

developmental events were obtained from species-specific references including: zebrafish

(Kimmel et al., 1995), chicken (Hamburger and Hamilton, 1992), softshell turtle (Tokita and

Kuratani, 2001), both *Xenopus* species (Nieuwkoop and Faber, 1994), and mouse (Graham et al.,

2015). For clusters containing more than one embryonic stage, an expression mean was used as

the representative expression for that cluster for the remaining analyses.

Ortholog calling

To identify orthologous genes across taxa we used the sequence-based ortholog calling software

package OrthoMCL (Li et al., 2003) for microarray data and FastOrtho (Wattam et al., 2013; an

implementation of OrthoMCL) for RNA-seq data. Predicted protein sequences of the reference

genomes were organized into orthologous gene groups based on sequence similarity. For each

reference proteome, protein and corresponding gene ids were grouped as paralogs when sequence

similarity was higher among genes within species than between species. To facilitate downstream

analysis of expression conservation across taxa, we removed any orthologous gene groups

containing paralogs or losses/absences in one or more species. The resulting one-to-one orthologs

were used for all downstream analyses (Supplementary Tables 4 and 5, microarray and RNA-seq,

respectively). To assess similarity in microarray and RNA-seq comparison, we compared the one-

to-one ortholog sets of three species (zebrafish, chicken, and the Western clawed frog).

Comparing transcriptomes through embryogenesis across species

For both microarray and RNA-seq data, we assessed transcriptomic similarity at early, middle and
late phases of embryogenesis across species by calculating the Spearman's rank correlation for all
pairwise comparisons of species for each of the five clusters of embryonic stages. For microarray
data, we excluded one frog (*X. laevis*) from the pairwise comparisons to prevent biasing the
outcome as a consequence of high correlations in gene expression between the two congeneric
anuran species at each cluster of embryonic stages. Due to the high correlations between these two
species, similar results were recovered when *X. tropicalis* was removed from the analysis instead
of *X. laevis*. We used permutation analysis to assess whether correlations are higher or lower than
expected by chance. Specifically, for each species we randomly assigned stages to a cluster
maintaining the original number of stages included in the observed cluster and computed the rank
correlation for all pairwise species comparison. We conducted 1000 permutations and assessed
significance by comparing the observed rank correlation to distribution of rank correlations
generated by permutation analysis. Permutation p-values were defined as the percentile of the
observed median Spearman's rho in the distribution of permuted Spearman's rho values. Because
correlations that are higher or lower than expected by chance were both of interest (i.e., a two-
tailed test) the both the percentile and 1-percentile were calculated using the empirical cumulative
distribution function in R, the lower of the two is reported.

Characterizing expression conservation of each gene through embryogenesis across vertebrates

To assess conservation of gene expression for each gene at each cluster of embryonic stages and
each node, we calculated a difference in expression rank scaled by the divergence time between
the groups (Supplementary Fig. S2). At each node of the phylogeny, we characterized patterns of
expression conservation across embryogenesis using the R package Clustering of Time Series
Gene Expression Data (*ctsGE*; Sharabi-Schwager and Ophir, 2019). Using *ctsGE*, gene
conservation scores of each gene were median-scaled and converted into conservation indices. For
each gene, at each cluster of embryonic stages, the standardized values indicate the median
absolute distance of that gene from its median conservation score. These standardized values were

then converted to index values that indicate whether gene expression conservation was above (1),

below (-1), or within (0) the cutoff range (+/- 0.7) of the median value at each time step (here

cluster of embryonic stages). Each index of expression conservation across embryogenesis was

assigned to a conservation pattern based on median transitions across the assigned significance

cutoff (+/- 0.7). For early conservation: similarity decreases through embryogenesis; hourglass:

similarity increases and then decreases: inverse hourglass: similarity decreases and then increases;

late conservation: similarity increases; invariant: similarity does not vary or does not follow other

conservation patterns through embryogenesis (Fig. 1A). Indices assignments are provided in

Supplementary Table 6.

To examine enrichment of genes in each conservation pattern, we first calculated the proportion

of indices in each conservation pattern and defined the expected number of genes as the equivalent

proportion of total genes. We determined significance of enrichment/depletion of genes exhibiting

each pattern using a permutation analysis. For each gene we first randomized the order of

conservation scores across the clusters of embryonic stages. Second, we characterized

conservation trajectory using ctsGE and the index assignment rules described above. Finally, for

each iteration we calculated enrichment/depletion of genes compared to the random expectation

for each conservation pattern. We repeated this permutation 1000 times for each node and gene

expression profiling technology and assessed by comparing the observed enrichment/depletion to

the null distribution generated by the permutation analysis. Permutation p-values were defined as

the probability of obtaining enrichment/depletion of genes at or above/below the observed number

in the permutation set as described above.

Results

Variation in gene sets and alignment of embryonic stage clusters across species and gene

expression profiling technologies

We identified 1626 and 1793 one-to-one orthologs for the microarray and RNA-seq data in our

analysis, respectively, consistent with other comparative gene expression studies spanning

vertebrates (3044 avian to human orthologs: Pfenning et al., 2014; 1979 orthologs across

Peromyscus mice, Microtus voles, Passeroid birds, Dendrobatid frogs, and Ectodini cichlids:

Young et al., 2019). Interestingly, one-to-one orthologs from the microarray and RNA-seq data set

were largely non-overlapping with a total of 255 overlapping genes (15.7% and 14.2% of each set

of orthologs, respectively). Similarly, one-to-one orthologs from microarray and RNA-seq data

sets from the same species were largely non-overlapping as well (~15-20%, Supplementary Table

7). Consistency of this overlap across species suggests that microarray and RNA-seq approaches

may target distinct features of the transcriptome.

When we clustered the embryonic stages according to their gene expression patterns we found

that, for each species and gene expression profiling technology, a $k = 5$ emerged as the number of

clusters at which the reduction in within cluster-variance begins to asymptote (Fig. 3). Importantly,

the temporal order of stages across embryogenesis was maintained at $k = 5$ clusters; however,

inconsistent sampling across species as well as heterochrony across species resulted in some

variation in the major events contained in each cluster of embryonic stages across species and

between microarray and RNA-seq platforms (Fig. 4). Stages contained in each cluster and a

biological description of embryonic events is provided in Fig. 5 and Supplementary Tables 2 and

3 (microarray and RNA-seq, respectively). The number of stages collapsed into each cluster also

varied across species and gene expression profiling technology. Because there was no apparent

bias between cluster timing in embryogenesis and the number of stages included (Fig. 5), we

concluded that $k = 5$ clusters was appropriate for downstream analyses.

Conservation of gene expression at the transcriptome level

We then quantified interspecific gene expression correlation for each of the $k = 5$ clusters of

embryogenesis identified in Fig. 4. Clusters that are correlated across species are more conserved.

We found that the pattern of pairwise rank correlations for all one-to-one orthologs (1626 and 1793

for microarray and RNA-seq, respectively) through embryogenesis differed for microarray and

RNA-seq comparisons (Fig. 6). In the microarray comparison, median pairwise rank correlations

increased mid-embryogenesis, with clusters 2, 3, and 4 having the highest median correlation, and

decreased early and late in embryogenesis, with cluster 5 having the lowest median, suggesting a

developmental hourglass (Fig. 6A). However, when compared to the null expectation of no gene

expression conservation above chance, generated by permuting the cluster assignment of each

stage, the observed pattern did not differ from the null (cluster 1: observed median Spearman's ρ_o

= 0.46, permutation Spearman's ρ_p = [0.37-0.54], p = 0.43; cluster 2: ρ_o = 0.49, ρ_p = [0.4-0.53], p

= 0.26; cluster 3: ρ_o = 0.49, ρ_p = [0.42=0.53], p = 0.47; cluster 4: ρ_o = 0.48, ρ_p = [0.44-0.53], p =

0.35; cluster 5: ρ_o = 0.46, ρ_p = [0.36-0.52], p = 0.49). In the RNA-seq comparison, median pairwise

rank correlations increased to its highest median score at cluster 2 and dropped through later stages

of embryogenesis 3, with cluster 5 having the lowest median (Fig. 6B). When compared to the null

expectation, median pairwise rank correlation of cluster 2 was significantly higher and correlations

of clusters 4 and 5 were significantly lower than expected by chance, providing strong support for

a developmental hourglass (cluster 1: ρ_o = 0.65, ρ_p = [0.56-0.68], p = 0.24; cluster 2: ρ_o = 0.67, ρ_p

= [0.52-0.69], p = 0.005; cluster 3: ρ_o = 0.62, ρ_p = [0.51-0.68], p = 0.49; cluster 4: ρ_o = 0.57, ρ_p =

[0.57-0.68], p = 0.003; cluster 5: ρ_o = 0.53, ρ_p = [0.59-0.69], p = 0.008).

Enrichment and overlap of expression conservation patterns across the phylogeny

Next, we used ctsGE time series analysis to characterize patterns of gene expression conservation

across embryogenesis at each phylogenetic node. For the microarray data set, gene conservation

scores yielded a total of 90, 90, 87, 89 patterns of expression conservation in anurans, amniotes,

tetrapods, and vertebrates, respectively. For the RNA-seq data set, gene conservation scores

yielded a total of 90, 51, and 49 patterns in amniotes, tetrapods, and vertebrates. We calculated the

expected number of genes for each pattern as equivalent to the proportion of total patterns (Fig. 7,

treemap insets; Tennekes, 2017). Across vertebrates, with the exception of inverse hourglass in

the RNA-seq analysis and time-invariance in both platforms, all patterns were exhibited by

significantly more genes than expected by chance (Fig. 7). This enrichment of genes across all

conservation patterns and the depletion of time-invariant genes deviated significantly from the null

expectation (microarray: early conservation p = 0, hourglass p = 0, inverse hourglass p = 0, late

conservation p = 0, time invariance p = 0; RNA-seq: early conservation p = 0, hourglass p = 0.02,

inverse hourglass p = 0.4, late conservation p = 0, time invariance p = 0) Enrichment/depletion of

genes for each conservation pattern across anurans (microarray only), amniotes, and tetrapods are

provided in Supplementary Fig. S3. p = 0 indicates that none of the 1000 permutation iterations

resulted in enrichment/depletion at or above/below the observed level.

Finally, we compared conservation pattern assignment for the 255 one-to-one orthologous gene
groups shared between microarray and RNA-seq gene sets and found that across platforms gene
expression patterns are not concordant at the major phylogenetic nodes. Specifically, only 25%
of genes (63/255) in amniotes, 19% (49/255) in tetrapods, and 20% (52/255) in vertebrates
exhibited concordant expression conservation patterns.

Discussion

In the present study, we examined patterns of expression diversification throughout embryogenesis
in six vertebrate species using a comparative analysis of 112 microarray and 74 RNA-seq data sets
(Fig. 1B). First, using an unbiased approach, we clustered stages of embryogenesis within species
and compared expression conservation of those clusters across species (Figs. 3-5; Supplementary
Tables 2 and 3). This approach allowed for the inclusion of all available stages of embryogenesis
and removed bias that could result from comparing only selected stages. Second, we used a
permutation analysis to generate a null expectation. We observed conservation estimates against
this null expectation to characterize transcriptome-level diversification through embryogenesis
across species (Fig. 4). Finally, we characterized the expression conservation of each gene at each
node of the phylogeny (Fig. 1B; Supplementary Fig. S2) to examine how expression conservation
patterns vary through evolutionary time (Fig. 6).

Over the past decade, the debate of whether diversification of embryogenesis follows generalizable
rules has been reinvigorated by the ability to test predictions of the hourglass, developmental
burden, and other hypotheses (Fig. 1A) on a genomic scale. Enabled by the increase in 'omics-
level data and next generation sequencing accessibility, a number of studies have by now explored
patterns of diversification in gene expression though embryogenesis across species (Fig. 2). These
studies have found mixed support for the hourglass and other models of diversification across
species (Fig. 2B-D; reviewed in: Irie, 2017; Liu and Robinson-Rechavi, 2018). Differences among
studies could reflect differences in species compared as some studies span phyla (de Mendoza et
al., 2013; Levin et al., 2016; Hu et al., 2017) and others are restricted to internal nodes of the
vertebrate phylogeny (e.g., amniotes; Wang et al., 2013). However, our quantitative literature
analysis did not indicate an effect of divergence time on the characterization of an hourglass model
of divergence (Fig. 2C). Alternatively, selection of developmental stages or specific gene sets to

compare may lead to different results and interpretations. Finally, studies that compare gene expression similarity at the transcriptome level rarely test against a null hypothesis (Young and Hofmann, 2019). Such a test is critical because the degree of variation expected through developmental stages across species is unknown (Church and Extavour, 2020).

Aligning stages of early animal development is complicated by taxon-specific sampling practices as well as pervasive heterochrony in developmental events across distantly related species, leading some researchers to question the validity of anatomical hourglass hypotheses (Bininda-Emonds et al., 2003). We found that while expression varies in similar ways through embryogenesis across species, stages of embryogenesis did not always consistently cluster together within species (Fig.3 and 4; Supplementary Tables 2 and 3). These differences likely reflect both biological variation in the molecular timing of developmental events, technical variation in sampling procedures across species, and a lack of available data sets particularly at early embryonic stages. Because clustering has the advantage of being unbiased, and no systematic bias in sampling was apparent, we moved forward by comparing gene expression at each embryonic stage cluster across species. Future studies with systematic sampling of embryogenesis across species could disentangle the source (biological or technical) of variation in stage clustering across species.

Consistent with the mixed support for the hourglass and other models of developmental divergence (e.g., early conservation or inverse hourglass) found across studies, our comparisons of expression variation in all one-to-one orthologs present across species in the microarray and RNA-seq data yielded significant but different patterns. Specifically, the observed gene expression correlations (as a measure of conservation) differed through embryogenesis for both microarray and RNA-seq, yet only the RNA-seq data showed a pattern that differed significantly from the null expectation that gene expression levels should be invariant of developmental time (Fig. 6). For the RNA-seq datasets, we found a significant increase in expression correlation over the null expectation in developmental time cluster 2 followed by a significant reduction in expression correlation in the later clusters 4 and 5. Though clusters 1 and 2 display similar gene expression correlations, suggesting an early conservation pattern, cluster 1 does not significantly differ from the null expectation. Inconsistencies between the microarray and RNA-seq data sets could result from differences in taxon sampling (Fig. 1B), the embryonic stages and resulting clusters that were included (Figs. 4 and 5), and/or systematic differences in which aspects of the transcriptome were

captured by these distinct platforms. In fact, transcriptome-level gene expression profiles

quantified using RNA-seq and microarray technologies have been shown to be correlated

especially for highly expressed genes, but variation between technologies is also commonly

reported (Marioni et al., 2008; Malone and Oliver, 2011; Trost et al., 2015). Whether these

differences reflect superiority of one technology over the other is unclear. Instead technical

differences between the two approaches may capture different elements of the transcriptome, in

which case the two approaches should be viewed as complementary (Kogenaru et al., 2012). In

our data sets, we found little overlap of one-to-one orthologs between microarray and RNA-seq,

with only ~15-20% of the genes contained in both analyses. Further, of those one-to-one orthologs

contained in both microarray and RNA-seq data sets, only ~20% shared conservation pattern

assignments (Fig. 8). This illustrates a major challenge for comparative analyses, like our present

study, that aim to capitalize on the vast amounts of publicly available transcriptome datasets to test

biological hypotheses.

Inconsistency across studies and gene expression profiling technology also suggests that

comparing the whole transcriptome may not be appropriate. Specifically, patterns of expression

conservation at the whole transcriptome-level may be biased by abundant and constitutively

expressed genes (e.g., see Piasecka et al., 2013). Whole genome approaches have the great

advantage of being unbiased, which allows for the identification of novel gene associations with

phenotypes and/or gene interactions that would be missed using a candidate gene approach.

However, we know that variation is not equivalent across levels of biological organization. For

example, studies comparing mRNA and protein levels have found only modest overall correlation

(Foss et al., 2007; Fu et al., 2009; Ghazalpour et al., 2011; Vogel and Marcotte, 2012). Further,

studies of character homology have shown that even anatomically and physiologically similar

homologies can differ in underlying developmental and molecular mechanisms (Wagner, 1989;

Wilkins, 2002; McGary et al., 2010; Young and Wagner, 2011). To address this issue, we also

used a time series analysis to characterize the conservation pattern of each gene at each node of

the phylogeny (Fig. 7), in addition to a transcriptome-level comparisons through embryogenesis

across species. Across vertebrates, we observed a significant enrichment of all patterned

conservation models (early conservation, hourglass, inverse hourglass, and late conservation)

above the expected number and a large depletion of genes whose expression was invariant of

developmental time (Fig. 7; Supplementary Fig. S3) in both microarray and RNA-seq data sets.

Both the enrichment of genes exhibiting different patterns and the proportion of overall genes with

distinct patterns varied across evolutionary nodes (e.g., tetrapods, amniotes, and anurans, Fig. 1B)

and between gene expression profiling technologies (Supplementary Fig. S3). These results

suggest that divergence in gene expression through embryogenesis may depend on the

evolutionary distance covered by any given analysis. Follow-up studies, including phylogenetic

comparative analyses of both closely and distantly related species are needed to better understand

these patterns and their implications for generating biological diversity. In addition, time series

analyses that characterize expression divergence of individual genes or gene sets should be used

to test hypothetical mechanisms of constraint. For example, we might ask whether regulatory

interactions or temporal and spatial expression patterns of a gene follow a correlated dynamic

pattern through embryogenesis.

Conclusions:

Taken together, our results provided strong support for a patterned embryonic gene expression

diversification across vertebrate species. However, the gene groups and evolutionary nodes under

which each pattern emerges remain unknown. By combining unbiased clustering of embryonic

stages and explicit tests against a null hypothesis our research demonstrates a critical need for

broad evolutionary sampling and systematic examination of developmental stages across species

to better characterize gene expression diversification in embryogenesis.

Acknowledgements

We thank D. Wylie for consulting on data analysis, S. Campbell for helping with quantitative

literature analyses, and, S. Caro, C. Friesen, I. Miller-Crews, M. Rodriguez Santiago, and T.

Solomon-Lane Santiago for discussions that improved this manuscript. This work was funded in

part by a University of Texas at Austin Undergraduate Research Fellowship to P.S.B. and a grant

from the NSF BEACON Center for the Study of Evolution in Action, DBI-0939454 to H.J.G,

A.H., H.A.H., and R.L.Y.

423
424

Literature Cited

Andrews, S. 2010. FastQC: a quality control tool for high throughput sequence data. Available online at: http://www.bioinformatics.babraham.ac.uk/projects/fastqc.

von Baer, K.E. 1828. Uber Entwickelungsgeschichte der Thiere (Königsberg: Bornträger).

Ballard, W.W. 1976. Problems of Gastrulation: Real and Verbal. *Bioscience* 26: 36–39.

Ballard, W.W. 1981. Morphogenetic movements and fate maps of vertebrates. *Am Zool* 21:.

de Beer, G. 1971. Homology, An Unsolved Problem, Volume 11 (Oxford: Oxford University Press).

Bininda-Emonds, O.R.P., Jeffery, J.E., and Richardson, M.K. 2003. Inverting the hourglass: quantitative evidence against the phylotypic stage in vertebrate development. *Proc. R. Soc. London B Biol. Sci.* 270: 341–346.

Bray, N.L., Pimentel, H., Melsted, P., and Pachter, L. 2016. Near-optimal probabilistic RNA-seq quantification. *Nat. Biotechnol.* 34: 525.

Church, S.H., and Extavour, C.G. 2020. Null hypotheses for developmental evolution. *Development* 147: dev178004.

Comte, A., Roux, J., and Robinson-Rechavi, M. 2010. Molecular signaling in zebrafish development and the vertebrate phylotypic period. *Evol. Dev.* 12: 144–156.

Davidson, E.H., and Erwin, D.H. 2006. Gene Regulatory Networks and the Evolution of Animal Body Plans. *Science (80-.).* 311: 796–800.

Domazet-Loso, T., and Tautz, D. 2010. A phylogenetically based transcriptome age index mirrors ontogenetic divergence patterns. *Nature* 468: 815–818.

Duboule, D. 1994. Temporal colinearity and the phylotypic progression: a basis for the stability of a vertebrate Bauplan and the evolution of morphologies through heterochrony. *Dev.* 1994: 135–142.

Dunn, C.W., Zapata, F., Munro, C., Siebert, S., and Hejnol, A. 2018. Pairwise comparisons across species are problematic when analyzing functional genomic data. *Proc. Natl. Acad. Sci.* .

Durinck, S., Spellman, P., Birney, E., and Huber, W. 2009. Mapping identifiers for the integration of genomic datasets with the R/Bioconductor package biomaRt. *Nat. Protoc.* 4: 1184–1191.

Elinson, R. 1987. Changes in developmental patterns: embryos of amphibians with large eggs. In Development as an Evolutionary Process, R.A. Raff, ed. (New York: Liss), pp. 1–21.

Foss, E.J., Radulovic, D., Shaffer, S.A., Ruderfer, D.M., Bedalov, A., Goodlett, D.R., and Kruglyak, L. 2007. Genetic basis of proteome variation in yeast. *Nat. Genet.* 39: 1369–1375.

Fu, J., Keurentjes, J.J.B., Bouwmeester, H., America, T., Verstappen, F.W.A., Ward, J.L., Beale, M.H., de Vos, R.C.H., Dijkstra, M., Scheltema, R.A., et al. 2009. System-wide molecular evidence for phenotypic buffering in Arabidopsis. *Nat. Genet.* 41: 166–167.

Galis, F., and Metz, J.A.J. 2001. Testing the vulnerability of the phylotypic stage: On modularity and evolutionary conservation. *J. Exp. Zool.* 291: 195–204.

Gerstein, M.B., Rozowsky, J., Yan, K.-K., Wang, D., Cheng, C., Brown, J.B., Davis, C.A., Hillier, L., Sisu, C., Li, J.J., et al. 2014. Comparative analysis of the transcriptome across distant species. *Nature* 512: 445–448.

Ghazalpour, A., Bennett, B., Petyuk, V.A., Orozco, L., Hagopian, R., Mungrue, I.N., Farber, C.R., Sinsheimer, J., Kang, H.M., Furlotte, N., et al. 2011. Comparative Analysis of Proteome and Transcriptome Variation in Mouse. *PLOS Genet.* 7: e1001393.

Graham, E., Moss, J., Burton, N., Armit, C., Richardson, L., and Baldock, R. 2015. The atlas of mouse development eHistology resource. *Development* 142: 1909 LP – 1911.

Hamburger, V., and Hamilton, H.L. 1992. A series of normal stages in the development of the chick embryo. *Dev Dyn* 88: 231–272.

Hu, H., Uesaka, M., Guo, S., Shimai, K., Lu, T.-M., Li, F., Fujimoto, S., Ishikawa, M., Liu, S., Sasagawa, Y., et al. 2017. Constrained vertebrate evolution by pleiotropic genes. *Nat. Ecol. Evol.* 1: 1722–1730.

Irie, N. 2017. Remaining questions related to the hourglass model in vertebrate evolution. *Curr. Opin. Genet. Dev.* 45: 103–107.

Irie, N., and Kuratani, S. 2011. Comparative transcriptome analysis reveals vertebrate phylotypic period during organogenesis. *Nat. Commun.* 2: 248.

Irie, N., and Kuratani, S. 2014. The developmental hourglass model: a predictor of the basic body plan? *Development* 141: 4649–4655.

Irizarry, R.A., Hobbs, B., Collin, F., Beazer-Barclay, Y.D., Antonellis, K.J., Scherf, U., and Speed, T.P. 2003. Exploration, normalization, and summaries of high density oligonucleotide array probe level data. *Biostatistics* 4: 249–264.

Kimmel, C.B., Ballard, W.W., Kimmel, S.R., Ullmann, B., and Schilling, T.F. 1995. Stages of embryonic development of the zebrafish. *Dev. Dyn.* 203: 253–310.

Kogenaru, S., Qing, Y., Guo, Y., and Wang, N. 2012. RNA-seq and microarray complement each other in transcriptome profiling. *BMC Genomics* 13: 629.

Kumar, S., Stecher, G., Suleski, M., and Hedges, S.B. 2017. TimeTree: A Resource for Timelines, Timetrees, and Divergence Times. *Mol. Biol. Evol.* 34: 1812–1819.

Levin, M., Anavy, L., Cole, A.G., Winter, E., Mostov, N., Khair, S., Senderovich, N., Kovalev, E., Silver, D.H., Feder, M., et al. 2016. The mid-developmental transition and the evolution of animal body plans. *Nature* advance on:

Levin, M., Hashimshony, T., Wagner, F., and Yanai, I. 2012. Developmental Milestones Punctuate Gene Expression in the Caenorhabditis Embryo. *Dev. Cell* 22: 1101–1108.

Li, J.J., Huang, H., Bickel, P.J., and Brenner, S.E. 2014. Comparison of D. melanogaster and C. elegans developmental stages, tissues, and cells by modENCODE RNA-seq data. *Genome Res.* 24: 1086–1101.

Li, L., Stoeckert, C.J., and Roos, D.S. 2003. OrthoMCL: Identification of Ortholog Groups for
Eukaryotic Genomes. *Genome Res.* 13: 2178–2189.

Liu, J., and Robinson-Rechavi, M. 2018. Developmental Constraints on Genome Evolution in
Four Bilaterian Model Species. *Genome Biol. Evol.* 10: 2266–2277.

Malone, J.H., and Oliver, B. 2011. Microarrays, deep sequencing and the true measure of the
transcriptome. *BMC Biol.* 9: 34.

Marioni, J.C., Mason, C.E., Mane, S.M., Stephens, M., and Gilad, Y. 2008. RNA-seq: an
assessment of technical reproducibility and comparison with gene expression arrays. *Genome*
Res 18: 1509–1517.

McGary, K.L., Park, T.J., Woods, J.O., Cha, H.J., Wallingford, J.B., and Marcotte, E.M. 2010.
Systematic discovery of nonobvious human disease models through orthologous phenotypes.
Proc. Natl. Acad. Sci. 107: 6544–6549.

de Mendoza, A., Sebé-Pedrós, A., Šestak, M.S., Matejčić, M., Torruella, G., Domazet-Lošo, T.,
and Ruiz-Trillo, I. 2013. Transcription factor evolution in eukaryotes and the assembly of the
regulatory toolkit in multicellular lineages. *Proc. Natl. Acad. Sci.* 110: E4858 LP-E4866.

Moher, D., Liberati, A., Tetzlaff, J., Altman, D.G., and Group, T.P. 2009. Preferred Reporting
Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PLOS Med.* 6:
e1000097.

Mungall, C.J., Torniai, C., Gkoutos, G. V, Lewis, S.E., and Haendel, M.A. 2012. Uberon, an
integrative multi-species anatomy ontology. *Genome Biol.* 13: R5.

Nieuwkoop, P.D., and Faber, J. 1994. Normal table of Xenopus laevis (Daudin) : a systematical
and chronological survey of the development from the fertilized egg till the end of
metamorphosis (New York: Garland Pub.).

Pfenning, A.R., Hara, E., Whitney, O., Rivas, M. V, Wang, R., Roulhac, P.L., Howard, J.T.,
Wirthlin, M., Lovell, P. V, Ganapathy, G., et al. 2014. Convergent transcriptional specializations
in the brains of humans and song-learning birds. *Sci.* 346:.

Piasecka, B., Lichocki, P., Moretti, S., Bergmann, S., and Robinson-Rechavi, M. 2013. The
Hourglass and the Early Conservation Models—Co-Existing Patterns of Developmental
Constraints in Vertebrates. *PLoS Genet* 9: e1003476.

Raff, R.A. 1996. The shape of life: Genes, development, and the evolution of animal form
(Chicago: University of Chicago Press).

Richardson, M.K. 1995. Heterochrony and the Phylotypic Period. *Dev. Biol.* 172: 412–421.

Richardson, M.K., Hanken, J., Gooneratne, M.L., Pieau, C., Raynaud, A., Selwood, L., and
Wright, G.M. 1997. There is no highly conserved embryonic stage in the vertebrates:
implications for current theories of evolution and development. *Anat Embryol* 196:.

Riedl, R. 1978. Order in Living Systems: A Systems Analysis of Evolution (New York: Wiley).

Ritchie, M.E., Phipson, B., Wu, D., Hu, Y., Law, C.W., Shi, W., and Smyth, G.K. 2015. limma
powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic*

Acids Res. 43: e47–e47.

Roux, J., and Robinson-Rechavi, M. 2008. Developmental Constraints on Vertebrate Genome
Evolution. *PLoS Genet* 4: e1000311.

Sander, K. 1983. The evolution of patterning mechanisms: gleanings from insect embryogenesis
and spermatogenesis BT - Development and Evolution. In Development and Evolution, B.C.
Goodwin, N. Holder, and C.C. Wylie, eds. (Cambridge: Cambridge University Press), pp. 137–
159.

Sharabi-Schwager, M., and Ophir, R. 2019. ctsGE: Clustering of Time Series Gene Expression
data. R package version 1.12.0, https://github.com/michalsharabi/ctsGE.

Tan, M.H., Au, K.F., Yablonovitch, A.L., Wills, A.E., Chuang, J., Baker, J.C., Wong, W.H., and
Li, J.B. 2013. RNA sequencing reveals a diverse and dynamic repertoire of the Xenopus
tropicalis transcriptome over development. *Genome Res.* 23: 201–216.

Tennekes, M. 2017. treemap: Treemap Visualization.

Tian, X., Strassmann, J.E., and Queller, D.C. 2013. Dictyostelium Development Shows a Novel
Pattern of Evolutionary Conservation. *Mol. Biol. Evol.* 30: 977–984.

Tokita, M., and Kuratani, S. 2001. Normal Embryonic Stages of the Chinese Softshelled Turtle
Pelodiscus sinensis (Trionychidae). *Zoolog. Sci.* 18: 705–
715.

Trost, B., Moir, C.A., Gillespie, Z.E., Kusalik, A., Mitchell, J.A., and Eskiw, C.H. 2015.
Concordance between RNA-sequencing data and DNA microarray data in transcriptome analysis
of proliferative and quiescent fibroblasts. *R. Soc. Open Sci.* 2: 150402.

True, J.R., and Haag, E.S. 2001. Developmental system drift and flexibility in evolutionary
trajectories. *Evol. Dev.* 3: 109–119.

Vogel, C., and Marcotte, E.M. 2012. Insights into the regulation of protein abundance from
proteomic and transcriptomic analyses. *Nat. Rev. Genet.* 13: 227–232.

Wagner, G.P. 1989. The Biological Homology Concept. *Annu. Rev. Ecol. Syst.* 20: 51–69.

Wagner, G.P. 2007. The developmental genetics of homology. *Nat Rev Genet* 8: 473–479.

Wang, Z., Pascual-Anaya, J., Zadissa, A., Li, W., Niimura, Y., Huang, Z., Li, C., White, S.,
Xiong, Z., Fang, D., et al. 2013. The draft genomes of soft-shell turtle and green sea turtle yield
insights into the development and evolution of the turtle-specific body plan. *Nat Genet* 45: 701–
706.

Wattam, A.R., Abraham, D., Dalay, O., Disz, T.L., Driscoll, T., Gabbard, J.L., Gillespie, J.J.,
Gough, R., Hix, D., Kenyon, R., et al. 2013. PATRIC, the bacterial bioinformatics database and
analysis resource. *Nucleic Acids Res.* 42: D581–D591.

Wilkins, A.S. 2002. The Evolution of Developmental Pathways (Sunderland, MA: Sinauer
Associates, Inc).

Wilson, C.L., and Miller, C.J. 2005. Simpleaffy: a BioConductor package for Affymetrix Quality

Control and data analysis. *Bioinformatics* 21: 3683–3685.

Wu, L., Ferger, K.E., and Lambert, J.D. 2019. Gene Expression Does Not Support the
Developmental Hourglass Model in Three Animals with Spiralian Development. *Mol. Biol. Evol.*
36: 1373–1383.

Xue, Z., Huang, K., Cai, C., Cai, L., Jiang, C., Feng, Y., Liu, Z., Zeng, Q., Cheng, L., Sun, Y.E.,
et al. 2013. Genetic programs in human and mouse early embryos revealed by single-cell
RNA[thinsp]sequencing. *Nature* 500: 593–597.

Yanai, I., Peshkin, L., Jorgensen, P., and Kirschner, M.W. 2011. Mapping Gene Expression in
Two Xenopus Species: Evolutionary Constraints and Developmental Flexibility. *Dev. Cell* 20:
483–496.

Yang, H., Zhou, Y., Gu, J., Xie, S., Xu, Y., Zhu, G., Wang, L., Huang, J., Ma, H., and Yao, J.
2013. Deep mRNA Sequencing Analysis to Capture the Transcriptome Landscape of Zebrafish
Embryos and Larvae. *PLoS One* 8: e64058.

Young, R.L., Ferkin, M.H., Ockendon-Powell, N.F., Orr, V.N., Phelps, S.M., Pogány, Á.,
Richards-Zawacki, C.L., Summers, K., Székely, T., Trainor, B.C., et al. 2019. Conserved
transcriptomic profiles underpin monogamy across vertebrates. *Proc. Natl. Acad. Sci.* 116: 1331
LP – 1336.

Young, R.L., and Hofmann, H.A. 2019. Reply to Jiang and Zhang: Parallel transcriptomic
signature of monogamy: What is the null hypothesis anyway? *Proc. Natl. Acad. Sci.* 116: 17629
LP – 17630.

Young, R.L., and Wagner, G.P. 2011. Why ontogenetic homology criteria can be misleading:
lessons from digit identity transformations. *J. Exp. Zool. B. Mol. Dev. Evol.* 316B: 165–170.

Zalts, H., and Yanai, I. 2017. Developmental constraints shape the evolution of the nematode
mid-developmental transition. *Nat. Ecol. Evol.* 1: 113.

Figures

Figure 1. Anatomical and gene expression similarity predicted under different models of

conservation through embryogenesis (A). Expression conservation was assessed using 186

publicly available microarray and RNA-seq data sets through embryogenesis across a phylogeny

of six vertebrate species (B). Divergence times at each node are shown in millions of years.

Figure 2. Studies testing the developmental hourglass and alternative hypotheses have increased considerably since the start of the new millennium (A, grey line: $r^2 = 0.44$, $F(1, 18) = 15.7$, $p = 0.0009$, mainly driven by an increase in the number of studies focused on gene expression patterns (A, black line: $r^2 = 0.6$, $F(1, 18) = 29.4$, $p = 3.729e-05$). Despite this increased attention to this question, whether variation in embryogenesis follows an hourglass pattern has remained unresolved (B, orange points: $r^2 = -0.06$, $F(1, 18) = 0.004$, $p = 0.95$;). Neither divergence time of the species compared (C), nor number of species included in any given study (D) affect whether an hourglass pattern is observed. Quantitative literature analysis of studies examining early embryonic development across species was carried out according to the PRISMA flow diagram (Supplementary Fig. 1; Moher et al., 2009). Detailed methods are provided in Supplementary Materials (Supplementary Table 1, Supplementary Fig. S1)

Figure 3. *k*-means clustering of the microarray (A) and RNA-seq (B) data sets analyzed.
Reduction of within cluster variance increases as the number of cluster (*k*) increases. Gains
asymptote at approximately *k* = 5 (dashed line).

627   **Figure 4.** Spearman Rank correlations were used to group stages into five clusters. Shown are all

628   pairwise correlations of stages for all species and both gene expression profiling technologies.

629   Grouping of stages is in indicated color.

Figure 5. Overlap of embryonic stage cluster and major developmental events. Variation in sampling and heterochrony among species result in differences of developmental events captured by the available data across species and platforms. For three species, gene expression datasets were obtained from both microarray (top) and RNA-seq (bottom) platforms.

Figure 6. Spearman rank correlations for pairwise comparisons of species at each cluster of embryogenesis for microarray (A) and RNA-seq data (B). Gene expression correlations (as a measure of conservation) vary through embryogenesis for both microarray and RNA-seq data and appear to show support for a developmental hourglass. Colored boxes indicate observed correlations; grey boxes indicate rank correlations after permutation analysis randomizing stage association with cluster. Asterisks indicate that the observed median correlation differs significantly from the null expectation at *p < 0.01*. Note that only the RNA-seq data showed a temporal pattern (consistent with the early conservation hypothesis) that differed from the null expectation (i.e., no temporal pattern present).

Figure 7. Enrichment of patterned gene expression conservation across vertebrates. The expected number of genes for a given temporal expression pattern is determined by the proportion of trajectories of each conservation pattern (A.2: microarray; B.2: RNA-seq). The proportion of trajectories differed across conservation patterns and between platforms (A.2 and B.2). The enrichment/depletion values from permutation analysis (1000 iterations) are shown as violin plots (A.1 and B.1). The number of genes associated with all conservation patterns were enriched compared to random expectation, except for the inverse hourglass (RNA-seq only), whereas the number of genes with no relationship between expression conservation across embryogenesis (A.1: microarray; B.1: RNA-seq) was significantly less than expected by chance. For all plots colors indicate the expression conservation pattern. Enrichment at other evolutionary nodes (tetrapods, amniotes, and anurans) are provided in Supplementary Fig. S3.

Figure 8. 255 one-to-one orthologs shared between microarray and RNA-seq data sets largely

differed in expression conservation pattern (A). Most genes with shared expression conservation

patterns were "no relationship" genes followed by hourglass or inverse hourglass genes (B).

Supplementary Tables and Figures

Supplementary Table 1. Studies and data included in literature analyses Fig. 2.

Supplementary Table 2. Description and clustering of embryonic stages in species with microarray
data

Supplementary Table 3. Description and clustering of embryonic stages in species with RNA-seq
data

Supplementary Table 4. Orthologous gene groups for microarray datasets

Supplementary Table 5. Orthologous gene groups for RNA-seq datasets

Supplementary Table 6. Gene conservations pattern indices

Supplementary Table 7. Number and percentage of overlapping one-to-one orthologs in
microarray and RNA-seq datasets

Species	# of overlapping orthologs	% of orthologs (microarray)	% of orthologs (RNA-seq)
Gallus gallus	264	18.7%	18.0%
Xenopus tropicalis	345	21.2%	19.2%
Danio rerio	335	20.6%	16.2%

Supplementary Figure S1. Modified PRIMSA (Moher et al., 2009) flow chart. 220 unique articles retrieved from Web of Science using the following search criteria topic (TS) equals "phylotypic" OR "developmental AND hourglass" on March 25, 2020. We removed 53 articles that were not about the developmental hourglass hypothesis (e.g., in microbiology, "phylotypic" is sometimes used to classify a group of organisms by their phenetic relationship). We removed an additional 64 methods and review articles that did not include data analysis, 30 articles that included intraspecific comparisons only, and 14 articles that compared development of individual phenotypes (e.g., craniofacial morphology, brain, limbs, and heart) rather than embryogenesis. For each of the remaining 59 unique articles (Supplementary Table 1), the pattern of variation (early conservation, hourglass, inverse hourglass, late conservation, or no relationship) was reported for the type of data analyzed (gene expression, molecular evolution, morphological, or regulatory). 11 studies included analyses of data from different biological levels of organization (e.g., molecular evolution and transcriptomics). These analyses were treated as independent resulting in a total of 70 entries. Finally, four studies compared species with unresolved divergence times. These articles were excluded from divergence time comparisons in Fig. 2C. Divergence time estimates were

692 obtained from TimeTree (Kumar et al., 2017). A linear regression model was used to assess

693 increase in research focus and gene expression-based studies and test whether an increased

694 proportion of studies find an hourglass.

a) anurans:
$$\frac{|Rank_{laevis} - Rank_{tropicalis}|}{sqrt(52 + 52)}$$

b) amniotes: (RNAseq)
$$\frac{|Rank_{Pelodiscus} - Rank_{Gallus}|}{sqrt(263 + 263)}$$

c) amniotes: (microarray)
$$\frac{|Rank_{Mus} - Rank_{Gallus}|}{sqrt(320 + 320)}$$

d) tetrapods:
$$\frac{| (Rank_{Mus}/2 + Rank_{Gallus}/2) - (Rank_{lavis}/2 + Rank_{tropicalis}/2) |}{sqrt(33 + 301)}$$

e) vertebrates:
$$\frac{| (Rank_{Mus}/4 + Rank_{Gallus}/4 + Rank_{lavis}/4 + Rank_{tropicalis}/4) - Rank_{Danio}|}{sqrt(79 + 432)}$$

Supplementary Figure S2. To characterize expression conservation of each gene at each
embryonic stage cluster at each node of the phylogeny we calculated the similarity in rank
expression scaled by the divergence times of the included species.

Supplementary Figure S3. Enrichment or depletion of gene expression conservation patterns at
available evolutionary nodes amniotes (A), tetrapods (B), and anurans (C) for microarray (1: left)
and RNA-seq (2: right). Proportion of patterns determines the expected proportion of genes for
each conservation pattern and are shown as associated treemap plots (Tennekes, 2017). The
enrichment/depletion values from permutation analysis (1000 iterations) are shown as overlaying
violin plots. For all plots colors indicate the expression conservation pattern. *** indicate
significance at $p = 0$.