1 **The genome of the zoonotic malaria parasite *Plasmodium simium* reveals**
2 **adaptations to host-switching**
3

Tobias Mourier[1], Denise Anete Madureira de Alvarenga[2], Abhinav Kaushik[1], Anielle de Pina-
Costa[3,4,5§], Francisco J. Guzmán-Vega[6§], Olga Douvropoulou[1§], Qingtian Guan[1§], Sarah
Forrester[7], Filipe Vieira Santos de Abreu[3,8], Cesare Bianco Júnior[3,9], Julio Cesar de Souza
Junior[10], Zelinda Maria Braga Hirano[10], Alcides Pissinatti[11], Silvia Bahadian Moreira[11], Maria
de Fátima Ferreira-da-Cruz[3,9], Ricardo Lourenço de Oliveira[3,8], Stefan T. Arold[6,12], Daniel C.
Jeffares[7], Patrícia Brasil[3,4], Cristiana Ferreira Alves de Brito[2], Richard Culleton[13], Cláudio
Tadeu Daniel-Ribeiro[3,9†] & Arnab Pain[1,14,15†]
11

1) Pathogen Genomics Laboratory, Biological and Environmental Sciences and Engineering
(BESE) Division, King Abdullah University of Science and Technology (KAUST), Thuwal,
Saudi Arabia
2) Grupo de Pesquisa em Biologia Molecular e Imunologia da Malária, Fundação Oswaldo
Cruz (Fiocruz), Belo Horizonte/MG, 30190-009, Brazil
3) Centro de Pesquisa, Diagnóstico e Treinamento em Malária (CPD-Mal), Fiocruz, Rio de
Janeiro/RJ, 21040-360, Brazil
4) Laboratório de Pesquisa Clínica em Doenças Febris Agudas, Instituto Nacional de
Infectologia Evandro Chagas, Fiocruz, Rio de Janeiro/RJ, 21040-360, Brazil
5) Centro Universitário Serra dos Órgãos (UNIFESO), Teresópolis/RJ, 25964-004, Brazil
6) Computational Bioscience Research Center, Biological and Environmental Sciences and
Engineering (BESE) Division, King Abdullah University of Science and Technology
(KAUST), Thuwal, Saudi Arabia
7) Department of Biology and York Biomedical Research Institute, University of York,
Wentworth Way, York, YO10 5DD, U.K.
8) Laboratório de Mosquitos Transmissores de Hematozoários. Instituto Oswaldo Cruz (IOC),
Fiocruz, Rio de Janeiro/RJ, 21040-360, Brazil
9) Laboratório de Pesquisa em Malária, IOC, Fiocruz, Rio de Janeiro/RJ, 21040-360, Brazil
10) Universidade Regional de Blumenau (FURB), Centro de Pesquisas Biológicas de Indaial
(CEPESBI)/ Projeto bugio, Blumenau and Indaial, SC, Brazil.
11) Centro de Primatologia do Rio de Janeiro (CPRJ/Inea), 25940-000, Guapimirim, RJ,
Brazil

12) Centre de Biochimie Structurale, CNRS, INSERM, Université de Montpellier, 34090

Montpellier, France

13) Malaria Unit, Department of Pathology, Institute of Tropical Medicine (NEKKEN),

Nagasaki University, 1-12-4 Sakamoto, Nagasaki, 852-8523, Japan

14) Global Station for Zoonosis Control, Global Institution for Collaborative Research and

Education (GI-CoRE), Hokkaido University, N20 W10 Kita-ku, Sapporo, Japan

15) Nuffield Division of Clinical Laboratory Sciences (NDCLS), University of

Oxford, Headington, Oxford, OX3 9DU, UK

§) Contributed equally, arranged alphabetically

†) Corresponding authors

Summary

Plasmodium simium, a malaria parasite of non-human primates in the Atlantic forest region of

Brazil was recently shown to cause zoonotic infection in humans in the region. Phylogenetic

analyses based on the whole genome sequences of six *P. simium* isolates infecting humans

and two isolates from brown howler monkeys revealed that *P. simium* is monophyletic within

the broader diversity of South American *Plasmodium vivax*, consistent with the hypothesis

that *P. simium* first infected non-human primates as a result of a host-switch from humans

carrying *P. vivax*. We provide molecular evidence that the current zoonotic infections of

people have likely resulted from multiple independent host switches, each seeded from a

different monkey infection. Very low levels of genetic diversity within *P. simium* genomes

and the absence of *P. simium-P. vivax* hybrids suggest that the *P. simium* population emerged

recently and has subsequently experienced a period of independent evolution in Platyrrhini

monkeys. We further find that Plasmodium Interspersed Repeat (PIR) genes, Plasmodium

Helical Interspersed Subtelomeric (PHIST) genes and Tryptophan-Rich Antigens (TRAg)

genes in *P. siumium* are genetically divergent from *P. vivax* and are enriched for non-

synonymous single nucleotide polymorphisms, consistent with the rapid evolution of these

genes. Analysis of genes involved in erythrocyte invasion revealed several notable differences

between *P. vivax* and *P. simium*, including large deletions within the coding region of the

Duffy Binding Protein 1 (DBP1) and Reticulocyte Binding Protein 2a (RBP2a) genes in *P.*

simium. Genotyping of *P. simium* isolates from non-human primates (NHPs) and zoonotic

human infections showed that a precise deletion of 38 amino acids in DBP1 is exclusively

present in all human infecting isolates, whereas non-human primate infecting isolates were

polymorphic for the deletion. We speculate that these deletions in the parasite-encoded key

erythrocyte invasion ligands and the additional rapid genetic changes have facilitated zoonotic

transfer to humans. Non-human primate malaria parasites can be considered a reservoir of

potential infectious human parasites that must be considered in any attempt of malaria

elimination. The genome of *P. simium* will thus form an important basis for future functional

characterizations on the mechanisms underlying malaria zoonosis.

Introduction

There are currently eight species of malaria parasites known to cause disease in humans;

Plasmodium falciparum, *Plasmodium vivax*, *Plasmodium malariae*, *Plasmodium ovale*

curtisi, *Plasmodium ovale wallikeri*, *Plasmodium knowlesi*, *Plasmodium cynomolgi* and

Plasmodium simium. The latter three species are more commonly parasitic on non-human

primates and have only relatively recently been shown to infect humans [1-3].

As interventions against human malaria parasites, particularly *P. falciparum* and *P. vivax,*

continue to reduce their prevalence, the increasing importance of zoonotic malaria is

becoming apparent. In countries currently moving towards the elimination of malaria, the

presence of populations of potentially zoonotic parasites in non-human primates constitutes a

significant obstacle.

The propensity of malaria parasites to switch hosts and the consequences of this for human

health are underlined by the fact that both *P. vivax* and *P. falciparum* first arose as human

pathogens as the result of host switches from great apes in Africa [4-6]. As contact between

humans and the mosquitoes that feed on non-human primates increases due to habitat

destruction and human encroachment into non-human primate habitats [7], there is increasing

danger of zoonotic malaria transmission leading to the emergence of novel human malaria

pathogens. Understanding how malaria parasites adapt to new hosts and new transmission

environments allows assessment of the risks posed by novel zoonotic malaria outbreaks.

The clinical epidemiology of zoonotic malaria varies according to the parasite species involved and the demographics of the human-host population infected. Severe and lethal outcomes have been reported in people infected with *P. knowlesi* in Malaysia [8], whilst infection with *P. cynomolgi* in the same region appears to cause moderate/mild clinical symptoms [9]. Interestingly, both *P. knowlesi* and *P. cynomolgi* infections in the Mekong region appear less virulent than in Malaysia, and are often asymptomatic [3,10], and this may be due to the relative virulence of the parasite strains circulating there and/or differences in the susceptibility of the local human populations. As the parasites of non-human primates have co-evolved with and adapted to their monkey hosts, it is impossible to predict their potential pathogenesis in zoonotic human infections. The virulence of *P. falciparum*, for example, has been attributed to its relatively recent emergence as a human pathogen [11], which appears to have occurred following a single host transfer from a gorilla in Africa [5].

Eighty nine percent of the malaria infections in Brazil are caused by *P. vivax,* with over 99% of these cases occurring in the Amazonian region. This region accounts for almost 60% of the area of Brazil, and is home to 13% of the population (https://www.ibge.gov.br/). Of the 0.4% of cases registered outside the Amazon, around 90% occur in the Atlantic Forest, a region of tropical forest that extends along the Atlantic coast of Brazil, and are caused by an apparently mild, vivax-like malaria parasite transmitted by *Anopheles (Kerteszia) cruzii*, a mosquito species that breeds in the leaf axils of bromeliad plants [12].

Following a malaria outbreak in the Atlantic Forest of Rio de Janeiro in 2015/2016, it was shown that these infections were caused by the non-human primate malaria parasite *P. simium* [1]. DNA samples collected both from humans and non-human primates (NHPs) in the same region shared identical mitochondrial genome sequences, distinct from *P. vivax* isolates from anywhere in the world and identical to that of a *P. simium* parasite isolated from a monkey in the same region in 1966, and to all isolates of *P. simium* recovered from NHPs since [13,14].

It was previously thought that *Plasmodium vivax* became a parasite of humans following a host switch from macaques in Southeast Asia, due to its close phylogenetic relationship with a clade of parasites infecting monkeys in this region and due to the high genetic diversity among *P. vivax* isolates from Southeast Asia [15]. We now know, however, that it became a human parasite following a host switch from great apes in Africa [6]. It is likely that it was introduced to the Americas by European colonisers following Columbus' journey to the New

World in 1492. Indeed, present-day *P. vivax* in South America is closely related to a strain of the parasite present, historically, in Spain [16]. The genetic diversity of extant *P. vivax* in the Americas suggests multiple post-Columbian colonising events associated with the passage of infected people from various regions throughout the world [17]. There is some evidence to suggest that *P. vivax* parasites may also have been introduced to South America in pre-Columbian times, and may have contributed to the extensive genetic diversity of the parasite on this continent [17].

Plasmodium simium, a parasite of various species of Platyrrhini monkeys whose range is restricted to the Atlantic Forest of south and southeast Brazil [18], is genetically and morphologically similar to *P. vivax* [1,19-22]. Based on this similarity, it appears likely that *P. simium* originated as a parasite of monkeys in Brazil following a host switch from humans carrying *P. vivax*. The recent 2015/2016 outbreak of *P. simium* in the local human population of Rio de Janeiro's Atlantic Forest raises questions about the degree of divergence that has occurred between *P. vivax* and *P. simium*, and whether adaptation to monkeys has led to the evolution of a parasite with clinical relevance to human health that differs from that of *P. vivax*.

It is unclear whether the current outbreak of *P. simium* in the human population of Rio de Janeiro was the result of a single transfer of the parasite from a monkey to a human and its subsequent transfers between people, or whether multiple independent host switches have occurred, each seeded from a different monkey infection. Furthermore, the degree and nature of adaptation to a non-human primate host and a sylvatic transmission cycle that has occurred in *P. simium* following its anthroponotic origin is of relevance to the understanding of how malaria parasites adapt to new hosts. It is also of interest to determine whether the current, human-infecting *P. simium* parasites have recently undergone changes at the genomic level that have allowed them to infect people in this region, as it has previously been suggested that *P. simium* has historically lacked the ability to infect man [23].

In order to resolve these questions, and so to better understand the epidemiology and natural history of this emerging zoonotic parasite, we analysed the whole genome sequences of *P. simium* parasites isolated from both humans and non-human primates in the Atlantic Forest region of Rio de Janeiro.

Results

Genome assembly and phylogeny

From a single *P. simium* sample collected from Rio de Janeiro state in 2016 [1] short read sequences were obtained and assembled into a draft genome (see Supplementary Materials). The assembled genome consists of 2,192 scaffolds over 1kb with a combined size of 29 Mb (Table S1). Two scaffolds corresponding to the apicoplast and mitochondrial organelles are also identified (Figure S1). Gene content analysis showed an annotation completeness comparable to previously published *Plasmodium* assemblies (Figure S35). A phylogenetic tree constructed from 3,181 of 1:1 orthologs of the annotated *P. simium* protein-coding genes with *Plasmodium vivax, P. cynomolgi, P. coatneyi, P. knowlesi, P. malariae, P. falciparum, P. reichenowi,* and *P. gallinaceum* confirmed that *P. simium* is very closely related to *P. vivax* (Figure S2).

P. simium-P. vivax diversity analysis

To detect single nucleotide polymorphisms (SNPs) within the *P. vivax*/*P. simium* clade, short Illumina paired-end sequence reads were mapped onto the *P. vivax* P01 reference genome [24]. Reads were collected from eleven human *P. simium* samples, two monkey *P. simium* samples, two *P. vivax* samples from Brazilian Amazon, and a range of *P. vivax* strains representing a global distribution retrieved from the literature [25]. Including only SNPs with a minimum depth of five reads, a total of 232,780 SNPs were initially called across 79 samples. Sixteen samples were subsequently removed from further analysis primarily due to low coverage resulting in a total of 63 samples (Table S2, Table S3). Few SNP loci are covered across all samples, and to enable diversity analysis, we restrict all further analysis to the 124,968 SNPs for which data is available from at least 55 samples (Figure S3).

P. simium-P. vivax population analysis

A Principal Component Analysis (PCA) plot constructed from these genome-wide SNP loci showed a clear separation between American and Asian *P. vivax* samples as well as a distinct grouping of *P. simium* samples (Figure S4). The latter observation suggesting that both human and monkey *P. simium* samples form a single population that is genetically differentiated from other American *P. vivax* populations. A similar pattern is observed when performing a multidimensional scaling analysis of the SNP data (Figure S5). To enable a

phylogenetic approach, we constructed an alignment from the 124,968 SNP sites. In the

resulting phylogenetic tree, *P. vivax* strains generally clustered according to their geographical

origin, and the Asian and American samples were clearly separated (Figure 1A, a tree with

sample IDs is available in Figure S6). *P. simium* samples clustered as a monophyletic group

with Mexican vivax samples (Figure 1A), consistent with a recent American origin for *P.*

simium.

To examine whether the *P. simium* isolates we obtained were part of a continuous population

with local *P. vivax,* we examined population ancestry with the ADMIXTURE program [26]

(Figure S7). This analysis is consistent with the PCA and MDS analysis (Figure S4 & Figure

S5) and the phylogenetic analysis of segregating SNPs (Figure 1), showing that *P. simium*

forms a genetically distinct population of *P. vivax*. The absence of *P. simium-P. vivax* hybrids

(introgression events) suggests that *P. simium* has undergone a period of independent

evolution in Platyrrhini monkeys.

***P. simium* genetic differentiation from *P. vivax* is enhanced in host-parasite interacting**

genes

To characterise the *P. simium* population further, we estimated the nucleotide diversity in *P.*

simium and *P. vivax* samples (see Materials and Methods). *P. simium* diversity (genome-

median: 1.3×10^{-4}) is more than five times lower than the diversity observed when comparing

all *P. vivax* samples (genome-median: 7.5×10^{-4}) (Figure 2). Diversity within coding sequences

in *P. vivax* is consistent with previous reports [6]. The median nucleotide diversity between *P.*

simium and *P. vivax* genomes of 8.4×10^{-4} and the low diversity within *P. simium* suggest that

the strains we examined are part of a relatively recent or isolated population.

We then examined the population differentiation over the entire genome using $F_{ST,}$ a measure

of the proportion of ancestry private to a population ($F_{ST}=0$ for completely intermixed

populations, $F_{ST}=1$ for populations with completely independent ancestry). Although our

analysis contains very few samples, F_{ST} estimates can be very accurate if multiple genomic

sites are used [27]. Consistent with phylogenetic and admixture analysis, we observed a high

level of differentiation between human *P. simium* and American *P. vivax* ($F_{ST}=0.46$). For

comparison, the differentiation between vivax from America and vivax from Myanmar and

Thailand (henceforth referred to as 'Asian vivax') is less than half of this ($F_{ST}=0.22$). To

examine whether there were any signals of adaptive change in *P. simium* that may have

occurred during its adaptation in monkeys upon anthroponotic transfer, we calculated the

fixation index for all individual genes. Clearly, the small number of samples renders this

analysis prone to false and incorrect signals, and F_{ST} values for individual genes should be

interpreted with caution. Nevertheless, we attempted to look for general patterns in F_{ST} values

across gene groups.

Amongst the 4,341 *P. vivax* genes with at least one SNP in our data set, we examined the top-

25% of the genes with highest F_{ST} values for enrichment in functional Gene Ontology (GO)

terms or metabolic pathways. No GO terms or pathways were significant at the 0.05 level

after Bonferroni correction (Table S4 & Table S5). Using the *P. falciparum* orthologs instead

– when available – gave similar results (not shown). We next tested if any of the gene families

(Figure S8, Figure S9, Table S6) were associated with high F_{ST} values. Genes belonging to

the Plasmodium Interspersed Repeat (PIR) family involved in antigenic variation [28], the

Plasmodium Helical Interspersed Subtelomeric (PHIST) genes, a family of exported proteins

[29], the merozoite surface proteins MSP7[30], and Tryptophan-rich antigens (TRAg) [31] were

enriched among the genes with high F_{ST} values (binomial distribution, PIR; $p=3.5\times10^{-3}$,

PHIST; $p=4.1\times10^{-4}$, MSP7; $p=0.034$, TRAg; $p=2.5\times10^{-3}$).

As these gene families are involved in parasite-host interactions, the observation of elevated

F_{ST} values may simply reflect a general pattern of rapid genetic divergence in *Plasmodium*

parasites. To test this, we repeated the F_{ST} analysis between American vivax and a selection of

Asian vivax isolates (Myanmar and Thailand samples only). Consistent with the phylogenetic

analysis (Figure 1A) gene F_{ST} was slightly higher overall between simium and American

vivax than between American and Asian vivax samples (Figure 3A). However, none of the

gene families were overrepresented among genes with high F_{ST} (top-25%) between American

and Asian vivax. To further examine if the elevated F_{ST} measures found for PIR, PHIST,

MSP7, and TRAg genes are exclusive to the comparison between simium and American

vivax, we calculated the ratio between the two F_{ST} measurements ('simium versus American

vivax' and ' American versus Asian vivax') (Figure 3B). The ratios for PIR, PHIST and TRAg

genes were significantly higher than observed for the remaining genes (Figure 3C), whereas

ratios for MSP7 genes were not (Mann-Whitney U, $p=0.12$). Although the *P. simium* and the

P. vivax P01 both genomes encode a high number of the gene family members, our analysis is

restricted to the *P. vivax* genes for which our *P. simium* short read sequences can map. For

example, only 408 out of the 1209 *P. vivax* PIR genes have coverage from *P. simium* reads

across at least 80% of their gene length (Figure S10). Further, an even smaller number of
these genes have detectable SNPs between simium and American vivax samples and are
included in the analysis (numbers shown below Figure 3B).

To test if the sequence redundancy among gene family loci could result in spurious cross-
mapping of short sequence reads we specifically tested the quality of SNPs in gene families,
and SNPs residing in gene families showed no signs of decreased calling, mapping, or base
qualities compared to other SNPs (Figure S11).

A phylogenetic analysis of PIR, PHIST and TRAg proteins harbouring genomic SNPs
revealed no apparent association between certain protein phylogenetic sub-groups and high
F_{ST} ratios (Figure S12-S14), consistent with a subtle signature of polygenic adaptation in
these gene families.

When testing all exported genes and genes involved in invasion and exported genes (Table
S8), the observed F_{ST} ratios were not significantly different from the background (Mann-
Whitney U, p=0.5473). Hence, the differences in F_{ST} observed for PIR, PHIST and TRAg
genes are not a general phenomenon amongst the genes known to be involved in interactions
with the host and red cell invasion.

The observed skew towards higher F_{ST} values when comparing simium and American vivax
(Figure 3A) could be a result of an inherent diversity between different American vivax
populations potentially stemming from multiple introductions of *P. vivax* to the American
continent [17]. To test if such founder effects and subsequent population bottlenecks could
explain the observations, we repeated the F_{ST} analysis using only Mexican vivax samples as
American representatives. Four Mexican samples (SRS693273, SRS694229, SRS694244,
SRS694267) were used. These clustered close together in both the SNP phylogeny (Figure 1)
and in the PCA and MDS plots (Figure S4 & Figure S5), and are assumed to share a recent
evolutionary history. This analysis revealed the same pattern of elevated F_{ST} values between
simium and Mexican vivax, and PIR genes did again display significantly higher F_{ST} ratios
(Figure S15). Although PHIST and TRAg genes also showed higher F_{ST} ratios, these were no
longer significant (Figure S15). We therefore conclude that the observed higher F_{ST} values
between simium and American vivax PIR genes are not solely a result of diversity within
American vivax populations, but rather appear specific to comparisons with *P. simium*.

Adaptive changes in PIR genes would be expected to produce stronger genetic divergence in

non-synonymous codon positions. To examine this, we divided genic SNPs into synonymous

and non-synonymous changes. In PIR genes, there are 353 non-synonymous and 185

synonymous SNPs (non-synonymous to synonymous SNP ratio = 1:1.91). Similarly, in

PHIST and TRAg genes we find 220 and 103 non-synonymous, respectively, and 67 and 41

synonymous SNPs, respectively (PHIST ratio = 1:3.28, TRAg ratio = 1:2.51). In all other

genes, the ratio between non-synonymous and synonymous SNPs is 1:1.49. Hence, the

proportion of non-synonymous SNPs in PIR, PHIST and TRAg genes is significantly higher

than in all other genes (chi-square, PIR; p= 0.0073, PHIST; p=9.4×10^{-9}, TRAg; p=0.0054).

Our finding that PIR, PHIST and TRAg genes overall display markedly higher F_{ST} values

between simium and vivax suggest that these gene groups are enriched for private alleles

consistent with natural selection acting upon these genes subsequent to the split between *P.*

simium and *P. vivax*.

***P. simium* invasome components**

In invading *P. vivax* merozoites, binding to host red blood cells is mediated by two gene

families: Duffy Binding Proteins (DBPs) bind the Duffy Antigen Receptor for Chemokines

(DARC) [32,33], which is present on both host normocytes and reticulocytes, whereas

Reticulocyte Binding Proteins (RBPs) preferentially bind host reticulocytes [34-36]. Recently, the

reported protein structure of *P. vivax* RBP2b revealed the evolutionary conservation of

residues involved in the invasion complex formation [36]. Two DBPs, DBP1 and DBP2, are

present in *P. vivax* P01 (Table S9). RBPs can be divided into three subfamilies, RBP1, RBP2,

and RBP3 [37]. The *P. vivax* P01 genome encodes 11 RBPs (including the reticulocyte binding

surface protein, RBSA), of which three are pseudogenes (Table S9).

The *P. vivax* DBP and RBP were used to search the *P. simium* proteins, resulting in the

detection of the two DBP proteins and RBP1a, RBP1b, RBP2a, RBP2b, and RBP3 and failure

to detect RBP2c and RBP2d (Figure 4; Table S9; Figure S16; Figure S17) across all

sequenced *P. simium* samples. As in other *P. vivax* genomes, the *P. simium* RBP3 is a

pseudogene [38], indicating that the pseudogenization event happened prior to the split between

P. vivax and *P. simium*.

To determine whether the apparent absences of individual RBP genes in *P. simium* were due

to incomplete genome assembly, we examined the coverage of *P. simium* reads mapped onto

P. vivax RBP gene loci. As expected, no *P. simium* coverage was observed at the RBP2c,

RBP2d, and RBP2e genes in *P. simium* samples, including the previously published CDC

strain deposited in GenBank (accession ACB42432) [39] (Figure S18).

Coverage of mapped reads across invasome gene loci revealed no apparent elevated coverage

in genes compared to their flanking genomic regions, which would otherwise be expected if

the *P. simium* genome contained multiple (duplicated) copies of non-assembled invasion

genes (Figure S19). Similarly, analysis of *P. simium* read mapping data using the DELLY

software [40] showed no large genomic duplications and deletions events occurring at loci

harbouring invasion genes (Table S10) although numerous short indels were detected within

protein-coding genes (Table S11).

Structural variation in *P. simium* Duffy Binding Protein 1

The simium assembly revealed that the invasion gene DBP1 contains a large deletion within

its coding sequence (Figure 4) (a full alignment is provided in Figure S20). Intriguingly, the

previously published *P. simium* CDC strain (originally isolated in 1966) DBP1 does not

contain the deletion ('simium CDC' in Figure 4B). A haplotype network confirms that this

previously published DBP1 gene is indeed a *P. simium* sequence (Figure S22), and the SNP

analyses consistently assign the CDC strain to the simium cluster (Figure 1, Figure S4, and

Figure S5). Compared to the *P. vivax* P01 reference genome the SalI reference harbours a 27

base pair deletion in DBP1, in contrast to the 115 bp deletion observed in all *P. simium*

samples isolated from humans (Figure 4). This deletion is also present in most *P. vivax*

isolates (Figure S23). Additional deletion patterns exist among isolates, and in a few cases

multiple versions are detected within samples (Figure S23).

The presence of repetitive sequences within the DBP1 gene could potentially result in

aberrant assembly across the DBP1 locus, which could appear as an apparent deletion in

subsequent bioinformatic analysis. We tested this possibility and the DBP1 gene does not

harbour any noticeable degree of repetitiveness (Figure S24). Several read mapping analyses

confirmed that the *P. simium*-specific 115 bp deletion was not an assembly artefact (Figure

S25-S27).

We next designed primers for PCR amplification of a genomic segment across the deleted

region in the *P. simium* DBP1 gene and tested the occurrence of these deletion events in a

range of *P. vivax* and *P. simium* field samples from Brazil. All *P. vivax* samples tested by

PCR produced bands consistent with absence of the deletion whereas all samples from

human-infecting *P. simium* produced bands consistent with the presence of the precise 115 bp

deletion (Figure S28, top & middle). Interestingly, non-human primate (NHP)-infecting *P.*

simium isolates were a mix of samples with and without deletions (Figure S28, bottom). If the

P. simium-specific deletion in DBP1 is a prerequisite for the ability to infect humans this

suggests that only a subset of NHP-infecting *P. simium* parasites currently possess the DBP1

allele required for zoonotic transfer to humans.

A large, additional deletion was observed in the *P. simium* RBP2a gene, the presence of

384
385   which was also supported by read mapping and PCR analysis (Figure 3, Figure S29-S32).

Potential structural implications of the deletion in DBP1 and RBP2a

We next investigated if the observed deletions render DBP1 and RBP2a dysfunctional. DBP1

contains a large extracellular region, which includes the N-terminal DBL region which is

mediating the association with DARC in *P. vivax* [41], followed by a largely disordered region

and a cysteine-rich domain (Figure 4c). DBP1 has a single-pass transmembrane helix and a

short cytoplasmic tail. The deletion observed in the human-infecting *P. simium* only affects

the disordered region, leaving the flanking domains intact. We produced homology models of

the DBL domains from the *P. vivax* strain P01, the human-infecting *P. simium* strain AF22,

and the *P. simium* CDC strain, based on the crystal structure of the >96% identical DBL

domain of *P. vivax* bound to DARC (PDB ID 4nuv). Whereas no significant substitutions

were found in the DBL domain between both *P. simium* sequences, our analysis showed that

residue substitutions between *P. simium* and *P. vivax* DBL domains cluster in proximity of the

DARC binding site (Figure S33). Based on our models, these substitutions are unlikely to

negatively affect the association with DARC, supporting that the DBL domains of both *P.*

simium would be capable of binding to human DARC. Hence, the human-infecting *P. simium*

sequence encodes for a protein that retains the capacity to bind to human DARC, but would

have the interacting domain positioned closer to the membrane than in the monkey-infecting

CDC strain.

The deletion we detected in human-infecting RBP2a was more severe, resulting in the loss of

1003 residues. These residues are predicted to form a mostly α-helical extracellular stem-like

structure that positions the reticulocyte binding domain away from the membrane (Figure 4d).

However, given that the deletion does neither affect the transmembrane region, nor the

receptor-binding domain, our analysis supports that the resulting truncated RBP2a protein can

still associate with the human receptor, but that the binding event would occur closer to the

plasmodium membrane.
412

Discussion

We present the genome of *Plasmodium simium*, the eighth malaria parasite species known to

infect humans in nature. In recent evolutionary time, *P. simium* has undergone both

anthroponosis and zoonosis making it unique for the study of the genetics underlying host-

switching in malaria parasites. The genome content confirmed the close phylogenetic

relationship between *P. simium* and *P. vivax*, and further analyses on single nucleotide

divergences support a very recent American origin for *P. simium*. This recent split between *P.*

vivax and *P. simium* precludes detection of genes under positive evolution [42], and we have

instead performed a general analysis of population differentiation between extant *P. simium*

and *P. vivax* isolates using F_{ST}. We find that members of three gene families involved in

antigenic variation, PIR, PHIST and TRAg, show significantly elevated F_{ST} levels between *P.*

simium and *P. vivax*. As higher F_{ST} values amongst these genes are not observed between

global vivax populations, their genetic differentiation appears to be associated with host-

switching between human and monkey.

429
430

Two proteins involved in host invasion, DBP1 and RBP2a, were found to harbour extensive

deletions in *P. simium* compared to *P. vivax*. Interestingly, experimental analysis of *P. simium*

samples revealed that isolates from human hosts all carried the DBP1 deletion, whereas

isolates from non-human primates displayed both absence and presence of the deletion. This

DBP1 deletion is not present in the *P. simium* isolated from a brown howler monkey in the

1960s, which was previously shown to be incapable of infecting humans [23], although some

degree of laboratory adaptation of this parasite may have affected its genome. However, this

deletion is also absent in *P. vivax*, so cannot in itself explain the ability of *P. simium* to infect

humans in the current outbreak. It is possible, however, that this deletion is required for *P. simium* to invade human red blood cells given the alterations that have occurred elsewhere in its invasome following adaptation to non-human primates since the split between *P. simium* and its human-infecting *P. vivax* ancestor.

Invasome proteins are obvious candidates for genetic factors underlying host-specificity, and an inactivating mutation in a *P. falciparum* erythrocyte binding antigen has recently been shown to underlie host-specificity [43]. Traditionally, functional studies on invasome proteins have focused on domains known to bind or interact directly with the host. Although the *P. simium*-specific DBP1 and RBP2a deletions reported here do not cover known structural motifs, these deletions could nevertheless affect host cell recognition as disordered protein regions have known roles in cellular regulation and signal transduction [44]. Further, a shorter, less flexible linker between the plasmodium membrane and the receptor-binding DBP1 domain may favour a more rigid and better oriented positioning of the dimeric DBP1, enhancing its capacity to engage the human receptor.

Phylogenetic analysis of the *P. simium* clade gives the geographical location of its most closely related *P. vivax* strain as Mexico, and not Brazil. In imported populations, the relationship between geographical and genetic proximity may be weak. Multiple introductions of diverse strains from founder populations may occur independently over large distances, so that two closely related strains may be introduced in distantly located regions. It may be postulated that there occurred the introduction of strains of *P. vivax* to Mexico from the Old World that were closely related, due to similar regions of origin, to strains introduced to the Atlantic Forest which went on to become *P. simium* in New World monkeys. Strains from a different point of origin were introduced to the Amazonian region of Brazil. This hypothesis necessitates reproductive isolation of the *P. simium* clade from the Brazilian *P. vivax* parasites following their initial introduction; an isolation that would be facilitated, presumably, by their separate host ranges.

Due to uncertainties regarding the number of individual genomes that were transferred during the original host switch from man to NHPs that resulted in the formation of the *P. simium* clade, it is impossible to perform dating analyses to determine a time for the split between *P. vivax* and *P. simium* with which we can be confident. The phylogeny shown in figure 1 is consistent with the hypothesis that all present-day *P. vivax*/*P. simium* originated from a now

extinct Old World population. The most parsimonious explanation for this is that today's New World *P. vivax/P. simium* originated from European *P. vivax*, which was itself a remnant of the original Eurasian/African *P. vivax* driven to extinction in Africa by the evolution of the Duffy negative condition in the local human populations, and from Europe by malaria eradication programmes in the latter half of the twentieth century. This hypothesis is supported by the evidence of a close relationship between historical Spanish *P. vivax* and South American strains of the parasite [16], and by previous analyses of the mitochondrial genome [45]. Therefore, we postulate that the host switch between humans and non-human primates that eventually led to establishment of *P. simium* in howler monkeys must have occurred subsequent to the European colonisation of the Americas, within the last 600 years.

We find no evidence from the nuclear genome, the mitochondrial genome or the apicoplast genome that any of the *P. vivax /P. simium* strains from the New World considered in our analyses are more closely related to Old World parasites than they are to each other, as previously contended [46]. However, our nuclear genome phylogeny is based on genome-wide SNPs, and so represents an "average" phylogeny across the genome. This cannot be considered to reflect a true history of parasite ancestry due to the effects of recombination, and it is possible that trees produced from individual genes might reveal different phylogenetic relationships.

Given the limited genetic diversity amongst the *P. simium* isolates considered here compared to that of *P. vivax*, it is almost certain that the original host switch occurred from humans to NHPs, and not the other way around [22]. Similarly, the larger amount of genetic diversity in the current NHP-infecting *P. simium* compared to those *P. simium* strains isolated from humans (as indicated by the higher degree of DBP1 polymorphism in the NHP-infecting *P. simium* compared to the strains infecting humans), suggests that humans are being infected from a pool of NHP parasites in a true zoonotic manner, as opposed to the sharing of a common parasite pool between humans and NHPs

The biological definition of a species is a group of organisms that can exchange genetic material and produce viable offspring. We have no way of knowing whether this is the case for *P. vivax* and *P. simium*, and genetic crossing experiments would be required to resolve this question. Our phylogenetic analysis, however, clearly shows *P. simium* forming a clade on its own within the broader diversity of *P. vivax*, and that strongly suggests, given what we know

about its biology, that allopatric speciation has been/is occurring. *Plasmodium simium* appears to have been reproductively isolated from other strains of *P. vivax* for long enough for significant genetic differentiation to occur (F_{ST} = 0.46), with some invasome genes showing even higher genetic differentiation.

Plasmodium simium is currently recognised as a species separate from *P. vivax*; it has been well characterised and described in the literature, and there is a type specimen available, with which all the strains sequenced here cluster in one monophyletic group. Therefore, we cannot at present overturn the species status of *P. simium* in the absence of conclusive proof from crossing experiments.

In summary, the recent outbreak of human malaria in the Atlantic Forest of Rio de Janeiro underlines the impact of zoonotic events on human health. In this sense non-human primate malaria parasites can be considered a reservoir of potential infectious human parasites that must be considered in any attempt of malaria eradication. Little is known about the genetic basis for zoonosis, yet the presented genome sequence of *P. simium* suggests a deletion within the DBP1 gene as a possible facilitator of zoonotic transfer. The genome of *P. simium* will thus form an important basis for future functional characterizations on the mechanisms underlying malaria zoonosis.

Methods

Sample Collection and Preparation

Human and primate samples of *P. simium* were collected and prepared as part of a previous study [1,14]. Additionally, two *P. vivax* samples from the Amazon area of Brazil were also collected from human patients (Table S2). All participants provided informed written consent. The *P. simium* CDC (Howler) strain (Catalog No. MRA-353) from ATCC was obtained via the BEI Resources Repository in NIAID-NIH (https://www.beiresources.org/).

DNA extraction and sequencing

DNA was extracted as described[1]. The genomic DNA for each sample was quantitated using the Qubit® 2.0 Fluorometer and was used for library preparation. The DNA for intact samples was sheared using a Covaris E220 DNA sonicator to fragments of 500bp. The DNA libraries

540 for intact samples were made using the TruSeq Nano DNA Library Prep kit (Illumina),

541 whereas the DNA libraries for degraded samples were made using Ovation Ultralow Library

542 System V2 kit (Nugen), according to the manufacturers' instructions. The amplified libraries

543 were stored in -20 °C. The pooled libraries were sequenced in an Illumina HiSeq4000

544 instrument (2 x 150 bp PE reads) (Illumina). A PhiX control library was applied to the

545 sequencing run as a base balanced sequence for the calibration of the instrument so that each

546 base type is captured during the entire run. Raw sequence reads were submitted to FastQC

547 v.0.11.5 and the quality score of the sequences generated was determined. Samples AF22,

548 AF26, AF36 were additionally sequenced and scaffolded by PacBio RS II platform (Pacific

549 Biosciences, California, US) using a SMRT library. Genomic DNA from the *P. vivax* samples

550 was extracted from filter paper as previously described [47].

552 **Illumina reads preparation and mapping**

553 Fastqc v 0.11.6 (http://www.bioinformatics.babraham.ac.uk/projects/fastqc) was used to

554 evaluate the quality of Illumina reads. Illumina adapters were removed, and reads were

555 trimmed using the trimmomatic v0.33 [48] software with the following conditions:

556 *LEADING:20 TRAILING:20 SLIDINGWINDOW:4:20 MINLEN:36*

557 To exclude human reads from our analysis, trimmed reads were mapped against the human

558 reference genome (v. hg38) and the *Plasmodium vivax* strain P01 reference genome (v. 36)

559 from PlasmoDB (www.plasmodb.org) with bowtie2 (v 2.3.3.1) [49]. Reads mapping against the

560 human genome were removed from further analysis.

562 **Genome Assembly**

564 *P. simium* sample AF22 was selected for genome assembly based on read quality and

 coverage. After removal of human contaminants, Illumina reads were assembled into contigs

 using the Spades (v 3.70) assembler [50]. Contigs assembled into scaffolds running SSPACE (v

3.0) [51] for 15 rounds and gaps filled with Gapfiller (v 1.10) [52]. Scaffolds were subsequently

correct with Illumina reads using the Pilon (v 1.22) software [53]. Blobtools (v 1.0) (DOI:

10.5281/zenodo.845347) [54] was used to remove any residual contaminant scaffolds. Genome

size and GC content is in line with that of *P. vivax* species (Table S1). Genomic scaffolds

representing the mitochondrial and apicoplast genome were identified through blastn searches

against the corresponding *P. falciparum* and *P. vivax* sequences (Figure S1). The *P. simium*

mitochondrial genome was aligned against a range of previously published *P. vivax* and *P.*

simium mitochondrial genomes [55,56]. A gap-filled region in the alignment where the distal

parts of the *P. simium* scaffold were merged was manually deleted. A minimum spanning
haplotype network was produced using PopART [57,58] confirming the authenticity of the *P.*
simium mitochondrial genome (Figure S34).

Genome Annotation

Two approaches were used to annotate the reference *P. simium* AF22 genome. Firstly, the
Maker pipeline (v 2.31.8) [59] was run for two rounds, using ESTs and protein evidence from *P.*
vivax and *P. cynomolgi* strain B and *P. falciparum* to generate Augustus gene models.
Secondly, a separate annotation was produced using the Companion web server [60].
Companion was run using the *P. vivax* P01 reference assembly and default parameters. Basic
annotation statistics are provided in Table S1. The relatively low number of genes (5966) is
lower due to the fragmented and incomplete nature of the *P. simium* assembly (Table S1).
Gene content was estimated using BUSCO [61,62] (v3.0) revealing an annotation completeness
comparable to other Plasmodium genome assemblies (Figure S35).

PlasmoDB Genome References and Annotations

Genome fasta files, as well as annotated protein and CDS files were obtained from PlasmoDB
for the following species: *P. gallinaceum* 8A, *P. cynomolgi* B and M, *P knowlesi* H, *P.*
falciparum 3D7, *P. reichenowi* G01, *P. malariae* UG01, *P. ovale curtisi* GH01, *P. coatneyi*
Hackeri, *P. vivax* P01 and *P. vivax* SalI. For each species, version 36 was used.

Orthologous group determination

Amino-acid sequences-based phylogenetic trees were prepared using protein sequences from
the *P. simium* annotation, as well as the protein annotations from 10 malaria species
downloaded from PlasmoDB: *P. vivax* P01, *P. cynomolgi* B, *P. knowlesi* H, *P. vivax*-like
Pvl01, *P. coatneyi* Hackeri, *P. falciparum* 3D7, *P. gallinaceum* 8A, *P. malariae* UG01, *P.*
ovale curtisi GH01, and *P. reichenowi* G01. *P. vivax*-like from PlasmoDB version 43, all
other annotations from version 41. A total of 3181 1:1 orthologous genes were identified
using the Proteinortho (v 6.0.3) software [63]. Approximately 88% of the predicted genes in *P.*
simium have orthologs in the *P. vivax* P01 (Figure S36).

Indels in genes

Structural variations were detected using DELLY [40] (v 0.7.9). Coordinates of structural
rearrangements their nearest genes are listed in Table S10. Shorter indels were detected from

soft-clipping information in read mapping (using the '-i' option in DELLY)(Table S11). Indels
in exons were further compared to indels present in the *P. simium* AF22 genome assembly,
suggesting a high false discovery rate of DELLY indels compared to assembly indels (Figure
S37).
612

613   **Protein phylogeny**
614   Protein sequences were aligned using mafft (v 7.222) [64] and alignments were subsequently
615   trimmed with trimAl (v 1.2rev59) [65] using the heuristic 'automated1' method to select the best
616   trimming procedure. Trimmed alignments were concatenated and a phylogenetic tree was
617   constructed using RAxML (v 8.2.3) [66] with the PROTGAMMALG model.
618

SNP calling and analysis
Short sequence reads from 15 simium samples (13 human and 2 monkey) and two vivax
samples, all from this study (Table S2), were aligned against a combined human (hg38) and
P. vivax (strain P01, version 39) genome using NextGenMap (v0.5.5) [67]. This was similarly
done for 30 previously published *P. vivax* strains [25] and the Sal1 reference. These data sets
were downloaded from ENA (https://www.ebi.ac.uk/ena) (Table S3).
Duplicate reads were removed using samtools (v 1.9) [68] and the filtered reads were realigned
using IndelRealigner from the GATK package (v 4.0.11) [69]. SNPs were called independently
with GATK HaplotypeCaller and freebayes (v 1.2.0) [70], keeping only SNPs with a QUAL
score above 30. The final SNP set were determined from the inter-section between GATK and
freebayes. Allele frequencies and mean coverage across SNP sites are shown in Figure S38.
PCA plot was constructed using plink (v 1.90) [71], and admixture analysis was done with
Admixture (v 1.3.0) [26]. F_{ST} values were estimated from nucleotide data with the PopGenome
R package [72,73] using the Weir & Cockerham method [74]. Non-synonymous and synonymous
SNPs were identified using snpeff [75].
634

SNP phylogeny
Alleles from SNP positions with data in 55 samples were retrieved, concatenated, and aligned
using mafft [64]. Tree was produced by PhyML [76,77] with the GTR substitution model selected
by SMS [78]. Branch support was evaluated with the Bayesian-like transformation of
approximate likelihood ratio test, aBayes [79]. Phylogenetic network was made in SplitsTree [80]
using the NeighborNet network [81].
641

Nucleotide diversity

Conventional tools calculating nucleotide diversity directly from the variant call files assumes

that samples are aligned across the entire reference sequence. But as read coverage across the

reference genome was highly uneven between samples (Figure S38), adjustment for this was

required. Coverage across the reference genome was thus calculated for each sample using

samtools mpileup (v 1.9) [68]. For each comparison between two samples, the nucleotide

divergence was calculated as number of detected bi-allelic SNPs per nucleotide with read

coverage of at least 5X in both samples.

Gene sequence deletions

Exploratory Neighbor-Joining phylogenies produced with CLUSTALW [82,83] and visualized

with FigTree (https://github.com/rambaut/figtree/) after alignment with mafft [82]. Pacbio reads

were aligned using Blasr (v 5.3.2) [84], short Illumina reads using NextGenMap (v0.5.5) [67].

Dotplots done with FlexiDot (v1.05) [85].

Gene families and groups

Exported gene sets were compiled from the literature [86-88]. Invasion genes were retrieved from

[89]. Gene families were assessed in seven Plasmodium genomes (*P. simium*, *P. vivax* SalI, *P.*

vivax P01, *P. vivax-like* Pvl01, *P. cynomolgi* M, *P. cynomolgi* B, and *P. knowlesi* H) using the

following pipeline: For all genomes annotated genes were collected for each gene families.

These 'seed' sequences were used to search all proteins from all genomes using BLASTP and

best hits for all proteins were recorded. For each gene family 'seed' sequences were then

aligned with mafft [64], trimmed with trimAl [65], and HMM models were then built using

HMMer (http://hmmer.org/). For PIR/VIR and PHIST genes, models were built for each

genome independently, for all other gene families a single model was built from all genomes.

These models were then used to search all proteins in all genomes. All proteins with best

BLASTP hit to a 'seed' sequence from a given genome were sorted according to their bit

score. The lowest 5% of hits were discarded and remaining proteins with best hits to a 'seed'

sequence were assigned one 'significant' hit. As all proteins were searched against 'seeds' from

the six annotated genomes (P. simium excluded), a maximum of six 'significant' BLAST hits

could be obtained. Similarly, for each HMM model the bottom 25% hits were discarded and

remaining hits were considered 'significant'. The final set of gene families consists of

previously annotated genes and un-annotated genes with at least two 'significant' hits (either

BLASTP or HMM).

PCR amplification of DBP1 and RBP2a genes

PCR primers were initially designed from alignments between *P. vivax* and *P. simium*

sequences and subsequent tested using Primer-BLAST [90] and PlasmoDB

(www.plasmodb.org). For DBP1, the reaction was performed in 10 μL volumes containing

0.5 μM of each oligonucleotide primer, 1 μL DNA and 5 μL of Master Mix 2x (Promega)

(0.3 units of Taq Polymerase, 200 μM each deoxyribonucleotide triphosphates and 1.5 mM

MgCl2). Samples were run with the following settings: 2 minutes of activation at 95°C,

followed by 35 cycles with 30 seconds denaturation at 95°C, 30 seconds annealing at 57°C

(ΔT=-0.2 °C from 2nd cycle) and 1 minute extension at 72°C, then 5 minutes final extension

at 72°C and hold in 4°C. For RBP2a PCR, the reaction was performed in 10 μL volumes

containing 0.5 μM of each oligonucleotide primer, 1 μL DNA, 0.1 μL PlatinumTaq DNA

Polymerase High Fidelity (Invitrogen, 5U/μL), 0.2 mM each deoxyribonucleotide

triphosphates and 2 mM MgSO$_4$. The PCR assays were performed with the following cycling

parameters: an initial denaturation at 94°C for 1.5 min followed by 40 cycles of denaturation

at 94°C for 15 sec, annealing at 65°C for 30 sec (ΔT=-0.2 °C from 2nd cycle) and extension at

68°C for 3.5 min. The temperature was then reduced to 4 °C until the samples were taken. All

Genotyping assays were performed in the thermocycler Veriti 96 wells, Applied Biosystems,

and the amplified fragments were visualized by electrophoresis on agarose gels (2% for DBP1

and 1% for RBP2a) in 1x TAE buffer (40 mM Tris-acetate, 1 mM EDTA) with 5 μg/ mL

ethidium bromide (Invitrogen) in a horizontal system (Bio-Rad) at 100 V for 30 min. Gels

were examined with a UV transilluminator (UVP - Bio-Doc System).

698
699

To prevent cross-contamination, the DNA extraction and mix preparation were performed in

"parasite DNA-free rooms" distinct from each other. Furthermore, each of these separate

areas has different sets of pipettes and all procedures were performed using plugged pipette

tips. DNA extraction was performed twice on different days. Positive (DNA extracted from

blood from patients with known *P. vivax* infection) and negative (no DNA and DNA extracted

from individuals who have never traveled to malaria-endemic areas) controls were used in

each round of amplification. DNA extracted from blood of a patient with high parasitemia for

P. vivax and DNA of *P. simium* of a non-human primate with an acute infection and

parasitemia confirmed by optical microscopy served as positive controls in the PCR assays.

Primer sequences are provided in Figure S28 and S32.

Structural modelling of DBP1 and RBP2a genes

RaptorX [91] was used for prediction of secondary structure and protein disorder. Homology models for the DBL domain of *P. vivax* P01 strain, *P. simium* AF22, and the previously published CDC *P. simium* strain were produced by SWISS-MODEL [92], using the crystallographic structure of the DBL domain from *Plasmodium vivax* DBP bound to the ectodomain of the human DARC receptor (PDB ID 4nuv), with an identity of 98%, 96% and 96% for *P. vivax*, *P. simium* AF22 and *P. simium* CDC, respectively. QMEAN values were -2.27, -2.04 and -2.03, respectively. The homology model for the reticulocyte binding protein 2 (RBP2a) of *P. vivax* strain P01 was produced based on the cryoEM structure of the complex between the *P. vivax* RBP2b and the human transferrin receptor TfR1 (PDB ID 6d05)[36], with an identity of 31% and QMEAN value of -2.46. The visualization and structural analysis of the produced models was done with PyMOL (https://pymol.org/2/).

Data availability

The reference genome assembly and short sequence reads have been uploaded to European Nucleotide Archive (https://www.ebi.ac.uk/ena/) under the Study accession number PRJEB34061.

Acknowledgements

We thank Prof. Xin-zhuan Su at the National Institute of Allergy and Infectious Diseases, NIH, for invaluable help in obtaining the parasite gDNA from the BEI Resources; Sidnei Silva and Graziela Zanini, for assistance on the parasitological diagnosis of the human samples; Aline Lavigne and Larissa Gomes for undertaking the PCR for *P vivax*; Alcides Pissinatti and Silvia Bahadian Moreira for the facilities provided at the Primate Centre of Rio de Janeiro; Orzinete Rodrigues Soares for non-human primates' blood slides'; Marcelo Quintela, Waldemir Paixão Vargas, Carlos Alberto C. da Silva, Alexandre B. de Souza, Vicente Klonowski, Romenique L. Araújo, Luis R. Nogueira, Fernando Barreto, Ana L. Quijada, Luiz P.P. Silva, Gelson Medeiros, Adilson B. Ramos, Marcilene B. Ramos, Carlos A.A. Júnior, Paulo G. Barbosa, Sérgio F. Fragoso, Adilson R. Silva, Cecília Cronemberger, Marcelo Rheingantz, Leonardo Nascimento and João Marins for the field support; *Grupo Técnico de Vigilância de Arboviroses* (GT-Arbo – Brazilian Ministry of Health) for field and

material supports; and Cassio Leonel Peterka from The Brazilian Ministry of Health for malaria epidemiological data. The following reagent was obtained through BEI Resources, NIAID, NIH: *Plasmodium simium*, Strain Howler, MRA-353, contributed by William E. Collins. The work was supported by the King Abdullah University of Science and Technology (KAUST) through the baseline fund BRF1020/01/01 to AP and BAS/1/1056-01-01 to STA, and the Award No. URF/1/1976-25 from the Office of Sponsored Research (OSR). The field work in the Atlantic Forest and laboratory analysis in Brazil received financial support from the Secretary for Health Surveillance of the Ministry of Health through the Global Fund (agreement IOC-005-Fio-13), *Programa Nacional de Excelência (PRONEX)* and contract 407873/2018-0 of the *Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq)*, the *Fundação de Amparo à Pesquisa do Estado de Minas Gerais (Fapemig* CBB-APQ-02620-15*)* and the *Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado do Rio de Janeiro (Faperj)*, Brazil. CNPq supports CFAB, CTDR, MFFC, PB and RLO, with a research productivity fellowship. CTDR (CNE: E-26/202.921/2018), MFFC, PB and RLO are also supported by Faperj as *Cientistas do nosso estado*. AdP-C was supported by a postdoctoral fellowship from the Faperj and DAMA by a fellowship from the CGZV-SVS (Brazilian Ministry of Health) TED 49/2018 grant. SF was supported by a Wellcome Seed Award in Science to DCJ (208965/Z/17/Z).

Author contributions

CTDR, PB, AdPC, RLdO, RC and AP conceived the study. CTDR, PB, CFAdB, MdFFdC, RC and AP supervised and co-ordinated the study. AdPC, FVSdA, DAMA, CBJ, JCdSJ and ZMBH collected materials. OD, QG, AdPC, CFAdB, MdFFdC, FVSdA, DAMA, CBJ, JCsSJ and ZMBH conducted wet-lab experiments. TM, AK, SF, DCJ, FJGV, SA, CTDR, PB, RLdO, CFAdB, MdFFdC, FVSdA, DAMA, CBJ, JCdSJ, ZMBH, RC and AP analysed and interpreted the data. TM, RC, AP, CTDR, PB, CFAdB and RLdO drafted and edited the manuscript. All authors read and approved the final manuscript.

References

1   Brasil, P. *et al.* Outbreak of human malaria caused by Plasmodium simium in the Atlantic
Forest in Rio de Janeiro: a molecular epidemiological investigation. *Lancet Glob Health* **5**,
e1038-e1046, doi:10.1016/S2214-109X(17)30333-9 (2017).

2   Cox-Singh, J. *et al.* Plasmodium knowlesi malaria in humans is widely distributed and
potentially life threatening. *Clinical infectious diseases : an official publication of the*
Infectious Diseases Society of America **46**, 165-171, doi:10.1086/524888 (2008).

3   Imwong, M. *et al.* Asymptomatic Natural Human Infections With the Simian Malaria
Parasites Plasmodium cynomolgi and Plasmodium knowlesi. *J Infect Dis* **219**, 695-702,
doi:10.1093/infdis/jiy519 (2019).

4   Loy, D. E. *et al.* Out of Africa: origins and evolution of the human malaria parasites
Plasmodium falciparum and Plasmodium vivax. *Int J Parasitol* **47**, 87-97,
doi:10.1016/j.ijpara.2016.05.008 (2017).

5   Liu, W. *et al.* Origin of the human malaria parasite Plasmodium falciparum in gorillas. *Nature*
467, 420-425, doi:10.1038/nature09442 (2010).

6   Liu, W. *et al.* African origin of the malaria parasite Plasmodium vivax. *Nature*
communications **5**, 3346, doi:10.1038/ncomms4346 (2014).

7   Marques, G. R., Condino, M. L., Serpa, L. L. & Cursino, T. V. [Epidemiological aspects of
autochthonous malaria in the Atlantic forest area of the northern coast of the State of Sao
Paulo, 1985-2006]. *Rev Soc Bras Med Trop* **41**, 386-389 (2008).

8   Cox-Singh, J. *et al.* Plasmodium knowlesi malaria in humans is widely distributed and
potentially life threatening. *Clin Infect Dis* **46**, 165-171, doi:10.1086/524888 (2008).

9   Ta, T. H. *et al.* First case of a naturally acquired human infection with Plasmodium
cynomolgi. *Malar J* **13**, 68, doi:10.1186/1475-2875-13-68 (2014).

10   Marchand, R. P., Culleton, R., Maeno, Y., Quang, N. T. & Nakazawa, S. Co-infections of
Plasmodium knowlesi, P. falciparum, and P. vivax among Humans and Anopheles dirus
Mosquitoes, Southern Vietnam. *Emerg Infect Dis* **17**, 1232-1239, doi:10.3201/eid1707.101551
(2011).

11   Otto, T. D. *et al.* Genomes of all known members of a Plasmodium subgenus reveal paths to
virulent human malaria. *Nature microbiology* **3**, 687-697, doi:10.1038/s41564-018-0162-2
(2018).

12   de Pina-Costa, A. *et al.* Malaria in Brazil: what happens outside the Amazonian endemic
region. *Mem Inst Oswaldo Cruz* **109**, 618-633, doi:10.1590/0074-0276140228 (2014).

13   Collins, W. E., Contacos, P. G. & Guinn, E. G. Observations on the sporogonic cycle and
transmission of Plasmodium simium Da Fonseca. *J Parasitol* **55**, 814-816 (1969).

14   de Alvarenga, D. A. M. *et al.* An assay for the identification of Plasmodium simium infection
for diagnosis of zoonotic malaria in the Brazilian Atlantic Forest. *Scientific reports* **8**, 86,
doi:10.1038/s41598-017-18216-x (2018).

15   Escalante, A. A. *et al.* A monkey's tale: the origin of Plasmodium vivax as a human malaria
parasite. *Proc Natl Acad Sci U S A* **102**, 1980-1985, doi:10.1073/pnas.0409652102 (2005).

16   Gelabert, P. *et al.* Mitochondrial DNA from the eradicated European Plasmodium vivax and P.
falciparum from 70-year-old slides from the Ebro Delta in Spain. *Proceedings of the National*
Academy of Sciences of the United States of America **113**, 11495-11500,
doi:10.1073/pnas.1611017113 (2016).

17   Rodrigues, P. T. *et al.* Human migration and the spread of malaria parasites to the New World.
Scientific reports **8**, 1993, doi:10.1038/s41598-018-19554-0 (2018).

18   Deane, L. M. Simian malaria in Brazil. *Mem Inst Oswaldo Cruz* **87 Suppl 3**, 1-20 (1992).

19   Fonseca, F. [Plasmodium of a primate of Brazil]. *Mem Inst Oswaldo Cruz* **49**, 543-553 (1951).

20   Leclerc, M. C. *et al.* Meager genetic variability of the human malaria agent Plasmodium
vivax. *Proceedings of the National Academy of Sciences of the United States of America* **101**,
14455-14460, doi:10.1073/pnas.0405186101 (2004).

21   Duarte, A. M. *et al.* Widespread occurrence of antibodies against circumsporozoite protein
and against blood forms of Plasmodium vivax, P. falciparum and P. malariae in Brazilian wild
monkeys. *J Med Primatol* **35**, 87-96, doi:10.1111/j.1600-0684.2006.00148.x (2006).

22   Tazi, L. & Ayala, F. J. Unresolved direction of host transfer of Plasmodium vivax v. P.
simium and P. malariae v. P. brasilianum. *Infect Genet Evol* **11**, 209-221,
doi:10.1016/j.meegid.2010.08.007 (2011).

23   Coatney, G. R., Collins, W. E., Warren, M. & Contacos, P. G. in *The Primate Malarias*
(National Institutes of Health, 1971).

24   Auburn, S. *et al.* A new Plasmodium vivax reference sequence with improved assembly of the
subtelomeres reveals an abundance of pir genes. *Wellcome Open Res* **1**, 4,
doi:10.12688/wellcomeopenres.9876.1 (2016).

25   Hupalo, D. N. *et al.* Population genomics studies identify signatures of global dispersal and
drug resistance in Plasmodium vivax. *Nat Genet* **48**, 953-958, doi:10.1038/ng.3588 (2016).

26   Alexander, D. H., Novembre, J. & Lange, K. Fast model-based estimation of ancestry in
unrelated individuals. *Genome research* **19**, 1655-1664, doi:10.1101/gr.094052.109 (2009).

27   Willing, E. M., Dreyer, C. & van Oosterhout, C. Estimates of genetic differentiation measured
by F(ST) do not necessarily require large sample sizes when using many SNP markers. *PloS*
one **7**, e42649, doi:10.1371/journal.pone.0042649 (2012).

28   Cunningham, D., Lawton, J., Jarra, W., Preiser, P. & Langhorne, J. The pir multigene family
of Plasmodium: antigenic variation and beyond. *Mol Biochem Parasitol* **170**, 65-73,
doi:10.1016/j.molbiopara.2009.12.010 (2010).

29   Warncke, J. D., Vakonakis, I. & Beck, H. P. Plasmodium Helical Interspersed Subtelomeric
(PHIST) Proteins, at the Center of Host Cell Remodeling. *Microbiol Mol Biol Rev* **80**, 905-
927, doi:10.1128/MMBR.00014-16 (2016).

30   Castillo, A. I., Andreina Pacheco, M. & Escalante, A. A. Evolution of the merozoite surface
protein 7 (msp7) family in Plasmodium vivax and P. falciparum: A comparative approach.
Infect Genet Evol **50**, 7-19, doi:10.1016/j.meegid.2017.01.024 (2017).

31   Ntumngia, F. B. *et al.* Characterisation of a tryptophan-rich Plasmodium falciparum antigen
associated with merozoites. *Mol Biochem Parasitol* **137**, 349-353,
doi:10.1016/j.molbiopara.2004.06.008 (2004).

32   Kanjee, U., Rangel, G. W., Clark, M. A. & Duraisingh, M. T. Molecular and cellular
interactions defining the tropism of Plasmodium vivax for reticulocytes. *Curr Opin Microbiol*
46, 109-115, doi:10.1016/j.mib.2018.10.002 (2018).

33   Miller, L. H., McAuliffe, F. M. & Mason, S. J. Erythrocyte receptors for malaria merozoites.
Am J Trop Med Hyg **26**, 204-208, doi:10.4269/ajtmh.1977.26.204 (1977).

34   Iyer, J., Gruner, A. C., Renia, L., Snounou, G. & Preiser, P. R. Invasion of host cells by
malaria parasites: a tale of two protein families. *Mol Microbiol* **65**, 231-249,
doi:10.1111/j.1365-2958.2007.05791.x (2007).

35   Chan, L. J., Dietrich, M. H., Nguitragool, W. & Tham, W. H. Plasmodium vivax Reticulocyte
Binding Proteins for invasion into reticulocytes. *Cell Microbiol*, e13110,
doi:10.1111/cmi.13110 (2019).

36   Gruszczyk, J. *et al.* Cryo-EM structure of an essential Plasmodium vivax invasion complex.
Nature **559**, 135-139, doi:10.1038/s41586-018-0249-1 (2018).

37   Carlton, J. M. *et al.* Comparative genomics of the neglected human malaria parasite
Plasmodium vivax. *Nature* **455**, 757-763, doi:10.1038/nature07327 (2008).

38   Gilabert, A. *et al.* Plasmodium vivax-like genome sequences shed new insights into
Plasmodium vivax biology and evolution. *PLoS Biol* **16**, e2006035,
doi:10.1371/journal.pbio.2006035 (2018).

39   Ntumngia, F. B. *et al.* Genetic variation among Plasmodium vivax isolates adapted to non-
human primates and the implication for vaccine development. *Am J Trop Med Hyg* **80**, 218-
227 (2009).

40   Rausch, T. *et al.* DELLY: structural variant discovery by integrated paired-end and split-read
analysis. *Bioinformatics* **28**, i333-i339, doi:10.1093/bioinformatics/bts378 (2012).

41   Batchelor, J. D., Zahm, J. A. & Tolia, N. H. Dimerization of Plasmodium vivax DBP is
induced upon receptor binding and drives recognition of DARC. *Nat Struct Mol Biol* **18**, 908-
914, doi:10.1038/nsmb.2088 (2011).

42   Jeffares, D. C., Tomiczek, B., Sojo, V. & dos Reis, M. A beginners guide to estimating the
non-synonymous to synonymous rate ratio of all protein-coding genes in a genome. *Methods*
Mol Biol **1201**, 65-90, doi:10.1007/978-1-4939-1438-8_4 (2015).

43   Proto, W. R. *et al.* Adaptation of Plasmodium falciparum to humans involved the loss of an
ape-specific erythrocyte invasion ligand. *Nature communications* **10**, 4512,
doi:10.1038/s41467-019-12294-3 (2019).

44   Wright, P. E. & Dyson, H. J. Intrinsically disordered proteins in cellular signalling and
regulation. *Nature reviews. Molecular cell biology* **16**, 18-29, doi:10.1038/nrm3920 (2015).

45   Culleton, R. & Carter, R. African Plasmodium vivax: distribution and origins. *Int J Parasitol*
42, 1091-1097, doi:10.1016/j.ijpara.2012.08.005 (2012).

46   Li, J. *et al.* Geographic subdivision of the range of the malaria parasite Plasmodium vivax.
Emerging infectious diseases **7**, 35-42, doi:10.3201/eid0701.010105 (2001).

47 Choi, E. H., Lee, S. K., Ihm, C. & Sohn, Y. H. Rapid DNA extraction from dried blood spots
on filter paper: potential applications in biobanking. *Osong Public Health Res Perspect* **5**,
351-357, doi:10.1016/j.phrp.2014.09.005 (2014).

48 Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114-2120, doi:10.1093/bioinformatics/btu170 (2014).

49 Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nature methods*
9, 357-359, doi:10.1038/nmeth.1923 (2012).

50 Bankevich, A. *et al.* SPAdes: a new genome assembly algorithm and its applications to single-
cell sequencing. *J Comput Biol* **19**, 455-477, doi:10.1089/cmb.2012.0021 (2012).

51 Boetzer, M., Henkel, C. V., Jansen, H. J., Butler, D. & Pirovano, W. Scaffolding pre-
assembled contigs using SSPACE. *Bioinformatics* **27**, 578-579,
doi:10.1093/bioinformatics/btq683 (2011).

52 Boetzer, M. & Pirovano, W. Toward almost closed genomes with GapFiller. *Genome biology*
13, R56, doi:10.1186/gb-2012-13-6-r56 (2012).

53 Walker, B. J. *et al.* Pilon: an integrated tool for comprehensive microbial variant detection and
genome assembly improvement. *PloS one* **9**, e112963, doi:10.1371/journal.pone.0112963
(2014).

54 Laetsch, D. & Blaxter, M. BlobTools: Interrogation of genome assemblies [version 1;
referees: 2 approved with reservations]. *F1000Research* **6**,
doi:10.12688/f1000research.12232.1 (2017).

55 Jongwutiwes, S. *et al.* Mitochondrial genome sequences support ancient population expansion
in Plasmodium vivax. *Molecular biology and evolution* **22**, 1733-1739,
doi:10.1093/molbev/msi168 (2005).

56 Rodrigues, P. T. *et al.* Using mitochondrial genome sequences to track the origin of imported
Plasmodium vivax infections diagnosed in the United States. *Am J Trop Med Hyg* **90**, 1102-
1108, doi:10.4269/ajtmh.13-0588 (2014).

57 Bandelt, H. J., Forster, P. & Rohl, A. Median-joining networks for inferring intraspecific
phylogenies. *Molecular biology and evolution* **16**, 37-48,
doi:10.1093/oxfordjournals.molbev.a026036 (1999).

58   Leigh, J. W. & Bryant, D. POPART: full-feature software for haplotype network construction.
Methods Ecol Evol **6**, 1110-1116, doi:10.1111/2041-210x.12410 (2015).

59   Holt, C. & Yandell, M. MAKER2: an annotation pipeline and genome-database management
tool for second-generation genome projects. *BMC Bioinformatics* **12**, 491, doi:10.1186/1471-
2105-12-491 (2011).

60   Steinbiss, S. *et al.* Companion: a web server for annotation and analysis of parasite genomes.
Nucleic acids research **44**, W29-34, doi:10.1093/nar/gkw292 (2016).

61   Waterhouse, R. M. *et al.* BUSCO applications from quality assessments to gene prediction
and phylogenomics. *Molecular biology and evolution*, doi:10.1093/molbev/msx319 (2017).

62   Simao, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V. & Zdobnov, E. M.
BUSCO: assessing genome assembly and annotation completeness with single-copy
orthologs. *Bioinformatics* **31**, 3210-3212, doi:10.1093/bioinformatics/btv351 (2015).

63   Lechner, M. *et al.* Proteinortho: detection of (co-)orthologs in large-scale analysis. *BMC*
Bioinformatics **12**, 124, doi:10.1186/1471-2105-12-124 (2011).

64   Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Molecular biology and evolution* **30**, 772-780,
doi:10.1093/molbev/mst010 (2013).

65   Capella-Gutierrez, S., Silla-Martinez, J. M. & Gabaldon, T. trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**, 1972-1973,
doi:10.1093/bioinformatics/btp348 (2009).

66   Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
phylogenies. *Bioinformatics* **30**, 1312-1313, doi:10.1093/bioinformatics/btu033 (2014).

67   Sedlazeck, F. J., Rescheneder, P. & von Haeseler, A. NextGenMap: fast and accurate read
mapping in highly polymorphic genomes. *Bioinformatics* **29**, 2790-2791,
doi:10.1093/bioinformatics/btt468 (2013).

68   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078-
2079, doi:btp352 [pii]
10.1093/bioinformatics/btp352 (2009).

69   McKenna, A. *et al.* The Genome Analysis Toolkit: a MapReduce framework for analyzing
next-generation DNA sequencing data. *Genome research* **20**, 1297-1303,
doi:10.1101/gr.107524.110 (2010).

70   Garrison, E. & Marth, G. Haplotype-based variant detection from short-read sequencing.
arXiv, 1207.3907 (2012).

71   Chang, C. C. *et al.* Second-generation PLINK: rising to the challenge of larger and richer
datasets. *Gigascience* **4**, 7, doi:10.1186/s13742-015-0047-8 (2015).

72   Pfeifer, B., Wittelsburger, U., Ramos-Onsins, S. E. & Lercher, M. J. PopGenome: an efficient
Swiss army knife for population genomic analyses in R. *Molecular biology and evolution* **31**,
1929-1936, doi:10.1093/molbev/msu136 (2014).

73   R Development Core Team. R: A language and environment for statistical computing. R
Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL
http://www.R-project.org.   (2007).

74   Weir, B. S. & Cockerham, C. C. Estimating F-Statistics for the Analysis of Population
Structure. *Evolution* **38**, 1358-1370, doi:10.1111/j.1558-5646.1984.tb05657.x (1984).

75   Cingolani, P. *et al.* A program for annotating and predicting the effects of single nucleotide
polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain w1118; iso-2;
iso-3. *Fly (Austin)* **6**, 80-92, doi:10.4161/fly.19695 (2012).

76   Guindon, S. *et al.* New algorithms and methods to estimate maximum-likelihood phylogenies:
assessing the performance of PhyML 3.0. *Syst Biol* **59**, 307-321, doi:10.1093/sysbio/syq010
(2010).

77   Guindon, S. & Gascuel, O. A simple, fast, and accurate algorithm to estimate large
phylogenies by maximum likelihood. *Syst Biol* **52**, 696-704, doi:10.1080/10635150390235520
(2003).

78   Lefort, V., Longueville, J. E. & Gascuel, O. SMS: Smart Model Selection in PhyML.
Molecular biology and evolution **34**, 2422-2424, doi:10.1093/molbev/msx149 (2017).

79   Anisimova, M., Gil, M., Dufayard, J. F., Dessimoz, C. & Gascuel, O. Survey of branch
support methods demonstrates accuracy, power, and robustness of fast likelihood-based
approximation schemes. *Syst Biol* **60**, 685-699, doi:10.1093/sysbio/syr041 (2011).

80　Huson, D. H. & Bryant, D. Application of phylogenetic networks in evolutionary studies.
Molecular biology and evolution **23**, 254-267, doi:10.1093/molbev/msj030 (2006).

81　Bryant, D. & Moulton, V. Neighbor-net: an agglomerative method for the construction of
phylogenetic networks. *Molecular biology and evolution* **21**, 255-265,
doi:10.1093/molbev/msh018 (2004).

82　Larkin, M. A. *et al.* Clustal W and Clustal X version 2.0. *Bioinformatics* **23**, 2947-2948,
doi:10.1093/bioinformatics/btm404 (2007).

83　Thompson, J. D., Higgins, D. G. & Gibson, T. J. CLUSTAL W: improving the sensitivity of
progressive multiple sequence alignment through sequence weighting, position-specific gap
penalties and weight matrix choice. *Nucleic acids research* **22**, 4673-4680 (1994).

84　Chaisson, M. J. & Tesler, G. Mapping single molecule sequencing reads using basic local
alignment with successive refinement (BLASR): application and theory. *BMC Bioinformatics*
13, 238, doi:10.1186/1471-2105-13-238 (2012).

85　Seibt, K. M., Schmidt, T. & Heitkam, T. FlexiDot: highly customizable, ambiguity-aware
dotplots for visual sequence analyses. *Bioinformatics* **34**, 3575-3577,
doi:10.1093/bioinformatics/bty395 (2018).

86　van Ooij, C. *et al.* The malaria secretome: from algorithms to essential function in blood stage
infection. *PLoS pathogens* **4**, e1000084, doi:10.1371/journal.ppat.1000084 (2008).

87　Boddey, J. A. *et al.* Role of plasmepsin V in export of diverse protein families from the
Plasmodium falciparum exportome. *Traffic* **14**, 532-550, doi:10.1111/tra.12053 (2013).

88　Schulze, J. *et al.* The Plasmodium falciparum exportome contains non-canonical PEXEL/HT
proteins. *Mol Microbiol* **97**, 301-314, doi:10.1111/mmi.13024 (2015).

89　Hu, G. *et al.* Transcriptional profiling of growth perturbations of the human malaria parasite
Plasmodium falciparum. *Nat Biotechnol* **28**, 91-98, doi:10.1038/nbt.1597 (2010).

90　Ye, J. *et al.* Primer-BLAST: a tool to design target-specific primers for polymerase chain
reaction. *BMC Bioinformatics* **13**, 134, doi:10.1186/1471-2105-13-134 (2012).

91　Wang, S., Li, W., Liu, S. & Xu, J. RaptorX-Property: a web server for protein structure
property prediction. *Nucleic acids research* **44**, W430-435, doi:10.1093/nar/gkw306 (2016).

92   Waterhouse, A. *et al.* SWISS-MODEL: homology modelling of protein structures and

1009   complexes. *Nucleic acids research* **46**, W296-W303, doi:10.1093/nar/gky427 (2018).
1010

1011

1012

Figure Legends

1013

Figure 1 - SNP phylogeny

A) Mid-point rooted maximum likelihood tree produced from 143,123 concatenated

SNP positions with data from at least 55 samples. The tree was produced using

PhyML with the GTR evolutionary model. Branch support was evaluated with the

Bayesian-like transformation of approximate likelihood ratio test (aBayes). Genetic

distance shown below tree. *P. vivax* isolates are denoted as colored circles by their

country of sample origin. A tree with specific sample IDs is available in Figure S7. B)

Magnification of the *P. simium* clade (as in panel A). C) Map denoting the geographic

location of *P. simium* samples.

1023

Figure 2 - Nucleotide diversity

Box plot showing the pair-wise nucleotide diversity between human-infecting *P.

simium* samples (left), *P. vivax* samples (middle), and between *P. simium* and *P. vivax*

samples (right). Diversity is shown for entire genome (left-most plots, blue) and

exonic regions only (right-most plots, orange). Individual values from pairwise

comparsions are shown as grey dots, boxes denote 25th and 75th percentiles, and

whiskers an additional 1.5 interquartile lengths. The observed nucleotide diversity

between *P. simium* and *P. vivax* samples is significantly higher than between *P. vivax*

samples (Mann-Whitney U, genome; $p=8.79\times10^{-20}$, exons; $p=3.72\times10^{-20}$).

1033

Figure 3 - F_{ST} ratios

A) Gene F_{ST} values were calculated between simium and American vivax samples (x-

axis) and American and Asian vivax samples (y-axis). Each dot corresponds to a

gene, and the distributions of the two FST measures are shown as bar charts above

and to the right of the scatter plot. B) The ratio between F_{ST} values between i) simium

and American vivax samples, and ii) American and Asian vivax samples were

calculated for each gene (top). A pseudo count of one was added to all F_{ST} values.

The distributions of log2-ratios are shown as violin plots (bottom) for all genes (grey),

PIR genes (red), PHIST genes (brown), TRAg genes (turquoise), invasion genes

(yellow), and exported genes (dark blue). Only genes with SNP differences between

the three populations are included in this analysis. F_{ST} values and ratios are provided

in Table S7. P-values from Mann-Whitney U tests for differences in medians between

PIR, PHIST, TRAg, invasion genes, exported genes, and all remaining genes are

indicated on plot (e.g. PIR genes are compared to all non-PIR genes).

1048

Figure 4 - Invasome deletions

A) Overview of the invasome gene groups, Reticulocyte Binding Proteins (RBPs) and

Duffy Binding Proteins (DBPs) in *Plasmodium vivax* and *P. simium*. The *P. vivax*

genome harbours two RBP2d genes, one of which is a pseudogene (Table S9). B)

Schematic depiction of samples with and without the deletion found in DBP1.

C) Left: Structural rendering of DBP1, showing known structural domains and motifs.

The two fragment molecules from the human DARC receptor are shown in grey. The

3-dimensional structure of the DBL-DARC complex was modeled based on the *P.*

vivax crystallographic model (PDB 4nuv). The region deleted in sequences from

human-infecting *P. simium*, as compared to *P. vivax* P01, is highlighted in red. Right:

Details of DBP1 protein alignments. A full alignment is available in Figure S20.

D) Similar to panel C) but for RBP2a. The complex between the reticulocyte binding

domain and the human receptor was modeled based on the cryoEM structure of the

complex between the *P. vivax* RBP2b and the human transferrin receptor TfR1 (PDB

6d05). A full alignment is available in Figure S21.
1064

1065

Figure 1

Figure 2

Figure 3

Figure 4